Comments to the Author:	
Dear authors,	
Thank you very much for your adjusted manuscript. After carefully reviewing the revised manuscript we come to the conclusions that a further revision is needed for the clarity of the paper for the readers benefit.	We appreciated your comments and a happy to provide an improved manuscript.
From the editor's perspective and from the suggestion from reviewer 2 the manuscript still needs to be reorganized to have a clearer structure and focus. Please clarify what each section focuses on, clarify to which results you refer to. Discuss each point which is valid (e.g. effect of properties, different databases, different wetlands,and clearly separate processing and results and discussion. The manuscript currently it's a mixture of methods, results and discussions for example found at L325. The authors refer to a method not described elsewhere, but at the same time referring to results and discussions in the following lines. Additionally, the editor suggest to make clearer	We reworded the end of the introduction to signal to the reader the layout of the paper. We restructured the paper into focused sections 1 Introduction—2 Comparison of databases— 3 Processing of HWSD—4 Spatial distribution—5 Discussion of SOC masses— 6 Conclusions. We marked "external"
throughout the whole manuscript where the authors refer to HWSD data and where to their own calculated data (e.g. by using specific acronym?). Currently there is too much interpretation freedom to which datasets is referred to (e.g. L271).	values with an asterisk in section 5.
The provision of the R-script for some calculations is certainly an asset to the paper. However, the current script cannot be used to reproduce the results as a) input data are not defined (e.g. URL), b) the content of the script contains R- commands while the heading suggest that the GRASS commands are described- the editor questions where is the GRASS part and where are the libraries in grass loaded to do some of the described processing. Are all parts of the analysis contained in the supplementary document for L229- 233? Please discuss and revise.	The script produces tables for import into GRASS. The script does not describe the elementary GIS procedures (in our view) for producing the individual results. We improved the script by adding a link to the HWSD for download and indicating which GRASS command is needed for loading the data into GRASS. We changed the wording within the script that may have implied that the script also includes GIS scripts.
Minor issues:	
L116 Suggestion: Review of Existing spatial databases	Chapter title changed to "Comparison of estimates

	of global SOC mass among existing spatial
	databases."
L25 BD - if first time us of acronym, please introduce. Please recheck manuscript for occurrences	Checked and changed.
L40-44. Please provide reference to each of this statements	Added.
L169 URL Spatial-analyst.net - while I certainly appreciate TH effort to provide data, a proper reference to the original data should be provided.	The reference to the original data is the one given (Reich 2000) and listed with its own URL (by now outdated, has been updated in our 2nd revision). The US NRC site provides only a JPG image of a wall map, hence the additional link to a source for the data.
L189 remove '(' before HWSD	Changed.
Table 1 - extra column with acronyms, revise table to ensure all definitions are included	Added column and acronyms; we omitted the terms for the equation in line 79 because they are used only in that instance.
Table 2 - extra column with the processing steps. What is processing step (-) ? Please clarify if the values are additive or singular. If singular, what happens if they are additive, and vice versa. Please discuss	We rephrased the text and and modified the table. In our opinion only the choice between applying the correction for organic soils and Histosols is meaningful to differentiate. We do so in what we now call step 3b, which was maybe not obvious in the previous text.
Please recheck all references in the manuscript. There them to be missing ones (e.g. ESRI, 2002)	Added.
Please clarify what is the goal of L415-420 and revise it where does the ³ / ₄ come from ?	Page et al. estimated the SOC down to a depth of 4 m, the HWSD covers the top 1 m, so we need only the lower 3 m here.

- Global distribution of soil organic carbon Part 1: Masses and
- ² frequency distributions of SOC stocks for the tropics,
- ³ permafrost regions, wetlands, and the world
- 4
- 5 Martin KÖCHY^{1,a}, Roland HIEDERER², Annette FREIBAUER³
- 6
- 7 [1]{Thünen Institute of Climate-Smart Agriculture, Bundesallee 50, 38116 Braunschweig,
- 8 Germany, Germany}
- 9 [2]{Joint Research Centre, Institute for Environment and Sustainability, Via E. Fermi 2749,

10 21027 Ispra (VA), Italy}

- 11 [3]{Thünen Institute of Climate-Smart Agriculture, Bundesallee 50, 38116 Braunschweig,
- 12 Germany}
- [a] {now at: Thünen Institute of Market Analysis, Bundesallee 50, 38116 Braunschweig,
- 14 Germany}
- 15
- 16 Correspondence to: Martin Köchy (office@martinkoechy.de)

18 Abstract

The global soil organic carbon (SOC) mass is relevant for the carbon cycle budget and thus 19 atmospheric carbon concentrations. We review current estimates of SOC stocks and mass 20 (stock × area) in wetlands, permafrost and tropical regions and the world in the upper 1 m of 21 soil. The Harmonized World Soil Database (HWSD) v.1.2 provides one of the most recent 22 and coherent global data sets of SOC, giving a total mass of 2476 Pg when using the original values for bulk density. Adjusting the HWSD's bulk density of soil high in organic carbon 24 results in a mass of 1230 Pg, and additionally setting the BD of Histosols to 0.1 g cm⁻³ (typical of peat soils) results in a mass of 1062 Pg. The uncertainty of bulk density of 26 Histosols alone introduces a range of -56 to +180 Pg C for the estimate of global SOC mass in the top 1 meter, larger than estimates of global soil respiration. We report the spatial 28 distribution of SOC stocks per 0.5 arc minutes, the areal masses of SOC and the quantiles of 29 SOC stocks by continents, wetland types, and permafrost types. Depending on the definition 30 of 'wetland', wetland soils contain between 82 and 158 Pg SOC. Incorporating more detailed estimates for permafrost from the Northern Circumpolar Soil Carbon Data Base (496 Pg SOC) and tropical peatland carbon, global soils contain 1325 Pg SOC in the upper 1 m including 421 Pg in tropical soils, whereof 40 Pg occur in tropical wetlands. Global SOC 34 amounts to just under 3000 Pg when estimates for deeper soil layers are included. Variability 35 in estimates is due to variation in definitions of soil units, differences in soil property 36 databases, scarcity of information about soil carbon at depths >1 m in peatlands, and variation 37 in definitions of 'peatland'. 38

39 1. Introduction

The global mass of soil organic carbon (SOC; for a list of term and acronyms see Table 1) is greater than the combined mass of carbon (C) contained in the atmosphere and in the living biomass (Ciais et al., 2013). Therefore, small relative changes in the mass of SOC can have profound effects on the concentration of atmospheric CO₂ and hence climate change (Myhre et al., 2013). Despite its importance, the global mass of SOC (Scharlemann et al, 2014) and its distribution in space and among land-use/land-cover classes is not well known (Jandl et al., 2014).

47 On the short to middle term (decades), variation in SOC mass is strongly related to the balance of input from net primary production and microbial decomposition. On longer time-48 scales, however, changes in the decomposable mass of SOC affect this balance. Globally, the 49 largest SOC stocks are located in wetlands and peatlands, most of which occur in regions of permafrost and in the tropics. Decomposition rates in wetlands and permafrost are low due to low availability of oxygen and low temperatures, respectively. This SOC is vulnerable to 53 changes in the hydrological cycle as well as to changes in permafrost dynamics. 54 A good knowledge of the global SOC mass and its spatial distribution is necessary for assessing, in an international context, where soils are most vulnerable to C losses or which land use/land cover types might provide the best opportunity for C sequestration to mitigate 56 increases in greenhouse gas concentrations. Since SOC mass is a product of several factors, uncertainty (or errors in measurement) in one of the factors affects all others. Consequently, 58

59 the measures to reduce the uncertainty of global SOC mass should be directed to those soils

 $_{60}$ that are associated with a large extent (area), high levels of C_{org} , low bulk density (BD) or

great depth. Variations at the lower end of BD are more consequential than at the high end of

BD because low BD is associated with organic soils (high C_{org}) and a change from, say, 0.1 to

3

Martin Köchy 2015·2·26 15:25 **Gelöscht:** productivity 64 0.2 leads to a doubling of SOC stock and mass. Variation within the range of BD typical of

65 mineral soils, <u>e.g.</u>, $1.2 - 1.8 \text{ g cm}^{-3}$, is less consequential.

66	The spatial distribution of SOC stocks is typically derived from maps (printed or electronic)
67	where areas with similar soil characteristics are aggregated to form soil units, and the SOC
68	mass of the area of the soil unit is calculated by multiplication of the area of the soil unit with
69	its unit-area SOC stock (Amundson, 2001). Historically, soil maps have been compiled
70	largely based on the experience of soil surveyors, taking into account topography, climate,
71	land use history, land management, vegetation, parent material, and soil typical characteristics
72	(McBratney et al., 2003b). The spatial soil units are linked to their defining properties, which
73	are based on measurements of soil profiles or an evaluation by experts. Typically,
74	measurements from several profiles within the same soil unit have been statistically
75	aggregated (e.g., averaged). Missing profile data may be estimated by pedotransfer functions
76	(PTFs) from other measured soil characteristics.
77	The SOC stock, $m_{\rm C}$, of a soil column is calculated by integrating the areal density of SOC
78	over all vertical depth layers (or within a specified depth). The areal density of SOC of a soil
79	layer is determined by measuring the organic carbon concentration (C_{org}) and the <u>BD</u> of
80	undisturbed soil samples in homogenous layers of thickness d (Table 1). The areal density,
81	$C_{org} \times BD \times d$, is reduced by the fractional volume f_G occupied by gravel, rocks, roots, and ice
82	in the soil layer, or $m_{\rm C} = C_{\rm org} \times BD \times (1 - f_{\rm G}) \times d$. The SOC mass of the area (A) is the product
83	of the soil unit's area and its SOC density (A $\times m_{\rm G}$). Lateral variation, temporal variation, and
84	methodological differences in measuring any of the necessary soil characteristics (BD, C_{org} ,
85	volume of gravel and roots, forms of C, depth) contribute to the variability of SOC stock and
86	mass estimates (Ellert et al., 2001).

4

Martin Köchy 2015-2-26 15:25 Gelöscht: Traditionally, maps of the Martin Köchy 2015-2-26 15:25 Gelöscht: are Martin Köchy 2015-2-26 15:25 Gelöscht: mapping Martin Köchy 2015-2-26 15:26 Gelöscht: mapping Martin Köchy 2015-2-26 15:26 Gelöscht: mapping

Martin Köchy 2015·2·26 15:26 Gelöscht: mapping

Martin Köchy 2015·2·27 09:08 Gelöscht: bulk density Martin Köchy 2015·2·19 15:59 Gelöscht: (BD)

Martin Köchy 2015·2·26 15:26 Gelöscht: stock Martin Köchy 2015·2·26 17:58 Gelöscht: × A Martin Köchy 2015·2·26 15:26 Gelöscht: variability Martin Köchy 2015·2·26 15:26 Gelöscht: variability

99	The accuracy of spatially interpolated maps of SOC stocks depends on how well the soil units
100	are represented by soil profiles with complete characteristics. The latest WISE database
101	(v.3.1) contains harmonized data of more than 10250 soil profiles (Batjes, 2009), which,
102	however, underrepresent the the non-agricultural areas of North America, the Nordic
103	countries, most parts of Asia (notably Iran, Kazakhstan, and Russia), Northern Africa, and
104	Australia. To calculate SOC stocks one needs Corg, BD, soil depth, and volumetric gravel
105	fraction. These are provided individually by 87%, 32%, 100%, and 22%, respectively, of the
106	profiles (Batjes, 2009). BD and gravel fraction have low representation because they are
107	seldom recorded during routine soil surveys. In numbers, 9970 profile descriptions include
108	C_{org} in at least one layer, but of these only 3655 also include BD. Gravel fraction is explicitly
109	indicated for 1100 of the 3655 profiles but earlier versions of the database could not
110	distinguish between zero and absence of value. BD is included for 806 profiles where $C_{org} >$
111	3% and for 74 profiles where $C_{org} > 20\%$. The temporal origin of profile descriptions ranges
112	from 1925 to 2005. The early data may no longer reflect current conditions where C input and
113	decomposition rates may have changed. Efforts to expand the database of data-rich soil
114	profiles and to use pedotransfer instead of taxotransfer functions has been going on since
115	1986 through the SOTER program (http://www.isric.org/projects/soil-and-terrain-database-
116	soter-programme, accessed 2014-07-07, Nachtergaele, 1999).
117	In this paper we review estimates of the global SOC mass in the top 1 m of soil derived from
118	spatial databases (maps) and additional sources. First, we compare the Harmonized World
119	Soil Database (HWSD, FAO et al., 2012) to earlier spatial databases. The HWSD was the
120	latest and most detailed inventory at the global scale when this study was begun and is still
121	widely used as an international reference (e.g., Wieder et al., 2014, Yan et al., 2014) Next. we
122	describe the adjustments, especially those of BDs of organic soils (Hiederer & Köchy 2011).
123	that are necessary for calculating the SOC stocks from the HWSD. Based on the adjusted

Martin Köchy 2015·2·27 09:10
Gelöscht: mapping
Martin Köchy 2015·2·27 09:10
Gelöscht: ISRIC-
Martin Köchy 2015·2·27 09:13
Gelöscht:
Martin Köchy 2015·2·27 09:13
Gelöscht: The profiles
Martin Köchy 2015·2·27 09:13
Gelöscht: do not yet
Martin Köchy 2015·2·27 09:13
Gelöscht: terrestrial surface equally. Gaps

include

Martin Köchy 2015·2·20 13:04

Martin Köchy 2015-2-20 13:04 Gelöscht: compare previous estimates of the global SOC mass in the top 1 m of soil derived from spatial databases (maps) to the mass derived from the Harmonized World Soil Database (HWSD, FAO et al., 2012), which was the latest and most detailed inventory at the global scale when this study was begun and is still widely used as an international reference (e.g., Wieder et al., 2014, Yan et al., 2014) but requires adjustment of bulk densities of organic soils (Hiederer & Köchy 2011). Based

- 143 HWSD, we report area-weighted frequency distributions of <u>SOC</u> stocks in the top 1m of soil,
- in particular for the large SOC stocks in wetlands, in the tropics, and in frozen soils.
- Frequency distributions can be used to improve the assessment of accuracy in studies where
- 146 SOC is an independent variable. <u>Finally, we update the HWSD-derived global SOC mass for</u>
- 147 the permafrost region and tropical peatlands for the top 1 m and complement it with estimates
- 148 of SOC below 1 m depth. Our conclusions provide recommendations for improving global
- soil mapping.

Martin Köchy 2015·2·26 18:34
Gelöscht: ,
Martin Köchy 2015·2·20 13:15
Gelöscht: our paper
Martin Köchy 2015·2·20 13:15
Gelöscht: s
Martin Köchy 2015·2·26 18:34
Gelöscht: for the first time
Martin Köchy 2015·2·26 18:34
Gelöscht: carbon
Martin Köchy 2015·2·26 18:35
Gelöscht: the
Martin Köchy 2015·2·26 18:35
Gelöscht: permafrost at high latitudes
Martin Köchy 2015·2·20 13:05
Gelöscht: We
Martin Köchy 2015·2·20 13:05
Gelöscht: using best estimates
Martin Köchy 2015·2·20 13:05
Gelöscht: ,
Martin Köchy 2015·2·26 18:36
Gelöscht: the
Martin Köchy 2015·2·26 18:36
Gelöscht: s
Martin Köchy 2015·2·26 18:36
Gelöscht: ,
Martin Köchy 2015·2·20 13:06
Gelöscht: soil
Martin Köchy 2015·2·20 13:06
Gelöscht: from several additional sources
Martin Köchy 2015·2·19 15:57
Gelöscht: <#>Spatial databases of global soil organic carbon mass

- 169 2. Comparison of estimates of global SOC mass among existing spatial databases
- 170 Historic estimates of global SOC mass compared among 27 studies range between 504 and
- 171 <u>3000 Pg with a median of 1461 Pg (Scharlemann et al., 2014). Here we concentrate on</u>
- 172 comparisons with the most recent ones.
- Before the publication of the HWSD, many global estimates were based on the Digital Soil
- 174 Map of the World (DSMW) (FAO, 2007) or its precursor, the Soil Map of the World (SMW,
- FAO, 1997). Batjes (1996), using information from 4353 WISE profiles, reported a range of
- 176 1462–1548 Pg for 0–1 m depth and 2376–2456 Pg for 0–2 m depth. Henry et al. (2009) report
- a global SOC mass of 1589 Pg for the top 1 m and 2521 Pg for the top 2 m (using an
- unspecified WISE version). Hiederer *et al.* (2010) report a slightly lower mass of 1455 Pg for
- 179 DSMW for the top 1 m.
- 180 The International Geosphere Biosphere Program (IGBP) (Global Soil Data Task Group, 2000)
- produced a map of SOC stock on a 5' by 5' grid derived from the DSMW in conjunction with
- WISE data (v.1, 1125 profiles). SOC mass (0–1 m) based on the IGBP map is 1550 Pg
- 183 (calculated as SOC stock \times grid cell area).
- 184 The US Natural Resources Conservation Services reclassified the SMW at 2' and combined it
- 185 with a soil climate map (Reich, 2000, data on a 3' grid -, downloaded from http://spatial-
- analyst.net/worldmaps/SOC.zip). This map shows the distribution of nine classes of SOC
- stocks that result in a global SOC mass (0–1 m) of 1463 Pg. Analyzing 2721 soil profiles
- grouped per biome, Jobbágy & Jackson (2000) estimated that the top 1 m contains 1502 Pg
- 189 SOC, with 491 Pg in 1-2 m and 351 Pg in 2-3 m depth.
- ¹⁹⁰ The recently published Global Soil Dataset for Earth System Models (Shangguan et al., 2014)
- with a resolution of 0.5', combined the DSMW with regional soil maps and global and

Martin Köchy 2015·2·26 16:13

Gelöscht: The global SOC mass calculated directly from the original HWSD (v.1.1) for the upper 1 m of soil is 2476 Pg. Henry *et al.* (2009), using an unspecified earlier version of HWSD, reported a mass of 1850 Pg for the first meter. These high values are, however, due to inconsistencies, gaps, and inaccuracies in the database (Hiederer & Köchy, 2011). The most consequential of the inaccuracies concerns the BD for soils high in C_{org} . In addressing these issues (see Methods), we calculated a global mass of SOC in the top 1 m of soil of 1230 Pg in a first step and 1062Pg in a second step.

206	regional profile databases from several sources beyond those used in the HWSD, including
207	the national databases of the USA, Canada, and Australia. Soil profile data and mapping units
208	were matched in several steps intended to result in the most reliable information. Several
209	harmonization steps are applied to the data to derive amongst others soil carbon
210	concentration, bulk soil density, and gravel content and depth for each soil mapping unit. The
211	global SOC stocks are reported as 1923, 1455 and 720 Pg for the upper 2.3, 1.0, and 0.3 m,
212	respectively.
213	The HWSD (vers. 1.2, FAO et al., 2012) is one of the most recent and most detailed databases
214	at the global scale and widely used as reference. The HWSD contains for the topsoil (0-30
215	cm) and the subsoil (30-100 cm) values for Core, BD and gravel content for dominant and
216	secondary soil types on a raster of 0.5 arc minutes (0.5'). Data sources for HWSD are earlier
217	global soil maps that were published by or in cooperation with FAO, the European Soil
218	Database, the Soil Map of China, SOTER regional studies, WISE profile data, and WISE
219	pedotransfer and taxotransfer functions. The HWSD does not yet include the extensive
220	national databases of USA, Canada, and Australia. The HWSD is the result of associating
221	existing maps of soil types (if necessary reclassified to FAO standards) with soil
222	characteristics derived from the WISE (v.2) database containing about 9600 soil profiles,
223	which is the largest number used for a global soil map until 2013.
224	The HWSD does not quantify variability or ranges of any soil properties within a soil unit. Its
225	description qualifies that "Reliability of the information contained in the database is variable:
226	the parts of the database that still make use of the Soil Map of the World such as North
227	America, Australia, West Africa and South Asia are considered less reliable, while most of
228	the areas covered by SOTER databases are considered to have the highest reliability (Central
229	and Southern Africa, Latin America and the Caribbean, Central and Eastern Europe)."

- 230 The global SOC mass calculated directly from the original HWSD (v.1.2) for the upper 1 m
- 231 of soil is 2476 Pg. Henry et al. (2009), using an unspecified earlier version of HWSD,
- reported a mass of 1850 Pg for the first meter. These high values are, however, due to
- 233 inconsistencies, gaps, and inaccuracies in the database (Hiederer & Köchy, 2011). The most
- 234 consequential of the inaccuracies concerns the BD for soils high in C_{org}. In addressing these
- 235 issues (see next section), we calculated a global mass of SOC in the top 1 m of soil of 1232

- 236 Pg after adjusting the BD of organic soils (SOC > 3%) and 1062 Pg after additionally
- 237 adjusting the BD of Histosols.

238	3. Processing and Adjustment of HWSD data for spatial analyses
239	Our analysis of SOC stocks and masses is based on HWSD vers. 1.1, (FAO et al., 2009)
240	because it was the latest version when this study was begun. Version 1.2 of the HWSD adds
241	two new fields for BD (one for topsoil and one for subsoil) based on the SOTWIS database
242	and addresses minor issues that are listed in detail on the HWSD's web site. Since the
243	resulting differences in global mass between HWSD versions were <0.3% (Table 2), we did
244	not recalculate the other values so that all values reported below are calculated based on
245	version 1.1 of the HWSD and a global mass of 1061 Pg unless explicitly mentioned
246	otherwise.
247	We calculated the SOC stocks for each soil type (s) within a grid cell as the areal density over
248	the thickness of the top and sub soil layer, accounting for the volume occupied by gravel, and
249	weighted it with the soil type's areal fraction in each cell or $m_{\underline{C},\underline{s}} \times \underline{A}_{\underline{s}} / \sum \underline{A}$. Consequently, SOC
250	mass of each cell is the sum over all soil types of the product of SOC stock of each soil type
251	and the fraction of cell area covered by each soil type or $\sum (m_{\underline{c},\underline{s}} \times \underline{A}_{\underline{s}}/\underline{A})$.
252	Despite the harmonization of spatial and attribute data, the HWSD suffers from some residual
253	inconsistencies in the data reported, gaps in some areas covered and errors in the values
254	reported (Hiederer and Köchy, 2011) that required pre-processing of the data. Here we
255	present a correction of overestimated BD values for Histosols contained in the HWSD that
256	was not specifically addressed by Shangguan et al. (2014), Hiederer & Köchy (2011), or
257	Scharlemann et al. (2014). For each processing step the resulting global SOC mass is used as
258	an indication of the magnitude of the data manipulation (Table 2).
259	(Step 1) We filled missing data for C_{ore} in top (4 cases) and subsoil layers (127 cases) with
260	data from cells characterized as the same soil unit and being closest in distance or most

 $\frac{10}{261} \frac{\text{similar in topsoil } C_{\text{org}}(2) \text{ In a similar way, we additionally filled missing values of BD for}{10}$

262	mineral soils in 27 cases. (3a) In HWSD v.1.1 high C_{org} values (>20%) are associated with a
263	BD of 1.1 to 1.4 kg/dm ³ although values of 0.05 to 0.3 kg/dm ³ would be typical of organic
264	soils (Boelter, 1968, Page et al., 2011). To address this issue, we set the topsoil BD to -0.31
265	$\ln(C_{org}[\%]) + 1.38 \ (R^2=0.69) \text{ and subsoil to } -0.32 \ln(C_{org}[\%]) + 1.38 \ (R^2=0.90) \text{ for } C_{org} > 3\%$
266	based on an analysis of the SPADE/M2 soil profile database (Hiederer, 2010). This results in
267	a global mass of 1230 Pg C for a soil depth of up to 1 m. (4) The maximum C _{ore} of Histosols
268	in the HWSD is 47%, resulting in a BD of 0.19 kg/dm ³ for topsoil and 0.15 kg/dm ³ for subsoil
269	using the mentioned equations. In contrast, the best estimate for the BD for tropical peatlands
270	is 0.09 kg/dm ³ (Page et al., 2011), for boreal and subarctic peatland the average BD is 0.112
271	kg/dm ³ (Gorham, 1991), and for Finnish agricultural peat soil the average value is 0.091
272	kg/dm ³ (Mäkkilä, 1994 in Turunen, 2008). Therefore, we finally set BD to 0.1 kg/dm ³ for all
273	Histosols in HWSD. After applying steps 1-4, i.e., the SPADE/M2-based corrections for BD
274	and the modification for Histosols, the global mass of SOC in the upper 1 m of soil is 1061
275	<u>Pg. (3b) If we had adjusted BD only for Histosols and not the other soils with $C_{org} > 3\%$, the</u>
276	global mass would be 1113 Pg. Hiederer & Köchy (2011) used WISE-based corrections for
277	<u>BD</u> with a threshold of $C_{org} > 12\%$ (BD _{top} = -0.285 ln(C_{org} [%])+1.457 and BD _{sub} = -0.291
278	$\ln(C_{org} [\%])+1.389$) that result in a higher a global C mass of 1376 Pg in step 3a but a very
279	similar mass (1062 Pg) after the additional BD correction for histosols in step 4. The
280	processing details for step 1 to 4 are contained in the Supplement.
281	A default reference soil depth of 100 cm is stipulated in the HWSD for each mapping unit as a
282	concession to harmonization of different soil databases. Only Rendzinas, Rankers, Leptosols,
283	and Lithosols are attributed reference soil depths of 30 cm or 10 cm. For most of the
284	remaining soil units the 25-percentile of lowest recorded depth of profiles in the WISE 3.1
285	database is equal to or greater than the reference depth, i.e., SOC stock within the top 1 m is
286	not underestimated by using the reference depth. The 25-percentiles of recorded depths of

287	Calcisols (95 cm, n=218), Cambisols (90 cm, n=1164), Cryosols (80 cm, n=6), Durisols (45
288	cm, n=1), Podsols (80 cm, n=222), Solonchaks (90 cm, n=165), and Umbrisols (49 cm,
289	n=173) are smaller than the reference depths so that C stocks may be overestimated. The
290	overestimate could be substantial for Cryosols, Podsols, and Umbrisols, which have high C_{org}
291	(median >10%). Even though the true soil depth of Cryosols and Podsols can be expected to
292	be deeper than the recorded depth in the databases, this would be of no consequence for the
293	estimated SOC mass of the top 1 m.
294	The HWSD database was pre-processed and analyzed with R (R Development Core Team,
295	2011). We summarized adjusted SOC stocks from HWSD globally and by geographic
296	regions, land cover types, and areas with specific soil characteristics (wetlands, peatlands,
297	permafrost soils). To achieve this we intersected raster maps of SOC with thematic maps in a
298	GIS (GRASS 6.4.2, GRASS Development Team, 2011), calculated SOC mass summed over
299	areas, and determined the 5th, 25th, 50th, 75th, and 95th percentiles of SOC stocks within
300	these areas.
301	

 Martin Köchy 2015-2-20 13:18

 Gelöscht: <#>Processing of HWSD data for spatial analyses .

306 **<u>4.</u>** Spatial distribution of SOC mass <u>based on the adjusted HWSD</u>

- 307 The total SOC mass derived from the unadjusted HWSD v1.2 database and using the
- 308 SOTWIS BD (when available for a soil mapping unit) is 2476 Pg and 1062 Pg after applying
- 309 the BD correction as described in the previous paragraph.

310 4.1. Continental distribution of SOC mass

- The distribution of SOC mass by continents (Table 3) follows the pattern of terrestrial
- ecological zones. A large areal fraction of deserts obviously reduces the continental mean
- 313 SOC stock, whereas a large fraction of frozen organic soil increases the continental mean
- 314 SOC stock (Fig. 1).

4.2. Carbon in frozen high-latitude soils

- Large SOC deposits exist in the frozen soils of the permafrost region and are vulnerable to the
- effects of global warming. The mass of these deposits, however, is not well known because
- 318 <u>the area and the stocks of the permafrost region are uncertain.</u> The uncertainty of the area is
- 319 characterized by the variation in the delineation and thus extent of the permafrost region
- among different maps and databases, which is due also to different definitions of "permafrost"
- and associated concepts.

<u>One permafrost delineation is directly defined by the HWSD.</u> The HWSD lists for each soil

- unit the presence of permafrost within the top 200 cm (a so-called 'gelic phase'). SOC mass
- in the top 1 m of soils with a gelic phase is 164 Pg for a 13.1 Mm^2 soil area (Table 4). <u>A</u>
- second delineation is given by the 'Supplementary data to the HWSD' (Fischer et al., 2008),
- 326 <u>This database indicates on a 5' grid</u> the presence of continuous or discontinuous (i.e.,
- excluding sporadic and isolated) permafrost that is based on the analysis of the snow-adjusted

13

Martin Köchy 2015·2·26 16:24 **Gelöscht:** indicate on a 5' grid air frost number (Harrij van Velthuizen, IIASA, pers. Comm. 2011) as used for the Global
Agro-ecological Zones Assessment v3.0 (Fischer et al., 2008). This extent (19.5 Mm² cell
area, Fig. 2) encompasses the area of soils with a gelic phase and contains 185 Pg SOC on
16.7 Mm² soil area according to the HWSD. A third permafrost delineation (24.9 Mm² cell
area) is described by the Circum-Arctic Map of Permafrost and Ground Ice Conditions
(CAMP, Heginbottom et al., 1993), which comprises 12 categories of permafrost and ground
ice prevalence without a defined depth limit for the occurrence of permafrost. The CAMP
permafrost region (including permafrost in the Alps and Central Asian ranges) represents 21.7
Mm² soil area of the HWSD with 249 Pg SOC in the top 1 m.

338 4.3. Carbon in global wetlands

SOC stocks in wetlands are considerable because water reduces the availability of oxygen and thus greatly reduces decomposition rates (Freeman et al., 2001). Draining of wetlands often 340 greatly increases the decomposition of dead plant material, which results in the release of 341 carbon dioxide into the atmosphere. This process can significantly affect the global C budget 342 343 when it happens on a large scale. There is, however, no consensus of what constitutes a wetland at the global scale (Mitra et al., 2005). Therefore, the volume of wetland soil and its 344 C mass are also uncertain (Joosten, 2010). 345 The most detailed and recent maps of global scope with detailed wetland classification 346 (Köchy and Freibauer, 2009) are the Global Land Cover Characteristics database, v 2.0 347

(GLCC, Loveland et al., 2000) that comprises up to 6 wetland types ('Wooded Wet Swamp',

'Rice Paddy and Field', 'Inland Water', 'Mangrove', 'Mire, Bog, Fen', 'Marsh Wetland') and

the Global Lakes and Wetland Database (GLWD, Lehner and Döll, 2004) that comprises 12

wetland categories. Both maps have a resolution of 0.5'. The GLCC originates from analysis

of remote sensing data in the <u>JGBP</u>. Lehner and Döll compiled their database from existing

Martin Köchy 2015·2·27 10:12 Gelöscht: extent

Martin Köchy 2015·2·27 10:12 Gelöscht: extent

Martin Köchy 2015-2-20 13:43 **Gelöscht:** <#>Tarnocai et al. (2009) used the CAMP's permafrost classification (excluding the Alps and Central Asian ranges, 20.5 Mm² grid cell area) together with SOC and soil information from the Northern Circumpolar Soil Carbon Data Base (NCSCDB,

http://wms1.agr.gc.ca/NortherCircumpolar/n orthercircumpolar.zip) to estimate SOC mass in the permafrost region. The NCSCDB includes soil profile data not incorporated into the HWSD. Data for calculating SOC stocks (C concentration, BD, depth, coarse fragments) in the upper 3 m were derived from 1038 pedons from northern Canada, 131 pedons from Alaska, 253 pedons from Russia, 90 peat cores from western Siberia, 266 mineral and organic soils from the Usa Basin database, and an unspecified number of profiles from the WISE database (v.1.1)for Eurasian soils. Extrapolations were used to estimate SOC mass in mineral soils and Eurasian peat soils >1 m depth. The spatial extent of soil classes was obtained from existing digital and paper maps. Tarnocai et al.'s (2009) estimate of 496 Pg for the 0-1 m depth is much higher than that of HWSD's mass in the permafrost region (185 Pg). The difference is partly due the limit of 2 m that HWSD uses for distinguishing the 'gelic phase', whereas the Circum-arctic Map of Permafrost does not refer to a limit (Heginbottom et al., 1993). The more important contribution to the difference in mass than arising from definitions and extent is the greater SOC stock calculated from the NCSCDB (Table 4). In NCSCDB the mean SOC areal density of soil in all permafrost classes is $>20 \text{ kg/m}^2$, whereas the mean SOC areal density is 11.4 kg/m² in the HWSD across all classes. The difference suggests that the BD of frozen organic soil is higher than assumed by us. In addition to the SOC mass in the top 1 m, Tarnocai et al. (2009) estimated that the permafrost region contains 528 Pg in 1 m to 3 m depth, and 648 Pg in depths greater than 3 m. ... [3] artin Köchy 2015-2-19 17:08

Gelöscht: International Geosphere Biosphere program

- ⁴⁰⁵ maps, including the GLCC, and inventories. Some wetland types are restricted geographically
- 406 due to the heterogeneous classification across the source materials. The categories "50-100%
- 407 wetland" and "25–50% wetland", for example, occur only in North America, "wetland
- 408 complex" occurs only in Southeast Asia. One consequence is that the global extent of 'bogs,
- 409 fens, and mires' in the GLWD, 0.8 Mm², is smaller than the Canadian area of peatlands, 1.1
- 410 Mm² (Tarnocai et al., 2002), which is dominated by bogs and fens.
- 411 The spatial overlap of the GLWD and the GLCC categories is rather small (Table 6). Only the
- "Mire bog, fen" category of the GLCC has been adopted completely by the GLWD (Lehner
- 413 & Döll, 2004). Even categories with similar names like "Freshwater Marsh" vs. "Marsh
- 414 Wetland" and "Swamp Forest, Flooded Forest" vs. "Wooded Wet Swamps" show little spatial
- 415 overlap. Despite the GLWD's overall larger wetland area it does not include the areas
- 416 identified as "rice paddies" in the GLCC.

Based on the intersection of GLWD and HWSD (Fig. 3), the global SOC mass in the top 1 m 417 of soil of permanent and non-permanent wetlands (excluding lakes, reservoirs, and rivers) is 418 140 Pg (on 117 Mm² soil area). Using the GLCC Global Ecosystems classification, the area 419 covered by wetlands (excluding inland waters) is much smaller (3 vs 12 Mm²) and contains 420 only 34 Pg SOC (Table 7). The difference is partly due to the classification of large parts of 421 North America (including the prairie) as temporary or patchy wetland in the GLWD; but even 422 wetlands in a stricter sense cover twice the area and contain nearly twice the mass of SOC in 423 the GLWD compared to the GLCC. Therefore, we combined both maps for the assessment of SOC stocks and masses (Table 7). 425

The differences in SOC mass estimates between the GLWD and the GLCC indicate that
 wetland types are defined heterogeneously and that especially the classification of swamp
 forests, marshes, mangroves, and rice paddies needs to be harmonized. The contrasting land

Martin Köchy 2015·2·20 13:53

Gelöscht: Wetlands with the highest Corg and highest SOC stocks are bogs, fens, mires, and marshes and the "25-50%" and "50-100%" wetlands in boreal North America. The latter two categories represent mostly bogs, fens, and small lakes. Due to their high C_{org} these wetland types can also be classified as peatland. When wet peatlands are drained, they may no longer qualify as wetlands, but remain peatlands with high Corg and a large SOC mass. Drainage exposes the carbon to oxygen and thus accelerates peat decomposition and, depending on circumstances, an increase in BD. The global area of peatland with a minimum peat depth of 30 cm is 3.8 Mm² based on the International Mire Conservation Group Global Peatland Database (GPD, Joosten, 2010). Total SOC mass of peatlands in the GPD is 447 Pg for their total depth. This estimate is considered conservative because mangroves, salt marshes paddies, paludified forests, cloud forests, dambos, and Cryosols were omitted because of lack of data. The information available in the database for peatlands is very heterogeneous. For some countries only the total area of peatland is known. When depth information was missing or not plausible, a depth of 2 m was assumed in the GPD, although most peatlands are deeper (Joosten, 2010). It is not clear, which default values were used for C or BD in the assessment. C content (organic C fraction of ash-free mass) varies from 0.48-0.52 in Sphagnum peat to 0.52-0.59 in Scheuchzeria and woody peat (Chambers et al., 2010). Values of BD show much stronger variation. Ash-free bulk density ranged from <0.01 to 0.23 kg dm-3 in 4697 samples (Chambers et al., 2010) with a median of 0.1 kg dm-3. The variation is due to water content, soil depth, plant material, and degree of decomposition (Boelter, 1968). The highest density is found in well-decomposed, deep peat of herbaceous or woody origin at low water content. The great variation demands that BD of peatlands actually be measured at several depths and at ambient soil moisture at the same time as the C concentration. If this is not possible, PTFs of BD for peat ought to include water content, decomposition status and plant material. ... [4]

481	cover classification could be overcome by using the more generic land cover classes
482	developed within the UN Framework Convention on Climate Change (di Gregorio and
483	Jansen, 2005). Remote sensing methods are being developed to improve the mapping of
484	wetlands, e.g., the GlobWetland project (http://www.globwetland.org, and Journal of
485	Environmental Management 90, special issue (7)) or the Wetland Map of China (Niu et al.,
486	2009),
487	4.4. Carbon in tropical wetlands
488	Soils in the tropical land area (50 Mm ² within 23.5°N–23.5°S) contain 355 Pg SOC in the top
489	<u>1 m (Table 8).</u> The high intensity of rain in some parts of the tropics contributes to the
490	presence of wetlands (union of GLWD and GLCC classes as in the previous section) in 9% of
491	the tropical land area (50 Mm ² within 23.5°N–23.5°S) containing 40 Pg SOC (Table 8,
492	excluding lakes, reservoirs, rivers). Most of the wetland SOC (27 Pg) is found in marshes and
493	floodplains, and swamp or flooded forests. The GLCC category with the highest SOC mass
494	(10 Pg) is "Rice Paddy and Field" (1.2 Mm ² soil and cell area) but only 14% of this area is
495	recognized as wetland in the GLWD.
496	

Martin Köchy 2015·2·20 13:57

Gelöscht: In situ measurements of soil C_{org}, soil depth, and BD, however, must still be improved, collected, and made available for calculating global SOC mass.

lartin Köchy 2015·2·20 14:25

Gelöscht: the tropics

Martin Köchy 2015·2·27 10:19

Gelöscht: carbon Martin Köchy 2015·2·26 17:2

Gelöscht: Only 6% of the area of each of the two C-richest tropical wetland types are categorized as Histosols in the HWSD, totaling 0.1 Mm². The total area of Histosol in the HWSD, 0.4 Mm², agrees with the most recent and detailed, independent estimate of tropical peatland area (Page et al., 2011, defining peatland as soil having >65% organic matter in a minimum thickness of 30 cm). The total mass of SOC in grid cells of the spatial layer with at least some fraction of Histosol is 24.2 Pg.

Martin Köchy 2015·2·20 13:59

Gelöscht: Page et al. (2011) used peatland area, thickness, BD and Corg to calculate the SOC mass for each country within the tropics of Cancer and Capricorn. They tried to trace the original data and used best estimates where data were missing. Most data was available for area, but less data was available for thickness. Page et al. (2011) used 25% of maximum thickness when only this information was reported instead of mean thickness and used 0.5 m when no thickness was reported. The percentiles of the frequency distribution of their best estimate of thickness weighted by their best estimate of area per country is 0-10%: 0.5 m, 25%: 1.75 m, 50%-90%: 5.5 m, 97.5%: 7.0 m, mean: 4.0 m ± 2.2 m SD. This distribution can be used for estimates of SOC mass and associated uncertainty in other tropical peatlands. Data on BD and SOC concentration were rare. When they were provided they often referred only to the subsurface, although these parameters vary with depth. When these data were missing, Page et al. (2011) used 0.09 g cm-3 and 56% as best estimates based on literature reviews Consequently, their best estimate of SOC mass for tropical peatlands is 88.6 Pg for the whole soil depth, with a minimum of 81.7 and a maximum of 91.9 Pg. If one assumes an average peat thickness of 4 m and uniform vertical mass distribution, the top 1 m contains 22 Pg of SOC, close to our HWSD-based estimate for grid cells containing Histosol (24 Pg). Joosten (2010) estimated SOC mass for individual tropical countries based on the Global Peatland Database. For some countries the difference between Joosten's and Page et al.'s estimates are large. For example, Joosten's estimate for Sudan is 1.98 Pg, whereas Page et al. have 0.457 Pg. These differences may be caused by different definitions of "peat" and variability in depth estimates, SOC concentration, and BD [... [5]

16

604 <u>5. Discussion of HWSD-based SOC masses</u>

- 605 In this section we compare values of SOC masses derived from the adjusted HWSD to those
- 606 given by other important sources for SOC-rich soils in the permafrost region and in peatlands.
- 607 The values of the other sources are marked in the text by an asterisk for clarity (e.g. 496 Pg*)

608 <u>5.1. Carbon in frozen high-latitude soils</u>

- ⁶⁰⁹ The permafrost region can be delineated according to different criteria (see previous section).
- ⁶¹⁰ Tarnocai *et al.* (2009) used the CAMP's permafrost classification (20.5 Mm² grid cell area,
- excluding the Alps and Central Asian ranges) together with SOC and soil information from
- 612 the Northern Circumpolar Soil Carbon Data Base (NCSCDB,
- 613 http://wms1.agr.gc.ca/NortherCircumpolar/northercircumpolar.zip) to estimate SOC mass in
- 614 the permafrost region. The NCSCDB includes soil profile data not incorporated into the
- 615 HWSD. Data for calculating SOC stocks (C concentration, BD, depth, coarse fragments) in
- 616 the upper 3 m were derived from 1038 pedons from northern Canada, 131 pedons from
- Alaska, 253 pedons from Russia, 90 peat cores from western Siberia, 266 mineral and organic
- soils from the Usa Basin database, and an unspecified number of profiles from the WISE
- database (v.1.1) for Eurasian soils. Extrapolations were used to estimate SOC mass in mineral
- 620 soils and Eurasian peat soils >1 m depth. The spatial extent of soil classes was obtained from
- existing digital and paper maps. Tarnocai et al.'s (2009) estimate of 496 Pg* for the 0-1 m
- depth is much higher than that of HWSD's mass in the CAMP's permafrost region (233 Pg).
- 623 The difference is partly due the limit of 2 m that HWSD uses for distinguishing the 'gelic
- 624 phase', whereas the CAMP does not refer to a depth limit (Heginbottom et al., 1993). The
- 625 difference in mass is not only due to contrasting definitions and extent but even more so due
- 626 to the greater SOC stock calculated from the NCSCDB (Table 4). In the NCSCDB the mean
- $\frac{\text{SOC areal density of soil in all permafrost classes is >20 kg/m^2, whereas the mean SOC areal}{\text{SOC areal density of soil in all permafrost classes is >20 kg/m^2, whereas the mean SOC areal}}$
 - 17

Martin Köchy 2015-2-20 13:41 Gelöscht: Conclusions

Martin Köchy 2015·2·26 17:41 Formatiert: Textkörper

Martin Köchy 2015·2·26 17:41 Formatiert: Schriftart:Times New Roman

density is 11.4 kg/m ² in the HWSD across all classes. The difference suggests that the BD of
frozen organic soil is higher than assumed by us.
Inaccuracies associated with the mass estimates arise from incomplete knowledge of the
spatial distribution of soil classes, soil depths, sparse distribution of soil profile data and lack
of soil profiles with a full complement of measured data. Tarnocai et al. discuss extensively
the uncertainty of their estimates. In terms of categories of confidence of the
Intergovernmental Panel's on Climate Change Fourth Assessment Report (IPCC AR4),
Tarnocai et al. have medium to high confidence (>66%) in the values for the North-American
stocks of the top 1 m, medium confidence (33-66%) in the values for the Eurasian stocks of
the top 1 m, and very low to low confidence (<33%) in the values for the other regional stocks
and stocks of layers deeper than 1 m. Here we note only that major uncertainty is linked to the
area covered by high latitude peatlands (published estimates vary between 1.2 and 2.7 Mm ²)
which alone results in a range of 94-215 Pg SOC. In addition to the SOC mass in the top 1 m,
Tarnocai et al. (2009) estimated that the permafrost region contains 528 Pg* in 1 m to 3 m
depth, and 648 Pg* in depths greater than 3 m. The C mass contained in >3 m depth of river
deltas is potentially great (241 Pg*, Tarnocai et al., 2009), but is based solely on extrapolation
on the SOC stock and area of the Mackenzie River delta. Yedoma (Pleistocene loess deposits
with high C_{org}) SOC mass (407 Pg*, >3 m depth) is also associated with great uncertainty.
The estimate (adopted from Zimov et al., 2006) is based on a sketched area of 1 Mm ² in
Siberia (thus excluding smaller Yedoma deposits in North America) and mean literature
values for depth (25 m) whose ranges extend > \pm 50% of the mean.

650 <u>5.2. Carbon in peatlands</u>

- 651 Wetlands with the highest C_{ore} and highest SOC stocks are bogs, fens, mires, and marshes and
- 652 the "25–50%" and "50–100%" wetlands in boreal North America. The latter two categories

18

Martin Köchy 2015·2·20 13:56 Formatiert: Überschrift 2 represent mostly bogs, fens, and small lakes. Due to their high C_{ore} these wetland types can
 also be classified as peatland.

The global area of peatland with a minimum peat depth of 30 cm is 3.8 Mm² based on the 655 656 International Mire Conservation Group Global Peatland Database (GPD, Joosten, 2010). 657 Total SOC mass of peatlands in the GPD is 447 Pg* for their total depth. This estimate is considered conservative because mangroves, salt marshes, paddies, paludified forests, cloud forests, dambos, and Cryosols were omitted because of lack of data. The information in the 659 660 GPD is very heterogeneous. Missing data for calculating SOC mass had to be be estimated. For some countries only the total area of peatland was known. When depth information was 661 missing or not plausible, a depth of 2 m was assumed in the GPD, although most peatlands 662 are deeper (Joosten, 2010). It is not clear, which default values were used for Core or BD in the 663 assessment. C content (organic C fraction of ash-free mass) varies from 0.48-0.52 in 664 665 Sphagnum peat to 0.52–0.59 in Scheuchzeria and woody peat (Chambers et al., 2010). Values of BD show much stronger variation. Ash-free bulk density ranged from <0.01 to 0.23 kg dm 666 ³ in 4697 samples (Chambers et al., 2010) with a median of 0.1 kg dm⁻³. The variation is due 667 668 to water content, soil depth, plant material, and degree of decomposition (Boelter, 1968). The highest density is found in well-decomposed, deep peat of herbaceous or woody origin at low 669 670 water content. When wet peatlands are drained, they may no longer qualify as wetlands, but remain peatlands with high Corg and a large SOC mass. Drainage exposes the carbon to 671 672 oxygen and thus accelerates peat decomposition and, depending on circumstances, an increase in BD. The great variation demands that BD of peatlands actually must be measured at several 673 674 depths and at ambient soil moisture at the same time as the C concentration. If this is not possible, PTFs of BD for peat ought to include water content, decomposition status, and plant 675 material.

677	Peatlands with a certain thickness of organic layer qualify as Histosols. HWSD adopted the
678	FAO definition "Soils having an H horizon of 40 cm or more of organic soil materials (60 cm
679	or more if the organic material consists mainly of sphagnum or moss or has a bulk density of
680	less than 0.1) either extending down from the surface or taken cumulatively within the upper
681	80 cm of the soil; the thickness of the H horizon may be less when it rests on rocks or on
682	fragmental material of which the interstices are filled with organic matter." (FAO, 1997). The
683	area covered by Histosols in the HWSD (Fig. 4) is 3.3 Mm ² (cell area multiplied by fraction
684	of Histosol), slightly lower than the area given by the GPD, and contains 113 Pg SOC. The
685	total area of cells with at least some fraction of Histosol, however, is 10 Mm ² containing 188
686	Pg SOC. The area of Histosol outside wetlands (1.7 Mm ²) might indicate that a large portion
687	of originally wet peatland has been drained and is exposed to decomposition.
688	5.3. Carbon in tropical peatlands
689	Six percent of the area of each of the two C-richest tropical wetland types are categorized as
690	Histosols in the HWSD, totaling only 0.1 Mm ² . Including non-wetlands, the total area of
691	Histosols in the HWSD, 0.4 Mm ² , agrees well with the most recent and detailed, independent
692	estimate of tropical peatland area (Page et al., 2011, defining peatland as soil having >65%
693	organic matter in a minimum thickness of 30 cm). The total mass of SOC in grid cells of the
694	spatial layer with at least some fraction of Histosol is 24.2 Pg.
605	Page at al. (2011) used peatland area, thickness, BD and C \perp to calculate the SOC mass for
606	each country within the transics of Cancer and Canricorn. They tried to trace the original data
090	and used best estimates where data were missing. Most data was evolable for area but less
697	and used best estimates where data were missing. Most data was available for area, but less
698	data was available for thickness. Page et al. (2011) used 25% of maximum thickness when
699	only this information was reported instead of mean thickness and used 0.5 m when no
699 700	only this information was reported instead of mean thickness and used 0.5 m when no thickness was reported. The percentiles of the frequency distribution of their best estimate of

701	thickness weighted by their best estimate of area per country is 0-10%: 0.5 m, 25%: 1.75 m,
702	50%-90%: 5.5 m, 97.5%: 7.0 m, mean: 4.0 m ± 2.2 m SD. This distribution can be used for
703	estimates of SOC mass and associated uncertainty in other tropical peatlands. Data on BD and
704	SOC concentration were rare. When they were provided they often referred only to the
705	subsurface, although these parameters vary with depth. When these data were missing, Page
706	et al. (2011) used 0.09 g cm ⁻³ and 56% as best estimates based on literature reviews. The best
707	estimate of SOC mass for tropical peatlands of Page et al. (2011) is 88.6 Pg* for the whole
708	soil depth, with a minimum of 81.7 and a maximum of 91.9 Pg*. If one assumes an average
709	peat thickness of 4 m and uniform vertical mass distribution, the top 1 m contains 22 Pg* of
710	SOC, close to our HWSD-based estimate for grid cells containing Histosol (24 Pg). Thus,
711	peatlands may contain about 6% of the tropical SOC mass within the first meter and
712	approximately 21% of the total tropical SOC mass (without depth limit). Obviously, the
713	uncertainty of these estimates is great.
	Lesster (2010) estimated SOC more for individual territori constring based on the Clabel
/14	Joosten (2010) estimated SOC mass for individual tropical countries based on the Global
715	Peatland Database. For some countries the difference between Joosten's and Page et al.'s
716	estimates are large. For example, Joosten's estimate for Sudan is 1.98 Pg*, whereas Page et
717	al. have 0.457 Pg*. These differences may be caused by different definitions of "peat" and

718 variability in depth estimates, SOC concentration, and BD in the data sources.

719 **6.** Conclusions

720 6.1. Global carbon mass – reprise

721	The estimate of the global SOC mass within the top 1 m based on the HWSD (1062 Pg) can
722	be improved if and where other sources provide better estimates. The HWSD estimate of SOC
723	mass for tropical peatlands agreed well with other sources. The SOC mass in the permafrost
724	region estimated by Tarnocai et al. (2009) appears to be more accurate than that of HWSD.
725	Therefore, we substitute for the permafrost region the HWSD-based estimate (- 233 Pg
726	[Table 4]) by Tarnocai et al.'s estimate (+ 496 Pg). This calculation (1062 –233 + 496 Pg)
727	updates the global SOC mass within the top 1 m to 1325 Pg.
728	For including deeper soils in an estimate of the global SOC mass, we consider first estimates
729	of deeper soil layers for the permafrost region and tropical peatlands. The best estimate of the
730	SOC mass below 1 m for the permafrost region known to us is 1176 Pg (calculated from
731	Tarnocai et al., 2009). In order to estimate the mass for 1-4 m depth of tropical peatlands, we
732	use 3/4 of Page et al.'s best estimate for the top 4 m (66.5 Pg). Additional 389 Pg SOC is
733	contained below 1 m outside the permafrost region and the tropics (Jobbágy and Jackson,
734	2000). In total, the mass of SOC in the soil is about 3000 Pg, but large uncertainties remain,
735	especially for depths >1 m.
736	Another source of uncertainty is the estimation of BD. The BD of peat varies between 0.05
737	and 0.26 kg/dm ^{3} (Boelter, 1968). If the same range holds for Histosols (3.3 Mm ^{2} Histosol
738	area, 1 m depth, 34% C_{org}), this variation alone introduces an uncertainty range of -56 to +180

- Pg for the estimate of global SOC in the top meter, which is larger than the estimated annual
- ⁷⁴⁰ global soil respiration (79.3–81.8 Pg C, Raich et al., 2002). The areal extent of <u>organic soils</u>,

Gelöscht: Assuming that the assessment of Tarnocai et al. (2009) of t Martin Köchy 2015-2-26 18:16 Gelöscht: is Martin Köchy 2015·2·20 14:38 Gelöscht:, Martin Köchy 2015·2·20 14:38 Gelöscht: update the global SOC mass within the top 1 m to 1325 Pg Martin Köchy 2015·2·20 14:39 Gelöscht: 1062 [HWSD global SOC mass] Martin Köchy 2015·2·20 14:39 Gelöscht: HWSD permafrost region SOC mass, Martin Köchy 2015·2·20 14:40 Gelöscht: + 496 [Martin Köchy 2015·2·20 14:40 Gelöscht:] Martin Köchy 2015-2-26 18:23 Gelöscht: 1672 Unknown Feldfunktion geändert Martin Köchy 2015·2· Gelöscht: We can use the best estimates of the total SOC mass for the permafrost region (1672 Pg - including deep carbon and high carbon content deposits -, Tarnocai et al.,2009) and the tropics (421 Pg) and add it to the SOC mass outside these areas (473 Pg). This sum (2567 Pg) does not yet comprise Feldfunktion geändert Martin Köchy 2015·2·20 14:55 Gelöscht: 389 Pg, Martin Köchy 2015·2·20 14:56 Gelöscht: Thus the Martin Köchy 2015·2·20 14:56 Gelöscht: in Martin Köchy 2015-2-20 14:56 Gelöscht: soil Martin Köchy 2015·2·26 18:28 Gelöscht: estimated at Unknown Feldfunktion geändert Martin Köchy 2015·2·26 18:01 Gelöscht: SO Martin Köchy 2015·2

Gelöscht: peatlands

Martin Köchy 2015·2·20 14:35

their depth, and the BD at different depths should therefore receive the greatest focus of future

⁷⁶⁸ soil mapping activities.

⁷⁶⁹ Soil monitoring is crucial for detecting changes in SOC stocks and as a reference for

projecting changes in the global carbon pool using models (Wei et al., 2014, Wieder et al.,

- 2014, Yan et al., 2014). The following conclusions from our study and a workshop of soil
- 772 experts (Köchy and Freibauer, 2011) with respect to improved soil monitoring agree with
- more comprehensive recommendations by an international group of experts (Jandl et al.,
- 2014). In situ measurements of soil C_{org}, soil depth, and BD must still be improved, collected,
- ⁷⁷⁵ and made available for calculating global SOC mass. Extra care is necessary to reduce
- variability of data because variability reduces the potential of detecting change. Classification
- of soils as it is currently used in mapping produces uncertainty in the reported C stock when
- the characteristics of soil classes are aggregated and then used in further calculations. The use
- 779 of pedotransfer rules and functions further increases the uncertainty of the real values. Since
- 780 **PTFs** are entirely empirical in nature, it is preferable that they be derived from soils that are
- ⁷⁸¹ similar in nature to the soils to which the functions will be applied. For purposes of detecting
- actual change in C stocks their uncertainty should be quantified. Of course it would be best if
- ⁷⁸³ C_{org}, BD, and coarse fragments were measured at the same point or sample to reduce effects of
- ⁷⁸⁴ spatial variability. Predictive mapping techniques, including geo-statistics, modelling, and
- other quantitative methods (McBratney et al., 2003a, Grunwald et al., 2011), especially in
- conjunction with proximal (radiometry, NIR spectroscopy) or hyperspectral remote-sensing of
- soil properties (Gomez et al., 2008, Stockmann et al., 2015) can potentially reduce
- ⁷⁸⁸ uncertainties in SOC mapping introduced by soil classification and help in interpreting spatio-
- temporal patterns. Whether soils are mapped in the classical way or by predictive methods,
- mapping of soils should be coordinated with the direct or indirect mapping of SOC input and

Martin Köchy 2015·2·20 15:00 **Gelöscht:** pedotransfer functions

- its controlling factors (land use, land cover, crop type, land use history and land management)
- and extent and soil depth of wetlands, peatlands, and permafrost.

Uncertainty of SOC stocks in current maps could further be reduced if all soil types and 794 regions were well represented by soil profile data with rich soil characteristics. Many soil 795 796 profile data collected by governments and publicly funded projects remain unused because they are not available digitally; their use is restricted because of data protection issues, or 797 because they are only known to a very limited number of soil scientists. Existing approaches 798 such as the NCSCB, the GlobalSoilMap.net project, and the Global Soil Partnership (coordinated by FAO), are important steps to improve the situation. These activities would 800 benefit further if all publicly funded, existing soil profile data were made publicly available to 801 the greatest possible extent. 802 Another source of uncertainty is introduced because profile data and soil maps have been 803 generated by a multitude of methods. Furthermore, if different methods are preferably used 804 for particular soil types or regions, small differences multiplied by large areas can result in 805 significant differences at the global level. Therefore, international activities to harmonize 806 807 methods of sampling, calculation, and scaling should be supported. The harmonized methods should then actually be applied in soil sampling. Preferably, samples should be archived so 808 that soils can be reanalyzed with improved or new methods or for checking data by more than 809

810

811 6.2. Implications

one laboratory.

- 812 The strong effect of BD values on the calculation of SOC stocks and regional or global
- masses should guide the focus of global observation networks to improve not only the
- observation of SOC concentrations but also on BD. Furthermore, our study describes for the
- first time the frequency distribution of SOC stocks within broad classes of land-use/land-

Martin Köchy 2015-2-26 18:30 Gelöscht: Northern Circumpolar Soil Carbon Base

818	cover and C-rich environments based on one of the most exhaustive, harmonized, spatially
819	explicit global databases available to date. The frequency distribution allows a more focused
820	spatial extrapolation and assessment of accuracy in studies where SOC is used as an
821	independent variable (e.g., Pregitzer and Euskirchen, 2004). The frequency distributions also
822	provide a foundation for targeting SOC conservation measures (Powlson et al., 2011) and for
823	improving carbon accounting methods with associated uncertainties as used in the UNFCCC
824	(García-Oliva and Masera, 2004).
825	CO_2 emissions from soils are used in calculations of the global carbon cycle. Direct
826	observations of CO_2 emissions from soils (e.g., by eddy-flux towers), however, cannot be
827	implemented in a spatially contiguous way. Indirect measurements by remote sensing can
828	improve the spatial coverage but require ground observations for conversion from observed
829	radiation to loss of CO_2 from soils and distinction from other CO_2 sources (Ciais et al., 2010).
830	At the global scale, in-situ measurements must be complemented by modelling activities,
831	which are greatly improved if variation in key factors like SOC can be accounted for. Thus,
832	more detailed information on the global distribution of SOC, horizontally and vertically,
833	including accounts of its accuracy and its variability, are necessary to improve estimates of

the global carbon flow.

835 Author contributions

- 836 MK designed and carried out the analyses and wrote the manuscript, RH contributed a
- thorough analysis of inconsistencies in the HWSD and alternative estimates, AF suggested the
- topic and provided valuable insights on the presentation of the data.
- 839 Acknowledgments. We thank Charles Tarnocai for comments on the manuscript. We
- appreciate the constructive comments of the reviewers. MK was funded by EU FP7 project
- 841 COCOS (grant 212196) and FACCE MACSUR (BMBF grant 031A103A).

842 References

- Amundson, R.: The carbon budget in soils, Annual Review of Earth and Planetary Sciences, 29, 535-843 562, doi:10.1146/annurev.earth.29.1.535, 2001. 844
- Batjes, N. H.: Total carbon and nitrogen in the soils of the world, European Journal of Soil Science, 845

47, 151-163, doi:10.1111/j.1365-2389.1996.tb01386.x, 1996.

Batjes, N. H.: Harmonized soil profile data for applications at global and continental scales: updates to 847

the WISE database, Soil Use and Management, 25, 124-127, doi:10.1111/j.1475-848

849 2743.2009.00202.x, 2009.

- Boelter, D. H.: Important physical properties of peat materials, Proceedings of the Third International 850 Peat Congress, Ottawa, Ontario, Canada, 18-23 August 1968, Canada. Dept. of Energy, Mines 851 852 and Resources and National Research Council of Canada, 1968.
- Carré, F., R. Hiederer, V. Blujdea, and R. Koeble: Background guide for the calculation of land 853
- 854 carbon stocks in the biofuels sustainability scheme drawing on the 2006 IPCC Guidelines for
- National Greenhouse Gas Inventories, JRC Scientific and Technical Reports, EUR 24573 EN, 855
- Office for Official Publications of the European Communities, Luxembourg, Office for Official 856
- Publications of the European Communities, available at: 857

http://eusoils.jrc.ec.europa.eu/ESDB_Archive/eusoils_docs/other/EUR24573.pdf, 2010. 858

- 859 Chambers, F. M., Beilman, D. W., and Yu, Z.: Methods for determining peat humification and for
- quantifying peat bulk density, organic matter and carbon content for palaeostudies of climate 860 and peatland carbon dynamics, Mires and Peat, 7, 7.1-7.10, 2010/2011. 861
- Ciais, P., A. J. Dolman, R. Dargaville, L. Barrie, A. Bombelli, J. Butler, P. Canadell, and T.
- 863 Moriyama: GEO Carbon Strategy, GEO Secretariat and FAO, Geneva and Rome, GEO
- Secretariat and FAO, available at: 864
- 865 http://www.earthobservations.org/documents/sbas/cl/201006_geo_carbon_strategy_report.pdf,
- 2010. 866

862

- Ciais, P., C. Sabine, G. Bala, L. Bopp, V. Brovkin, J. Canadell, A. Chhabra, R. DeFries, J. Galloway, 867
- M. Heimann, C. Jones, C. Le Quere, R. B. Myneni, S. Piao, and P. Thornton: Carbon and other 868

869	biogeochemical cycles, in: Climate Change 2013: The Physical Science Basis. Contribution of	
870	Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate	
871	Change, Stocker, T. F., D. Qin, GK. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels,	
872	Y. Xia, V. Bex, and P. M. Midgley (Eds), Cambridge University Press, Cambridge, UK, and	
873	New York, NY, U.S.A., 465–570, doi: 0.1017/CBO9781107415324.015., 2013.	
874	Digital Soil Map of the World, Rome, Italy, available at:	
875	http://www.fao.org/geonetwork/srv/en/metadata.show?id=14116, 2007.	
876	Ellert, B. H., H. H. Janzen, and B. G. McConkey: Measuring and comparing soil carbon storage, in:	
877	Assessment methods for soil carbon, Advances in Soil Science, Lal, R., J. M. Follett, and B. A.	
878	Stewart (Eds), Lewis, Boca Raton, Florida, 131-146, 2001.	
879	ESRI (Environmental Systems Research Institute): World Continents, ESRI Data & Maps 2002, CD 1,	
880	Environmental Systems Research Institute, Inc. (ESRI), Redlands, California, U.S.A., 2002.	
881	FAO: FAO/Unesco Soil Map of the World, Revised Legend, with corrections and updates. Originally	
882	published in 1988 as World Soil Resources Report 60, FAO, Rome. Reprinted with updates,	
882 883	published in 1988 as World Soil Resources Report 60, FAO, Rome. Reprinted with updates, Technical Paper, 20, ISRIC, Wageningen, ISRIC, available at:	
882 883 884	published in 1988 as World Soil Resources Report 60, FAO, Rome. Reprinted with updates, Technical Paper, 20, ISRIC, Wageningen, ISRIC, available at: http://library.wur.nl/isric/fulltext/isricu_i9264_001.pdf, 1997.	
882 883 884 885	 published in 1988 as World Soil Resources Report 60, FAO, Rome. Reprinted with updates, Technical Paper, 20, ISRIC, Wageningen, ISRIC, available at: http://library.wur.nl/isric/fulltext/isricu_i9264_001.pdf, 1997. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.1), FAO and 	
882 883 884 885 886	 published in 1988 as World Soil Resources Report 60, FAO, Rome. Reprinted with updates, Technical Paper, 20, ISRIC, Wageningen, ISRIC, available at: http://library.wur.nl/isric/fulltext/isricu_i9264_001.pdf, 1997. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.1), FAO and IIASA, Rome, Italy, and Laxenburg, Austria, 2009. 	N
882 883 884 885 886 886	 published in 1988 as World Soil Resources Report 60, FAO, Rome. Reprinted with updates, Technical Paper, 20, ISRIC, Wageningen, ISRIC, available at: http://library.wur.nl/isric/fulltext/isricu_i9264_001.pdf, 1997. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.1), FAO and IIASA, Rome, Italy, and Laxenburg, Austria, 2009. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.2), FAO and 	N
882 883 884 885 886 887 888	 published in 1988 as World Soil Resources Report 60, FAO, Rome. Reprinted with updates, Technical Paper, 20, ISRIC, Wageningen, ISRIC, available at: http://library.wur.nl/isric/fulltext/isricu_i9264_001.pdf, 1997. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.1), FAO and IIASA, Rome, Italy, and Laxenburg, Austria, 2009. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.2), FAO and IIASA, Rome, Italy, and Laxenburg, Austria, 2012. 	Λ
882 883 884 885 886 887 888 888	 published in 1988 as World Soil Resources Report 60, FAO, Rome. Reprinted with updates, Technical Paper, 20, ISRIC, Wageningen, ISRIC, available at: http://library.wur.nl/isric/fulltext/isricu_i9264_001.pdf, 1997. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.1), FAO and IIASA, Rome, Italy, and Laxenburg, Austria, 2009. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.2), FAO and IIASA, Rome, Italy, and Laxenburg, Austria, 2012. Fischer, G., F. Nachtergaele, S. Prieler, H. T. van Velthuizen, L. Verelst, and D. Wiberg: Global agro- 	N
882 883 884 885 886 887 888 889 889	 published in 1988 as World Soil Resources Report 60, FAO, Rome. Reprinted with updates, Technical Paper, 20, ISRIC, Wageningen, ISRIC, available at: http://library.wur.nl/isric/fulltext/isricu_i9264_001.pdf, 1997. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.1), FAO and IIASA, Rome, Italy, and Laxenburg, Austria, 2009. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.2), FAO and IIASA, Rome, Italy, and Laxenburg, Austria, 2012. Fischer, G., F. Nachtergaele, S. Prieler, H. T. van Velthuizen, L. Verelst, and D. Wiberg: Global agro- ecological zones assessment for agriculture (GAEZ 2008), IIASA, Laxenburg, Austria and 	ſ
882 883 884 885 886 887 888 889 890 891	 published in 1988 as World Soil Resources Report 60, FAO, Rome. Reprinted with updates, Technical Paper, 20, ISRIC, Wageningen, ISRIC, available at: http://library.wur.nl/isric/fulltext/isricu_i9264_001.pdf, 1997. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.1), FAO and IIASA, Rome, Italy, and Laxenburg, Austria, 2009. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.2), FAO and IIASA, Rome, Italy, and Laxenburg, Austria, 2012. Fischer, G., F. Nachtergaele, S. Prieler, H. T. van Velthuizen, L. Verelst, and D. Wiberg: Global agro- ecological zones assessment for agriculture (GAEZ 2008), IIASA, Laxenburg, Austria and FAO, Rome, Italy., IIASA, Laxenburg, Austria and FAO, Rome, Italy., available at: 	Γ
882 883 884 885 886 887 888 889 890 891 892	 published in 1988 as World Soil Resources Report 60, FAO, Rome. Reprinted with updates, Technical Paper, 20, ISRIC, Wageningen, ISRIC, available at: http://library.wur.nl/isric/fulltext/isricu_i9264_001.pdf, 1997. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.1), FAO and IIASA, Rome, Italy, and Laxenburg, Austria, 2009. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.2), FAO and IIASA, Rome, Italy, and Laxenburg, Austria, 2019. Fischer, G., F. Nachtergaele, S. Prieler, H. T. van Velthuizen, L. Verelst, and D. Wiberg: Global agro- ecological zones assessment for agriculture (GAEZ 2008), IIASA, Laxenburg, Austria and FAO, Rome, Italy., IIASA, Laxenburg, Austria and FAO, Rome, Italy., available at: http://www.iiasa.ac.at/Research/LUC/External-World-soil- 	
882 883 884 885 886 887 888 889 890 890 891 892 893	 published in 1988 as World Soil Resources Report 60, FAO, Rome. Reprinted with updates, Technical Paper, 20, ISRIC, Wageningen, ISRIC, available at: http://library.wur.nl/isric/fulltext/isricu_i9264_001.pdf, 1997. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.1), FAO and IIASA, Rome, Italy, and Laxenburg, Austria, 2009. FAO, IIASA, ISRIC, ISSCAS, and JRC: Harmonized World Soil Database (version 1.2), FAO and IIASA, Rome, Italy, and Laxenburg, Austria, 2012. Fischer, G., F. Nachtergaele, S. Prieler, H. T. van Velthuizen, L. Verelst, and D. Wiberg: Global agro- ecological zones assessment for agriculture (GAEZ 2008), IIASA, Laxenburg, Austria and FAO, Rome, Italy., IIASA, Laxenburg, Austria and FAO, Rome, Italy., available at: http://www.iiasa.ac.at/Research/LUC/External-World-soil- database/HTML/SoilQualityData.html?sb=11, 2008. 	

895 doi:10.1038/35051650, 2001. Martin Köchy 2015·2·20 08:59

Gelöscht: Digital Soil Map of the World, Rome, Italy, available at: http://www.fao.org/geonetwork/srv/en/metad ata.show?id=14116, 2007. .

- 900 García-Oliva, F., and Masera, O. R.: Assessment and Measurement Issues Related to Soil Carbon
- 901 Sequestration in Land-Use, Land-Use Change, and Forestry (LULUCF) Projects under the
- 902 Kyoto Protocol, Clim. Change, 65, 347-364, doi:10.1023/B:CLIM.0000038211.84327.d9, 2004.
- 903 Global Soil Data Task Group: Global Soil Data Products CD-ROM (IGBP-DIS), Available from Oak
- Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A.
 [http://www.daac.ornl.gov], 2000.
- Gomez, C., Viscarra, R., Raphael A., and McBratney, A. B.: Soil organic carbon prediction by
- 907 hyperspectral remote sensing and field vis-NIR spectroscopy: An Australian case study,
- 908 Geoderma, 146, 403-411, doi:10.1016/j.geoderma.2008.06.011, 2008.
- Gorham, E.: Northern peatlands: role in the carbon cycle and probable responses to climatic warming,
 Ecol. Appl., 1, 182-195, doi:10.2307/1941811, 1991.
- 911 GRASS Development Team: Geographic Resources Analysis Support System (GRASS) Software,
- 912 Version 6.4.2, Open Source Geospatial Foundation, http://grass.osgeo.org, 2011.
- 913 di Gregorio, A., and L. J. M. Jansen: Land cover classification system (LCCS). Classification concepts
- and user manual. Software version (2), Food and Agriculture Organization, Rome, Italy, Food
- 915 and Agriculture Organization, available at:
- 916 http://www.fao.org/docrep/008/y7220e/y7220e00.htm#Contents, 2005.
- 917 Grunwald, S., Thompson, J. A., and Boettinger, J. L.: Digital soil mapping and modeling at
- continental scales: finding solutions for global issues, Soil Sci. Soc. Am. J., 75, 1201-1213,
 doi:10.2136/sssaj2011.0025, 2011.
- 920 Heginbottom, J. A., J. F. Brown, E. S. Melnikov, and O. J. Ferrians: Circum-arctic map of permafrost
- 921 and ground ice conditions, Proceedings of the Sixth International Conference on Permafrost,
- 922 Wushan, Guangzhou, China, 2, 1132-1136, South China University Press, 1993.
- 923 Henry, M., Valentini, R., and Bernoux, M.: Soil carbon stocks in ecoregions of Africa, Biogeosciences
- 924 Discussions, 6, 797-823, doi:10.5194/bgd-6-797-2009, 2009.
- 925 Hiederer, R.: Data update and model revision for soil profile analytical database of Europe of
- 926 measured parameters (SPADE/M2), JRC Scientific and Technical Reports, EUR 24333 EN,



- 927 Office for Official Publications of the European Communities, Luxembourg, Office for Official
- 928 Publications of the European Communities, doi:10.2788/85262, 2010.
- 929 Hiederer, R., and M. Köchy: Global soil organic carbon estimates and the Harmonized World Soil
- Database, JRC Scientific and Technical Reports, 68528/EUR 25225 EN, Joint Research Centre,
 Ispra, Italy, Joint Research Centre, doi:10.2788/13267, 2011.
- 932 Hiederer, R., F. Ramos, C. Capitani, R. Koeble, V. Blujdea, O. Gomez, D. Mulligan, and L. Marelli:
- 933 Biofuels: a new methodology to estimate GHG emissions from global land use change, JRC
- 934 Scientific and Technical Reports, EUR 24483 EN, Office for Official Publications of the
- 935 European Communities, Luxembourg, Office for Official Publications of the European
- 936 Communities, doi:10.2788/48910, 2010.
- 937 IUSS Working Group WRB: World reference base for soil resources 2006. A framework for
- 938 international classification, correlation and communication, World Soil Resources Reports, 103,
- Food and Agricultre Organization of the United Nations, Rome, 2006.
- Jandl, R., Rodeghiero, M., Martinez, C., Cotrufo, M. F., Bampa, F., van Wesemael, B., Harrison, R.,
- 941 B., Guerrini, I. A., Richter, D. d., Jr., Rustad, L., Lorenz, K., Chabbi, A., and Miglietta, F.:
- 942 Current status, uncertainty and future needs in soil organic carbon monitoring, Science of the
- 943 Total Environment, 468, 376-383, doi:10.1016/j.scitotenv.2013.08.026, 2014.
- Jobbágy, E. G., and Jackson, R. B.: The vertical distribution of soil organic carbon and its relation to
- 945 climate and vegetation, Ecol. Appl., 10, 423-436, doi:10.1890/1051-
- 946 0761(2000)010[0423:TVDOSO]2.0.CO;2, 2000.
- 947 Joosten, H.: The global peatland CO₂ picture. Peatland status and emissions in all countries of the
- world, Wetlands International, Ede, 2010.
- 949 Köchy, M., and A. Freibauer: Global spatial distribution of wetlands, COCOS Report, D4.3a, Johann
- 950 Heinrich von Thünen-Institut, Braunschweig, Germany, available at: http://www.cocos-
- 951 carbon.org/docs/D4.3a_wetlands_report.pdf, 2009.
- 952 Köchy, M., and A. Freibauer: Workshop on mapping of soil carbon stocks at the global scale. COCOS
- 953 Report D1.4.3. Johann Heinrich von Thünen Institut, Braunschweig, Germany, available at
- 954 http://literatur.ti.bund.de/digbib_extern/dn049595.pdf, 2011 (last accessed 13 July 2011).

Martin Köchy 2015·2·27 10:59 **Gelöscht:** , Johann Heinrich von Thünen-Institut

- 957 Lehner, B., and Döll, P.: Development and validation of a global database of lakes, reservoirs and
- 958 wetlands, J. Hydrol., 296, 1-22, doi:10.1016/j.jhydrol.2004.03.028, 2004.
- 959 Loveland, T. R., Reed, B. C., Brown, J. F., Ohlen, D. O., Zhu, J., Yang, L., and Merchant, J. W.:
- 960 Development of a Global Land Cover Characteristics Database and IGBP DISCover from 1-km
- 961 AVHRR Data, International Journal of Remote Sensing, 21, 1303-1330,
- 962 doi:10.1080/014311600210191,2000.
- 963 Mäkilä, M.: Calculation of the energy content of mires on the basis of peat properties [In Finnish with
- 964 English summary], Report of Investigation, 121, Geological Survey of Finland, Geological
 965 Survey of Finland, 1994.
- McBratney, A. B., Mendonça, S., M.L, and Minasny, B.: On digital soil mapping, Geoderma, 117, 3 52, doi:10.1016/S0016-7061(03)00223-4, 2003/11//a.
- 968 McBratney, A. B., Mendonça Santos, M. L., and Minasny, B.: On digital soil mapping, Geoderma,
- 969 117, 3-52, doi:10.1016/S0016-7061(03)00223-4, 2003b.
- 970 Mitra, S., Wassmann, R., and Vlek, P.: An appraisal of global wetland area and its organic carbon
- 971 stock, Current Science, 88, 25-35, 2005.
- 972 Myhre, G., D. Shindell, F.-M. Breon, W. Collins, J. Fuglestvedt, J. Huang, D. Koch, J.-F. Lamarque,
- 973 D. Lee, B. Mendoza, T. Nakajima, A. Robock, G. Stephens, T. Takemura, and H. Zhang:
- 974 Anthropogenic and Natural Radiative Forcing, in: Climate Change 2013: The Physical Science
- 975 Basis. Contribution of Working Group I to the Fifth Assessment Report of the
- 976 Intergovernmental Panel on Climate Change, Stocker, T. F., D. Qin, G.-K. Plattner, M. Tignor,
- 977 S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P. M. Midgley (Eds), Cambridge
- 978 University Press, Cambridge, UK, and New York, NY, U.S.A., 659–740, doi: 10.1017/
- 979 <u>CBO9781107415324.018, 2013.</u>
- 980 Nachtergaele, F. O.: From the Soil Map of the World to the Digital Global Soil and Terrain Database:
- 1960–2002, in: Handbook of Soil Science, Sumner, M. E. (Ed.), CRC Press, Boca Raton, H5 17, 1999.
- 983 Niu, Z. G., Gong, P., Cheng, X., Guo, J. H., Wang, L., Huang, H. B., Shen, S. Q., Wu, Y. Z., Wang,
- 984 X. F., Wang, X. W., Ying, Q., Liang, L., Zhang, L. N., Wang, L., Yao, Q., Yang, Z. Z., Guo, Z.
 - 30

- 985 Q., and Dai, Y. J.: Geographical analysis of China's wetlands preliminarily derived from
- 986 remotely sensed data, Science in China Series D: Earth Sciences, 39, 188-203, 2009.
- 987 Page, S. E., Rieley, J. O., and Banks, C. J.: Global and regional importance of the tropical peatland
- 988 carbon pool, Glob. Change Biol., 17, 798-818, doi:10.1111/j.1365-2486.2010.02279.x, 2011.
- 989 Powlson, D. S., Whitmore, A. P., and Goulding, K. W. T.: Soil carbon sequestration to mitigate
- 990 climate change: a critical re-examination to identify the true and the false, European Journal of
- 991 Soil Science, 62, 42-55, doi:10.1111/j.1365-2389.2010.01342.x, 2011.
- 992 Pregitzer, K. S., and Euskirchen, E. S.: Carbon cycling and storage in world forests: biome patterns
- 993 related to forest age, Glob. Change Biol., 10, 2052-2077, doi:10.1111/j.1365-
- 994 2486.2004.00866.x, 2004.
- *R* Development Core Team: *R*: A language and environment for statistical computing, R Foundation
 for Statistical Computing, Vienna, 2011.
- 997 Raich, J. W., Potter, C. S., and Bhagawati, D.: Interannual variability in global soil respiration, 1980-
- 998 94, Glob. Change Biol., 8, 800-812, doi:10.1046/j.1365-2486.2002.00511.x, 2002.
- 999 Reich, P.: Soil organic carbon map, USDA-NRCS,
- 1000
 http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/use/?cid=nrcs142p2_054018
 last access:

 1001
 2015-02-19, 2000.
- Scharlemann, J. P. W., Tanner, E., V. J., Hiederer, R., and Kapos, V.: Global soil carbon:
 understanding and managing the largest terrestrial carbon pool, Carbon Management, 5, 81-91,
- 1004 doi:10.4155/CMT.13.77, 2014.
- 1005 Shangguan, W., Dai, Y., Duan, Q., Liu, B., and Yuan, H.: A global soil data set for earth system
- 1006 modeling, Journal of Advances in Modeling Earth Systems, 6, 249-263,
- 1007 doi:10.1002/2013MS000293, 2014.
- 1008 Stockmann, U., Malone, B. P., McBratney, A. B., and Minasny, B.: Landscape-scale exploratory
- radiometric mapping using proximal soil sensing, Geoderma, 239–240, 115-129,
- doi:10.1016/j.geoderma.2014.10.005, 2015.

Gelöscht: http://soils.usda.gov/use/worlds oils/mapindex/soc.html Martin Köchy 2015-2-19 16:44 Gelöscht: 2011-09-20



- 1014 Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., and Zimov, S.: Soil organic
- carbon pools in the northern circumpolar permafrost region, Glob. Biogeochem. Cycles, 23,
 GB2023, doi:10.1029/2008GB003327, 2009.
- Tarnocai, C., I. M. Kettles, and B. Lacelle: Peatlands of Canada database, Open File, 4002, Geological
 Survey of Canada, doi:10.4095/213529, 2002.
- Turunen, J.: Development of Finnish peatland area and carbon storage 1950–2000, Boreal
 Environment Research, 13, 319-334, 2008.
- Wei, X., Shao, M., Gale, W., and Li, L.: Global pattern of soil carbon losses due to the conversion of
 forests to agricultural land, Scientific Reports, 4, 4062, 2014.
- Wieder, W. R., Boehnert, J., and Bonan, G. B.: Evaluating soil biogeochemistry parameterizations in
 Earth system models with observations, Glob. Biogeochem. Cycles, 28, 211-222,
 doi:10.1002/2013GB004665, 2014.
- 1026 Yan, Y., Luo, Y., Zhou, X., and Chen, J.: Sources of variation in simulated ecosystem carbon storage
- 1027 capacity from the 5th Climate Model Intercomparison Project (CMIP5), Tellus Series B –
- 1028 Chemical and Physical Meteorology, 66, 22568, doi:10.3402/tellusb.v66.22568, 2014.
- 1029 Zimov, S. A., Davydov, S. P., Zimova, G. M., Davydova, A. I., Schuur, E. A. G., Dutta, K., and
- 1030 Chapin, F. S.: Permafrost carbon: Stock and decomposability of a globally significant carbon
- pool, Geophys. Res. Let., 33, L20502, doi:10.1029/2006GL027484, 2006.



1034 Table 1. Definition of terms with respect to organic soil carbon.

Term	Abbreviation/Ac	ronymDefinition	
Concentration	Corg	organic carbon mass/soil dry mass, Corg	Martin Köchy 2015-2-19 16:50
Areal density (of		$C_{org} \times depth \times (1 - fractional volume of rocks,$	Martin Köchy 2015:2:26 17:50
fine soil)		coarse roots, and ice)	Formatierte Tabelle
Stock	m _C	areal density of fine soil integrated over all layers	Martin Köchy 2015·2·19 16:54
		to a specified depth	Formatiert: Tiefgestellt
Mass		stock integrated over a specified area	Martin Köchy 2015-2-19 16:58
	BD	bulk density	Formatiert: Theigestellt
	CAMP	Circum-Arctic Map of Permafrost and Ground Ice	
		Conditions	
	DSMW	Digital Soil Map of the World	
	<u>GLCC</u>	Global Land Cover Characteristics database	
	GLWD	Global Lakes and Wetland Database	
	<u>GPD</u>	Global Peatland Database	Martin Köchy 2015·2·26 17:51
	HWSD	Harmonized World Soil Database	Formatiente Tabelle
	IGBP	International Geosphere Biosphere Program	
	NCSCDB	Northern Circumpolar Soil Carbon Data Base	
	PTF	pedotransfer function	
	SMW	Soil Map of the World	Martin Köchy 2015·2·26 17:50
	SOC	soil organic carbon	Formatiente Tabelle
	SOTER	Soil and Terrain Database	Martin Köchy 2015·2·26 17:52
	SOTWIS	Harmonized continental SOTER-derived database	Formalierte Tabelle
	WISE	World Inventory of Soil Emission Potentials	

- Table 2. Changes to the global SOC mass in the top 1 m after each adjustment to the HWSD 1037
- v.1.1 database. 1038

1040

v.1.1 database.			Gelöscht: after
			Martin Köchy 2015·2·27 11:07
			Gelöscht: modifications
processing step	SOC mass (Pg)	•	
no adjustment	2469.5		Martin Köchy 2015·2·27 08:57
(1) filling of missing values for C _{org}	2470.6		Formatierte Tabelle
(2) and filling of missing values for BD	2471.3		Martin Köchy 2015·2·27 11:07
$(3a)$ and adjusting BD values when $C_{org} > 3\%$	1230.2		Gelöscht: modification
(3b) or replacing BD values only for Histosols	1113.3		
(4) = (3a) and (3b)	1060.9		Martin Köchy 2015·2·19 17:16
			Gelöscht: (–)
			Martin Köchy 2015·2·19 17:19

Martin Köchy 2015·2·27 11:04

Gelöscht: adjusting BD values for C_{org}>3% &

Martin Köchy 2015·2·19 17:19 Gelöscht: replacing BD for Histosols

1048 Table 3. Soil organic carbon masses by continent. For the definition of 'continents' we used

the ESRI (2002) map of continents with coastlines extended by 2 pixels to increase the

Martin Köchy 2015·2·27 11:05 Gelöscht: stocks

1050 overlap. 1 $Mm^2 = 10^6 km^2$

Continent	Soil area (Mm ²)	SOC mass, 0-1 m (Pg)
converted to 30" raster		HWSD v.1.1-modified
Asia, incl. Malay Archipelago	42.0	369
North America, incl. Greenland, Central America	21.3	223
Europe, incl. Iceland, Svalbard, Novaya Zemlya	9.4	110
Africa, incl. Madagascar	27.2	148
South America	17.7	163
Australia, New Zealand, Pacific Islands	8.0	46
non-overlapping pixels	0.2	2
total (90°N – 60°S)	125.8	1061

Martin Köchy 2015·2·27 11:05 Gelöscht: Carbon Martin Köchy 2015·2·27 11:05 Gelöscht: stock

1051

1049

1055 Table 4. Organic carbon mass (top 1 m) of soils with gelic properties in HWSD v.1.1-

1056 adjusted. (All areas north of 60°S). Percentiles refer to the distribution of C stocks in each cell

within the soil area mentioned. 1 $Mm^2 = 10^6 \text{ km}^2$. Hist/soil: fraction of soil area covered by

Martin Köchy 2015·2·27 11:08 Gelöscht: modified

1058 Histosols.

Gelic phase	Cell area (Mm ²)	Soil area (Mm ²)	Hist/ soil	C sto	C stock (kg m ⁻²), percentiles				
				5%	25%	50%	75%	95%	-
continuous, >90% of area	5.46	5.30	12%	5.9	7.4	7.6	12.6	38	65.2
discontinuous, 50–90%	4.11	4.07	12%	6.4	6.5	9.5	15.8	28.9	51.8
sporadic, 10–50%	3.79	3.68	6%	3.8	8.3	12.5	15.6	19	45.3
isolated, 0–10%	0.05	0.05	86%	8.4	27.9	32.8	32.8	32.8	1.5
whole area	13.41	13.10	11%	5.3	6.9	9.8	15.6	30.6	163.8

1059

1061 Table 5. Comparison of organic carbon stocks (top 1 m) between HWSD v.1.1-<u>adjusted</u> and

1062 NCSCDB (Tarnocai et al. 2009). Permafrost contingency refers to the Circumarctic

1063 Permafrost Map. NCSDB used different soil areas <u>within grid cells</u> than HWSD. Percentiles

- refer to the distribution of C stocks in each grid cell within the soil area mentioned. $1 \text{ Mm}^2 =$
- 1065 10^6 km^2 .

	HWSD								NCSDE	3		
Permafrost contingency,	Cell area (Mm ²)	Soil area (Mm ²)	C	C stock (kg m ⁻²), percentiles					Soil area (Mm ²)	C stock (kg m ⁻²), mean	C mass (Pg)	
			5%	25%	50%	75%	95%					
continuous, >90% of area	10.64	9.97	4.1	6.5	8	14.6	18.8	105.8	10.1	29.5	299	
discontinuous, 50–90%	3.17	3.05	4.4	6.9	12.9	16.9	32.6	41.3	3.1	21.8	67	
sporadic, 10–50%	3.08	2.94	4.9	7.4	12.7	17	35.5	40.3	2.6	24.3	63	
isolated, 0–10%	3.67	3.55	5.6	7.8	10.1	16	32.3	45.4	3.0	22.6	67	
whole area	20.55	19.52	4.4	6.9	9.4	15.5	28	232.7	18.8	26.4	496	

Martin Köchy 2015·2·26 17:00 Gelöscht: of NCSDB

Martin Köchy 2015-2-27 Gelöscht: modified

1066

Table 6. Area and spatial overlap of wetland types in GLWD and GLCC (grid cell area, Mm²) within the extent of the HWSD.

GLWD		GLCC, ec	osystems l	egend				
	area	14 Inland Water	45 Marsh Wetland	13 Wooded Wet Swamps	72 Man- grove	44 Mire, Bog, Fen	36 Rice Paddy and Field	Dry- land
	(Mm)	2.339	0.062	0.083	0.048	0.797	2.406	128.033
1–3 Lake, Reservoir, River	2.370	1.437	0.000	0.002	0.006	0.027	0.008	0.845
4 Freshwater Marsh, Floodplain	2.487	0.077	0.015	0.003	0.006	0.058	0.167	2.155
5 Swamp Forest, Flooded Forest	1.154	0.041	-	0.013	0.001	-	0.006	1.090
6 Coastal Wetland	0.413	0.015	0.001	0.007	0.011	0.002	0.026	0.321
7 Pan, Brackish/ Saline Wetland	0.433	0.002	< 0.001	< 0.001	< 0.001	_	0.001	0.429
8 Bog, Fen, Mire	0.710	-	-	_	-	0.710	_	· _
9 Intermittent Wetland/Lake	0.689	0.004	< 0.001	< 0.001	< 0.001	-	0.003	0.681
10 50-100% Wetland	1.762	0.045	-	0.005	-	_		1.693
11 25-50% Wetland	3.153	0.065	-	< 0.001	-	-	· –	3.077
12 Wetland Complex (0-25% Wetland)	0.898	<0.001	-	-	-	_	0.046	0.846
Dryland	120.433	0.646	0.045	0.052	0.024	_	2.149	116.896

Table 7. Organic carbon stocks and masses in the top 1 m of global wetland soils derived 1073

from the HWSD v1.1-<u>adjusted</u>. Wetland extent is primarily <u>defined</u> according to the Global Lake and Wetlands Database (1–12), augmented by wetland in the GLCC (13–72). 1074

1075

Percentiles refer to the distribution of C stocks in each grid cell within the soil area 1076

mentioned. SOC mass of permanent wetlands (types B-I) is 81.8 Pg, that of all wetlands 1077 except open waters (types B–K) is 158.1 Pg. 1 $Mm^2 = 10^6 km^2$. Hist/soil: fraction of soil area 1078 covered by Histosols.

1079 1080

W	Wetland type		Soil area	Hist./ soil		C stock (kg m ⁻²), percentiles					C mass (Pg)
	GLWD and GLCC category	(Mm ²)	(Mm ²)	%		5%	25%	50%	75%	95%	-
A	1–3 Lake, Reservoir, River 14 Inland Water	3.01	2.11		7	4.2	6.5	9	14.2	24.6	22.8
В	4 Freshwater Marsh, Floodplain 45 Marsh Wetland	2.53	2.48		17	4.4	7	10	19.1	38	32.3
С	5 Swamp Forest, Flooded Forest 13 Wooded Wet Swamps	1.21	1.21		6	3.6	5.6	8.6	13.6	33.8	13.2
D	8/44 Bog, Fen, Mire	0.71	0.68		14	4.4	8.4	14.9	18.3	35.4	10.3
Е	7 Pan, Brackish/ Saline Wetland	0.43	0.31		<1	2.8	4	4.7	5.4	7.5	1.5
F	6 Coastal Wetland 72 Mangrove	0.44	0.43		4	3.9	6.1	7.3	11.8	21.9	4.4
G	36 Rice Paddy and Field	2.15	2.14		<1	4.7	6	7.1	8.9	12.1	17.1
Н	9 Intermittent Wetland/Lake	0.69	0.60		<1	2.3	3.6	4.4	5.9	9.6	3
Ι	10 50-100% Wetland	1.75	1.74		33	6.9	12.5	13.7	24.4	38	31.1
J	11 25-50% Wetland	3.14	3.11		10	5.6	8.8	12.3	14.6	28	38.5
K	12 Wetland Complex (0-25% Wetland)	0.9	0.89		1	5.8	5.9	5.9	7.3	12.6	6.7
Dr	yland	117.24	110.15		2	2.5	4.9	7.1	10.3	18.1	880

1081

1082

Martin Köchy 2015·2·27 11:08

Gelöscht: modified

Table 8. Organic carbon stocks and masses in the top 1 m of tropical wetland soils derived 1084

from HWSD v.1.1-adjusted. Wetlands are classified primarily according to the Global Lake 1085

and Wetlands Database (1-12), augmented by wetland classes in the GLCC (13-72).

Martin Köchy 2015. Gelöscht: modified

1086

Percentiles refer to the distribution of C stocks in each grid cell within the soil area 1087

mentioned. C mass of permanent wetlands (types B-H) is 38.3 Pg, that of all wetlands except 1088

open waters (types_B–K) is 39.9 Pg. 1 $Mm^2 = 10^6 km^2$. Hist/soil: fraction of soil area covered 1089

1090

by Histosols.

Wetland type		Cell Soil area area		Hist./ soil		C stock (kg m ⁻²), percentiles				C mass (Pg)
	GLWD and GLCC category	(Mm ²)	(Mm ²)	%	5%	25%	50%	75%	95%	-
A	1–3 Lake, Reservoir, River 14 Inland Water	0.76	0.49	2%	3.9	5.9	7.9	10.6	18.8	4.5
В	4 Freshwater Marsh, Floodplain 45 Marsh Wetland	1.27	1.26	6%	3.7	6.2	7.7	10.3	24.2	12.0
С	5 Swamp Forest, Flooded Forest 13 Wooded Wet Swamps	1.21	1.20	6%	3.6	5.6	8.6	13.6	33.8	13.2
D	8/44 Bog, Fen, Mire	0.0	0.00	0%	2.5	6.0	6.0	11.9	12.0	0.0
Е	7 Pan, Brackish/ Saline Wetland	0.12	0.10	0%	2.5	3.2	4.3	5.3	7.5	0.5
F	6 Coastal Wetland 72 Mangrove	0.31	0.31	4%	4.0	6.1	8.5	13.7	25.7	3.4
G	36 Rice Paddy and Field	1.06	1.06	1%	5.1	6.2	6.9	8.1	13.2	8.4
Н	9 Intermittent Wetland/Lake	0.22	0.20	0%	2.2	3.3	4.1	5.0	6.4	0.8
K	12 Wetland Complex (0-25% Wetland)	0.2	0.20	3%	5.0	5.9	6.5	8.2	13.2	1.6
Dryland		44.71	43.06	1%	2.2	4.3	6.1	8.5	15.2	310.6
Tropical area		49.87	47.88	1%						354.9

1092 Figure captions

- ¹⁰⁹³ Figure 1. Global stock (a) and mass (b, per 5° latitude) of organic carbon in the top 1 m of the
- 1094
 terrestrial soil calculated from HWSD v.1.1-adjusted.

 1095
 Martin Köchy 2015-2-27 11:10

 Gelöscht: modified

Martin Köchy 2015-2-27 11:10

Gelöscht: 30"

- 1096 Figure 2. Extent of permafrost in HWSD v.1.1. Colour scale: fraction of soil units within a
- 1097 **0.5** grid cell with 'gelic phase' (averaged for display to 30' resolution); red outline:
- 1098 permafrost attribute in HWSD supplementary data sets SQ1–7 at 5' resolution.

1099

- Figure 3. (a) Global distribution of important wetlands (by carbon mass) according to the
- Global Lakes and Wetlands Database and Global Land Cover Characterization. The most
- frequent wetland type is displayed within a 0.5° grid cell. Wetland types A-K are explained in
- Table 6. (b) Carbon mass in wetland soils (top 1 m) in bands of 5° latitude (calculated from
- HWSD v.1.1-modified). (c) Carbon mass in aggregated types of wetland soils (panel b).

1105

Figure 4. Fraction of Histosol area per 0.5° grid cell according to HWSD v.1.1.

1109 Figures



1110



terrestrial soil calculated from HWSD v.1.1-<u>adjusted</u>.

1113

Martin Köchy 2015·2·27 11:11 Gelöscht: modified



- Figure 2. Extent of permafrost in HWSD v.1.1. Colour scale: fraction of soil units within a
- 1117 0.5' grid cell with 'gelic phase' (averaged for display to 30' resolution); red outline:
- permafrost attribute in HWSD supplementary data sets SQ1–7 at 5' resolution.

Martin Köchy 2015·2·27 11:11 Gelöscht: 30"

1119







frequent wetland type is displayed within a 0.5° grid cell. Wetland types A-K are explained in

Table 6. (b) Carbon mass in wetland soils (top 1 m) in bands of 5° latitude (calculated from

1126 HWSD v.1.1-<u>adjusted</u>). (c) Carbon mass in aggregated types of wetland soils (panel b).

1127

1121

Martin Köchy 2015·2·27 11:11 Gelöscht: modified



Figure 4. Fraction of Histosol area per 0.5° grid cell according to HWSD v.1.1.