

**Overview of terroir  
research**

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This discussion paper is/has been under review for the journal SOIL. Please refer to the corresponding final paper in SOIL if available.

# An overview of the recent approaches for terroir functional modelling, footprinting and zoning

**E. Vaudour<sup>1,2</sup>, E. Costantini<sup>3</sup>, G. V. Jones<sup>4</sup>, and S. Mocali<sup>3</sup>**

<sup>1</sup>AgroParisTech, UMR 1091 INRA/AgroParisTech “ Environnement et Grandes Cultures”, Equipe Sol, avenue Lucien Brétignières 78850 Thiverval-Grignon, France

<sup>2</sup>INRA, UMR 1091 INRA/AgroParisTech “ Environnement et Grandes Cultures”, Equipe Sol, avenue Lucien Brétignières 78850 Thiverval-Grignon, France

<sup>3</sup>Consiglio per la Ricerca e la Sperimentazione in Agricoltura, Centro di Ricerca per l’Agrobiologia e la Pedologia, CRA-ABP D’Azeglio 30 Firenze 50121, Italy

<sup>4</sup>Department of Environmental Studies 101A Taylor Hall, Ashland, OR 97520, USA

Received: 29 October 2014 – Accepted: 31 October 2014 – Published: 20 November 2014

Correspondence to: E. Vaudour (emmanuelle.vaudour@agroparistech.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

## Abstract

Notions of terroir and their conceptualization through agri-environmental sciences have become popular in many parts of world. Originally developed for wine, terroir now encompasses many other crops including fruits, vegetables, cheese, olive oil, coffee, cacao and other crops, linking the uniqueness and quality of both beverages and foods to the environment where they are produced, giving the consumer a sense of place. Climate, geology, geomorphology, and soil are the main environmental factors which compose the terroir effect at different scales. Often considered immutable at the cultural scale, the natural components of terroir are actually a set of processes, which together create a delicate equilibrium and regulation of its effect on products in both space and time. Due to both a greater need to better understand regional to site variations in crop production and the growth in spatial analytic technologies, the study of terroir has shifted from a largely descriptive regional science to a more applied, technical research field. Furthermore, the explosion of spatial data availability and sensing technologies has made the within-field scale of study more valuable to the individual grower. The result has been greater adoption but also issues associated with both the spatial and temporal scales required for practical applications, as well as the relevant approaches for data synthesis. Moreover, as soil microbial communities are known to be of vital importance for terrestrial processes by driving the major soil geochemical cycles and supporting healthy plant growth, an intensive investigation of the microbial organization and their function is also required. Our objective is to present an overview of existing data and modelling approaches for terroir functional modelling, footprinting and zoning at local and regional scales. This review will focus on three main areas of recent terroir research: (1) quantifying the influences of terroir components on plant growth, fruit composition and quality, mostly examining climate-soil-water relationships and/or using new tools to unravel the biogeochemical cycles of both macro- and micronutrients, the functional diversity of terroirs and the chemical signature of products for authentication (the metagenomic approach and the regional fingerprinting); (2) terroir zoning at

## SOILD

1, 827–906, 2014

### Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1994). In that sense, the shaping of terroirs results from a long heuristic process (likely hundreds of years) through History marked with discontinuities due to wars, spread of plagues, and wine market opportunities. For centuries, such heuristic processes have mostly been carried out on less fertile soils (fertile soils being reserved to annual crop cultivation) and without resorting to irrigation, thus accentuating multi-year variability due to vintage weather (“millésime” effect). As an entity distributed over space and time, terroir has cultural aspects that have heritage, landscape, and reputation value-added components (Tomasi et al., 2013), that come from historical empirically-derived technical adjustments, the transmission of taste typicity over generations, and strong gastronomic traditions. On the other hand, the agri-environmental aspects of terroir are likely to be conceptualized, in order to characterize, delineate and monitor zones with homogeneous or outstanding grape and/or wine, soil, geomorphological, geological, landscape, and climate characteristics at a given spatial level and over a given duration. This process may be in the nascent stages of understanding vineyard spatial management in young winegrowing regions, or refined in those winegrowing regions with long-lived wine traditions.

At the international scale, a definition focused on the agri-environmental facets of terroir was adopted in 2010 by the International Vine and Wine Organization (resolution OIV/VITI 333/2010):

an area in which collective knowledge of the interactions between the identifiable physical and biological environment and applied vitivicultural practices develops, providing distinctive characteristics of the products originating from this area. “Terroir” includes specific soil, topography, climate, landscape characteristics and biodiversity features.

Originally developed for wine, approaches for defining terroirs are now being carried to other specialty crops such as coffee (e.g., De Assis Silva et al., 2014), tea (e.g., Besky, 2014), tequila (Bowen and Zapata, 2009), honey, maple syrup, cacao, olive oil, fruits, vegetables, and cheese (Trubek, 2008; Jacobsen, 2010), ultimately linking the uniqueness and quality of both beverages and foods to the environment where they

# SOILD

1, 827–906, 2014

## Overview of terroir research

E. Vaudour et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

are produced. The trend for providing the consumer a sense of place has historically developed alongside the legal protection of these products, through either “Protected Designations of origin” (PDOs) or Protected Geographical Indications (PGIs). However, although both rely on the assumption of a deterministic relationship between food quality and agri-environmental features, legal definitions of PDOs and PGIs are not always spatially representative of the terroirs that they come from. In the European Union for instance, PDOs refer to the names of regions, given areas or even countries assigned to agricultural crops or value-added products which are produced, processed or prepared in a region according to traditional methods. In France, Italy and Spain, official PDOs that provide terroir wines legal protection from falsified wines coming from other areas date back to 1935 (Vaudour, 2003). However, there have been many historical precursors of PDOs, such as, Chianti in the XVIIth century (Tomasi et al., 2013) or in the XIXth century Jerez (Cabral Chamorro, 1987) and Champagne (Marre, 2004). Under a PDO, wine-growers, wine-makers and experts jointly define those zones that are warranted to produce wines named after their most renowned places, under common producing rules considered as traditional. These zones are generally based on pre-existing boundaries of administrative districts or easily demarcating patterns derived from hydrological networks, roads or railways. PGIs refer to the names of areas with some linkage to product quality with at least one of the stages of production, processing or preparation occurring in the considered area. Despite differing definitions, terroir is sometimes confused with PDOs or may even be confused with PGIs (Barham, 2003), when one or, more likely, several terroir units may constitute the delimited areas within the PDO, be included in them, or even intersect them.

Whatever winegrowing region in the world, either valuing inherited management zones, or attempting to construct them, the so-called “natural” components of terroir actually result from a set of processes, which together with viticultural practices create a delicate equilibrium and regulate its effect on products in both space and time (Van Leeuwen et al., 2004; Deloire et al., 2005; Van Leeuwen and Seguin, 2006; Costantini and Bucelli, 2014). At its most basic understanding, the so-called “concept of terroir”

relates the sensory attributes of wine to the environmental conditions in which the grapes are grown (Van Leeuwen and Seguin, 2006; White et al., 2007; Tempesta et al., 2010).

Given the economic importance of the wine industry worldwide, there is clearly a need to better understand the spatial and temporal variability of grape composition and which spatial and cultural scales and resolutions are best suited to manage the production of terroir wines that reveal the typical qualities of terroir units across a given territory, together with minimizing the environmental impacts of this production. Underlying notions for such questioning stem from two main research areas: first, the concepts and knowledge on agro-ecosystems raised and revisited for present-day agriculture, which faces an increasing number of challenges, and the prospect to ensure various ecosystem services implementing agro-ecological practices (Doré et al., 2011; Wezel et al., 2014); second, concepts and approaches on digital soil assessments based on pedometrics and/or proxy-remote sensing techniques (Carré et al., 2007; Minasny et al., 2012; Werban et al., 2013; Hartemink and Minasny, 2014). There is a greater need to understand regional to site and site to regional variations in crop production and the growth in spatial analytic technologies is likely to facilitate downscaling and up-scaling approaches to address these needs. Together with the emergence of precision viticulture, the explosion of spatial data availability and geospatial technologies in the past 15 years have made the within-field/farm scale of study more valuable to the individual grower, resulting in greater adoption and application (Tomasi et al., 2013). Furthermore, the study of terroir has shifted from a largely descriptive regional science back to the 1990s to a more applied, technical research field at the beginning of the XXIth century. Confusion between PDOs and terroir may occur, and the strict observance of no irrigation according to historical heuristic processes of terroir, results in misleading questions such as the possible compatibility of terroir and precision viticulture (Bramley and Hamilton, 1997). However, the long process of terroir identification over time questions those practices that enhance or diminish terroir sustainability, particularly in recent times with the advent of modern viticulture. For example, viticultural soils appear

---

**Overview of terroir research**E. Vaudour et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to be exposed to degradation processes, perhaps more than ever because of improper practices of land management (Blavet et al., 2009; Follain et al., 2012; Costantini and Lorenzetti, 2013). In addition, soil contamination by copper resulting for the cumulated use of Bordeaux mixture and other copper fungicides is an increasing issue (Pieztrak and McPhail, 2004; Fernandez-Calvino et al., 2008; Chopin et al., 2008; Mirlean et al., 2007, 2009; El Hadri et al., 2012; El Azzi et al., 2013) and appears to result in modifying the spatial distribution of soil micro-organisms (Jacobson et al., 2007; Mackie et al., 2013). Because of water scarcity, irrigation may be practiced with saline water or saline effluent with possible deleterious effects on plant growth (Paranychianakis and Angelakis, 2008; Walker et al., 2009; Stevens et al., 2011) and on soil salinity, structure and quality (Crescimanno et al., 2007; Urdanoz and Aragüés, 2009). Thus, an intensive investigation of the microbial and fungal organization and their function is required, as soil microbial and fungal communities are known to be of vital importance for terrestrial processes by driving the major soil geochemical cycles and supporting healthy plant growth (Nannipieri et al, 2003; Bokulich et al, 2014).

Our objective is therefore to present an overview of existing data and modelling approaches for the functional understanding of terroir and its footprinting and zoning at local and regional scales. This review will focus on three main areas of recent terroir research: (1) quantifying the influences of terroir components on plant growth, fruit composition and quality, mostly examining climate-soil-water relationships and/or using new tools to unravel the biogeochemical cycles of both macro- and micronutrients, the functional diversity of terroirs and the chemical signature of products for authentication (the metagenomic approach and the regional fingerprinting); (2) terroir zoning at different scales including mapping terroirs and using remote and proxy sensing technologies to manage the crop system for higher quality and the monitoring of soil quality; and (3) terroir sustainability assessment and new preservation practices.



## 2 Quantifying the influences of terroir components on plant growth, fruit composition and wine quality

### 2.1 Climate-soil relationships

Grapevines are a perennial plant characterized by a great ecological plasticity adapting to a wide range of environmental conditions (Galet, 1988). *Vitis vinifera* is the main grapevine species cultivated for wine production and the latitudinal boundaries for cultivation are strongly temperature dependent and are roughly between 52° North (Southern England) and 45° South (New Zealand) (Winkler, 1962; Vaudour, 2003; Tonietto and Carbonneau, 2004), although climate change is pushing these bounds in both hemispheres (Jones et al., 2012). Grapevine functioning relies on radiation, energy, nutrient and water balances, with the following environmental drivers: incident solar radiation, temperature regime, water regime, soil mineral nutrient availability, thermal and energetic soil properties, and soil obstacles to rooting. While numerous weather and climate factors influence grapevine growth and productivity, broad suitability to climate finds that *Vitis vinifera* is limited to zones where the average growing season temperatures are 13–21 °C (Jones et al., 2012). Conditions in these regions allow for enough heat accumulation to drive phenological development and ripen numerous varieties to produce various wine styles (Winkler et al., 1974; Huglin, 1978; Huglin and Schneider, 1998; Jones et al., 2010). In all wine-growing areas, numerous mechanisms govern the soil effect on terroir with important influences on vine functioning, grape composition and consequently wine typicality. Interacting with climate and technical choices in crop management, the availability of water, nitrogen and oxygen are the main drivers of the effects of soil on terroir (Seguin, 1970, 1983, 1986; Morlat and Jacquet, 1993; Morlat, 2001; White, 2003; Deloire et al., 2005; Santesteban and Royo, 2006; Van Leeuwen and Seguin, 2006; Carbonneau et al., 2007; Zsófi et al., 2009; Costantini and Bucelli, 2014). Given a suitable climate with adequate heat accumulation and a long enough growth period to ripen, the best terroirs for wine production (in particular for red wine varieties) are often situated where soil limiting factors create mild water stress, induce

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





# SOILD

1, 827–906, 2014

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



slow but complete ripening by reducing vine vigour and berry size and increasing grape skin phenolics (Seguin, 1986; Deloire et al., 2005; Van Leeuwen and Seguin, 2006; Carbonneau et al., 2007). Moreover, in cool climate areas, soil temperature, more than air temperature, may differentiate terroir behaviour between years (Morlat and Jacquet, 1993; Barbeau et al., 1998; Morlat, 2001). According to Seguin (1986), in the highest quality terroirs, soils can compensate for extreme climatic conditions or heavy rainfall, so that their influence can become more apparent in years with poor weather.

Beyond empirical evidence recorded for such terroirs, the shaping of quality components due to specific soil components such as calcium or potassium is still confusing. Rather than the specific influence of single soil components, some ecophysiological mechanisms have been demonstrated to be highly influential in relationship to nitrogen nutrition and water supply at key stages of the vegetative cycle. As early as 1906, based on observations collected in France, Tunisia, Algeria, and California, Louis Ravaz, an agronomist in Montpellier, France suggested that grape quality and vine health relied on a balance between the flux of accumulated compounds towards either vegetative growth or grape berries, and proposed an ideal ratio between fruit and vegetation production (Ravaz, 1906). Soil water availability was later proved to influence the hormonal equilibrium of each grapevine variety, in turn regulating the expression of the genotype (Loveys et al., 1973; Hardie and Considine, 1976; Loveys and Düring, 1984; Düring, 1984; Champagnol, 1984; Davies and Robinson, 1996; Ojeda et al., 2001, 2002; Patakas et al., 2002; Ageorges et al., 2006; Santesteban and Royo, 2006; Castellarin et al., 2007, 2011; Zsófi et al., 2009; Carbonneau et al., 2011; Teixeira et al., 2013). The effect comes when auxins, gibberellins and cytokinins (growth hormones) dominate over ethylene and abscisic acid (aging hormones), and competition for sugars occurs between vegetative and reproductive sinks, resulting in perturbed secondary metabolism (which conditions the biosynthesis of anthocyanins and tannins), degradation of leaf and fruit microclimate and disease development, and thus low potential for high quality grapes (Champagnol, 1984). While severe water deficit results in low sugar concentration as a result of restricted carbon assimilation (Roby

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Matthews, 2004; Santesteban and Royo, 2006), mild to moderate water deficit has a beneficial effect on wine phenolic composition as well as sensory characteristics (Zsófi et al., 2009, 2011). Beneficial water deficit retards shoot growth without notably affecting photosynthetic activity, which facilitates the distribution of sugar in the berries and perennial organs during ripening (Deloire et al., 2005).

In the last decade, progress has been made towards formalizing water-grapevine relationships and better predicting and diagnosing water stress, either from soil water measurements (Trambouze and Voltz, 2001; Pellegrino et al., 2004, 2006), plant water measurements (Acevedo-Opazo et al., 2008a,b, 2010a,b, 2013) and/or carbon isotope discrimination (Gaudillère et al., 2002; de Souza et al., 2005; Gómez-Alonso and García-Romero, 2010; Costantini et al., 2013). Water stress diagnosis is commonly performed through the measurement of predawn leaf water potential ( $\Psi_b$ ) using pressure chamber methodology (Scholander et al., 1965), but this measurement is time-consuming and not easy to replicate over both time and space. Alternatives approaches can be based on geospatial technologies (e.g. Acevedo-Opazo et al., 2008; Galleguillos et al., 2011a, b, Sect. 3) or rely on simulation of the soil water deficit with a simple balance (Lebon et al., 2003; Costantini et al., 2009) or more sophisticated models, such as the soil-water-atmosphere-plant (SWAP) model (Bonfante et al., 2011). The latter enables one to calculate the fraction of transpirable soil water (FTSW), which was shown to be tightly linked to  $\Psi_b$  (Pellegrino et al., 2004). FTSW is the ratio of the available soil water on a given date to total transpirable soil water (TTSW), however the difficulty is to retrieve TTSW, as it implies laborious measurements of soil moisture content over the rooting profile of the vine (to a depth of at least 2.5 m or up to 3.5 m in rainfed vineyards, insofar as no changes to soil water content are detected below that depth, according to Pellegrino et al., 2004 and Guilpart et al., 2014).

As a tool to compensate for severe water deficits in vineyards in regions with semi-arid climates, irrigation is likely to modify terroir expression (Van Leeuwen and Seguin, 2006). Under dry climate conditions in numerous wine regions worldwide, irrigation is to maintain yields, however this practice needs to be managed throughout the vegetative

# SOILD

1, 827–906, 2014

## Overview of terroir research

E. Vaudour et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

cycle in order to maintain a mild water deficit, by means of deficit irrigation or through the so-called “partial rootzone drying” (PRD) method which consists in giving a partial amount of water to one side of the root system, while the remainder is left to dry (Stoll et al., 2000; de Souza et al., 2003, 2005; Intrigliolo and Castel, 2009; Buchetti et al., 2011). PRD relies on hormonal signals, probably the abscisic acid (ABA) originating from the roots in response to the low water potential within the dry zone (Loveys and During, 1984; Stoll et al., 2000). The ABA concentration in leaves thus increases, resulting in reducing stomatal conductance and an overall improvement of water use efficiency without significant crop yield reduction.

In addition to soil water functioning, important grapevine processes are driven by the main macronutrients of nitrogen, potassium and phosphorus which influence quality components in the grape and the microbiological processes of fermentation. Along with water, nitrogen may be supplied to the grapevine through different pathways according to the development stage (Conradie, 1980, 1992; Zapata et al., 2004; Nendel and Kersebaum, 2004): (i) starting at budbreak and up to flowering, the grapevine relies on the mobilization of N reserves in the woody tissues; (ii) from flowering to veraison, the plant supplied N comes from the uptake of soil N through the roots and/or the mobilization of the N reserves in the roots; (iii) after veraison, because of possible dry conditions, N can be supplied through chemical manuring via irrigation water (fertigation). Low nitrogen constrains vegetative growth and favours berry development, nevertheless it should be made available to grapevines during fruit maturation to obtain well-developed berries with an amino acid content (of  $120 \text{ mg L}^{-1}$  ammonium N at least) that sustains vinification (Rodriguez-Lovelle and Gaudillère, 2000; Carbonneau et al., 2007). In addition to total N, grapevine nitrogen nutrition has an impact on grape berry and/or juice components, such as total amino acids, arginine, praline and ammonium, and therefore yeast-assimilable nitrogen (YAN) (Bell and Hencke, 2005). Amount and type of nitrogen supplement can substantially modulate wine volatiles composition and perceived aroma, as demonstrated for Sauvignon blanc (Choné et al., 2006), Chardonnay (Torrea et al., 2011), several Vinhos Verdes cultivars (Moreira et

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



al., 2011), and the aromatic Albariño variety characterised by terpenes (Vilanova et al., 2012). Albariño varietal compounds in the free fraction (limonene, linalool,  $\alpha$ -terpineol,  $\alpha$ -ionone and  $\beta$ -damascenone) and bound (limonene, linalool and  $\alpha$ -terpineol), as well as most yeast-derived fermentation products (including esters, higher alcohols and volatile acids) are associated with moderate nitrogen concentrations ( $\sim 350 \text{ mg N L}^{-1}$ ), while free  $\beta$ -ionone, bound geraniol, bound  $\alpha$ -ionone, bound  $\beta$ -damascenone and 1-hexanol are associated with high nitrogen concentrations (Vilanova et al., 2012). For the Sauvignon blanc variety, improving the nitrogen supply of the vine after bloom increases its aromatic potential in terms of thiol-based aromas, through the biosynthesis of cystein precursors (Choné et al., 2006).

The water regime strongly impacts the nitrogen dynamics, particularly under Mediterranean climate (Celette and Gary, 2013), even more than rainfall erosivity may result in N losses by runoff (Ramos and Casasnovas, 2006). Very poor eroded soils can show low or very low nitrogen content, as a consequence of low soil organic matter. The nitrogen deficiency can be enhanced in moderate to severe water stress conditions (Costantini et al., 2013). When soils are biologically very poor, even organic farming may be not enough to restore soil functionality, at least over a short-time period (Costantini et al., 2014).

In the case of competition with an inter-row cover crop, the grapevine may redistribute its root system to better explore deeper layers of soil (Celette et al., 2005, 2008). Considering that grapevine varieties differ in their reactivity and stability regarding the cultivation site, in the case of competition for resources terroir dictates the phenology of cultivar (Bucelli and Costantini, 2009) and some varieties, such as Sangiovese, are capable of readily responding to small pedological and climatic variations (Palliotti et al., 2008).

Given that grapevine clusters are a very important sink for potassium, particularly during ripening, this element constitutes as much as 70 % of the mineral cations and ultimately influences the pH value of juices (Hale, 1977; Carbonneau et al., 2007).

Potassium in excess tends to result in decreasing free tartaric acid, and thus the ratio tartrate to malate, one criterion of grape quality.

In addition to soil water physical properties or the content of macronutrients, some other qualities of soil implied in these variations have been experienced through a series of empirical relationships. While the specific influences are not comprehensively understood presently, it appears that optical and thermic properties of the soil surface (in particular, colour and stoniness), lime content, soil salinity, soil micronutrients such as sulphur, zinc and manganese, soil micro- and macrofauna and the biological mechanisms all influence the terroir effect (see Sect. 2.2). All these properties are expected to play an important role in shaping grape composition and thus wine typicity, but scant evidence exist of their precise determinism.

Active lime content and salinity are known to affect vine physiology and particularly nutrient balance. Active lime, which is  $\text{CaCO}_3$  in the clay and silt fraction (0–0.05 mm) is measured by extraction in ammonium oxalate (Drouineau, 1942) and has long been identified as determining the iron chlorosis in grapevines (Houdaille and Sémichon, 1894). It can be related to the free iron content in soil extracted by ethylenediaminetetraacetic acid (EDTA) to give a chlorosis-risk index (Juste and Pouget, 1972), a useful tool for selecting most rootstock varieties, which, except those derived from *Vitis Berlandieri*, originate from American species of *Vitis* that are more sensitive to chlorosis than *Vitis vinifera*. From studies carried out in Alto Adige (Northern Italy) with Schiava vines, Fregoni (2005) reports that total polyphenols of grapes increased with the increase of active lime in soil, but possibly when this active lime does not result in severe chlorosis. In addition, Martín et al. (2010) have shown that grape composition parameters are negatively affected by a decrease in chlorophyll leaf pigments due to severe chlorosis.

Conversely, soils deprived of calcium carbonate and with pH lower than 5 are likely to give rise to metal phytotoxicity, particularly due to Cu accumulation (higher than 500 ppm in some vineyard soils) resulting from the long-term application of fungicidal Bordeaux mixture (Pieztrak and McPhail, 2004; Fernandez-Calvino et al., 2008; Chopin

## SOILD

1, 827–906, 2014

### Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 2008; Mirlean et al., 2007, 2009; El Hadri et al., 2012; El Azzi et al., 2013). In all cases, those nutrient imbalances result in weakening the grapevine.

Soil salinity, as assessed through the electrical conductivity of the soil saturation extract (EC<sub>e</sub>), is likely to affect vine performance, either slightly (EC<sub>e</sub> of 2–4 dS m<sup>-1</sup>), moderately (EC<sub>e</sub> of 4–8 dS m<sup>-1</sup>), strongly (EC<sub>e</sub> of 8–16 dS m<sup>-1</sup>) or lethally (EC<sub>e</sub> > 16 dS m<sup>-1</sup>) (White, 2003). Grapevine response to salinity involves, potentially producing a reduction in transpiration and growth, due to the osmotic effect, or toxic ion uptake and altered physiological processes that may result in mortality when sodium and chlorine contents sharply rise in leaves (Shani and Ben-Gal, 2005). An increase in soil salinity results in reduced berry weight (Walker et al., 2002), and increased leaf δ<sup>13</sup>C (Costantini et al., 2010; Stevens et al., 2010). In the Chianti area, Costantini et al. (2010) demonstrated that Sangiovese performance was significantly improved by slightly saline deeper horizons, which increased plant water stress during berry ripening and reduced production. In Sicily, the presence of deeper saline horizons (7.6 dS m<sup>-1</sup> in average from 55 to up to a depth of 105 cm) was found to favour colour intensity and “salty”, “citrus”, and “fruit” sensory characteristics in the Nero D’avola wines (Scacco et al., 2010).

In addition to flavour and aroma characteristics of wines reviewed through sensory analysis, recent studies have demonstrated that variations in vine performance at the within-vineyard scale can be also monitored through the direct analysis of specific aromatic molecules found in the skins of berries. This has been shown for compounds like rotundone, which is responsible for the wine style and “pepperness” of Shiraz (Scarlett et al., 2014). Scarlett et al. (2014) recently demonstrated that within-vineyard spatial variability of rotundone biosynthesis within a vineyard in the region of Victoria (Australia) was related to soil properties and topography, which ultimately affect water availability, air and soil temperature.

## SOILD

1, 827–906, 2014

### Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.2 New tools for assessing terroir footprints: metabolomics, metagenomic approach and microbial/chemical fingerprinting

New technologies such as metabolomics and metagenomics, are promising for understanding the functioning of the terroir effect in the vine. Metabolomics seek to identify the chemical fingerprint of a particular cell, organ, organism or tissue-type that result from and are thus indicative of the chemical processes that occur within the specimen of analysis (Tarr et al., 2013). Using metabolite and gene transcript profiling, Castellarin et al. (2007) highlighted the biosynthesis pathway of flavonols with relationships to water deficit and nitrogen nutrition. The first reaction involves the deamination of phenylalanine by the enzyme phenylalanine ammonia lyase (PAL), into cinnamic acid, thus diverting phenylalanine from the pathway that relates carbohydrates to the synthesis of proteins. Metabolomic studies have also been conducted through  $^1\text{H}$  NMR spectroscopy which revealed that grapes grown in regions with high sun exposure and low rainfall showed higher levels of sugar, proline, Na, and Ca together with lower levels of malate, citrate, alanine, threonine, and trigonelline than those grown in regions with relatively low sun exposure and high rainfall (Son et al., 2009). The sensitivity of this method has allowed it to be successfully applied to classify wines according to their phenolic profile and allowed the discrimination between wines from different wineries of the same wine-producing zone and between different vintages for wines of the same variety in Greece (Anastasiadi et al., 2009) and Spain (López-Rituerto et al., 2012).

Wines from distinct countries or regions can be also discriminated through their chemical composition and/or sensorial profiles. A recent study quantifying a large number of elements (33) in wine and soil samples analysed by quadrupole inductively plasma mass spectrometry (Q-ICPMS) in addition to  $^{87}\text{Sr}/^{86}\text{Sr}$  isotopic analysis allowed for differentiation among wine-producing regions of Argentina (Di Paola-Naranjo et al., 2011). In this case the  $^{87}\text{Sr}/^{86}\text{Sr}$  ratio was amongst the best discriminators and the value ranges of which were similar between soils and wines. This corroborates the results obtained by Marchionni et al. (2013) across six distinct wine appellations

SOILD

1, 827–906, 2014

### Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Overview of terroir  
research**

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in Italy, but in their study they focused on parent material and this ratio matched those observed for homogeneous soil parental material such as volcanic rocks. Studying the whole parent material-soil-plant-wine chain at a single rainfed vineyard located in the volcanic area of Campi Flegrei (Southern Italy), and considering both a set of putative rare earth elements geotracers and the Sr isotope ratio, Mercurio et al. (2014) found that only this isotopic ratio was “consistently and inherently transferred and maintained from geologic parent material to wine, through soil horizons, branches, leaves, and grapes”. However, considering several vintages, Marchionni et al. (2013) observed that wines showing Sr isotopic ratios matching those of the underlying substrates were mostly originating from vineyards grown on volcanic rocks, unlike wines from vineyards on sedimentary or granitic rocks. As isotopic ratios of geological substrates belonging to different geological districts may partially overlap, Mercurio et al. (2014) recommend assessing the isotopic ratios together with a pertinent soil classification for a reliable assurance of wine provenance. Geographical fingerprinting of wines may also apply at distances as short as 4 km. Research by Tarr et al. (2013) found that more than a thousand components were identified from high performance liquid chromatography of juices from two varieties (Syrah and Grenache) originating from two distinct terroirs roughly 4 km from each other. Hierarchical clustering of data peaks suggested that terroir played a large part in the final composition of the grape berry metabolome (Tarr et al., 2013).

Another emerging research area refers to the exploitation of vine-associated microorganisms. In fact the commensal microbial flora that coexists with the plant may be one of the key factors that influence the plant traits. To date the role of microbes has been largely ignored, except for the microbial communities present at the surface of the grape berry, which are known to be very large and change according to the stage of grape development (Barata et al., 2012; Pinto et al., 2014). The microbiological life of wine starts before reception and fermentation of the grapes at the winery. In particular the yeast population and bacterial and fungal consortia inhabiting grape surfaces could reflect a wine region, as reported in some recent studies (Renouf et

**Overview of terroir  
research**

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



al., 2005; Setati et al., 2012; Bokulich et al., 2013). However, determinants of regional wine characteristics have not been identified. Renouf et al. (2007) identified 52 yeast species and 40 bacteria. The majority of the bacterial groups were present in the study, in particular the proteobacteria, which are not commonly described in oenology, while the most common oenological yeast (*S. cerevisiae*, *B. bruxellensis*) and bacteria (*O. oeni*, *P. parvulus*, *G. oxydans*) were detected on grape skins from the first stages of development.

Bokulich et al. (2013) surveyed 273 grape musts from two vintages of Chardonnay, Zinfandel, and Cabernet Sauvignon, demonstrating that the grape surface microbial communities present were significantly different between regions. However, the authors also show that the degree of significant differentiation between regions is increased dramatically when they look at the biogeography within a grape variety of a given vintage. This finding suggests that other factors also play a significant role, including host genotype, phenotype (grape variety), local and inter-annual climate variation (vintage) and soil quality.

Information on the impact of soil microbial communities on soil functions, grapevine plants, and wine quality is still one of the biggest challenges of soil science (Gilbert et al., 2014). The interface between roots and soil (rhizosphere) is often considered the key point of interaction between a plant and its environment. However, microbes colonizing at the root can also migrate through the plant to colonize aerial tissues, either internally (endophytes) or externally (epiphytes) (Compant et al., 2011; Bulgarelli et al., 2013). In a recent work Martins et al. (2013) observed the interaction between telluric bacterial communities and the epiphytic bacteria present on the different grapevine parts. Yet, the ecological interactions and the role of such organisms are still not clear. Indeed, most of the available literature reports studies concerning the bacterial structure and plant-associated microbiomes (Gilbert et al., 2014) but the functional diversity of such microbiomes are still largely ignored. The use of high-throughput culture-independent methods, such as microarrays or metagenomics, might elucidate such aspects. For example, considering soils from two close sites in Central Tuscany

cultivated with the same variety Sangiovese, but with contrasting wine quality, and water stress, Mocali et al. (2013) used functional GeoChip microarrays to explore both microbial composition and functions. Preliminary results revealed different amounts of Actinobacteria and Proteobacteria among the two sites and an overrepresentation of sulfur-oxidation genes in samples where both the increased level of sulfates and the abundance of Firmicutes such as *Sulfobacillus thermosulfidooxidans* occurred. Furthermore, sequencing the entire microbial genomes present in a vineyard soil sample and referred to as the “metagenome”, can provide a cultivation-independent assessment of the largely unexplored genetic reservoir of soil microbial communities and their functions (Daniel, 2005; Mocali and Benedetti, 2010). The importance of this approach has led to the establishment of “Terragenome”, an international consortium for the exploitation of the soil metagenome (Vogel et al., 2009). While these studies represent an initial examination of these relationship, it is an essential step which may potentially help to revolutionize how sites for agriculture are chosen, or indeed how they could be manipulated by probiotics designed to select suitable bacterial species, which could improve soil quality and, hence, crop productivity. Further research is needed to study the degree to which it could enhance some wine characteristics of given terroirs in sites a priori not suitable for generating such characteristics.

### 2.3 The perspective of climate change

Given the importance of climate in terroir influences on grapevine growth, productivity and the resulting wine quality, changes in climate have the potential to alter numerous aspects of terroir functioning and zoning at different scales (Jones et al., 2012). Potential responses to changing climates reflect the interactions between temperature, water availability and timing, increasing soil salinity and nutrient stresses, and increasing carbon dioxide concentrations (Bisson et al., 2002). As such, understanding impacts on viticulture and wine production from climate change necessitates integrated information and research examining the combined effects of these and other factors. Recent research on aspects of global environmental change on viticulture and wine production

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**Overview of terroir  
research**


---

 E. Vaudour et al.
 

---

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


reveal significant changes and many unknowns (Fraga et al., 2012). From a general climate perspective, wine regions worldwide have seen changes in average climate structure producing warmer and longer growing and dormant periods (Jones et al., 2005). Depending on the underlying emission scenario, climate models predict continued increases in global temperature of 1.3 to 4.8 °C by the end of this century along with increased variance in weather and climate extremes (IPCC, 2013). For wine regions, projected changes in climate point to a progressive warming in all bioclimatic indices as well as increasing water deficit, earlier phenological events, notably over the European domain (Fraga et al., 2012; Moriondo et al., 2013) but with strong regional contrasts (Santos et al., 2012), and quality changes (Jones et al., 2005; Santos et al., 2013). The possible impacts that climate change may have on viticultural suitability in the next 40 years is controversial. Some research has predicted a large decrease (up to 70–80 %) in the area suitable for viticulture in major wine producing regions worldwide by 2050 (White et al., 2006; Hannah et al., 2013). Other research has argued that both the implementation of adaptive strategies by growers and winemakers along with the evolution of consumer's preferences will keep balance the projected changes in climate (Van Leeuwen et al., 2014). The truth is likely somewhere in between, however a better understanding of terroir influences will arguably lead to lower vulnerability and greater adaptive capacity in the wine industry.

As a prominent aspect of terroir soil strongly influences yield and quality in wine production. Soil erosion, degradation and salinity are likely to be major indirect effects of climate change on viticulture and wine production (Anderson et al., 2008). Climate change is expected to lead to a more vigorous hydrological cycle, including more total rainfall and more frequent high intensity rainfall events (IPCC, 2013). Observations show that rainfall amounts and intensities increased on average globally during the 20th century (Donat et al., 2013), and according to climate change models they are expected to continue to increase during the 21st century (IPCC, 2013). These rainfall changes, along with expected changes in temperature, solar radiation, and atmospheric CO<sub>2</sub> concentrations, will likely have significant impacts on soil erosion rates (Nearing et al.,

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2004). Even in cases where annual rainfall is projected to decrease, system feedbacks related to decreased biomass production could lead to greater susceptibility for soil erosion. There will likely be non-linear effects of climate change caused by interactions between soil, climate and nutrition, in part dependent on adaptation in vine management where responses could potentially exacerbate, or ameliorate, the potential changes in erosion rates (Anderson et al., 2008; Hayman et al., 2009). In regions with very arid climates and saline water irrigation there is often a decline in soil structure and increased salinity (Clark et al., 2002; Richards et al., 2008), particularly in cracking clay soils (Crescimanno and Garofalo, 2006). Soil structure decline and increased sodicity can occur when saline water is used for summer irrigation and then subsequently the soil receives high quality rainwater during winter (Clark et al., 2002). Decreased flows in arid regions in the western US and Australia will likely result in increased salinity of irrigation water in many viticulture regions. Furthermore, a changing climate and other demands on water will increase the pressure to restrict irrigation, ultimately increasing rootzone salinity with greater impacts on wine quality over time.

Much work has been done examining the likely impacts of climate change on water and/or nitrogen dynamics through models such as Lebons's (adapted from Riou's models, 1989, 1994; Lebon et al., 2003) Lin and Host's (Costantini et al., 2009), SWAT (Ramos and Casasnovas, 2014), and SWAP (Bonfante et al., 2011) hydropedological models that aim to simulate the dynamics in seasonal soil water balance. Kersebaum's model was proposed to simulate nitrogen dynamics (Nendel and Kersebaum, 2004), while the STICS model simulates crop growth, soil water and nitrogen balances driven by daily climatic data (Brisson et al., 2009), the WaLIS model simulates water partitioning in intercropped vineyards (Celette et al., 2010). WaLIS was also used to quantify the effects of water deficit and nitrogen stress on yield components (Guilpart et al., 2014). To model the seasonal pattern of evaporation from a grassed Mediterranean vineyard, Montes et al. (2014) recently elaborated a Soil-Vegetation-Atmosphere-Transfer (SVAT) model coupling an evaporation formulation together with a reservoir-type soil water balance model. Attempts have also been made to use micromorphology to characterize

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and monitor soil internal drainage of vineyards and olive tree groves (Costantini et al., 2006). Bonfante et al. (2011) have integrated the outputs of the SWAP model with a set of regional spatial data in order to map crop water stress indices together with soil map units, bioclimatic indices, and potential radiation onto terroirs with simulated crop water status. Some of these models have been tested for predicting budburst date (De Cortazar-Atauri et al., 2009), reconstructing past climate through harvest dates (De Cortazar-Atauri et al., 2010) or could be used in order to anticipate the phenological/physiological impacts of climate change on terroir typicality, along with changes in vineyard practices such as adoption of intercrops, irrigation, and cultivar changes.

In summary, the research areas that are delineating strategies to face current global changes include numerous issues that span many aspects of the terroir continuum, including: (i) selection/introduction of new varieties in a defined winegrowing region; (ii) introduction of genes in present varieties cultivated in specific regions; (iii) selection of clones and rootstocks; (iv) selection of most suitable pedoclimatic conditions within a winegrowing region; (v) enhancing vineyard soils to specifically address new climatic conditions; and (vi) adapting or enhancing modified viticultural and/or soil management practices, and/or agroforestry practices with a given region.

### 3 Terroir zoning at different scales using geospatial technologies

#### 3.1 General perspectives

Differentiation and mapping of regions of grape/wine quality need comprehensive spatial modelling of climatic, soil, and agronomical properties, including their changes through time (Vaudour, 2002; Costantini and Bucelli, 2014). The construction of such spatial models for demarcating terroir units and predicting their viticultural/wine response is undergoing a methodological revolution as new technologies and analytical methods enable the capture of detailed spatial and temporal variability of grapevines according to functional properties in the soil. Recent developments in the combined

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



use of several geospatial technologies including geographic information systems (GIS), global positioning systems (GPS), remote sensing (RS) and direct measurements in the field (proxy sensing) are likely to open many new areas in the spatial modelling of winegrowing terroirs. A considerable amount of research dealing with terroir zoning (> 120 journal papers, book chapters and books) has been published since 2002 (Fig. 2). Most of the published research has been carried out at the within-field scale in the context of the so-called “site-specific” or “precision viticulture” (Bramley, 2004, 2005; Bramley and Hamilton, 2004; Bramley et al., 2011a, b, c, d, e; Tisseyre and McBratney, 2008; Roudier et al., 2008; Arnó et al., 2009, 2012; Pedroso et al., 2010) and their median study area covered only 0.12 ha, while between-field or regional studies covering more than 11 ha only represented 1/4 of the total (Fig. 3). Twenty or so viticultural, environmental or oenological target properties have been considered over the last decade (Table 1), mainly focusing on grape/canopy characteristics and yield or biomass, and secondarily on soil properties.

While most former terroir-studies dating back the 1970s–1990s mainly relied on conventional soil mapping and sparse but very time-consuming soil/vine/grape observations at field sites, new techniques are promising for both capturing the detailed spatial variability of vineyard areas and collecting a large amount of soil/vine/grape data. A number of terroir-related studies of the last decade have relied upon a large amount of yield data collected by means of on-the-go yield sensors mounted on mechanical harvesters, which were made commercially available beginning with the 1999 harvest. Pioneering research by Bramley and Hamilton (2004) and Bramley (2005) not only highlighted the magnitude and extent of spatial variability of yield in some Australian vineyards (of between 4.5 and 7.3 ha), but also emphasized the significant influence of soils in driving yield differences. The need to assess spatial variations in soil properties has driven development and application of direct (so-called “proximal”) geophysical sensing, particularly for measuring soil apparent electrical conductivity by means of either electrical resistivity surveys and/or electro-magnetic induction scans (EMI) (Lamb et al., 2005; Morari et al., 2009; Taylor et al., 2009; Trought and Bramley, 2011; Fulton



et al., 2011; André et al., 2012; Martini et al., 2013; Andrenelli et al., 2013; Priori et al., 2013; Rossi et al., 2013; Brillante et al., 2014). Moreover, the need to assess spatial variation in grapevine biomass and canopy properties, or to map terroir units or to identify vines, has driven the development, acquisition and processing of remote sensing data. Figure 4 shows that the development of very high spatial resolution airborne acquisitions for the purpose of characterizing grapevine physiological status has represented about 1/3 of the terroir-related studies over the last decade (Hall et al., 2002, 2003; Dobrowski et al., 2003; Lamb et al., 2004; Zarco-Tejada et al., 2005; Martín et al., 2007; Hall et al., 2008, 2013; Gil-Pérez et al., 2010; Meggio et al., 2010).

### 3.2 Remote sensing of terroir

The use of remote sensing data in the context of defining terroir and/or within-field management zones in the last decade has given rise to either straightforward use of spectral indices aimed at characterizing vigour or physiological condition (e.g. Hall et al., 2003; Johnson et al., 2003; Lamb et al., 2004; Zarco-Tejada et al., 2005; Mathews and Jensen, 2012; Hall and Wilson 2013), grape quality (Martín et al., 2007; Martínez-Casasnovas et al., 2012; Primicerio et al., 2014), or more sophisticated image processing aimed at either mapping terroir units (Pedroso et al., 2010; Vaudour et al., 2010; Urretavizcaya et al., 2013) or identifying vineyards (e.g. Wassenaar et al., 2002; Warner and Steinmaus, 2005; Rabatel et al., 2006, 2008; Delenne et al., 2010) (Table 1). Other remote sensing studies have dealt with the incorporation of remote sensing information into the prediction of soil properties and their monitoring in vineyards (Vaudour, 2008; Corbane et al., 2010; Lagacherie et al., 2012), the assessment of soil erosion patterns (Quiquerez et al., 2014; Chevigny et al., 2014) or with the prediction of vineyard evapotranspiration (Galleguillos et al., 2010a, b; Bellvert et al., 2014).

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.2.1 Use of vegetation indices for assessing vine vigour/physiology: the NDVI

Typically, the information retrieved from remote sensing has solely relied on the calculation of the most commonly used normalized difference vegetation index (NDVI) defined by Eq. (1) (Tucker, 1979) from images acquired at or near veraison.

$$5 \quad \text{NDVI} = \frac{(R_{\text{NIR}} - R_{\text{R}})}{(R_{\text{NIR}} + R_{\text{R}})}, \quad (1)$$

where  $R_{\text{NIR}}$ , is reflectance in the near infrared spectral band;  $R_{\text{R}}$ , is reflectance in the red spectral band.

A number between  $-1$  and  $+1$ , quantifying the relative difference between the near infrared reflectance “peak” of vegetation tissues and the red reflectance “trough” due to chlorophyll/carotenoids pigment absorption, the NDVI is the most widely used indicator of plant canopy vigour and relates to the leaf area index (LAI: the ratio of leaf surface area to ground area), fractional cover, biomass and shaded area (Hall et al., 2002; Johnson, 2003; Johnson et al., 2003; Hall et al., 2003; Dobrowski et al., 2008), grape quality (Fiorillo et al., 2012). In VHRS vineyard imagery, canopy-only pixels tend to have very high NDVI values, commonly ranging between 0.75 and 0.85 (Hall et al., 2002, 2003, 2008). Interpretation of NDVI threshold values requires adaptation to each case study region. This is because typical multispectral images with coarser resolution, like 4-m-IKONOS, may have values that integrate mixed signals from both vine row and inter-row, either bare soil or vegetated with an inter-row crop. However, many studies do not give sufficient detail on the retrieval of NDVI, which basically requires that the digital numbers of the raw images be atmospherically corrected into surface reflectance values prior to NDVI computation, in order to minimize differences in illumination conditions (Vaudour et al., 2014b). In an attempt to address the issue of variable environmental conditions when acquiring multi-temporal imagery, the digital numbers are either normalized (e.g. Vaudour et al., 2010) or alternatively sliced into high, medium and low values for each image (e.g., Dobrowski et al., 2003). However,

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



more often than not the relationships between NDVI and grapevine vegetation/terroir parameters are assessed on an individual image basis, using raw digital numbers (e.g., Johnson et al., 2003; Lamb et al., 2004) or even ignoring atmospheric effects. Using airborne hyperspectral sensors, such as the Compact Airborne Spectrographic Imager (CASI), the Reflective Optics Imaging Spectrometer (RODIS) and the Digital Airborne Imaging Spectrometer (DAIS-7915) and atmospherically corrected images, Zarco-Tejada et al. (2005) demonstrated that narrow band hyperspectral indices in the 700–750 nm spectral region performed better than NDVI for the purpose of estimating chlorophyll content of leaves, and then detecting iron deficiency chlorosis. Using the Airborne Hyperspectral Scanner (AHS) at a 1000 m-altitude flight enabling a spatial resolution of 2.5 m, Meggio et al. (2010) elaborated other physiological indices for predicting carotenoid and anthocyanin leaf contents, which were related to grape quality in a previous study (Martín et al., 2007).

### 3.2.2 Identification and/or characterization of vineyards

Some researchers have developed change detection classifications in order to extract regional land use changes including vineyards from multitemporal, multispectral images with medium spatial resolution such as 30 m-Landsat (Lanjeri et al., 2004; Rodríguez-Pérez et al., 2007; Manandhar et al., 2010). However, much of the work in this area over the last decade is especially marked by the advent of VHSR images, which have favoured innovative approaches for retrieving specific patterns of vineyard arrangements from helicopter colour images with  $\sim 0.25$  m resolution (Wassenaar et al., 2002), airborne multispectral images with  $\sim 2$  m-resolution (Gong et al., 2003) or 0.5 m-resolution (Rabatel et al., 2006), satellite panchromatic 1 m-IKONOS images (Warner and Steinmaus, 2005), satellite panchromatic 0.6 m-Quickbird images (Rabatel et al., 2006), and ultra-light motorized (ULM) colour 0.5 m-images (Rabatel et al., 2008; Delenne et al., 2010). Approaches for vineyard identification include grapevine field detection (Wassenaar et al., 2002; Rabatel et al., 2006), grapevine field delineation (Da Costa et al., 2007), grapevine row extraction (Hall et al., 2003; Delenne et al., 2010;

---

**Overview of terroir  
research**

 E. Vaudour et al.
 

---

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Matthews and Jensen, 2013; Puletti et al., 2014), and detection of missing plants (Chanussot et al., 2005; Delenne et al., 2010). These approaches have mostly used grey-level images and either relied on frequency analysis (Wassenaar et al., 2002; Rabatel et al., 2006; Delenne et al., 2010) or developed textural analysis, a branch of image processing focused on the spatial statistics of the grey levels of images, the variations of which are perceived as homogeneous areas by the human eye (Haralick et al., 1973). The textural analysis is based on co-occurrences matrices, i.e. “the histogram, in a given neighbourhood for each pixel (e.g.  $7 \times 7$ ), of the grey-level transitions when considering a given translation in a given direction, from which various parameters can be computed”; for instance energy, correlation, directivity, entropy, and contrast (Rabatel et al., 2006). Provided that the boundaries of each field are available, one can apply a local Fourier transform to extract information on the type of vineyard planting as well as crop spacing and orientation (Wassenaar et al., 2002) or isolate each individual field by selecting the corresponding frequencies in the Fourier spectrum using a specific Gabor filter applied recursively (Rabatel et al., 2008; Delenne et al., 2010). Another approach is that of spatial autocorrelation, requiring that grapevines be equally spaced and that the spatial resolution be fine enough that individual grapevines are discriminated (Warner and Steinbaus, 2005).

### 3.2.3 Characterization of soil types, soil properties, soil surface condition, erosion

As vineyard inter-rows are frequently left bare, particularly in Mediterranean regions, remote sensing satellite images acquired in Spring, before budburst, have been used for the purpose of mapping soil surface type and condition. For instance, high resolution multispectral 20 m-SPOT images during Spring have been processed using supervised Bayesian maximum likelihood classifier to map red Mediterranean soils originating from Plio-Pleistocene fluvial deposits in the Southern Rhone Valley, with an overall accuracy of 60–70 % (Vaudour, 2008). Being important factors in runoff and infiltration processes, soil surface characteristics (SSC) have been mapped at the within-field

scale using VHSR images with 0.1 m-resolution acquired by means of Unmanned Aerial Vehicles (UAVs) equipped with a colour camera: Pixy<sup>®</sup> (Corbane et al., 2008) or DRELIO<sup>®</sup> (Quiquerez et al., 2014). Depending on SSC classes and surface conditions, overall accuracy ranged from 63 to 84 % (Corbane et al., 2008). Clay content of viticultural soils of the La Peyne catchment in Languedoc (Southern Mediterranean France) has been successfully estimated from the 2206 nm wavelength of an airborne hyperspectral HYMAP image with 5 m spatial resolution, then spatially predicted from a co-kriging model using this image as co-variable (Lagacherie et al., 2012).

### 3.2.4 Incorporation of remote sensing information into the spatial modelling of terroirs

Remote sensing images have been used for the purpose of mapping terroir units at regional scales, facilitating selection of plant material, the assemblage of harvest and the monitoring of vine phenology and status across a number of individual fields spread over a regional viticultural area. A terroir concept adapted to spatial modelling from remote sensing considers soil landscape units as base elements for defining terroir units, jointly with climate series and grape composition series (Vaudour, 2003) (Fig. 5). A soil landscape, also referred to as soilscape (Hole, 1978) or “pédopaysage” (Girard, 1983), can be defined as a landscape unit including a limited number of soil classes that are geographically distributed according to an identifiable pattern (Lagacherie et al., 2001). Soil landscape units were also defined as “a set of pedological horizons and landscape features (vegetation, effects of human activities, geomorphology, hydrology, and parent material) whose spatial organisation allows for defining a soil mantle (or a subset of it)” (Girard, 1983; Carré and Girard, 2002). Their identifiable pattern can be retrieved from visual interpretation of several geographical data layers including image classification results and stereoscopic photograph examination (Vaudour et al., 1998; Vaudour, 2003). The visual interpretation process follows a set of rules describing a conceptual model of soil landscape organization, which relies on the assumption that soil landscape units may be inferred from the geomorphological identification

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



# SOILD

1, 827–906, 2014

## Overview of terroir research

E. Vaudour et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

of surficial formations (e.g., mainly glacio-alluvial terraces, in the Rhone Valley), the age and relative elevation of which correspond to distinct durations of pedogenesis (“fersiallitzation”) and thus distinct soil layer depths and properties, including soil surface stoniness and colour (Vaudour, 2008). In this pioneering approach, visual interpretation was digitally performed within a GIS along with description and recording of several soil landscape attributes at each delineated polygon area. Potential terroir units were then obtained from the Ward’s clustering of these soil landscape attributes and were further validated against a considerable set of grape composition data over 17 years. In an attempt to reduce the time-consuming stage of visual interpretation, alternate approaches which solely relied on automatic processing of remote sensing and/or morphometric data were proposed, either based on combining per-pixel plus textural classifier (Vaudour, 2003) or on bootstrapped regression trees (Vaudour et al., 2010). In these studies, the resulting map units were termed “terroir” and “viticultural” because they were tested against a considerable set of grape composition data over a long-term period (Vaudour et al., 1998; Vaudour, 2003), or relied on ~ 50 reference vineyards, the oenological properties of which were known from previous research (Vaudour et al., 2010). In another study, the spatial units of which were termed “terrons”, because they were not tested against viticultural data (Hughes et al., 2012; Malone et al., 2014), Landsat bands and several Landsat band ratios including NDVI have been included as co-variates of soil profile data in a number of geostatistical models in order to spatially predict several soil properties which include soil pH, clay percentage, soil mineralogy (clay types and presence of iron oxides), continuous soil classes, and presence or absence of marl: the predicted soil properties were then combined with landscape attributes (derived from a digital elevation model) through fuzzy  $k$  means in order to predict 10 non-marl “terron units” and 2 marl “terron units” depending on the presence of marl (i.e., active lime) at 0.5 m depth. Such an approach of defining “terrons” was initially proposed by Carré and McBratney (2005) and is meant as an initial stage prior to defining viticultural terroirs. A similar study using the term “natural terroir units” and derived from the geostatistical methods described by Castrignanò et

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



al. (2009) was carried out in Tuscany, but did not include remote sensing layers (Priori et al., 2014). “Natural terroir units” (NTU) were first proposed by Laville (1990), as “a volume of the Earth’s biosphere that is characterized by a stable configuration and values of the environmental factors”, and built from morphometric data and lithological units.

Because of the data availability, topography, climate, substrate, and soil are the most commonly used land features in digital terroir zoning (e.g., Herrera-Nuñez et al., 2011). Actually, there is a conceptual similarity, if not filiation, between the NTU as constructed by Priori et al. (2014) (see Sect. 3.3.3), terroir units and soil landscape units.

At the within-field scale, “management zones” originating from a set of soil and/or vegetation proxy and remotely-sensed attributes, typically NDVI, were also obtained using either fuzzy  $k$  means (Bramley and Hamilton, 2004; Pedroso et al., 2010; Taylor et al., 2013; Tagarakis et al., 2013; Priori et al., 2013; Urretavizcaya et al., 2013) or Ward’s clustering (Santesteban et al., 2013). Similar clustering approaches have been performed using a multivariate set of spatial layers including apparent electrical conductivity but without remote sensing images (e.g. Martini et al., 2013).

In order to be applicable in an operational manner, management zones need (Bramley and Hamilton, 2004): (i) to provide stable, constant patterns from year to year; (ii) be related to yield; (iii) be manageable; and (iv) be more economically beneficial than conventional uniform management. An objection to using remote sensing images for delineating management zones is that they “provide only a within-season snapshot and may not relate to final crop yield” (Taylor et al., 2007). According to Tisseyre et al. (2007), and later confirmed by Trought and Bramley (2011), “yield or vigour (pruning weight, size of the canopy) maps of the previous years are relevant in designing site-specific management strategies in the year “ $n + 1$ ” and subsequent years” and conversely to produce maps of quality parameters (sugar content, titratable acidity, pH) which present no temporal stability of within-field variability. This is in compliance with the observations made previously at the regional scale, in the terroir units mapped in the Southern Rhone Valley, for which a long-term frequency analysis of Grenache





a NDVI airborne ADS40 image with 5 m resolution, aiming at optimising variance partitioning of vine circumference data, removing small unmanageable polygons covering less than 0.1 ha.

Zoning based on automatic procedures should also enable to assess prediction error and/or error uncertainty along with predictions. In the approach elaborated by Pedroso et al. (2010), the effectiveness of the management unit delineations was determined from the adjusted  $R^2$  from ANOVA using the management units as the independent variable and grapevine circumference data as dependent variable. Most terroir units/management zones are validated following a similar variance testing procedure, i.e. demonstrating how well does delineation explain a key growth or grape/berry composition parameter (e.g. Priori et al., 2013; Santesteban et al., 2013). However, while the agronomical/soil property model is verified and/or validated, its spatial prediction error is not or seldom assessed except for geostatistical approaches (Castrignanò et al., 2009; Lagacherie et al., 2012) and for the bootstrapped approach by Vaudour et al. (2010).

### 3.2.5 Prediction of evapotranspiration and management of irrigation from remote sensing information

Recent developments of remote sensing applications to terroir-related studies have dealt with the prediction of evapotranspiration using multispectral ASTER images through the thermal infrared bands with 90 m resolution (Galleguillos et al., 2011a, b). Using the approach developed by Galleguillos et al. (2011a, b) and a linear downscaling based on land use at smaller pixel sizes, Taylor et al. (2013) demonstrated that the ASTER-derived evapotranspiration-based covariates were of particular significance in the soil depth modelling while water table depth was better explained by models that used digital terrain attributes at smaller pixel size.

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.2.6 Contribution of LiDAR and UAV remote sensing

The use of UAVs in precision viticulture is very recent and promising, as the time of acquisition is tightly controlled and adapted to the user's needs. In particular, promising approaches for mapping vine water stress were presented by Baluja et al. (2012a) and by Bellvert et al. (2014), based on 0.3 m-thermal UAV images acquired around noon (solar time), the pixels of which were significantly linked to midday water potential (MWP) measurements. UAV equipped with LiDAR sensors also enabled detection of rows, the 3D-reconstruction of a vine plantation (Llorens et al., 2011) and the quantifying of vineyard canopy (Llorens et al., 2011; Matthews and Jensen, 2013), while airborne LiDAR images enabled the mapping of landscape linear features (ditches) in a viticultural catchment (Bailly et al., 2011). UAVs equipped with visible-near infrared cameras also performed well in LAI estimation (Arnó et al., 2013a).

UAVs allow for acquiring spectral or thermal measurements that are very comparable to proxy measurements and generally carried out along with field proxy measurements. Their use, however, requires a perfect mastering of a chain of image series acquisition, acute georeferencing, spectral calibration, mosaicking and processing, which is the subject of ongoing technical developments, as shown by Verger et al. (2014) for predicting the green area index of annual crops.

### 3.3 Proxy measurements of terroir and their statistical processing

#### 3.3.1 Geophysical proxy measurements

Observations should account for the entire depth of the root-systems of vines, which may explore the parental material of the soil, often being surficial formations (Vaudour, 2002, 2003). Geophysical techniques applied to soil (Samouëlian et al., 2005; Doolittle and Brevik, 2014) offer a unique opportunity to explore deep horizons and it may be expected that key soil properties related to soil-vine water balance be retrieved from EMI or ground penetrating radar (GPR) measurements, in possible conjunction with remote

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





### 3.3.3 The issue of big data handling and the statistical processing of the varied spatial data collected

In addition to remote and proxy data collected from distinct sensors, the use of mobile devices with multitag technologies (Cunha et al., 2010) facilitates the recording of a great wealth of data. These numerous spatial data have stimulated new developments in both software and hardware, jointly with statistical processing stemming from geostatistic, image pattern recognition, satellite image processing, which includes machine learning. However the most common pattern adopted for within-field spatial data and observed for  $\sim 40$  of the last-decade terroir-related research (Table 2), relied on geostatistical analysis as emphasised by Baveye and Laba (2014). Considering the target parameter as a random property following a random process with assumption of stationarity (i.e. there is the same degree of variation from place to place), Mathéron (1962, 1965) formalized the approach to predict target properties from spatially correlated sample data through the computation of the semi-empirical variogram, to which are fitted a number of standard parametric models (Oliver and Webster, 2014). “Kriging is a generic term for a range of least-squares methods to provide the best linear unbiased predictions in the sense of minimum variance”, through solving a set of linear equations (the kriging system) from the fitted variogram function and the available data (Oliver and Webster, 2014). Ordinary kriging based on primary spatial information such as yield or ECa is the most popular method used in terroir-related studies (Table 2, e.g., Unamunzaga et al., 2014), though more sophisticated spatial models using ancillary spatial information have also been built, such as block co-kriging with an hyperspectral image (Lagacherie et al., 2012) or factorial kriging of several soil variables and ECa (Morari et al., 2009). To single out NTU-based viticultural terroir units at the province scale (1 : 125 000 scale), Priori et al. (2014) combined a multivariate and geostatistical approach showing the variability of the soilscares within the DOC and DOCG territories in Italy. A first Principal Component Analyses (PCA) was performed to relate climate, pedoclimate and morphometric features with the viticultural

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



data, and a second PCA linked the main soil features with viticultural data. The two PCAs revealed which environmental features were better related to the viticultural parameters of the experimental farms. Several geostatistical models (Castrignanò et al., 2009) were then used to spatialize the selected environmental features: (i) regression kriging to interpolate rooting depth; (ii) simple kriging with varying local means, to interpolate coarse fragments and redoximorphic mottles depth; and (iii) multicolocated simple cokriging with varying local mean to interpolate AWC, clay and sand content. Finally, a k-means cluster analysis was performed in the viticultural areas, using the selected variable maps (mean annual temperature, mean annual precipitation, mean annual soil temperature, elevation, clay, sand, coarse fragment content, available water capacity, rooting and redoximorphic mottles depths) to determine the NTU-based viticultural terroir units.

The map results obtained from digital terroir zoning performed at regional scales over some thousands of km<sup>2</sup> lead needing to better understand the issue of the relevant scales for defining and managing terroirs. It is important to both address the usefulness of both spatial and temporal resolutions that provide insights in terroir zoning according to the aim of the research or management practice. Studies need to examine the nature of within-field zones and the regional terroir units as the relationship between the number of map units and the surveyed area is log-linear (Fig. 6).

The underlying motivations for detailed assessments of the within-field spatial variation of yield/biomass/soil properties are related to the variable-rate application of inputs and selective harvesting at parcel level (Arnó et al., 2012). However considering the within-field scale only may be questionable in the case of highly-parcelled vineyards of less than 1 ha each such as in European countries with secular viticulture. More than ever, the issue of information synthesis and spatial scale is at the very heart of the spatial modelling of terroir (Vaudour, 2002). Considering a whole 90 ha-vineyard composed of 27 contiguous fields, Santesteban et al. (2013) showed that a per-field, within-field study would have resulted in missing the spatial trend due to slope. Therefore the usefulness of the multiple-field approach becomes one of a “whole-vineyard”

# SOILD

1, 827–906, 2014

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



or regional approach. Such issues similarly arises when attempting to predict the vine water status from a set of predawn water potential (PLWP) or MWP measurements, which are very time-consuming, require accurate methods to collect with a pressure chamber, and should be made at several fields managed by the vineyard operation.

5 Using PLWP (Acevedo-Opazo et al., 2008, 2010b) or MWP (Acevedo-Opazo et al., 2010a, 2013) measurements, Acevedo-Opazo et al. (2008, 2010a, b, 2013), elaborated a linear model applicable at the within-field level but requiring a time-consuming calibration set; in the case of PLWP, it corresponded to rainfed vineyards with high water stress, while in the case of MWP, it was aimed at scheduling irrigation. A similar approach using  $\delta^{13}\text{C}$  measurements as ancillary variables enabled extrapolation of stem water potentials for a rainfed vineyard with moderate water stress (Herrero-Langreo et al., 2013). These approaches may be transferable to a whole-vineyard scale (~ 29 km<sup>2</sup>), according to Baralon et al. (2012), who constructed an important measurement database (58 fields monitored in all) over the course of three consecutive vin-  
10 tages. However, several limitations hinder its practical application, namely those due to spatial sampling optimization and improvement of the temporal resolution of the model, using real-time monitoring sensors (such as sap flow sensors) (Herrero-Langreo et al., 2013). According to farmers, a model should be capable of predicting water stress at least two weeks before severe irreversible water stress damage occurs (J. C. Viaud,  
20 personal communication, 2014).

Arnó et al. (2012) stated that overall within-field variability of grape yield and quality raises important questions concerning whether site-specific crop management could be used in vineyards. Problems arise, in particular, when looking for causes of this variability, especially those related to the presence of soil carbonates which may lead to Mn deficiencies, with deleterious effects on grape colour. Another problem pointed out by Baveye and Laba (2014) related to N or P manuring is the frequent possibility  
25 that P or N deficiency in a management zone may be due to a higher leaching rate in the area, leading to an increased rather than decreased risk of groundwater contamination if more fertilizer is applied in this management zone.



### 3.4 Modelling and depicting climate at the region to vineyard scale

Another growth area in terroir zoning studies is the development of spatial climate data products. Historically, a region's climate and suitability for viticulture were assessed via climate station analyses, which seldom depict the spatial variation of climate in actual or prospective vineyard sites within wine-producing regions (Jones et al., 2010). As a result, reference vineyard networks were developed within regions to better capture the spatial climate characteristics (e.g., Jones and Davis, 2000). However, the low network density even in reference vineyard networks does not account for the range in meso-climates found within regions. To overcome these issues, spatial climate data products providing robust, validated, and more spatially appropriate climate data have been developed through the interpolation of existing long-term, quality-controlled data sources. Numerous techniques such as kriging and smoothing splines have been used to produce interpolated surfaces from hundreds to thousands of stations containing valuable meteorological inputs. The results are spatial climate products at daily or monthly time scales and at a range of spatial and temporal scales such as Daymet (Thornton et al., 1997), PRISM (parameter-elevation relationships on independent slopes model) (Daly et al., 2008), and WorldClim (Hijmans et al., 2005). Most of these approaches use elevation data (digital elevation data) and station climate data to calculate a climate-elevation regression for each grid cell, and stations entering the regression are assigned weights based primarily on the physiographic similarity of the station to the grid cell (Daly et al., 2008). These models attempt to account for location, elevation, coastal proximity, aspect, vertical differences in atmospheric layers, and orographic effects; although they can vary in the number and complexity of the factors involved. Validation procedures have shown that these products are generally robust at the scales for which they have been developed (Daly et al., 2008) and even at sub-grid resolutions, showing accuracy in regions characterized by sparse station data coverage, large elevation gradients, rain shadows, inversions, cold air drainage, and coastal effects.

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**Overview of terroir  
research**E. Vaudour et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Using these new gridded climate data sets, previous studies have examined viticulture region climate characteristics at various resolutions including 18 wine regions in Europe (1 KM; Jones et al., 2009), 50 DOs and sub-DOs in Portugal (1 KM; Jones and Alves 2012), 35 AOQS and AOCs in Greece (1 KM; Anderson et al., 2014), 135 AVAs in the western United States (400 m; Jones et al., 2010), 63 GIs in Australia (500 m; Hall and Jones, 2010), and 21 wine regions in New Zealand (500 m; Anderson et al., 2012) providing more holistic measures to help understand the range of climates within viticulture regions. Various climate parameters such as heat accumulation indices, frost timing, evapotranspiration, and dryness indices are often used along with mapping and spatial summaries over delimited winegrowing regions, ultimately helping to define the climate component of terroir over time and space. However, scale issues need to be further addressed for both terroir zoning and other applications. In vineyards, micro-variations in weather and climate often produce the greatest risk (e.g., frost or freeze zones, heat stress areas, etc.) and to truly address climate change we will need finer scales to assess the potential impacts (Quénol and Bonnardot, 2014).

#### 4 Terroir sustainability assessment and new preservation practices

The above-mentioned issues lead to considerations in an emerging and complex study area: the sustainability assessment of terroirs and the design of new preservation practices. Terroir sustainability started to be accounted for in the late 1990s, through some growers unions' initiatives in both California and Champagne, and also through the growing awareness of the soil contamination by copper due to the cumulated use of Bordeaux mixture (Vaudour, 2003). Mediterranean viticultural areas are amongst those most exposed and aware of the sustainability issue, because of the soil losses due to intense erosional processes (e.g. Le Bissonnais et al., 2007; Martínez-Casasnovas et al., 2009), which may lead to the destruction of a vineyard field (Fig. 7). Another issue is that of abandonment and land use change due to either urban pressure, aging of farmers or declines in profitability (Fig. 8).

---

**Overview of terroir  
research**


---

 E. Vaudour et al.
 

---

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Depletion of soil fertility in general, along with the concomitant problems of weeds, pests and diseases, is the fundamental root cause of low agricultural production at the global level (Tan et al., 2005). Even if grapevines are not nutrient demanding, it nevertheless requires an adequate supply, which may no longer be the case in terroirs where the soils are undergoing degradation processes, especially including soil losses (Le Bissonnais et al., 2007; Martínez-Casasnovas et al., 2009; Paroissien et al., 2010; Quiquerez et al., 2014; Chevigny et al., 2014), nutrient depletion (Ramos et al., 2006), compaction (Lagacherie et al., 2006), salinization/sodization (Clark et al., 2002; Crescimanno and Garofalo, 2006; Crescimanno et al., 2007; Costantini and Lorenzetti, 2013), pesticide runoff and deposition (Landry et al., 2005; Louchart and Voltz, 2007; Lacas et al., 2012; Daouk et al., 2013; Lefrancq et al., 2013), and copper contamination (Pieztrak and McPhail, 2004; Fernandez-Calvino et al., 2008; Chopin et al., 2008; Wightwick et al., 2008; Mirlean et al., 2007, 2009; Rusjan et al., 2007; El Hadri et al., 2012; El Azzi et al., 2013). In the Pénèdes viticultural region, Ramos et al. (2006) estimated that runoff processes exported significant amounts of nutrients, which represented as much as  $8 \text{ kg ha}^{-1} \text{ year}^{-1}$  of N and  $6.5 \text{ kg ha}^{-1} \text{ year}^{-1}$  of P. In Burgundy, using vine-stock unearthing–burying measurements (Brenot et al., 2008), Chevigny et al. (2014) estimated that the erosion rate had increased significantly over the last decade, and also that spatial distribution of erosion had changed and was now basically controlled by slope steepness and present-day vineyard management practices. Using the same method in Languedoc, Paroissien et al. (2010) estimated that the average soil loss reached  $10.5 \text{ t ha}^{-1} \text{ year}^{-1}$  and was much higher than the average erosion rates established around  $3 \text{ t ha}^{-1} \text{ year}^{-1}$  for cultivated soils. The high vulnerability of vineyard soil to erosion may be partly explained by the steep slopes where vineyards are established, but also by the generally low content in organic matter ( $< 2\%$ ) and the low microbial activity of soils which leads to a reduced aggregate stability that increases soil crusting and soil erosion (Le Bissonnais et al., 2007). According to Blavet et al. (2009), chemically weeded vineyards result in the highest runoff rates and soil losses, but the losses can be reduced when the prunings are left on

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the soil, when straw mulching is used, when rock fragments are left, and when grass intercrops are used. The corollary issue of runoff is that, in addition to soil erosion and soil nutrient loss, it also leads to fertilizer and pesticide residue loss to surface waters, depending on the aging of the applied pesticide (Louchart and Voltz, 2007).

In order to mitigate environmental damages and foster soil conservation and restoration, studies over the past decade have focused on the specific effect of one or several soil surface management techniques, such as cover cropping vs. chemical weeding or tillage and/or mulching, on soil structural stability (Goulet et al., 2004), pesticide leaching (Landry et al., 2005), soil nutrient and water management and vine growth and yield (Steenwerth and Belina, 2008, 2010; Ripoche et al., 2010, 2011), soil C dynamics (Steenwerth et al., 2010; Agnelli et al., 2014), and grape quality (Lee and Steenwerth, 2013). As recently described by Agnelli et al. (2014), a greater organic matter accumulation is fostered by the presence of the grass cover and the absence of tillage. In the Vosne-Romanée area in Burgundy, Landry et al. (2005) showed that glyphosate and its metabolite (AMPA) leached in greater amounts through a chemically treated bare calcosol than through a vegetated calcosol. In an upslope vineyard in the Beaujolais area, Lacas et al. (2012) demonstrated the usefulness of a grass buffer strip on a coarsely textured soil to limit the dispersal of diuron losses by runoff towards surface and sub-surface water. Jacobson et al. (2005) studied the leaching of the herbicide diuron jointly with that of copper, through vineyard soils contaminated with copper, and found no direct interaction between the metal and herbicide, but interpreted that Cu was possibly affecting microbial activity, resulting in slight increases in diuron persistence.

Despite the varied environmental problems that viticultural terroirs are facing currently that put their functioning at risk, the design of new preservation and mitigation practices has just begun to be addressed in the literature of the past decade. If some methods to globally assess the sustainability of agricultural systems have emerged (Bockstaller et al., 2009), they have seldom addressed that of viticultural systems (Abbona et al., 2007) and not only below the environmental point of view, but also for the attractiveness of the landscape, that in many viticultural areas conveys a remarkable

added value to the wine produced and the region (Tempesta et al., 2010; Costantini and Barbetti, 2008).

The global water crisis, particularly water scarcity (Hanjra and Qureshi, 2010), that threatens those viticultural areas under semi-arid or arid climates, particularly in the Mediterranean area (Iglesias et al., 2007; Plan Bleu, 2013), questions the relevancy of irrigating, in addition to the degradation effects that this practise may have on soil properties over a long-term period and its high remediation costs (Hajkowicz and Young, 2005). To address the issue of terroir sustainability in the years to come, one of the greatest challenges is the design of efficient soil restoration practices along with crop and/or intercrop management plans, taking into account the possible effects of climate change (see Sect. 2.3). This implies a complex multicriteria decision analysis, as attempted by Ripoche et al. (2010) in order to evaluate a range of intercrop management plans. The monitoring of soil quality as potentially obtainable from remote/proxy techniques is likely to identify when soil degradation is occurring in order to allow management intervention. Amongst the key biological indicators are soil organic carbon, potentially mineralisable nitrogen and microbial biomass (Riches et al., 2013), although further research is still needed to identify a suite of biological indicators for viticultural soils. Application of organic amendments is likely to improve soil quality but its effects are seldom studied, particularly over long-term experiments (Tatti et al., 2012). In rainfed Marchesi Antinori vineyards observed over two consecutive growing seasons, Baronti et al. (2014) suggested that biochar amendment could be used to improve soil water content, but other possible negative effects of changes in surface albedo or accumulation of polycyclic aromatic hydrocarbons still need to be studied. In contrast, considering a 3 year period in a Valais vineyard (Switzerland), Schmidt et al. (2014) observed only small and mostly-non-significant effects of either biochar or biochar-compost amendments. However, over the same duration in a vineyard in Jumilla (Spain), other organic inputs such as winery and distillery waste composts induced “an increase in the activity of the soil microorganisms and in the soil macro and micronutrient contents, as well as a slow release of inorganic N” (Bustamante et al.,

## SOILD

1, 827–906, 2014

### Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2011), in a soil characterized by a highly calcareous sandy-loam soil (Torriorthent). Coll et al. (2011) evaluated the long-term effect of organic viticulture on soil quality in commercial vineyards of Languedoc where plots which had been organically managed for 7, 11 and 17 years, comparing them to conventionally managed plots. The results emphasised that a transition period of 7–11 years, depending on the considered indicator, was needed to clearly separate conventional and organic farming. The overall benefits of organic farming were an increase of soil organic matter, potassium content, soil microbial biomass, plant-feeding, fungal feeding, nematode densities, while its drawbacks were increased soil compaction, decreased endogeic earthworm density (due to reduced soil porosity), both consequences of the increase of the traffic for tillage and phytosanitary treatments in organic management. The grapevines studied by these authors were not intercropped with grass, and knowledge is actually scarce about the joint long term environmental effects of the various possible sets of practices, such as cover/intercrop or not, mulching, mouldboard ploughing or/and tillage depth and frequency, type/quantity of applied organic amendments, and the use of chemical fertilizer or not.

Further studies should also address the possible reintroduction of agroforestry systems jointly with vineyards, as traditionally practised in Mediterranean regions, such as in Ancient Greece (Vaudour and Boulay, 2013) or in Italy and Provence till the XIXth century. Those systems are known to limit water consumption, fertilizers and pest diseases, particularly *Botrytis cinerea* bunch rot in Portugal (Altieri and Nicholls, 2002), and have positive impacts on phytoseid mite species, known for their ability to control mite pests, as shown in the last decade for vines co-planted with *Sorbus domestica* L. or *Pinus pinea* L. in Languedoc (Liguori et al., 2011).

## 5 Conclusions

Recent studies based on models of soil water and nitrogen dynamics and fingerprinting studies based either on metabolomics or the Sr isotopic ratio lead to a strengthening

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of the assumption that geographical origin does leave a footprint in wines and that both soil and substrate, in interaction with climate and cultural choices, influence the shaping of grapevine phenology and grape and wine quality. Differentiation and mapping of viticultural terroirs meant as homogeneous regions of grape/wine quality need comprehensive spatial modelling of soil, agronomical and climatic properties, including their changes through time. As such the development of a myriad of either remote or proxy sensing techniques and the corollary challenge of processing large quantities of data acquired at a very fine spatial resolution and/or at several spatial resolutions, scales, and organisational levels. These techniques in data collection and processing are needed to produce easy-to-update decision maps with associated uncertainties that allow users to make appropriate and timely management decisions. Another great challenge in the years to come is to address the issue of terroir sustainability and construct efficient strategies for assessing and applying them across numerous scales. These include the design of efficient soil restoration practices along with crop and/or intercrop management plans, and/or agroforestry viticultural systems, that take into account the possible effects of climate change. Therefore, terroirs are more and more likely to be addressed through the concept of ecosystem services, as viticultural agroecosystems, the services of which need to be constantly evaluated and rationalized.

## References

- Abbona, E. A., Sarandón, S. J., Marasas, M. E., and Astier, M.: Ecological sustainability evaluation of traditional management in different systems in Berisso, Argentina, *Agriculture, Ecosyst. Environ.*, 119, 335–345, 2007.
- Acevedo-Opazo, C., Tisseyre, B., Guillaume, S., and Ojeda, H.: The potential of high spatial resolution information to define within-vineyard zones related to vine water status, *Precis. Agric.*, 9, 285–302, 2008a.
- Acevedo-Opazo, C., Tisseyre, B., Ojeda, H., Ortega-Farias, S., and Guillaume, S.: Is it possible to assess the spatial variability of vine water status?, *J. Int. Sci. Vigne Vin*, 42, 203–219, 2008b.



## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acevedo-Opazo, C., Ortega-Farias, S., and Fuentes, S.: Effects of grapevine (*Vitis vinifera* L.) water status on water consumption, vegetative growth and grape quality: an irrigation scheduling application to achieve regulated deficit irrigation, *Agric. Water Manage.*, 97, 956–964, 2010a.

5 Acevedo-Opazo, C., Tisseyre, B., Ojeda, H., and Guillaume, S.: Spatial extrapolation of the vine (*Vitis vinifera* L.) water status: a first step towards a spatial prediction model, *Irrig. Sci.*, 28, 143–155, 2010b.

Acevedo-Opazo, C., Valdés-Gómez, H., Taylor, J. A., Avalo, A., Verdugo-Vásquez, N., Araya, M., Jara-Rojas, F., and Tisseyre, B.: Assessment of an empirical spatial prediction model of  
10 vine water status for irrigation management in a grapevine field, *Agric. Water Manage.*, 124, 58–68, 2013.

Ageorges, A., Fernandez, L., Vialet, S., Merdinoglu, D., Terrier, N., and Romieu, C.: Four specific isogenes of the anthocyanin metabolic pathway are systematically co-expressed with the red colour of grape berries, *Plant Science*, 170, 372–383, 2006.

15 Agnelli, A., Bol, R., Trumbore, S. E., Dixon, L., Cocco, S., and Corti, G.: Carbon and nitrogen in soil and vine roots in harrowed and grass-covered vineyards, *Agriculture, Ecosyst. Environ.*, 193, 70–82, 2014.

Anastasiadi, M., Zira, A., Magiatis, P., Haroutounian, S. A., Skaltsounis, A. L., and Mikros, E.: 1H NMR-based metabonomics for the classification of Greek wines according to variety, region, and vintage. Comparison with HPLC data, *J. Agr. Food Chem.*, 57, 11067–11074, 2009.

Anderson, J. D., Jones, G. V., Tait, A., Hall, A., and Trought, M. T. C.: Analysis of viticulture region climate structure and suitability in New Zealand, *International Journal of Vine and Wine Sciences*, 46, 149–165, 2012.

25 Anderson, J. D., Dimou, P., Jones, G. V., Kalivas, D., Koufos, G., Mavromatis, T., Koundouras, S. and Fyllas, N. M.: Harvest dates, climate, and viticultural region zoning in Greece. Proceedings of the 10th International Terroir Congress, 7–10 July, 2014, Tokaj, Hungary Vol. 2, 55–60, 2014.

Anderson, K., Findlay, C., Fuentes, S., and Tyerman, S.: Viticulture, wine and climate change, Commissioned report for the Garnaut Climate Change Review, accessible at: [www.garnautreview.org.au](http://www.garnautreview.org.au) (last access: 26 October 2014), 2008.

30 André, F., Van Leeuwen, Saussez, S., Van Durmen, R., Bogaert, P., Moghadas, D., de Rességuier, L., Delvaux, B., Vereecken, H., and Lambot, S.: High resolution imaging of a

**SOILD**

1, 827–906, 2014

**Overview of terroir  
research**

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



vineyard in south of France using ground-penetrating radar, electromagnetic induction and electrical resistivity tomography, *J. Appl. Geophys.*, 78, 113–122, 2012.

Andrenelli, M. C., Magini, S., Pellegrini, S., Perria, R., Vignozzi, N., and Costantini, E. A. C.: The use of the ARPt system to reduce the costs of soil survey for precision viticulture, *J. Appl. Geophys.*, 99, 24–34, 2013.

Arnó, J., Martínez-Casasnovas, J. A., Ribes-Dasi, M., and Rosell, J. R.: Review. Precision viticulture. Research topics, challenges and opportunities in site-specific vineyard management, *Spanish J. Agr. Res.*, 7, 779–790, 2009.

Arnó, J., Escolà, A., Vallès, J. M., Llorens, J., Sanz, R., Masip, J., Palacín, J., and Rosell-Polo, J. R.: Leaf area index estimation in vineyards using a ground-based LiDAR scanner, *Precis. Agric.*, 14, 290–306, 2012a.

Arnó, J., Rosell, J. R., Blanco, R., Ramos, M. C., and Martínez-Casasnovas, J. A.: Spatial variability in grape yield and quality influenced by crop and soil nutrition characteristics, *Precis. Agric.*, 13, 393–410, 2012b.

Bailly, J. S., Lagacherie, P., Millier, C., Puech, C., and Kosuth, P.: Agrarian landscapes linear features detection from LiDAR: application to artificial drainage networks, *Int. J. Remote Sens.*, 29, 3489–3508, 2008.

Bailly, J. S., Levavasseur, F., and Lagacherie, P.: A spatial stochastic algorithm to reconstruct artificial drainage networks from incomplete network delineations, *Int. J. Appl. Earth Obs.*, 13, 853–862, 2011.

Baluja, J., Diago, M. P., Balda, P., Zorer, R., Meggio, F., Morales, F., and Tardaguila, J.: Assessment of vineyard water status variability by thermal and multispectral imagery using an unmanned aerial vehicle (UAV), *Irrig. Sci.*, 30, 511–522, 2012a.

Baluja, J., Diago, M. P., Goovaerts, P., and Tardaguila, J.: Assessment of the spatial variability of anthocyanins in grapes using a fluorescence sensor: relationships with vine vigour and yield, *Precis. Agric.*, 13, 457–472, 2012b.

Baluja, J., Tardaguila, J., Ayestaran, B., and Diago, M. P.: Spatial variability of grape composition in a Tempranillo (*Vitis vinifera* L.) vineyard over a 3-year survey, *Precis. Agric.*, 14, 40–58, 2013.

Baralon, K., Payan, J. C., Salançon, E., and Tisseyre, B.: SPIDER: spatial extrapolation of the vine water status at the whole denomination scale from a reference site, *J. Int. Sci. Vigne Vin*, 46, 167–175, 2012.

---

**Overview of terroir  
research**

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Barata, A., Malfeito-Ferreira, M., and Loureiro, V.: The microbial ecology of wine grape berries, *Int. J. Food Microbiol.*, 153, 243–259, 2012.
- Barbeau, G., Asselin, C., and Morlat, R. : Estimation du potentiel viticole des terroirs en Val de Loire selon un indice de précocité du cycle de la vigne, *Bull. O.I.V.*, 805–806, 247–262, 1998 (in French).
- Barham, E.: Translating terroir: the global challenge of French AOC labelling, *J. Rural Stud.*, 19, 127–138, 2003.
- Baronti, S., Vaccari, F. P., Miglietta, F., Calzolari, C., Lugato, E., Orlandini, S., Pini, R., Zulian, C., and Genesio, L.: Impact of biochar application on plant water relations in *Vitis vinifera* (L.), *European J. Agronomy*, 53, 38–44, 2014.
- Ben Ghazlen, N., Cerovic, Z. G., Germain, C., Toutain, S., and Latouche, G.: Non-destructive optical monitoring of grape maturation by proximal sensing, *Sensors*, 10, 10040–10068, 2010.
- Bellvert, J., Zarco-Tejada, P. J., Girona, J., and Fereres, E.: Mapping crop water stress index in a ‘Pinot-noir’ vineyard: comparing ground measurements with thermal remote sensing imagery from an unmanned aerial vehicle, *Precis. Agric.*, 15, 361–376, 2014.
- Besky, S.: The labor of terroir and the terroir of labor: Geographical Indication and Darjeeling tea plantations, *Agric. Hum. Values*, 31, 83–96, 2014.
- Beucher, S. and Lantuéjoul, C.: Use of watersheds in contour detection, International Workshop on image processing, Real time edge and motion detection/estimation, Rennes, France, 17–21 September 1979, available at: <http://cmm.enscm.fr/~beucher/publi/watershed.pdf>, 1979.
- Bisson, L. F., Waterhouse, A. L., Ebeler, S. E., Walker, M. A. and Lapsley, J. T.: The present and future of the international wine industry, *Nature*, 418, 696–699, 2002.
- Blavet, D., De Noni, G., Le Bissonnais, Y., Léonard, M., Maillo, L., Laurent, J. Y., Asseline, J., Leprun, J. C., Arshad, M. A., and Roose, E.: Effect of land use and management on the early stages of soil water erosion in French Mediterranean vineyards, *Soil Till. Res.*, 106, 124–136, 2009.
- Bockstaller, C., Guichard, L., Keichinger, O., Girardin, P., Galan, M. B., and Gaillard, G.: Comparison of methods to assess the sustainability of agricultural systems. A review, *Agron. Sustain. Dev.*, 29, 223–235, 2009.

---

**Overview of terroir  
research**

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Bokulich, N. A., Thorngate, J. H., Richardson, P. M., and Mills, D. A.: Microbial biogeography of wine grapes is conditioned by cultivar, vintage and climate, *Proc. Natl. Aca. Sci.*, 111, E139–E148, 2013.
- Bonfante, A., Basile, A., Langella, G., Manna, P., and Terribile, F.: A physically oriented approach to analysis and mapping of terroirs, *Geoderma*, 167–168, 103–117, 2011.
- Bowen, S., and Zapata, A. V.: Geographical indications, terroir, and socioeconomic and ecological sustainability: the case of tequila, *J. Rural Stud.*, 25, 108–119, 2009.
- Bramley, R. G. V.: Understanding variability in winegrape production systems. 2. Within vineyard variation in quality overall several vintages, *Aust. J. Grape Wine Res.*, 11, 33–42, 2005.
- Bramley, R. G. V. and Hamilton, R. P.: Understanding variability in winegrape production systems. 1. Within vineyard variation in yield overall several vintages, *Aust. J. Grape Wine Res.*, 10, 32–45, 2004.
- Bramley, R. G. V. and Hamilton, R. P.: Terroir and precision viticulture: are they compatible?, *J. Int. Sci. Vigne Vin*, 41, 1–8, 2007.
- Bramley, R. G. V., Evans, K. J., Dunne, K. J., and Gobbett, D. L.: Spatial variation in response to “reduced input” spray programs for powdery mildew and botrytis identified through whole-of-block experimentation, *Aust. J. Grape Wine Res.*, 17, 341–350, 2011a.
- Bramley, R. G. V., Le Moigne, M., Evain, S., Ouzman, J., Florin, L., Fadaili, E. M., Hinze, C. J., and Cerovic, Z. G.: On-the-go sensing of grape berry anthocyanins during commercial harvest: development and prospects, *Aust. J. Grape Wine Res.*, 17, 316–326, 2011b.
- Bramley, R. G. V., Ouzman, J., and Boss, P. K.: Variation in vine vigour, grape yield and vineyard soils and topography as indicators of variation in the chemical composition of grapes, wine and wine sensory attributes, *Aust. J. Grape Wine Res.*, 17, 217–229, 2011c.
- Bramley, R. G. V., Ouzman, J., and Thornton, C.: Selective harvesting is a feasible and profitable strategy even when grape and wine production is geared towards large fermentation volumes, *Aust. J. Grape Wine Res.*, 17, 298–305, 2011d.
- Bramley, R. G. V., Trought, M. C. T., and Praat, J. P.: Vineyard variability in Malborough, New Zealand: characterizing variation in vineyard performance and options for the implementation of precision viticulture, *Aust. J. Grape Wine Res.*, 17, 72–78, 2011e.
- Brenot, J., Quiquerez, A., Petit, C., and Garcia, J. P.: Erosion rates and sediment budgets in vineyards at 1-m resolution based on stock unearthing (Burgundy, France), *Geomorphology*, 100, 345–355, 2008.

# SOILD

1, 827–906, 2014

## Overview of terroir research

E. Vaudour et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Brillante, L., Bois, B., Mathieu, O., Bichet, V., Michot, D., and Lévêque, J.: Monitoring soil volume wetness in heterogeneous soils by electrical resistivity. A field-based pedotransfer function, *J. Hydrol.*, 516, 56–66, 2014.

Brisson, N., Launay, M., Mary, B., Beaudoin, N.: Conceptual basis, formalisations and parameterization of the STICS crop model, Quae, Paris, 2009.

Bucelli, P. and Costantini, E. A. C.: Wine Grape and Vine zoning. Manual of methods for soil and land evaluation, edited by: Costantini, E. A. C., Science Publishers, Enfield (NH), (USA), 353-400, 2009.

Bulgarelli, D., Schlaeppli, K., Spaepen, S., Ver Loren van Themaat, E., and Schulze-Lefert, P.: Structure and functions of the bacterial microbiota of plants, *Annu. Rev. Plant Biol.*, 64, 807–838, 2013.

Bustamante, M. A., Said-Pullicino, D., Agulló, E., Andreu, E., Paredes, C., and Moral, R.: Application of winery and distillery waste composts to a Jumilla (SE Spain) vineyard: Effects on the characteristics of a calcareous sandy-loam soil, *Agr. Ecosyst. Environ.*, 140, 80–87, 2011.

Cabral-Chamorro, A.: Observaciones sobre la regulación y ordenación del mercado del vino en Jerez de la Frontera 1850–1935: antecedentes del consejo regulador de la denominación de origen Jerez-Xérès-Cherry, *Agricultura y Sociedad*, 44, 171–197, 1987 (in Spanish).

Carbonneau, A., Deloire, A., and Jaillard, B.: La vigne. Physiologie, terroir, culture. Dunod, Paris, 2007, (in French).

Carré, F. and McBratney, A. B.: Digital terroir mapping, *Geoderma*, 128, 340–353, 2005.

Carré, F., McBratney, A. B., Mayr, T., and Montanarella, L.: Digital soil assessments: beyond DSM, *Geoderma*, 142, 69–79, 2007.

Castellarin, S. D., Matthews, M. A., Di Gaspero, G., and Gambetta, G. A.: Water deficits accelerate ripening and induce changes in gene expression regulating flavonoid biosynthesis in grape berries, *Planta*, 227, 101–112, 2007.

Castrignanò, A., Costantini, E. A. C., Barbetti, R., and Sollitto, D.: Accounting for extensive topographic and pedologic secondary information to improve soil mapping, *Catena*, 77, 28–38, 2009.

Celette, F. and Gary, C.: Dynamics of water and nitrogen stress along the grapevine cycle as affected by cover cropping, *Europ. J. Agronomy*, 45, 142–152, 2013.

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Celette, F., Wéry, J., Chantelot, E., Celette, J., and Gary, C.: Belowground interactions in a vine (*Vitis vinifera* L.)-tall fescue (*Festuca arundinacea* Shreb.) intercropping system: water relations and growth, *Plant Soil*, 276, 205–217, 2005.

Celette, F., Gaudin, R., and Gary, C.: Spatial and temporal changes to the water regime of a Mediterranean vineyard due to the adoption of cover cropping, *Euro. J. Agronomy*, 29, 153–162, 2008.

Celette, F., Findeling, A., and Gary, C.: Competition for nitrogen in an unfertilized intercropping system: The case of an association of grapevine and grass cover in a Mediterranean climate, *Euro. J. Agronomy*, 30, 41–51, 2009.

Celette, F., Ripoche, A., and Gary, C.: WaLIS. A simple model to simulate water partitioning in a crop association: the example of intercropped vineyard, *Agr. Water Manage.*, 97, 1749–1759, 2010.

Champagnol, F.: *Éléments de physiologie de la vigne et de viticulture générale*, ISBN 978-2-950061-40-0, Saint-Gély-du-Fesc, 1984 (in French).

Chanussot, J., Bas, P., and Bombrun, L.: Airborne remote sensing of vineyards for the detection of dead vine trees, *Geoscience and Remote Sensing Symposium, 2005. IGARSS '05, Proceedings, 2005 IEEE International*, 5, 3090–3093, 2005.

Chevigny, E., Quiquerez, A., Petit, C., and Curmi, P.: Lithology, landscape structure and management practice changes: Key factors patterning vineyard soil erosion at metre-scale spatial resolution, *Catena*, 121, 354–364, 2014.

Choné, X., Lavigne-Cruège, V., Tominaga, T., Van Leeuwen, C., Castagnède, C., Saucier, C., and Dubourdieu, D.: Effect of vine nitrogen status on grape aromatic potential: flavor precursors (s-cysteine conjugates), glutathione and phenolic content in *Vitis Vinifera* L. cv. Sauvignon blanc grape juice, *Journal International des Sciences de la Vigne et du Vin*, 40, 1–6, 2006.

Chopin, E. I. B., Marin, B., Mkoungafoko, R., Rigaux, A., Hopgood, M.J., Delannoy, E., Cancès, B., and Laurain, M.: Factors affecting distribution and mobility of trace elements (Cu, Pb, Zn) in a perennial grapevine (*Vitis vinifera* L.) in the Champagne region of France, *Environ. Pollut.*, 156, 1092–1098, 2008.

Chuine, I., Yiou, P., Viovy, N., Seguin, B., Daux, V., and Le Roy Ladurie, E. : Grape ripening as a past climate indicator, Summer temperature variations are reconstructed from harvest dates since 1370, *Nature*, 432, 289–290, 2004.

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Clark, L. R., Fitzpatrick, R. W., Murray, R. S., and McCarthy, G.: Vineyard soil degradation following irrigation with saline groundwater for twenty years, 17th World Congress of Soil Science Bangkok, Thailand, 14–20, 2002.

Coll, P., Le Cadre, E., Blanchart, E., Hinsinger, P., and Villenave, C.: Organic viticulture and soil quality, a long-term study in Southern France, *Appl. Soil Ecol.*, 50, 37–44, 2011.

Compant, S., Mitter, B., Colli-Mull, J. G., Gangl, H., and Sessitsch, A.: Endophytes of grapevine flowers, berries, and seeds: identification of cultivable bacteria, comparison with other plant parts, and visualization of niches of colonization, *Microb. Ecol.*, 62, 188–197, 2011.

Conradie, W. J.: Seasonal Uptake of Nutrients by Chenin blanc in Sand Culture: I. Nitrogen, *S. Afr. J. Enol. Vitic.*, 1, 59–65, 1980.

Conradie, W. J.: Partitioning of nitrogen in grapevines during autumn and the utilisation of nitrogen reserves during the following growing season, *S. Afr. J. Enol. Vitic.*, 13, 45–51, 1992.

Corbane, C., Andrieux, P., Voltz, M., Chadoeuf, J., Albergel, J., Robbez-Masson, J. M., and Zante, P.: Assessing the variability of soil surface characteristics in row-cropped fields: the case of Mediterranean vineyards in Southern France, *Catena*, 72, 79–90, 2008a.

Corbane, C., Raclot, D., Jacob, F., Albergel, J., and Andrieux, P.: Remote sensing of soil surface characteristics from a multiscale classification approach, *Catena*, 75, 308–318, 2008b.

Corbane, C., Jacob, F., Raclot, D., Albergel, J., and Andrieux, P.: Multitemporal analysis of hydrological soil surface characteristics using aerial photos: A case study on a Mediterranean vineyard, *Int. J. Appl. Earth Obs. Geoinf.*, 18, 356–367, 2012.

Costantini E. A. C. and Barbetti, R.: Environmental and visual impact analysis of viticulture and olive tree cultivation in the Province of Siena (Italy), *Euro. J. Agronomy*, 28, 412–426, 2008.

Costantini, E. A. C. and Lorenzetti, M. : Soil degradation processes in the Italian agricultural and forest ecosystems, *Italian J. Agronomy* 8:e28, 233–243, 2013.

Costantini, E. A. C., Pellegrini, S., Vignozzi, N., and Barbetti, R.: Micromorphological characterization and monitoring of internal drainage in soils of vineyards and olive groves in Central Italy, *Geoderma*, 131, 388–403, 2006.

Costantini, E. A. C., Pellegrini, S., Bucelli, P., Storchi, P., Vignozzi, N., Barbetti, R., and Campagnolo, S.: Relevance of the Lin's and Host hydroopedological models to predict grape yield and wine quality, *Hydrol. Earth Syst. Sci.*, 13, 1635–1648, doi:10.5194/hess-13-1635-2009, 2009.



---

**Overview of terroir  
research**

E. Vaudour et al.

---

 Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Costantini, E. A. C., and Bucelli, P.: Soil and terroir, in: Soil security for ecosystem management, edited by: Kapur, S. and Erşahin, S., SpringerBriefs in Environment, Security, Development and Peace, 8, 97–133, doi:10.1007/978-3-319-00699-4\_6, 2014.

Costantini, E. A. C., Bucelli, P., and Priori, S.: Quaternary landscape history determines the soil functional characters of terroir, *Quaternary Int.*, 265, 63–73, 2012.

Costantini, E. A. C., Agnelli, A., Bucelli, P., Ciambotti, A., Dell'Oro, V., Natarelli, L., Pellegrini, S., Perria, R., Priori, S., Storch, P., Tsolakis, C., and Vignozzi, N.: Unexpected relationships between  $\delta^{13}\text{C}$  and grape performance in organic farming, *J. Int. Sci. Vigne Vin*, 47, 269–285, 2013.

Costantini, E. A. C., Agnelli, A., Fabiani, A., Gagnarli, E., Mocali, S., Priori, S., Simoni, S., and Valboa, G.: Short term recovery of soil biological functions in a new vineyard cultivated in organic farming, *EGU, Geophys. Res. Abstr.*, Vienna, Austria, vol. 16, 3691-1, 2014.

Crescimanno, G. and Garofalo, P.: Management of irrigation with saline water in cracking clay soils, *Soil Sci. Soc. Am. J.*, 70, 1774–1787, 2006.

Crescimanno, G., De Santis, A., and Provenzano, G.: Soil structure and bypass flow processes in a Vertisol under sprinkler and drip irrigation, *Geoderma*, 138, 110–118, 2007.

Cunha, C. R., Peres, E., Morais, R., Oliveira, A. A., Matos, S. G., Fernandes, M. G., Ferreira, P. J. S. G., and Reis, M. J. C. S.: The use of mobile devices with multi-tag technologies for an overall contextualized vineyard management, *Comput. Electron. Agric.*, 73, 154–164, 2010.

Da Costa, J. P., Michelet, F., Germain, C., Laviolle, O., and Grenier, G.: Delineation of vine parcels by segmentation of high resolution remote sensed images, *Precis. Agric.*, 8, 95–110, 2007.

Daly, C., Halbleib, M., Smith, J. I., Gibson, W. P., Doggett, M. K., Taylor, G. H., Curtis, J., and Pasteris, P. A.: Physiographically-sensitive mapping of temperature and precipitation across the conterminous United States, *Int. J. Climatol.*, 28, 2031–2064, 2008.

Daniel, R.: The metagenomics of soil, *Nat. Rev. Microbiol.*, 3, 470–478, 2005.

Davies, C. and Robinson, S. P.: Sugar accumulation in grape berries. Cloning of two putative vacuolar invertase cDNAs and their expression in grapevine tissues, *Plant Physiol.*, 111, 275–283, 1996.

De Assis Silva, S., Marçal de Queiroz, D., De Assis De Carvalho Pinto, F., and Terra Santos, N.: Characterization and delimitation of the terroir coffee in plantations in the municipal district of Araponga, Minas Gerais, Brazil, *Revista Ciência Agronômica*, 45, 18–26, 2014.

---

**Overview of terroir  
research**

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Delenne, C., Durrieu, S., Rabatel, G., Deshayes, M., Bailly, J. S., Lelong, C., Couteron, P.: Textural approaches for vineyard detection and characterization using very high spatial resolution remote sensing data, *Int. J. Remote Sens.*, 29, 1153–1167, 2008.

Delenne, C., Durrieu, S., Rabatel, G., and Deshayes, M.: From pixel to vine parcel: a complete methodology for vineyard delineation and characterization using remote-sensing data, *Comput. Electron. Agric.*, 70, 78–83, 2010.

Deloire, A., Vaudour, E., Carey, V., Bonnardot, V., and Van Leeuwen, C.: Grapevine responses to terroir: a global approach, *J. Int. Sci. Vigne Vin*, 39, 149–162, 2005.

De Souza, C. R., Maroco, J. P., Dos Santos, T. P., Rodrigues, M. L., Lopes, C. M., Pereira, J. S., and Chaves, M. M.: Partial rootzone drying: regulation of stomatal aperture and carbon assimilation in field-grown grapevines, *Funct. Plant Biol.*, 30, 653–662, 2003.

De Souza, C. R., Maroco, J. P., Dos Santos, T. P., Rodrigues, M. L., Lopes, C. M., Pereira, J. S., and Chaves, M. M.: Impact of deficit irrigation on water use efficiency and carbon isotope composition ( $\delta^{13}\text{C}$ ) of field-grown grapevines under Mediterranean climate, *J. Experiment. Botany*, 56, 2163–2172, 2005.

Dion, R.: *Le paysage et la vigne: essai de géographie historique*, Payot, Paris, 1990 (in French).

Di Paola-Naranjo, R. D., Baroni, M. V., Podio, N. S., Rubinstein, H. R., Fabani, M. P., Badini, R. G., Inga, M., Ostera, H. A., Cagnoni, M., Gallegos, E., Gautier, E., Peral-García, P., Hoogewerff, J., and Wunderlin, D. A.: Fingerprints for main varieties of Argentinean wines: terroir differentiation by inorganic, organic, and stable isotopic analyses coupled to chemometrics, *J. Agr. Food Chem.*, 59, 7854–7865, 2011.

Dobrowski, S. Z., Ustin, S. L., and Wolpert, J. A.: Grapevine dormant pruning weight prediction using remotely sensed data, *Aust. J. Grape Wine Res.*, 9, 177–182, 2003.

Dobrowski, S. Z., Ustin, S. L., and Wolpert, J. A.: Remote estimation of vine canopy density in vertically shoot-positioned vineyards: determining optimal vegetation indices, *Aust. J. Grape Wine Res.*, 8, 117–125, 2008.

Donat, M. G., Alexander, L. V., Yang, H., Durre, I., Vose, R., and Caesar, J.: Global land-based datasets for monitoring climatic extremes, *Bull. Amer. Meteor. Soc.*, 94, 997–1006, 2013.

Doolittle, J. A. and Brevik, E. C.: The use of electromagnetic induction techniques in soil studies, *Geoderma*, 223–225, 33–45, 2014.

Doré, T., Makowski, M., Malézieux, E., Munier-Jolain, N., Tchamitchian, M., and Titonell, P.: Facing up to the paradigm of ecological intensification in agronomy: revisiting methods, concepts and knowledge, *Euro. J. Agronomy*, 34, 197–210, 2011.

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Düring, H.: Evidence of osmotic adjustment to drought in grapevines (*Vitis vinifera* L.), *Vitis*, 23, 1–10, 1984.
- El Azzi, D., Viers, J., Guiesse, M., Probst, A., Aubert, D., Caparros, J., Charles, F., Guizien, K., and Probst, J. L.: Origin and fate of copper in a small Mediterranean vineyard catchment, new insights from combined chemical extraction and  $\delta^{65}\text{Cu}$  isotopic composition, *Sci. Total Environ.*, 463–464, 91–101, 2013.
- El Hadri, H., Chéry, P., Jalabert, S., Lee, A., Potin-Gautier, M., and Lespes, G.: Assessment of diffuse contamination of agricultural soil by copper in Aquitaine region by using French national databases, *Sci. Total Environ.*, 441, 239–247, 2012.
- Fernández-Calviño, D., Garrido-Rodríguez, B., López-Periago, J. E., Paradelo, M., and Arias-Estévez, M.: Spatial distribution of copper fractions in a vineyard soil, *Land Degrad. Develop.*, 24, 556–563, 2013.
- Fiorillo, E., Crisci, A., De Filippis, T., Di Gennaro, S. F., Di Blasi, S., Matese, A., Primicerio, J., Vaccari, F. P., and Genesio, L.: Airborne high-resolution images for grape classification: changes in correlation between technological and late maturity in a Sangiovese vineyard in Central Italy, *Aust. J. Grape Wine Res.*, 18, 80–90, 2012.
- Follain, S., Ciampalini, R., Crabit, A., Coulouma, G., and Garnier, F.: Effects of redistribution processes on rock fragment variability within a vineyard topsoil in Mediterranean France, *Geomorphology*, 175–176, 45–53, 2012.
- Fraga, H., Malheiro, A. C., Moutinho-Pereira, J., and Santos, J. A.: An overview of climate change impacts on European viticulture, *Food and Energy Security*, 1, 94–110, 2012.
- Fregoni, M.: *Viticultura di qualità*. Phytoline, Affi (VR), 2005 (in Italian).
- Fulton, A., Schwankl, L., Lynn, K., Lampinen, B., Edstrom, J., and Prichard, T.: Using EM and VERIS technology to assess land suitability for orchard and vineyard development, *Irrig. Sci.*, 29, 497–512, 2011.
- Galet, P.: *Cépages et vignobles de France. Tome II : L'ampélographie française*, ISBN-13: 978-2902771042, 1993 (in French).
- Galleguillos, M., Jacob, F., Prévot, L., French, A., and Lagacherie, P.: Comparison of two temperature differencing methods to estimate daily evapotranspiration over a Mediterranean vineyard watershed from ASTER data, *Remote Sens. Environ.*, 115, 1326–1340, 2011a.
- Galleguillos, M., Jacob, F., Prévot, L., Lagacherie, P., and Liang, S.: Mapping daily evaporation over a Mediterranean vineyard watershed, *IEEE Geosci. Remote Sens. Lett.*, 8, 168–172, 2011b.

---

**Overview of terroir  
research**


---

 E. Vaudour et al.
 

---

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- García de Cortázar-Atauri, I., Brisson, N., and Gaudillère, J. P.: Performance of several models for predicting budburst date of grapevine (*Vitis vinifera* L.), *Int. J. Biometeorol.*, 53, 317–326, 2009.
- 5 Gilbert, J. A., Van der Lelie, D., and Zarraonaindia, I.: Microbial terroir for wine grapes, *Proc. Natl. Aca. Sci.*, 111, 5–6, 2014.
- Gil-Pérez, B., Zarco-Tejada, P. J., Correa-Guimaraes, A., Relea-Gangas, E., Navas-Gracia, L. M., Hernández-Navarro, S., Sanz-Requena, J. F., Berjón, A., and Martín-Gil, J.: Remote sensing detection of nutrient uptake in vineyards using narrow-band hyperspectral imagery, *Vitis*, 49, 167–173, 2010.
- 10 Girard, M. C.: Recherche d'une modélisation en vue d'une représentation spatiale de la couverture pédologique. Application à une région des plateaux jurassiques de Bourgogne, Thèse de doctorat d'Etat, Institut National Agronomique Paris-Grignon, Sols, 13, 1983 (in French).
- Gómez-Alonso, R. and García-Romero, E.: Effect of irrigation and variety on oxygen ( $\delta^{18}\text{O}$ ) and carbon ( $\delta^{13}\text{C}$ ) stable isotope composition of grapes cultivated in a warm climate, *Aust. J. Grape Wine Res.*, 16, 283–289, 2010.
- 15 Gong, P., Mahler, S. A., Biging, G. S., and Newburn, D. A.: Vineyard identification in a oak woodland landscape with airborne digital camera imagery, *Int. J. Remote Sens.*, 24, 1303–1315, 2003.
- Guilpart, N., Métay, A., and Gary, C.: Grapevine bud fertility and number of berries per bunch are determined by water and nitrogen stress around flowering in the previous year, *Euro. J. Agronomy*, 54, 9–20, 2014.
- 20 Guix-Hébrard, N., Voltz, M., Trambouze, W., Garnier, F., Gaudillère, J. P., and Lagacherie, P.: Influence of watertable depths on the variation of grapevine water status at the landscape scale, *Eur. J. Agron.*, 27, 187–196, 2007.
- 25 Hajkowicz, S. and Young, M.: Costing yield loss from acidity, sodicity, and dryland salinity to Australian agriculture, *Land Degrad. Dev.*, 16, 417–433, 2005.
- Hall, A. and Jones, G. V.: Spatial analysis of climate in winegrape growing regions in Australia, *Aust. J. Grape Wine Res.*, 16, 389–404, 2010.
- Hall, A. and Wilson, M.A.: Object-based analysis of grapevine canopy relationships with winegrape composition and yield in two contrasting vineyards using multitemporal high spatial resolution optical remote sensing, *Int. J. Remote Sens.*, 34, 1772–1797, 2013.
- 30 Hall, A., Lamb, D. W., Holzapfel, B. P., and Louis, J. P.: Optical remote sensing applications in viticulture – a review, *Aust. J. Grape Wine Res.*, 8, 36–47, 2002.

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Hall, A., Louis, J., and Lamb, D.: Characterizing and mapping vineyard canopy using high-spatial-resolution aerial multispectral images, *Comput. Geosci.*, 29, 813–822, 2003.
- Hall, A., Louis, J. P., and Lamb, D. W.: Low-resolution remotely sensed images of winegrape vineyards map spatial variability in planimetric canopy instead of leaf area index, *Aust. J. Grape Wine Res.*, 14, 9–17, 2008.
- Hall, A., Lamb, D. W., Holzapfel, B. P., and Louis, J. P.: Within-season temporal variation in correlations between vineyard canopy and winegrape composition and yield, *Precis. Agric.*, 12, 103–107, 2011.
- Hanjra, M. A. and Qureshi, E.: Global water crisis and future food security in an era of climate change, *Food Policy*, 35, 365–377, 2010.
- Hannah, L., Roehrdanz, P. R., Ikegami, R., Shepard, A. V., Shaw, M. R., Tabor, G., Zhi, L., Marquet, P. A., and Hijmans, R. J.: Climate change, wine, and conservation, *Proc. Natl. Acad. Sci. USA*, 110, 6907–6912, 2013.
- Haralick, R. M., Shanmugam, K., and Dinstein, I.: Textural features for image classification, *IEEE Trans. Syst. Man Cybernet.*, 3, 610–621, 1973.
- Hardie, W. and Considine, J.: Response of grapes to water deficit stress in particular stages of development, *Am. J. Enol. Vitic.*, 27, 55–61, 1976.
- Hartemink, A. E. and Minasny, B.: Towards digital soil morphometrics, *Geoderma*, 230–231, 305–317, 2014.
- Herrera-Núñez, J. C., Ramazotti, S., Stagnari, F., and Pisante, M.: A multivariate clustering approach for characterization of the Montepulciano d'Abruzzo Colline Teramane area, *Am. J. Enol. Vitic.*, 62, 239–244, 2011.
- Herrero-Langreo, A., Tisseyre, B., Goutouly, J. P., Scholasch, T., and Van Leeuwen, C.: Mapping grapevine (*Vitis vinifera* L.) water status during the season using carbon isotope ratio ( $\delta^{13}\text{C}$ ) as ancillary data, *Am. J. Enol. Vitic.*, 64, 307–315, 2013.
- Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., and Jarvis, A.: Very high resolution interpolated climate surfaces for global land areas, *Int. J. Climatol.*, 25, 1965–1978, 2005.
- Hole, F. D.: An approach to landscape analysis with emphasis on soil, *Geoderma*, 21, 1–23, 1978.
- Houdaille, F. and Sémichon, L.: Influence de l'état physique du calcaire considéré comme cause déterminante de la chlorose, in: *Revue de viticulture*, Paris, rue Gay-Lussac, 8, 243–326, 1894 (in French).

---

**Overview of terroir  
research**

E. Vaudour et al.

---

 Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Huglin, P.: Nouveau mode d'évaluation des possibilités héliothermiques d'un milieu viticole, *Comptes-Rendus de l'Académie d'Agriculture de France*, 64, 1117–1126, 1978 (in French).
- Huglin, P. and Schneider, C.: *Biologie et écologie de la vigne*, Lavoisier Tec&Doc, Paris, 2nd Edn., 1998 (in French).
- 5 Hugues, P., McBratney, A. B., Malone, B. P., and Minasny, B.: Development of terrons for the Lower Hunter Valley wine-growing region, in: *Digital soil assessments and beyond*, edited by: Minasny, B., Malone, B. P., and McBratney, A. B., Taylor & Francis, London, 31–36, 2012.
- Iglesias, A., Garrote, L., Flores, F., and Moneo, M.: Challenges to manage the risk of water scarcity and climate change in the Mediterranean, *Water Resour. Manage.*, 21, 775–788, 10 2007.
- IPCC: *Climate Change 2013, The Physical Science Basis, Contribution of Working Group I to the fifth Assessment Report of the intergovernmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Lignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, United Kingdom and 15 New York, NY, USA, 2013.
- Jacobson, A. R., Dousset, S., Andreux, F., and Baveye, P. C.: Electron microprobe and synchrotron X-ray fluorescence mapping of the heterogeneous distribution of copper in high-copper vineyard soils, *Environ. Sci. Technol.*, 14, 6343–6349, 2007.
- Jacobsen, R.: *American Terroir: Savoring the Flavors of Our Woods, Waters, and Fields*, Bloomsbury USA, New York, 2010.
- 20 Johnson, L. F.: Temporal stability of an NDVI-LAI relationship in a Napa Valley vineyard, *Aust. J. Grape Wine Res.*, 9, 96–101, 2003.
- Johnson, L. F., Roczen, D. E., Youkhana, S. K., Nemani, R. R., and Bosch, D. F.: Mapping vineyard leaf area with multispectral satellite imagery, *Comput. Electron. Agric.*, 38, 33–44, 25 2003.
- Jones, G. V. and Alves, F.: Spatial Analysis of Climate in Winegrape Growing Regions in Portugal. *Proceedings of the 9th International Terroir Congress*, 25–29 June, 2012, Burgundy and Champagne, France, Volume 1, 3, 1–4, 2012.
- Jones, G. V. and Davis, R. E.: Climate influences on grapevine phenology, grape composition, and wine production and quality for Bordeaux, France, *Am. J. Enol. Vitic.*, 51, 249–261, 30 2000.
- Jones, G. V., White, M. A., Cooper, O. R., and Storchmann, K.: Climate change and global wine quality, *Climatic Change*, 73, 319–343, 2005.

---

**Overview of terroir  
research**

 E. Vaudour et al.
 

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Jones, G. V., Moriondo, M., Bois, B., Hall, A., and Duff, A.: Analysis of the spatial climate structure in viticulture regions worldwide, *Bull. OIV*, 944–946, 507–518, 2009.
- Jones, G. V., Duff, A. A., Hall, A., and Myers, J. W.: Spatial analysis of climate in winegrape growing regions in the Western United States, *Am. J. Enol. Vitic.*, 61, 313–326, 2010.
- 5 Jones, G. V., Reid, R., and Vilks, A.: Climate, grapes, and wine: structure and suitability in a variable and changing climate, 109–133, in: *The Geography of Wine: Regions, Terroir, and Techniques*, edited by: Dougherty, P., Springer Press, 255 pp., 2012.
- Komárek, M., Száková, J., Rohošková, M., Javorská, H., Chrastrný, V., and Balík, J.: Copper contamination of vineyard soils from small wine producers: a case study from the Czech Republic, *Geoderma*, 147, 16–22, 2008.
- 10 Lagacherie, P., Coulouma, G., Ariagno, P., Virat, P., Boizard, H., and Richard, G.: Spatial variability of soil compaction over a vineyard region in relation with soils and cultivation operations, *Geoderma*, 134, 207–216, 2010.
- Lagacherie, P., Robbez-Masson, J. M., Nguyen-The, N., and Barthès, J. P.: Mapping of reference area representativity using a mathematical soilscape distance, *Geoderma*, 101, 105–118, 2001.
- 15 Lagacherie, P., Bailly, J. S., Monestiez, P., and Gomez, C.: Using scattered hyperspectral imagery data to map the soil properties of a region, *Euro. J. Soil Sci.*, 63, 110–119, 2012.
- Lai, H. Y., Juang, K. W., and Chen, B. C.: Copper concentrations in grapevines and vineyard soils in central Taiwan, *Soil Sci. Plant Nutr.*, 56, 601–606, 2013.
- 20 Lamb, D. W., Weedon, M. M., and Bramley, R. G. V.: Using remote sensing to predict grape phenolics and colour at harvest in a Cabernet-Sauvignon vineyard: timing observations against vine phenology and optimising image resolution, *Aust. J. Grape Wine Res.*, 10, 46–54, 2004.
- Lamb, D. W., Mitchell, A., and Hyde, G.: Vineyard trellising with steel posts distorts data from EM soil surveys, *Aust. J. Grape Wine Res.*, 11, 24–32, 2005.
- 25 Lanjeri, S., Segarra, D., and Melia, J.: Interannual vineyard crop variability in the Castilla-La Mancha region during the period 1991–1996 with Landsat Thematic Mapper images, *Int. J. Remote Sens.*, 25, 2441–2457, 2004.
- Laville, P.: Le terroir, un concept indispensable à l’élaboration et à la protection des appellations d’origine comme à la gestion des vignobles: le cas de la France, *Bull. OIV*, 709–710, 217–241, 1990.
- 30



## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Le Bissonnais, Y., Blavet, D., De Noni, G., Laurent, J.-Y., Asseline, J., and Chenu, C.: Erodibility of Mediterranean vineyard soils: relevant aggregate stability methods and significant soil variables, *Euro. J. Soil Sci.*, 58, 188–195, 2007.
- Lebon, E., Dumas, V., Pieri, P., and Schultz, H. R.: Modelling the seasonal dynamics of the soil water balance of vineyards, *Funct. Plant Biol.*, 30, 699–710, 2003.
- Lefrancq, M., Imfeld, G., Payraudeau, and Millet, M.: Kresoxim methyl deposition, drift and runoff in a vineyard catchment, *Sci. Total Environ.*, 442, 503–508, 2013.
- Liguori, M., Tixier, M. S., Fabio Hernandez A., Douin, M., and Kreiter, S.: Agroforestry management and phytoseiid communities in vineyards in the South of France, *Exp. Appl. Acarol.*, 55, 167–181, 2011.
- Llorens, J., Gil, E., Llop, J., and Queraltó, M: Georeferenced LiDAR 3D vine plantation map generation, *Sensors*, 11, 6237–6256, doi:10.3390/s110606237, 2011.
- López-Rituerto, E., Savorani, F., Avenzoza, A., Busto, J. H., Peregrina, J. M., and Balling Engelsen, S.: Investigations of la Rioja terroir for wine production using  $^1\text{H}$  NMR metabolomics, *J. Agr. Food Chem.*, 60, 3452–3461, 2012.
- Loveys, B. R. and Düring, H.: Diurnal changes in water relations and abscisic acid in field-grown *Vitis vinifera* cultivars. II. Abscisic acid changes under semi-arid conditions, *New Phytol.*, 97, 37–47, 1984.
- Loveys, B. R., Kriedemann, P. E., and Törökfalvy, E.: Is abscisic acid involved in stomatal response to carbon dioxide?, *Plant Sci. Lett.*, 1, 335–338, 1973.
- Mackie, K. A., Müller, T., and Kandeler, E.: Remediation of copper in vineyards. A mini-review, *Environ. Pollut.*, 167, 16–26, 2012.
- Mackie, K. A., Müller, T., Zikeli, S., and Kandeler, E.: Long-term copper application in an organic vineyard modifies spatial distribution of micro-organisms, *Soil Biol. Biochem.*, 65, 245–253, 2013.
- Malone, B. P., Hugues, P., McBratney, A. B., and Minasny, B.: A model for the identification of terrons in the Lower Hunter Valley, Australia, *Geoderma Reg.*, 1, 31–47, 2014.
- Marchionni, S., Braschi, E., Tommasini, S., Bollati, A., Cifelli, F., Mulinacci, N., Mattei, M., and Conticelli, S.: High precision  $^{87}\text{Sr}/^{86}\text{Sr}$  analyses in wines and their use as geological fingerprint for tracing geographic provenance, *J. Agric. Food. Chem.*, 61, 6822–6831, 2013.
- Marre, A.: Existe-t-il des terroirs en Champagne?, *Revue Géographique de l'Est*, 1–2, 17–30, 2004 (in French).

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Martín, P., Zarco-Tejada, P. J., González, M. R., and Berjón, A.: Using hyperspectral remote sensing to map grape quality in “Tempranillo” vineyards affected by iron deficiency chlorosis, *Vitis*, 46, 7–14, 2007.
- Martínez-Casasnovas, J. A., Agelet-Fernandez, J., Arnó, J., and Ramos, M. C.: Analysis of vineyard differential management zones and relation to vine development, grape maturity and quality, *Span. J. Agric. Res.*, 10, 326–337, 2012.
- Martínez-Casasnovas, J. A., Ramos, M. C., and García-Hernández, D.: Effects of land-use changes in vegetation cover and sidewall erosion in a gully head of the Penedès region (northeast Spain), *Earth Surf. Proc. Landf.*, 34, 1927–1937, 2009.
- Martins, G., Lauga, B., Miot-Sertier, C., Mercier, A., Lonvaud, A., Soulas, M. L., Soulas, G., and Masneuf-Pomarède, I.: Characterization of epiphytic bacterial communities from grapes, leaves, bark and soil of grapevine plants grown, and their relations, *Plos one*, 8, e73013, 1–9, 2013.
- Matheron, G.: *Traité de géostatistique appliquée*, Tome 1, Editions Technip, Paris, France, 1962.
- Matheron, G.: *Les variables régionalisées et leur estimation: une application de la théorie des fonctions aléatoires aux sciences de la nature*, Masson, Paris, 1965.
- Mathews, A. J. and Jensen, J. L. R.: An airborne LiDAR-based methodology for vineyard parcel detection and delineation, *Int. J. Remote Sens.*, 33, 5251–5267, 2012.
- Mathews, A. J. and Jensen, J. L. R.: Visualizing and quantifying vineyard canopy LAI using an Unmanned Aerial Vehicle (UAV) collected high density structure from motion point cloud, *Remote Sens.*, 5, 2164–2183, 2013.
- Meggio, F., Zarco-Tejada, P. J., Núñez, L. C., Sepulcre-Cantó, G., González, M. R., and Martín, P.: Grape quality assessment in vineyards affected by iron deficiency chlorosis using narrow-band physiological remote sensing indices, *Remote Sens. Environ.*, 114, 1968–1986, 2010.
- Mercurio, M., Grilli, E., Odierna, P., Morra, V., Prohaska, T., Coppola, E., Grifa, C., Buondonno, A., and Langella, A.: A “Geo-Pedo-Fingerprint” (GPF) as a tracer to detect univocal parent material-to-wine production chain in high quality vineyard districts, Campi Flegrei (Southern Italy), *Geoderma*, 230–231, 64–78, 2014.
- Mirlean, N., Roisenberg, A., and Chies, J. O.: Metal contamination of vineyard soils in wet subtropics (Southern Brazil), *Bull. Environ. Contam. Toxicol.*, 82, 373–377, 2007.

---

**Overview of terroir  
research**


---

 E. Vaudour et al.
 

---

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Mirlean, N., Baisch, P., and Medeanic, S.: Copper bioavailability and fractionation in copper-contaminated sandy soils in the wet subtropics (Southern Brazil), *Bull. Environ. Contam. Toxicol.*, 82, 373–377, 2009.

Mocali, S. and Benedetti, A.: Exploring research frontiers in microbiology: the challenge of metagenomics in soil microbiology, *Res. Microbiol.*, 161, 497–505, 2010.

Mocali, S., Fabiani, A., Kuramae, E., de Hollander, M., Kowalchuk, G. A., Vignozzi, N., Valboa, G., and Costantini, E.: Genetic and functional diversity of soil microbial communities associated to grapevine plants and wine quality, EGU Abstract no 1187-1, Vienna, Austria, 2013.

Morari, F., Castrignanò, A., and Pagliarin, C.: Application of multivariate geostatistics in delineating management zones within a gravelly vineyard using geo-electrical sensors, *Comput. Electron. Agric.*, 68, 97–107, 2009.

Moreira, N., Guedes de Pinho, P., Santos, C., and Vasconcelos, I.: Relationship between nitrogen content in grapes and volatiles, namely heavy sulphur compounds, in wines, *Food Chem.*, 26, 1599–1607, 2011.

Moriondo, M., Jones, G. V., Bois, B., Dibari, C., Ferrise, R., Trombi, G., and Bindi, M.: Projected shifts of wine regions, *Climatic Change*, 119, 825–839, 2013.

Morlat, R. (Coord.): *Terroirs viticoles: étude et valorisation*. Editions Oenoplurimedia, Chaintré, France, 2001 (in French).

Morlat, R. and Jacquet, A.: The soil effects on the grapevine root system in several vineyards of the Rhone valley (France), *Vitis*, 32, 35–42, 1993.

Nannipieri, P., Ascher, J., Ceccherini, M., Landi, L., Pietramellara, G., and Renella, G.: Microbial diversity and soil functions, *Euro. J. Soil Sci.*, 54, 655–670, 2003.

Nearing, M. A., Pruski, F. F., and O’Neal, M. R.: Expected climate change impacts on soil erosion rates: A review, *J. Soil Water Cons.*, 59, 43–50, 2004.

Nendel, C. and Kersebaum, K. C.: A simple model approach to simulate nitrogen dynamics in vineyard soils, *Ecol. Model.*, 177, 1–15, 2004.

Ojeda, H., Deloire, A., and Carbonneau, A.: Influence of water deficits on grape berry growth, *Vitis*, 40, 141–145, 2001.

Ojeda, H., Andary, C., Kraeva, E., Carbonneau, A., and Deloire, A.: Influence of pre- and post-veraison water deficit on synthesis and concentration of skin phenolic compounds during berry growth of *Vitis vinifera* cv. Shiraz, *Am. J. Enol. Vitic.*, 53, 261–267, 2002.

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Oliver, M. A. and Webster, R.: A tutorial guide to geostatistics: computing and modelling variograms and kriging, *Catena*, 113, 56–69, 2014.
- Palliotti, A., Silvestroni, O., Petoumenou, D., Vignaroli, S., and Berrios, J. G.: Evaluation of low-energy demand adaptative mechanisms in Sangiovese grapevine during drought, *J. Int. Sci. Vigne Vin*, 42, 41–47, 2008.
- Paoli, J. N., Tisseyre, B., Strauss, O., and McBratney, A. B.: A technical opportunity index based on the fuzzy footprint of a machine for site-specific management: an application to viticulture, *Precis. Agric.*, 11, 379–396, 2010.
- Paranychanakis, N. V. and Angelakis, A. N.: The effect of water stress and rootstock on the development of leaf injuries in grapevines irrigated with saline effluent, *Agr. Water Manage.*, 95, 375–382, 2008.
- Paroissien, J. B., Lagacherie, P., and Le Bissonnais, Y.: A regional-scale study of multi-decennial erosion of vineyard fields using vine-stock unearthing–burying measurements, *Catena*, 82, 159–168, 2010.
- Pedroso, M., Taylor, J., Tisseyre, B., Charnomordic, B., and Guillaume, S.: A segmentation algorithm for the delineation of agricultural management zones, *Comput. Electron. Agric.*, 70, 199–208, 2010.
- Pellegrino, A., Lebon, E., Voltz, M., and Wery, J.: Relationships between plant and soil water status in vine (*Vitis vinifera* L.), *Plant Soil*, 266, 129–142, 2004.
- Pellegrino, A., Gozé, E., Lebon, E., and Wery, J.: A model-based diagnosis tool to evaluate the water stress experienced by grapevine in field sites, *Euro. J. Agronomy*, 25, 49–59, 2006.
- Pereira, G. E., Gaudillère, J. P., Van Leeuwen, C., Hilbert, G., Maucourt, M., Deborde, C., Moing, A., and Rolin, D.:  $^{1}H$  NMR metabolite fingerprints of grape berry: comparison of vintage and soil effects in Bordeaux grapevine growing areas, *Anal. Chim. Acta*, 563, 346–352, 2006.
- Pietrzak, U. and McPhail, D. C.: Copper accumulation, distribution and fractionation in vineyard soils of Victoria, Australia, *Geoderma*, 122, 151–166, 2004.
- Pinto, C., Pinho, D., Sousa, S., Pinheiro, M., Egas, C., and Gomes, A. C.: Unraveling the diversity of the grapevine microbiome, *PLOS-One*, 9, e85622, 2014.
- Plan Bleu: Mediterranean strategy for sustainable development follow-up, Main indicators, 2013 update, 2013.

**SOILD**

1, 827–906, 2014

**Overview of terroir research**

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Primicerio, J., Di Gennaro, S. F., Fiorillo, E., Genesio, L., Lugato, E., Matese, A., and Vaccari, F. P.: A flexible unmanned aerial vehicle for precision agriculture, *Precis. Agric.*, 13, 517–523, 2012.
- Priori, S., Fantappiè, M., Magini, S., and Costantini, E. A. C.: Using the ARP-03 for high-resolution mapping of calcic horizons, *Int. Agrophys.*, 27, 313–321, 2013a.
- Priori, S., Martini, E., Andrenelli, M. C., Magini, S., Agnelli, A. E., Bucelli, P., Biagi, M., Pellegrini, S., and Costantini, E. A. C.: Improving wine quality through harvest zoning and combined use of remote and proximal sensing, *Soil Sci. Soc. Am. J.*, 77, 1338–1348, 2013b.
- Priori, S., Barbetti, R., L'Abate, G., Bucelli, P., Storchi, P., and Costantini, E. A. C.: Natural terroir units, Siena province, Tuscany, *Journal Maps*, 10, 466–477, 2014.
- Puletti, N., Perria, R., and Storchi, P.: Unsupervised classification of very high remotely sensed images for grapevine rows detection, *Eur. J. Remote Sens.*, 47, 45–54, 2014.
- Quénot, H. and Bonnardot, V.: A multi-scale climatic analysis of viticultural terroirs in the context of climate change: the TERADCLIM Project, *J. Int. Sci. Vigne Vin*, Special Issue Laccave, 25–34, 2014.
- Quiquerez, A., Chevigny, E., Allemand, P., Curmi, P., Petit, C., and Grandjean, P.: Assessing the impact of soil surface characteristics on vineyard erosion from very high spatial resolution aerial images (Côte de Beaune, Burgundy, France), *Catena*, 116, 163–172, 2014.
- Rabatel, G., Debain, C., Delenne, C., and Deshayes, M.: “Bacchus” methodological approach for vineyard inventory and management, edited by: Montesinos Aranda, S. and Quintanilla, A., European Commission, DG Research, RTD action: energy, environment and sustainable development, 67–87, 2006.
- Rabatel, G., Delenne, C., and Deshayes, M.: A non-supervised approach using Gabor filters for vine-plot detection in aerial images, *Comput. Electron. Agric.*, 62, 159–168, 2008.
- Ramos, M. C. and Martínez-Casasnovas, J. A.: Nutrient losses by runoff in vineyards of the Mediterranean Alt Penedès region (NE Spain), *Agriculture, Ecosyst. Environ.*, 113, 356–363, 2006.
- Ramos, M. C. and Martínez-Casasnovas, J. A.: Effects of precipitation patterns and temperature trends on soil water available for vineyards in a Mediterranean climate area, *Agr. Water Manage.*, 97, 1495–1505, 2010.
- Ramos, M. C. and Martínez-Casasnovas, J. A.: Soil water variability and its influence on transpirable soil water fraction with two grape varieties under different rainfall regimes, *Agr. Ecosyst. Environ.*, 185, 253–262, 2014.

---

**Overview of terroir  
research**

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Ravaz, L.: Influence de la surproduction sur la végétation de la vigne, *Annales de l'Ecole Nationale d'Agriculture de Montpellier*, 6, 1–79, 1906 (in French).
- Renouf, V., Claisse, O., and Lonvaud-Funel, A.: Understanding the microbial ecosystem on the grape berry surface through numeration and identification of yeast and bacteria, *Aust. J. Grape Wine Res.*, 11, 316–327, 2005.
- Richards, A. L., Hutson, J. L., and McCarthy, G.: Monitoring and modelling transient rootzone salinity in drip irrigated viticulture, in: *Proceedings of the 13th AWITC*, edited by: Blair, R., Williams, P., and Pretorius, S., 212–217, 2008.
- Riches, D., Porter, I. J., Oliver, D. P., Bramley, R. G. V., Rawnsley, B., Edwards, J., and White, R. E.: Review: soil biological properties as indicators of soil quality in Australian viticulture, *Aust. J. Grape Wine Res.*, 19, 311–323, 2013.
- Riou, C., Valancogne, C., and Pieri, P.: Un modèle simple d'interception du rayonnement solaire par la vigne. Vérification expérimentale, *Agronomie*, 9, 441–450, 1989 (in French).
- Riou, C., Pieri, P., and Le Clech, B.: Consommation d'eau de la vigne en conditions hydriques non limitantes. Formulation simplifiée de la transpiration, *Vitis*, 33, 109–115, 1994 (in French).
- Ripoche, A., Celette, F., Cinna, J. P., and Gary, C.: Design of intercrop management plans to fulfil production and environmental objectives in vineyards, *Euro. J. Agronomy*, 32, 30–39, 2010.
- Roby, G. and Matthews, M. A.: Relative proportions of seed, skin and flesh, in ripe berries from Cabernet Sauvignon grapevines grown in a vineyard either well irrigated or under water deficit, *Aust. J. Grape Wine Res.*, 10, 74–82, 2004.
- Rodriguez-Lovelle, B. and Gaudillère, J. P.: Carbon and nitrogen partitioning in either fruiting or non-fruiting grapevines: effects of nitrogen limitation before and after veraison, *Aust. J. Grape Wine Res.*, 8, 86–94, 2002.
- Rodríguez-Pérez, J. R., Riaño, D., Carlisle, E., Ustin, S., and Smart, D. R.: Evaluation of hyperspectral reflectance indexes to detect grapevine water status in vineyards, *Am. J. Enol. Vitic.*, 58, 302–317, 2007.
- Rodríguez-Pérez, J. R., Plant, R. E., Lambert, J. J., and Smart, D. R.: Using apparent soil electrical conductivity (ECa) to characterize vineyard soils of high clay content, *Precis. Agric.*, 12, 775–794, 2011.

---

**Overview of terroir  
research**

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Rossi, R., Pollice, A., Diago, M. P., Oliveira, M., Millan, B., Bitella, G., Amato, M., and Tardaguila, J.: Using an automatic resistivity profiler soil sensor on-the-go in precision viticulture, *Sensors*, 13, 1121–1136, 2013.
- Roudier, P., Tisseyre, B., Poilvé, H., and Roger, J. M.: A technical opportunity index adapted to zone-specific management, *Precis. Agric.*, 12, 130–145, 2011.
- Rusjan, D., Strlič, M., Pucko, D., and Korošec-Koruza, Z.: Copper accumulation regarding the soil characteristics in sub-Mediterranean vineyards of Slovenia, *Geoderma*, 141, 111–118, 2007.
- Samouëlian, A., Cousin, I., Tabbagh, A., Bruand, A., and Richard, G.: Electrical resistivity survey in soil science: a review, *Soil Till. Res.*, 83, 173–193, 2005.
- Santesteban, L. G. and Royo, J. B.: Water status, leaf area and fruit load influence on berry weight and sugar accumulation of cv. “Tempranillo” under semiarid conditions, *Scientia Horticulturae*, 127, 494–499, 2006.
- Santesteban, L. G., Guillaume, S., Royo, J. B., and Tisseyre, B.: Are precision agriculture tools and methods relevant at the whole-vineyard scale?, *Precis. Agric.*, 14, 2–17, 2013.
- Santos, J. A., Malheiro, A. C., Pinto, J. G., and Jones, G. V.: Macroclimate and viticultural zoning in Europe: observed trends and atmospheric forcing, *Clim. Res.*, 51, 89–103, 2012.
- Santos, J. A., Grätsch, S. D., Karremann, M. K., Jones, G. V., and Pinto, J. G.: Ensemble projections for wine production in the Douro Valley of Portugal, *Climatic Change*, 117, 211–225, 2013.
- Scacco, A., Verzera, A., Lanza, C. M., Sparacio, A., Genna, G., Raimondi, S., Tripodi, G., and Dima, G.: Influence of soil salinity on sensory characteristics and volatile aroma compounds of Nero d’Avola wine, *Am. J. Enol. Vitic.*, 61, 498–505, 2010.
- Scarlett, N. J., Bramley, R. G. V., and Siebert, T. E.: Within-vineyard variation in the “pepper” compound rotundone is spatially structured and related to variation in the land underlying the vineyard, *Aust. J. Grape Wine Res.*, 20, 214–222, 2014.
- Schmidt, H. P., Kammann, C., Niggli, C., Evangelou, M. W. H., Mackie, K. A., and Abiven, S.: Biochar and biochar-compost as soil amendments to a vineyard soil: influences on plant growth, nutrient uptake, plant health and grape quality, *Agriculture, Ecosyst. Environ.*, 191, 117–123, 2014.
- Scholander, P. F., Bradstreet, E. D., Hemmingsen, E. A., and Hammel, H. T.: Sap pressure in vascular plants. Negative hydrostatic pressure can be measured in plants, *Science*, 148, 339–346, 1965.



## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Seguin, G.: Les sols viticoles du Haut-Médoc. Influence sur l'alimentation en eau de la vigne et sur la maturation du raisin, PhD thesis, University of Bordeaux II, 1970 (in French).
- Seguin, G.: Influence des terroirs viticoles sur la constitution et la qualité des vendanges, Bull. O.I.V., 56, 3–18, 1983 (in French).
- 5 Seguin, G.: “Terroirs” and pedology of wine growing, *Experientia*, 42, 861–873, 1986.
- Setati, M. E., Jacobson, D., Andong, U. C., and Bauer, F.: The vineyard yeast microbiome, a mixed model microbial map, *PLoS one*, 7, e52609, 2012.
- Shani, U. and Ben-Gal, A.: Long-term response of grapevines to salinity: osmotic effects and ion toxicity, *Am. J. Enol. Vitic.*, 56, 148–154, 2005.
- 10 Son, H. S., Hwang, G. S., Kim, K. M., Ahn, H. J., Park, W. M., Van Den Berg, F., Hong, Y. S., and Lee, C. H.: Metabolomic studies on geographical grapes and their wines using  $^1\text{H}$  NMR analysis coupled with multivariate statistics, *J. Agr. Food Chem.*, 57, 1481–1490, 2009.
- Stamatiadis, S., Taskos, D., Tsadila, E., Christofides, C., Tsadilas, C., and Schepers, J. S.: Comparison of passive and active canopy sensors for the estimation of vine biomass production, *Precis. Agric.*, 11, 306–315, 2010.
- 15 Steenwerth, K. L. and Belina, K. M.: Vineyard weed management practices influence nitrate leaching and nitrous oxide emissions, *Agr. Ecosyst. Environ.*, 138, 127–131, 2010.
- Steenwerth, K. L., Pierce, D. L., Carlisle, E. A., Spencer, R. G. M., and Smart, D. R.: A vineyard agroecosystem. Disturbance and precipitation affect soil respiration under Mediterranean conditions, *Soil Sci. Soc. Am. J.*, 74, 231–239, 2010.
- 20 Stevens, R. M., Harvey, G., Norton, S., and Frahn, W.: Over-canopy saline sprinkler irrigation of grapevines during different growth stages, *Agr. Water Manage.*, 101, 62–70, 2011.
- Stevens, R. M., Pech, J. M., Gibberd, M. R., Walker, R. R., and Nicholas, P. R.: Reduced irrigation and rootstock effects on vegetative growth, yield and its components, and leaf physiological responses of Shiraz, *Aust. J. Grape Wine Res.*, 16, 413–425, 2010.
- 25 Stoll, M., Loveys, B., and Dry, P.: Hormonal changes induced by partial rootzone drying of irrigated grapevine, *J. Experiment. Botany*, 51, 350, 1627–1634, 2000.
- Tagarakis, A., Liakos, V., Fountas, S., Koundouras, S., and Gemtos, T. A.: Management zones delineation using fuzzy clustering techniques in grapevines, *Precis. Agric.*, 140, 18–39, 2013.
- 30 Tan, Z. X., Lal, R., and Wiebe, K. D.: Global soil nutrient depletion and yield reduction, *J. Sustain. Agr.*, 26, 123–146, 2005.

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Tarr, P. T., Dreyer, M. L., Athanas, M., Shahgholi, M., Saarloos, K., and Second, T. P.: A metabolomics based approach for understanding the influence of terroir in *Vitis Vinifera* L., *Metabolomics*, 9, S170–S177, 2013.
- 5 Tartaguila, J., Baluja, J., Arpon, L., and Balda, P.: Variations of soil properties affect the vegetative growth and yield components of “Tempranillo” grapevines, *Precis. Agric.*, 12, 762–773, 2011.
- Tatti, E., Goyer, C., Zebarth, B. J., Burton, D. L., Giovanetti, L., and Viti, C.: Short-term effects of mineral and organic fertilizer on denitrifiers, nitrous oxide emissions and denitrification in long-term amended vineyard soils, *Soil Sci. Soc. Am. J.*, 77, 113–122, 2012.
- 10 Taylor, J. A., Acevedo-Opazo, C., Ojeda, H., and Tisseyre, B.: Identification and significance of sources of spatial variation in grapevine water status, *Aust. J. Grape Wine Res.*, 16, 218–226, 2010.
- Taylor, J. A., Coulouma, G., Lagacherie, P., and Tisseyre, B.: Mapping soil units within a vineyard using statistics associated with high-resolution apparent soil electrical conductivity data and factorial discriminant analysis, *Geoderma*, 153, 278–284, 2009.
- 15 Taylor, J. A., Jacob, F., Galleguillos, M., Prévot, L., Guix, N., and Lagacherie, P.: The utility of remotely-sensed vegetative and terrain covariates at different spatial resolutions in modelling soil and watertable depth (for digital soil mapping), *Geoderma*, 193–194, 83–93, 2013.
- Teixeira, A., Eiras-Dias, J., Castellarin, S. D., and Gerós, H.: Berry phenolics of grapevine under challenging environments, *Int. J. Mol. Sci.*, 14, 18711–18739, 2013.
- 20 Tempesta, T., Giancristofaro, R. A., Corain, L., Salmaso, L., Tomasi, D., and Boatto, V.: The importance of landscape in wine quality perception: An integrated approach using choice-based conjoint analysis and combination-based permutation tests, *Food Qual. Pref.*, 21, 827–836, 2010.
- 25 Thornton, P. E., Running, S. W., and White, M. A.: Generating surfaces of daily meteorology variables over large regions of complex terrain, *J. Hydrol.*, 190, 214–251, 1997.
- Tisseyre, B. and McBratney, A. B.: A technical opportunity index based on mathematical morphology for site-specific management: an application to viticulture, *Precis. Agric.*, 9, 101–103, 2008.
- 30 Tisseyre, B., Mazzone, C., and Fonta, H.: Within-field temporal stability of some parameters in viticulture: potential toward a site specific management, *J. Int. Sci. Vigne Vin*, 42, 27–39, 2008.

---

**Overview of terroir  
research**

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Tomasi, D., Gaiotti, F., and Jones, G. V.: The power of the terroir: the case study of Prosecco wine, Springer, Basel, 2013.
- Tonietto, J. and Carbonneau, A.: A multicriteria climatic classification system for grape-growing regions worldwide, *Agr. Forest Meteorol.*, 124, 81–87, 2004.
- 5 Torrea, T., Varela, C., Ugliano, M., Ancin-Azpilicueta, C., Francis, I. L., and Henschke, P. A.: Comparison of inorganic and organic nitrogen supplementation of grape juice. Effect on volatile composition and aroma profile of a Chardonnay wine fermented with *Saccharomyces cerevisiae* yeast, *Food Chem.*, 127, 1072–1083, 2011.
- Trambouze, W. and Voltz, M.: Measurement and modelling of the transpiration of a Mediterranean vineyard, *Agr. Forest Meteorol.*, 107, 153–166, 2001.
- 10 Trought, M. C. T. and Bramley, R. G. V.: Vineyard variability in Marlborough, New Zealand: characterizing spatial and temporal changes in fruit composition and juice quality in the vineyard, *Aust. J. Grape Wine Res.*, 17, 79–89, 2010.
- Trubek, A. B.: *Taste of Place: A Cultural Journey into Terroir*, University of California Press, Berkeley and Los Angeles, California, 2008.
- 15 Tucker, C. J.: Red and photographic infrared linear combinations for monitoring vegetation, *Remote Sens. Environ.*, 8, 127–150, 1979.
- Unamunzaga, O., Besga, G., Castellón, A., Usón, M. A., Chéry, P., Gallejones, P., and Aizpurua, A.: Spatial and vertical analysis of soil properties in a Mediterranean vineyard soil, *Soil Use Manage.*, 1–12, doi:10.1111/sum.12110, online first, 2014.
- 20 Unwin, T.: *Vine and the wine: an historical geography of viticulture and the wine trade*, Routledge, London, 1991.
- Urdanoz, V. and Aragüés, R.: Three-year field response of drip-irrigated grapevine (*Vitis vinifera* L., cv. Tempranillo) to soil salinity, *Plant Soil*, 324, 219–230, 2009.
- 25 Urretavizcaya, I., Santesteban, L. G., Tisseyre, B., Guillaume, S., Miranda, C., and Royo, J. B.: Oenological significance of vineyard management zones delineated using early grape sampling, *Precis. Agric.*, 9, 101–103, 2013.
- Van Leeuwen, C. and Seguin, G.: The concept of terroir in viticulture, *J. Wine Res.*, 17, 1–10, 2006.
- 30 Van Leeuwen, C. F. P., Chonei, X., Tregouat, O., Koundouras, S., and Dubourdieu, D.: The influence of climate, soil and cultivar on terroir, *Am. J. Enol. Vitic.*, 55, 207–217, 2004.
- Vaudour, E.: The quality of grapes and wine in relation to geography: notions of terroir at various scales, *J. Wine Res.*, 13, 117–141, 2002.

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vaudour, E.: Les terroirs viticoles, Dunod, Paris, 2003 (in French).

Vaudour, E.: Remote sensing of Red Mediterranean soils: a case study in the viticultural Southern Rhone Valley (France) using SPOT satellite imagery, *Geocarto Int.*, 23, 197–216, 2008.

Vaudour, E. and Boulay, T.: Towards a spatial analysis of ancient viticultural areas: the case study of Amos (Turkey), *Food History*, 11, 157–176, 2013.

Vaudour, E., Girard, M. C., Brémond, L. M., and Lurton, L.: Caractérisation spatiale et constitution des raisins en AOC Côtes-du-Rhône méridionales (Bassin de Nyons-Valréas), *J. Int. Sci. Vigne Vin*, 32, 169–182, 1998 (in French).

Vaudour, E., Morlat, R., Van Leeuwen, C., and Dolédec, A. F.: Terroirs viticoles et sols, in: *Sols et Environnement*, Dunod, Paris, 105–126, (in French), 2005.

Vaudour, E., Carey, V. A., and Gilliot, J. M.: Digital zoning of South African viticultural terroirs using bootstrapped decision trees on morphometric data and multitemporal SPOT images, *Remote Sens. Environ.*, 114, 2940–2950, 2010.

Vaudour, E., Carey, V. A., and Gilliot, J. M.: Multidate remote sensing approaches for digital zoning of terroirs at regional scales: case studies revisited and perspectives images, EGU Abstract no 11533, Vienna, Austria, 2014a.

Vaudour, E., Gilliot, J. M., Bel, L., Bréchet, L., Hadjar, D., Hamiache, J., Lemonnier, Y.: Uncertainty of soil reflectance retrieval from SPOT and RapidEye multispectral satellite images using a per-pixel bootstrapped empirical line atmospheric correction over an agricultural region, *Int. J. Appl. Earth Obs. Geoinf.*, 26, 217–234, 2014b.

Verger, A., Vigneau, N., Chéron, C., Gilliot, J. M., Comar, A., and Baret, F.: Green area index from an unmanned aerial system over wheat and rapeseed crops, *Remote Sens. Environ.*, 152, 654–664, 2014.

Vilanova, M., Siebert, T. E., Varela, C., Pretorius, I. S., and Henschke, P. A.: Effect of ammonium nitrogen supplementation of grape juice on wine volatiles and non-volatiles composition of the aromatic grape variety Albariño, *Food Chem.*, 133, 124–131, 2012.

Vincent, L. and Soille, P.: Watershed in digital spaces – An efficient algorithm based on immersion simulations, *IEEE T. Pattern Anal.*, 13, 583–598, 1991.

Vogel, T. M., Simonet, P., Jansson, J. K., Hirsch, P. R., Tiedje, J. M., Van Elsas, J. D., Bailey, M. J., Nalin, R., and Philippot, L.: TerraGenome: a consortium for the sequencing of a soil metagenome, *Nature Reviews Microbiology*, 7(4), 252, 2009.

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Warner, T. A. and Steinmaus, K.: Spatial classification of orchards and vineyards with high spatial resolution panchromatic imagery, *Photogramm. Eng. Remote Sens.*, 71, 179–187, 2005.
- Wassenaar, T., Robbez-Masson, J. M., Andrieux, P., and Baret, F.: Vineyard identification and description of spatial crop structure by per-field frequency analysis, *Int. J. Remote Sens.*, 23, 3311–3325, 2002.
- Wassenaar, T., Andrieux, P., Baret, F., and Robbez-Masson, J. M.: Soil surface infiltration capacity classification based on the bi-directional reflectance distribution function sampled by aerial photographs. The case of vineyards in a Mediterranean area, *Catena*, 62, 94–110, 2005.
- Werban, U., Bartholomeus, H., Dietrich, P., Grandjean, G., and Zacharias, S.: Digital soil mapping: approaches to integrate sensing techniques to the prediction of key soil properties, *Vadose Zone J.*, Special section: Digital soil mapping, 1–4, doi:10.2136/vzj2013.10.0178, 2013.
- West, J. B., Ehrlinger, J. R., and Cerling, T. E.: Geography and Vintage Predicted by a Novel GIS Model of Wine  $\delta^{18}\text{O}$ , *J. Agric. Food Chem.*, 55, 7075–7083, 2007.
- Wezel, A., Casagrande, M., Celette, F., Vian, J. F., Ferrer, A., and Peigné, J.: Agroecological practices for sustainable agriculture. A review, *Agron. Sustain. Dev.*, 34, 1–20, 2014.
- White, M. A., Diffenbaugh, N. S., Jones, G. V., Pal, J. S., and Giorgi, F.: Extreme heat reduces and shifts United States premium wine production in the 21st century, *Proc. Natl. Acad. Sci. USA*, 103, 11217–11222, 2006.
- White, R., Balachandra, L., Edis, R., and Chen D.: The soil component of terroir, *J. Int. Sci. Vigne Vin*, 41, 9–18, 2007.
- White, R. E.: *Soils for fine wines*. Oxford University Press, New York, 2003.
- Wightwick, A. M., Mollah, M. R., Partington, D. L., and Allinson, G.: Copper fungicide residues in Australian vineyard soils, *J. Agric. Food Chem.*, 56, 2457–2464, 2008.
- Winkler, A. J.: *General Viticulture*, 4th Edn., University of California Press, Berkeley and Los Angeles, 1962.
- Zarco-Tejada, P. J., Berjón, A., López-Lozano, R., Miller, J. R., Martín, P., Cachorro, V., González, M. R., and de Frutos, A.: Assessing vineyard condition with hyperspectral indices: leaf and canopy reflectance simulation on, a row-structured discontinuous canopy, *Remote Sens. Environ.*, 99, 271–287, 2005.

Zarco-Tejada, P. J., Guillén-Climent, M. L., Hernández-Clemente, R., Catalina, A., González, M. R., and Martín, P.: Estimating leaf carotenoid content in vineyards using high resolution hyperspectral imagery acquired from an unmanned aerial vehicle (UAV), *Agric. Forest. Meteorol.*, 171–172, 281–294, 2013.

# SOILD

1, 827–906, 2014

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** Target variables of the zoning studies carried out over the 2002–2013 period (see Fig. 2).

viticultural	<i>N</i>	environmental	<i>N</i>	oenological	<i>N</i>
canopy characteristics	20	soil properties	14	grape composition	24
yield	17	soil units	6	wine composition	3
biomass	15	soil surface condition	3	wine sensory attributes	3
vine water status	12	temperatures	3		
vineyard identification	11	erosion	3		
management zones	8	climatic zones	2		
terroir units	6	land use changes	2		
pest/weed management	6	artificial drainage network	2		
evapotranspiration	4	landslide	1		
vine rows	4				
vineyard characteristics	3				

*N*: number of papers







# SOILD

1, 827–906, 2014

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Figure 2.** Increase of the number of papers ( $N$ ) dealing with terroir zoning from 2002 to March 2014. Source: Web of Science (v.5.14, 2014).



**Figure 3.** Plot of the number of map units vs. the log of the study area in studies published from 2002 to 2014 (see Fig. 2).


**Overview of terroir research**

E. Vaudour et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	







**Figure 5.** Terroir concept adapted to remote sensing. Adapted from Vaudour (2003).

**Overview of terroir research**


E. Vaudour et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



**Overview of terroir  
research**

E. Vaudour et al.



**Figure 6.** Relationship between the number of map units vs. the study area (log-transformed variables). 2002-early 2014 period (see Fig. 2).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





# SOILD

1, 827–906, 2014

## Overview of terroir research

E. Vaudour et al.



**Figure 7.** Former vineyard field destroyed by erosion in the Southern Rhone Valley, with dead vines still visible (Photograph by E. Vaudour).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Figure 8.** Abandoned vineyard field colonized by native Mediterranean plant species in the Languedoc region (France), with some vines still visible (Photograph by E. Vaudour).

# SOILD

1, 827–906, 2014

## Overview of terroir research

E. Vaudour et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

