

The effects of four irrigation regimes on vineyard vigour using proximal multi-spectral active sensors

Terrón J.M.¹, Blanco J.², Moral F.J.², Mancha L.A.¹, Uriarte D.¹, Marques da Silva J.R.³

¹ Centro de Investigaciones Científicas y Tecnológicas de Extremadura (CICYTEX) - Instituto de Investigaciones Agrarias Finca “La Orden-Valdesequera”, Gobierno de Extremadura, Autovía A-5 p.k. 372, 06187, Guadajira (Badajoz). e-mail: jose.terron@gobex.es

² Universidad de Extremadura, Escuela de Ingenierías Industriales, Departamento de Expresión Gráfica. Avda. de Elvas, s/n, 06071, Badajoz.

³ Universidade de Évora, Instituto de Ciências Agrárias e Ambientais Mediterrânicas (ICAAM), Escola de Ciências e Tecnologia, Apartado 94, 7002-554, Évora, Portugal. Applied Management and Space Centre for Interdisciplinary Development and Research on Environment (DREAMS), Lisboa, Portugal. Centro de Inovação em Tecnologias de Informação (CITI). Évora, Portugal.

Abstract

Precision agriculture is a useful tool to assess plant growth and development in vineyards. The present study was focused in the spatial and temporal analysis of vegetation growth variability analysis; considering four irrigation treatments with four replicates. The research was carried out in a vineyard located in the southwest of Spain during 2012 and 2013 growing seasons. Two multispectral sensors mounted on an All-Terrain Vehicle (ATV) were used in the different growing seasons/stages in order to calculate the vineyard Normalized Difference Vegetation Index (NDVI). Soil apparent Electrical Conductivity (ECa) was also measured up to 0.8 m soil depth using a geophysical sensor. All measured data were analysed by means of Principal Component Analysis (PCA). The spatial and temporal NDVI and ECa variations showed relevant differences between irrigation treatments and climatological conditions.

Keywords: Precision agriculture, Vineyard, apparent Electrical Conductivity, NDVI, Irrigation.

1 Introduction

Terroir is a French concept that says - “*there are unique aspects of a place that shape the quality of grapes and wine*”. Those aspects that impact on grapes and wine quality are usually associated with topography, soil, climate, plant management and plant genetics (Vaudour, 2002). According to several authors, the study of the plant vegetative vigour is an essential parameter to successfully manage yield and grapes/wines quality because of the fact that plant growth integrates climate, soil, topography, available water and other plant controlling factors (Smart, 1985; Carbonneau, 1995; Cortell et al., 2005; Deloire et al., 2005). Consequently, the appropriate management of soil and the consideration of the main climatic variables are key factors to obtain good yields and, finally, quality wines. Vineyard canopy management such as pruning systems, shoot orientation, shoot thinning or leaf removal, has the capacity to modify climate factors around the plant and therefore, modifying grape and wine quality (Dry, 2000).

34 Water management in vineyards and their responses have been studied since last decades in a high range of
35 environments and vineyard varieties due to the irrigation implications in yield and final product quality (Smart and
36 Coombe, 1983; Bravdo and Hepner, 1986; Mullins et al., 1992; Williams and Araujo, 2002; Intrigliolo and Castel, 2010).
37 Previous authors also indicate that vine vegetative development is highly influenced by available water, up to the extent
38 that it may become a limiting factor. However, under the same irrigation depth, sometime the response between two
39 closer plants is not the same. This point should be considered when selecting methods to estimate crop water status in
40 order to achieve a better management and the production objectives defined at the beginning of the growing season. On
41 one hand, to cover all the water needs is not recommended because it creates management problems, reduces crop quality
42 and overall unnecessarily increases the cost of cultivation. On the other hand, to increase the water availability to the
43 vineyard, the grape production rises as well, but also the canopy, increasing the cost of pruning, plant protection
44 treatments and usually reduces the quality of the grapes. Thus, water stress had to be controlled to achieve a good yield /
45 quality of grapes and balanced growth while avoiding the problems of excess water. Therefore, it is essential to know the
46 right way to manage this crop.

47 Some studies related to spectral Vegetation Indices (VI) performed different analysis of vine canopy, shape, size and
48 functional capacity, in order to manage spatially and temporally vegetation and other productions factors such as water.
49 Spectral VI, have the possibility to predict a large number of plant features, such as Leaf Area Index (LAI), vegetation
50 fraction cover, fraction of Absorbed Photosynthetically Active Radiation (fAPAR), chlorophyll pigment concentration,
51 plant stress and other related parameters (Jordan, 1969; Baret and Guyot, 1991; Peñuelas et al., 1993; Rondeaux et al.,
52 1996; Gitelson and Merzlyak, 2004). These spectral vegetation indexes, which are mathematical combinations of two or
53 more electromagnetic bands reflectance, can be used in vine growth site-specific management enabling the optimization
54 of grape yield and grape yield-quality (Lamb and Bramley, 2001).

55 Nowadays, it is possible to obtain a plant spectral signature with a multispectral proximal sensor (Tardáguila and
56 Diago, 2008), which is relevant to study vine vegetation *terroir*. . The Normalized Difference Vegetation Index (NDVI),
57 developed by Rouse et al. (1973) is one of the most extensively vegetation index used for analysis of vegetation growth.
58 It can be calculated as:

$$59 \quad NDVI = \frac{NIR - Red}{NIR + Red} \quad (1)$$

60 where *Red* and *NIR* parameters are the reflectance in the Red and NIR electromagnetic radiation bands, respectively.
61 When electromagnetic radiation (natural or man-made) impacts on living green leaves, part of it is absorbed, other part is
62 transmitted and the rest is reflected. The electromagnetic radiation spectral range that can be absorbed by plants is the

63 Photosynthetically Active Radiation (PAR), being between 0.4 μm to 0.7 μm (similar to the visible range). In this range,
64 chlorophyll is efficient in capturing the Red and Blue ranges and normally reflects the Green, the Infrared (IR) and the
65 Near Infrared (NIR) ranges. Thus, on the basis of NDVI, the greater amount of vegetative cover or canopy, the greater
66 value in the index. However, the ability to absorb and reflect both bands depends not only on the health plant status but
67 also on its size. In this way, a plant with water stress or any other kind of stress (pests, diseases, nutritional deficiencies
68 ...), will have less capacity to absorb the red band by their photosynthetic apparatus and to reflect the NIR band by the
69 cell walls, resulting a lower value of the NDVI. Therefore, the expression of the vineyard vegetative development can be
70 related to NDVI. Several studies have shown the relationship between parameters related to the amount of vegetative
71 canopy vineyard, such as LAI and fAPAR, and physiological factors, such as crop production, quality the grape on
72 harvest or plant water status (Jackson et al., 1983; Smart and Coombe, 1983; Dry, 2000). Furthermore, NDVI is also
73 largely related to the density of vegetative canopy vineyard (Dobrowski et al., 2002; Johnson, 2003; Hall et al., 2008), so
74 that a change in the factors affecting growth and vineyard development could be estimated by the NDVI.

75 Additionally, *terroir* is affected by physical, chemical and biological soil properties and as a tool to interpret these
76 soil properties variations, soil apparent Electrical Conductivity (ECa) may be used. Soil ECa measurements may
77 characterize the soil spatial variability, mainly the soil physical features and have been used by other authors in order to
78 delineate soil homogeneous management zones (Corwin and Lesch, 2003; Moral et al., 2010; Terrón et al., 2013). Soil
79 ECa measurements can be obtained through geoelectric sensors and this can be an easy and economical way of sampling
80 the soil and guiding soil evaluators in their soil properties analysis (Terrón et al., 2011).

81 According to Hall et al. (2002) the implementation of vineyard site-specific tools are needed in order to better manage
82 vineyards. Thus, considering the previous, the present work makes use of precision agriculture tools to determine: i) the
83 effects of different irrigation treatments in the vine vegetation growth considering two different climatic seasons; and ii)
84 the soil influence in the vegetation growth expression.

85 **2 Material and methods**

86 **2.1 Study area and experimental design**

87 The study was carried out during 2012 and 2013 growing seasons, in a field belonging to the Agrarian Research
88 Institute “La Orden - Valdesequera” , in Extremadura (Spain) (38° 51′ N; 6° 40′ E). Study area is located in a vineyard of
89 1.8 ha, varietal Tempranillo (*Vitis vinifera* L.) grafted on Richter-110. It was planted in 2001 by vertical trellis in bilateral
90 cordon system, with 60 cm stem height and 12 buds per plant. Cultivar Tempranillo is a vigorous variety adaptable to all
91 types of soils, preferably slightly acid, and oriented towards the sunny noon terrain.

92 The climate is characterized by mild winters and hot summers, with maximum temperatures reaching 40°C. Rainfall
93 is irregular, with dry summers and often with an annual average below 500 mm. The soil is typical from the Guadiana
94 Valley, with a uniform profile, poorly differentiated. According to the soil survey staff (2006) is into the order Alfisol,
95 suborder Xeralf and the great group is Haploxeralf (Aquic), generally are soils slightly leached, with scarce of calcium
96 and with low sand-adherence value. The upper soil has slight humus content, while the lower soil is poor in it and has
97 weak nitrogen content as well. According to Olsen method, the field content in available phosphorus was satisfactory,
98 while in case of potassium, an essential monovalent cation, was unsatisfactory and sodium content in this soil was low
99 too. It also had lower electrical conductivity and exchangeable cation levels, with a relatively low CEC.

100 The experimental design was a randomised completely blocks, with 4 replicates (plots) per treatment. Each plot had
101 108 vines in 6 rows with 18 vines per row, where the distance between plants and rows were 1.20 m and 2.50 m
102 respectively, placed on a trellis with East-West row direction. Watered treatments were dependent on the growing season
103 (Fig. 1): i) 2012 treatments were divided into four levels of irrigation, corresponding to four levels of Crop
104 Evapotranspiration (ETc) rates: a) Fully watered, based on the application of the 100% of the ETc; b) RDI 50-20, based
105 on the regulated deficit irrigation technic, with a 50% of ETc before veraison and 20% of ETc after it; c) RDI 50-0, based
106 on the regulated deficit irrigation technic, with a 50% of ETc before veraison and 0% of ETc after it; d) Non – watered,
107 based on a rainfed treatment; and ii) 2013 treatments were reduced to three levels of irrigation, corresponding to three
108 levels of ETc rates: a) Fully watered, based on the application of the 100% of the ETc; b) RDI 30, based on the regulated
109 deficit irrigation technic, with a 30% of ETc throughout the season; and c) Non – watered, based on a rainfed treatment.

110 The irrigation system is characterized by drip irrigation with one emitter of 4 l h⁻¹ every 0.6 m (two emitters per vine)
111 attached to a wire suspended 0.4 m above the ground. Full ETc was calculated by means of the weight differences
112 recorded in a weighing lysimeter installed in the centre of the assay, corresponding to a fully watered treatment plot
113 (Yrissarry and Naveso, 1999). Two grapevine plants were planted into the lysimeter container in order to provide the
114 water balance along their canopy development. Precipitation was collected by an agro-meteorological station located
115 over a reference prairie nearby the vineyard.

116 Soil management was characterized by two annual cultivator treatments, one in winter dormancy and another in bud
117 break phenological stage. Later on, spontaneous vegetation was controlled by herbicide treatments. Furthermore, it is
118 added to soil 250-350 kg/ha. of NPK fertilization (9-18-27). Figure 2 shows the spatial distribution of some soil
119 physicochemical parameters analyzed by official laboratory procedures. Regarding to the canopy management, a spring
120 pruning was realized to adjust the potential yield to the 12 initial shoots. Subsequently, before veraison stage, growing

121 shoots are introduced into the trellis to facilitate passage of agricultural machinery. From veraison to harvest, plant
122 protect treatments were done against cryptogamic diseases in cycles of 15 to 20 days.

123 **2.2 Vegetation index and soil apparent electrical conductivity**

124 The NDVI estimation was performed with two active proximal multi-spectral sensors mounted on All – Terrain
125 Vehicle (ATV). These sensors (OptRx ACS-430, Ag Leader Technology, USA) report directly the vineyard canopy
126 NDVI calculated with Red (0.67 μm) and NIR (0.78 μm) wavelengths. Datasets were collected using a PDA data logger
127 connected to the sensors with the TopView software (Betop Topografía SL, Seville - Spain). Geographical coordinates
128 were obtained by a dual frequency GPS (GGD Maxor JAVAD Javad GNSS Inc., U.S.A.) with Real – Time Kinematic
129 (RTK) differential corrections that reached a planimetric accuracy lower than 0.03 m. To obtain the vineyard canopy
130 reflectance the active multi-spectral sensors were placed at nadir position and at a distance, from the top of the
131 grapevines rows, of 0.80 m (\pm 0.20 m, depending on the vineyard height) (Fig. 3). The number of intra-year spectral
132 datasets was fixed to 5 and, according to the season: i) in 2012, they were started on 29 May and ended on 6 September;
133 and ii) in 2013, they were started on 30 May and ended on 2 September.

134 To validate the NDVI with the LAI, several measurements of the latter was carried out throughout the ripening stage
135 of the crop in both years. Measurements was recorded by a Plant Canopy Analyser LAI-2000 (LI-COR, Inc, U.S.A.),
136 following the procedure of Mabrouk and Carbonneau (1996).

137 ECa measurements were conducted on 18 February 2011, with a VERIS 3150 Surveyor sensor (Fig 4.), obtaining
138 simultaneously in two different soil levels: i) Shallow or ECs – in to a depth of 0.30 m from the soil surface and, ii) Deep
139 or ECd – in to a depth of 0.80 m from the surface. Sampling details can be consulted at Moral et al. (2010).

140 **2.3 Geostatistical and statistical data processing**

141 The samplings showed in this work, corresponding to each dataset of both growing seasons, were statistically
142 analyzed by means of some tools contained in the ArcGIS v.10.1 software (ESRI, U.S.A), for those geostatistically
143 analyses, and SPSS v.17 software (SPSS Inc., U.S.A.), for inferential statistics analyses.

144 The geostatistical analysis of the multi – temporal NDVI samplings included the followings phases: i) Voronoi map –
145 it was performed a previous exploratory analysis of the samplings to take out outliers; ii) Ordinary Kriging interpolation
146 – the parameters used in the semivariograms of each sampling to generate the corresponding maps are showed in Table 1.
147 Once obtained these maps, they were rasterized using a pixel size of 2 m; iii) Principal Component Analysis (PCA) – at
148 this work, a PCA process was established separately for each of the years of study. At each analysis, input raster dataset
149 included the five NDVI sampling of the growing seasons, and the output data were distributed in 5 principal components.

150 Thus, results of PCA analyses obtained were composed of five principal components for each year, where the first
151 principal component shows the spatial variability of NDVI for the whole of all mapping dates of each year.

152 Meanwhile, the samplings belonging to the ECa were also geostatistically analyzed. In this case, only ordinary kriging
153 interpolation tool was used, from which it was obtained the ECs and ECd maps of 2011. The parameters used to
154 interpolate the samplings of ECa are shown in Table 1.

155 Furthermore, NDVI samplings of both growing seasons and samplings of ECa of both depths acquired by kriging
156 were statistically analyzed in two phases: i) On the one hand, it was acquired descriptive parameters of each water
157 treatment in each sampling date to get a global knowledge of the behavior of each component that make up the statistical
158 design; ii) On the other hand, it was done variance analyses of each treatment in each sampling date too. These analyses
159 let compare the behavior the spatial and temporal behavior previously mentioned.

160 In addition, with the aim of determine the importance of the local soil characteristics over the vegetative expression of
161 the vineyard, given by the ECa and NDVI parameters respectively, it was used the Geographically Weighted Regression
162 tool (GWR), included in the ArcGIS v.10.1 software (ESRI, U.S.A.). The relationship between both variables resulted in
163 maps of coefficient of determination (R^2) of each water treatment, growing season and depth. The chosen geometric
164 resolution was of 4 m of spatial resolution, which led the goodness of fit in the influence of soil characteristics on the
165 vegetative growth of vines in each of the irrigation treatments of the assay.

166 **3 Results and Discussions**

167 Climatic variables logged by the weather station sited in a reference prairie nearby the tested vineyard, recorded a
168 diverse behaviour during the two – years test, with drier conditions in the first growing season. Figures 5a and 5b show
169 the cumulative annual rainfall, the cumulative annual ETc, temperature parameters and growing degree days (GDD) on
170 both years. Focusing in the accumulation of precipitation, the total amount on the second year trial (2013) was more than
171 double when compared with the first year trial, where only in its first quarter had the same amount of rainfall that the
172 whole previous season. However, during final stages of vegetative development and within the whole ripening
173 phenologic stages, both years had a similar low accumulation of precipitation. On the contrary, temperature was not very
174 different between both years. The observed climatological differences on both seasons influenced differentially the
175 vineyard vegetative development when considering the different irrigation treatments analysed on this study.

176 On the other hand, in spite of the large differences in precipitations between the two growing seasons, it is observed
177 how, being the wettest, the second year of the test presented a similar hydric demand that the previous year. This result
178 allowed to compare the vegetative response of two consecutive years that were very different on their climatology.
179 Furthermore, if this premise is constant over the years, it could be possible to know the total needs of the culture of

180 vineyards at any annual climatological quality and make appropriate reductions in ETc for a watering schedule based on
181 precipitation occurred in every moment of the campaign. Obviously, and according to Wample and Smithyman (2002), it
182 must be taken into account the increases of hydric necessities of each phenological stage, which are showed in the slope
183 changes of the accumulation curve of ETc (Fig. 5a), paying more attention in dry seasons to not producing unwanted
184 water stresses to the vineyard.

185 In this study, Figure 6 shows the relationship between LAI estimations and NDVI measurements, which
186 measurements of LAI recorded throughout the period ripening of the crop in both years confirm that they are well related
187 ($R^2 = 0.81$), indicating the ability to estimate the degree of development of the vineyard crops by NDVI determinations
188 obtained by proximal active sensors. These results are coincident with several authors, which has been stated a good
189 relationship NDVI – LAI (Johnson, 2003).

190 Regarding to the temporal variability, Fig. 7 shows the obtained results in the first principal component (PC1) of
191 each PCA made to the different mapping dates in each growing season. According to the results, there were differences
192 in plant development even when the same doses of irrigation and cultural practices were received into the different plots
193 of each type of treatment of irrigated. In this way, it was estimated the spatial variability of the soil properties by means
194 of laboratory analyses (Fig. 2) and the geographical determination of ECa, shallow and deep, which are represented in
195 Fig 8. There seems to be a pattern consisted in a variation of ECa from the northern and southern boundaries of the assay
196 up to its centre and, on the other hand, from east to west, coincident with some physicochemical parameters of soil. Then,
197 exist a pattern in the soil characteristics variability due to the good relationship that ECa keeps with some of them,
198 mainly with the clay content, and soil pH (Moral et al., 2010). The spatial variability of ECa, shallow and deep, also had
199 shown significant differences among the locations of the plots of the different irrigation treatments (Table 2), designating
200 different values in the soil properties that influenced the vegetative growth of grapevines. It is observed how the plots of
201 each treatment shown, in general, the spatial variability pattern above discussed, presenting higher values of ECs or ECd
202 in plots near to the northern and southern boundaries of the vineyard test site. Because of this spatial variability, even
203 within plots of the same treatment, it was necessary the geostatistical analysis between NDVI and ECa to know how
204 much influence the soil properties on the vegetative growth of the vineyard in each irrigation treatments and their
205 respective plots.

206 **3.1 Intra-year variability**

207 **3.1.1 2012 growing season**

208 Figure 9 shows both temporal a spatial evolution of NDVI index of the irrigation treatments and their respective plots
209 in the 2012 growing season. At first glance, the results of NDVI mapping of this year show how all the treatments had a

210 temporal evolution similar to Gaussian function, increasing the mean value of the index as the campaign went, reaching a
211 maximum value around the phenological stage of veraison, from which the index went lower up to the harvest. In spite of
212 this sigmoidal evolution, a positive relationship between the NDVI and the water dose was produced, in which the Fully
213 watered treatment kept the mean value of NDVI higher for all the mapping dates, and the Non-watered treatment the
214 lower mean value being this differences, furthermore, significant (Table 3). These results indicate that the more quantity
215 of water in vineyard the more vegetative development of its canopy.

216 The intermediate RDI 50-20 and RDI 50-0 irrigation treatments also had significant differences between the NDVI
217 values regarding to the previous ones, positioning itself at intermediate values. Both RDI treatments kept similar their
218 NDVI values up to January and then they were differentiated because of the change on the water dose of the
219 experimental design. At that moment, the RDI 50-0 treatment had a higher decreasing in the NDVI mean value and,
220 consequently, in the vegetative expression of the vineyard. Taking into account these aspects, and knowing the existing
221 relationship between the vegetative growth of the vines and the NDVI value, it can be considered that the last one
222 increase its value when the water doses are higher and variations on that dose will result in changes on the vegetative
223 expression of the vineyard.

224 On the other hand, despite the relationship given among the water doses applied in the assay and the vegetative
225 development of the vines, significant differences were given among the several plot of each one of them (data not
226 shown), indicating that exist a spatial variability of the NDVI index, thus the vegetative growth too, that is dependent of
227 other factors, but the characteristics of the management were identical. At this way, it is observed in Fig. 9 how the
228 vegetative expression was not homogeneous at the whole of plots within a specific water treatment, but it was found
229 variations in the NDVI value dependent on the geographical location of each one of those plots. Thus, for a specific
230 mapping date, some plots of different water treatments had similar mean values of NDVI, even among plots of Fully
231 watered and Non-watered treatments. Then, it was occurred an associated factor to the geographical location that
232 provided some influence over the vegetative growth. The *terroir* effect, in which are included the physicochemical
233 parameters of soil, could be one of the factors that caused a certain influence on the vegetative development, as indicated
234 by Van Leeuwen and Seguin (2006).

235 A priori, the global results about the relationship between NDVI and ECa indicated a low association if it is compared
236 the first 0.3 m of soil depth (ECs, Table 4), and relatively high when it is considered a large section of soil (ECd, Table
237 4). This results suggest that the soil surface layer is not very much influent over the vegetative expression of the
238 vineyard, which a pivoting conformation of the roots cause no effect substantially to their development, but it does in
239 other crops with shallow roots (Fortes et al., 2014). Furthermore, in the year where the climatic quality involves drought

240 (2012), ECa and NDVI values were lower, suggesting that the soil properties seem to be an influent factor but not a
241 limiting one over the vegetative expression and it does the availability of water resources.

242 3.1.2 2013 growing season

243 Figure 10 shows the spatial and temporal evolution of NDVI in the watered treatments and their respective plots in
244 the 2013 growing season. At the same way the previous year, an increase in the water doses applied to the vineyard still
245 being associated to a higher mean value of NDVI index. However, in this season, the differences occurred in this mean
246 value were closer, being no higher than 0.1 points of index value. The intense precipitations given between post-harvest
247 of 2012 and flowering of 2013 decreased the possibility of water stress in the vines, so its vegetative development it was
248 presented very similar at the beginning of the NDVI mappings, differing only the RDI 30 treatment that coming from the
249 RDI 50-20 of the previous growing season (Table 3). On the other hand, at this 2013 season, the temporal evolution of
250 mean value of NDVI of the whole treatments was more homogeneous during most of the season. Generally speaking, it
251 was an initial increment of the NDVI value in all treatments up to the phenological stage of veraison, from which that
252 value was constant up to the harvest. Both results, higher and constant values of NDVI than the previous season could be
253 caused by the high groundwater recharge, which could provide water available to plants almost without limitations
254 during the early stages of vineyard growth.

255 Related to the temporal behavior of the NDVI among water treatments, the mean value of the index resulted in
256 significant differences slightly higher according as the season went, establishing around the veraison two different groups
257 of treatments (Table 3): i) Fully watered and RDI 30a, and; ii) Non-watered and RD 30b. Since that moment, and up to
258 harvest, the irrigation treatments of the first group shown significant differences in the mean value of NDVI, while
259 treatments of the second one had a similar value. In general, at the same way that the previous season, there were some
260 factors, in this case climatological ones, that modified the expected trend of a vineyard managed under specific water
261 conditions.

262 The irrigation treatments of 2013 growing season also had spatial significant differences in mean value of NDVI
263 among their respective plots (data not shown), following a reduction pattern of its value from north to south of the
264 vineyard test area. Thus, for the same water treatment and mapping date, the mean value of NDVI of each plot decreased
265 the further south was located that plot, existing in addition, significant differences among them. This result was already
266 shown by Blanco et al. (2012), indicating that vegetative growth of the vines under the same management had different
267 behaviors due to spatial changes in some influent factor, such us the spatial variability of the physic – chemical properties
268 of soil. On the other hand, the influence of *terroir*, taken into account its climatic and edaphic factors, was so high in the
269 2013 season that caused that closed plots of different irrigation treatments had similar mean values of NDVI, with some

270 exceptions. Thus, for example, northern plots of Fully watered and Non-watered treatments had shown a similar value of
271 NDVI, at the same way the southern plots, but being statistically different between both geographical locations. This
272 behavior can be observed in Fig. 10.

273 Figure 11 shows the local relationship between the PC1 of NDVI of each growing season and ECa of 2011, shallow
274 and deep, along the test area, which it is given the level of influence of the soil features over the vegetative development
275 in each water treatment. The highest ratios prevailed, again, in the northern and southern limits of the test area, agreeing
276 with those zones where ECa reached the lower values. Thus, the maximum values in the relationship between soil
277 properties and vegetative growth were given during the 2013 season, with values of R^2 in the relationship between soil
278 properties and vegetative growth were given during the 2013 season, with values of R^2 of 0.55 and 0.64 points of ECs
279 and ECd respectively, compared to the 0.56 and 0.47 points reached in 2012. Nevertheless, this latter growing season
280 shows high relationship in a large area of the assay, suggesting that, in drier seasons with lower amount of available
281 groundwater, the variability of soil were influent over a great vegetation surface, but soils with limits on water in zones
282 where ECa has low values, and lower clay content expected (Sudduth et al., 2005; Terrón et al., 2011), tend to have
283 higher availability to the plant of the water that contain versus soils or zones with higher clay content (higher values of
284 ECa).

285 **3.2 Between-year variability**

286 The results of each mapping date of NDVI of both growing seasons, in Figs. 9 and 10, shown the behavior of the
287 vegetative development of the whole treatments established in the experimental designs. As said before, NDVI values
288 and, accordingly, the vegetative growth of the vineyard were influenced by means of the soil properties (included the
289 level of waterground), in its spatial component, and climatic features, in its temporal ones.

290 Regarding to the temporal variability, Fig. 7 shows the obtained results in the first principal component (PC1) of each
291 PCA made to the different mapping dates in each growing season. This PC1 shows the spatial variability of NDVI for the
292 whole of NDVI mapping dates of each year. Thus, each PC1 map of 2012 explains an 80.57% of the temporal variability
293 of each geographical location within the assay area, and an 85.92% for the 2013 growing season. Thus, PC1 of each year
294 shows more than an 80% of the mean variability of the NDVI values throughout both seasons in each irrigation
295 treatments and their respective plots. In general, PC1 map of 2013 shows higher and homogeneous values than the 2012
296 one, indicating a higher and homogeneous vegetative growth of grapevines.

297 On the other hand, Table 5 shows the level of relationship of NDVI values among the different mapping dates for
298 each irrigation treatment. Generally speaking, both 2012 and 2013 got an increase of the correlation coefficient (R) given
299 by the NDVI values as the season went, indicating that the continuous development of the vineyard canopy it was

300 slowing, i.e., the development rate or evolution of that canopy was increasingly smaller up to reach the harvest stage.
301 However, the behavior of the different irrigation treatments did not equally evolve neither intra-year nor inter year ways.
302 So, in 2012, the treatment with higher water doses (Fully watered), had low values of correlation (R lower than 0.65) in
303 all NDVI mapping dates due to a higher development rate versus the rest of water and rainfed treatments during the later
304 phenological stages of the vineyard, indicating higher change rates. On the other side, Non-watered treatment had
305 correlation coefficients above 0.65 points, suggesting a low development rate due to the lower hydric availability, as
306 limiting factor. Meanwhile, the 2013 season had shown a similar behavior pattern in the extremes water treatments.
307 Obviously, the correlation coefficients were shown higher and homogeneous than the previous season among the
308 different mapping dates due to the intense precipitations, being $R < 0.77$ for Fully watered and $R > 0.73$ for Non-watered
309 treatments. These results point out a lower canopy development than 2012 and, within the 2013 season, the differences
310 among treatments were less pronounced.

311 Respecting to differences on the spatial variability of the vegetative growth between years tested, the 2013 season
312 shown a higher homogeneity, where the higher rise was given in the northern half of the test area, independently of the
313 water dose applied. On the other hand, this vegetative development was lower the further south, where the southern plot
314 of Non-watered treatment had not the lower vegetative growth, but responded to a spatial pattern. Thus, the response of
315 vegetation in 2012 was more dependent of the irrigation treatments, meanwhile in 2013 it was more dependent of the soil
316 characteristics or other edaphic – climatic variables. In 2013, RDI 50-20 and RDI 50-0 treatments became RDI 30a and
317 RDI 30b respectively, with water dose of 30% of ETc during the whole irrigation period. At the same way that the rest of
318 the treatments had higher values of NDVI in 2013, RDI 30 also shown higher values of NDVI than the RDI treatments of
319 the previous season. However, despite to have the same water dose, RDI 30b resulted in lower values than RDI 30a
320 during most of the season (data not shown), suggesting one more time that the water dose must be redefined considering
321 the climate and the soil properties.

322 According to Howell (2001), there must be an optimal method of management of a crop at any situation, with the
323 goal to obtain yields and qualities searched and, but the intra – year and between – year management must be performed
324 depending on the *terroir* features of each year or a group of them.

325 **4 Conclusions**

326 Water level and vegetative growth are clearly related, where a higher availability of water resources gave way to a
327 higher vegetative development of the vineyard. However, changes spatio – temporal in the climatic quality or in the soil
328 properties also affect to its vegetative expression. At the already estimated differences in the vegetative growth of
329 grapevines among different water doses, it must be applied the effects that the climate and soil properties perform over

330 the plants. Due to that, the application of the same cultural practices in each growing season makes unfeasible the
331 attainment of stable goals during them, i.e., the same level of quality in grapes and wines or similar yields every season.
332 The application of some precision agriculture techniques to the vineyard crop, through real-time measurements of the
333 NDVI and ECa, makes possible the determination of homogeneous zones of growth and development of the vineyard as
334 function of the climatic and soil characteristics for a specific irrigation treatment. Thus, according to the results of this
335 study: i) in global terms, the higher water doses the higher values of NDVI and, hence, the higher vegetative growth of
336 the vineyard; ii) nevertheless, the vegetative development is not homogeneous, even when the same cultural practices are
337 being used, but it is shown a spatial and temporal variability as function of the climatic and soil characteristics, and the
338 interaction among them; iii) so, it is necessary that the crop management fits to the variability of the agronomic factors to
339 reach an homogeneous vegetative growth even in zones where the soil characteristics are different. The irrigation
340 schedule as function of the real-time results of the NDVI, and the knowledge of the variability of the soil characteristics
341 could be the basis to improve the vineyard management.

342

343

344 **Acknowledgements**

345 This work was carried out with funding the RITECA Project, Transboundary Research Network Extremadura, Center
346 and Alentejo, co-financed by the European Regional Development Fund (ERDF) by the Spain-Portugal Border
347 Cooperation Operational Programme (POCTEP) 2007-2013 and by the Government of Extremadura.

348 This research was also co-financed by the Government of Extremadura and the European Regional Development
349 Fund (ERDF) through the project GR10038 (Research Group TIC008).

350 The vineyard irrigation project with which has complemented this work is INIA RTA2009-00026-C02-02 and it was
351 co-financed by the European Regional Development Fund (ERDF).

352

353 **References**

- 354 Baret, F., and Guyot, G.: Potentials and Limits of Vegetation Indices for Lai and Apar Assessment, Remote
355 Sensing of Environment, 35, 161-173, 1991.
- 356 Blanco, J., Terrón, J. M., Pérez, F. J., Galea, F., Salgado, J. A., Moral, F. J., Marques da Silva, J. R., and Silva, L.
357 L.: Variabilidad Espacial Y Temporal Del Vigor Vegetativo En Viñedo Sin Restricciones Hídricas En La Demanda
358 Evapotranspirativa, VII Congreso Iberico de Agroingeniería y Ciencias Hortícolas, Madrid, 2012.
- 359 Bravdo, B., and Hepner, Y.: Irrigation Management and Fertigation to Optimize Grape Composition and Vine
360 Performance, Symposium on Grapevine Canopy and Vigor Management, XXII IHC 206, 1986, 49-68,
- 361 Carbonneau, A.: General Relationship within the Whole-Plant: Examples of the Influence of Vigour Status,
362 Crop Load and Canopy Exposure on the Sink" Berry Maturation" for the Grapevine, Strategies to Optimize
363 Wine Grape Quality 427, 99-118, 1995.
- 364 Cortell, J. M., Halbleib, M., Gallagher, A. V., Righetti, T. L., and Kennedy, J. A.: Influence of Vine Vigor on
365 Grape (*Vitis Vinifera* L. Cv. Pinot Noir) and Wine Proanthocyanidins, Journal of Agricultural and Food
366 Chemistry, 53, 5798-5808, 10.1021/jf0504770, 2005.
- 367 Corwin, D., and Lesch, S.: Application of Soil Electrical Conductivity to Precision Agriculture, Agronomy
368 Journal, 95, 455-471, 2003.
- 369 Deloire, A., Vaudour, E., Carey, V., Bonnardot, V., and Van Leeuwen, C.: Grapevine Responses to Terroir: A
370 Global Approach, J. Int. Sci. Vigne Vin, 39, 149-162, 2005.
- 371 Dobrowski, S. Z., Ustin, S. L., and Wolpert, J. A.: Remote Estimation of Vine Canopy Density in Vertically
372 Shoot-Positioned Vineyards: Determining Optimal Vegetation Indices, Australian Journal of Grape and Wine
373 Research, 8, 117-125, 10.1111/j.1755-0238.2002.tb00220.x, 2002.
- 374 Dry, P. R.: Canopy Management for Fruitfulness, Australian Journal of Grape and Wine Research, 6, 109-115,
375 10.1111/j.1755-0238.2000.tb00168.x, 2000.
- 376 Fortes, R., Prieto, M., Terron, J., Blanco, J., Millan, S., and Campillo, C.: Using Apparent Electric Conductivity
377 and Ndvi Measurements for Yield Estimation of Processing Tomato Crop, Transactions of the Asabe, 57, 827-
378 835, 2014.
- 379 Gitelson, A. A., and Merzlyak, M. N.: Non-Destructive Assessment of Chlorophyll Carotenoid and
380 Anthocyanin Content in Higher Plant Leaves: Principles and Algorithms, in: Remote Sensing for Agriculture
381 and the Environment, edited by: Stamatiadis, S., Lynch, J. M., and Schepers, J. S., Ella, Greece, 78-94, 2004.
- 382 Hall, A., Louis, J., and Lamb, D. W.: Low-Resolution Remotely Sensed Images of Winegrape Vineyards Map
383 Spatial Variability in Planimetric Canopy Area Instead of Leaf Area Index, Australian Journal of Grape and
384 Wine Research, 14, 9-17, 2008.
- 385 Howell, G. S.: Sustainable Grape Productivity and the Growth-Yield Relationship: A Review, American Journal
386 of Enology and Viticulture, 52, 165-174, 2001.
- 387 Intrigliolo, D. S., and Castel, J. R.: Response of Grapevine Cv. 'Tempranillo' to Timing and Amount of Irrigation:
388 Water Relations, Vine Growth, Yield and Berry and Wine Composition, Irrigation Science, 28, 113-125, 2010.
- 389 Jackson, R. D., Slater, P. N., and Pinter Jr, P. J.: Discrimination of Growth and Water Stress in Wheat by
390 Various Vegetation Indices through Clear and Turbid Atmospheres, Remote Sensing of Environment, 13, 187-
391 208, [http://dx.doi.org/10.1016/0034-4257\(83\)90039-1](http://dx.doi.org/10.1016/0034-4257(83)90039-1), 1983.
- 392 Johnson, L. F.: Temporal Stability of an Ndvi-Lai Relationship in a Napa Valley Vineyard, Australian Journal of
393 Grape and Wine Research, 9, 96-101, 10.1111/j.1755-0238.2003.tb00258.x, 2003.
- 394 Jordan, C. F.: Derivation of Leaf-Area Index from Quality of Light on Forest Floor, Ecology, 50, 663-&, 669,
395 10.2307/1936256, 1969.
- 396 Lamb, D., and Bramley, R. G. V.: Managing and Monitoring Spatial Variability in Vineyard Variability, Natural
397 Resource Management, 4, 25-30, 2001.
- 398 Mabrouk, H., and Carbonneau, A.: A Simple Method for Determination of Grapevine *Vitis Vinifera* L. Leaf
399 Area, Progres Agricole et Viticole, 113, 392-398, 1996.
- 400 Moral, F. J., Terrón, J. M., and Marques da Silva, J. R.: Delineation of Management Zones Using Mobile
401 Measurements of Soil Apparent Electrical Conductivity and Multivariate Geostatistical Techniques, Soil and
402 Tillage Research, 106, 335-343, <http://dx.doi.org/10.1016/j.still.2009.12.002>, 2010.
- 403 Mullins, M. G., Bouquet, A., and Williams, L. E.: Biology of the Grapevine, Cambridge University Press, 1992.

404 Peñuelas, J., Filella, I., Biel, C., Serrano, L., and Save, R.: The Reflectance at the 950–970 Nm Region as an
405 Indicator of Plant Water Status, *International Journal of Remote Sensing*, 14, 1887-1905,
406 10.1080/01431169308954010, 1993.

407 Rondeaux, G., Steven, M., and Baret, F.: Optimization of Soil-Adjusted Vegetation Indices, *Remote Sensing of*
408 *Environment*, 55, 95-107, 10.1016/0034-4257(95)00186-7, 1996.

409 Smart, R., and Coombe, B.: *Water Relations of Grapevines [Vitis]*, Water deficits and plant growth, 1983.

410 Smart, R. E.: Principles of Grapevine Canopy Microclimate Manipulation with Implications for Yield and
411 Quality. A Review, *American Journal of Enology and Viticulture*, 36, 230-239, 1985.

412 Sudduth, K., Kitchen, N., Wiebold, W., Batchelor, W., Bollero, G., Bullock, D., Clay, D., Palm, H., Pierce, F., and
413 Schuler, R.: Relating Apparent Electrical Conductivity to Soil Properties across the North-Central USA,
414 *Computers and electronics in agriculture*, 46, 263-283, 2005.

415 Tardáguila, J., and Diago, M.: *Viticultura De Precisión: Principios Y Tecnologías Aplicadas En El Viñedo*, VI
416 *World Wine Forum*", Logroño. Spain, 2008,

417 Terrón, J., da Silva, J. M., Moral, F., and García-Ferrer, A.: Soil Apparent Electrical Conductivity and
418 Geographically Weighted Regression for Mapping Soil, *Precision Agriculture*, 12, 750-761, 2011.

419 Terrón, J., Moral, F., da Silva, J. M., and Rebollo, F.: Analysis of Spatial Pattern and Temporal Stability of Soil
420 Apparent Electrical Conductivity and Relationship with Yield in a Soil of High Clay Content, 3rd Global
421 Workshop on Proximal Soil Sensing 2013, 2013, 302,

422 Van Leeuwen, C., and Seguin, G.: The Concept of Terroir in Viticulture, *Journal of Wine Research*, 17, 1-10,
423 2006.

424 Vaudour, E.: The Quality of Grapes and Wine in Relation to Geography: Notions of Terroir at Various Scales,
425 *Journal of Wine Research*, 13, 117-141, 10.1080/0957126022000017981, 2002.

426 Wample, R., and Smithyman, R.: Regulated Deficit Irrigation as a Water Management Strategy in *Vitis*
427 *Vinifera* Production, in: *Deficit Irrigation Practices*, 2002.

428 Williams, L. E., and Araujo, F. J.: Correlations among Predawn Leaf, Midday Leaf, and Midday Stem Water
429 Potential and Their Correlations with Other Measures of Soil and Plant Water Status in *Vitis Vinifera*, *Journal*
430 *of the American Society for Horticultural Science*, 127, 448-454, 2002.

431 Yrissarry, J. J. B., and Naveso, F. S.: Use of Weighing Lysimeter and Bowen-Ratio Energy-Balance for
432 Reference and Actual Crop Evapotranspiration Measurements, III International Symposium on Irrigation of
433 Horticultural Crops 537, 1999, 143-150,

434

435

436

Tables**Table 1.** Parameters corresponding to the theoretical semivariograms for NDVI samplings in 2012 and 2013 growing seasons.

| Dataset | Variable | Model | Lag size (m) | Nugget | Range (m) | Partial Sill | RMSE |
|-------------------------|-----------------|--------------|---------------------|---------------|------------------|---------------------|-------------|
| 29 May 2012 | NDVI | Spherical | 6 | 0.009 | 36.5 | 0.003 | 0.098 |
| 6 July 2012 | NDVI | Spherical | 6 | 0.007 | 32.7 | 0.005 | 0.091 |
| 24 July 2012 | NDVI | Spherical | 6 | 0.005 | 31.0 | 0.007 | 0.083 |
| 14 August 2012 | NDVI | Spherical | 6 | 0.005 | 28.7 | 0.007 | 0.078 |
| 6 September 2012 | NDVI | Spherical | 6 | 0.003 | 33.2 | 0.005 | 0.063 |
| 30 May 2013 | NDVI | Spherical | 6 | 0.008 | 72.0 | 0.003 | 0.092 |
| 8 July 2013 | NDVI | Spherical | 6 | 0.004 | 72.0 | 0.003 | 0.068 |
| 22 July 2013 | NDVI | Spherical | 6 | 0.006 | 72.0 | 0.002 | 0.083 |
| 12 August 2013 | NDVI | Spherical | 6 | 0.005 | 72.0 | 0.003 | 0.074 |
| 2 September 2013 | NDVI | Spherical | 6 | 0.002 | 72.0 | 0.002 | 0.051 |
| 18 February 2011 | ECs | Spherical | 7 | 0.321 | 70.6 | 0.808 | 0.601 |
| 18 February 2011 | ECd | Spherical | 7 | 0.594 | 67.3 | 2.264 | 0.943 |

Table 2. Statistic descriptive analyses of shallow and deep soil apparent electrical conductivity interpolated data; sampling was carried out on 18 February 2011.

| Dataset | Treatment | Plot* | Mean (mS m ⁻¹)** | Std. Deviation (mS m ⁻¹) | Minimum (mS m ⁻¹) | Maximum (mS m ⁻¹) | Range (mS m ⁻¹) | Skewness |
|---------|---------------------|-------|---------------------------------|---|----------------------------------|----------------------------------|--------------------------------|----------|
| ECs | Fully watered | 1 | 5.57cd | 0.29 | 4.95 | 6.30 | 1.35 | 0.32 |
| | Fully watered | 2 | 5.49d | 0.33 | 4.84 | 6.21 | 1.37 | 0.01 |
| | Fully watered | 3 | 6.63a | 0.55 | 5.54 | 7.69 | 2.15 | -0.02 |
| | Fully watered | 4 | 5.94b | 0.48 | 5.05 | 6.83 | 1.78 | 0.16 |
| | RDI 50-20 - RDI 30a | 1 | 4.55h | 0.17 | 4.23 | 5.06 | 0.83 | 0.40 |
| | RDI 50-20 - RDI 30a | 2 | 5.52d | 0.42 | 4.60 | 6.75 | 2.15 | -0.08 |
| | RDI 50-20 - RDI 30a | 3 | 6.59a | 0.30 | 5.82 | 7.39 | 1.57 | 0.14 |
| | RDI 50-20 - RDI 30a | 4 | 5.61de | 0.43 | 4.81 | 6.51 | 1.70 | 0.24 |
| | RDI 50-0 - RDI 30b | 1 | 5.29ef | 0.49 | 4.50 | 6.26 | 1.76 | 0.27 |
| | RDI 50-0 - RDI 30b | 2 | 5.25f | 0.23 | 4.72 | 5.63 | 0.91 | -0.52 |
| | RDI 50-0 - RDI 30b | 3 | 5.14f | 0.49 | 4.31 | 6.40 | 2.09 | 0.51 |
| | RDI 50-0 - RDI 30b | 4 | 5.72c | 0.74 | 4.33 | 7.10 | 2.77 | -0.16 |
| | Non Watered | 1 | 4.80g | 0.30 | 4.39 | 5.71 | 1.32 | 0.66 |
| | Non Watered | 2 | 5.4de | 0.45 | 4.27 | 6.45 | 2.18 | -0.29 |
| | Non Watered | 3 | 5.60cd | 0.51 | 4.61 | 6.50 | 1.89 | 0.19 |
| | Non Watered | 4 | 5.49d | 0.30 | 5.03 | 6.60 | 1.57 | 0.76 |
| ECd | Fully watered | 1 | 9.90cd | 0.77 | 8.79 | 13.81 | 5.02 | 2.69 |
| | Fully watered | 2 | 10.01c | 0.39 | 9.06 | 10.83 | 1.77 | 0.21 |
| | Fully watered | 3 | 10.96b | 0.77 | 8.95 | 12.49 | 3.54 | -0.03 |
| | Fully watered | 4 | 9.96c | 0.64 | 8.76 | 12.06 | 3.30 | 0.51 |
| | RDI 50-20 - RDI 30a | 1 | 8.62gh | 0.33 | 8.07 | 9.62 | 1.55 | 0.76 |
| | RDI 50-20 - RDI 30a | 2 | 9.97c | 0.49 | 9.00 | 11.23 | 2.23 | 0.15 |
| | RDI 50-20 - RDI 30a | 3 | 11.37a | 0.36 | 9.62 | 12.00 | 2.38 | -1.38 |
| | RDI 50-20 - RDI 30a | 4 | 8.82fg | 1.05 | 7.11 | 10.82 | 3.71 | 0.23 |
| | RDI 50-0 - RDI 30b | 1 | 8.91f | 0.48 | 7.75 | 10.15 | 2.40 | 0.35 |
| | RDI 50-0 - RDI 30b | 2 | 9.68d | 0.37 | 9.01 | 10.33 | 1.32 | -0.21 |
| | RDI 50-0 - RDI 30b | 3 | 9.76cd | 0.48 | 8.73 | 10.55 | 1.82 | -0.43 |
| | RDI 50-0 - RDI 30b | 4 | 8.88fg | 1.51 | 6.08 | 11.50 | 5.42 | -0.04 |
| | Non Watered | 1 | 8.53h | 0.37 | 7.82 | 9.47 | 1.65 | 0.27 |
| | Non Watered | 2 | 9.40e | 0.77 | 7.41 | 11.14 | 3.73 | -0.20 |
| | Non Watered | 3 | 9.77cd | 0.28 | 9.11 | 10.49 | 1.38 | 0.35 |
| | Non Watered | 4 | 8.72fgh | 0.57 | 7.90 | 10.53 | 2.63 | 0.99 |

* Plots are numbered in a North – South orientation.

** Variance analyses among treatments are made for each dataset independently; a, b, c, d means significant difference at p-value ≤ 0.05 in Tukey post-hoc analysis.

Table 3. Statistic descriptive analyses of NDVI interpolated datasets for 2012 and 2013 growing seasons (dimensionless).

| Dataset | Treatment | Mean* | Std. Deviation | Minimum | Maximum | Range** | Skewness |
|-------------------|--------------------------|--------------|----------------|--------------|--------------|--------------|----------|
| 29 May 2012 | Fully Watered | 0.643a | 0.036 | 0.502 | 0.713 | 0.211 | -0.319 |
| | RDI 50-20 | 0.608b | 0.039 | 0.507 | 0.691 | 0.184 | -0.548 |
| | RDI 50-0 | 0.597c | 0.046 | 0.472 | 0.706 | 0.234 | -0.265 |
| | Non-watered | 0.572d | 0.044 | 0.446 | 0.677 | 0.231 | -0.302 |
| | MEAN | 0.605 | 0.041 | 0.482 | 0.697 | 0.215 | |
| 06 July 2012 | Fully Watered | 0.729a | 0.050 | 0.586 | 0.807 | 0.221 | -0.535 |
| | RDI 50-20 | 0.708b | 0.042 | 0.579 | 0.780 | 0.201 | -0.591 |
| | RDI 50-0 | 0.714b | 0.054 | 0.569 | 0.817 | 0.248 | -0.311 |
| | Non-watered | 0.624c | 0.060 | 0.453 | 0.766 | 0.313 | -0.210 |
| | MEAN | 0.694 | 0.052 | 0.547 | 0.793 | 0.246 | |
| 24 July 2012 | Fully Watered | 0.750a | 0.041 | 0.597 | 0.813 | 0.216 | -0.998 |
| | RDI 50-20 | 0.718b | 0.046 | 0.452 | 0.789 | 0.337 | -1.300 |
| | RDI 50-0 | 0.721b | 0.055 | 0.554 | 0.803 | 0.249 | -0.767 |
| | Non-watered | 0.618c | 0.064 | 0.430 | 0.730 | 0.300 | -0.448 |
| | MEAN | 0.702 | 0.052 | 0.508 | 0.784 | 0.276 | |
| 14 August 2012 | Fully Watered | 0.742a | 0.039 | 0.483 | 0.803 | 0.320 | -1.853 |
| | RDI 50-20 | 0.712b | 0.048 | 0.577 | 0.794 | 0.217 | -0.475 |
| | RDI 50-0 | 0.696c | 0.070 | 0.512 | 0.800 | 0.288 | -0.828 |
| | Non-watered | 0.613d | 0.054 | 0.404 | 0.731 | 0.327 | -0.568 |
| | MEAN | 0.691 | 0.053 | 0.494 | 0.782 | 0.288 | |
| 06 September 2012 | Fully Watered | 0.701a | 0.032 | 0.575 | 0.761 | 0.186 | -0.825 |
| | RDI 50-20 | 0.673b | 0.045 | 0.534 | 0.740 | 0.206 | -0.681 |
| | RDI 50-0 | 0.647c | 0.070 | 0.445 | 0.750 | 0.305 | -0.917 |
| | Non-watered | 0.600d | 0.056 | 0.417 | 0.707 | 0.290 | -0.647 |
| | MEAN | 0.655 | 0.051 | 0.493 | 0.740 | 0.247 | |
| 30 May 2013 | Fully Watered | 0.671b | 0.039 | 0.570 | 0.749 | 0.179 | -0.454 |
| | RDI 30a (previous 50-20) | 0.680a | 0.045 | 0.570 | 0.749 | 0.179 | -0.728 |
| | RDI 30b (previous 50-0) | 0.665b | 0.053 | 0.518 | 0.747 | 0.229 | -0.573 |
| | Non-watered | 0.671b | 0.050 | 0.528 | 0.761 | 0.233 | -0.547 |
| | MEAN | 0.672 | 0.047 | 0.547 | 0.752 | 0.205 | |
| 08 July 2013 | Fully Watered | 0.779a | 0.040 | 0.655 | 0.831 | 0.176 | -0.827 |
| | RDI 30a (previous 50-20) | 0.766b | 0.052 | 0.597 | 0.833 | 0.236 | -1.000 |
| | RDI 30b (previous 50-0) | 0.754bc | 0.069 | 0.555 | 0.832 | 0.277 | -1.138 |
| | Non-watered | 0.761c | 0.050 | 0.614 | 0.823 | 0.209 | -0.808 |
| | MEAN | 0.769 | 0.053 | 0.605 | 0.830 | 0.225 | |
| 22 July 2013 | Fully Watered | 0.737a | 0.034 | 0.646 | 0.794 | 0.148 | -0.429 |
| | RDI 30a (previous 50-20) | 0.738a | 0.049 | 0.607 | 0.792 | 0.185 | -1.200 |
| | RDI 30b (previous 50-0) | 0.724b | 0.063 | 0.547 | 0.802 | 0.255 | -1.238 |
| | Non-watered | 0.728b | 0.043 | 0.617 | 0.792 | 0.175 | -0.659 |
| | MEAN | 0.732 | 0.047 | 0.604 | 0.795 | 0.191 | |
| 12 August 2013 | Fully Watered | 0.749a | 0.042 | 0.632 | 0.822 | 0.190 | -0.366 |
| | RDI 30a (previous 50-20) | 0.734b | 0.053 | 0.570 | 0.797 | 0.227 | -0.986 |
| | RDI 30b (previous 50-0) | 0.721c | 0.071 | 0.542 | 0.810 | 0.268 | -0.989 |
| | Non-watered | 0.718c | 0.050 | 0.583 | 0.796 | 0.213 | -0.735 |
| | MEAN | 0.731 | 0.054 | 0.582 | 0.806 | 0.225 | |
| 02 September 2013 | Fully Watered | 0.753a | 0.030 | 0.656 | 0.795 | 0.139 | -0.766 |
| | RDI 30a (previous 50-20) | 0.742b | 0.035 | 0.624 | 0.790 | 0.166 | -1.076 |
| | RDI 30b (previous 50-0) | 0.731c | 0.054 | 0.564 | 0.791 | 0.227 | -1.133 |
| | Non-watered | 0.725d | 0.037 | 0.609 | 0.781 | 0.172 | -0.543 |
| | MEAN | 0.738 | 0.039 | 0.613 | 0.789 | 0.176 | |

* Variance analyses among treatments are made for each dataset independently; a, b, c, d means significant difference at p-value \leq 0.05 in Tukey post-hoc analysis.

** Statistical range of NDVI values (max – min)

443

Table 4. Correlation matrix (R) between 1st principal components of 2012 and 2013 growing seasons and apparent electrical conductivities, shallow and deep, interpolated data of 2011.


| Variable | 1st PC NDVI 2012 | 1st PC NDVI 2013 | ECs 2011 | ECd 2011 |
|------------------------------------|------------------------------------|------------------------------------|-----------------|-----------------|
| 1st PC NDVI 2012 | 1.00 | | | |
| 1st PC NDVI 2013 | 0.58 | 1.00 | | |
| ECs 2011 | 0.18 | 0.16 | 1.00 | |
| ECd 2011 | 0.59 | 0.70 | 0.83 | 1.00 |

444

Table 5. Correlation matrices among 2012 and 2013 NDVI surfaces of each irrigation treatment.

| | | 2012 | | | | |
|---------------|-----------|--------|--------|---------|-----------|---------|
| Treatment | Dataset | 29 May | 6 July | 24 July | 14 August | 6 Sept. |
| Fully watered | 29 May | 1 | | | | |
| | 6 July | 0.47 | 1 | | | |
| | 24 July | 0.33 | 0.65 | 1 | | |
| | 14 August | 0.42 | 0.35 | 0.47 | 1 | |
| | 6 Sept. | 0.28 | 0.57 | 0.59 | 0.57 | 1 |
| RDI 50-20 | 29 May | 1 | | | | |
| | 6 July | 0.74 | 1 | | | |
| | 24 July | 0.61 | 0.72 | 1 | | |
| | 14 August | 0.69 | 0.79 | 0.70 | 1 | |
| | 6 Sept. | 0.70 | 0.81 | 0.66 | 0.84 | 1 |
| RDI 50-0 | 29 May | 1 | | | | |
| | 6 July | 0.59 | 1 | | | |
| | 24 July | 0.69 | 0.86 | 1 | | |
| | 14 August | 0.69 | 0.89 | 0.86 | 1 | |
| | 6 Sept. | 0.68 | 0.86 | 0.83 | 0.95 | 1 |
| Non-watered | 29 May | 1 | | | | |
| | 6 July | 0.70 | 1 | | | |
| | 24 July | 0.68 | 0.83 | 1 | | |
| | 14 August | 0.66 | 0.83 | 0.81 | 1 | |
| | 6 Sept. | 0.65 | 0.82 | 0.79 | 0.90 | 1 |
| | | 2013 | | | | |
| | | 30 May | 8 July | 22 July | 12 August | 2 Sept. |
| Fully watered | 30 May | 1 | | | | |
| | 8 July | 0.76 | 1 | | | |
| | 22 July | 0.61 | 0.61 | 1 | | |
| | 12 August | 0.58 | 0.66 | 0.67 | 1 | |
| | 2 Sept. | 0.64 | 0.79 | 0.63 | 0.76 | 1 |
| RDI 30a | 30 May | 1 | | | | |
| | 8 July | 0.85 | 1 | | | |
| | 22 July | 0.82 | 0.86 | 1 | | |
| | 12 August | 0.83 | 0.85 | 0.93 | 1 | |
| | 2 Sept. | 0.83 | 0.87 | 0.91 | 0.90 | 1 |
| RDI 30b | 30 May | 1 | | | | |
| | 8 July | 0.90 | 1 | | | |
| | 22 July | 0.87 | 0.93 | 1 | | |
| | 12 August | 0.88 | 0.94 | 0.95 | 1 | |
| | 2 Sept. | 0.89 | 0.95 | 0.95 | 0.96 | 1 |
| Non-watered | 30 May | 1 | | | | |
| | 8 July | 0.80 | 1 | | | |
| | 22 July | 0.77 | 0.88 | 1 | | |
| | 12 August | 0.84 | 0.89 | 0.88 | 1 | |
| | 2 Sept. | 0.73 | 0.83 | 0.85 | 0.86 | 1 |


449 **Figures**



450

451 **Figure 1.** Maps of treatments and respective plots: a) Map of treatments of 2012 growing season; b) Map of treatment of
452 2013 growing season, where “a” and “b” replicates of RDI 30 are in the same emplacement of the respective replicates of
453 RDI 50-20 and RDI 50-0 of the previous season.

454



455

456 **Figure 2.** Maps of spatial distribution of some soil components in the vineyard site: a) Clay; b) Sand; c) Silt; d) pH of
 457 2012 growing season; e) pH of 2013 growing season; f) Organic matter; g) Total Nitrogen; h) Assimilable phosphorus; i)
 458 Potassium ion (K).

459




460
461 **Figure 3.** ATV with two multi-spectral sensors for NDVI mapping of vineyard canopy
462



463

464 **Figure 4.** Mobile sensor platform Veris 3150 for ECa mapping.


465



466

467 **Figure 5a.** Accumulation of rainfall and ETc of 2012 and 2013 growing seasons.


468



469

470 **Figure 5b.** Temperature components recorded on both 2012 and 2013 growing seasons: Tmean, Tmax and Tmin are the
 471 monthly average, maximum and minimum temperature respectively; GDD is the growing degree day reached the last day
 472 of the month.


473



474

475 **Figure 6.** NDVI – LAI relationship of both 2012 and 2013 years.


476



477

478 **Figure 7.** NDVI First principal component of: a) 2012; and b) 2013

479




480

481 **Figure 8.** Interpolated apparent electrical conductivity maps of 2011 growing season: a) shallow ECa map; b) deep ECa

482 map.

483




484

485 **Figure 9.** Interpolated NDVI maps of 2012 growing season: a) 29 May; b) 6 July; c) 24 July; d) 14 August; and e) 6

486 September.


487



488

489 **Figure 10.** NVDI maps year 2013: a) May 30th; b) July 8th; c) July 22nd; d) August 12th; and e) September 2nd.

490



491

492 **Figure 11.** Local R^2 of GWR analyses: a) 1st principal component of NDVI in 2012 and ECs of 2011; b) 1st principal
 493 component of NDVI in 2013 and ECs of 2011; c) 1st principal component of NDVI in 2012 and ECd of 2011; d) 1st
 494 principal component of NDVI in 2013 and ECd 2011.