

**Litter decomposition rate and soil organic matter quality in a patchwork heathland of
Southern Norway**

Giacomo Certini^{1*}, Live S. Vestgarden^{2,3}, Claudia Forte⁴, Line Tau Strand²

¹*Dipartimento di Scienze delle Produzioni Agroalimentari e dell'Ambiente (DISPAA),
Università degli Studi di Firenze, Firenze, Italy*

²*Department of Environmental Sciences, Norwegian University of Life Sciences, Ås, Norway*

³*Department of Environmental and Health Studies, Telemark University College, Bø, Norway*

⁴*Istituto di Chimica dei Composti OrganoMetallici (ICCOM), [UOS Pisa](#), CNR, Pisa, Italy*

* Corresponding author: certini@unifi.it

14 Abstract

15 Norwegian heathland soils, although scant and shallow, are major reservoirs of carbon (C).
16 We aimed at assessing whether vegetation cover and, indirectly, its driving factor soil
17 drainage are good proxies for soil organic matter (SOM) composition and dynamics in a
18 typical heathland area of Southern Norway consisting in a patchwork of three different types
19 of vegetation, dominated by Calluna (*Calluna vulgaris* (L.) Hull), Molinia (*Molinia caerulea*
20 (L.) Moench), or Sphagnum (*Sphagnum capillifolium* (Ehrh.) Hedw.). Such vegetation covers
21 were clearly associated to microtopographic differences, which in turn dictated differences in
22 soil moisture regime, Calluna growing in the driest sites, Sphagnum in the wettest, and
23 Molinia in sites with intermediate moisture.

24 Litter decomposition was followed over a period of 1 year, by placing litterbags filled with
25 biomass from each dominant species in each type of vegetation cover. The composition of the
26 plant material and SOM were investigated by chemical methods and solid-state ^{13}C nuclear
27 magnetic resonance (NMR) spectroscopy.

28 Litter decomposition was faster for Molinia and Calluna, irrespective of the vegetation cover
29 of the site where they were placed. Sphagnum litter decomposed very slowly, especially
30 under Calluna, where the soil environment is by far more oxidising than under itself. In terms
31 of SOM quality, Calluna covered areas showed the greatest differences from the others, in
32 particular a much higher contribution from lipids and aliphatic biopolymers, apparently
33 related to biomass composition.

34 Our findings showed that in the studied environment litter decomposition rate and SOM
35 composition are actually dependent on vegetation cover and/or soil drainage. On this basis,
36 monitoring changes in the patchwork of vegetation types in boreal heathlands could be a
37 reliable cost-effective way to account for climate change induced modifications to SOM and
38 its potential to last.

Line Tau Strand 22/9/y 19:29

Eliminato: under

Line Tau Strand 22/9/y 19:34

Eliminato: living biomass

giacomo 27/9/y 20:33

Eliminato: ,

giacomo 27/9/y 20:34

Eliminato: the bulk

giacomo 27/9/y 20:33

Eliminato: and some extractable fractions of SOM

giacomo 27/9/y 21:45

Eliminato: moisture

giacomo 27/9/y 21:15

Eliminato: in the

giacomo 27/9/y 21:15

Eliminato: induced by climate change

48 **1. Introduction**

49 Heathland vegetation covers approximately 60% of Norway's land area. Norwegian

50 heathland soils, although scant and shallow, are so rich in organic matter that they represent a

51 stock of carbon (C) at least one order of magnitude larger than the aboveground vegetation

52 they sustain (Rosberg et al., 1981). To predict the ecological effects of climate and land use

53 changes, it is essential to understand the nature and environmental dependencies of soil

54 organic matter (SOM) in these widespread systems. In fact, any change influencing their

55 SOM stocks and dynamics may have major consequences for both C balance and the water

56 quality of lakes and rivers (Stuanes et al., 2008).

57 Following changes in SOM stocks is not a simple task, and several approaches have been

58 proposed for this purpose (e.g., Johnson and Curtis, 2001; Trumbore, 2009; Chiti et al.,

59 2011). In some environments, vegetation cover is a good proxy for soil C dynamics, since it

60 controls the input and quality of litter (De Deyn et al., 2008). In turn, vegetation depends,

61 among other factors, on soil drainage, which also influences litter decay and humification

62 processes (Wickland et al., 2010), so representing another possible proxy for SOM storage.

63 Although present-day vegetation may be different from the one the underlying SOM

64 originated from (Chambers et al., 1999; Hjelle et al., 2010), many studies have demonstrated

65 that the most active part of SOM is the youngest (e.g., Leavitt et al., 1996; Trumbore, 2000;

66 Chiti et al., 2009). Trumbore (2000) found that the average age of the carbon dioxide (CO₂)

67 released by decomposition processes in boreal forest soils is 30 years, and 50-60% of total

68 soil respiration arises from SOM with mean residence time less than 1 year. The dominant

69 contribution of recently synthesized organic matter to soil respiration was also assessed by

70 Certini et al. (2003) for forests in temperate regions. Theoretically, the moister and colder the

71 pedoclimate, the better preserved the dead biomass in soil (Hobbie et al., 2000; Hicks Pries et

giacomo 27/9/y 21:16

Eliminato: —interruzione pagina—

Line Tau Strand 22/9/y 19:37

Eliminato: ice-free

giacomo 27/9/y 21:26

Eliminato: the Norwegian

Line Tau Strand 22/9/y 19:38

Eliminato: This consideration is valid also at a larger scale, since heathlands represent a significant portion of the northern regions of America, Europe, and Asia.

Line Tau Strand 15/9/y 15:40

Eliminato: however,

81 | al., 2013). Hence, the wet boreal heathlands are environments where the investigation of a
82 | possible relationship between vegetation covers and SOM dynamics is particularly
83 | meaningful. Here, due to the intense leaching, lost dissolved organic C (DOC) may be much
84 | older than the respired C (Karlun et al., 2005), rendering any possible relationship between
85 | present day vegetation and bulk SOM quality less clear. Nonetheless, in the uppermost soil,
86 | where SOM is younger and less degraded than below, such relationship is expected to be
87 | strong enough.

88 | In Southern Norway, heathland areas are in most cases characterised by the alternate
89 | occurrence – essentially dictated by the soil drainage, in turn controlled by topography,
90 | particle size distribution, and soil depth to bedrock – of three vegetation types, which are
91 | dominated by the heather Calluna (*Calluna vulgaris* (L) Hull), the moor grass Molinia
92 | (*Molinia caerulea* (L) Moench), and the peat moss Sphagnum (*Sphagnum capillifolium*
93 | (*Ehrh.*) Hedw.). Such different vegetation types are cause and effect of the properties and
94 | behaviour of the underlying soil. This is undoubtedly true for the soil profile morphology and
95 | the sequence of horizons, generally ranging from the O-E-Bhs soil sequum of Calluna-
96 | sustaining podzols to multiple H horizons forming histosols where Sphagnum grows (Strand
97 | et al., 2008).

98 | In this study we report an *in situ* investigation of the relationships between vegetation cover,
99 | litter decay rate and soil organic matter composition for a typical montane heathland area in
100 | Southern Norway where the alternation between Calluna, Molinia, and Sphagnum occurs on
101 | decametric scale. The objective of the study was to assess whether in this environment the
102 | current vegetation cover is a good proxy for SOM quality and dynamics. For this purpose,
103 | litter decomposition was followed over a period of 1 year, by placing litterbags filled with
104 | biomass from each dominant species under each type of vegetation cover, so as to simulate
105 | the effects of possible climate change induced shift of vegetation on litter decomposition rate.

Line Tau Strand 15/9/y 15:41

Eliminato: water rich

giacomo 24/9/y 03:41

Eliminato:

Line Tau Strand 22/9/y 19:47

Eliminato: *Calluna*, the moor grass *Molinia*, and the peat moss *Sphagnum*.

Line Tau Strand 22/9/y 19:44

Eliminato: Field studies on the 0-10 cm depth interval revealed that DOC and dissolved organic nitrogen (DON) concentrations increased in the order Sphagnum<Molinia<Calluna in spite of the similar SOM content (Vestgarden et al., 2010). Laboratory experiments with intact soil columns finally showed that milder winters cause a decrease in the release of CO₂, DOC, DON and ammonium (NH₄⁺) compared to winters with severe frost, and that the soil loss of CO₂, DOC, DON and NH₄⁺ is highest under Molinia and lowest under Sphagnum, with Calluna in between (Vestgarden and Austnes, 2009). Relatively little work has focused on the solid phase of SOM in these environments, most of the research having chiefly focused on SOM storage (e.g., Berendse et al., 1994; Kopittke et al., 2013).

giacomo 27/9/y 22:34

Eliminato: e present

giacomo 27/9/y 22:45

Eliminato: To this end

128 Furthermore, the composition of the aboveground biomass, and the bulk SOM, were
129 investigated by chemical methods and solid-state ¹³C nuclear magnetic resonance (NMR)
130 spectroscopy.

giacomo 27/9/y 22:43

Eliminato: ,

giacomo 27/9/y 22:43

Eliminato: and some extractable fractions of SOM

131

132 2. Materials and Methods

133 2.1. Study site

134 The study area, Storgama (59°02'47"N, 8°39'37"E), is located in the Telemark county,
135 southern Norway, at an elevation of 560-m above sea level. The mean annual precipitation in
136 Storgama for the period 1961-1990 was 994 mm, the mean annual air temperature for the
137 same period was 5.0° C. Approximately 30% of the area is barren granite bedrock and
138 boulders, and soil often occurs as pockets in depressions in the bedrock surface (Fig. 1a). The
139 average soil depth generally varies between 10 and 35 cm but greater thicknesses, up to 100
140 cm, do occur. According to the U.S. Soil Taxonomy (Soil Survey Staff, 2010) and moving
141 from drier to wetter locations, soils are Lithic Haplorthods, Lithic Udipsamments, Lithic
142 Endoaquents, and Lithic Haplosaprists. Although there are some scattered or vaguely
143 grouped Scots pines (*Pinus sylvestris* L) and Downy birch trees (*Betula pubescens* Ehrh), the
144 vegetation is largely dominated by heather (*Calluna vulgaris* (L) Hull) at well drained sites,
145 peat moss (*Sphagnum capillifolium* (Ehrh.) Hedw.) at poorly drained sites, and moor grass
146 (*Molinia caerulea* (L) Moench) at intermediately drained sites (Figs. 1a and 1b). These
147 dominant vegetation types are interspersed in the area, forming a patchwork dictated by
148 topography, which in turn is a driving factor of water supply. At the *Calluna* sites *Calluna*
149 *vulgaris* was virtually 100% of the vegetation cover. At the *Molinia* sites some *Calluna*, *Erica*
150 (*Erica tetralix* L), and *Nartecium* (*Narthecium ossifragum* (L) Huds) were associated with
151 *Molinia caerulea* but, on a visual basis, amounting to no more than 5% of the total cover. At
152 the *Sphagnum* sites, *Sphagnum capillifolium*, covered the entire surface except for a few

giacomo 27/9/y 22:56

Eliminato: small

Line Tau Strand 22/9/y 19:27

Eliminato: *Sphagnum* spp. L

giacomo 27/9/y 23:07

Eliminato: soil

Line Tau Strand 17/9/y 15:37

Eliminato: *Sphagnum*

giacomo 24/9/y 03:48

Eliminato: (

giacomo 24/9/y 03:48

Eliminato: (Ehrh.) Hedw.)

161 scattered individuals of *Molinia*, *Erica* and *Calluna*. Hereafter, we will refer to such
 162 vegetation assemblages simply as *Calluna*, *Molinia* and *Sphagnum*, respectively. Further
 163 pictures and information on vegetation and soils at Storgama are reported in Strand et al.
 164 (2008).

165

166 2.2. *Vegetation sampling and analysis*

167 Three sampling sites per dominant vegetation were chosen within an area of approximately
 168 one hectare. At each location, three soil pits were dug down to bedrock, which was 35 to 50
 169 cm deep. All the vegetation above the pit had been previously sampled and divided according
 170 to species. In the case of *Calluna*, the woody stems and branches were separated from the
 171 leaves and flowers. Capitula and the five upper centimetres were used to represent the whole
 172 *Sphagnum* material. Visible roots were picked out from the soil samples and separated
 173 according to species when possible. The aboveground biomass and the roots were analysed
 174 for C and N by dry combustion on oven-dried (60 °C to constant weight) and finely ground
 175 samples using a LECO® CHN1000 Analyser. The aboveground biomass also underwent
 176 NMR investigation.

177

178 2.3. *Soil sampling and analysis*

179 We focused our attention on the uppermost soil layer, where we expected the closest
 180 relationship between SOM quality and current vegetation. Two undisturbed soil samples, to
 181 be used for soil solution extraction, were taken by completely inserting 7.0 cm high and 4.6
 182 cm in inner diameter, rigid cylinders at about five cm depth in each soil profile. The filled
 183 cylinders were carefully extracted from the soil and placed in a cooling box after sealing the
 184 ends with plastic lids. The samples were stored at 4°C, for a maximum of one week, until
 185 they were processed further. Two standard disturbed soil samples were taken near the holes

giacomo 27/9/y 23:21

Eliminato: *Sphagnum*

giacomo 27/9/y 23:21

Eliminato: *Molinia*

giacomo 28/9/y 08:07

Eliminato: and soil

Line Tau Strand 18/9/y 10:58

Eliminato: about a couple of hectares.

giacomo 24/9/y 03:41

Eliminato:

giacomo 24/9/y 03:49

Eliminato: , all

giacomo 28/9/y 01:16

Eliminato: was

Line Tau Strand 18/9/y 11:03

Eliminato: we sampled the living biomass of the dominant vegetation within approximately a square meter.

giacomo 28/9/y 00:33

Eliminato: plant samples

giacomo 28/9/y 00:32

Eliminato: carbon

giacomo 28/9/y 00:32

Eliminato: nitrogen

Claudia Forte 9/10/y 17:30

Eliminato: ,

Claudia Forte 9/10/y 17:30

Eliminato: while just

giacomo 14/10/y 22:04

Eliminato: NMR analyses were performed on

giacomo 28/9/y 00:34

Eliminato: .

giacomo 14/10/y 21:59

Eliminato: analysis

giacomo 28/9/y 00:34

Eliminato: only were performed on the aboveground plant material

giacomo 28/9/y 08:12

Eliminato: water

giacomo 28/9/y 08:12

Eliminato: of cold and hot water extracts

Claudia Forte 9/10/y 17:33

Eliminato: to occur

Claudia Forte 9/10/y 17:33

Eliminato: the

Claudia Forte 9/10/y 17:36

Eliminato: , after sealing the ends with plastic lids,

211 | left by the cylinders and used for C, N, and pH determination, and NMR analysis. As for the
212 | plant material, soil C and N concentrations were measured by dry combustion on oven-dried
213 | and ground samples, while soil pH was determined potentiometrically in a 1:2.5 V/V distilled
214 | water suspension.

Claudia Forte 9/10/y 17:37

Eliminato: s

215 | The soil-containing cylinders were inserted in two-compartment buckets and centrifuged at
216 | 4620 g for 20 min, following the method described by Giesler et al. (1996). The obtained
217 | solution was filtered through a 0.45 µm filter. One aliquot of the filtrate was analysed for
218 | total C (Shimadzu TOC-V element analyser) and, after oxidation by peroxodisulphate
219 | (NS4743 1975), for total N (FiaSTAR, Tecator Spectrophotometer system). Another aliquot
220 | of the filtrate was used to measure hydrophobicity, by determining the ratio between the
221 | absorbances of the solution at 285 and 254 nm using an UV-VIS spectrophotometer (UV-
222 | 1201 Shimadzu). These two absorbances are, in fact, correlated to hydrophobic C (π - π^*
223 | electron transitions occur at ~285 nm for a number of aromatic substances, as described in
224 | Chin et al. 1994) and total C (Brandstetter et al., 1996), respectively.

giacomo 28/9/y 20:06

Eliminato: These C and N fractions were assumed to represent DOC – since inorganic C was not compatible with the low pH of these soils – and total dissolved N (TDN), respectively.

225 | After centrifugation the soil was immediately passed through a 2 mm-mesh sieve. Two grams
226 | of the moist sieved soil was treated as in the second step of the procedure proposed by Ghani
227 | et al. (2003) to obtain hot-water extract (80 °C for 16 h). After centrifugation for 20 min at
228 | 2,000 g and filtration through 0.45 µm filters, the extract was analysed for total C (HWC),
229 | total N (HWN), and carbohydrate C (Carb-C). HWC and HWN were determined by the same
230 | method as DOC and total dissolved nitrogen (TDN), while the analysis of Carb-C was done
231 | according to the “direct determination” method proposed by Safarik and Santrucková (1992).
232 | In brief, 1 mL of the extract was combined in a polyethylene tube with 1 mL 5% phenol
233 | solution and 5 mL concentrated sulphuric acid and immediately shaken on a vortex mixer.
234 | The absorbance of the mixture was read after 1 h at 485 nm on a UV-VIS spectrophotometer

Line Tau Strand 18/9/y 11:23

Eliminato:

241 (UV-1201 Shimadzu). A calibration curve was built with the following standards: 0.00, 0.05,
242 0.10, 0.25, 0.40 g L⁻¹ of α-D glucose (R²=0.9907).

243
244 2.5. Nuclear magnetic resonance spectroscopy

245 The chemical structure of the aboveground vegetation (one composite sample per dominant
246 species, after removal of stems and coarse branches, in the case of Calluna) and SOM (one
247 composite sample each soil pit, hence three samples per vegetation type) was investigated by
248 solid-state ¹³C nuclear magnetic resonance (NMR) spectroscopy using the cross polarization
249 with magic angle spinning (CP MAS) technique. Prior to analysis, soil samples underwent
250 2% HF treatment according to Skjemstad et al. (1994) in order to remove possible
251 paramagnetic oxides, which cause broadened resonances and signal loss. NMR spectra were
252 obtained by a Bruker AMX 300-WB spectrometer equipped with a 4 mm CP MAS probe.
253 The operating frequencies were 300.13 and 75.47 MHz for ¹H and ¹³C, respectively; the π/2
254 pulse was 3.4 μs on the ¹H channel. A contact time of 2 ms and a relaxation delay of 4 s were
255 used. The MAS speed was set to 8 kHz and the number of scans recorded ranged between
256 4,800 and 40,000, depending on the sample. The chemical shifts were referenced to
257 tetramethylsilane (TMS) using adamantane as external standard. Seven chemical-shift regions
258 of the NMR-spectra, corresponding to the main C forms, were integrated and expressed as
259 per cent contribution to total area subtended by the spectrum between 0 and 220 ppm. The
260 seven regions account for alkyl C (0-45 ppm, mainly comprising lipids, waxes, resins,
261 suberin), methoxyl and N-alkyl C (45-60 ppm, comprising the methoxy group of guaiacyl
262 and the two methoxy groups of syringyl lignin moieties at ~56 ppm), O-alkyl C (60-90 ppm,
263 carbohydrates, mainly cellulose and hemicellulose, with contributions from carbohydrate
264 carbons bonded to one oxygen), di-O-alkyl C (90-110 ppm, mainly from polysaccharides,
265 with contributions from anomeric carbons of carbohydrates, i.e., bonded to two oxygens), H-

Line Tau Strand 18/9/y 11:32

Eliminato: 2.4. Total C, N and pH analysis ... [1]

giacomo 27/9/y 23:26

Eliminato: -

Claudia Forte 10/10/y 12:08

Eliminato: ignoring

Claudia Forte 10/10/y 12:08

Eliminato: of the

Line Tau Strand 18/9/y 11:38

Eliminato: and

Line Tau Strand 18/9/y 11:33

Eliminato: one composite bulk soil sample per profile

giacomo 28/9/y 08:56

Eliminato: from the O-horizon of

giacomo 28/9/y 08:56

Eliminato: sampled

Claudia Forte 10/10/y 12:12

Eliminato: iron

277 and C-substituted aromatic C (110-140 ppm), O-substituted aromatic C (140-162 ppm,
 278 mainly from lignin structures, tannins, polyphenols), and carboxyl C (162-190 ppm, esters,
 279 acids and amides); no carbonyl intensity in the 190-220 ppm region, ascribable to aldehydes
 280 and ketones, was detected.

281

282 2.6. Litter decomposition

283 Litter decomposition was determined *in situ* by the litterbag technique. Recently formed
 284 aboveground biomass of Calluna, Molinia, and Sphagnum (approximately, the top 5 cm),
 285 were collected at the end of the growing season in late September. This material, was oven
 286 dried (35 °C to constant weight) and used for filling 10 x 12 cm nylon mesh bags (0.5-1 mm
 287 mesh), with 3.0 g Calluna, 2.0 g Molinia, or 1.0 g Sphagnum, respectively. In November, 32
 288 litterbags of each vegetation type were installed on the surface of each sampling site, except
 289 Calluna under Sphagnum, since a substitution of Calluna by Sphagnum was judged to be
 290 highly improbable. Eight to ten litterbags per type of content were sampled from each site
 291 after 6, 9 and 12 months of decomposition. The removed litterbags were cleaned of plant
 292 remnants and other foreign material, oven dried (35 °C to constant weight), and weighed for
 293 determining mass loss. Their content was thus ground and analysed for carbon and nitrogen
 294 as described for the vegetation and soil samples.

295

296 2.7. Statistics

297 All statistical analyses were performed using the software program SAS (SAS Institute, Inc.,
 298 1990, Cary, NC). After checking the dataset for normality and variance heterogeneity, effects
 299 of vegetation and decomposition site on soil pH and SOM were tested by one-way analysis of
 300 variance (General Linear Model, GLM). Two separate one-way ANOVAs were performed on
 301 litterbags data: one assessing differences in decomposition rate between the three litter types

giacomo 6/10/y 18:31

Eliminato: 0...ppm, mainly from lignin str... [2]

giacomo 2/10/y 11:37

Eliminato: Twigs, last... recently formed ye... [3]

Line Tau Strand 17/9/y 16:09

Eliminato: capitula

giacomo 2/10/y 11:38

Eliminato: upper... approximately, the top 5... [4]

Line Tau Strand 17/9/y 16:10

Eliminato: which represented the most recently formed biomass

giacomo 28/9/y 09:07

Eliminato: ,

Line Tau Strand 15/9/y 15:43

Eliminato: dried at 40 °C

giacomo 28/9/y 10:03

Eliminato: air

Line Tau Strand 17/9/y 16:15

Eliminato: netting

giacomo 28/9/y 09:55

Eliminato: The Calluna and Molinia material had to be cut into pieces smaller than 6 cm to fit into the bags.

Claudia Forte 10/10/y 12:17

Eliminato: we installed

giacomo 28/9/y 09:55

Eliminato: litter ... vegetation type were inst... [5]

Line Tau Strand 22/9/y 20:11

Eliminato: dried at 40

giacomo 28/9/y 10:05

Eliminato: air dried

Line Tau Strand 22/9/y 20:11

Eliminato: °C to

giacomo 28/9/y 10:05

Eliminato: initial moisture

giacomo 28/9/y 17:06

Spustato (inserimento) [8]

giacomo 28/9/y 17:13

Eliminato: The... after checking the dataset ... [6]

giacomo 28/9/y 17:01

Spustato in giù [7]: Pairwise comparisons were done by the Tukey's Simultaneous test.

giacomo 28/9/y 15:57

Eliminato: For the litterbag study ... wt... < ... [7]

disregarding the dominant vegetation effect, the other, assessing differences between the dominant vegetations, disregarding the litter type effect. Pairwise comparisons were done by the Tukey's Simultaneous test.

3. Results

The experimental data set indicated marked differences in composition among the dominant plant species. The C concentration in the aboveground biomass increased in the order Sphagnum<Molinia<Calluna, whereas the C/N ratio increased in the order Molinia<Sphagnum<Calluna (Table 1), suggesting a parallel increase in intrinsic recalcitrance to decomposition. Belowground, Calluna, and Molinia showed similar composition to that of the aboveground biomass (Table 1), which supports the use of the above biomass only, and not the roots as well, without introducing major errors in the litterbags experiment.

Concerning the soil, the measured pH values, all much below neutrality (Table 2), ensured that all C there present was in organic forms. The C/N ratio of SOM was higher under Calluna than under the other two types of vegetation, thus reflecting the trend observed for the aboveground biomass. However, it must be noted that the differences among the soils were much smaller than those among the dominating plant species. For the rest, soils under the three vegetation types were fairly similar, although Calluna differed significantly from Molinia and Sphagnum in terms of SON, HWN, and HW-C/N ratio (Table 2).

There was a large variability in soil DOC and TDN concentrations, and vegetation types did not show any significant difference with respect to these two variables (Table 2). On the contrary, the hydrophobicity index was significantly different in the three types of vegetation, being highest for Calluna and lowest for Molinia. This difference indicates that a greater proportion of DOC under Calluna was hydrophobic.

Claudia Forte 10/10/y 12:19	Eliminato: (
giacomo 28/9/y 16:06	giacomo 28/9/y 16:06
Eliminato: no...isregarding the site	... [8]
Claudia Forte 10/10/y 12:19	Eliminato:)
giacomo 14/10/y 22:10	Eliminato: and one with...assessing differ
Eliminato: as factor for each litter type	... [9]
giacomo 28/9/y 16:09	giacomo 28/9/y 16:09
Eliminato: as factor for each litter type	giacomo 28/9/y 17:06
giacomo 28/9/y 17:06	Spostato in su [8]: The dataset was test
Spostato (inserimento) [7]	giacomo 28/9/y 17:01
Claudia Forte 10/10/y 12:21	Eliminato: (
giacomo 28/9/y 16:09	Eliminato: dominating vegetation
Eliminato: dominating vegetation	Claudia Forte 10/10/y 12:21
Eliminato:)	giacomo 28/9/y 17:15
giacomo 28/9/y 17:15	Eliminato: .
Eliminato: .	giacomo 28/9/y 17:17
giacomo 28/9/y 17:17	Eliminato: Analysis of the initial
Eliminato: Analysis of the initial	... [12]
Claudia Forte 10/10/y 12:23	Eliminato: rT
giacomo 14/10/y 22:43	Eliminato: clearly
Eliminato: clearly	Claudia Forte 10/10/y 12:23
Claudia Forte 10/10/y 12:23	Eliminato: here were ...arked differences
Eliminato: here were ...arked differences	... [13]
giacomo 24/9/y 03:52	Eliminato: dominate
Eliminato: dominate	Line Tau Strand 22/9/y 20:21
Line Tau Strand 22/9/y 20:21	Eliminato: vegetation types.... The C
Eliminato: vegetation types.... The C	... [14]
Claudia Forte 10/10/y 12:34	Eliminato: followed ...howed
Eliminato: followed ...howed	... [15]
Line Tau Strand 15/9/y 15:50	Eliminato: is a sort of guarantee about...u
Eliminato: is a sort of guarantee about...u	... [16]
Claudia Forte 10/10/y 12:37	Eliminato: using ...he above biomass only
Eliminato: using ...he above biomass only	... [17]
giacomo 28/9/y 19:44	Eliminato: measured ...resent was in orga
Eliminato: measured ...resent was in orga	... [18]
Line Tau Strand 22/9/y 20:28	Eliminato: Assuming that all N is organic, t
Eliminato: Assuming that all N is organic, t	giacomo 28/9/y 19:44
giacomo 28/9/y 19:44	Eliminato: the ...OM was higher under C
Eliminato: the ...OM was higher under C	... [19]
Line Tau Strand 22/9/y 20:33	Eliminato: in the case of SOM
Eliminato: in the case of SOM	Claudia Forte 10/10/y 12:40
Claudia Forte 10/10/y 12:40	Eliminato: between
Eliminato: between	giacomo 28/9/y 19:45
giacomo 28/9/y 19:45	Eliminato: .
Eliminato: .	Line Tau Strand 22/9/y 20:34
Line Tau Strand 22/9/y 20:34	Eliminato: the differences among vegetation types
Eliminato: the differences among vegetation types	Claudia Forte 10/10/y 12:40
Claudia Forte 10/10/y 12:40	Eliminato: between
Eliminato: between	Line Tau Strand 22/9/y 20:30
Line Tau Strand 22/9/y 20:30	Eliminato: ir...biomasses
Eliminato: ir...biomasses	... [20]
giacomo 24/9/y 03:52	Eliminato: species...pecies. For the rest,
Eliminato: species...pecies. For the rest,	... [21]
Line Tau Strand 22/9/y 20:32	Eliminato: and, anyway, they were not sta
Eliminato: and, anyway, they were not sta	... [22]
giacomo 28/9/y 19:59	Eliminato: .
Eliminato: [23]
Claudia Forte 12/10/y 08:05	... [24]

488 The ^{13}C CPMAS NMR spectra of the aboveground biomass and soil are shown in Fig. 2, and
 489 the relative contributions of the different chemical shift regions are reported in Table 3. The
 490 NMR spectra of the aboveground vegetation suggested more similar compositions for
 491 Molinia and Sphagnum with respect to Calluna. The spectrum of the Calluna biomass was
 492 dominated by signals between 60 and 104 ppm, characteristic of polysaccharides; the
 493 relatively high intensity in the alkyl C region (0-50 ppm) ~~was~~ due to lipids and aliphatic
 494 biopolymers. The spectrum also revealed the presence of lignin and tannins, as indicated by
 495 the lignin methoxyl carbon signal at 56 ppm, and the distinct aromatic peaks at 145 and 155
 496 ppm, typical of condensed tannins. The sharp peak at 172 ppm is normally assigned to the
 497 carboxyl C of hemicellulose esters, but may also have contributions from amides (Forte et al.,
 498 2006). The spectra of Molinia and Sphagnum aboveground biomasses showed the same
 499 dominant polysaccharide features of Calluna in the 50-110 ppm range, but a significantly
 500 lower intensity of signal in the alkyl and aromatic C regions, which means lower contribution
 501 of lipids and lignin/tannins, respectively. In the case of Molinia, the slightly narrower signals
 502 in the 60-100 ppm region and the relatively smaller peak shoulder at about 103 ppm
 503 ~~compared with~~ both Calluna and Sphagnum, suggested ~~the occurrence of~~ less hemicellulose
 504 and some crystalline cellulose, ~~respectively~~. Sphagnum did not show the typical lignin
 505 signals, in agreement with the common lignin-free composition of bryophytes (Klaviņa et al.,
 506 2012). The only aromatic signals in the sphagnum spectrum were due to unsubstituted or C-
 507 substituted aryl C at 130 and 117 ppm, while the signal at 158 ppm was ascribable to
 508 phenolic structures. In the case of Calluna, the two sharp tannin peaks at 145 and 155 ppm
 509 observed in the aboveground biomass spectrum were totally absent in the SOM spectrum. In
 510 the case of Molinia, differently from the other two vegetation types, the relative contribution
 511 of aromatic C significantly increased in soil compared to that observed in the aboveground
 512 biomass. In the case of Sphagnum, no major changes occurred in the aromatic region, except

giacomo 28/9/y 22:36

Eliminato: i

giacomo 28/9/y 22:21

Eliminato: with respect

Claudia Forte 10/10/y 12:56

Eliminato: to

giacomo 28/9/y 22:22

Eliminato: respectively

Claudia Forte 10/10/y 12:57

Eliminato: s

giacomo 29/9/y 06:19

Spostato in giù [9]: Overall, the NMR investigation revealed that Calluna and the related SOM were richer in alkyl C and poorer in O-alkyl C than the corresponding specimens from Molinia and Sphagnum (Fig. 2 and Table 3). NMR spectra also revealed that the residues of all three dominant plants, once in soil, experienced a significant increase in the alkyl C contribution and a concomitant decrease in the O-alkyl C one, most probably as a result of a faster decay of carbohydrates than of other C forms and the synthesis of alkyl carbon from the biodegradation of carbohydrate and aromatic fractions (Baldock et al., 1992). Noteworthy differences in the spectral features between the aboveground biomass and soil were observed in the aromatic region as well.

534 for the absence in the soil spectrum of the signal at 158 ppm detected for the aboveground
535 vegetation (Fig. 2). The alkyl C/O-alkyl C ratio increased for all vegetation types on passing
536 from the intact biomass to its decomposition products in soil (Table 3), with large differences
537 in absolute values between Calluna, on the one side, and Molinia and Sphagnum, on the
538 other.

539 The *in situ* decomposition study using litterbags showed that the litter mass remaining after
540 one year of decomposition varied between 62 and 66% in the case of Molinia and Calluna
541 and 83 and 94% for Sphagnum (Fig. 3). The discrepancy between Calluna and Molinia, on the
542 one hand, and Sphagnum, on the other, was lower but significant in the intermediate stages of
543 the experiment. After six months, Calluna showed significantly lower mass loss than Molinia
544 under itself, while at the end of the experiment, Calluna resulted to be better preserved than
545 Molinia only, under Molinia (Fig. 3).

546 In terms of relative C content of the residual litter, Calluna did not change throughout the 12
547 months of the experiment, while Molinia and Sphagnum experienced a marked decrease
548 compared to the original value (Fig. 4). Relative concentrations of N in the litter changed
549 more than the C ones. Except for Sphagnum under itself or under Molinia, all litters increased
550 their N content from November to May; later, all of them increased until August, with the
551 exception of Sphagnum under Calluna and Molinia under itself; finally, in the period from
552 August to November, N concentration continued to increase in Calluna, whereas it decreased
553 in Molinia and showed an irregular trend in Sphagnum (Fig. 4). These C and N trends
554 implied progressive, although slight, decrease in C/N ratio for Calluna and Sphagnum, and a
555 sharper decrease for the same ratio for Molinia until August, after which it increased (Fig. 4).

556 Some site effect in terms of C/N ratio resulted for the Molinia litter only, with significantly
557 higher values under Sphagnum than under Molinia and Calluna. At the end of the experiment,
558 in November, the C/N ratio in Molinia under Sphagnum was even higher than the original

Line Tau Strand 22/9/y 21:09

Eliminato:

giacomo 29/9/y 06:34

Eliminato: Sphagnum

giacomo 29/9/y 06:34

Eliminato: Calluna and Molinia

giacomo 6/10/y 22:25

Eliminato: ,

giacomo 29/9/y 06:34

Eliminato: were

giacomo 29/9/y 06:35

Eliminato: experiment;

giacomo 29/9/y 06:36

Eliminato: if it was

Line Tau Strand 17/9/y 14:28

Eliminato: drastic

giacomo 29/9/y 06:46

Eliminato: Significant

giacomo 29/9/y 06:49

Eliminato: were observed for

569 value.

570

571 4. Discussion

572 In the heathland environment of Storgama, the composition of SOM appeared to partly reflect

573 that of the parent vegetation. Hence, for example, the abundance in alkyl C in the Calluna

574 biomass relative to the other two vegetation types was transferred to the SOM. Nevertheless,

575 SOM accumulated over a long period of time; hence, it could be the result of multiple

576 changes in vegetation cover in the area and, consequently, partly unrelated to the current

577 vegetation cover, although there is no direct or indirect evidence in this regard. Furthermore,

578 input of wind-blown or water-transported material may not be excluded at any site.

579 Sphagnum showed a composition potentially more prone to decay than Calluna and Molinia.

580 Nevertheless, there were no significant differences in the SOM content of the topsoil of the

581 three vegetation covers. Evidently, the prevailing anoxic conditions limited decomposition at

582 the Sphagnum sites. This is in accordance with several studies that used the type of

583 vegetation cover as a proxy for carbon dynamics, based on the consideration that vegetation

584 chiefly reflects the soil moisture regime (Bridgham et al., 2008; Couwenberg et al., 2011;

585 Delarue et al., 2011), which is in turn a driving factor of litter decomposition (Hobbie et al.,

586 2000; Laiho 2006). Large variability in DOC concentrations and no significant effect of

587 vegetation was observed (Table 2). It must be noted, however, that our study shows the

588 conditions only at one sampling occasion, *i.e.* at the end of the growing season, when DOC

589 concentrations are affected by a considerable contribution from senescing plant material. The

590 measured DOC concentrations were generally in agreement with those recorded in autumn

591 using zero tension lysimeters in soils at Storgama and other Norwegian heathland areas

592 (Strand et al., 2002; Vestgarden et al., 2010), although DOC concentrations in centrifuged

593 and freely drained soil solutions are not directly comparable (Giesler et al., 1996). Similarly

giacomo 29/9/y 08:46

Eliminato: SOM chemical structure

giacomo 24/9/y 03:41

Eliminato:

Line Tau Strand 15/9/y 15:58

Eliminato: maintain memory

giacomo 24/9/y 03:42

Eliminato:

Line Tau Strand 15/9/y 15:58

Eliminato: of

giacomo 29/9/y 08:47

Eliminato: original composition of the

giacomo 29/9/y 08:51

Eliminato: reflected

giacomo 29/9/y 08:52

Eliminato: in

Claudia 12/10/y 07:40

Eliminato: s

giacomo 29/9/y 08:53

Eliminato: the accumulated

Claudia 12/10/y 07:43

Eliminato: in so many years

Claudia 12/10/y 07:44

Eliminato: that it

Claudia 12/10/y 07:44

Eliminato: ,, hence, it could be

giacomo 29/9/y 08:56

Eliminato: could

Line Tau Strand 22/9/y 21:12

Eliminato: the

giacomo 29/9/y 08:59

Eliminato: T

giacomo 29/9/y 08:59

Eliminato: clearly prevented

giacomo 29/9/y 09:00

Eliminato: in

giacomo 24/9/y 03:42

Eliminato:

Line Tau Strand 22/9/y 21:15

Eliminato: associated to

Line Tau Strand 17/9/y 14:36

Eliminato: was richer in SOM than the ones supporting the other vegetation types, which is clearly a result of the prevailing anoxic conditions under Sphagnum.

giacomo 4/10/y 18:22

Eliminato: . Soil drainage

giacomo 4/10/y 18:22

Eliminato: of course also

giacomo 4/10/y 17:27

Eliminato: processes and, thus, of both SOM quantity and quality

Claudia 12/10/y 07:48

Eliminato: As for

623 to DOC, TDN showed a large variability and no apparent correlation with vegetation. The
 624 relatively small amount of water extracted by centrifugation limited the number of possible
 625 analyses, preventing N speciation. TDN therefore included both organic N and inorganic N,
 626 the latter amounting to 25-50% of TDN in soil water from southern Norway (Austnes et al.,
 627 2008; Kaste et al., 2008).

628 The hydrophobicity index of soil water differed significantly among vegetation types.
 629 Apparently, Calluna released DOC with the highest proportion of hydrophobic organic
 630 compounds, perhaps mostly arising from tannins and decomposition of lignin (Dilling and
 631 Kaiser, 2002), which are indeed important components of the Calluna litter (Fig. 2).

632 Hot water C approximately amounted to 4.5% of SOC in all samples, irrespective of
 633 vegetation. This percentage is in the range reported by von Lützow et al. (2007). Significantly
 634 lower amounts of HWN were extracted from the Calluna soils compared to the Molinia and
 635 Sphagnum ones, which also implied significantly higher HWC / HWN-ratio for Calluna
 636 (Table 2). We did not partition HWN, however Curtin et al. (2006) demonstrated that it is
 637 mainly organic and, in suborder, NH₄-N generated by hydrolysis of heat-labile organic N.
 638 The quality of the hot water extract rather well discriminated Calluna from Molinia and
 639 Sphagnum. Some authors have proposed hot water extraction of SOM as a method to
 640 measure the labile SOM pool (Chodak et al., 2003; Ghani et al., 2003; Curtin et al., 2006);
 641 however, other authors consider this method not selective enough for this purpose (Landgraf
 642 et al., 2006; von Lützow et al., 2007). In our case, approximately half the C extracted by hot
 643 water belonged to carbohydrates.

644 The NMR spectra showed clear structural differences in aboveground plant material (Fig. 2
 645 and Table 3). These led us to expect differences in decomposition rates among the three
 646 vegetation types. Overall, the NMR investigation revealed that Calluna and the related SOM
 647 were richer in alkyl C and poorer in O-alkyl C than the corresponding specimens from

giacomo 28/9/y 08:53

Eliminato: total dissolved nitrogen (

giacomo 28/9/y 08:53

Eliminato:)

Claudia 12/10/y 07:50

Eliminato: between

giacomo 29/9/y 09:11

Eliminato: , but is much lower than that reported by Wieder and Starr (1998) for sphagnum peat soils

giacomo 6/10/y 09:59

Eliminato: ,

Line Tau Strand 18/9/y 09:38

Eliminato: except in the Sphagnum soils, where this proportion was lower.

giacomo 6/10/y 09:59

Eliminato: -

Line Tau Strand 22/9/y 20:39

Eliminato: the initial litter quality

Claudia 12/10/y 07:52

Eliminato:

Line Tau Strand 22/9/y 20:39

Eliminato: ,

giacomo 24/9/y 03:40

Eliminato:

Claudia 12/10/y 07:55

Eliminato: The clear differences

Claudia 12/10/y 07:52

Eliminato: in the NMR spectra of the aboveground plant materials

giacomo 24/9/y 03:40

Eliminato:

Line Tau Strand 22/9/y 20:42

Eliminato: which

Claudia 12/10/y 07:55

Eliminato: some discrepancy

Claudia 12/10/y 07:55

Eliminato: between

giacomo 24/9/y 03:45

Eliminato: '

giacomo 29/9/y 06:22

Eliminato: In particular, the Calluna biomass was characterised by an intense signal of lipids, which has been correlated to slow decomposition rates in heathland ecosystems (van Vuuren and van der Eerden, 1992; van Vuuren and Berendse, 1993).

giacomo 29/9/y 06:19

Spotato (inserimento) [9]

674 Molinia and Sphagnum. Actually, the apparent richness in alkyl C has been correlated to slow
 675 decomposition rates of Calluna biomass in heathland ecosystems (van Vuuren and van der
 676 Eerden, 1992; van Vuuren and Berendse, 1993). NMR spectra also revealed that the residues
 677 of all three dominant plants, once in soil, experienced a significant increase in the alkyl C
 678 contribution and a concomitant decrease in the O-alkyl C one, most probably as a result of a
 679 faster decay of carbohydrates than of other C forms and the synthesis of lipids from the
 680 biodegradation of carbohydrate and aromatic fractions (Baldock et al., 1992). Less important
 681 but noteworthy differences in the spectral features between the aboveground biomass and soil
 682 were observed in the aromatic region as well (Fig. 2 and Table 3). The alkyl C / O-alkyl C
 683 ratio, which generally increases as decomposition proceeds, was significantly higher under
 684 Calluna than under Molinia and Sphagnum. Hence, on the basis of the NMR spectra, Calluna
 685 appeared to be potentially more recalcitrant to decomposition than Molinia and Sphagnum.
 686 However, this was not confirmed by our litterbags experiment, where there were little and
 687 variable differences between the mass losses of Calluna and Molinia, and both of them were
 688 much higher than the one in Sphagnum wherever the latter was placed (Fig. 3). A possible
 689 explanation for the intrinsic resistance of Sphagnum could be that this type of vegetation is
 690 particularly rich in sphagnum pectin-like polysaccharides, which, unlike the other types of
 691 polysaccharides, induce processes that prevent organic matter decay (Hájek et al., 2011;
 692 Ballance et al., 2012). Moreover, it must be considered that Sphagnum might have
 693 experienced a “non-additive” pattern of mass loss, i.e. a decomposition behaviour sometimes
 694 observed in litter mixes that deviates from the response predicted for the individual species
 695 because of the influence of the other species present in the mix (Gartner and Cardon, 2004).
 696 In this case, the necromasses of Molinia and, in particular, Calluna could have partly
 697 inhibited the decomposition of the Sphagnum in the litterbags.

giacomo 29/9/y 06:29
Eliminato: (Fig. 2 and Table 3)

giacomo 30/9/y 05:55
Eliminato: alkyl carbon
 giacomo 30/9/y 06:05
Eliminato: N

giacomo 30/9/y 06:09
Eliminato: due to both a prevailing decrease in carbohydrates (O-alkyl C) and a release of hydrophobic by-products of decomposition (alkyl C)

giacomo 2/10/y 11:43
Eliminato: it

giacomo 2/10/y 11:45
Eliminato: , in

giacomo 2/10/y 11:46
Eliminato: , a consistent portion of polysaccharides are

Claudia 12/10/y 07:58
Eliminato: rest of

Line Tau Strand 17/9/y 14:38
Eliminato: , are hard to decompose

Line Tau Strand 17/9/y 14:49
Eliminato: Verhoeven and Toth, 1995; Scheffer et al., 2001

giacomo 24/9/y 03:37
Eliminato: ;

Line Tau Strand 22/9/y 20:44
Eliminato: Hájek

giacomo 24/9/y 03:37
Eliminato: (

giacomo 24/9/y 03:37
Eliminato:)

Claudia 12/10/y 07:59
Eliminato: has to be minded

Claudia 12/10/y 08:00
Eliminato: that type of

Claudia 12/10/y 08:03
Eliminato: -

Claudia 12/10/y 08:04
Eliminato: a single

Claudia 12/10/y 08:06
Eliminato: Therefore

721 In addition to a “vegetation effect”, the litterbags experiment showed some “site effect” *i.e.*
 722 more rapid decomposition when litter was placed beneath the parent vegetation rather than
 723 beneath other species (Ayres et al., 2009; Perez et al., 2013; Wang et al., 2013). In fact, for
 724 Sphagnum the mass loss was significantly lower when it decayed under Calluna than under
 725 Molina or Sphagnum (Fig. 3). Calluna was better preserved under Molina than under itself at
 726 the end of the trial, while after six months only, Molina litter showed significant
 727 environment-induced advantage under Sphagnum compared to under itself (Fig. 3).
 728 Unexpectedly, the well drained Calluna soils preserved Sphagnum and Molina from decay
 729 better than the moister soils where they were growing perhaps as an effect of a seasonal
 730 drought.
 731 In our litterbag experiment, Molina showed an initial C/N ratio much higher than the ones of
 732 Calluna and, especially, Sphagnum (Fig. 4), which suggested a more marked intrinsic
 733 resistance of its tissues to decay. Noteworthy is the difference in C/N ratio between the
 734 aboveground Molina biomass analysed for basic characterisation (data of Table 1) and the
 735 Molina used in the litterbags experiment (30 vs. circa 80). Actually, Molina is a grass that
 736 wilts at the end of the growing season, when we sampled the material to be inserted in the
 737 bags, while the Molina sampled for basic characterisation was still with active
 738 photosynthesis, when the C/N ratio is relatively low (Taylor et al., 2001). On the other hand,
 739 Calluna is an evergreen and no great seasonal changes in C and N concentrations occur, while
 740 sphagnum, although not an evergreen, does not wilt and its C/N ratio is rather constant
 741 throughout the year. The litterbags experiment showed that the C/N ratio is not a highly
 742 reliable predictor of decay in this environment. The anoxic conditions imposed by prolonged
 743 water saturation, commonly occurring in the Sphagnum soils and expected to have
 744 considerable influence in slackening litter decomposition, on the contrary appeared to be
 745 irrelevant in preserving organic residues during our one-year long experiment (Fig. 3). In this

Line Tau Strand 17/9/y 14:48

Eliminato: or “home-field advantage”

giacomo 24/9/y 03:46

Eliminato: , ... *i.e.* more rapid decomposi ... [25]

Line Tau Strand 17/9/y 14:57

Eliminato: (Taylor et al. 2001)

giacomo 2/10/y 11:55

Eliminato: just after six months of trial ... [26]

Line Tau Strand 22/9/y 21:39

Eliminato: This finding leads to hypothesise the action of some antibiotic substances from Calluna. In this regard, Handley (1963) in his investigations on the suppression of tree growth on Calluna heathland found in the raw humus some water soluble-substances that inhibited the development of mycorrhizal hymenomycetes. Since the inhibiting factor seemed to be associated with the Calluna roots, the author suggested an endophyte in the roots as the excretory agent of the antibiotic substances

Claudia 12/10/y 08:11

Eliminato: actually

giacomo 2/10/y 19:44

Eliminato: (Taylor et al. 2001)... while t ... [27]

Claudia 12/10/y 08:12

Eliminato: difference ...easonal changes ... [28]

giacomo 2/10/y 19:27

Eliminato: but...does not wilt and its C/N ... [29]

Line Tau Strand 17/9/y 14:53

Eliminato: Anyway, an outcome of t...he ... [30]

giacomo 2/10/y 19:31

Eliminato: powerful enough...eliable pred ... [31]

810 regard, during a 3-year study in heathlands on *Molinia caerulea* and *Erica tetralix*, van
811 Vuuren and Berendse (1993) did not find any site effect and litter quality appeared to be the
812 sole driving factor. Also Scheffer et al. (2001), studying the decomposition process in fens
813 dominated by Sphagnum species or without Sphagnum, concluded that decomposition was
814 controlled more by intrinsic differences in litter quality than by the environment.

816 5. Conclusions

817 We found that in the varied heathland of Storgama there were many significant differences in
818 terms of SOM composition between the Calluna dominated areas and the interspersed
819 Sphagnum-covered areas. Most differences were clearly due to the quality of the parent
820 vegetation. A “vegetation effect” on litter decomposition rate was clear, Sphagnum remnants
821 being much more refractory independently of the environmental conditions they underwent,
822 which varied especially in terms of soil drainage. Hence, overall, vegetation appeared to be a
823 good proxy for SOM quality. On this basis, monitoring the distribution of vegetation types in
824 heathlands of Norway and elsewhere could be of particular interest for assessing the
825 consequences of environmental changes such as global warming and higher concentration of
826 rainfall on litter stocks and dynamics. In the plausible scenario of a less continuous rainfall
827 supply and a consequent contraction of Sphagnum-covered areas, the Sphagnum-released
828 SOM seems to have good short-term ability to resist decomposition under the two replacing
829 types of vegetation, Molinia and Calluna. Medium to long-term experiments addressing this
830 issue are needed.

833 Acknowledgements

giacomo 2/10/y 19:39

Eliminato: ied

Claudia 12/10/y 08:17

Eliminato: how

Line Tau Strand 18/9/y 09:58

Eliminato: is influenced by the transition from a vascular plant-dominated system to a Sphagnum-dominated system. To this end, they carried out a two-year long reciprocal litterbag experiment using *Carex diandra*, *C. lasiocarpa*, *Sphagnum papillosum* and *S. squarrosum* in a fen

giacomo 2/10/y 19:37

Eliminato: and

Line Tau Strand 18/9/y 09:58

Eliminato: a fen

Line Tau Strand 18/9/y 09:58

Eliminato: .

Line Tau Strand 18/9/y 09:58

Eliminato: The decomposition rate hardly differed between the two sites and was highest for the *Carex* litter types and lowest for the *Sphagnum* ones, indicating

Line Tau Strand 18/9/y 09:58

Eliminato:

giacomo 2/10/y 19:44

Eliminato: egat

giacomo 2/10/y 19:52

Eliminato: , at least over a relatively short term

852 We thank Irene Eriksen Dahl, Grete Bloch, and Ivan Digernes for laboratory assistance at the
 853 Department for Plants and Environmental Sciences, Norwegian University of Life Sciences.
 854 We also thank Silvia Pizzanelli of ICCOM-CNR, for performing part of the NMR analyses.
 855 The study was carried out in close cooperation with the CLUE project (NFR 155826/S30).
 856 This specific investigation was made possible by a grant from the Research Council of
 857 Norway (NFR 164903/S30) enabling the first author to cooperate with researchers from the
 858 Norwegian University of Life Sciences.

860 **References**

861 Austnes, K., Kaste, O., Vestgarden, L. S., and Mulder, J.: Manipulation of snow in small
 862 headwater catchments at Storgama, Norway: Effects on leaching of total organic carbon
 863 and total organic nitrogen. *Ambio* 37: 38–47, 2008.

864 Ayres, E., Steltzer, H., Simmons, B. L., Simpson, R. T., Steinweg, J. M., Wallenstein, M. D.,
 865 Mellor, N., Parton, W. J., Moore J. C., and Wall, D.H.: Home-field advantage accelerates
 866 leaf litter decomposition in forests. *Soil Biol. Biochem.* 41: 606–610, 2009.

867 Baldock, J. A., Oades, J. M., Waters, A. G., Peng, X., Vassallo, A. M., and Wilson, M. A.:
 868 Aspects of the chemical structure of soil organic materials as revealed by solid-state ¹³C
 869 NMR spectroscopy. *Biogeochemistry* 16: 1–42, 1992.

870 Ballance, S., Kristiansen, K. A., Skogaker, N. T., Tvedt, K. E., and Christensen, B. E.: The
 871 localisation of pectin in Sphagnum moss leaves and its role in preservation. *Carbohydr.*
 872 *Polym.* 87: 1326–1332, 2012.

873 Brandstetter, A., Sletten, R. S., Mentler, A., and Wenzel, W. W.: Estimating dissolved
 874 organic carbon in natural waters by UV absorbance (254 nm). *Z. Pflanz. Bodenkunde*
 875 159: 605–607, 1996.

giacomo 24/9/y 03:31

Spostato in giù [1]: K. A.

Line Tau Strand 18/9/y 08:59

Formattato ... [32]

giacomo 24/9/y 03:31

Spostato in giù [2]: N. T.

giacomo 24/9/y 03:31

Spostato (inserimento) [1] ... [33]

giacomo 24/9/y 03:31

Spostato (inserimento) [3] ... [37]

Line Tau Strand 18/9/y 08:59

Formattato ... [34]

Line Tau Strand 18/9/y 08:59

Formattato ... [36]

giacomo 24/9/y 03:31

Spostato (inserimento) [2] ... [35]

giacomo 24/9/y 03:31

Spostato in giù [3]: K. E.

Line Tau Strand 18/9/y 08:59

Formattato ... [38]

giacomo 24/9/y 03:32

Spostato in giù [4]: B. E.

giacomo 24/9/y 03:32

Spostato (inserimento) [4] ... [39]

giacomo 24/9/y 03:31

Eliminato: .

Line Tau Strand 18/9/y 08:59

Formattato ... [40]

giacomo 24/9/y 03:32

Eliminato: .

Line Tau Strand 18/9/y 08:59

Formattato ... [41]

giacomo 24/9/y 11:30

Eliminato: rate

giacomo 24/9/y 11:30

Eliminato: ers

Line Tau Strand 18/9/y 08:59

Formattato ... [42]

Line Tau Strand 18/9/y 08:59

Formattato ... [43]

giacomo 24/9/y 03:29

Eliminato: -

Line Tau Strand 18/9/y 08:59

Formattato ... [44]

giacomo 24/9/y 03:29

Eliminato: .

Line Tau Strand 18/9/y 08:59

Formattato ... [45]

Line Tau Strand 18/9/y 08:59

Formattato ... [46]

Line Tau Strand 18/9/y 08:59

Formattato ... [47]

giacomo 27/9/y 23:24

Eliminato: Berendse, F., Schmitz, M., at ... [48]

891 Bridgman, S. D., Pastor, J., Dewey, B., Weltzin, J. F., and Updegraff, K.: Rapid carbon
 892 response of peatlands to climate change. *Ecology* 89: 3041–3048, 2008.

893 Certini, G., Corti, G., Agnelli, A., and Sanesi, G.: Carbon dioxide efflux and concentrations
 894 in two soils under temperate forests. *Biol. Fert. Soils* 37: 39–46, 2003.

895 Chambers, F. M., Mauquoy, D., and Todd, P. A.: Recent rise to dominance of *Molinia*
 896 *caerulea* in environmentally sensitive areas: new perspectives from palaeoecological data.
 897 *J. Appl. Ecol.* 36: 719–733, 1999.

898 Chin, Y. P., Aiken, G., and Oloughlin, E.: Molecular-weight, polydispersity, and
 899 spectroscopic properties of aquatic humic substances. *Environ. Sci. Technol.* 28: 1853–
 900 1858, 1994.

901 Chiti, T., Certini, G., Perugini, L., Mastrodonato, G., Papale, D., and Valentini, R.: Soil
 902 carbon dynamics in a Mediterranean forest during the Kyoto Protocol commitment
 903 periods. *Reg. Environ. Change* 11: 371–376, 2011.

904 Chiti, T., Neubert, R. E. M., Janssens, I. A., Certini, G., Curiel Yuste, J., and Sirignano, C.:
 905 Radiocarbon dating reveals different past managements of adjacent forest soils in the
 906 Campine region, Belgium. *Geoderma* 149: 137–142, 2009.

907 Chodak, M., Khanna, P., and Beese, F.: Hot water extractable C and N in relation to
 908 microbiological properties of soils under beech forests. *Biol. Fert. Soils* 39: 123–130,
 909 2003.

910 Couwenberg, J., Thiele, A., Tanneberger, F., Augustin, J., Barisch, S., Dubovik, D.,
 911 Liashchynskaya, N., Michaelis, D., Minke, M., Skuratovich, A., and Joosten, H.:
 912 Assessing greenhouse gas emissions from peatlands using vegetation as a proxy.
 913 *Hydrobiologia* 674: 67–89, 2011.

914 Curtin, D., Wright, C. E., Beare, M. H., and McCallum F. M.: Hot water-extractable nitrogen
 915 as an indicator of soil nitrogen availability. *Soil Sci. Soc. Am. J.* 70: 1512–1521, 2006.

916 De Deyn, G. B., Cornelissen, J. H. C., and Bardgett, R. D.: Plant functional traits and soil
 917 carbon sequestration in contrasting biomes. *Ecol. Lett.* 11: 516–531, 2008.

918 Delarue, F., Laggoun-Defarge, F., Disnar, J. R., Lottier, N., and Gogo, S.: Organic matter
 919 sources and decay assessment in a Sphagnum-dominated peatland (Le Forbonnet, Jura
 920 Mountains, France): impact of moisture conditions. *Biogeochemistry* 106: 39–52, 2011.

921 Dilling, J. and Kaiser, K.: Estimation of the hydrophobic fraction of dissolved organic matter
 922 in water samples using UV photometry. *Water Res.* 36: 5037–5044, 2002.

923 Forte, C., Piazzzi, A., Pizzanelli, S., and Certini, G.: CP MAS C-13 spectral editing and
 924 relative quantitation of a soil sample. *Solid State Nucl. Mag.* 30: 81–88, 2006.

925 [Gartner, T. B. and Cardon, Z. G.: Decomposition dynamics in mixed-species leaf litter. *Oikos*](#)
 926 [104: 230–246, 2004.](#)

927 Ghani, A., Dexter, M., and Perrott, K. W.: Hot-water extractable carbon in soils: a sensitive
 928 measurement for determining impacts of fertilisation, grazing and cultivation. *Soil Biol.*
 929 *Biochem.* 35: 1231–1243, 2003.

930 Giesler, R., Lundstrom, U. S., and Grip, H.: Comparison of soil solution chemistry
 931 assessment using zero-tension lysimeters or centrifugation. *Eur. J. Soil Sci.* 47: 395–405,
 932 1996.

933 [Hájek, T., Ballance, S., Limpens, J., Zijlstra, M., and Verhoeven, J. T. A.: Cell-wall](#)
 934 [polysaccharides play an important role in decay resistance of Sphagnum and actively](#)
 935 [depressed decomposition in vitro. *Biogeochemistry* 103: 45–57, 2011.](#)

936 [Hicks Pries, C. E., Schuur, E. A. G., Vogel, J. G., and Natali, S. M.: Moisture drives surface](#)
 937 [decomposition in thawing tundra. *J. Geophys. Res.-Biogeo.* 118: 1133–1143, 2013.](#)

938 Hjelle, K. L., Halvorsen, L. S., and Overland, A.: Heathland development and relationship
 939 between humans and environment along the coast of western Norway through time.
 940 *Quatern. Int.* 220: 133–146, 2010.

Line Tau Strand 22/9/y 21:42

Eliminato: Hajek

giacomo 24/9/y 03:38

Eliminato: (2011)

Line Tau Strand 22/9/y 21:42

Eliminato: Handley, W. R. C.: Mycorrhizal Associations and Calluna Heathland Afforestation. Forestry Commission bulletin, no. 36. Her Majesty's Stationary Office, London, 1963. .

947 Hobbie, S. E., Schimel, J. P., Trumbore, S. E., and Randerson, J. R.: Controls over carbon
 948 storage and turnover in high-latitude soils. *Glob. Change Biol.* 6 (Suppl. 1): 196–210,
 949 2000.

950 Johnson, D. W. and Curtis, P. S.: Effects of forest management on soil C and N storage:
 951 meta-analysis. *Forest Ecol. Manag.* 140: 227–238, 2001.

952 Karlton, E., Harrison, A. F., Alriksson, A., Bryant, C., Garnett, M. H., and Olsson, M. T.: Old
 953 organic carbon in soil solution DOC after afforestation – evidence from C-14 analysis.
 954 *Geoderma* 127: 188–195, 2005.

955 Kaste, O., Austnes, K., Vestgarden, L.S., and Wright, R. F.: Manipulation of snow in small
 956 headwater catchments at Storgama, Norway: Effects on leaching of inorganic nitrogen.
 957 *Ambio* 37: 29–37, 2008.

958 Kļaviņa, L., Bikovens, O., Šteinberga, I., Maksimova, V., and Eglīte, L.: Characterization of
 959 chemical composition of some bryophytes common in Latvia. *Environ. Exp. Biol.* 10: 27–
 960 34, 2012.

961 [Laiho, R.: Decomposition in peatlands: Reconciling seemingly contrasting results on the](#)
 962 [impacts of lowered water levels. *Soil Biol. Biochem.* 638: 2011–2024, 2006.](#)

963 [Landgraf, D., Leinweber, P., and Makeschin, F.: Cold and hot water-extractable organic](#)
 964 [matter as indicators of litter decomposition in forest soils. *J. Plant Nutr. Soil Sci.* 169: 76–](#)
 965 [82, 2006.](#)

966 Leavitt, S. W., Follett, R. F., and Paul, E. A.: Estimation of slow- and fast-cycling soil
 967 organic carbon pools from 6N HCl hydrolysis. *Radiocarbon* 38: 231–239, 1996.

968 Perez, G., Aubert, M., Decaens, T., Trap, J., and Chauvat, M.: Home-Field Advantage: A
 969 matter of interaction between litter biochemistry and decomposer biota. *Soil Biol.*
 970 *Biochem.* 67: 245–254, 2013.

giacomo 27/9/y 23:25

Eliminato: Kopittke, G. R., Tietema, A., van Loon, E. E., and Kalbitz, K.: The age of managed heathland communities: implications for carbon storage? *Plant Soil* 369: 219–230, 2013. .

975 Rosberg, I., Ovstedal, D. O., Seljelid, R., Schreiner, O., and Goksoyr, J.: Estimation of
 976 carbon flow in a Calluna Heath System. Oikos 37: 295–305, 1981.

977 Safarik, I., and Santrucková, H.: Direct determination of total soil carbohydrate content. Plant
 978 Soil 143: 109–114, 1992.

979 Scheffer, R. A., van Logtestijn, R. S. P., and Verhoeven, J. T. A.: Decomposition of Carex
 980 and Sphagnum litter in two mesotrophic fens differing in dominant plant species. Oikos
 981 92: 44–54, 2001.

982 Skjemstad, J. O., Clarke, P., Taylor, J. A., Oades, J. M., and Newman, R. H.: The removal of
 983 magnetic-materials from surface soils – A solid-state C-13 Cp/Mas nmr-study. Aust. J.
 984 Soil Res. 32: 1215–1229, 1994.

985 Soil Survey Staff: Keys to Soil Taxonomy. 11th Edition. USDA Natural Resources
 986 Conservation Service. Washington, DC., 2010.

987 Strand, L. T., Abrahamsen, G., and Stuanes, A. O.: Leaching from organic matter-rich soils
 988 by rain of different qualities: I. Concentrations. J. Environ. Qual. 31: 547–556, 2002.

989 Strand, L. T., Haaland, S., Kaste, Ø., and Stuanes, A. O.: Natural variability in soil and runoff
 990 from small headwater catchments at Storgama, Norway. Ambio 37: 18–28, 2008.

991 Stuanes, A. O., de Wit, H. A., Hole, L. R., Kaste, O., Mulder, J., Riise, G., and Wright, R. F.:
 992 Effect of climate change on flux of N and C: Air-land-freshwater-marine links: Synthesis.
 993 Ambio 37: 2–8, 2008.

994 Taylor, K., Rowland, A. P., and Jones, H. E.: *Molinia caerulea* (L.) Moench. J. Ecol. 89:
 995 126–144, 2001.

996 Trumbore, S.: Age of soil organic matter and soil respiration: Radiocarbon constraints on
 997 belowground C dynamics. Ecol. Appl. 10: 399–411, 2000.

998 Trumbore, S.: Radiocarbon and soil carbon dynamics. Annu. Rev. Earth Pl. Sci. 37: 47–66,
 999 2009.

Line Tau Strand 17/9/y 14:55
Eliminato: c
Line Tau Strand 17/9/y 14:55
Eliminato: h
Line Tau Strand 17/9/y 14:55
Eliminato: s
giacomo 24/9/y 03:40
Eliminato: t
Line Tau Strand 22/9/y 19:49
Formattato ... [49]
Line Tau Strand 22/9/y 19:49
Formattato ... [50]
Line Tau Strand 22/9/y 19:49
Formattato ... [51]
giacomo 6/10/y 22:19
Eliminato:
giacomo 24/9/y 03:40
Eliminato: s
Line Tau Strand 22/9/y 19:49
Formattato ... [52]
Line Tau Strand 22/9/y 19:49
Formattato ... [53]
Line Tau Strand 22/9/y 19:49
Formattato ... [54]
giacomo 6/10/y 22:20
Eliminato:
Line Tau Strand 18/9/y 08:59
Formattato ... [55]
giacomo 24/9/y 03:33
Spostato in giù [5]: A. P.
giacomo 24/9/y 03:33
Spostato (inserimento) [5] ... [56]
giacomo 24/9/y 03:33
Spostato in giù [6]: H. E.
giacomo 24/9/y 03:33
Eliminato: .
giacomo 24/9/y 03:33
Spostato (inserimento) [6] ... [57]
giacomo 24/9/y 03:36
Eliminato: ournal of
Line Tau Strand 18/9/y 08:59
Formattato ... [58]
giacomo 24/9/y 03:36
Eliminato: ogy
Line Tau Strand 18/9/y 08:59
Formattato ... [59]
Line Tau Strand 18/9/y 08:59
Formattato ... [60]
giacomo 24/9/y 03:34
Eliminato: -
giacomo 24/9/y 03:34
Eliminato: .
Line Tau Strand 18/9/y 08:59
Formattato ... [61]
Line Tau Strand 18/9/y 08:59
Formattato ... [62]

1014 van Vuuren, M. M. I. and Berendse, F.: Changes in soil organic-matter and net nitrogen
 1015 mineralization in heathland soils, after removal, addition or replacement of litter from
 1016 *Erica-tetralix* or *Molinia-caerulea*. Biol. Fert. Soils 15: 268–274, 1993.

1017 van Vuuren, M. M. I. and van der Eerden, L. J.: Effects of 3 rates of atmospheric nitrogen
 1018 deposition enriched with n-15 on litter decomposition in a heathland. Soil Biol. Biochem.
 1019 24: 527–532, 1992.

1020 Verhoeven, J. T. A. and Toth, E.: Decomposition of carex and sphagnum litter in fens → effect
 1021 of litter quality and inhibition by living tissue-homogenates. Soil Biol. Biochem. 27: 271–
 1022 275, 1995.

1023 Vestgarden, L. S. and Austnes, K.: Effects of freeze-thaw on C and N release from soils
 1024 below different vegetation in a montane system: a laboratory experiment, Glob. Chang.
 1025 Biol. 15: 876–887, 2009.

1026 Vestgarden, L. S., Austnes, K., and Strand, L. T.: Vegetation control on DOC, DON and DIN
 1027 concentrations in soil water from a montane system, southern Norway. Boreal Environ.
 1028 Res. 15: 565–578, 2010.

1029 von Lützow, M., Kögel-Knabner, I., Ekschmitt, K., Flessa, H., Guggenberger, G., Matzner,
 1030 E., and Marschner, B.: SOM fractionation methods: Relevance to functional pools and to
 1031 stabilization mechanisms. Soil Biol. Biochem. 39: 2183–2207, 2007.

1032 Wang, Q. K., Zhong, M., He, T.: Home-field advantage of litter decomposition and nitrogen
 1033 release in forest ecosystems. Biol. Fert. Soils 49: 427–434, 2013.

1034 Wickland, K. P., Neff, J. C., and Harden, J. W.: The role of soil drainage class in carbon
 1035 dioxide exchange and decomposition in boreal black spruce (*Picea mariana*) forest
 1036 stands. Can. J. Forest Res. 40: 2123–2134, 2010.

1037 ▼

giacomo 8/10/y 08:28

Formattato: Tipo di carattere:Corsivo

giacomo 8/10/y 08:28

Formattato: Tipo di carattere:Corsivo

giacomo 8/10/y 08:28

Eliminato: -

giacomo 8/10/y 08:28

Eliminato: ,

giacomo 8/10/y 08:28

Eliminato: ,

giacomo 8/10/y 08:29

Formattato: Tipo di carattere:Corsivo

giacomo 29/9/y 09:12

Eliminato: Wieder, R. K. and Starr, S. T.: Quantitative determination of organic fractions in highly organic, Sphagnum peat soils. Commun. Soil Sci. Plant An. 29: 847–857, 1998. .

1045 **Figures captions:**

1046

1047 **Figure 1. a)** A general view of the study area, Storgama, showing soil occurring in pockets
1048 and small depressions at the bedrock surface; note that close up vegetation at the bottom right
1049 is dominated by *Molinia caerulea* (L), the understorey of pines beyond is *Calluna vulgaris*
1050 (L) Hull, the basin in the background is covered by *Sphagnum* spp. **b)** A rare coalescence of
1051 the three dominant species, *Calluna vulgaris*, on the left, *Sphagnum* spp. L, at the bottom,
1052 and *Molinia caerulea*, on the right.

1053

1054 **Figure 2.** ¹³C CPMAS NMR spectra of the aboveground biomass of the dominant plant
1055 species and the underlying soil.

1056

1057 **Figure 3.** Residual mass in the litterbags as a function of time for different combinations of
1058 litter and vegetation cover. Cal in Cal means *Calluna* litter decomposing under *Calluna*, Cal
1059 in Mol means *Calluna* litter decomposing under *Molinia*, and so on. Lower case letters
1060 indicate significant differences (p<0.05) between same litters decomposing under different
1061 types of vegetation. The trial was one year long.

1062

1063 **Figure 4.** Carbon and nitrogen concentrations and C/N ratio in decaying biomass in the
1064 litterbags as a function of time for different combinations of litter and vegetation cover. Cal
1065 in Cal means *Calluna* litter decomposing in soil under *Calluna*, Cal in Mol means *Calluna*
1066 litter decomposing under *Molinia*, and so on. Upper case letters indicate significant
1067 differences (p<0.05) between different litters, whereas lower case letters indicate significant
1068 differences between same litters decomposing in soils covered by different types of
1069 vegetation.

Line Tau Strand 22/9/y 21:42

Eliminato: buried

Line Tau Strand 22/9/y 21:43

Eliminato: in soil

Line Tau Strand 22/9/y 21:43

Eliminato: in soil

Line Tau Strand 22/9/y 21:44

Eliminato: in soil

giacomo 6/10/y 18:38

Eliminato: of the trial of Figure 3

Line Tau Strand 22/9/y 21:44

Eliminato: in soil