

SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

This discussion paper is/has been under review for the journal SOIL. Please refer to the corresponding final paper in SOIL if available.

Interactions between organisms and parent materials of a constructed Technosol shape its hydrostructural properties

M. Deeb^{1,2}, M. Grimaldi², T. Z. Lerch¹, A. Pando^{1,2}, A. Gigon¹, and M. Blouin¹

¹UPEC, Institute of Ecology and Environmental Sciences of Paris (UMR 7618),
61 avenue du Général de Gaulle, 94010 Créteil, France

²IRD, Institute of Ecology and Environmental Sciences of Paris (UMR 7618),
32 avenue Henri Varagnat, 93142 Bondy CEDEX, France

Received: 25 October 2015 – Accepted: 27 November 2015 – Published: 17 December 2015

Correspondence to: M. Deeb (mahadeeb.y@gmail.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



little understood (Paton, 1978; Amundson et al., 2007). The influence of bioturbation (physical displacement by organisms) on soil formation is not straightforward (Amundson et al., 2007; Wilkinson et al., 2009). Some authors consider biotic mixing agents as a secondary cause of soil formation (Carson and Kirkby, 1972), while others argue that bioturbation plays a major role in forming soil (Paton, 1978; Wilkinson and Humphreys, 2005).

Soils developed on non-traditional substrates and largely influenced by human activity are now referenced as Technosols in the World Reference Base for Soil Resources. When Technosols are technogenic materials or artifacts assembled deliberately to create soils, they are called constructed Technosols (IUSS Working Group WRB, 2015). Many urban planners and greenspace enterprises are interested in constructed Technosols because they are an alternative to topsoil uptake from the countryside, which can be costly and harmful for the environment due to CO₂ emitted during material transport. Moreover, Technosols offer an opportunity to recycle urban wastes, such as excavated deep horizons/backfills from enterprises of the building sector, sewage sludge from waste water plants or green wastes from greenspaces enterprises or local authorities. In this way, urban wastes are used to improve urban ecosystem services (Morel et al., 2014), and form a closed loop that reduces the impact of cities on the environment. Constructed Technosols are different from other soils, because they are designed assemblages of technogenic materials. The evolution of Technosols is thus different from the pedogenesis of a natural soils. However, Technosols exhibit some formation processes similar to those observed in natural soil pedogenesis, such as decarbonization and aggregation (Séré et al., 2010; Jangorzo et al., 2014). Humans are an agent with an increasing importance in the evolution of the Biosphere. This has led to consider that we are living on a planet dominated by humans (Vitousek et al., 1997) and to define our geological era as the Anthropocene (Zalasiewicz et al., 2008). In soil science too, this importance of humans has also to be acknowledged. In this paper, we will thus consider the evolution of constructed

SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

aggregation and aeration (Blanchart et al., 1997), which influence the hydric properties of soil (Schrader and Zhang, 1997; Shipitalo and Butt, 1999). These modifications of hydro-structural properties by earthworms have tremendous consequences for plant growth (Scheu, 2003; van Groenigen et al., 2014). Plant roots and rhizosphere inhabitants also have a significant influence on aggregates and their stability (Jastrow et al., 1998; Rillig et al., 2002), sometimes more significant than that of earthworms (Blanchart et al., 2004). Roots penetrate the soil and create macropores which favor fluid transport (Beven and Germann, 1982). Roots also create weak zones that fragment the soil and form aggregates, whose formation is strengthened by wetting-drying cycles due to water uptake by the plant (Angers and Caron, 1998). In addition, plant root residues provide a food source for microorganisms and fauna, which contribute to soil structure formation and stabilization (Innes et al., 2004). In return, microorganism-mediated changes in soil structure affect plant growth, mostly by modifying the root's physical environment (Dorioz et al., 1993).

In this study, we were interested in the effect of two soil-forming factors, i.e. parent materials and organisms, on hydro-structural parameters via measurements of soil shrinkage curves (SSC) (Haines, 1923). The influence of parent materials (especially clay content and type) (Boivin et al., 2004), organic matter (Boivin et al., 2009) and organisms (Kohler-Milleret et al., 2013; Milleret et al., 2009) on shrinkage properties has already been studied in natural soils. This study addresses the question of materials-organisms interaction on the hydrostructural properties of a constructed Technosols in a five-months microcosm experiment with four “organism” treatments (control, plants, earthworms, plants and earthworms) combined with six percentages of green waste compost/excavated deep horizons under controlled climatic conditions.

2 Materials and methods

2.1 Parent materials

The main urban waste used in this study was earth material excavated from deep horizons, hereafter called “excavated deep horizons” (EDH). This material is typically what is found during the foundation works in the Ile-de-France. It is mainly made of calcareous strata from the Eocene. For our study, we sampled 500 kg of EDH at several locations at the basis of the urban waste dump of the ECT company (Villeneuve-sous-Dammartin, France) in the aim to have a composite sample representative of what is susceptible to be used for Technosol construction around Paris. To avoid the presence of calcareous blocks of big size, this material was sieved at 4 mm. Carbonate was 33 % ($w : w$) of total mass of the EDH fraction < 4 mm. After carbonate removing in the aim to measure the granulometry, remaining material was made of 2 % clay, 10 % silt and 88 % sand. Without removing carbonate, it had 11 % of particles < 2 μm , 30 % from 2 to 50 μm , and 59 % from 50 μm to 2 mm, indicating that carbonates were mainly in the finer particle class. Powder X-ray diffraction performed with a Siemens D500 (Cu-K α , 40 kV, 30 mA) diffractometer identified quartz, calcite and dolomite. EDH had very low levels of total organic carbon (0.38 %) and total nitrogen (0.035 %). They had a basic $\text{pH}_{\text{H}_2\text{O}}$ of 8.3 and a pH_{KCl} of 8.2. Cation-exchange capacity was as low as $3.12 \text{ cmol}^+ \text{ kg}^{-1}$. We measured a particle density of 2.75 g cm^{-3} , relatively high compared to that of natural soils, and a bulk density of 1.33 g cm^{-3} . Green waste compost (GWC) was retrieved from the Biodepe company (Ahuy, France). It contained 21.41 % total organic carbon, 1.47 % total nitrogen, a $\text{pH}_{\text{H}_2\text{O}}$ of 7.93 and a pH_{KCl} of 7.44, with a particle density of 2.06 g cm^{-3} and a bulk density of 0.61 g cm^{-3} . As EDH, GWC was sieved at 4 mm to avoid the presence of big wood pieces in the microcosm.

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Experimental design and conditions

EDH and GWC were mixed using a concrete mixer for ten minutes to prepare six different mixtures with specific volumetric percentages of GWC at 0, 10, 20, 30, 40, and 50%. One liter of each mixture was placed in a microcosm of 13 × 13 × 12.5 cm with full capacity of 1.2 L. Water retention capacity of each mixture was measured at the beginning of the experiment by using a pressure plate apparatus (Richards, 1948) with a water potential of -31 kPa. During the experiment, microcosms were moistened two to three times a week with deionized water to maintain soil moisture at 80 % of field capacity for each mixture (Table S1 in the Supplement).

Plants were sown 24 h after watering the pots, and earthworms were introduced 24 h after sowing. Each percentage of GWC was combined with four treatments: a control without organisms (C), a treatment with two individuals (0.5 ± 0.1 g each) of the endogeic earthworm species *Aporrectodea caliginosa* (E), a treatment with *Lolium perenne* plants (50 seeds with a 80 % germination rate scattered homogeneously on the microcosm surface) (P), and a treatment with both earthworms and plants (EP). In total, 96 microcosms were divided into 24 treatments, each with four replicates.

Microcosms were kept 21 weeks in a climate chamber (S10H, Conviron, Canada) under the following conditions: photoperiod of 12 h, luminosity of $500 \pm 20 \mu\text{mol photons m}^{-2} \text{s}^{-1}$, temperature at $22/20 \pm 0.2^\circ\text{C}$ day/night respectively and $75 \pm 2\%$ air humidity.

2.3 Shrinkage analysis

Technosol samples were collected from the surface of each microcosm at the end of experiment using a 5 cm high, 5 cm diameter cylinder and were placed on a wet porous plate for saturation with deionized water according to manufacturer instructions (Sandbox, Eijkelkamp, Netherlands) for seven days by applying a water potential of 0 kPa at the base of the sample. The shrinkage curve was continuously measured according to Braudeau et al. (1999) by using the retractometer[®] apparatus. Water-

SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



saturated Technosol samples were placed in an oven at a constant temperature (30 °C) to provide continuous and rapid evaporation. An electronic scale (0.01 g precision) ensured accurate measurement of water loss during drying. Each sample's volume (diameter, height) was determined with laser beams and recorded along with its mass every 10 min.

At the end of the measurement, samples were dried in an oven at 105 °C for 48 h to measure dry mass and bulk density. These data were converted into soil specific volume (V , $\text{cm}^3_{\text{soil}} \text{g}_{\text{dry soil}}^{-1}$) and water content (W , $\text{g}_{\text{water}} \text{g}_{\text{soil}}^{-1}$). We then determined the soil shrinkage curve (SSC) to describe hydro-structural properties, as proposed by Assi et al. (2014). The data obtained by shrinkage measures were fitted according to the pedostructure model (Braudeau et al., 2004). In this model, the SSC is subdivided into a maximum of four shrinkage phases (interpedal/saturated (ip), structural (st), basic (bs) and residual (re) shrinkage phases) that are due to four types of water (W_{ip} , W_{st} , W_{bs} , W_{re}) (Fig. 1). The pedostructure is considered an assembly of primary peds (aggregates made of the clayey particles) that determines two nested levels of organization: the macropore level (containing $W_{\text{ma}} = W_{\text{ip}} + W_{\text{st}}$) and the micropore level (containing $W_{\text{mi}} = W_{\text{re}} + W_{\text{bs}}$). These levels do not refer to pore size by itself, but to water pore behavior during soil drying. Based on this distinction, the two pore systems were also called plasma (micropores) and structural properties (macropores) (Boivin et al., 2004; Schäffer et al., 2008).

The three transition points separating the four pseudo linear shrinkage phases (Fig. 1) are points L, M and N, which are at the intersection of the tangent straight lines of the linear phases. According to this model of SSC (Braudeau et al., 1999b, 2004), the value of the water content at each point is equal to the value of $\max(W_{\text{st}})$ for W_{L} , $\max(W_{\text{mi}}) = \max(W_{\text{re}}) + \max(W_{\text{bs}})$ for W_{M} , and $\max(W_{\text{re}})$ for W_{N} . The other hydro-structural parameters are: slope of the saturated phase (K_{ip}), slope of the structural phase (K_{st}), slope of the basic shrinkage phase (K_{bs}), slope of the residual phase (K_{re}), and three parameters (k_{L} , k_{M} and k_{N}) related to the SSC shape at points L, M

SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and N , respectively. Finally, according to Braudeau et al. (2001):

$$\text{Max}(W_{re}) = W_N \quad (1)$$

$$\text{Max}(W_{bs}) = W_M - W_N \quad (2)$$

$$\text{Max}(W_{st}) = W_L - W_M \quad (3)$$

Specific volume V as a function of the water content W obtained from the Braudeau model was converted into a void ratio, (e , $\text{cm}^3_{\text{pore}} \text{cm}^{-3}_{\text{solid}}$) as a function of the moisture ratio (v , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{solid}}$). This step makes it easier to compare Technosols that have different compositions and thus different particle densities. Considering Eqs. (4) and (5):

$$v = (\rho_s / \rho_w) W \quad (4)$$

$$e = V \rho_s - 1 \quad (5)$$

With ρ_w the water density and ρ_s the particle density (g cm^{-3}) calculated for all mixtures from measurements of GWC and EDH using a pycnometer on materials sieved at 2 mm (ISO 17892-3:2004).

All hydro-structural parameters were transformed with Eqs. (4) and (5) and thus became the moisture ratio at macropore saturation (v_L), the moisture ratio at micropore saturation (v_M), the moisture ratio at the shrinkage limit (v_N), the four slopes (K_L , K_{st} , K_{bs} , K_{re}), parameters related to the SSC shape (k_L , k_M , k_N) and the void ratio at the end of the shrinkage period (e_0).

Considering these hydro-structural parameters (Braudeau et al., 2004), the ratio of the maximum available water for plants from macropores (v_{ma} , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{solid}}$) and the ratio of the maximum available water for plants from micropores (v_{mi} , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{solid}}$) can be calculated from the Eqs. (2) and (3) as follows: $v_{ma} = v_L - v_M$, $v_{mi} = v_M - v_N$. The sum of both is the total moisture ratio (v_{Total} in $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{solid}}$).

SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Finally, volumetric water content (Θ , $\text{cm}^3_{\text{water}} \text{cm}^3_{\text{soil}}$) was calculated to compare available water reservoirs (holding capacities) for plants.

$$\Theta = v(\rho_d/\rho_s) = v(\rho_d/\rho_s) \quad (6)$$

With ρ_d the bulk density ($\text{g}_{\text{solid}} \text{cm}^{-3}_{\text{soil}}$). Similarly, we calculated the volumetric water content from macropores (θ_{ma}) and micropores (θ_{mi}), by applying the following equations: $\theta_{\text{ma}} = \theta_L - \theta_M$, $\theta_{\text{mi}} = \theta_M - \theta_N$, and eventually the sum of both known as the total volumetric water content for plants (θ_{Total}).

2.4 Plant harvest and root size distribution

Plants were cut at the soil surface 21 weeks after sowing. Fresh leaves were weighed, dried in an oven at 50°C for two days and weighed again. Root mass was estimated from one quarter of the pot, since other quarters were used for physico-chemical and shrinkage analyses, which requires not disturbing soil physical characteristics (i.e. root or earthworm sampling).

Dry root biomass distribution among diameter classes was determined according to the method of Blouin et al. (2007). It is based on the granulometry method used to assess soil texture: roots are dried, cut transversely with a mixer and placed on a column of sieves with decreasing mesh size. During the shaking of the sieve column, root fragments with a section diameter smaller than the mesh size pass through this mesh and stop on the first sieve with a mesh size below that of the root section diameter. Biomass distribution is assessed by weighing the biomass recovered in each sieve. Five diameter classes were chosen according to sieve mesh size: 0–100, 100–200, 200–400, 400–800 and $> 800 \mu\text{m}$.

2.5 Data analysis

We calculated means and standard errors of hydro-structural parameters for all treatments by fitting the curves with the hydro-structural model (Table S2). The

SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a positive effect on belowground biomass only at 50 % GWC, whereas aboveground biomass was affected only in the 0–30 % GWC range. As a result total biomass was always significantly higher in the presence of earthworms, except at 40 % GWC. In overall, earthworms increased total plant biomass of 21 % (Fig. 2c). The best treatment for plant growth was clearly the mixture of 50 % GWC with earthworms, with a total dried plant biomass of 8.07 g, which was significantly higher than that of all other mixtures, except for 40 % GWC with earthworms. There was no interaction between the effects of GWC and earthworms on plant biomasses, which means that these two effects are additive. All parameters describing biomass allocation inside the plant, such as the root : shoot ratio, the thick ($\geq 400 \mu\text{m}$) and fine ($< 400 \mu\text{m}$) root percentages, were not affected by the presence of GWC, earthworms or their interaction (Table 1); we thus concluded that GWC percentage and presence of earthworms had a quantitative influence but not a qualitative one, as growth was affected but not development.

3.2 Specific influence of organisms and parent materials on hydrostructural parameters

All our Technosols exhibited the classical sigmoid shape of the shrinkage curve reported for most natural soils (Laurizen, 1948; Braudeau et al., 1999; Peng and Horn, 2005) (Figs. 3, 4); thus, shrinkage phases (residual, basic, structural and the saturating shrinkage phase) were easy to recognize. All the parameters deduced from SSC are given in Table S2.

High GWC percentage caused moisture ratio v and void ratio e to increase (Fig. 3). The positive effect of GWC was particularly important in treatments with plants at 50 % GWC and in treatment with earthworms and plants at 40 and 50 % (Fig. 3). Earthworms had a positive influence on the void ratio in the 0–30 % GWC range, but this positive effect disappeared at 40 and 50 % GWC (Fig. 4). The influence of plants on void ratio was positive for at 10, 20, 30 and 50 % GWC but not at 0 and 40 % GWC (Fig. 4). The simultaneous presence of plants and earthworms resulted in a positive effect on void ratio for all mixtures (Fig. 4). For example e_0 varied from 0.91–

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



that predicting variations in hydro-structural parameters of our Technosols requires taking into account variation in parent materials and organisms simultaneously.

The LDA explained 76% of hydro-structural properties observed variance ($P < 0.001$; Wilks and Pillai tests) (Fig. 5). Axis 1, which explained 42% of the total variance, distinguished treatment “earthworms” from treatment “earthworms and plants” whereas axis 2, which explained 26% of the total variance, separated the “control” and the “plants” treatments. By relating the correlation circle (Fig. 5a) to the factorial plan (Fig. 5b) we found that: (i) the parameter related to the shape of shrinkage curves between interpedal and structural phases (K_L) was higher for the control than for organism treatments; (ii) earthworms increased moisture ratio at the shrinkage limit (v_N); (iii) plants increased the slope of the structural phase (K_{st}); (iv) the simultaneous presence of plants and earthworms increased the moisture ratio at saturated macropores (v_L), minimum void ratio (e_0), and a parameter related to the shape of shrinkage curves (K_N).

Additional PCA were performed to characterize the effect of organisms on hydro-structural properties for each GWC percentage. The effect of plants was not significant at 0, 10, and 20% GWC ($P > 0.05$, Monte Carlo test), while it was significant at 30, 40, and 50% of GWC ($P < 0.05$, Monte Carlo test). In contrast, combined influences of plants and earthworms were always significant ($P < 0.05$, Monte Carlo test).

3.4 Influence of organisms and parent materials on moisture ratio and available water for plants

Hydro-structural parameters (v_L , v_M , v_N) provide information about moisture ratio and available volumetric water content in soil. The influences of GWC percentage and the presence/absence of earthworms and plants on macropore, micropore and total moisture ratios, as well as micropore, macropore and total available volumetric water contents were tested in a three-way ANOVA. The complete model with GWC percentage, earthworms and plants had a significant effect ($P < 0.001$) on micropore, macropore and total moisture ratios and available volumetric water contents (Table 2).

SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Considering single factors, increasing the GWC percentage had a positive influence on micropore, macropore (GWC < 40 %) and total moisture ratios and available volumetric water contents ($P < 0.001$). Plants had an influence on all of the previous variables, except for micropore volumetric available water content. Earthworms affected micropore and total moisture ratios, but not the macropores moisture ratio; they affected only micropore volumetric available water content (Table 2).

Regarding interactions between factors, the influence of GWC on moisture ratio and volumetric available water content depended on the presence of plants for micropore, macropore and total volumetric available water content, on earthworms for macropore and total volumetric available water content and the interaction between plants and earthworms for total volumetric available water content (Table 2). The presence of earthworms influenced the effect of GWC on moisture ratio and total volumetric available water contents at macropore and micropore. For example, in the absence of earthworms, GWC had a positive influence on moisture ratio at macropore for 0–40 % GWC, while in the presence of earthworms, moisture ratio at macropore decreased at percentages of 30–50 %. The presence of plants modified the influence of GWC on moisture ratios at micropore and macropore, and total volumetric available water at macropore and micropore. For example, in the absence of plants, the influence of GWC on moisture ratio at macropore was positive at percentages of 0–40 % and became negative at 50 %, whereas in the presence of plants, the influence of GWC was positive regardless of its percentage (Fig. 4a). A similar influence was observed for the interaction between plants and GWC on macropore volumetric available water (Fig. 6d). The interaction between earthworms and plants had a significant effect only for moisture ratios in micropore and macropore but not for total moisture ratio, suggesting opposite effect on micropores and macropores (Table 2). Indeed, v_{ma} was higher in the plants and earthworms treatment as compared with the plant treatment and the earthworm treatment, but v_{mi} was higher in the earthworm treatment or the plant treatment as compared with the plants and earthworms treatment. The triple interaction had a significant influence on moisture ratio and volumetric available water

SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in pedogenesis (Jenny, 1941; Carson and Kirkby, 1972) versus those stressing their importance (Paton, 1978; Wilkinson and Humphreys, 2005). Indeed, if the influence of organisms is particularly important in interaction with parent materials, its observation may be random. Pedogenesis, especially Technosol pedogenesis, thus appears as an interdisciplinary field of study that needs to include ecological. We found that biological activity improved Technosol properties by increasing aggregation, porosity and water-retention capacity, with potential consequences on resistance to drought and erosion. An original research perspective could be to investigate benefits of these changes caused by plants and earthworms for their own survival and reproduction to determine if these biological activities increase the fitness of these organisms and could thus be considered as niche construction (Odling-Smee et al., 1996).

5 Conclusions

In a nutshell, we found that compost and plants play a positive role in macroporosity and microporosity in Technosols, while earthworms affect only microporosity. GWC positively affected macroporosity up to a percentage of 30% and plants were responsible for extending this positive influence at 40 and 50% GWC. The simultaneous presence of earthworms and plants was responsible for a synergistic positive influence on macroporosity. These observations highlighted the need to consider plants not only as an output indicating the level of fertility, but also as an actor in Technosol construction, in the same way than earthworms. Organisms that physically modify their environment by creating, destroying or maintaining ecological niches have been called “ecosystem engineers” (Jones et al., 1994). These ecosystem engineers can help restore ecosystems (Byers et al., 2006) and create new ecosystems such as constructed Technosols by assisting managers, who could “sub-contract” one aspect of management. Therefore, instead of increasing the amount of compost, which is usually expensive, managers could avoid the difficult-to-explain negative influence of high percentages of compost by favoring conservation, recolonization or inoculation

of ecosystem engineers such as plants and earthworms, especially in combination (Blouin et al., 2013).

**The Supplement related to this article is available online at
doi:10.5194/-15-1309-2015-supplement.**

5 *Acknowledgements.* This study was conducted in collaboration with the Departmental Council
of the Seine-Saint-Denis department, France, and the company Enviro Conseil et Travaux.
The authors wish to thank the University of Damas, Syria, for financial support via a Ph.D.
scholarship. We also thank Thierry Desjardins, Gaghik Hovhannissian and Pascal Podwojewski
for their scientific advice and Florence Dubs for her help with statistical analyses. Michael
10 Corson was responsible for post-editing the English.

References

- Amundson, R., Richter, D. D., Humphreys, G. S., Jobbágy, E. G., and Gaillardet, J.:
Coupling between biota and earth materials in the critical zone, *Elements*, 3, 327–332,
doi:10.2113/gselements.3.5.327, 2007.
- 15 Angers, D. A. and Caron, J.: Plant-induced changes in soil structure: Processes and feedbacks,
Biogeochemistry, 42, 55–72, doi:10.1023/A:1005944025343, 1998.
- Arocena, J. M., van Mourik, J. M., Schilder, M. L. M., and Faz Cano, A.: Initial soil development
under *Pioneer* plant species in metal mine waste deposits, *Restor. Ecol.*, 18, 244–252,
doi:10.1111/j.1526-100X..2009.00582.x, 2010.
- 20 Assi, A. T., Braudeau, E. F., Accola, J. J. O., Hovhannissian, G., and Mohtar, R.: Physics of the
soil medium organization Part 2: Pedostructure characterization through measurement and
modeling of the soil moisture characteristic curves, *Frontiers in Environmental Science* 2, 5,
doi:10.3389/fenvs.2014.00005, 2014.
- 25 Beven, K. and Germann, P.: Macropores and water flow in soils, *Water Resour.*, 18, 1311–1325,
doi:10.1029/WR018i005p01311, 1982.

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Blanchart, E., Lavelle, P., Braudeau, E., Le Bissonnais, Y., and Valentin, C.: Regulation of soil structure by geophagous earthworm activities in humid savannas of Côte d'Ivoire, *Soil Biol. Biochem.*, 29, 431–439, doi:10.1016/S0038-0717(96)00042-9, 1997.

Blanchart, E., Albrecht, A., Chevallier, T., and Hartmann, C.: The respective roles of roots and earthworms in restoring physical properties of Vertisol under a *Digitaria decumbens* pasture (Martinique, WI), *Agr. Ecosyst. Environ.*, 103, 343–355, doi:10.1016/j.agee.2003.12.012, 2004.

Blouin, M., Hodson, M. E., Delgado, E. A., Baker, G., Brussaard, L., Butt, K. R., Dai, J., Dendooven, L., Peres, G., Tondoh, J. E., Cluzeau, D., and Brun, J.-J.: A review of earthworm impact on soil function and ecosystem services, *Eur. J. Soil Sci.*, 64, 161–182, doi:10.1111/ejss.12025, 2013.

Blouin, M., Lavelle, P., and Laffray, D.: Drought stress in rice (*Oryza sativa* L.) is enhanced in the presence of the compacting earthworm *Millsonia anomala*, *Environ. Exp. Bot.*, 60, 352–359, doi:10.1016/j.envexpbot.2006.12.017, 2007.

Braudeau, E., Costantini, J. M., Bellier, G., and Colleuille, H.: New device and method for soil shrinkage curve measurement and characterization, *Soil Sci. Soc. Am. J.*, 63, 525, doi:10.2136/sssaj1999.03615995006300030015x, 1999.

Braudeau, E., Frangi, J. P., and Mohtar, R. H.: Characterizing nonrigid aggregated soil-water medium using its shrinkage curve, *Soil Sci. Soc. Am. J.*, 68, 359–370, 2004.

Byers, J. E., Cuddington, K., Jones, C. G., et al: Using ecosystem engineers to restore ecological systems, *Trends Ecol. Evol.*, 21, 493–500, doi:10.1016/j.tree.2006.06.002, 2006.

Caron, J., Espindola, C. R., and Angers, D. A.: Soil structural stability during Rapid Wetting: Influence of land use on some aggregate Properties, *Soil Sci. Soc. Am. J.*, 60, 901–908, doi:10.2136/sssaj1996.03615995006000030032x, 1996.

Carson, M. A. and Kirkby, M. J.: Hillslope form and process, Cambridge University Press, New York, 1972.

Charman, P. E. V. and Murphy, B. W.: Soil conservation service of New South Wales: Soils?: their properties and management, edited by: Charman, P. E. V. and Murphy, B. W., Oxford University Press, Melbourne; Oxford, 2000.

Deeb, M., Grimaldi, M., Lerch, T. Z., Pando, A., Podwojewski, P., and Blouin, M.: Effect of organic content on the water retention and shrinkage properties of Constructed Technosols, *Pedosphere*, accepted, 2015.

SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Dorioz, J. M., Robert, M., and Chenu, C.: The role of roots, fungi and bacteria on clay particle organization. An experimental approach, *Geoderma*, 56, 179–194, doi:10.1016/0016-7061(93)90109-X, 1993.
- 5 Dray, S. and Dufour, A. B.: The ade4 package: implementing the duality diagram for ecologists, *J. Stat. Softw.*, 22, 1–20, 2007.
- Edwards, C. A: *Earthworm ecology*, CRC Press, USA, 20 pp., 2004.
- Eisenhauer, N., Partsch, S., Parkinson, D., and Scheu, S.: Invasion of a deciduous forest by earthworms: Changes in soil chemistry, microflora, microarthropods and vegetation, *Soil. Biol. Biochem.*, 39, 1099–1110, doi:10.1016/j.soilbio.2006.12.019, 2007.
- 10 Haines, W. B: The volume-changes associated with variations of water content in soil, *J. Agric. Sci.*, 13, 296–310, doi:10.1017/S0021859600003580, 1923.
- Innes, L., Hobbs, P. J., and Bardgett, R. D.: The impacts of individual plant species on rhizosphere microbial communities in soils of different fertility, *Biol. Fert. Soils*, 40, 7–13, doi:10.1007/s00374-004-0748-0, 2004.
- 15 IUSS Working Group WRB: World Reference Base for Soil Resources 2014, update 2015 International soil classification system for naming soils and creating legends for soil maps, World Soil Resources Reports No. 106, FAO, Rome, 2015.
- Jangorzo, N. S., Watteau, F., and Schwartz, C.: Evolution of the pore structure of constructed Technosols during early pedogenesis quantified by image analysis, *Geoderma*, 207–208, 180–192, doi:10.1016/j.geoderma.2013.05.016, 2013.
- 20 Jangorzo, N. S., Watteau, F., Hajos, D., and Schwartz, C.: Nondestructive monitoring of the effect of biological activity on the pedogenesis of a Technosol, *J. Soil. Sediment.*, 15, 1705–1715, doi:10.1007/s11368-014-1008-z, 2014.
- Jastrow, J. D., Miller, R. M., and Lussenhop, J.: Contributions of interacting biological mechanisms to soil aggregate stabilization in restored prairie, *Soil. Biol. Biochem.*, 30, 905–916, doi:10.1016/S0038-0717(97)00207-1, 1998.
- 25 Jenny, H.: *Factors of soil formation*, McGraw-Hill, New York, London, 1941.
- Jones, C. G., Lawton, J. H., and Shachak, M.: Organisms as ecosystem engineers, in: *Ecosystem Management*, edited by: Samson, F. B., and Knopf, F. L., Springer, New York, 130–147, 1994.
- 30 Kohler-Milleret, R., Le Bayon, R.-C., Chenu, C., Gobat, J.-M., and Boivin, P.: Impact of two root systems, earthworms and mycorrhizae on the physical properties of an unstable silt loam Luvisol and plant production, *Plant Soil*, 370, 251–265, 2013.

SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Olszewski, M. W., Holmes, M. H., and Young, C. A.: Assessment of Physical Properties and Stonecrop growth in green roof substrates amended with compost and hydrogel, *HortTechnology*, 20, 438–444, 2010.

Paradelo, R. and Barral, M. T.: Influence of organic matter and texture on the compactability of Technosols, *Catena*, 110, 95–99, doi:10.1016/j.catena.2013.05.012, 2013.

Paton, T. R.: The formation of soil material, George Allen and Unwin, London, UK, 1978.

Peng, X. and Horn, R.: Modeling soil shrinkage curve across a wide range of soil types, *Soil Sci. Soc. Am. J.*, 69, 584–592, doi:10.2136/sssaj2004.0146, 2005.

R Core Team: R: A Language and environment for statistical computing, 2014.

Reid, J. B. and Goss, M. J.: Interactions between soil drying due to plant water use and decreases in aggregate stability caused by maize roots, *J. Soil Sci.*, 33, 47–53, doi:10.1111/j.1365-2389.1982.tb01746.x, 1982.

Rillig, M. C., Wright, S. F., and Eviner, V. T.: The role of arbuscular mycorrhizal fungi and glomalin in soil aggregation: comparing effects of five plant species, *Plant Soil*, 238, 325–333, doi:10.1023/A:1014483303813, 2002.

Rokia, S., Séré, G., Schwartz, C., et al: Modelling agronomic properties of Technosols constructed with urban wastes, *Waste Manage.*, 34, 2155–2162, doi:10.1016/j.wasman.2013.12.016, 2014.

Santos, G. G., Da Silva, E. M., Marchão, R. L., Da Silveira, P. M., Bruand, A., James, F., and Becquer, T.: Analysis of physical quality of soil using the water retention curve: validity of the S-index, *C. R. Geosci.*, 343, 295–301, 2011.

Scheu, S.: Effects of earthworms on plant growth: patterns and perspectives: The 7th international symposium on earthworm ecology · Cardiff · Wales · 2002, *Pedobiologia*, 47, 846–856, doi:10.1078/0031-4056-00270, 2003.

Schrader, S. and Zhang, H.: Earthworm casting: Stabilization or destabilization of soil structure?, *Soil Biol Biochem.*, 29, 469–475, doi:10.1016/S0038-0717(96)00103-4, 1997.

Séré, G., Schwartz, C., Ouvrard, S., Renat, J.-C., Watteau, F., Villemin, G., and Morel, J.-L.: Early pedogenic evolution of constructed Technosols, *J. Soil Sediment.*, 10, 1246–1254, doi:10.1007/s11368-010-0206-6, 2010.

Shipitalo, M. and Butt, K.: Occupancy and geometrical properties of *Lumbricus terrestris* L. burrows affecting infiltration, *Pedobiologia*, 43, 782–794, 1999.

SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Tejada, M. and Gonzalez, J. L.: Effects of the application of compost originating from crushed cotton gin residues on wheat yield under dryland conditions, *Eur. J. Agron.*, 19, 357–368, doi:10.1016/S1161-0301(02)00089-8, 2003.

5 Van Groenigen, J. W., Lubbers, I. M., Vos, H. M. J., Brown, G. G., De Deyn, G. B., and van Groenigen, K. J.: Earthworms increase plant production: a meta-analysis, *Scientific Reports*, 4, 6365, doi:10.1038/srep06365, 2014.

Venables, W. N. and Ripley, B. D.: *Modern Applied Statistics with S*, Fourth Edition, Springer, New York, 2002.

10 Wilkinson, M. T. and Humphreys, G. S.: Exploring pedogenesis via nuclide-based soil production rates and OSL-based bioturbation rates, *Soil Res.*, 43, 767–779, 2005.

Wilkinson, M. T., Richards, P. J., and Humphreys, G. S.: Breaking ground: Pedological, geological, and ecological implications of soil bioturbation, *Earth-Sci. Rev.*, 97, 257–272, doi:10.1016/j.earscirev.2009.09.005, 2009.

SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Two-ways ANOVA showing the effects of the presence/absence of earthworms (E) and the proportion of green waste compost (GWC) in the mixtures on plant dry biomasses, shoot: root ratio and root system structure (thick root $\geq 400 \mu\text{m}$ and fine root $<400 \mu\text{m}$) ($n = 48$). The number in the table are the F values, significance codes: *: $P \leq 0.05$, **: $P \leq 0.01$, ***: $P \leq 0.001$, ns: $P > 0.05$.

	d.f.	Aboveground biomass (g)	Belowground biomass (g)	Totalbiomass (g)	Shoot : root ratio	Thick root proportion	Fine root proportion
Complete model	11	11.29***	5.85***	13.33***	1.27 ^{ns}	0.78 ^{ns}	0.95 ^{ns}
GWC	5	10.27***	8.73***	16.22***	2.08 ^{ns}	0.49 ^{ns}	0.72 ^{ns}
E	1	65.65***	15.24***	60.12***	0.14 ^{ns}	0.62 ^{ns}	1.59 ^{ns}
GWC \times E	5	1.43 ^{ns}	1.08 ^{ns}	0.39 ^{ns}	0.68 ^{ns}	0.56 ^{ns}	1.05 ^{ns}

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Table 2. Three ways ANOVA testing the effect of green waste compost (GWC), earthworms (E) and plants (P) on the maximum moisture ratio from macropores (v_{ma} , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{solid}}$), maximum moisture from micropores (v_{mi} , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{solid}}$), total moisture ratio (v_{Total} , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{solid}}$), macro available water (θ_{ma} , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{solid}}$), micro available water (θ_{mi} , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{solid}}$) and finally total available water (θ_{Total} , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{solid}}$) ($n = 96$). The number in the table are the F values, significance codes: *: $P \leq 0.05$, **: $P \leq 0.01$, ***: $P \leq 0.001$, ns: $P > 0.05$.

	d.f.	v_{ma}	v_{mi}	v_{Total}	θ_{ma}	θ_{mi}	θ_{Total}
Complete model	23	13.68***	18.63***	34.91***	10.73***	26.77***	23.8***
GWC	5	34.35***	122.36***	124.30***	13.89***	103.01***	98.61***
P	1	66.16***	23.97***	43.06***	35.47***	0.07 ^{ns}	16.88***
E	1	0.42 ^{ns}	31.62***	19.59***	0.36 ^{ns}	4.26*	1.51 ^{ns}
P × E	1	5.63*	7.25**	1.88 ^{ns}	2.28 ^{ns}	3.09 ^{ns}	0.23 ^{ns}
GWC × P	5	27.64***	4.87***	1.46 ^{ns}	17.97***	16.16***	2.64*
GWC × E	5	3.55**	0.96 ^{ns}	1.78 ^{ns}	2.41*	2.02 ^{ns}	2.73*
GWC × P × E	5	11.47***	1.42 ^{ns}	1.26 ^{ns}	7.44***	0.45 ^{ns}	1.80 ^{ns}

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Interactions between organisms and parent materials of Technosol

M. Deeb et al.




Figure 1. Configurations of water partitioning in macropores and micropores related to the shrinkage phases of a standard shrinkage curve (water content W , specific volume V) (adapted from Braudeau et al., 2004).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Interactions between organisms and parent materials of Technosol

M. Deeb et al.




Figure 2. (a) Belowground, (b) aboveground and (c) total biomasses production of *Lolium perenne* according to different ratios of green waste compost in the presence/absence of the earthworm *Aporrectodea caliginosa*; mean \pm s.e., $n = 4$ per treatment. Tukey test, significant differences are indicated by different letters, $P < 0.05$.

Interactions between organisms and parent materials of Technosol

M. Deeb et al.




Figure 3. Averaged shrinkage curves ($n = 4$ per curve) for the six mixtures of green waste compost (GWC) and excavated deep horizons (0, 10, 20, 30, 40, 50 % of GWC) reported as the void ratio as a function of the moisture ratio. Each panel represents one of the four treatments: **(a)** control, **(b)** earthworms, **(c)** plants, **(d)** earthworms and plants. The dashed line represents the saturation line.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Interactions between organisms and parent materials of Technosol

M. Deeb et al.




Figure 4. Averaged shrinkage curves ($n = 4$ per curve) for the four treatments (control, earthworms, plants, earthworms and plants) reported as the void ratio as a function of the moisture ratio. Each panel represents one mixture of green waste compost (GWC) and excavated deep horizons: **(a)** 0% GWC, **(b)** 10% GWC, **(c)** 20% GWC, **(d)** 30% GWC, **(e)** 40% GWC, **(f)** 50% GWC. The dashed line represents the saturation line.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Interactions between organisms and parent materials of Technosol

M. Deeb et al.





Figure 5. Linear Discriminant Analysis of the influence of control, earthworm, plant, and both earthworm and plant on hydro-structural parameters. The first and the second axes explained 42 and 26% of the variance, respectively. v_L , moisture ratio at saturated macropores, v_M moisture ratio at saturated micropores, v_N limit of shrinkage, e_0 void ratio at the end of the shrinkage curve, K_{st} the slope of structural phase K_{bs} the slope of the basic phase and K_L , K_N parameters related to shape form.

SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.



Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Figure 6. Moisture ratio at **(a)** maximum saturated macropores (v_{ma} , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{solid}}$), **(b)** maximum saturated micropores (v_{mi} , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{solid}}$), **(c)** total moisture ratio (v_{Total} , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{solid}}$); available water of **(d)** macropores (θ_{ma} , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{soil}}$), **(e)** micropores (θ_{mi} , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{soil}}$), and **(f)** total available water (θ_{Total} , $\text{cm}^3_{\text{water}} \text{cm}^{-3}_{\text{soil}}$) according to the proportion of compost for the four organism treatments (presence/absence of earthworms and/or plants). Mean \pm s.e., $n = 4$ per treatment.

SOILD

2, 1309–1344, 2015

Interactions between organisms and parent materials of Technosol

M. Deeb et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Interactions between organisms and parent materials of Technosol

M. Deeb et al.




Figure 7. Linear regression between total dry plant biomass and available water ($\text{cm}^3_{\text{water}} \cdot \text{cm}^{-3}_{\text{soil}}$) with earthworm (dotted line) or without earthworm (plain line). Plant biomass was higher with earthworms than without ($P < 0.001$). Total plant biomass increased with available water, but the difference in slope of the two linear regressions was not significant. The best equations fitting the relation between total dried plant biomass (X , g) and plant available water (θ_{Total} , $\text{cm}^3_{\text{water}} \cdot \text{cm}^{-3}_{\text{soil}}$) are: $X = 8.97 \cdot \theta_{\text{Total}} + 4.07$ and $X = 8.97 \cdot \theta_{\text{Total}} + 2.69$ with and without earthworms, respectively ($P < 0.001$, adjusted $r^2 = 0.65$).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

