

1 **Exploring the linkage between spontaneous grass cover biodiversity and soil degradation**
2 **in two olive orchard microcatchments with contrasting environmental and management**
3 **conditions**

4 E.V. Taguas^(*), C. Arroyo⁽¹⁾, A. Lora⁽¹⁾, G. Guzmán⁽¹⁾, K. Vanderlinden⁽²⁾ and J.A. Gómez⁽³⁾

5 (1) School of Agronomy and Forestry Engineering- University of Cordoba. Campus Rabanales. Leonardo
6 Da Vinci building. 14071 Córdoba (Spain). Ph. +34 957 218533; E-mail: o72arbac@uco.es,
7 evtaguas@uco.es(*), cr1logoa@uco.es, g92gudim@uco.es.

8 (2) IFAPA, Centro Las Torres-Tomejil-Ctra. Sevilla-Cazalla, km 12.2- 41200 Alcalá del Río (Seville),
9 Spain. E-mail: karl.vanderlinden@juntadeandalucia.es.

10 (3) Institute of Sustainable Agriculture (CSIC). Avenida Alameda del Obispo s/n 14004. Córdoba
11 (Spain). E-mail: joseagomez@ias.csic.es.

12
13 **Abstract.** Spontaneous grass covers are an inexpensive soil erosion control measure in olive orchards.
14 Olive farmers allow grass to grow on sloping terrain to comply with the basic environmental standards
15 derived from the Common Agricultural Policy (CAP- European Commission). However, to date there are
16 few studies assessing the environmental quality considering such covers. In this study, we measured
17 biodiversity indices for spontaneous grass cover in two olive orchards with contrasting site conditions and
18 management regimes in order to evaluate the potential for biodiversity metrics to serve as an indicator of
19 soil degradation. In addition, the differences and temporal variability of biodiversity indicators and their
20 relationships with environmental factors such as soil type and properties, precipitation, topography and
21 soil management were analyzed.

22 Different grass cover biodiversity indices were evaluated in two olive orchard catchments under
23 conventional tillage and no tillage with grass cover, during 3 hydrological years (2011-2013). Seasonal
24 samples of vegetal material and photographs in a permanent grid (4 samples/ha) were taken to
25 characterize the temporal variations of the number of species, frequency of life forms, diversity and
26 modified Shannon's and Pielou's indices.

27 Sorensen's index showed strong differences in species composition for the grass covers in the two olive
28 orchard catchments probably linked with the different site conditions. The catchment (CN) with the best
29 site conditions (deeper soil and higher precipitation) and most intense management presented the highest
30 biodiversity indices as well as the highest soil losses (over 10 t.ha⁻¹). In absolute terms, the diversity
31 indices of vegetation were reasonably high for agricultural systems in both catchments, despite the fact
32 that management activities usually severely limit the landscape and the variety of species. Finally, a
33 significantly higher content of organic matter in the first 10 cm of soil was found in the catchment with
34 worse site conditions in terms of water deficit, average annual soil losses of 2 t.ha⁻¹ and the least intense
35 management. Therefore, the biodiversity indices considered in this study to evaluate spontaneous grass
36 cover were not found to be suitable for describing the soil degradation in the study catchments.

37 **Key words:** olive orchard; spontaneous grass cover, biodiversity, management; soil degradation.

38 **1. Introduction**

39 Soil degradation is defined as the deterioration and loss of soil functions, involving processes such as
40 soil erosion, sedimentation problems in flood plains and reservoirs, climate change, watershed functions
41 and changes in natural habitats leading to loss of genetic stock and biodiversity (Chen et al., 2002). The
42 agricultural intensification of 20th century Europe has led in general terms to a widespread decline in
43 farmland biodiversity across many taxa (Benton et al., 2003). The new 2020 Biodiversity Strategy
44 (European Commission, 2011; 2011/2307 INI) aims to improve the contribution of fisheries and
45 agricultural and forestry sectors to biodiversity. In addition, the Multi-annual Financial Framework for
46 2014–2020 offers significant opportunities to improve synergies not only in soil biodiversity but also with
47 respect to other degradation processes such as soil loss (Cross-compliance. Agriculture and Rural
48 Development; European Commission, 2014).

49 An area of over 2.5 Mha is dedicated to olive cultivation in Spain (MAGRAMA, 2013), which represents
50 about 41% of the world olive production. Olive harvesting and its associated agri-food industries are
51 especially important in rural areas from a socio-economical viewpoint. Over 60% of the area dedicated to
52 olives is located in Andalusia, the southernmost region of the country. A high risk of soil degradation has
53 been described by multiple authors such as Gómez-Limón et al., (2009) and Gómez et al., (2014a) as the
54 result of the interaction of climatological and topographical factors and/or inappropriate soil management.
55 Olive trees have traditionally been cropped under rainfed conditions and on sloping areas where other
56 crops are difficult to grow; they usually provide very low yields or require large investments in order to
57 exploit them properly. The characteristics of the Mediterranean type of climate, where long dry periods
58 alternate with intense rainfall events, in conjunction with soil management systems that pursue bare soils
59 to minimize water competition by weeds entail a high susceptibility to severe water erosion of the soil
60 (Gómez et al., 2014a). Therefore, the use of cover crops has been promoted for soil protection, given their
61 proven effectiveness in controlling water erosion (Gómez et al., 2004; Gómez et al. 2009a, 2009b;
62 Márquez-García et al., 2013; Taguas et al., 2013 among others). In fact, growing in between the olive tree
63 rows is currently a compulsory requirement if the mean slope of the plot is over 15%, according to cross-
64 compliance rules (European Commission, 2014). Spontaneous covers are usually irregular and develop
65 slowly, but tend to achieve a significant growth during spring which may result in greater competition for
66 water and nutrients during the most critical periods of the olive growing cycle. However, due to its zero
67 cost, it is a common alternative in low production olive farms (e.g. Taguas et al., 2013). Furthermore,
68 additional advantages of spontaneous covers in terms of biodiversity, carbon sequestration and the
69 aesthetic improvement of the landscape might make it worth to study their potential contribution.

70 The study of spontaneous grass cover and their interactions with soil have been traditionally associated
71 with the improvement in crop yield (e.g. Graziani et al., 2012; Kamoshita et al., 2014;) or habitat and
72 species conservation (e.g. Albrecht, 2003; Hyvönen and Huusela-Veistola, 2008; Aavik and Liira, 2009)
73 in agronomical and ecological terms, respectively. However, their importance as indicators of soil
74 degradation has scarcely been explored.

75 The bio-indicators of soil quality are commonly associated to the biological activity of their
76 microorganisms; however, spontaneous grass cover biodiversity may be a simpler way to indicate the risk
77 of soil degradation, given that richer and more complex ecological niches might produce more vegetal
78 biomass, efficient cover and eventually, soil protection, as well as habitat and food opportunities for other

79 elements of the trophic chain, such as birds or reptiles. In addition, one key drawback for the proper
80 implementation of environmental protection policies is the lack of a well-defined quantitative measure or
81 indicator of biodiversity which was suitable to describe, compare or measure possible changes (Büchs,
82 2003; Spangenberg, 2007; Moonen and Barberi, 2008). The use of biological indices –in this case
83 associated to grass spontaneous cover- might be helpful because they are more sensitive to changes than
84 chemical and because physical soil indicators and that they could give a broader picture of soil quality
85 (Bastida et al, 2008).

86 The main hypothesis of this study was that richer ecological niches mean lower risks of soil degradation
87 in terms of indicators such as organic matter decline, bulk density and runoff coefficients and soil losses.
88 This would be associated to an optimum space taking derived from the presence of distinct species. In
89 addition, we postulate that the interactions of soil and management explain better the diversity of
90 spontaneous grass covers than the environmental site conditions (annual/seasonal patterns) due to minor
91 soil disturbances which might produce conditions which bring it closer to natural systems.

92 The specific objectives of this work were 1) to describe and compare the biodiversity indices for
93 spontaneous grass covers in two olive orchards with contrasting management intensities, environmental
94 conditions and yields; 2) to analyze the temporal patterns of these indices, relative to meteorological
95 conditions and soil management; and 3) to evaluate the relevance of biodiversity indices as indicators for
96 soil quality, in terms of soil degradation.

97 **2. Materials and methods**

98 **2.1. Study sites**

99 The study catchments are located in the province of Córdoba (Fig. 1, Table 1), in Southern Spain. Both
100 were described in detail by Gómez et al. (2014b) and Taguas et al. (2013) to evaluate the erosive patterns
101 for the periods 2006-2011 and 2005-2011, respectively. The results of those studies were considered an
102 accurate representation of the soil degradation state.

103 The “Conchuela” catchment (CN; 37.6 °N, -5.0 °W, Spain) is situated in a fertile area along the old
104 terraces of the River Guadalquivir (Gómez et al. (2014b). The drainage area of the catchment is 8.0 ha,
105 and it presents an average elevation of 142 m and a mean slope equal to 9%. The climate is classified as
106 Mediterranean with an average annual precipitation of 642 mm, which is mainly concentrated from
107 October to March (about 76% of the precipitation). The average annual temperature is 17.5 °C. The
108 maximum daily mean temperature is usually recorded in July (27.8 °C) while the minimum is generally
109 observed in January (8.1 °C). The soil is a Vertisol, according to the FAO classification (FAO, 2006). It is
110 a deep soil, very plastic when wet, but when dry, the presence of cracks induces high infiltration rates.
111 The predominant soil texture is clay-loam (Table 1). The olive trees were planted in 1993 with 6 × 7 m
112 tree spacing. The mean olive yield in the catchment is 8000 kg·ha⁻¹ During the study period, the farmer
113 allowed the growth of grass spontaneous cover in the lanes from the end of winter until April. Herbicide
114 (glyphosate and oxifluorfen) treatments were applied to control their growth in the tree line from March
115 to September (Table 2). Occasionally surface tillage was made at selected locations within the catchment
116 to cover rills and small gullies obstructing machinery traffic within the orchard. Mowing in the tree lane
117 was performed in areas of excessive grass cover from late winter to early spring. Harvesting is semi-

118 mechanized using tree-vibrators from late autumn to mid-winter, depending on weather conditions and
119 when the fruit ripens (Gómez et al., 2014b; Table 2).

120 The “Puente Genil” catchment (PG; 37.4 °N, -4.8 °W) represented a marginal olive orchard with a very
121 low production. Management operations are kept to a minimum in order to reduce costs. It is located in an
122 area with a long tradition of olive cropping in the upper reaches of the Guadalquivir Valley (Taguas et al.
123 2013). The catchment has a drainage area of 6.1 ha and the mean elevation is 239 m. The average slope is
124 equal to 15 %. As for the climate type, the catchment is located in a Mediterranean area with a mean
125 annual precipitation is of 400 mm. The average temperature in the hottest month (July) is 26.5 °C, while
126 in the coldest month (January) it is 8.4 °C. The main soil category of the catchment is Cambisol (FAO
127 classification; FAO, 2006) with sandy-loam texture (Table 1 and 3). Calcic parental material is located at
128 different points of the catchment with a very shallow soil, mainly on the Western hillslope (Fig. 1b). In
129 contrast, on the Eastern hillslope, soil depth is more than 3 m. The areas closer to the catchment outlet are
130 old terraces with abundant coarse calcarean material. The mean olive yield is 1300 kg·ha⁻¹. The olive
131 trees’ age is 17 years. They were planted on a 7 m × 7 m grid. No-tillage with spontaneous grass cover
132 growing from winter to spring was the management type corresponding with the first few years.
133 Spontaneous grass is removed once (only in spring) or twice a year (September or October and March,
134 April or May), mechanically or using phytosanitary products under the canopies (or combining both; see
135 also Taguas et al., 2013). The details of the management applied during the study period are summarized
136 in Table 2.

137

138 **2.2. Spontaneous grass cover sampling**

139 Four spontaneous grass cover surveys were performed per year (1 per season) during 2011, 2012 and
140 2013. Survey dates were based on the preceding meteorological conditions that determined the
141 germination periods, as well as the development of the spontaneous grass cover. A grid was established in
142 each catchment (Fig. 1) with a sampling density between 4 and 6 points/ha. In each geo-referenced grid
143 point, a 0.5 × 0.5-m frame was used to delimit the survey area (Fig. 2). These sampling points were
144 always placed in the lanes between the lines of trees, away from the olive canopy and the areas of drip
145 irrigation and herbicide application. Plant samples were taken in order to identify the species present at
146 each grid point. In addition, photographs of each point were taken (Reflex Olympus E-420, ED 14-42
147 mm; height 1.4 m-1.7 m; Fig. 2) to observe the annual and seasonal differences of the grass spontaneous
148 cover.

149

150 **2.3. Data Analyses: biodiversity indices, meteorological variables and soil quality** 151 **indicators**

152 *2.3.1. Biodiversity indices*

153 The indices considered to evaluate the biodiversity associated to the grass spontaneous grass cover were
154 richness (*R*), Sorensen’s index (*Is*), transformed Shannon’s (*Hmod*) and Pielou’s indices (*Jmod*), absolute
155 frequency of occurrence and biological spectrum. *R* was determined for the total number of grasses and
156 forbs found per catchment per season and per point. Firstly, in each sample point of the grid (Fig. 1 and

157 Fig 2a and 2b), the species present were identified with pictures and vegetal material, and then the total
158 number species in each catchment (on a seasonal and annual scale) were calculated.

159 I_s indicates the degree of similarity of two samples (study sites) as regards the species composition (Eq.
160 1). It ranges from 0 to 1, where 0 means that the two samples are completely different and 1 completely
161 equal.

$$162 \quad I_s = \frac{2 \cdot C}{A+B} \quad (\text{Eq. 1})$$

163 Where: A is the number of species identified in PG, B the number of species identified in CN, and C is the
164 number of species common to both study sites.

165 Shannon's index, H , (Eq. 2; Shannon and Weaver, 1949) represents the uncertainty associated to the
166 prediction of species identity of an individual taken from a sample. It usually produces values of between
167 1.5 and 4.5. Minimum values are obtained when most of the individuals belong to the same species or to a
168 small group of (less diverse) species, while the highest values are produced in communities where all the
169 species have the same number of individuals. If there is only one group of species Shannon's index is
170 equal to 0.

$$171 \quad H = -\sum_{i=1..n} (p_i \cdot \ln(p_i)) \quad (\text{Eq. 2})$$

173 Where: $p_i = n_i / N$; n_i is the number of individuals corresponding to the species i , and N is the total
174 number of individuals. In this case, a modification of Shannon's index, H_{mod} , was used to simplify the
175 analysis, based on the evaluation of pictures that presented each grid point of the considered in the
176 catchment sample (see Fig. 1 and 2).
177 Therefore, n_i was substituted by the number of grid points where a species was present and N , the total
178 number of grid points considered. The suitability of the transformations associated to H_{mod} was verified
179 with the samples taken in spring 2013 in both catchments.

180 Pielou's equity index (Eq. 3; Pielou, 1969) measures the ratio of the observed diversity and the maximum
181 expected diversity. It varies between 0 and 1, with 1 describing systems where all species are equally
182 abundant.

$$183 \quad J = \frac{H}{\ln(S)} \quad (\text{Eq. 3})$$

185 Where:
186 H is Shannon's index and S is the number of species. If H (Eq. 3) is substituted for H_{mod} , then J_{mod} is
187 obtained.

188
189 Finally, the biological spectrum or life-form (Raunkiaer, 1934) was identified for each species according
190 to its behavior during the unfavorable season (June-September): Epiphytes; Phanerophytes
191 Chamaephytes; Hemicryptophytes; Therophytes; Cryptophytes.

192
193 *2.3.2. Meteorological variables to describe temporal variability of biodiversity indicators.*
194 The cumulative precipitation (P), cumulative reference evapotranspiration (ETP) and average minimum
195 daily temperatures (Tm) were considered in order to evaluate their influence on the biodiversity indices.

196 The daily precipitation was recorded in the gauging stations of the catchments, while the daily values of
197 *ETP* and *Tm* were collected from “La Reina” and “Santaella-CSIC” meteorological stations for CN and
198 PG, respectively (CSIC, 2014).

199

200 *2.3.3. Soil degradation indicators: soil loss, runoff, organic matter and bulk density.*

201 The relationships between the mean values of soil losses, runoff coefficients and organic matter content
202 (0-10 cm) in the catchments with *R*, *Jmod* and *Hmod* were explored to discuss the role of biodiversity
203 indices as a proxy of soil quality indicators. Soil loss (*SL*) and runoff coefficient (*Rc*) were measured in
204 the catchments over 5 years (Taguas et al., 2013; Gómez et al., 2014b).

205 The samples for organic matter (*OM*) analysis were taken between 0-10 cm combining the inter-row and
206 the area under the tree canopies obtained on regular grids with a density of 6-10 samples/ha. The number
207 of samples was 90 and 65 in CN and PG, respectively. The Walkley-Black procedure (Nelson and
208 Sommers, 1982) with samples (2 mm sieve) was followed to determine the organic matter content. Bulk
209 density (*BD*) was measured on the same grid using undisturbed soil cores of approximately 250 cm³. The
210 differences in grid and number of samples are due to the tree spacing in the catchments.

211

212 *2.3.4. Statistical analyses*

213 Basic statistics (mean, standard deviation and coefficient of variation) were evaluated for the annual
214 values of *R*, *Jmod*, *Hmod*, *Is* as well as *Tm*, *ETP* and *P*. In the case of *Is*, the average seasonal values were
215 calculated to observe the possible differences in the study sites over the year. The histograms of the
216 biological spectrum measured in the catchments for the study period were also compared.

217 In addition, in order to evaluate the influence of the meteorological variables on the biodiversity indices
218 *Hmod*, *Jmod* and *R*, a correlation analysis was carried out with meteorological features: *P*, *ETP* and *Tm*.
219 The analysis was carried out with the mean values of the variables *P*, *ETP* and *Tm* corresponding to the 5,
220 15, 30, 60, 365 days previous to the sample date. As for soil properties *OM* and *BD*, box and whisker
221 plots and t-test for independent samples were used to determine whether there were significant
222 differences between the study sites. For *SL* and *Rc*, only box and whisker plots were represented because
223 the number of samples was 5. These properties were compared with the biodiversity indices to
224 qualitatively describe the correlation degree.

225

226

227 **3. Results and discussion**

228 **3.1. Variability of the biodiversity indicators**

229 The mean values of *R*, *Hmod* and *Jmod*, were higher in CN than in PG, which probably shows that site-
230 specific conditions have greater importance than long term management effects (Table 4). A lower
231 diversity was identified in PG, which was probably associated with worse environmental conditions in
232 terms of water deficit, as compared to CN (Table 4), coupled with coarser soil texture and lower soil
233 water holding capacity (Table 3). Precipitation was on average 25% lower in PG while *ETP* was slightly
234 higher, with respect to CN (Table 4). The soils at PG were also shallower than at CN and of coarser

235 texture (Table 3), leading to a smaller water storage capacity which might limit the development of
236 vegetation in PG.

237 With the exception of *Jmod*, the highest coefficients of variation were also observed in PG (Table 4).
238 Despite the extremely simplified landscapes of both catchments, *Hmod*-values were notably high for
239 agricultural systems, particularly in the driest year (2011) with values near to 2.2 and 1.9 in CN and PG,
240 respectively (Table 4). As references, Guzmán and Forester (2007) observed for olive orchards with
241 leguminous cover crops *H*-values close to 1.2, whereas in natural systems of Mediterranean semi-arid
242 areas, *H*-values were approximately equal to 1 (Kawada et al.,2012). Under conventional cereal crops,
243 Armengot et al. (2013) quantified a mean *H*-value of 1.5 for 11 fields in Catalonia (Spain) while for a
244 pine afforestation located in a semi-arid catchment in Southwestern Spain, Bonet et al. (2004) came up
245 with *H*-values of 2.8.

246 On the other hand, *Jmod*-values closed to 1, indicated that there were no dominant species in either of the
247 catchments. The lack of a dominant species is frequent in Mediterranean agricultural areas, where a high
248 inter-annual and intra-annual variability of precipitation and temperature produce a wide range of
249 colonizing species awaiting their optimal development conditions. In spite of the selective herbicide
250 treatments (Table 2), differences in *Jmod* between both catchments were small.

251 Sorensen's index numerically illustrated the notable differences of species existing in the catchments
252 (Tables 4-5 and Fig. 3). It is worth noting how winter was the period when the floristic composition was
253 the most similar ($I_s=0.378$) while the spring, the most different ($I_s=0.139$). Although similar distributions
254 of life forms were found (Fig. 3), a different floristic catalogue of species was observed in both
255 catchments, where the lack of Monocotyledonous in PG is remarkable (Table 5). From the soil protection
256 point of view, the current spectrum is not appropriate because most of the species are not permanently
257 present for a long period of the year. However, most of the species constitute the nutritional base for
258 insects and birds. Enrichment of the biological spectrum with Hemicryptophytes and Chamaephytes is
259 suggested in locations where e.g. hedges are compatible with agricultural operations (Guzmán and
260 Foraster, 2007).

261 The coefficients of correlation between weather variables (*Tm*, *ETP* and *P*) and seasonal biodiversity
262 indicators (*Hmod*, *Jmod* and *R*) were in general low (Table 6). Significant correlations were only found
263 for PG as a result of the shallow sandy soil with short-term water availability controlling vegetation. In
264 contrast, the deeper clay soil at CN (Table 1, 3) enhanced long-term water availability and weakened the
265 correlations between weather variables and biodiversity indicators. Significant negative correlations for
266 *ETP15*, *ETP60* (and *Tm60*) are related to water stress, whereas the positive correlations for short-term
267 indicators such as *Tm15* and *ETP5* might indicate optimal conditions for the seed germination and the
268 growth of grass.

269

270 **3.2. Relationships between biodiversity indices and indicators of soil quality**

271 In addition to R , J_{mod} and H_{mod} , the mean annual values of SL and R_c , measurements of OM and BD are
272 also shown in Table 7 and Figure 4. R , J_{mod} and H_{mod} were not correlated with soil indicators. The
273 highest values of soil losses and the lowest values of organic matter were found in CN. The differences in
274 OM between the catchments were significant as is shown in Table 7 and Fig. 4a (average $OM-CN=1.249$
275 $g.cm^{-3}$; average $OM-PG=1.479 g.cm^{-3}$). A large quantity of coarse elements was found in PG, which must
276 be taken into account when understanding the differences in BD (Table 7), although they were not
277 significant (Table 7 and Fig. 4b; $BD-CN= 1.57 g.cm^{-3}$ and $BD-PG=1.50 g.cm^{-3}$). Substantial higher mean
278 soil loss in CN($16.1 t\cdot ha^{-1}$) was found with respect to PG ($1.8 t\cdot ha^{-1}$; Fig. 4c), Likewise, the mean R_c in
279 CN(15.3%) tripled the value of PG (5.1% ; Fig. 4d),

280

281 3.3. General Discussion

282 Indicators of spontaneous grass cover biodiversity were not correlated with soil losses and organic matter.
283 The role of cover crops in soil erosion is related with the dissipation of energy from rainfall and runoff
284 and with the increase of infiltration, which reduces the sediment transport. It was expected that a wider
285 ecological niche would allow for a more efficient occupation of space with probably more biomass, as
286 well as a higher efficiency in the runoff control on the hillslopes. However, in CN, other factors such as
287 precipitation, soil hydrologic characteristics and the possible dominance of concentrated flow (gullies and
288 rills; Gómez et al., 2014b) accounted for higher soil losses and runoff coefficient (much higher than PG
289 values). Lewis et al. (2013) highlighted the potential for soil erosion to disseminate the spontaneous grass
290 cover seedbank and to improve the biodiversity indicators in agro-ecosystems of Northern Europe. In
291 natural Mediterranean systems Cerdá and García-Fayos (2002) and García-Fayos et al. (2010) described
292 the susceptibility to seed removal by water erosion according to seed and landscape features. In this
293 context, an annual sediment delivery ratio of 4% was found in PG using the SEDD model (Taguas et al.,
294 2011) while in Conchuela, the value was over 90% indicating an efficient rate of transport, as calculated
295 by Burguet (2015). Both the different values of soil losses and the annual sediment delivery ratios might
296 illustrate the very different sediment dynamics which contribute towards explaining the greater
297 biodiversity in CN.

298 As for the values of organic matter content, these might be explained by the management systems. No
299 tillage operations were applied in PG from 2005 and machinery traffic was usually minimal (Table 2),
300 which implies less mechanical soil disturbance than in CN, where productive farm management is carried
301 out. In two sites with a silt loam texture in the Ebro Valley in Spain, Fernández-Ugalde et al. (2009) also
302 described an increase in soil organic carbon content associated with non-tillage practices.

303 It is important not to confuse non-tillage allowing spontaneous grass cover vegetation, as used in PG,
304 with non-tillage management with herbicide to maintain bare soil in olive orchards. The later led to larger
305 soil losses, runoff coefficients and soil compaction as compared to conventional tillage and cover crops as
306 was described by Gómez et al., (2004), however, larger carbon and organic matter contents were found in
307 the topsoil, particularly under the canopy (Gómez et al., 1999). As for surface tillage operations in CN,
308 Márquez-García (2013) also found lower values of organic carbon in the topsoil of olive orchards under

309 conventional tillage as compared to cover crops (spontaneous and sown). Near the study catchments, in
310 other agricultural land uses under conservation agriculture, smaller amounts of crop residues, lower soil
311 water contents and larger CO₂ emissions were observed in managements where tillage operations were
312 applied (Cid, 2013).

313 Despite the annual and seasonal variations of meteorological conditions, overall a larger availability of
314 water was observed in CN, as a result of the higher annual precipitation and the notably deeper soil. More
315 extensive management did not lead to greater spontaneous grass cover biodiversity in PG compared to
316 CN. Benton et al. (2003) highlighted the importance of differential seed or edaphic factors contributing
317 distinctly to plant growth and to patchiness in the presence of insects. Similarly, Albrecht and Mattheis
318 (1998) found that a management change from conventional to integrated farming in dicotyledonous crops
319 in Germany did not lead to a substantial increment of rare species number of spontaneous grass cover.
320 Hyvönen et al. (2003) described that differences in spontaneous grass cover species numbers between
321 organically and conventionally cropped fields in Finland were small. Similar results were highlighted
322 under Mediterranean conditions by Graziani et al. (2012) for a sequence of six rotations in Italy. They
323 found that the number of spontaneous grass cover species was only slightly higher in organic systems as
324 compared to low-input conventional systems.

325 Although single steps, such as the application of fertilizers or certain herbicides, may lead to the
326 dominance of some species such as in the case of monocotyledoneous in CN (Table 5), no clear
327 sensitivity to the management was found, as described by Albrecht (2003) in Germany or Pysek et al.
328 (2005) in Central Europe for different crops. This is likely to be a result of the site conditions in CN being
329 substantially better for vegetation growth, which becomes evident from the olive yields at both
330 catchments (CN, 5000-8000 kg ha⁻¹ and PG < 2000 kg ha⁻¹). In fact, crop yield was also used with other
331 soil properties (such as bulk density, water retention, pH, electrical conductivity, plant-available nutrients,
332 organic matter, microbial biomass, soil enzymes) by Mastro et al. (2007) to define a soil quality index in
333 an agricultural area with a rotation of maize, pearl millet, wheat and cowpea in India. In fact, the yield is a
334 common agronomical factor of soil quality for farmers, which may be well-correlated with biodiversity
335 indices of spontaneous grass cover. On the other hand, the traditional metrics used in this study to
336 measure biodiversity - widely used in ecological studies since they are simple to calculate and understand
337 and has been used for a long time (Lamb et al., 2009) – have been criticized because they provide a
338 limited part of the information (Magurran, 2004) and may be unsuitable for monitoring biodiversity
339 intactness (Lamb et al., 2009). These traditional indices, for example, cannot indicate the presence of non-
340 native species or rare plants. Additionally to the yield, *R*, *Hmod*, *Jmod* and *Is*, the group of species
341 shown in Table 5 support short-term environmental advantages of the vegetation growth found in CN,
342 which is likely to be linked to greater water availability despite a more intense management.

343

344 **4. Conclusions**

345 Sorensen's index for two olive orchard catchments in the province of Cordoba (Spain) showed notable
346 differences in composition, which were probably associated with the different site conditions. Although
347 CN had a more intense management, its better site conditions (higher precipitation, deeper soils and less

348 steep slopes) can explain the higher values in richness, Pielou's index and Shannon's index. Water stress
349 is a limiting factor for the development of vegetation in the Mediterranean area, so the notable differences
350 in annual precipitation (400 mm in PG versus 600 mm in CN) account for the differences observed. In
351 addition, a more active sediment transport dynamic might contribute to seed dispersal and to increasing
352 the biodiversity indices.

353 Shannon's index and Pielou's index were relatively high in both catchments, in spite of the major
354 simplifications derived from the agricultural systems. This can be related with the typical Mediterranean
355 dynamics where temporal variability allows different individual species to be incorporated each year
356 according to certain climatological features. The impact of land-use and management in both catchments
357 explains the dominance of short cycle Therophytes, Hemicryptophytes and Cryptophytes, which are
358 extremely resistant to mechanical/chemical treatments, since their buds are kept underground. On the
359 other hand, Therophytes and Hemicryptophytes do not provide efficient soil protection, since their aerial
360 parts are not present during autumn and winter seasons. However, these species are ecologically
361 important for feeding numerous insects and local birds such as partridge (*Alectoris rufa* L).

362 Higher contents of organic matter were determined in PG, the catchment with the worst site conditions in
363 terms of water availability and the least intense management. Additionally, low soil losses have been
364 measured in this catchment. Therefore, biodiversity indicators associated to spontaneous grass cover were
365 not appropriate to describe the soil degradation state in the study areas. More effort to increase the
366 number of study sites should be applied to evaluate if under more similar environmental conditions, the
367 weight of the management in the olive orchards might determine the biodiversity indices of grass
368 spontaneous cover.

369 **Acknowledgments**

370 This study was supported by Projects AGL2009-12936-C03-01, AGL2012-40128-C03-01, AGL2012-
371 40128-C03-02, AGL2012-40128-C03-03 (Spanish Ministry of Economy and Competitiveness), RESEL
372 (Spanish Ministry for Science and Environment) and by FEDER funds. We are very grateful to Francisco
373 Natera and Antonio Muñoz, the owners of the farms who allowed us to carry out all the surveys and to
374 Laura Aguilera and Enrique Gimeno who helped with the field work and provided numerous pictures and
375 material to identify the species.

376

377 **5. References**

378

379 Aavik, T. and Liira, J.: Agrotolerant and high nature-value species—Plant biodiversity indicator groups
380 in agroecosystems. *Ecol. Indic.*, 9 (5), 892-901, 2007.

381 Albrecht, H.: Suitability of arable weeds as indicator organisms to evaluate species conservation effects
382 of management in agricultural ecosystems. *Agric. Ecosyst. Environ.*, 98, 201-211, 2003.

383 Albrecht, H. and Mattheis, A.: The effects of organic and integrated farming on rare arable weeds on the
384 Forschungsverbund Agrarökosysteme München (FAM) research station in southern Bavaria. *Biol.*
385 *Conserv.*, 86, 347-356, 1998.

386 Armengot L.J.M., Chamorro L., Sans F.X. Weed harrowing in organically grown cereal crops avoids
387 yield losses without reducing weed diversity. *Agronomy for Sustainable Development* 33 (2), 405-411,
388 2013.

389 Bastida F., Zsolnay A., Hernández T And García C. Past, present and future of soil quality indices: a
390 biological perspective. *Geoderma*, 147, 159-171, 2008.

391 Benton T.G., Vickery J.A. and Wilson J.D: Farmland biodiversity: is habitat heterogeneity the key?
392 *Trends Ecol. Evol*, 18(4), 182-188, 2003.

393 Bonet, A., Bellot J., Peña J. Landscape dynamics in a semiarid Mediterranean catchment (SE Spain)". In:
394 Mazzoleni, S., G. Di Pasquale, P.De Martino and F. Rego Eds. *Recent dynamics of Mediterranean*
395 *vegetation landscape*. John Wiley & Sons (UK), pp 47-56, 2004.

396 Büchs, W.: Biotic indicators for biodiversity and sustainable agriculture- introduction and background.
397 *Agr. Ecosys. Env.*98, 1-16, 2003.

398 Burguet, M.: Estimating soil losses in two Mediterranean olive catchments using a sediment distributed
399 model. PhD thesis. University of Cordoba, Spain, 1998.

400 Cerdá, A. and García-Fayos, P.,: The influence of seed size and shape on their removal by water erosion.
401 *Catena* 48, 293-301, 2002.

402 Chen J., Chen J., Tan M. and Gong Z: Soil degradation: a global problem endangering sustainable
403 development. *J. Geo. Sci.* 12, 243-252, 2002.

404 Cid, P., Pérez-Priego, O., Orgaz F. and Gómez-Macpherson H.: Short and mid-term tillage-induced soil
405 CO₂ efflux on irrigated permanent and conventional bed planting systems with controlled traffic in
406 southern Spain. *Soil Sci.*, 51, 447-458., 2013.

407 CSIC, Estaciones Agrometeorológicas del Instituto de Agricultura Sostenible.
408 <http://www.uco.es/grupos/meteo/>. Accessed in November 2014.

409 European Commission. 2014. Cross-compliance. Agriculture and Rural Development. Policy areas.
410 <http://ec.europa.eu/agriculture/direct-support/cross-compliance/>, Accessed in January 2014.

411 European Commision, EU Biodiversity Strategy to 2020 – towards implementation.
412 <http://ec.europa.eu/environment/nature/biodiversity/comm2006/2020.htm>, Accessed June 2014.

413 European Commision, 2011. European Parliament resolution of 20 April 2012 on our life insurance, our
414 natural capital: an EU biodiversity strategy to 2020 (2011/2307(INI)), Accessed June 2014.

415 FAO-Food and Agriculture organization of the United Nations. IUSS Working Group WRB. 2006.
416 World reference base for soil resources 2006. World Soil Resources Reports No. 103. FAO, Rome.

417 Fernández-Ugalde, O., Virto, I., Bescansa, P., Imaz, M.J., Enrique, A. and Karlen, D.L: No-tillage
418 improvement of soil physical quality in calcareous, degradation-prone, semiarid soils. *Soil Till. Res.* 106,
419 29-35. 2009.

420 García-Fayos, P., Bochet, E. and Cerdá, A.: Seed removal susceptibility through soil erosion shapes
421 vegetation composition. *Plant Soil* 334, 289-297, 2010.

422 Graziani, F., Onofri, A., Pannacci, E., Tei, F. and Guiducci, M.: Size and composition of weed seedbank
423 in long-term organic and conventional low-input cropping systems. *Eur. J. Agron.* 39, 52-61, 2012.

424 Gómez, J.A. and Giráldez, J.V.: Erosión y degradación de suelos. En: *Sostenibilidad de la producción de*
425 *olivar en Andalucía*". Junta de Andalucía. Sevilla, Spain, pp. 45-86, 2009.

426 Gómez, J.A., Giráldez, J.V, Pastor, M. and Fereres, E. Effects of tillage method on soil physical
427 properties, infiltration and yield in an olive orchard. *Soil Till. Res.* 52, 167-175, 1999.

428 Gómez, J.A., Infante-Amate, J., González de Molina, M., Vanwalleghem, T., Taguas, E.V. and Lorite I.
429 Olive Cultivation, its Impact on Soil Erosion and its Progression into Yield Impacts in Southern Spain in
430 the Past as a Key to a Future of Increasing Climate Uncertainty. *Agriculture*, 4 (2), 170-198, 2014a.

431 Gómez, J.A., Vanwalleghem, T., De Hoces, A., Taguas, E.V. 2014. Hydrological and erosive response of
432 a small catchment under olive cultivation in a vertic soil during a five-year period: Implications for
433 sustainability. *Agriculture, Ecosystems and Environment* 188: 229–244, 2014b. Gómez, J.A., Guzmán,
434 M.G., Giráldez J.V., and Fereres, E.: The influence of cover crops and tillage on water and sediment
435 yield, and on nutrient, and organic matter losses in an olive orchard on a sandy loam soil. *Soil Till. Res.*,
436 106, 137–144, 2009b.

437 Gómez, J.A., Romero, P., Giráldez J.V. and Fereres, E.: Experimental assessment of runoff and soil
438 erosion in an olive grove on a Vertic soil in southern Spain affected by soil management. *Soil Use*
439 *Manage* 20, 426–431, 2004.

440 Gómez, J.A., Sobrinho, T., Giráldez, J.V. and Fereres, E.: Soil management effects on runoff, erosion and
441 soil properties in an olive grove of Southern Spain. *Soil Till.Res.*, 102, 5-13, 2009a.

442 Gómez-Limón, J.A., Picazo-Tadeo, A.J. and Reig-Martínez, E.: Eco-efficiency assessment of olive farms
443 in Andalusia. *Land Use Policy* 29, 395- 406, 2011.

444 Guzmán G and Foraster L.: Manejo de la cubierta vegetal en el olivar ecológico en Andalucía: siembra de
445 leguminosas entre calles. *Consejería de Agricultura y Pesca, Junta de Andalucía*, pp 78, 2007.

446 Hyvönen, T., and Huusela-Veistola, E. Arable weeds as indicators of agricultural intensity – A case study
447 from Finland. *Biol. Conserv.*, 141 (11), 2857-2864, 2008.

448 Kamoshita, A., Araki, Y. and Nguyen, Y.T.B. Weed biodiversity and rice production during the irrigation
449 rehabilitation process in Cambodia. *Agr. Ecosys. Environ.*, 194 (1), 1-6. 2014.

450 Kawada, K., Suzuki K., Suganuma H., Smaoui A., Isoda H. Biodiversity in the Semi-arid Zone of
451 Tunisia. *Journal of Arid Land Studies*, 22 (1), 83 -86, 2012.

452 Lamb, E.C., Bayne E., Holloway G., Schieck J., Boutin S., Herbers J. and Haughland D.L.: Indices for
453 monitoring biodiversity change: Are some more effective than others? *Ecol. Ind.* 9, 423-444, 2009.

454 Lewis, T.D., Rowan, J.S., Hawes, C., McKenzie B.M.: Assessing the significance of soil erosion for
455 arable weed seedbank diversity in agro-ecosystems. *Prog. Phys. Geog.*, 37 (5), 622-641, 2013.

456 MAGRAMA- Ministry of Agriculture, Food and Environment of Spain. Líneas Estratégicas para la
457 Internacionalización del Sector Agroalimentario. [http://www.magrama.gob.es/es/ministerio/planes-
458 estrategias/lineas-estrategicas-para-la-
459 internacionalizaciondelsectoragroalimentario/lineas_estrat%C3%A9gicas_internacionalizaci%C3%B3n_t
460 cm7-278627.pdf](http://www.magrama.gob.es/es/ministerio/planes-estrategias/lineas-estrategicas-para-la-internacionalizaciondelsectoragroalimentario/lineas_estrat%C3%A9gicas_internacionalizaci%C3%B3n_tcm7-278627.pdf) . Accessed in December 2013.

461 Magurran, A.E. *Measuring Biological Biodiversity*. Blackwell, Oxford, pp:18-71, 2004.

462 Márquez-García, F., González-Sánchez, E.J., Castro-García, S. and Ordóñez-Fernández, R. Improvement
463 of soil carbon sink by cover crops in olive orchards under semiarid conditions. Influence of the type of
464 soil and weed. *Spanish J. Agr. Res.*, 11(2), 335-346, 2013.

465 Masto, R.E., Chhonkar, P.K., Singh, D. and Patra, A.K. Soil quality response to long-term nutrient and
466 crop management on a semi-arid Inceptisol. *Agric. Ecosyst. Environ.*, 118 (1-4), 130-142, 2007.

467 Nelson, D.W., and Sommers, L.E., Total carbon, organic carbon, and organic matter. In: Page, A. L., H.
468 Miller, and D. R. Keeney (eds). *Methods of Soil Analysis: Part 2. Chemical and Microbiological*
469 *Properties*. ASA Monograph No. 9. Soil Science Society of America, Madison WI, USA. pp: 539-577,
470 1982.

471 Pysek, P., Jarosik, V., Kropac, Z., Chytrý, M., Wild, J. and Tichý, L: Effects of abiotic factors on species
472 richness and cover in Central European weed communities. *Agr. Ecosys. Environ.*, 109, 1-8, 2005.

473 Spangenberg, J.H.: Biodiversity pressure and the driving forces behind. *Ecol. Econ.* 61, 146-158, 2007.

474 Taguas, E.V., Ayuso, J.L., Pérez, R., Giráldez, J.V. and Gómez J.A: Intra and inter-annual variability of
475 runoff and sediment yield of an olive micro-catchment with soil protection by natural ground cover in
476 Southern Spain. *Geoderma*, 206, 49-62, 2013.

477 Taguas, E.V., Moral, C., Ayuso, J.L., Pérez, R. and Gómez, J.A.: Modeling the spatial distribution of
478 water erosion within a Spanish olive orchard microcatchment using the SEDD model. *Geomorphology*,
479 133, 47-56, 2011.

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497 **TABLES**

498 Table 1. Summary of the main environmental features in the study catchments

Name	La Conchuela	Arroyo Blanco
Location	Córdoba	Puente Genil (Córdoba)
Drainage area (ha)	8.0	6.1
Mean elevation (m)	142	239
Mean slope (%)	9	15
Mean annual precipitation (mm)	642	400
Max. and min. daily average temperatures	27.8° July/8.1° January	26.5° July/ 8.4° January
Soil type (FAO; see details in Table 3)	Vertisol	Cambisol
Texture	Clay-loam	Sandy-loam
OM content (% topsoil)	1.1	1.4
Mean olive yield (kg/ha)	8000	1300
Management (see details in Table 2)	Spontaneous grass cover controlled with a combination of mowing, and occasional herbicide application	Extensive, non-tillage with a spontaneous grass cover

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518 **Table 2.** Management operations applied during the study periods in both catchments.

Catchment	Month	2011	2012	2013
CN	January		Harvesting: Mechanical vibrators combined with a buggy with an umbrella to collect the olives.	Harvesting: Mechanical vibrators combined with a buggy with an umbrella to collect the olives.
	February			
	March	Herbicide treatments around trees (glyphosate and oxifluorfen in infested areas)		Herbicide treatments around trees (glyphosate and oxifluorfen in infested areas)
	April	Mowing of lane areas	Herbicide treatments around trees (glyphosate and oxifluorfen in infested areas) Mowing of lane areas	Mowing of lane areas
	May	Drip irrigation	Drip irrigation	Drip irrigation
	June	Drip irrigation	Drip irrigation	Drip irrigation
	July	Drip irrigation Herbicide treatments around trees (glyphosate and oxifluorfen in infested areas)	Drip irrigation	Drip irrigation Herbicide treatments around trees (glyphosate and oxifluorfen in infested areas)
	August	Drip irrigation	Drip irrigation Herbicide treatments around trees (glyphosate and oxifluorfen in infested areas)	Drip irrigation
	September	Drip irrigation	Drip irrigation	Drip irrigation
	October			
	November			
	December	Harvesting: Mechanical vibrators combined with a buggy with an umbrella to collect the olives.	Harvesting: Mechanical vibrators combined with a buggy with an umbrella to collect the olives.	
PG	January			
	February			
	March			
	April	4 tractor passes to mechanically clear the spontaneous grass cover.		
	May	Foliar fertilization (N, Mg & Fe)	4 tractor passes to mechanically clear the spontaneous grass cover Herbicide treatments around trees (glyphosate)	
	June			
	July			
	August			
	September			4 tractor passes to mechanically clear the spontaneous grass cover. Herbicide treatments around trees (glyphosate)
	October			
	November	Harvesting: Mechanical vibrators combined with a buggy with an umbrella to collect the olives.	Harvesting: Mechanical vibrators combined with a buggy with an umbrella to collect the olives.	Harvesting: Mechanical vibrators combined with a buggy with an umbrella to collect the olives.
	December			

519

520

521

522
523

Table 3. Soil properties in two profiles of the catchments (PG= Puente Genil; CN= Conchuela; OM= organic matter content)

<i>Catchment</i>	<i>Horizon</i>	<i>Width (cm)</i>	<i>Coarse elements (%)</i>	<i>Sand (%)</i>	<i>Silt (%)</i>	<i>Clay (%)</i>	<i>Texture class</i>	<i>pH</i>	<i>OM (%)</i>
PG	A	10	22.7	59.5	35.2	5.3	Sandy-loam	8.8	1.59
	C	40	24.4	60.8	34.3	4.9	Sandy-loam	8.8	1.59
CN	A	0-56	0.36	5.9	45.1	49.0	Clay	8.6	0.96
	B	56-110	0.00	5.9	46.4	47.7	Clay	8.7	0.53
	BC	110-138	0.00	-	-	-	Clay-loam	-	-
	C	>138	0.00	-	-	-	Clay-loam	-	-

524
525
526
527

528

529 Table 4. Annual values of biodiversity indices: Richness (*R*), modified Shannon's (*Hmod*) and
 530 Pielou's indices (*Jmod*) and seasonal Sorensen's indices (*Is*); and meteorological attributes:
 531 average minimum temperature (*Tm*), annual evapotranspiration (*ETP*) and precipitation (*P*) for
 532 both catchments. (CV=coefficient of variation).

Index	Catchment/ Season	2011	2012	2013	Mean	St. Dev.	CV(%)
<i>R</i>	CN	23	26	28	25.7	2.5	9.7
	PG	24	14	24	20.7	5.8	28.0
<i>Hmod</i>	CN	2.194	1.947	1.826	1.989	0.187	9.4
	PG	1.880	1.213	1.751	1.614	0.354	21.9
<i>Jmod</i>	CN	0.897	0.839	0.850	0.862	0.031	3.6
	PG	0.840	0.834	0.817	0.830	0.012	1.4
<i>Is</i>	Winter	0.231	0.571	0.333	0.378	0.174	46.0
	Spring	0.231	0.100	0.087	0.139	0.080	57.6
	Summer	0.320	0.000	0.363	0.228	0.198	86.8
	Sutumn	0.166	0.333	0.000	0.166	0.167	100.6
<i>Tm</i> (°C)	CN	11.7	11.6	11.1	11.5	0.3	2.6
	PG	12.4	11.6	11.7	11.9	0.4	3.4
<i>ETP</i> (mm)	CN	1270.5	1310.2	1230.4	1270.4	39.9	3.1
	PG	1383.7	1359.8	1355.1	1366.2	15.3	1.1
<i>P</i> (mm)	CN	401	610	621.1	544	124	22.8
	PG	376.8	434.4	423.8	411.7	30.7	7.5

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547 Table 5. Species identified in the study catchments present in Puente Genil (PG), Conchuela
 548 (CN) or both catchments (Both) for the study period.

Species Scientific name	Biological Spectrum	Location
Dicotyledonous		
APIACEAE(UMBELLIFERAE)		
<i>Daucus carota</i> L.	Hemicryptophites	CN
ASTERACEAE(COMPOSITAE)		
<i>Anacyclus clavatus</i> (Desf.) Pers.	Therophytes	Both
<i>Anthemis arvensis</i> L.	Therophytes	Both
<i>Calendula arvensis</i> L.	Therophytes	CN
<i>Centaurea melitensis</i> L.	Therophytes	Both
<i>Cirsium arvense</i> (L.) Scop.	Geophytes	Both
<i>Cichorium intybus</i> L.	Hemicryptophites	CN
<i>Conyza sumatrensis</i> (Retz) E. Walker	Therophytes	PG
<i>Chrysanthemum segetum</i> L.	Therophytes	Both
<i>Picris echioides</i> L.	Hemicryptophites, Therophytes	Both
<i>Senecio vulgaris</i> L.	Therophytes	Both
<i>Silybum marianum</i> (L.) Gaerth	Hemicryptophites	CN
<i>Sonchus asper</i> (L.) Hill	Hemicryptophites, Therophytes	Both
<i>Sonchus oleraceus</i> L.	Hemicryptophites, Therophytes	Both
<i>Taraxacum officinale</i> Weber ex F.H. Wiss	Hemicryptophites	Both
<i>Taraxacum obovatum</i> (Willd) D.C	Hemicryptophites	PG
<i>Pulicaria paludosa</i> Link	Hemicryptophites, Therophytes	Both
BORAGINACEAE		
<i>Anchusa azurea</i> Mill	Hemicryptophites	PG
<i>Echium plantagineum</i> L.	Hemicryptophites, Therophytes	Both
<i>Heliotropium europaeum</i> L.	Therophytes	Both
BRASICACEAE(CRUCIFERAE)		
<i>Diplotaxis virgata</i> (Cav) DC	Therophytes	PG
<i>Raphanus raphanistrum</i> L.	Geophytes, Therophytes	Both
<i>Rapistrum rugosum</i> (L.) Bergeret	Therophytes	Both
<i>Sinapis arvensis</i> L.	Therophytes	CN
CARYOPHYLLACEAE		
<i>Spergula arvensis</i> L.	Therophytes	PG
<i>Stellaria media</i> (L.) Vill	Therophytes	Both
CISTACEAE		
<i>Fumana ericoides</i> (cav) Gand. In Magnier	Chamaephytes	PG
CONVOLVULACEAE		
<i>Convolvulus arvensis</i> L.	Geophytes, Hemicryptophites	CN
CRASSULACEAE		
<i>Umbilicus rupestris</i> (Salisb.) Dandy	Hemicryptophites	PG
CUCURBITACEAE		
<i>Ecballium elaterium</i>	Hemicryptophites	CN

FABACEAE(LEGUMINOSAE)		
<i>Ononis punescens</i> L.	Therophytes	PG
<i>Trifolium repens</i> L.	Hemicryptophites	CN
<i>Trifolium campestre</i> Scrb.	Therophytes	CN
GERANIACEAE		
<i>Erodium cicutarium</i> (L.) L'Her	Therophytes	Both
<i>Erodium moschatum</i> (L.) L'Her	Therophytes	CN
<i>Erodium malacoides</i> (L.) L'Her	Therophytes, Hemicryptophites	PG
<i>Geranium molle</i> L.	Therophytes	CN
LAMIACEAE		
<i>Lamium amplexicaule</i> L.	Therophytes	Both
MALVACEAE		
<i>Malva sylvestris</i> L.	Hemicryptophites	Both
PAPAVERACEAE		
<i>Fumaria officinalis</i> L.	Therophytes	CN
POLYGONACEAE		
<i>Polygonum aviculare</i> L.	Therophytes	PG
PRIMULACEAE		
<i>Anagallis arvensis</i> L.	Therophytes	Both
RANUNCULACEAE		
<i>Ranunculus arvensis</i> L.	Therophytes	Both
RUBIACEAE		
<i>Galium aparine</i> L.	Therophytes	Both
SCROPHULARIACEAE		
<i>Veronica arvensis</i> L.	Therophytes	PG
<i>Veronica heredifolia</i> L.	Therophytes	PG
URTICACEAE		
<i>Urtica urens</i> L.	Therophytes	PG
Monocotyledonous		
LILIACEAE		
<i>Muscari comosum</i> (L.) Miller	Geophytes	PG
POACEAE		
<i>Bromus hordaceus</i> L.	Therophytes	CN
<i>Bromus madritensis</i> L.	Therophytes	CN
<i>Bromus squarrosus</i> L.	Therophytes	CN
<i>Hordeum murimum</i> L.	Therophytes	CN
<i>Hordeum leporinum</i> (Link)	Therophytes	CN
<i>Lolium rigidum</i> Gaudin	Therophytes	CN
<i>Poa annua</i> L.	Therophytes	CN

550 Table 6. Matrix of correlation between diversity indices (seasonal values) and climatological
 551 features: *Hmod* = Shannon's modified index; *Jmod* = Pielou's modified index; *R* = richness; *P* =
 552 cumulative precipitation; *Tm*= average of minimum daily temperatures; *ETP* = cumulative
 553 evapotranspiration. Numbers indicate the interval of previous days (5, 15, 30 and 60).

		<i>P5</i>	<i>P15</i>	<i>P30</i>	<i>P60</i>	<i>Tm5</i>	<i>Tm15</i>	<i>Tm30</i>	<i>Tm60</i>	<i>ETP5</i>	<i>ETP15</i>	<i>ETP30</i>	<i>ETP60</i>
	<i>Hmod</i>	0.12	0.33	0.40	0.39	-0.28	-0.26	-0.25	-0.31	-0.35	-0.36	-0.42	-0.43
CN	<i>Jmod</i>	-0.19	-0.25	-0.20	-0.10	0.55	0.52	0.41	0.17	0.29	0.54	0.55	0.44
	<i>R</i>	0.35	0.52	0.49	0.45	-0.16	-0.17	-0.20	-0.29	-0.25	-0.32	-0.36	-0.37
	<i>Hmod</i>	0.23	0.29	0.11	0.39	-0.12	-0.05	-0.42	-0.64	-0.27	-0.58	-0.39	-0.58
PG	<i>Jmod</i>	-0.19	-0.29	-0.42	-0.18	0.40	0.60	0.29	-0.01	0.61	0.26	0.51	0.36
	<i>R</i>	0.29	0.38	0.16	0.36	-0.22	-0.09	-0.42	-0.61	-0.35	-0.62	-0.46	-0.61

554 N=12 – Bold indicates correlations are significant at $p < 0.05$

555

556

557

558 Table 7. Means and standard deviations of the annual biodiversity indicators and parameters of
 559 soil quality: *Hmod* = Shannon's modified index; *Jmod* = Pielou's modified index; *R* = richness;
 560 *OM*= organic matter content in upper horizon (0-10 cm); *BD*= bulk density of upper horizon (0-
 561 10 cm); *SL* = annual soil loss; *Rc* = runoff coefficient (ratio of the annual values of precipitation
 562 and runoff).


Catchment	Stat.	<i>R</i>	<i>Jmod</i>	<i>Hmod</i>	<i>OM</i> * (%)	<i>BD</i> ** (g.cm ⁻³)	<i>SL</i> + (t.ha ⁻¹)	<i>Rc</i> + (%)
CN	Mean	25.7	0.86	1.99	1.25	1.57	16.1	15.3
	St. Dev.	2.5	0.03	0.19	0.37	0.19	20.8	12.7
PG	Mean	20.7	0.83	1.61	1.48	1.50	1.8	5.1
	St. Dev.	5.8	0.01	0.35	0.53	0.25	2.3	4.2

563 (*) T-test showed p=0.00054; CN (n= 95); PG (n=65) (See also Fig. 4a)

564 (**) T-test showed p=0.07764; CN (n= 95); PG (n=65) (See also Fig. 4b)

565 (+) See Figures 3c-d, T-test was not carried out because the number of samples was very
 566 low. CN(n=5 years), PG (n=6 years)

567



569

570 Figure 1. Locations of the study catchments and sample grids: a) La Conchuela (CN); b) Arroyo Blanco
 571 in Puente Genil (PG).

572

573

574

575

576

a)



b)



c)



d)



57

578 Figure 2. Examples of grass spontaneous cover sample plots and view of the catchments: a) a plot in La
579 Conchuela; b) a plot in Puente Genil; c) gully with cover crop in CN; d) view of a hillslope in PG.


580

581

582

583

584



585

586 Figure 3. Number of species by life forms (biological spectrum) in the study catchments (CN= La
 587 Conchuela; PG= Puente Genil).

588

589

590

591

592

593

594


595

596

597

598

599



600
601
602
603
604
605

Figure 4. Box and whisker plots of the measurements of soil degradation indicators: (a) organic matter content in the upper horizon (b) bulk density in the upper horizon; (c) annual soil losses in the catchment outlets; (d) annual runoff coefficients (PG= Puente Genil; CN=La Conchuela; SE= Standard error; For (a) and (b), the sample size was 65 in PG and 95 in CN; For (c) and (d) the sample size was 6 in PG and 5 in CN; The data of (c) and (d) were described in Taguas et al. (2013) and Gómez et al. (2014b)..