

---

Interactive  
comment

# ***Interactive comment on “Low molecular weight organic anions (carboxylates) increase microbial activity and alter microbial community composition in uncontaminated and diesel contaminated soil” by B. C. Martin et al.***

**B. C. Martin et al.**

belinda.martin@research.uwa.edu.au

Received and published: 21 July 2016

Response to referee #2

We thank the reviewer for their useful and positive comments.

You have raised a similar point as referee #1, in that a variety of compounds should have been tested (i.e. not just carboxylates). However, previous work in this area has already been conducted using a mixture of root exudate compounds (Joner and Leyval, 2003; Miya and Firestone, 2001; Xie et al., 2012). We instead attempted to focus on a single group of exudates, as these exudates have been studied extensively for their

potential role in plant nutrient acquisition and metal detoxification, but impact on the soil microbial community has received less attention (Pang et al., 2010; Pearse et al., 2006; Shane and Lambers, 2005). As far as we are aware, there are no studies examining the biodegradation of carboxylate compounds in polluted soils, and we therefore agree with you that it is still a useful contribution to this area of research.

## References

- Joner, E. J. and Leyval, C.: Rhizosphere gradients of polycyclic aromatic hydrocarbon (PAH) dissipation in two industrial soils and the impact of arbuscular mycorrhiza, *Environ. Sci. Technol.*, 37(11), 2371–2375, doi:10.1021/es020196y, 2003.
- Miya, R. K. and Firestone, M. K.: Enhanced phenanthrene biodegradation in soil by slender oat root exudates and root debris., *J. Environ. Qual.*, 30(6), 1911–1918, doi:10.2134/jeq2001.1911, 2001.
- Pang, J., Ryan, M. H., Tibbett, M., Cawthray, G. R., Siddique, K. H. M., Bolland, M. D. A., Denton, M. D. and Lambers, H.: Variation in morphological and physiological parameters in herbaceous perennial legumes in response to phosphorus supply, *Plant Soil*, 331(1), 241–255, doi:10.1007/s11104-009-0249-x, 2010.
- Pearse, S. J., Veneklaas, E. J., Cawthray, G. R., Bolland, M. D. A. and Lambers, H.: Carboxylate release of wheat, canola and 11 grain legume species as affected by phosphorus status, *Plant Soil*, 288(1-2), 127–139, doi:10.1007/s11104-006-9099-y, 2006.
- Shane, M. W. and Lambers, H.: Cluster roots: A curiosity in context, *Plant Soil*, 274(1-2), 101–125, doi:10.1007/s11104-004-2725-7, 2005.
- Xie, X. M., Liao, M., Yang, J., Chai, J. J., Fang, S. and Wang, R. H.: Influence of root-exudates concentration on pyrene degradation and soil microbial characteristics in pyrene contaminated soil, *Chemosphere*, 88(10), 1190–1195, doi:10.1016/j.chemosphere.2012.03.068, 2012.

Printer-friendly version

Discussion paper



Interactive  
comment

