

1 **Soil Denitrifier Community Size Changes with Land Use Change to Perennial**
2 **Bioenergy Cropping Systems**

3

4 *Running Head: Soil Denitrifiers associated with Perennial Grasses*

5 Karen A. Thompson¹, Bill Deen² and *Kari E. Dunfield¹

6 ¹School of Environmental Sciences, University of Guelph, 50 Stone Road East, Guelph, Ontario
7 N1G 2W1, Canada; ²Department of Plant Agriculture, University of Guelph, 50 Stone Road
8 East, Guelph, Ontario N1G 2W1, Canada.

9 *Correspondence to Kari E. Dunfield:

10 tel: (519) 824-4120 Ext.58088, dunfield@uoguelph.ca, fax:519-837-0756

11 Original Research Article

12 **Abstract**

13 Dedicated biomass crops are required for future bioenergy production. However, the effects of
14 large-scale land use change (LUC) from traditional annual crops, such as corn-soybean rotations
15 to the perennial grasses (PGs) switchgrass and miscanthus on soil microbial community
16 functioning is largely unknown. Specifically, ecologically significant denitrifying communities,
17 which regulate N₂O production and consumption in soils, may respond differently to LUC due to
18 differences in carbon (C) and nitrogen (N) inputs between crop types and management systems.
19 Our objective was to quantify bacterial denitrifying gene abundances as influenced by corn-
20 soybean crop production compared to PG biomass production. A field trial was established in
21 2008 at the Elora Research Station in Ontario, Canada (n=30), with miscanthus and switchgrass
22 grown alongside corn-soybean rotations at different N rates (0 and 160 kg N ha⁻¹) and biomass
23 harvest dates within PG plots. Soil was collected on four dates from 2011-2012 and quantitative
24 PCR was used to enumerate the total bacterial community (16S rRNA), and communities of
25 bacterial denitrifiers by targeting nitrite reductase (*nirS*) and N₂O reductase (*nosZ*) genes.
26 Miscanthus produced significantly larger yields and supported larger *nosZ* denitrifying
27 communities than corn-soybean rotations regardless of management, indicating large-scale LUC
28 from corn-soybean to miscanthus may be suitable in variable Ontario climatic conditions and
29 under varied management, while potentially mitigating soil N₂O emissions. Harvesting
30 switchgrass in the spring decreased yields in N-fertilized plots, but did not affect gene
31 abundances. Standing miscanthus overwinter resulted in higher 16S rRNA and *nirS* gene copies
32 than in fall-harvested crops. However, the size of the total (16S rRNA) and denitrifying bacterial
33 communities changed differently over time and in response to LUC, indicating varying controls
34 on these communities.

35 **Key Words: biomass, bioenergy, miscanthus, switchgrass, corn, soy**

36

37 **1. Introduction**

38 Future energy needs require dedicated biomass crop production for bioethanol and combustion-
39 based electricity generation. Corn (*Zea mays* L.) –soybean (*Glycine max* L.) rotations currently
40 dominate the landscape across Ontario and the northern US Corn Belt (Gaudin et al., 2015), and
41 corn grain is currently the primary feedstock for bioethanol production in Canada (Jayasundara
42 et al., 2014). The C4 perennial grasses (PGs) switchgrass (*Panicum virgatum* L.) and miscanthus
43 (*Miscanthus* spp.) have been proposed as alternate feedstock crops to corn for biomass-based
44 bioenergy production due to their large biomass yields, reduced nitrogen (N) and water
45 requirements, decreased nutrient leaching and potential for increased soil carbon (C) storage
46 (Blanco-Canqui and Lal, 2009; Foster et al., 2013). Large scale production of C4 PGs in Ontario
47 and the northern Corn Belt would require land use change (LUC) from existing corn-soybean
48 rotations to PG biomass cropping systems (Deen et al., 2011; Kludze et al., 2013; Liang et al.,
49 2012; Sanscartier et al., 2014).

50 Few studies have assessed how this LUC may influence soil microbial community functioning.
51 In particular, soil denitrifying communities represent an ideal subset of the soil microbial
52 community to target to assess changes in ecosystem functioning due to agricultural management
53 and LUC. Denitrifying bacteria represent approximately 5% of the total soil microbial biomass
54 (Braker and Conrad, 2011) and have been identified in over 60 genera (Philippot, 2006),
55 encompassing a wide range of phylogenetic and functional diversity. Multiple studies have
56 linked changes in denitrifier communities with plant types or development stage (Bremer et al.,
57 2007; Hai et al., 2009; Petersen et al., 2012), N fertilization (Hallin et al., 2009; Yin et al., 2014),
58 organic or conventional crop management (Reeve et al., 2010), perennial vs. annual crop land

59 use (Bissett et al., 2011) and C and N inputs (Bastian et al., 2009). These studies suggest that
60 LUC from corn-soybean rotations to PG species may influence the soil bacterial communities
61 which drive soil N₂O production and consumption.

62 N₂O is a potent greenhouse gas with a global warming potential 296x that of CO₂ (IPCC 2007).
63 However, measuring N₂O directly in the field is often difficult with chamber methods in
64 cropping systems that produce large aboveground biomass. Additionally, including multiple field
65 treatments (eg: as in a RCBD design) make micrometeorological methods of N₂O flux
66 impossible to obtain. Instead, relative abundances of denitrifier genes can be used to assess a
67 soil's potential to produce (e.g. *nirS* or *nirK*) and consume (e.g. *nosZ*) N₂O via denitrification,
68 representing a qualitative proxy of relative N₂O emission potential of a soil (Butterbach-Bahl et
69 al., 2013; Hallin et al., 2009; Morales et al., 2010; Petersen et al., 2012; Philippot, 2002).

70 Denitrifier community size has been correlated with denitrification process rates (Hallin et al.,
71 2009; Wu et al., 2012), and denitrification potential (Attard et al., 2011; Cuhel et al., 2010;
72 Enwall et al., 2010). Potential denitrifying activity and denitrifying community size have also
73 been shown to be correlated with each other in some studies (Hallin et al., 2009; Morales et al.,
74 2010; Szukics et al., 2010; Throback et al., 2007); suggesting community size may indicate
75 potential differences in soil N processes after LUC. Particularly, the *nosZ*-bearing community
76 may act as a N₂O sink and counter high N₂O production rates (Braker and Conrad, 2011;
77 Philippot et al., 2011), therefore influencing N₂O emissions (Cuhel et al., 2010; Morales et al.,
78 2010; Philippot et al., 2011).

79 Denitrification *nirS* and *nosZ* gene targets represent the two most important steps in the
80 denitrification pathway that produce gaseous by-products, and account for a large proportion of
81 functional N genes in some studies (Stone et al., 2015). The first step in denitrification that

82 produces a gaseous N product is the reduction of nitrite (NO₂⁻) to nitric oxide (NO), catalyzed by
83 nitrite reductases either encoded by the cytochrome cd₁ (*nirS*) or copper-containing (*nirK*) genes,
84 which are equivalent but have not been detected within the same species (Zumft, 1997). We
85 chose to quantify *nirS* because ¾ of cultured denitrifiers possess the *nirS* gene (Zumft, 1997) and
86 some molecular reports indicate *nirS* may dominate in abundance over *nirK* in some natural
87 environments (Deslippe et al., 2014; Nogales et al., 2002), indicating it may be a better-suited
88 target for relative characterization of potential nitrite-reducing communities than *nirK*.
89 Additionally, *nirK* has been recently identified in autotrophic ammonia-oxidizing species
90 (Cantera and Stein, 2007; Casciotti and Ward, 2001), calling into question its utility in
91 specifically targeting denitrifying communities. The *nosZ* target codes for nitrous oxide
92 reductase, which catalyzes the reduction of N₂O to N₂ in the denitrification pathway, indicating
93 *nosZ*-bearing communities help to complete the N cycle and determine the N₂O:N₂ balance. 16S
94 rRNA was chosen as a molecular target for the total bacterial community size; although 16S
95 rRNA gene copies vary from 1-15 copies per genome, its use has continued to be the ‘gold
96 standard’ for microbial ecology (Case et al., 2007; Vos et al., 2012). Although an alternate target,
97 such as *rpoB*, which is a single copy gene would be valuable if assessing phylogenetic diversity,
98 there are no universal primers for it (Adékambi et al., 2009) as *rpoB* is not conserved enough to
99 be of use as a universal marker and only a subset of the microbial community can be targeted
100 (Vos et al., 2012). Many studies have used 16S rRNA gene copy numbers as a proxy for the total
101 bacterial community size, and some have found that the total estimated numbers of
102 proteobacteria species was not significantly different if using *rpoB* or 16S rRNA markers (Vos et
103 al., 2012). As this study has not assessed phylogenetic relationships of the microbial

104 communities, 16S rRNA is an appropriate target for the relative comparison of the overall
105 bacterial community size between environmental treatments/variables.

106 LUC resulting from displacement of corn-soybean rotations by PG production may alter soil
107 microhabitats and therefore soil microbial N-cycling due both to extensive root and rhizome
108 biomass and to large leaf litter inputs to soils in perennial vs. annual systems (Dohleman et al.,
109 2012). Within studies targeting soil microbial communities in biomass cropping systems to date
110 (Hedenec et al., 2014; Liang et al., 2012; Mao et al., 2013a, 2011; Orr et al., 2015), the effects of
111 various management practices (e.g. N fertilization and harvest) on soil microbial community
112 functioning have not been an area of focus. The effect this type of LUC may have on soil
113 microbial communities may depend on PG management practices in these systems.

114 Currently, there is no consensus regarding optimal N fertilization practices for increased yields in
115 PG production as yield responses can be highly variable depending on environmental conditions
116 and crop species (Deen et al., 2011). Depending on downstream use, miscanthus can be either
117 harvested in the fall pre-frost, harvested post-frost kill, or left to overwinter as standing biomass
118 for harvest in the spring. Switchgrass is commonly harvested in the fall, and is often windrowed
119 (cut, swathed and left on soil) over winter due to producers' limitations in collecting and storing
120 harvested biomass in winter (Resource Efficient Agricultural Production (REAP)-Canada, 2008;
121 Sokhansanj et al., 2009). Differences in N fertilizer requirements and harvest regimes may alter
122 C and N inputs (Attard et al., 2011) and may influence LUC impacts on soil denitrifier
123 community sizes.

124 Our objective was to compare the effects of LUC from corn-soybean to PG biomass production
125 on the relative abundances of total (16S rRNA gene target) and denitrifier (*nirS* and *nosZ* gene

126 targets) soil bacterial communities 3-4 years after PG planting. Soil was collected on four dates
127 from 2011 to 2012 from a field trial established in Ontario in 2008. This study is unique in that
128 it consists of two PG biomass crops produced in parallel with the existing common land use of
129 corn-soybean rotation within the same field trial site. It also includes unfertilized and fertilized
130 plots in both annual and perennial systems, and varied harvest practices within PG plots. We
131 hypothesized that soils from PG plots would support larger total bacterial and denitrifier
132 communities than soils from corn-soybean plots due to increased shoot residue return and root
133 inputs to soils in PG systems, and that soils from PG plots with biomass harvested in the spring
134 would support larger total bacterial and denitrifier communities than supported by soils from
135 PGs harvested in the fall due to increased root inputs and leaf loss to soil over winter.

136

137 **2. Materials and Methods**

138 **2.1 Site Description and Experimental Design**

139 A field trial was established in 2008 at the University of Guelph Research Station in Elora, ON
140 (43°38'46.73" N and 80°24'6.66" W). The field site was cultivated on May 16th and June 6th,
141 2008. Switchgrass (*Panicum virgatum* L., Shelter variety) was planted on June 6th, 2008.
142 Miscanthus (*M. sinensis* x *M. sacchariflorus*, Nagara-116 variety) was planted on June 12th,
143 2008, and soybean (*Glycine max* L.) was planted on June 24th, 2008 and annually rotated with
144 corn (*Zea mays* L.). Corn was planted on May 5th, 2010; soy was planted on June 3rd, 2011, and
145 corn was re-planted on May 18th, 2012, with annual light cultivation to prepare seedbeds for
146 planting. In 2007, prior to trial establishment, the experimental area was planted to barley
147 (*Hordeum vulgare* L.). The soil type is a London silt loam (Gray Brown Luvisol).

148 The field trial was a split-split strip plot design with three replicates. The main plot factor was
149 PG crop or annual rotation (miscanthus, switchgrass, and corn-soybean). Main treatment plots
150 measured 6.2 m x 26.0 m. Nitrogen fertilizer (0 or 160 kg N ha⁻¹) was applied in strips
151 randomly within replicates. 160 kg N ha⁻¹ subplots received hand-broadcast urea fertilizer (46-0-
152 0; N-P-K) in May 2011 or hand-broadcast ammonium nitrate fertilizer (34-0-0; N-P-K) in May
153 2012, after soil sampling procedures described below. Main treatments were split into two
154 harvest timings (fall or spring) within the PG fertilizer strips only. Miscanthus plots were either
155 harvested in the late fall season after post-frost kill, or left standing to overwinter until spring
156 harvest. Switchgrass plots were harvested in the fall, or cut and assembled into windrows in the
157 field for biomass removal in the spring. Spring-harvest of PGs occurred before N fertilizer was
158 applied. Harvest methods of PG yields (dry harvested biomass (tonnes) ha⁻¹) are described in
159 Deen et al. (2011). Figure 1 illustrates the seasonal and annual variation in daily average air
160 temperature (°C), and daily precipitation (mm) measured at the Elora Research Station.

161

162 **2.2 Soil Sampling and Analysis**

163 Baseline bulk density and carbon measurements were measured for each main plot on October
164 23rd, 2010. For bulk density, two soil cores per plot were collected at 0-5 cm depth using 2.5
165 cm-diameter cylindrical aluminum cores. Cores were weighed before and after drying for 24h at
166 105°C (Maynard & Curran, 2007). For soil carbon analysis, ten soil cores per plot were
167 collected from both 0-15 cm and 15-30 cm depths using a 5 cm-diameter soil corer on a zed-
168 shaped transect, and then composited per treatment plot for each depth. Total soil carbon and
169 inorganic carbon were analyzed with a Leco[®] Carbon Determinator CR-12 (Model No. 781-700,

170 Leco Instruments Ltd.) following the dry combustion technique (Périé and Ouimet, 2008) on
171 approximately 0.300 g of dried, ground and homogenized soil (Table 1).

172 For molecular analyses, soil was sampled on 4 dates (May 9th, 2011; October 30th, 2011; May
173 2nd, 2012 and October 20th, 2012). October sampling dates occurred before fall harvest of PG
174 crops, while May sampling dates occurred before N fertilizer application and after spring PG
175 biomass removal (Fig. 1). Ten soil cores per plot were sampled aseptically to 15 cm depth using
176 a 5 cm-diameter soil corer on a zed-shaped transect, composited and kept on ice until transport
177 back to the laboratory. The transect shape was chosen to encompass plot heterogeneity; at a pre-
178 trial study date initial analysis indicated gene abundances were not significantly different
179 between bulk or rhizosphere soils in corn-soybean or PG plots, possibly due to the large root
180 biomass/leaf loss to soils in perennial plots and residual soy/corn residue cover on soil in corn-
181 soybean plots. Soil samples were divided for storage at 4°C and -20°C.

182 Mean values of gravimetric soil moisture (g g^{-1}) are shown in Figure 1. Soil exchangeable NO_3^- -
183 N and NH_4^+ -N were determined for each of the soil samples by KCl extraction. Soil samples
184 (10.0 g) were placed into 125 mL flasks and 100 mL of 2.0M KCl was added to each flask.
185 Flasks were stoppered and shaken for 1h at 160 strokes per minute; solutions were allowed to
186 settle and were then filtered through Whatman no. 42 filter paper (Whatman plc, ME, U.S.A).
187 Extractable NO_3^- -N and NH_4^+ -N were determined colourmetrically with segmented flow
188 analyses (AA3, SEAL Analytical, Wisconsin, USA) via a cadmium reduction (Technician
189 Instrument Corporation, 1971), and a Berthelot reaction respectively (Fig. 2).

190 **2.3 Soil DNA Extraction**

191 Total DNA was extracted from field-moist soil sampled from each plot (3 field replicates, n=3;
192 total plots n=30). DNA was extracted in duplicate (ca. 0.250g) within 48 h of sampling as per
193 manufacturer's protocol using the DNA PowerSoil Kit (Mobio, Carlsbad, USA). Duplicate
194 extracts were then pooled, separated into aliquots, and stored at -80°C until use in downstream
195 analyses.

196 **2.4 Quantification of total bacteria and functional genes**

197 Quantitative PCR (qPCR) assays were used to enumerate the total bacterial communities (16S
198 rRNA gene), and communities of denitrifiers by targeting nitrite reductase (*nirS*) and nitrous
199 oxide reductase (*nosZ*) genes, using primer pairs 338f/518r (16S rRNA, Fierer et al., 2005),
200 Cd3af/R3Cd (*nirS*, Throbäck et al., 2004) and 1F/1R (*nosZ*, Henry et al., 2006).

201 For each gene target analyzed, duplicate replicates were run in parallel on an IQ5 thermocycler
202 (Bio-Rad Laboratories, Hercules, CA, USA). qPCR reaction mixtures contained 12.5 µL of 1x
203 SYBR Green Supermix, each forward and reverse primer at a final concentration of 400 nM, 1
204 µL of DNA template and RNase/DNase-free water to a final volume of 25 µL. The SYBR Green
205 Supermix contained 100 nM KCl, 40 mM Tris-HCl, 0.4 mM dNTPs, 50 units mL⁻¹ iTaq DNA
206 polymerase, 6 mM MgCl₂, SYBR Green 20 nM fluorescein, and stabilizer (Bio-Rad
207 Laboratories, Hercules, CA, USA).

208 Conditions for qPCR were an initiation step at 94°C for two minutes, followed by 35 cycles of
209 denaturing at 94°C for thirty seconds, annealing at 57°C for thirty seconds (16S rRNA) or at
210 55°C for sixty seconds (*nirS*), followed by elongation at 72°C for thirty (16S rRNA) or sixty
211 (*nirS*) seconds. For *nosZ*, a touchdown protocol adapted from Henry et al., (2006) was used.

212 Amplicon specificity was screened by running qPCR products on an ethidium bromide-stained
213 gel (1% agarose, 80 volts for 20 minutes) with a 100bp ladder, which resulted in clean bands for
214 all gene targets. The 16S rRNA primers used are degenerate and have been cited as having 89-
215 91% matching efficiency to all bacteria (Bergmark et al., 2012). The primers amplify one of two
216 conserved regions in V3 of the SSU rRNA gene, resulting in a ca. 200 bp amplicon that is within
217 small enough to amplify via qPCR methodology and amplifies for most bacterial taxa (Bakke et
218 al., 2011)

219 Known template standards were made from cloned PCR products from pure culture genomic
220 DNA (*Clostridium thermocellum* (16S), *Pseudomonas aeruginosa* (*nirS*), and *Pseudomonas*
221 *fluorescens* (*nosZ*)) and transformed into *Escherichia coli* plasmids (TOPO TA cloning kit);
222 plasmids were sequenced to confirm successful cloning and transformation of the target genes.

223 Amplicon specificity was screened by running PCR products on an ethidium bromide-stained gel
224 (1% agarose, 80 volts for 20 minutes) with a 100bp ladder. PCR amplicons of cloned gene
225 targets were sequenced by the Laboratory Services Department at the University of Guelph using
226 an ABI Prism 3720 (Applied Biosystems, Foster City, CA, USA) to confirm target identity.

227 In all qPCR assays, all unknown samples were amplified in parallel with a triplicate serial
228 dilution (10^1 - 10^8 gene copies per reaction) of control plasmids. PCR assays were optimized to
229 ensure efficiencies ranging from 93.0-106.4%, with R^2 's ranging from 0.990-0.999 and standard
230 curve slopes of -3.177 to -3.408 by testing serial dilutions of DNA extracts in order to minimize
231 inhibition of amplification due to humic and fulvic contaminants. Duplicate no-template
232 controls were run for each qPCR assay, which gave null or negligible values. Melt curve
233 analysis was used to confirm amplicon specificity. Normalization of DNA concentrations to

234 gram of dry soil was used to give results on a biologically significant scale, which assumes
235 similar DNA isolation efficiency across samples..

236 **2.5 Statistical Analysis**

237 Analysis of variance was conducted in SAS 9.3 (Carlsbad, NC, USA) using a generalized linear
238 mixed model (PROC GLIMMIX). The Shapiro-Wilks test was used to test for normality of data;
239 studentized test for residuals confirmed the absence of outliers. The probability distributions of
240 gene abundance data sets were log normal or highly skewed and were analyzed using an
241 overdispersed Poisson distribution for count data (Ver Hoef and Boveng, 2007). Bulk density,
242 organic carbon, total carbon, nitrate and ammonium data were log transformed when required
243 and fitted to the normal distribution.

244 Within each data set, sampling time was a repeated measure; independent and interactive fixed
245 effects were associated with crop/crop rotation, nitrogen application rate and harvest timing
246 within perennial grasses, while field replicate and its associated interactions were random effects.
247 The residual maximum likelihood method was employed to fit the model for all data sets.
248 Several covariance structures were entertained before the variance components structure was
249 chosen based on convergence and model fitting criteria. Individual treatment means within data
250 sets were compared using a post-hoc Tukey's test for all pairwise comparisons, with significance
251 denoted at $p < 0.05$.

252 Correlation analysis was used to assess nonparametric measures of statistical dependence
253 between gene abundances and H_2O , NO_3^- -N and NH_4^+ -N measured over time (Supplementary
254 Table 1). Correlation analysis resulted in multiple significant correlations between variables; as

255 such a principal component analysis was conducted in SAS (PROC FACTOR) on 120 samples
256 using a VARIMAX rotation.

257 3. Results

258 3.1 Environmental and Soil Conditions

259 Environmental conditions varied during the periods prior to the four soil sampling dates (Figure
260 1). Average air temperatures over the growing seasons (May-October) were 16.9°C and 17.3°C
261 in 2011 and 2012 respectively (Roy et al. 2014); average air temperatures in spring 2012 were
262 warmer than normal and resulted in earlier emergence of PG crops compared to 2011.

263 Cumulative monthly precipitation was above average prior to the May 2011 sampling date (101
264 mm vs. 72 mm 30-year average in April 2011 and 113 mm vs. 82 mm 30-year average in May
265 2011) (Roy et al., 2014). In comparison, Southern Ontario received very low cumulative
266 precipitation in April 2012 (30 mm vs. 72 mm 30-year average) and May 2012 (28 mm vs. 82
267 mm 30-year average) (Roy et al., 2014). Cumulative precipitation levels were lower in 2012
268 compared to 2011 from May-August (391 mm in 2011 vs. 186 mm in 2012), however higher
269 than normal precipitation levels occurred in October of 2011 (129 mm vs. 77 mm 30-year
270 average) and both September (106 mm vs. 77 mm 30-year average) and October (127 mm vs. 77
271 mm 30-year average) of 2012 (Roy et al., 2014). Environmental conditions prior to soil
272 sampling directly impact soil gravimetric content measured at time of sampling (Fig. 1), and
273 could also impact soil N and soil bacterial communities.

274 Soil physical and chemical properties were assessed in October 2010, after only two years of
275 contrasting management since crop establishment in 2008. The corn-soybean rotation had higher
276 soil bulk density than soils from both miscanthus and switchgrass plots harvested in the fall. No
277 differences in total or organic soil carbon were detected between the corn-soybean rotation and
278 the PG treatments at either the 0-15cm or 15-30cm depth (Table 1). Soil NH₄-N levels did not

279 differ significantly between the corn-soybean rotation and the PG soils, however N fertilization
280 significantly increased $\text{NH}_4\text{-N}$ levels in soils from fall-harvested miscanthus plots ($p < 0.05$) (Fig.
281 2a). N fertilization also significantly increased $\text{NO}_3\text{-N}$ levels in spring-harvested switchgrass
282 ($p < 0.05$) (Fig. 2b). From May to October 2011, soil $\text{NH}_4\text{-N}$ levels increased significantly and
283 soil $\text{NO}_3\text{-N}$ levels decreased significantly in PG soils (data not shown); a similar trend was not
284 observed in 2012 or for soils from the corn-soybean rotation in either year.

285 **3.2 Biomass Yields**

286 Despite significant differences in precipitation between 2011 and 2012, biomass yields of
287 miscanthus and switchgrass did not differ between years. In comparison, corn grain yields were
288 higher in 2011 vs 2012 (Table 1). Miscanthus produced higher yields (12.7-18.3 dry tonnes ha^{-1})
289 than either switchgrass or corn grain, regardless of N fertilization rate or harvest timing (Table
290 1). When harvested in the fall and N-fertilized, switchgrass yields were not significantly lower
291 (10.5-11.1 dry tonnes ha^{-1}) than miscanthus yields. Switchgrass yields from unfertilized plots
292 were not significantly different if harvested in the fall or spring; however, switchgrass yields
293 from fertilized plots were significantly higher (ca. 3-4 dry tonnes ha^{-1}) when harvested in the fall
294 compared to yields obtained when switchgrass was windrowed over winter.

295 **3.3 Bacterial Responses to Annual and Perennial Crops and their Management**

296 There was no statistically significant effect of N fertilization or any significant interactions
297 between cropping system and sampling time on any of the targeted gene abundances. Therefore
298 we analyzed the impact of each biomass crop under specific harvest management on soil
299 bacterial gene abundances (Table 2). Denitrifying (*nosZ*) gene copy abundances were affected
300 by LUC; regardless of harvest or N management, mean *nosZ* gene copies were higher in

301 miscanthus plots than in the corn-soybean rotation, and *nirS:nosZ* ratios were higher in the corn-
302 soybean soils than in miscanthus or switchgrass soils ($p < 0.05$) (Table 2). Under fall harvesting
303 management, biomass crop had no impact on total bacterial 16S rRNA gene copies or *nirS* gene
304 copies. However, leaving miscanthus biomass standing overwinter until spring resulted in
305 significantly higher 16S rRNA gene copies than observed in soils from fall-harvested biomass
306 crops and significantly higher *nirS* gene copies than in fall-harvested switchgrass or the corn-
307 soybean rotation (Table 2).

308 **3.3 Temporal Changes in Bacterial Gene Abundances**

309 Sampling date had a significant impact on gene abundances for all genes quantified (Fig. 3).
310 Over both sampling years, 16S rRNA gene copies were significantly higher ($5.2\text{-}5.4 \times 10^9$ gene
311 copies g^{-1} dry soil) at fall (October) sampling dates compared to the ca. $5.5\text{-}6.4 \times 10^8$ gene copies
312 g^{-1} dry soil quantified at spring (May) sampling dates (Fig. 3). Populations of *nirS* and *nosZ*
313 denitrifiers represented ca. 1.58% and 0.26% on a gene-to-gene basis (*nirS* or *nosZ* to 16S) of the
314 total bacterial community (data not shown), and did not follow similar trends with time of
315 sampling (Fig. 3). The abundance of *nirS* gene copies was significantly higher in 2012 (4.0×10^6
316 $\text{-} 1.6 \times 10^7$ gene copies g^{-1} dry soil) compared to 2011 ($2.5\text{-}6.3 \times 10^5$ gene copies g^{-1} dry soil),
317 with no significant differences between May and October sampling dates within each year (Fig.
318 3). The abundance of *nosZ* gene copies were approximately $1.3\text{-}3.2 \times 10^5$ gene copies g^{-1} dry
319 soil, but increased significantly in May 2012 to approximately 3.2×10^6 gene copies g^{-1} dry soil
320 and dropped back to previous levels by October 2012 (Fig. 3). Higher relative proportions of
321 denitrifiers (*nirS* or *nosZ* to 16S) were observed at spring sampling dates, when total bacterial
322 16S rRNA gene abundances decreased in comparison to fall sampling dates (Fig. 3).

323 Two factors were selected in the principal components analysis, which accounted for 67.73%
324 cumulative variance. A scree plot was examined for breaks and factors were retained when
325 eigenvalues ≥ 1 . Soil $\text{NH}_4\text{-N}^+$, soil $\text{NO}_3\text{-N}$, *nirS* and *nosZ* loaded on factor 1, which accounted
326 for 43.89% variance while soil gravimetric H_2O and 16S rRNA loaded on factor 2, which
327 accounted for 23.84% variance (Fig.4 a and b). Differences in soil $\text{NO}_3\text{-N}$ and $\text{NH}_4\text{-N}$ were
328 strongly related to differences in *nirS* and *nosZ* gene abundances observed between May 2011
329 and May 2012 sampling dates (Fig. 3 and Fig. 4), while the size of the total bacterial community
330 (16S rRNA) was related to soil gravimetric moisture levels (Fig. 4).

331 **4. Discussion**

332 Denitrification is an important process contributing to the production and consumption of N_2O in
333 soils, and mitigation of GHGs such as N_2O is required to create sustainable biomass cropping
334 systems (Miller et al., 2008; Schlesinger, 2013). Changes in the potential functional abilities of
335 the soil microbial community may reflect changes in LUC or agricultural management and
336 should be considered to assess the ecological impact of biomass crop production (Hedenec et al.,
337 2014). Currently, few studies have assessed soil microbial community responses to PG biomass
338 production systems (Hedenec et al., 2014; Liang et al., 2012; Mao et al., 2013a, 2011; Orr et al.,
339 2015). The highest potential to reduce GHG emissions from biomass cropping systems is to
340 produce crops with high yields, such as PGs (Sanscartier et al., 2014), which offset the amount of
341 land required for crop production (Kludze et al., 2013). However, if PG biomass production
342 negatively affects soil health as indicated by changes in the potential functioning of microbial
343 communities, large-scale LUC from annual to perennial biomass production may not be as
344 sustainable as originally proposed. As such it is necessary to identify biomass cropping systems

345 that not only result in large biomass yields, but also ensure agroecosystem sustainability by
346 maintaining or improving ecosystem services (Orr et al., 2015), such as soil N-cycling.

347 **4.1 Biomass Yields of Annual and Perennial Crops**

348 Miscanthus and switchgrass biomass yields were within the typical range of values reported
349 previously in Ontario (Kludze et al., 2013; Resource Efficient Agricultural Production (REAP)-
350 Canada, 2008) and Europe (Christian et al., 2008; Himken et al., 1997), despite differences in
351 temperature and precipitation between the two study years. Corn grain yields were within the
352 lower range for reported Ontario yields (Munkholm et al., 2013), potentially due to wetter (2011)
353 and drier (2012) field conditions than normal over the two growing seasons (Roy et al., 2014).
354 Deen et al. (2011) showed increases in PG biomass yields between the second and third years
355 after PG planting at our site, whereas we measured similar yields in 2011 and 2012, indicating
356 the PGs may have reached maximum yield potential.

357 Nitrogen fertilization significantly increased corn grain yields and fall-harvested switchgrass
358 biomass yields, however no significant increases due to N fertilization were observed in
359 miscanthus or spring-harvested switchgrass biomass yields. Potential yield increases from N
360 fertilization in spring-harvested switchgrass may have been offset due to leaf loss over the winter
361 season, as increases in switchgrass yields to N fertilization have been previously observed
362 (Nikièma et al., 2011; Vogel et al., 2002). Similar to the present study, European and US field
363 trials have also found no response of miscanthus yields to N (Lewandowski et al., 2000;
364 Lewandowski et al., 2003; Behnke et al., 2012; Christian et al., 2008), and PG yields were
365 minimally impacted by differences in growing season conditions compared to corn grain yields
366 (Table 1).

367 Despite significant differences in biomass yields between miscanthus and corn-soybean systems,
368 there were no significant differences in either total or organic soil carbon between any of
369 cropping systems assessed (Table 1). Sampling of soil carbon occurred only two years after PG
370 planting; PGs are expected to be productive for 20+ years, indicating future changes in soil
371 carbon levels may occur. Additionally, Ontario-based land conversion modelling scenarios have
372 estimated a soil carbon decrease of 2.5% upon miscanthus establishment (Sanscartier et al.,
373 2014), which may have negated potential increases in soil organic carbon. However, high
374 miscanthus yields most likely resulted in increases in above and below-ground plant residue
375 return to soils (Mutegi et al., 2010; Soil Quality National, 2006); therefore our carbon measures
376 may not have reflected short-term changes in labile carbon sources that had occurred.
377 Regardless of management or climatic conditions, miscanthus consistently produced large yields,
378 emphasizing its potential as a bioenergy crop suitable for production in variable Ontario
379 conditions.

380 **4.2 Bacterial Responses to Annual and Perennial Crops and their Management**

381 Some studies in biomass cropping systems have not observed differences in soil microbial
382 responses between perennial and annual crop types (Mao et al., 2011), while others have
383 measured significant differences in microbial abundance, diversity and community structure
384 between these cropping types (Liang et al., 2012; Morales et al., 2010; Watrud et al., 2013).
385 Currently, we observed significantly higher *nosZ* gene copies in miscanthus soils compared to
386 corn-soybean soils, illustrating a distinct effect of LUC from corn-soybean to miscanthus
387 production on soil N cycling (Table 2).

388 Due to the large biomass produced by miscanthus compared to corn, a large amount of plant
389 residues are returned to the soil; these residues provide surface cover, decrease soil bulk density,
390 increase water retention and regulate temperatures (Blanco-Canqui and Lal, 2009). Previous
391 work at the Elora Research Station found an inverse correlation between field-scale N₂O fluxes
392 and *nosZ* transcript abundance in conventionally-tilled corn plots with residues returned to soils
393 (Németh et al., 2014), and increased *nosZ* activity after residue amendment has also been
394 observed in lab studies (Henderson et al., 2010). High C:N plant residues have been negatively
395 correlated with cumulative N₂O emissions (Huang, 2004), and may encourage complete
396 reduction of N₂O to N₂ as soil available NO₃-N is limiting, so bacterial populations with the
397 ability to reduce N₂O to N₂ are favoured (Miller et al., 2008). Presently, the primers used for
398 *nosZ* gene target amplification provided good coverage of γ -Proteobacteria (Henry et al., 2006),
399 which are stimulated by surface-applied residues (Pascault et al, 2010). Increased residue return
400 in miscanthus plots may have selected for bacterial populations harbouring enhanced catabolic
401 capabilities, such as N₂O reduction (Pascault et al., 2010). This implies that producing biomass
402 crops with large yields may indirectly alter soil N cycling and potentially mitigate soil N₂O
403 emissions due to increased residue return influencing the soil microbial community. It is likely
404 that differences in environmental conditions (e.g. temperature, H₂O and O₂ availability) and
405 resource quality and availability between corn-soybean and miscanthus soils related to
406 differences in microbial community structure (Cusack et al., 2011) and selected for different
407 dominant taxa that filled different ecological niches (Stone et al., 2015).

408 N fertilization did not affect targeted gene abundances, however studies in other cropping
409 systems have found that N fertilization affected the size of denitrifying communities (Hallin et
410 al., 2009), nitrifying communities (He et al., 2007), and proportions of *nirS* to *nirK* communities

411 (ratio of *nirS:nirK* genes) (Hai et al., 2009). Elevated 16S rRNA and *nirS* gene copies were
412 observed in soils from spring-harvested miscanthus and windrowed switchgrass (Table 2).
413 Increased N return via senescent leaf loss in PG plots over winter contributes to the soil organic
414 matter pool (Heaton et al., 2009), and may have contributed to elevated total (16S rRNA)
415 bacterial populations in these soils, concomitantly increasing *nirS* abundances (Huang et al.,
416 2011).

417 **4.3 Temporal Changes in Bacterial Gene Abundances**

418 Total soil bacterial communities (16S rRNA) followed a seasonal trend, with elevated 16S rRNA
419 gene copies at fall (October) compared to spring (May) sampling dates, possibly due to an
420 increase in the availability and diversity of resources for microbial metabolism and growth over
421 the growing season (Habekost et al., 2008). Denitrifying abundances changed differently than
422 the total bacterial community, suggesting denitrifiers were influenced by different proximal
423 regulators than the total bacterial community (Fig. 3 and 4). Seasonal dynamics of N-cycling
424 microbial communities have been previously characterized (Boyer et al., 2006; Nemeth et al.,
425 2014; Wolsing and Priemé, 2004; Dandie et al., 2008; Bremer et al., 2007), and are tightly
426 coupled with seasonal changes in labile C and N pools, temperature and soil H₂O (Butterbach-
427 Bahl et al. 2013; Rasche et al., 2011), indicating that local edaphic drivers may often take
428 precedence over crop-specific drivers (Mao et al., 2013).

429 **5.0 Conclusions**

430 Miscanthus consistently produced large yields and supported larger *nosZ*-bearing communities
431 than the corn-soybean rotation, emphasizing its influence on soil N cycling and its potential to
432 mitigate soil N₂O emissions while being suitable for production in variable Ontario conditions.

433 Additionally, miscanthus yields were not increased with N fertilization, indicating a lower N
434 input requirement for biomass production compared to switchgrass. Higher 16S rRNA and *nirS*
435 gene copies were associated with reduced yields in spring-harvested PGs, indicating that
436 harvesting PGs in the spring may increase populations of denitrifiers capable of producing N₂O
437 emissions while simultaneously decreasing biomass yields. The size of both denitrifying (*nirS*
438 and *nosZ*) and total bacterial (16S rRNA) communities changed over the sampling period,
439 however changes in denitrifying gene abundances did not parallel changes in the total soil
440 bacterial community, indicating denitrifying communities were regulated differently than the
441 total bacterial community. Future work measuring N₂O emissions and denitrifier activity
442 (mRNA) and community structure in these systems is required to link the effects of LUC on
443 these communities directly with N₂O fluxes.

444 **Author Contributions:**

445 K. Thompson was the primary researcher and author on this study, conducting all field work, lab
446 work, and manuscript preparation. Author B. Deen is credited for the use of his OMAFRA-
447 funded field plots for this research, and valuable advice on experimental design, statistical
448 analyses and manuscript focus. Author K. Dunfield is credited for her invaluable mentorship on
449 molecular analyses and trouble-shooting on field sampling and sample preservation techniques,
450 manuscript focus and preparation, and data interpretation.

451

452 **Acknowledgements**

453 Funding for this research was awarded through grants from the Natural Science and Engineering
454 Research Council of Canada (NSERC), the Canada Research Chairs (CRC) Program and the
455 Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs (OMAF/MRA).

456 K.T. was also supported through an Ontario Graduate Scholarship (OGS) and an Ontario
457 Graduate Fellowship (OGF). Special thanks to Henk Wichers (University of Guelph) for his
458 expertise in managing the Alternate Renewable Fuels trial at the Elora Research Station.

459 **References**

- 460 Adékambi, T., Drancourt, M., Raoult, D., 2009. The *rpoB* gene as a tool for clinical
461 microbiologists. *Trends Microbiol.* 17, 37–45. doi:10.1016/j.tim.2008.09.008
- 462 Attard, E., Recous, S., Chabbi, A., De Berranger, C., Guillaumaud, N., Labreuche, J., Philippot,
463 L., Schmid, B., Le Roux, X., 2011. Soil environmental conditions rather than denitrifier
464 abundance and diversity drive potential denitrification after changes in land uses. *Glob.*
465 *Chang. Biol.* 17, 1975–1989. doi:10.1111/j.1365-2486.2010.02340.x
- 466 Bakke, I., De Schryver, P., Boon, N., Vadstein, O., 2011. PCR-based community structure
467 studies of Bacteria associated with eukaryotic organisms: A simple PCR strategy to avoid
468 co-amplification of eukaryotic DNA. *J. Microbiol. Methods* 84, 349–351.
469 doi:10.1016/j.mimet.2010.12.015
- 470 Bastian, F., Bouziri, L., Nicolardot, B., Ranjard, L., 2009. Impact of wheat straw decomposition
471 on successional patterns of soil microbial community structure. *Soil Biol. Biochem.* 41,
472 262–275. doi:10.1016/j.soilbio.2008.10.024
- 473 Bergmark, L., Poulsen, P.H.B., Al-Soud, W.A., Norman, A., Hansen, L.H., Sørensen, S.J., 2012.
474 Assessment of the specificity of Burkholderia and Pseudomonas qPCR assays for detection
475 of these genera in soil using 454 pyrosequencing. *FEMS Microbiol. Lett.* 333, 77–84.
476 doi:10.1111/j.1574-6968.2012.02601.x
- 477 Bissett, A., Richardson, A.E., Baker, G., Thrall, P.H., 2011. Long-term land use effects on soil
478 microbial community structure and function. *Appl. Soil Ecol.* 51, 66–78.
479 doi:10.1016/j.apsoil.2011.08.010
- 480 Blanco-Canqui, H., Lal, R., 2009. Crop Residue Removal Impacts on Soil Productivity and
481 Environmental Quality. *CRC. Crit. Rev. Plant Sci.* 28, 139–163.
482 doi:10.1080/07352680902776507
- 483 Boyer, E.W., Alexander, R.B., Parton, W.J., Li, C., Butterbach-Bahl, K., Donner, S.D., Skaggs,
484 R.W., Del Grosso, S.J., 2006. Modeling denitrification in terrestrial and aquatic ecosystems
485 at regional scales. *Ecol. Appl.* 16, 2123–42.
- 486 Braker, G., Conrad, R., 2011. Diversity, structure, and size of N₂O-producing microbial
487 communities in soils-what matters for their functioning?, *Advances in applied*
488 *microbiology.* Elsevier. doi:10.1016/B978-0-12-387046-9.00002-5
- 489 Bremer, C., Braker, G., Matthies, D., Reuter, A., Engels, C., Conrad, R., 2007. Impact of plant
490 functional group, plant species, and sampling time on the composition of nirK-type
491 denitrifier communities in soil. *Appl. Environ. Microbiol.* 73, 6876–84.
492 doi:10.1128/AEM.01536-07
- 493 Butterbach-Bahl, K., Baggs, E.M., Dannenmann, M., Kiese, R., Zechmeister-Boltenstern, S.,
494 2013. Nitrous oxide emissions from soils: how well do we understand the processes and
495 their controls? *Philos. Trans. R. Soc. B Biol. Sci* 368, 1–13. doi:10.1098/rstb.2013.0122
- 496 Cantera, J.J.L., Stein, L.Y., 2007. Role of nitrite reductase in the ammonia-oxidizing pathway of
497 *Nitrosomonas europaea*. *Arch. Microbiol.* 188, 349–54. doi:10.1007/s00203-007-0255-4
- 498 Casciotti, K.L., Ward, B.B., 2001. Dissimilatory Nitrite Reductase Genes from Autotrophic
499 Ammonia-Oxidizing Bacteria. *Appl. Environ. Microbiol.* 67, 2213–2221.

500 doi:10.1128/AEM.67.5.2213

501 Case, R.J., Boucher, Y., Dahllöf, I., Holmström, C., Doolittle, W.F., Kjelleberg, S., 2007. Use of
502 16S rRNA and rpoB genes as molecular markers for microbial ecology studies. *Appl.*
503 *Environ. Microbiol.* 73, 278–88. doi:10.1128/AEM.01177-06

504 Christian, D.G., Riche, a. B., Yates, N.E., 2008. Growth, yield and mineral content of
505 *Miscanthus×giganteus* grown as a biofuel for 14 successive harvests. *Ind. Crops Prod.* 28,
506 320–327. doi:10.1016/j.indcrop.2008.02.009

507 Cuhel, J., Simek, M., Laughlin, R.J., Bru, D., Cheneby, D., Watson, C.J., Philippot, L., Chèneby,
508 D., 2010. Insights into the effect of soil pH on N₂O and N₂ emissions and denitrifier
509 community size and activity. *Appl. Environ. Microbiol.* 76, 1870–8.
510 doi:10.1128/AEM.02484-09

511 Cusack, D.F., Silver, W.L., Torn, M.S., Burton, S.D., Firestone, M.K., 2011. Changes in
512 microbial community characteristics and soil organic matter with nitrogen additions in two
513 tropical forests. *Ecology* 92, 621–632. doi:10.1890/10-0459.1

514 Dandie, C.E., Burton, D.L., Zebarth, B.J., Henderson, S.L., Trevors, J.T., Goyer, C., 2008.
515 Changes in bacterial denitrifier community abundance over time in an agricultural field and
516 their relationship with denitrification activity. *Appl. Environ. Microbiol.* 74, 5997–6005.
517 doi:10.1128/AEM.00441-08

518 Deen, B.B., Young, D., Rowsell, J., Tubeileh, A., Engbers, H., Rosser, B., 2011. A comparative
519 study assessing variety and management effects on C 4 perennial grasses in a northern
520 climate. *Asp. Appl. Biol.* 112, 205–212.

521 Deslippe, J.R., Jamali, H., Jha, N., Saggari, S., 2014. Soil Biology & Biochemistry Denitrifier
522 community size , structure and activity along a gradient of pasture to riparian soils. *Soil*
523 *Biol. Biochem.* 71, 48–60. doi:10.1016/j.soilbio.2014.01.007

524 Dohleman, F.G., Heaton, E. a., Arundale, R. a., Long, S.P., 2012. Seasonal dynamics of above-
525 and below-ground biomass and nitrogen partitioning in *Miscanthus × giganteus* and
526 *Panicum virgatum* across three growing seasons. *GCB Bioenergy* 4, 534–544.
527 doi:10.1111/j.1757-1707.2011.01153.x

528 Enwall, K., Throback, I.N., Stenberg, M., Söderström, M., Hallin, S., 2010. Soil resources
529 influence spatial patterns of denitrifying communities at scales compatible with land
530 management. *Appl. Environ. Microbiol.* 76, 2243–50. doi:10.1128/AEM.02197-09

531 Fierer, N., Jackson, J.A., Vilgalys, R., Jackson, R.B., 2005. Assessment of Soil Microbial
532 Community Structure by Use of Taxon-Specific Quantitative PCR Assays. *Appl. Environ.*
533 *Microbiol.* 71, 4117–4120. doi:10.1128/AEM.71.7.4117

534 Foster, J.L., Guretzky, J.A., Huo, C., Kering, M.K., Butler, T.J., 2013. Effects of Row Spacing,
535 Seeding Rate, and Planting Date on Establishment of Switchgrass. *Crop Sci.* 53, 309–314.
536 doi:10.2135/cropsci2012.03.0171

537 Gaudin, A.C.M., Janovicek, K., Deen, B., Hooker, D.C., 2015. Wheat improves nitrogen use
538 efficiency of maize and soybean-based cropping systems. *Agric. Ecosyst. Environ.* 210, 1–
539 10. doi:10.1016/j.agee.2015.04.034

540 Habekost, M., Eisenhauer, N., Scheu, S., Steinbeiss, S., Weigelt, A., Gleixner, G., 2008.
541 Seasonal changes in the soil microbial community in a grassland plant diversity gradient

542 four years after establishment. *Soil Biol. Biochem.* 40, 2588–2595.
543 doi:10.1016/j.soilbio.2008.06.019

544 Hai, B., Diallo, N.H., Sall, S., Haesler, F., Schauss, K., Bonzi, M., Assigbetse, K., Chotte, J.-L.,
545 Munch, J.C., Schloter, M., 2009. Quantification of key genes steering the microbial nitrogen
546 cycle in the rhizosphere of sorghum cultivars in tropical agroecosystems. *Appl. Environ.*
547 *Microbiol.* 75, 4993–5000. doi:10.1128/AEM.02917-08

548 Hallin, S., Jones, C.M., Schloter, M., Philippot, L., 2009. Relationship between N-cycling
549 communities and ecosystem functioning in a 50-year-old fertilization experiment. *ISME J.*
550 3, 597–605. doi:10.1038/ismej.2008.128

551 He, J.-Z., Shen, J.-P., Zhang, L.-M., Zhu, Y.-G., Zheng, Y.-M., Xu, M.-G., Di, H., 2007.
552 Quantitative analyses of the abundance and composition of ammonia-oxidizing bacteria and
553 ammonia-oxidizing archaea of a Chinese upland red soil under long-term fertilization
554 practices. *Environ. Microbiol.* 9, 2364–74. doi:10.1111/j.1462-2920.2007.01358.x

555 Hedenc, P., Novotny, D., Ustak, S., Cajthaml, T., Slejska, A., Simackova, H., Honzik, R.,
556 Kovarova, M., Frouz, J., 2014. The effect of native and introduced biofuel crops on the
557 composition of soil biota communities. *Biomass and Bioenergy* 60, 137–146.
558 doi:10.1016/j.biombioe.2013.11.021

559 Henderson, S.L., Dandie, C.E., Patten, C.L., Zebarth, B.J., Burton, D.L., Trevors, J.T., Goyer, C.,
560 2010. Changes in denitrifier abundance, denitrification gene mRNA levels, nitrous oxide
561 emissions, and denitrification in anoxic soil microcosms amended with glucose and plant
562 residues. *Appl. Environ. Microbiol.* 76, 2155–64. doi:10.1128/AEM.02993-09

563 Henry, S., Bru, D., Stres, B., Hallet, S., Philippot, L., 2006. Quantitative detection of the *nosZ*
564 gene, encoding nitrous oxide reductase, and comparison of the abundances of 16S rRNA,
565 *narG*, *nirK*, and *nosZ* genes in soils. *Appl. Environ. Microbiol.* 72, 5181–9.
566 doi:10.1128/AEM.00231-06

567 Himken, M., Lammel, J., Neukirchen, D., Olf, H., 1997. Cultivation of *Miscanthus* under West
568 European conditions : Seasonal changes in dry matter production , nutrient uptake and
569 remobilization. *Plant Soil* 189, 117–126.

570 Huang, S., Chen, C., Yang, X., Wu, Q., Zhang, R., 2011. Distribution of typical denitrifying
571 functional genes and diversity of the *nirS*-encoding bacterial community related to
572 environmental characteristics of river sediments. *Biogeosciences* 8, 3041–3051.
573 doi:10.5194/bg-8-3041-2011

574 Huang, Y., 2004. Nitrous oxide emissions as influenced by amendment of plant residues with
575 different C:N ratios. *Soil Biol. Biochem.* 36, 973–981. doi:10.1016/j.soilbio.2004.02.009

576 Intergovernmental Panel on Climate Change (IPCC) (2007) *Climate Change 2007: The Physical*
577 *Science Basis. Working Group I Contribution to the Fourth Assessment Report of the*
578 *Inter- governmental Panel on Climate Change. Solomon et al. (eds). Cambridge*
579 *University Press, Cambridge.*

580 Jayasundara, S., Wagner-Riddle, C., Dias, G., Kariyapperuma, K. a., 2014. Energy and
581 greenhouse gas intensity of corn (*Zea mays* L.) production in Ontario: A regional
582 assessment. *Can. J. Soil Sci.* 94, 77–95. doi:10.4141/cjss2013-044

583 Kludze, H., Deen, B., Weersink, A., van Acker, R., Janovicek, K., De Laporte, A., 2013. Impact
584 of land classification on potential warm season grass biomass production in Ontario,
585 Canada. *Can. J. Plant Sci.* 93, 249–260. doi:10.4141/cjps2012-143

586 Lewandowski, I., Andersson, B., Basch, G., Christian, D.G., Jørgensen, U., Jones, M.B., Riche,
587 A.B., Schwarz, K.U., Tayebi, K., Teixeira, F., 2003. Environment and Harvest Time Affects
588 the Combustion Qualities of of *Miscanthus* Genotypes. *Agron. J.* 95, 1274–1280.

589 Lewandowski, I., Clifton-brown, J.C., Scurlock, J.M.O., Huisman, W., 2000. *Miscanthus* :
590 European experience with a novel energy crop. *Biomass and Bioenergy* 19, 209–227.

591 Liang, C., Jesus, E.D.C., Duncan, D.S., Jackson, R.D., Tiedje, J.M., Balsler, T.C., 2012. Soil
592 microbial communities under model biofuel cropping systems in southern Wisconsin, USA:
593 Impact of crop species and soil properties. *Appl. Soil Ecol.* 54, 24–31.
594 doi:10.1016/j.apsoil.2011.11.015

595 Mao, Y., Yannarell, A.C., Davis, S.C., Mackie, R.I., 2013a. Impact of different bioenergy crops
596 on N-cycling bacterial and archaeal communities in soil. *Environ. Microbiol.* 15, 928–42.
597 doi:10.1111/j.1462-2920.2012.02844.x

598 Mao, Y., Yannarell, A.C., Davis, S.C., Mackie, R.I., 2013b. Impact of different bioenergy crops
599 on N-cycling bacterial and archaeal communities in soil: Impact of bioenergy crops on soil
600 N-cycling archaea and bacteria. *Environ. Microbiol.* 15, 928–942. doi:10.1111/j.1462-
601 2920.2012.02844.x

602 Mao, Y., Yannarell, A.C., Mackie, R.I., 2011. Changes in N-transforming archaea and bacteria
603 in soil during the establishment of bioenergy crops. *PLoS One* 6, 1–12.
604 doi:10.1371/journal.pone.0024750

605 Maynard, D.G. and Curran, M.P. 2007. Bulk Density Measurement in Forest Soil. Chapter 66.
606 In: *Soil Sampling and methods of analysis.* (Carter M.R. & Gregorich E.G.) Canadian
607 Society of Soil Science. CRC Press, Boca Raton, FL. 863-869.

608 Miller, M.N., Zebarth, B.J., Dandie, C.E., Burton, D.L., Goyer, C., Trevors, J.T., 2008. Crop
609 residue influence on denitrification, N₂O emissions and denitrifier community abundance in
610 soil. *Soil Biol. Biochem.* 40, 2553–2562. doi:10.1016/j.soilbio.2008.06.024

611 Morales, S.E., Cosart, T., Holben, W.E., 2010. Bacterial gene abundances as indicators of
612 greenhouse gas emission in soils. *ISME J.* 4, 799–808. doi:10.1038/ismej.2010.8

613 Munkholm, L.J., Heck, R.J., Deen, B., 2013. Long-term rotation and tillage effects on soil
614 structure and crop yield. *Soil Tillage Res.* 127, 85–91. doi:10.1016/j.still.2012.02.007

615 Mutege, J.K., Munkholm, L.J., Petersen, B.M., Hansen, E.M., Petersen, S.O., 2010. Nitrous
616 oxide emissions and controls as influenced by tillage and crop residue management
617 strategy. *Soil Biol. Biochem.* 42, 1701–1711. doi:10.1016/j.soilbio.2010.06.004

618 Németh, D.D., Wagner-Riddle, C., Dunfield, K.E., 2014. Abundance and gene expression in
619 nitrifier and denitrifier communities associated with a field scale spring thaw N₂O flux
620 event. *Soil Biol. Biochem.* 73, 1–9. doi:10.1016/j.soilbio.2014.02.007

621 Nogales, B., Timmis, K.N., Nedwell, D.B., Osborn, A.M., 2002. Detection and Diversity of
622 Expressed Denitrification Genes in Estuarine Sediments after Reverse Transcription-PCR
623 Amplification from mRNA. *Appl. Environ. Microbiology* 68, 5017–5025.

624 doi:10.1128/AEM.68.10.5017
625 Ontario Ministry of Energy, The Government of Ontario (2014). Achieving Balance: Ontario's
626 Long-Term Energy Plan. [Online] Available: <http://www.energy.gov.on.ca/>

627 Orr, M.-J., Gray, M.B., Applegate, B., Volenec, J.J., Brouder, S.M., Turco, R.F., 2015.
628 Transition to second generation cellulosic biofuel production systems reveals limited
629 negative impacts on the soil microbial community structure. *Appl. Soil Ecol.* 95, 62–72.
630 doi:10.1016/j.apsoil.2015.06.002

631 Pascault, N., Nicolardot, B., Bastian, F., Thiébeau, P., Ranjard, L., Maron, P.-A.A., 2010. In situ
632 dynamics and spatial heterogeneity of soil bacterial communities under different crop
633 residue management. *Microb. Ecol.* 60, 291–303. doi:10.1007/s00248-010-9648-z

634 Périé, C., Ouimet, R., 2008. Organic carbon, organic matter and bulk density relationships in
635 boreal forest soils. *Can. J. Soil Sci.* 88, 315–325. doi:10.4141/CJSS06008

636 Petersen, D.G., Blazewicz, S.J., Firestone, M., Herman, D.J., Turetsky, M., Waldrop, M., 2012.
637 Abundance of microbial genes associated with nitrogen cycling as indices of
638 biogeochemical process rates across a vegetation gradient in Alaska. *Environ. Microbiol.*
639 14, 993–1008. doi:10.1111/j.1462-2920.2011.02679.x

640 Philippot, L., 2002. Denitrifying genes in bacterial and Archaeal genomes. *Biochim. Biophys.*
641 *Acta* 1577, 355–76.

642 Philippot, L., 2006. Use of functional genes to quantify denitrifiers in the environment. *Biochem.*
643 *Soc. Trans.* 34, 101–3. doi:10.1042/BST0340101

644 Philippot, L., Andert, J., Jones, C., Bru, D., Hallin, S., 2011. Importance of denitrifiers lacking
645 the genes encoding the nitrous oxide reductase for N₂O emissions from soil. *Glob. Chang.*
646 *Biol.* 17, 1497–1504. doi:10.1111/j.1365-2486.2010.02334.x

647 Rasche, F., Knapp, D., Kaiser, C., Koranda, M., Kitzler, B., Zechmeister-Boltenstern, S.,
648 Richter, A., Sessitsch, A., 2011. Seasonality and resource availability control bacterial and
649 archaeal communities in soils of a temperate beech forest. *ISME J.* 5, 389–402.
650 doi:10.1038/ismej.2010.138

651 Reeve, J.R., Schadt, C.W., Carpenter-boggs, L., Kang, S., Zhou, J., Reganold, J.P., 2010. Effects
652 of soil type and farm management on soil ecological functional genes and microbial
653 activities. *ISME J.* 4, 1099–1107. doi:10.1038/ismej.2010.42

654 Resource Efficient Agricultural Production (REAP)-Canada, 2008. ARF07 P ROJECT
655 Optimization of Switchgrass Management for Commercial Fuel Pellet Production.
656 OMAFRA-Alternative Renew. Fuels Res. Dev. Fund.

657 Roy, A.K., Wagner-Riddle, C., Deen, B., Lauzon, J., Bruulsema, T., 2014. Nitrogen application
658 rate, timing and history effects on nitrous oxide emissions from corn (*Zea mays* L.). *Can. J.*
659 *Soil Sci.* 94, 563–573. doi:10.4141/cjss2013-118

660 Sanscartier, D., Deen, B., Dias, G., MacLean, H.L., Dadfar, H., McDonald, I., Kludze, H., 2014.
661 Implications of land class and environmental factors on life cycle GHG emissions of
662 *Miscanthus* as a bioenergy feedstock. *GCB Bioenergy* 6, 401–413. doi:10.1111/gcbb.12062

663 Schlesinger, W.H., 2013. An estimate of the global sink for nitrous oxide in soils. *Glob. Chang.*
664 *Biol.* 19, 2929–2931. doi:10.1111/gcb.12239

665 Soil Quality National, 2006. Crop Residue Removal For Biomass Energy Production: Effects on
666 Soils and Recommendations. Tech. Note 19, 1–7.

667 Sokhansanj, S., Mani, S., Turhollow, A., Kumar, A., Bransby, D., Laser, L.L., Laser, M., 2009.
668 Large-scale production, harvest and logistics of switchgrass (*Panicum virgatum* L .) –
669 current technology and envisioning a mature technology. *Biofuels Bioprod. Bioprocess.*
670 124–141. doi:10.1002/bbb

671 Stone, M.M., Kan, J., Plante, A.F., 2015. Parent material and vegetation influence bacterial
672 community structure and nitrogen functional genes along deep tropical soil profiles at the
673 Luquillo Critical Zone Observatory. *Soil Biol. Biochem.* 80, 273–282.
674 doi:10.1016/j.soilbio.2014.10.019

675 Szukics, U., Abell, G.C.J., Hödl, V., Mitter, B., Sessitsch, A., Hackl, E., Zechmeister-
676 Boltenstern, S., 2010. Nitrifiers and denitrifiers respond rapidly to changed moisture and
677 increasing temperature in a pristine forest soil. *FEMS Microbiol. Ecol.* 72, 395–406.
678 doi:10.1111/j.1574-6941.2010.00853.x

679 Throbäck, I.N., Enwall, K., Jarvis, A., Hallin, S., 2004. Reassessing PCR primers targeting nirS,
680 nirK and nosZ genes for community surveys of denitrifying bacteria with DGGE. *FEMS*
681 *Microbiol. Ecol.* 49, 401–17. doi:10.1016/j.femsec.2004.04.011

682 Ver Hoef, J.M. and Boveng, P.L., 2007. Quasi-Poisson vs. Negative Binomial Regression: How
683 should we model overdispersed count data? *Ecology* 88, 2766–2772.

684 Vos, M., Quince, C., Pijl, A.S., Hollander, M. De, Kowalchuk, G.A., 2012. A Comparison of
685 rpoB and 16S rRNA as Markers in Pyrosequencing Studies of Bacterial Diversity 7, 1–8.
686 doi:10.1371/journal.pone.0030600

687 Watrud, L.S., Reichman, J.R., Bollman, M. a., Smith, B.M., Lee, E.H., Jastrow, J.D., Casler,
688 M.D., Collins, H.P., Fransen, S., Mitchell, R.B., Owens, V.N., Bean, B., Rooney, W.L.,
689 Tyler, D.D., King, G. a., 2013. Chemistry and Microbial Functional Diversity Differences
690 in Biofuel Crop and Grassland Soils in Multiple Geographies. *BioEnergy Res.* 6, 601–619.
691 doi:10.1007/s12155-012-9279-y

692 Wolsing, M., Priemé, A., 2004. Observation of high seasonal variation in community structure of
693 denitrifying bacteria in arable soil receiving artificial fertilizer and cattle manure by
694 determining T-RFLP of nir gene fragments. *FEMS Microbiol. Ecol.* 48, 261–71.
695 doi:10.1016/j.femsec.2004.02.002

696 Wu, L., Osmond, D.L., Graves, A.K., Burchell, M.R., Duckworth, O.W., 2012. Relationships
697 Between Nitrogen Transformation Rates and Gene Abundance in a Riparian Buffer Soil.
698 *Environ. Manage.* 50, 861–874. doi:10.1007/s00267-012-9929-z

699 Yin, C., Fan, F., Song, A., Li, Z., Yu, W., Liang, Y., 2014. Different denitrification potential of
700 aquic brown soil in Northeast China under inorganic and organic fertilization accompanied
701 by distinct changes of nirS - and nirK -denitrifying bacterial community. *Eur. J. Soil Biol.*
702 65, 47–56. doi:10.1016/j.ejsobi.2014.09.003

703 Zumft, W.G., 1997. Cell biology and molecular basis of denitrification. *Microbiol. Mol. Biol.*
704 *Rev.* 61, 533–616.

705

06 Table 1. Mean soil properties measured at the Elora Research Station.

Cropping System/ Harvest	N Rate (kg ha ⁻¹)	*Bulk Density (g cm ⁻³)	% Organic Carbon		% Total Carbon		Yield (dry tonnes ha ⁻¹)			
			0-15cm	15-30cm	0-15cm	15-30cm	2011	2012	†Mean	
Corn-soybean	Fall	0	1.21 AB	1.88	1.06	2.22	1.86	5.341	2.912	E
Corn-soybean	Fall	160	1.27 A	1.79	1.47	2.25	2.11	9.92	7.882	BC
Miscanthus	Fall	0	1.10 B	2.06	1.44	2.27	1.72	17.62	12.77	A
Miscanthus	Fall	160	1.10 B	2.13	1.63	2.36	1.84	17.43	18.32	A
Miscanthus	Spring	0	1.13 AB	2.09	1.53	2.31	1.69	12.66	13.38	AB
Miscanthus	Spring	160	1.13 AB	2.24	1.42	2.47	1.89	14.33	14.56	A
Switchgrass	Fall	0	1.11 B	2.12	1.43	2.33	1.61	7.648	6.458	CD
Switchgrass	Fall	160	1.09 B	2.12	1.34	2.32	1.73	11.1	10.45	AB
Switchgrass	Spring	0	1.11 B	2.09	1.23	2.32	1.55	6.33	4.146	DE
Switchgrass	Spring	160	1.21 AB	1.92	1.33	2.23	1.7	6.905	6.441	CD

07 *Means of bulk density (n=6) followed by the same letter within one column are not significantly different according to a post-hoc Tukey's means
 08 comparison (p<0.05); carbon measurements (n=3) were not significantly different between treatments. †Mean yields (n=3) followed by the same letter
 09 are not significantly different according to a post-hoc Tukey's means comparison (p< 0.05).

718 Table 2. Mean gene abundance responses to crop and harvest management, averaged over nitrogen application rate and time at the Elora Research
 719 Station.

Cropping System	Management	†Total soil bacteria (gene copy g ⁻¹ soil)		†Soil denitrifying bacteria (gene copy g ⁻¹ soil)	
		16S	<i>nirS</i>	<i>nosZ</i>	<i>nirS:nosZ</i> (x10 ⁻²)
Corn-Soybean	Fall Harvest	1.35 x 10 ⁹ b	1.95 x 10 ⁶ b	2.63 x 10 ⁵ b	7.42
Miscanthus	Fall Harvest	1.38 x 10 ⁹ b	2.30 x 10 ⁶ ab	4.47 x 10 ⁵ a	5.15
Miscanthus	Spring Harvest	1.91 x 10 ⁹ a	3.02 x 10 ⁶ a	5.25 x 10 ⁵ a	5.75
Switchgrass	Fall Harvest	1.41 x 10 ⁹ b	2.19 x 10 ⁶ b	3.55 x 10 ⁵ ab	6.17
Switchgrass	Spring Windrow	1.48 x 10 ⁹ ab	2.46 x 10 ⁶ ab	3.98 x 10 ⁵ ab	6.18

†Means followed by the same letter within one column are not significantly different according to post-hoc Tukey's means comparison at p < 0.05 (n = 24).


720 **Figure Captions**


721 **Figure 1.** Mean daily air temperature ($^{\circ}\text{C}$) and daily precipitation (mm) at the Elora Research Station from
722 January 2011 to November 2012. Soil gravimetric H_2O was measured on a per-sample basis and is shown as
723 crop means ($\pm\text{SE}$) for each sampling date (May 9th, 2011; October 30th, 2011; May 2nd, 2012 and October 20th,
724 2012) (n=12 in perennial grasses, n=6 in corn-soybean rotation).


725 **Figure 2.** Mean soil $\text{NH}_4\text{-N}$ and $\text{NO}_3\text{-N}$ (mg g^{-1} dry soil $\pm\text{SE}$) in annual and perennial biomass cropping
726 systems under varied harvest and N management at the Elora Research Station. CS = corn-soybean, SF = fall-
727 harvested switchgrass, SS = spring-harvested switchgrass, MF = fall-harvested miscanthus and MS = spring-
728 harvested miscanthus. Different letters within panels indicate significant differences according to a post-hoc
729 Tukey's test ($p < 0.05$).

730 **Figure 3.** Mean log gene copies (g^{-1} dry soil $\pm\text{SE}$) in annual and perennial biomass cropping systems under
731 varied harvest management at the Elora Research Station (n=6) over time. Different letters within panels
732 indicate significant differences according to a post-hoc Tukey's test ($p < 0.05$).


733 **Figure 4a.** Principal Component Analysis; factor 1 accounted for 43.89% variance and factor 2 accounted for
734 23.84% variance. **4b.** Loading plot for principal components of response variables (*nirS*, *nosZ* and 16S rRNA
735 gene copies, and soil $\text{NO}_3\text{-N}$, soil $\text{NH}_4\text{-N}$, gravimetric soil H_2O).








742 Figure 4a.



743

744 Figure 4b.



745