



1 **Soil Denitrifier Community Size Changes with Land Use Change to Perennial**
2 **Bioenergy Cropping Systems**

3

4 *Running Head: Soil Denitrifiers associated with Perennial Grasses*

5 Karen A. Thompson¹, Bill Deen² and *Kari E. Dunfield¹

6 ¹School of Environmental Sciences, University of Guelph, 50 Stone Road East, Guelph, Ontario
7 N1G 2W1, Canada; ²Department of Plant Agriculture, University of Guelph, 50 Stone Road
8 East, Guelph, Ontario N1G 2W1, Canada.

9 *Correspondence to Kari E. Dunfield:

10 tel: (519) 824-4120 Ext.58088, dunfield@uoguelph.ca, fax:519-837-0756

11 Original Research Article



12 **Abstract**

13 Dedicated biomass crops are required for future bioenergy production. However, the effects of
14 large-scale land use change (LUC) from traditional annual crops, such as corn-soybean rotations
15 to the perennial grasses (PGs) switchgrass and miscanthus on soil microbial community
16 functioning is largely unknown. Specifically, ecologically significant denitrifying communities,
17 which regulate N₂O production and consumption in soils, may respond differently to LUC due to
18 differences in carbon (C) and nitrogen (N) inputs between crop types and management systems.
19 Our objective was to quantify bacterial denitrifying gene abundances as influenced by corn-
20 soybean crop production compared to PG biomass production. A field trial was established in
21 2008 at the Elora Research Station in Ontario, Canada (n=30), with miscanthus and switchgrass
22 grown alongside corn-soybean rotations at different N rates (0 and 160 kg N ha⁻¹) and biomass
23 harvest dates within PG plots. Soil was collected on four dates from 2011-2012 and quantitative
24 PCR was used to enumerate the total bacterial community (16S rRNA), and communities of
25 bacterial denitrifiers by targeting nitrite reductase (*nirS*) and N₂O reductase (*nosZ*) genes.
26 Miscanthus produced significantly larger yields and supported larger *nosZ* denitrifying
27 communities than corn-soybean rotations regardless of management, indicating large-scale LUC
28 from corn-soybean to miscanthus may be suitable in variable Ontario conditions while
29 potentially mitigating soil N₂O emissions. Harvesting switchgrass in the spring decreased yields
30 in N-fertilized plots, but did not affect gene abundances. Standing miscanthus overwinter
31 resulted in higher 16S rRNA and *nirS* gene copies than in fall-harvested crops. However, the
32 size of the total (16S rRNA) and denitrifying communities changed differently over time and in
33 response to LUC, indicating varying controls on these communities.

34 **Key Words: biomass, bioenergy, miscanthus, switchgrass, corn, soy**

35



36 1. Introduction

37 Future energy needs require dedicated biomass crop production for bioethanol and combustion-
38 based electricity generation. Corn (*Zea mays* L.) –soybean (*Glycine max* L.) rotations currently
39 dominate the landscape across Ontario and the northern US Corn Belt (Gaudin et al., 2015), and
40 corn grain is currently the primary feedstock for bioethanol production in Canada (Jayasundara
41 et al., 2014). The C4 perennial grasses (PGs) switchgrass (*Panicum virgatum* L.) and miscanthus
42 (*Miscanthus* spp.) have been proposed as alternate feedstock crops to corn for biomass-based
43 bioenergy production due to their large biomass yields, reduced nitrogen (N) and water
44 requirements, decreased nutrient leaching and potential for increased soil carbon (C) storage
45 (Blanco-Canqui and Lal, 2009; Foster et al., 2013). Large scale production of C4 PGs in Ontario
46 and the northern Corn Belt would require land use change (LUC) from existing corn-soybean
47 rotations to PG biomass cropping systems (Deen et al., 2011; Kludze et al., 2013; Liang et al.,
48 2012; Sanscartier et al., 2014).

49 Few studies have assessed how this LUC may influence soil microbial community functioning.
50 In particular, soil denitrifying communities represent an ideal subset of the soil microbial
51 community to target to assess changes in ecosystem functioning due to agricultural management
52 and LUC. Denitrifying bacteria represent approximately 5% of the total soil microbial biomass
53 (Braker and Conrad, 2011) and have been identified in over 60 genera (Philippot, 2006),
54 encompassing a wide range of phylogenetic and functional diversity. Multiple studies have
55 linked changes in denitrifier communities with plant types or development stage (Bremer et al.,
56 2007; Hai et al., 2009; Petersen et al., 2012), N fertilization (Hallin et al., 2009; Yin et al., 2014),
57 organic or conventional crop management (Reeve et al., 2010), perennial vs. annual crop land
58 use (Bissett et al., 2011) and C and N inputs (Bastian et al., 2009). These studies suggest that



59 LUC from corn-soybean rotations to PG species may influence the soil bacterial communities
60 which drive soil N₂O production and consumption.

61 Denitrifier community size has been correlated with denitrification process rates (Hallin et al.,
62 2009; Wu et al., 2012), and denitrification potential/potential activity (Attard et al., 2011; Cuhel
63 et al., 2010; Enwall et al., 2010). Potential denitrifying activity and denitrifying community size
64 have also been shown to be correlated in some studies (Hallin et al., 2009; Morales et al., 2010;
65 Szukics et al., 2010; Throback et al., 2007); suggesting community size may indicate potential
66 differences in soil N processes after LUC. Particularly, the *nosZ*-bearing community may act as
67 a N₂O sink and counter high N₂O production rates (Braker and Conrad, 2011; Philippot et al.,
68 2011), therefore influencing N₂O emissions (Cuhel et al., 2010; Morales et al., 2010; Philippot et
69 al., 2011). Relative abundances of denitrifier genes have been used to assess a soil's potential to
70 produce (e.g. *nirS* or *nirK*) and consume (e.g. *nosZ*) N₂O via denitrification, representing a
71 qualitative proxy of relative N₂O emission potential of a soil (Butterbach-Bahl et al., 2013;
72 Hallin et al., 2009; Morales et al., 2010; Petersen et al., 2012; Philippot, 2002).

73 LUC resulting from displacement of corn-soybean rotations by PG production may alter soil
74 microhabitats and therefore soil microbial N-cycling due both to extensive root and rhizome
75 biomass and to large leaf litter inputs to soils in perennial vs. annual systems (Dohleman et al.,
76 2012). Within studies targeting soil microbial communities in biomass cropping systems to date
77 (Hedenec et al., 2014; Liang et al., 2012; Mao et al., 2013a, 2011; Orr et al., 2015), the effects of
78 various management practices (e.g. N fertilization and harvest) on soil microbial community
79 functioning have not been an area of focus. The effect this type of LUC may have on soil
80 microbial communities may depend on PG management practices in these systems.

81



82 Currently, there is no consensus regarding optimal N fertilization practices for increased yields in
83 PG production as yield responses can be highly variable depending on environmental conditions
84 and crop species (Deen et al., 2011). Depending on downstream use, miscanthus can be either
85 harvested in the fall pre-frost, harvested post-frost kill, or left to overwinter as standing biomass
86 for harvest in the spring. Switchgrass is commonly harvested in the fall, and is often windrowed
87 (cut, swathed and left on soil) over winter due to producers' limitations in collecting and storing
88 harvested biomass in winter (Resource Efficient Agricultural Production (REAP)-Canada, 2008;
89 Sokhansanj et al., 2009). Differences in N fertilizer requirements and harvest regimes may alter
90 C and N inputs (Attard et al., 2011) and may influence LUC impacts on soil denitrifier
91 community sizes.

92 Our objective was to compare the effects of LUC from corn-soybean to PG biomass production
93 on the relative abundances of total (16S rRNA gene target) and denitrifier (*nirS* and *nosZ* gene
94 targets) soil bacterial communities 3-4 years after PG planting. Soil was collected on four dates
95 from 2011 to 2012 from a field trial established in Ontario in 2008. The field trial is unique in
96 that it consists of two PG biomass crops produced in parallel with the existing common land use
97 of corn-soybean rotation. It also includes multiple N fertilization rates in both annual and
98 perennial systems, and varied harvest practices within PG plots. We hypothesized that soils from
99 PG plots would support larger total bacterial and denitrifier communities than soils from corn-
100 soybean plots due to increased shoot residue return and root inputs to soils in PG systems, and
101 that soils from PG plots with biomass harvested in the spring would support larger total bacterial
102 and denitrifier communities than supported by soils from PGs harvested in the fall due to
103 increased root inputs and leaf loss to soil over winter.

104



105 2. Materials and Methods

106 2.1 Site Description and Experimental Design

107 A field trial was established in 2008 at the University of Guelph Research Station in Elora, ON
108 (43°38'46.73" N and 80°24'6.66" W). The field site was cultivated on May 16th and June 6th,
109 2008. Switchgrass (*Panicum virgatum* L., Shelter variety) was planted on June 6th, 2008.
110 Miscanthus (*M. sinensis* x *M. sacchariflorus*, Nagara-116 variety) was planted on June 12th,
111 2008, and soybean (*Glycine max* L.) was planted on June 24th, 2008 and annually rotated with
112 corn (*Zea mays* L.). Corn was planted on May 5th, 2010; soy was planted on June 3rd, 2011, and
113 corn was re-planted on May 18th, 2012, with annual light cultivation to prepare seedbeds for
114 planting. In 2007, prior to trial establishment, the experimental area was planted to barley
115 (*Hordeum vulgare* L.) The soil type is a London silt loam (Gray Brown Luvisol).

116 The field trial was a split-split strip plot design with three replicates. The main plot factor was
117 PG crop or annual rotation (miscanthus, switchgrass, and corn-soybean). Main treatment plots
118 measured 6.2 m x 26.0 m. Nitrogen fertilizer (0 or 160 kg N ha⁻¹) was applied in strips
119 randomly within replicates. 160 kg N ha⁻¹ subplots received hand-broadcast urea fertilizer (46-0-
120 0) in May 2011 or hand-broadcast ammonium nitrate fertilizer (34-0-0) in May 2012, after soil
121 sampling procedures described below. Main treatments were split into two harvest timings (fall
122 or spring) within the PG fertilizer strips only. Miscanthus plots were either harvested in the late
123 fall season after post-frost kill, or left standing to overwinter until spring harvest. Switchgrass
124 plots were harvested in the fall, or cut and assembled into windrows in the field for biomass
125 removal in the spring. Spring-harvest of PGs occurred before N fertilizer was applied. Harvest
126 methods of PG yields (dry harvested biomass (tonnes) ha⁻¹) are described in Deen et al. (2011).



127 2.2 Soil sampling and analysis

128 Baseline bulk density and carbon measurements were measured for each main plot on October
129 23rd, 2010. For bulk density, two soil cores per plot were collected at 0-5 cm depth using 2.5
130 cm-diameter cylindrical aluminum cores. Cores were weighed before and after drying for 24h at
131 105°C (Maynard & Curran, 2007). For soil carbon analysis, ten soil cores per plot were
132 collected from both 0-15 cm and 15-30 cm depths using a 5 cm-diameter soil corer on a zed-
133 shaped transect, and then composited per treatment plot for each depth. Total soil carbon and
134 inorganic carbon were analyzed with a Leco[®] Carbon Determinator CR-12 (Model No. 781-700,
135 Leco Instruments Ltd.) following the dry combustion technique (Périé and Ouimet, 2008) on
136 approximately 0.300 g of dried, ground and homogenized soil (Table 1).

137 For molecular analyses, soil was sampled on 4 dates (May 9th, 2011; October 30th, 2011; May
138 2nd, 2012 and October 20th, 2012). October sampling dates occurred before fall harvest of PG
139 crops, while May sampling dates occurred before N fertilizer application and after spring PG
140 biomass removal (Fig. 1). Ten soil cores per plot were sampled aseptically to 15 cm depth using
141 a 5 cm-diameter soil corer on a zed-shaped transect, composited and kept on ice until transport
142 back to the laboratory. The transect shape was chosen to encompass plot heterogeneity; at a pre-
143 trial study date initial analysis indicated gene abundances were not significantly different
144 between bulk or rhizosphere soils in corn-soybean or PG plots, possibly due to the large root
145 biomass/leaf loss to soils in perennial plots and residual soy/corn residue cover on soil in corn-
146 soybean plots. Soil samples were divided for storage at 4°C and -20°C.

147 Mean values of gravimetric soil moisture (g g^{-1}) are shown in Figure 1. Soil exchangeable NO_3^- -
148 N and NH_4^+ -N were determined for each of the soil samples by KCl extraction. Soil samples
149 (10.0 g) were placed into 125 mL flasks and 100 mL of 2.0M KCl was added to each flask.



150 Flasks were stoppered and shaken for 1h at 160 strokes per minute; solutions were allowed to
151 settle and were then filtered through Whatman no. 42 filter paper (Whatman plc, ME, U.S.A).
152 Extractable NO_3^- -N and NH_4^+ -N were determined colourmetrically with segmented flow
153 analyses (AA3, SEAL Analytical, Wisconsin, USA) via a cadmium reduction (Technician
154 Instrument Corporation, 1971), and a Berthelot reaction respectively (Fig. 2).

155 Figure 1 illustrates the seasonal and annual variation in daily average air temperature ($^{\circ}\text{C}$), and
156 daily precipitation (mm) measured at the Elora Research Station.

157 **2.3 Soil DNA Extraction**

158 Total DNA was extracted from field-moist soil sampled from each plot (3 field replicates, $n=3$;
159 total plots = 30). DNA was extracted in duplicate (ca. 0.250g) within 48 h of sampling as per
160 manufacturer's protocol using the DNA PowerSoil Kit (Mbio, Carlsbad, USA). Duplicate
161 extracts were then pooled, separated into aliquots, and stored at -80°C until use in downstream
162 analyses.

163 **2.4 Quantification of total bacteria and functional genes**

164 Quantitative PCR (qPCR) assays were used to enumerate the total bacterial communities (16S
165 rRNA gene), and communities of denitrifiers by targeting nitrite reductase (*nirS*) and nitrous
166 oxide reductase (*nosZ*) genes, using primer pairs 338f/518r (16S, Fierer et al., 2005),
167 Cd3af/R3Cd (*nirS*, Throbäck et al., 2004) and 1F/1R (*nosZ*, Henry et al., 2006). Denitrification
168 *nirS* and *nosZ* gene targets represent the two most important steps in the denitrification pathway
169 that produce gaseous by-products, and account for a large proportion of functional N genes in
170 some studies (Stone et al., 2015). The first step in denitrification that produces a gaseous N
171 product is the reduction of nitrate (NO_2^-) to nitric oxide (NO), catalyzed by nitrite reductases



172 either encoded by the cytochrome cd_1 (*nirS*) or copper-containing (*nirK*) genes, which are
173 equivalent but have not been detected within the same species (Zumft, 1997). We chose to
174 quantify *nirS* because $\frac{3}{4}$ of cultured denitrifiers possess the *nirS* gene (Zumft, 1997) and some
175 molecular reports indicate *nirS* may dominate in abundance over *nirK* in some natural
176 environments (Deslippe et al., 2014; Nogales et al., 2002), indicating it may be a better-suited
177 target for relative characterization of potential nitrite-reducing communities than *nirK*.
178 Additionally, *nirK* has been recently identified in autotrophic ammonia-oxidizing species
179 (Cantera and Stein, 2007; Casciotti and Ward, 2001), calling into question its utility in
180 specifically targeting denitrifying communities. The *nosZ* target codes for nitrous oxide
181 reductase, which catalyzes the reduction of N_2O to N_2 in the denitrification pathway, indicating
182 *nosZ*-bearing communities help to complete the N cycle and determine the $N_2O:N_2$ balance.

183 For each gene target analyzed, duplicate replicates were run in parallel on an IQ5 thermocycler
184 (Bio-Rad Laboratories, Hercules, CA, USA). qPCR reaction mixtures contained 12.5 μ L of 1x
185 SYBR Green Supermix, each forward and reverse primer at a final concentration of 400 nM, 1
186 μ L of DNA template and RNase/DNase-free water to a final volume of 25 μ L. The SYBR Green
187 Supermix contained 100 nM KCl, 40 mM Tris-HCl, 0.4 mM dNTPs, 50 units mL^{-1} iTaq DNA
188 polymerase, 6 mM $MgCl_2$, SYBR Green 20 nM fluorescein, and stabilizer (Bio-Rad
189 Laboratories, Hercules, CA, USA).

190 Conditions for qPCR were an initiation step at 94°C for two minutes, followed by 35 cycles of
191 denaturing at 94°C for thirty seconds, annealing at 57°C for thirty seconds (16S) or at 55°C for
192 sixty seconds (*nirS*), followed by elongation at 72°C for thirty (16S) or sixty (*nirS*) seconds. For
193 *nosZ*, a touchdown protocol adapted from Henry et al., (2006) was used. Amplicon specificity
194 was screened by running qPCR products on an ethidium bromide-stained gel (1% agarose, 80



195 volts for 20 minutes) with a 100bp ladder, which resulted in clean bands for all gene targets. The
196 16S rRNA primers used are degenerate and have been cited as having 89-91% matching
197 efficiency to all bacteria (Bergmark et al., 2012). The primers amplify one of two conserved
198 regions in V3 of the SSU rRNA gene, resulting in a ca. 200 bp amplicon that is within small
199 enough to amplify via qPCR methodology and amplifies for most bacterial taxa (Bakke et al.,
200 2011)

201 Known template standards were made from cloned PCR products from pure culture genomic
202 DNA (*Clostridium thermocellum* (16S), *Pseudomonas aeruginosa* (*nirS*), and *Pseudomonas*
203 *fluorescens* (*nosZ*)) and transformed into *Escherichia coli* plasmids (TOPO TA cloning kit);
204 plasmids were sequenced to confirm successful cloning and transformation of the target genes.
205 Amplicon specificity was screened by running PCR products on an ethidium bromide-stained gel
206 (1% agarose, 80 volts for 20 minutes) with a 100bp ladder. PCR amplicons of cloned gene
207 targets were sequenced by the Laboratory Services Department at the University of Guelph using
208 an ABI Prism 3720 (Applied Biosystems, Foster City, CA, USA) to confirm target identity.

209 In all qPCR assays, all unknown samples were amplified in parallel with a triplicate serial
210 dilution (10^1 - 10^8 gene copies per reaction) of control plasmids. PCR assays were optimized to
211 ensure efficiencies ranging from 93.0-106.4%, with R^2 s ranging from 0.990-0.999 and standard
212 curve slopes of -3.177 to -3.408 by testing serial dilutions of DNA extracts in order to minimize
213 inhibition of amplification due to humic and fulvic contaminants. Duplicate no-template
214 controls were run for each qPCR assay, which gave null or negligible values. Melt curve
215 analysis was used to confirm amplicon specificity. Normalization of DNA concentrations to
216 gram of dry soil was used to give results on a biologically significant scale; this assumes similar



217 DNA isolation efficiency across samples, which is only appropriate when measuring relative (vs.
218 absolute) quantification, as in this study.

219 **2.5 Statistical Analysis**

220 Analysis of variance was conducted in SAS 9.3 (Carlsbad, NC, USA) using a generalized linear
221 mixed model (PROC GLIMMIX). The Shapiro-Wilks test was used to test for normality of data;
222 studentized test for residuals confirmed the absence of outliers. The probability distributions of
223 gene abundance data sets were log normal or highly skewed and were analyzed using an
224 overdispersed Poisson distribution for count data (Ver Hoef and Boveng, 2007). Bulk density,
225 organic carbon, total carbon, nitrate and ammonium data were log transformed when required
226 and fitted to the normal distribution.

227 Within each data set, sampling time was a repeated measure; independent and interactive fixed
228 effects were associated with crop/crop rotation, nitrogen application rate and harvest timing
229 within perennial grasses, while field replicate and its associated interactions were random effects.
230 The residual maximum likelihood method was employed to fit the model for all data sets.
231 Several covariance structures were entertained before the variance components structure was
232 chosen based on convergence and model fitting criteria. Individual treatment means within data
233 sets were compared using a post-hoc Tukey's test for all pairwise comparisons. Significant
234 differences among and between least-square means were determined by p-values; H_0 was
235 rejected at $p < 0.05$.

236 Correlation analysis was used to assess nonparametric measures of statistical dependence
237 between gene abundances and H_2O , NO_3^- -N and NH_4^+ -N measured over time (Supplementary
238 Table 1). Correlation analysis resulted in multiple significant correlations between variables; as



239 such a principal component analysis was conducted in SAS (PROC FACTOR) on 120 samples
240 using a VARIMAX rotation.



241 **3. Results**

242 **3.1 Environmental and Soil Conditions**

243 Environmental conditions varied during the periods prior to the four soil sampling dates (Figure
244 1). Average air temperatures over the growing seasons (May-October) were 16.9°C and 17.3°C
245 in 2011 and 2012 respectively (Roy et al. 2014); air temperatures in spring 2012 were warmer
246 than normal and resulted in earlier emergence of PG crops compared to 2011. Precipitation was
247 above average prior to the May 2011 sampling date (101 mm vs. 72 mm 30-year average in April
248 2011 and 113 mm vs. 82 mm 30-year average in May 2011) (Roy et al., 2014). In comparison,
249 S. Ontario received very low precipitation in April 2012 (30 mm vs. 72 mm 30-year average) and
250 May 2012 (28 mm vs. 82 mm 30-year average) (Roy et al., 2014). Precipitation levels were
251 lower in 2012 compared to 2011 from May-August (391 mm in 2011 vs. 186 mm in 2012),
252 however higher than normal precipitation levels occurred in October of 2011 (129 mm vs. 77
253 mm 30-year average) and both September (106 mm vs. 77 mm 30-year average) and October
254 (127 mm vs. 77 mm 30-year average) of 2012 (Roy et al., 2014). Environmental conditions prior
255 to soil sampling directly impact soil gravimetric content measured at time of sampling (Fig. 1),
256 and could also impact soil N and soil bacterial communities.

257 Soil physical and chemical properties were assessed in October 2010, after only two years of
258 contrasting management since crop establishment in 2008. The corn-soybean rotation had higher
259 soil bulk density than soils from both miscanthus and switchgrass plots harvested in the fall. No
260 differences in total or organic soil carbon were detected between the corn-soybean rotation and
261 the PG treatments at either the 0-15cm or 15-30cm depth (Table 1). Soil NH₄-N levels did not
262 differ significantly between the corn-soybean rotation and the PG soils, however N fertilization



263 significantly increased $\text{NH}_4\text{-N}$ levels in soils from fall-harvested miscanthus plots ($p < 0.05$) (Fig.
264 2a). N fertilization also significantly increased $\text{NO}_3\text{-N}$ levels in spring-harvested switchgrass
265 ($p < 0.05$) (Fig. 2b). From May to October 2011, soil $\text{NH}_4\text{-N}$ levels increased significantly and
266 soil $\text{NO}_3\text{-N}$ levels decreased significantly in PG soils (data not shown); a similar trend was not
267 observed in 2012 or for soils from the corn-soybean rotation in either year.

268 **3.2 Biomass Yields**

269 Despite significant differences in precipitation between 2011 and 2012, biomass yields of
270 miscanthus and switchgrass did not differ between years. In comparison, corn grain yields were
271 higher in 2011 vs 2012 (Table 1). Miscanthus produced higher yields (12.7-18.3 dry tonnes ha^{-1})
272 than either switchgrass or corn grain, regardless of N fertilization rate or harvest timing (Table
273 1). When miscanthus was left standing overwinter and harvested in the spring a slight biomass
274 reduction was observed in 2011 (12.7-14.6 dry tonnes ha^{-1}); however this was not statistically
275 significant at $p < 0.05$. When harvested in the fall and N-fertilized, switchgrass yields were not
276 significantly lower (10.5-11.1 dry tonnes ha^{-1}) than miscanthus yields. Switchgrass yields from
277 unfertilized plots were not significantly different if harvested in the fall or spring; however,
278 switchgrass yields from fertilized plots were significantly higher (ca. 3-4 dry tonnes ha^{-1}) when
279 harvested in the fall compared to yields obtained when switchgrass was windrowed over winter.

280 **3.3 Bacterial Responses to Annual and Perennial Crops and their Management**

281 There was no statistically significant effect of N fertilization or any significant interactions
282 between cropping system and sampling time on any of the targeted gene abundances. Therefore
283 we analyzed the impact of each biomass crop under specific harvest management on soil
284 bacterial gene abundances (Table 2). Denitrifying (*nosZ*) gene copy abundances were affected



285 by LUC; regardless of harvest or N management, mean *nosZ* gene copies were higher in
286 miscanthus plots than in the corn-soybean rotation, and *nirS:nosZ* ratios were higher in the corn-
287 soybean soils than in miscanthus or switchgrass soils ($p < 0.05$) (Table 2). Under fall harvesting
288 management, biomass crop had no impact on total (16S rRNA) gene copies or *nirS* gene copies.
289 However, leaving miscanthus biomass standing overwinter until spring resulted in significantly
290 higher 16S rRNA gene copies than observed in soils from fall-harvested biomass crops and
291 significantly higher *nirS* gene copies than in fall-harvested switchgrass or the corn-soybean
292 rotation (Table 2).

293 3.3 Temporal Changes in Bacterial Gene Abundances

294 Sampling date had a significant impact on gene abundances for all genes quantified (Fig. 3).
295 Over both sampling years, 16S rRNA gene copies were significantly higher ($5.2\text{--}5.4 \times 10^9$ gene
296 copies g^{-1} dry soil) at fall (October) sampling dates compared to the ca. $5.5\text{--}6.4 \times 10^8$ gene copies
297 g^{-1} dry soil quantified at spring (May) sampling dates (Fig. 3). Populations of *nirS* and *nosZ*
298 denitrifiers represented ca. 1.58% and 0.26% on a gene-to-gene basis (*nirS* or *nosZ* to 16S) of the
299 total bacterial community (data not shown), and did not follow similar trends with time of
300 sampling (Fig. 3). The abundance of *nirS* gene copies was significantly higher in 2012 (4.0×10^6
301 $\text{--} 1.6 \times 10^7$ gene copies g^{-1} dry soil) compared to 2011 ($2.5\text{--}6.3 \times 10^5$ gene copies g^{-1} dry soil),
302 with no significant differences between May and October sampling dates within each year (Fig.
303 3). The abundance of *nosZ* gene copies were approximately $1.3\text{--}3.2 \times 10^5$ gene copies g^{-1} dry
304 soil, but increased significantly in May 2012 to approximately 3.2×10^6 gene copies g^{-1} dry soil
305 and dropped back to previous levels by October 2012 (Fig. 3). Higher relative proportions of
306 denitrifiers (*nirS* or *nosZ* to 16S) were observed at spring sampling dates, when total (16S) gene
307 abundances decreased in comparison to fall sampling dates (Fig. 3).



308 Two factors were selected in the principal components analysis, which accounted for 67.73%
309 cumulative variance. A scree plot was examined for breaks and factors were retained when
310 eigenvalues ≥ 1 . Soil $\text{NH}_4\text{-N}^+$, soil $\text{NO}_3^-\text{-N}$, *nirS* and *nosZ* loaded on factor 1, which accounted
311 for 43.89% variance while soil gravimetric H_2O and 16S rRNA loaded on factor 2, which
312 accounted for 23.84% variance (Fig.4 a and b). Differences in soil $\text{NO}_3^-\text{-N}$ and $\text{NH}_4^+\text{-N}$ were
313 strongly related to differences in *nirS* and *nosZ* gene abundances observed between May 2011
314 and May 2012 sampling dates (Fig. 3 and Fig. 4), while the size of the total bacterial community
315 (16S rRNA) was related to soil gravimetric moisture levels (Fig. 4).

316 4. Discussion

317 Denitrification is an important process contributing to the production and consumption of N_2O in
318 soils, and mitigation of GHGs such as N_2O is required to create sustainable biomass cropping
319 systems (Miller et al., 2008; Schlesinger, 2013). Changes in the potential functional abilities of
320 the soil microbial community may reflect changes in LUC or agricultural management and
321 should be considered to assess the ecological impact of biomass crop production (Hedenec et al.,
322 2014). Currently, few studies have assessed soil microbial community responses to PG biomass
323 production systems (Hedenec et al., 2014; Liang et al., 2012; Mao et al., 2013a, 2011; Orr et al.,
324 2015). The highest potential to reduce GHG emissions from biomass cropping systems is to
325 produce crops with high yields, such as PGs (Sanscartier et al., 2014), which offset the amount of
326 land required for crop production (Kludze et al., 2013). However, if PG biomass production
327 negatively affects soil health as indicated by changes in the potential functioning of microbial
328 communities, large-scale LUC from annual to perennial biomass production may not be as
329 sustainable as originally proposed. As such it is necessary to identify biomass cropping systems



330 that not only result in large biomass yields, but also ensure agroecosystem sustainability by
331 maintaining or improving ecosystem services (Orr et al., 2015), such as soil N-cycling.

332 **4.1 Biomass Yields of Annual and Perennial Crops**

333 Miscanthus and switchgrass biomass yields were within the typical range of values reported
334 previously in Ontario (Kludze et al., 2013; Resource Efficient Agricultural Production (REAP)-
335 Canada, 2008) and Europe (Christian et al., 2008; Himken et al., 1997), despite differences in
336 temperature and precipitation between the two study years. Corn grain yields were within the
337 lower range for reported Ontario yields (Munkholm et al., 2013), potentially due to wetter (2011)
338 and drier (2012) field conditions than normal over the two growing seasons (Roy et al., 2014).
339 Deen et al. (2011) showed increases in PG biomass yields between years 2 and 3 at our site,
340 whereas we measured similar yields in 2011 and 2012, indicating the PGs may have reached
341 maximum yield potential.

342 Nitrogen fertilization significantly increased corn grain yields and fall-harvested switchgrass
343 biomass yields, however no significant increases due to N fertilization were observed in
344 miscanthus or spring-harvested switchgrass biomass yields. Potential yield increases from N
345 fertilization in spring-harvested switchgrass may have been offset due to leaf loss over the winter
346 season, as increases in switchgrass yields to N fertilization have been previously observed
347 (Nikièma et al., 2011; Vogel et al., 2002). Similar to the present study, European and US field
348 trials have also found no response of miscanthus yields to N (Lewandowski et al., 2000;
349 Lewandowski et al., 2003; Behnke et al., 2012; Christian et al., 2008), and PG yields were
350 minimally impacted by differences in growing season conditions compared to corn grain yields
351 (Table 1).



352 Despite significant differences in biomass yields between miscanthus and corn-soybean systems,
353 there were no significant differences in either total or organic soil carbon between any of
354 cropping systems assessed (Table 1). Sampling of soil carbon occurred only two years after PG
355 planting; PGs are expected to be productive for 20+ years, indicating future changes in soil
356 carbon levels may occur. Additionally, Ontario-based land conversion modelling scenarios have
357 estimated a soil carbon decrease of 2.5% upon miscanthus establishment (Sanscartier et al.,
358 2014), which may have negated potential increases in soil organic carbon. However, high
359 miscanthus yields most likely resulted in increases in above and below-ground plant residue
360 return to soils (Mutegi et al., 2010; Soil Quality National, 2006); therefore our carbon measures
361 may not have reflected short-term changes in labile carbon sources that had occurred.
362 Regardless of management or climatic conditions, miscanthus consistently produced large yields,
363 emphasizing its potential as a bioenergy crop suitable for production in variable Ontario
364 conditions.

365 **4.2 Bacterial Responses to Annual and Perennial Crops and their Management**

366 Some studies in biomass cropping systems have not observed differences in soil microbial
367 responses between perennial and annual crop types (Mao et al., 2011), while others have
368 measured significant differences in microbial abundance, diversity and community structure
369 between these cropping types (Liang et al., 2012; Morales et al., 2010; Watrud et al., 2013).
370 Currently, we observed significantly higher *nosZ* gene copies in miscanthus soils compared to
371 corn-soybean soils, illustrating a distinct effect of LUC from corn-soybean to miscanthus
372 production on soil N cycling (Table 2).



373 Due to the large biomass produced by miscanthus compared to corn, a large amount of plant
374 residues are returned to the soil; these residues provide surface cover, decrease soil bulk density,
375 increase water retention and regulate temperatures (Blanco-Canqui and Lal, 2009). Previous
376 work at the Elora Research Station found an inverse correlation between field-scale N₂O fluxes
377 and *nosZ* transcript abundance in conventionally-tilled corn plots with residues returned to soils
378 (Németh et al., 2014), and increased *nosZ* activity after residue amendment has also been
379 observed in lab studies (Henderson et al., 2010). High C:N plant residues have been negatively
380 correlated with cumulative N₂O emissions (Huang, 2004), and may encourage complete
381 reduction of N₂O to N₂ as soil available NO₃-N is limiting, so bacterial populations with the
382 ability to reduce N₂O to N₂ are favoured (Miller et al., 2008). Presently, the primers used for
383 *nosZ* gene target amplification provided good coverage of γ -Proteobacteria (Henry et al., 2006),
384 which are stimulated by surface-applied residues (Pascault et al, 2010). Increased residue return
385 in miscanthus plots may have selected for bacterial populations harbouring enhanced catabolic
386 capabilities, such as N₂O reduction (Pascault et al., 2010). This implies that producing biomass
387 crops with large yields may indirectly alter soil N cycling and potentially mitigate soil N₂O
388 emissions due to increased residue return influencing the soil microbial community. It is likely
389 that differences in environmental conditions (e.g. temperature, H₂O and O₂ availability) and
390 resource quality and availability between corn-soybean and miscanthus soils related to
391 differences in microbial community structure (Cusack et al., 2011) and selected for different
392 dominant taxa that filled different ecological niches (Stone et al., 2015).

393 N fertilization did not affect targeted gene abundances, however studies in other cropping
394 systems have found that N fertilization affected the size of denitrifying communities (Hallin et
395 al., 2009), nitrifying communities (He et al., 2007), and proportions of *nirS/nirK* communities



396 (Hai et al., 2009). Elevated 16S and *nirS* gene copies were observed in soils from spring-
397 harvested miscanthus and windrowed switchgrass (Table 2). Increased N return via senescent
398 leaf loss in PG plots over winter contributes to the soil organic matter pool (Heaton et al., 2009),
399 and may have contributed to elevated total (16S) bacterial populations in these soils,
400 concomitantly increasing *nirS* abundances (Huang et al., 2011).

401 **4.3 Temporal Changes in Bacterial Gene Abundances**

402 Total soil bacterial communities (16S rRNA) followed a seasonal trend, with elevated 16S rRNA
403 gene copies at fall (October) compared to spring (May) sampling dates, possibly due to an
404 increase in the availability and diversity of resources for microbial metabolism and growth over
405 the growing season (Habekost et al., 2008). Denitrifying abundances changed differently than
406 the total bacterial community, suggesting denitrifiers were influenced by different proximal
407 regulators than the total bacterial community (Fig. 3 and 4). Seasonal dynamics of N-cycling
408 microbial communities have been previously characterized (Boyer et al., 2006; Nemeth et al.,
409 2014; Wolsing and Priemé, 2004; Dandie et al., 2008; Bremer et al., 2007), and are tightly
410 coupled with seasonal changes in labile C and N pools, temperature and soil H₂O (Butterbach-
411 Bahl et al. 2013; Rasche et al., 2011), indicating that local edaphic drivers may often take
412 precedence over crop-specific drivers (Mao et al., 2013).

413 **5.0 Conclusions**

414 Miscanthus consistently produced large yields and supported larger *nosZ*-bearing communities
415 than the corn-soybean rotation, emphasizing its influence on soil N cycling and its potential to
416 mitigate soil N₂O emissions while being suitable for production in variable Ontario conditions.
417 Additionally, miscanthus yields were not increased with N fertilization, indicating a lower N



418 input requirement for biomass production compared to switchgrass. Higher 16S rRNA and *nirS*
419 gene copies were associated with reduced yields in spring-harvested PGs, indicating that
420 harvesting PGs in the spring may increase populations of denitrifiers capable of producing N₂O
421 emissions while simultaneously decreasing biomass yields. The size of both denitrifying (*nirS*
422 and *nosZ*) and total bacterial (16S rRNA) communities changed over the sampling period,
423 however changes in denitrifying gene abundances did not parallel changes in the total soil
424 bacterial community, indicating denitrifying communities were regulated differently than the
425 total bacterial community. Future work measuring N₂O emissions and denitrifier activity
426 (mRNA) and community structure in these systems is required to link the effects of LUC on
427 these communities directly with N₂O fluxes.

428 **Author Contributions:**

429 K. Thompson was the primary researcher and author on this study, conducting all field work, lab
430 work, and manuscript preparation. Author B. Deen is credited for the use of his OMAFRA-
431 funded field plots for this research, and valuable advice on experimental design, statistical
432 analyses and manuscript focus. Author K. Dunfield is credited for her invaluable mentorship on
433 molecular analyses and trouble-shooting on field sampling and sample preservation techniques,
434 manuscript focus and preparation, and data interpretation.

435

436 **Acknowledgements**

437 Funding for this research was awarded through grants from the Natural Science and Engineering
438 Research Council of Canada (NSERC), the Canada Research Chairs (CRC) Program and the
439 Ontario Ministry of Agriculture and Food and the Ministry of Rural Affairs (OMAF/MRA).
440 K.T. was also supported through an Ontario Graduate Scholarship (OGS) and an Ontario



441 Graduate Fellowship (OGF). Special thanks to Henk Wichers (University of Guelph) for his
442 expertise in managing the Alternate Renewable Fuels trial at the Elora Research Station.

443 **References**

- 444 Attard, E., Recous, S., Chabbi, A., De Berranger, C., Guillaumaud, N., Labreuche, J., Philippot,
445 L., Schmid, B., Le Roux, X., 2011. Soil environmental conditions rather than denitrifier
446 abundance and diversity drive potential denitrification after changes in land uses. *Glob.*
447 *Chang. Biol.* 17, 1975–1989. doi:10.1111/j.1365-2486.2010.02340.x
- 448 Bastian, F., Bouziri, L., Nicolardot, B., Ranjard, L., 2009. Impact of wheat straw decomposition
449 on successional patterns of soil microbial community structure. *Soil Biol. Biochem.* 41,
450 262–275. doi:10.1016/j.soilbio.2008.10.024
- 451 Bakke, I., De Schryver, P., Boon, N., Vadstein, O., 2011. PCR-based community structure
452 studies of Bacteria associated with eukaryotic organisms: A simple PCR strategy to avoid
453 co-amplification of eukaryotic DNA. *J. Microbiol. Methods* 84, 349–351.
454 doi:10.1016/j.mimet.2010.12.015
- 455 Bergmark, L., Poulsen, P.H.B., Al-Soud, W.A., Norman, A., Hansen, L.H., Sørensen, S.J., 2012.
456 Assessment of the specificity of Burkholderia and Pseudomonas qPCR assays for detection
457 of these genera in soil using 454 pyrosequencing. *FEMS Microbiol. Lett.* 333, 77–84.
458 doi:10.1111/j.1574-6968.2012.02601.x
- 459 Bissett, A., Richardson, A.E., Baker, G., Thrall, P.H., 2011. Long-term land use effects on soil
460 microbial community structure and function. *Appl. Soil Ecol.* 51, 66–78.
461 doi:10.1016/j.apsoil.2011.08.010
- 462 Blanco-Canqui, H., Lal, R., 2009. Crop Residue Removal Impacts on Soil Productivity and
463 Environmental Quality. *CRC. Crit. Rev. Plant Sci.* 28, 139–163.
464 doi:10.1080/07352680902776507
- 465 Boyer, E.W., Alexander, R.B., Parton, W.J., Li, C., Butterbach-Bahl, K., Donner, S.D., Skaggs,
466 R.W., Del Grosso, S.J., 2006. Modeling denitrification in terrestrial and aquatic ecosystems
467 at regional scales. *Ecol. Appl.* 16, 2123–42.
- 468 Braker, G., Conrad, R., 2011. Diversity, structure, and size of N₂O-producing microbial
469 communities in soils-what matters for their functioning?, *Advances in applied*
470 *microbiology.* Elsevier. doi:10.1016/B978-0-12-387046-9.00002-5
- 471 Bremer, C., Braker, G., Matthies, D., Reuter, A., Engels, C., Conrad, R., 2007. Impact of plant
472 functional group, plant species, and sampling time on the composition of nirK-type
473 denitrifier communities in soil. *Appl. Environ. Microbiol.* 73, 6876–84.
474 doi:10.1128/AEM.01536-07
- 475 Butterbach-Bahl, K., Baggs, E.M., Dannenmann, M., Kiese, R., Zechmeister-Boltenstern, S.,
476 2013. Nitrous oxide emissions from soils: how well do we understand the processes and
477 their controls? *Philos. Trans. R. Soc. B Biol. Sci* 368, 1–13. doi:10.1098/rstb.2013.0122
- 478 Canera, J.J.L., Stein, L.Y., 2007. Role of nitrite reductase in the ammonia-oxidizing pathway of
479 *Nitrosomonas europaea.* *Arch. Microbiol.* 188, 349–54. doi:10.1007/s00203-007-0255-4
- 480 Casciotti, K.L., Ward, B.B., 2001. Dissimilatory Nitrite Reductase Genes from Autotrophic
481 Ammonia-Oxidizing Bacteria. *Appl. Environ. Microbiol.* 67, 2213–2221.
482 doi:10.1128/AEM.67.5.2213
- 483 Christian, D.G., Riche, a. B., Yates, N.E., 2008. Growth, yield and mineral content of



- 484 *Miscanthus* × *giganteus* grown as a biofuel for 14 successive harvests. *Ind. Crops Prod.* 28,
485 320–327. doi:10.1016/j.indcrop.2008.02.009
- 486 Cuhel, J., Simek, M., Laughlin, R.J., Bru, D., Cheneby, D., Watson, C.J., Philippot, L., Chèneby,
487 D., 2010. Insights into the effect of soil pH on N₂O and N₂ emissions and denitrifier
488 community size and activity. *Appl. Environ. Microbiol.* 76, 1870–8.
489 doi:10.1128/AEM.02484-09
- 490 Cusack, D.F., Silver, W.L., Torn, M.S., Burton, S.D., Firestone, M.K., 2011. Changes in
491 microbial community characteristics and soil organic matter with nitrogen additions in two
492 tropical forests. *Ecology* 92, 621–632. doi:10.1890/10-0459.1
- 493 Dandie, C.E., Burton, D.L., Zebarth, B.J., Henderson, S.L., Trevors, J.T., Goyer, C., 2008.
494 Changes in bacterial denitrifier community abundance over time in an agricultural field and
495 their relationship with denitrification activity. *Appl. Environ. Microbiol.* 74, 5997–6005.
496 doi:10.1128/AEM.00441-08
- 497 Deen, B.B., Young, D., Rowsell, J., Tubeileh, A., Engbers, H., Rosser, B., 2011. A comparative
498 study assessing variety and management effects on C 4 perennial grasses in a northern
499 climate. *Asp. Appl. Biol.* 112, 205–212.
- 500 Deslippe, J.R., Jamali, H., Jha, N., Saggarr, S., 2014. Soil Biology & Biochemistry Denitrifier
501 community size, structure and activity along a gradient of pasture to riparian soils. *Soil
502 Biol. Biochem.* 71, 48–60. doi:10.1016/j.soilbio.2014.01.007
- 503 Dohleman, F.G., Heaton, E. a., Arundale, R. a., Long, S.P., 2012. Seasonal dynamics of above-
504 and below-ground biomass and nitrogen partitioning in *Miscanthus* × *giganteus* and
505 *Panicum virgatum* across three growing seasons. *GCB Bioenergy* 4, 534–544.
506 doi:10.1111/j.1757-1707.2011.01153.x
- 507 Enwall, K., Throbäck, I.N., Stenberg, M., Söderström, M., Hallin, S., 2010. Soil resources
508 influence spatial patterns of denitrifying communities at scales compatible with land
509 management. *Appl. Environ. Microbiol.* 76, 2243–50. doi:10.1128/AEM.02197-09
- 510 Fierer, N., Jackson, J.A., Vilgalys, R., Jackson, R.B., 2005. Assessment of Soil Microbial
511 Community Structure by Use of Taxon-Specific Quantitative PCR Assays. *Appl. Environ.
512 Microbiol.* 71, 4117–4120. doi:10.1128/AEM.71.7.4117
- 513 Foster, J.L., Guretzky, J.A., Huo, C., Kering, M.K., Butler, T.J., 2013. Effects of Row Spacing,
514 Seeding Rate, and Planting Date on Establishment of Switchgrass. *Crop Sci.* 53, 309–314.
515 doi:10.2135/cropsci2012.03.0171
- 516 Gaudin, A.C.M., Janovicek, K., Deen, B., Hooker, D.C., 2015. Wheat improves nitrogen use
517 efficiency of maize and soybean-based cropping systems. *Agric. Ecosyst. Environ.* 210, 1–
518 10. doi:10.1016/j.agee.2015.04.034
- 519 Habekost, M., Eisenhauer, N., Scheu, S., Steinbeiss, S., Weigelt, A., Gleixner, G., 2008.
520 Seasonal changes in the soil microbial community in a grassland plant diversity gradient
521 four years after establishment. *Soil Biol. Biochem.* 40, 2588–2595.
522 doi:10.1016/j.soilbio.2008.06.019
- 523 Hai, B., Diallo, N.H., Sall, S., Haesler, F., Schauss, K., Bonzi, M., Assigbetse, K., Chotte, J.-L.,
524 Munch, J.C., Schloter, M., 2009. Quantification of key genes steering the microbial nitrogen
525 cycle in the rhizosphere of sorghum cultivars in tropical agroecosystems. *Appl. Environ.*



- 526 Microbiol. 75, 4993–5000. doi:10.1128/AEM.02917-08
- 527 Hallin, S., Jones, C.M., Schloter, M., Philippot, L., 2009. Relationship between N-cycling
528 communities and ecosystem functioning in a 50-year-old fertilization experiment. ISME J.
529 3, 597–605. doi:10.1038/ismej.2008.128
- 530 He, J.-Z., Shen, J.-P., Zhang, L.-M., Zhu, Y.-G., Zheng, Y.-M., Xu, M.-G., Di, H., 2007.
531 Quantitative analyses of the abundance and composition of ammonia-oxidizing bacteria and
532 ammonia-oxidizing archaea of a Chinese upland red soil under long-term fertilization
533 practices. Environ. Microbiol. 9, 2364–74. doi:10.1111/j.1462-2920.2007.01358.x
- 534 Hedeneč, P., Novotny, D., Ustak, S., Cajthaml, T., Slejska, A., Simackova, H., Honzik, R.,
535 Kovarova, M., Frouz, J., 2014. The effect of native and introduced biofuel crops on the
536 composition of soil biota communities. Biomass and Bioenergy 60, 137–146.
537 doi:10.1016/j.biombioe.2013.11.021
- 538 Henderson, S.L., Dandie, C.E., Patten, C.L., Zebarth, B.J., Burton, D.L., Trevors, J.T., Goyer, C.,
539 2010. Changes in denitrifier abundance, denitrification gene mRNA levels, nitrous oxide
540 emissions, and denitrification in anoxic soil microcosms amended with glucose and plant
541 residues. Appl. Environ. Microbiol. 76, 2155–64. doi:10.1128/AEM.02993-09
- 542 Henry, S., Bru, D., Stres, B., Hallet, S., Philippot, L., 2006. Quantitative detection of the *nosZ*
543 gene, encoding nitrous oxide reductase, and comparison of the abundances of 16S rRNA,
544 *narG*, *nirK*, and *nosZ* genes in soils. Appl. Environ. Microbiol. 72, 5181–9.
545 doi:10.1128/AEM.00231-06
- 546 Himken, M., Lammel, J., Neukirchen, D., Olf, H., 1997. Cultivation of *Miscanthus* under West
547 European conditions : Seasonal changes in dry matter production , nutrient uptake and
548 remobilization. Plant Soil 189, 117–126.
- 549 Huang, S., Chen, C., Yang, X., Wu, Q., Zhang, R., 2011. Distribution of typical denitrifying
550 functional genes and diversity of the *nirS*-encoding bacterial community related to
551 environmental characteristics of river sediments. Biogeosciences 8, 3041–3051.
552 doi:10.5194/bg-8-3041-2011
- 553 Huang, Y., 2004. Nitrous oxide emissions as influenced by amendment of plant residues with
554 different C:N ratios. Soil Biol. Biochem. 36, 973–981. doi:10.1016/j.soilbio.2004.02.009
- 555 Jayasundara, S., Wagner-Riddle, C., Dias, G., Kariyapperuma, K. a., 2014. Energy and
556 greenhouse gas intensity of corn (*Zea mays* L.) production in Ontario: A regional
557 assessment. Can. J. Soil Sci. 94, 77–95. doi:10.4141/cjss2013-044
- 558 Kludze, H., Deen, B., Weersink, A., van Acker, R., Janovicek, K., De Laporte, A., 2013. Impact
559 of land classification on potential warm season grass biomass production in Ontario,
560 Canada. Can. J. Plant Sci. 93, 249–260. doi:10.4141/cjps2012-143
- 561 Lewandowski, I., Andersson, B., Basch, G., Christian, D.G., Jørgensen, U., Jones, M.B., Riche,
562 A.B., Schwarz, K.U., Tayebi, K., Teixeira, F., 2003. Environment and Harvest Time Affects
563 the Combustion Qualities of of *Miscanthus* Genotypes. Agron. J. 95, 1274–1280.
- 564 Lewandowski, I., Clifton-brown, J.C., Scurlock, J.M.O., Huisman, W., 2000. *Miscanthus* :
565 European experience with a novel energy crop. Biomass and Bioenergy 19, 209–227.
- 566 Liang, C., Jesus, E.D.C., Duncan, D.S., Jackson, R.D., Tiedje, J.M., Balsler, T.C., 2012. Soil
567 microbial communities under model biofuel cropping systems in southern Wisconsin, USA:



- 568 Impact of crop species and soil properties. *Appl. Soil Ecol.* 54, 24–31.
569 doi:10.1016/j.apsoil.2011.11.015
- 570 Mao, Y., Yannarell, A.C., Davis, S.C., Mackie, R.I., 2013a. Impact of different bioenergy crops
571 on N-cycling bacterial and archaeal communities in soil. *Environ. Microbiol.* 15, 928–42.
572 doi:10.1111/j.1462-2920.2012.02844.x
- 573 Mao, Y., Yannarell, A.C., Davis, S.C., Mackie, R.I., 2013b. Impact of different bioenergy crops
574 on N-cycling bacterial and archaeal communities in soil: Impact of bioenergy crops on soil
575 N-cycling archaea and bacteria. *Environ. Microbiol.* 15, 928–942. doi:10.1111/j.1462-
576 2920.2012.02844.x
- 577 Mao, Y., Yannarell, A.C., Mackie, R.I., 2011. Changes in N-transforming archaea and bacteria
578 in soil during the establishment of bioenergy crops. *PLoS One* 6, 1–12.
579 doi:10.1371/journal.pone.0024750
- 580 Maynard, D.G. and Curran, M.P. 2007. Bulk Density Measurement in Forest Soil. Chapter 66.
581 In: *Soil Sampling and methods of analysis.* (Carter M.R. & Gregorich E.G.) Canadian
582 Society of Soil Science. CRC Press, Boca Raton, FL. 863-869.
- 583 Miller, M.N., Zebarth, B.J., Dandie, C.E., Burton, D.L., Goyer, C., Trevors, J.T., 2008. Crop
584 residue influence on denitrification, N₂O emissions and denitrifier community abundance in
585 soil. *Soil Biol. Biochem.* 40, 2553–2562. doi:10.1016/j.soilbio.2008.06.024
- 586 Morales, S.E., Cosart, T., Holben, W.E., 2010. Bacterial gene abundances as indicators of
587 greenhouse gas emission in soils. *ISME J.* 4, 799–808. doi:10.1038/ismej.2010.8
- 588 Munkholm, L.J., Heck, R.J., Deen, B., 2013. Long-term rotation and tillage effects on soil
589 structure and crop yield. *Soil Tillage Res.* 127, 85–91. doi:10.1016/j.still.2012.02.007
- 590 Mutegi, J.K., Munkholm, L.J., Petersen, B.M., Hansen, E.M., Petersen, S.O., 2010. Nitrous
591 oxide emissions and controls as influenced by tillage and crop residue management
592 strategy. *Soil Biol. Biochem.* 42, 1701–1711. doi:10.1016/j.soilbio.2010.06.004
- 593 Németh, D.D., Wagner-Riddle, C., Dunfield, K.E., 2014. Abundance and gene expression in
594 nitrifier and denitrifier communities associated with a field scale spring thaw N₂O flux
595 event. *Soil Biol. Biochem.* 73, 1–9. doi:10.1016/j.soilbio.2014.02.007
- 596 Nogales, B., Timmis, K.N., Nedwell, D.B., Osborn, A.M., 2002. Detection and Diversity of
597 Expressed Denitrification Genes in Estuarine Sediments after Reverse Transcription-PCR
598 Amplification from mRNA. *Appl. Environ. Microbiology* 68, 5017–5025.
599 doi:10.1128/AEM.68.10.5017
- 600 Orr, M.-J., Gray, M.B., Applegate, B., Volenec, J.J., Brouder, S.M., Turco, R.F., 2015.
601 Transition to second generation cellulosic biofuel production systems reveals limited
602 negative impacts on the soil microbial community structure. *Appl. Soil Ecol.* 95, 62–72.
603 doi:10.1016/j.apsoil.2015.06.002
- 604 Pascault, N., Nicolardot, B., Bastian, F., Thiébeau, P., Ranjard, L., Maron, P.-A.A., 2010. In situ
605 dynamics and spatial heterogeneity of soil bacterial communities under different crop
606 residue management. *Microb. Ecol.* 60, 291–303. doi:10.1007/s00248-010-9648-z
- 607 Périé, C., Ouimet, R., 2008. Organic carbon, organic matter and bulk density relationships in
608 boreal forest soils. *Can. J. Soil Sci.* 88, 315–325. doi:10.4141/CJSS06008



- 609 Petersen, D.G., Blazewicz, S.J., Firestone, M., Herman, D.J., Turetsky, M., Waldrop, M., 2012.
 610 Abundance of microbial genes associated with nitrogen cycling as indices of
 611 biogeochemical process rates across a vegetation gradient in Alaska. *Environ. Microbiol.*
 612 14, 993–1008. doi:10.1111/j.1462-2920.2011.02679.x
- 613 Philippot, L., 2002. Denitrifying genes in bacterial and Archaeal genomes. *Biochim. Biophys.*
 614 *Acta* 1577, 355–76.
- 615 Philippot, L., 2006. Use of functional genes to quantify denitrifiers in the environment. *Biochem.*
 616 *Soc. Trans.* 34, 101–3. doi:10.1042/BST0340101
- 617 Philippot, L., Andert, J., Jones, C., Bru, D., Hallin, S., 2011. Importance of denitrifiers lacking
 618 the genes encoding the nitrous oxide reductase for N₂O emissions from soil. *Glob. Chang.*
 619 *Biol.* 17, 1497–1504. doi:10.1111/j.1365-2486.2010.02334.x
- 620 Rasche, F., Knapp, D., Kaiser, C., Koranda, M., Kitzler, B., Zechmeister-Boltenstern, S.,
 621 Richter, A., Sessitsch, A., 2011. Seasonality and resource availability control bacterial and
 622 archaeal communities in soils of a temperate beech forest. *ISME J.* 5, 389–402.
 623 doi:10.1038/ismej.2010.138
- 624 Reeve, J.R., Schadt, C.W., Carpenter-boggs, L., Kang, S., Zhou, J., Reganold, J.P., 2010. Effects
 625 of soil type and farm management on soil ecological functional genes and microbial
 626 activities. *ISME J.* 4, 1099–1107. doi:10.1038/ismej.2010.42
- 627 Resource Efficient Agricultural Production (REAP)-Canada, 2008. ARF07 P ROJECT
 628 Optimization of Switchgrass Management for Commercial Fuel Pellet Production.
 629 OMAFRA-Alternative Renew. Fuels Res. Dev. Fund.
- 630 Roy, A.K., Wagner-Riddle, C., Deen, B., Lauzon, J., Bruulsema, T., 2014. Nitrogen application
 631 rate, timing and history effects on nitrous oxide emissions from corn (*Zea mays* L.). *Can. J.*
 632 *Soil Sci.* 94, 563–573. doi:10.4141/cjss2013-118
- 633 Sanscartier, D., Deen, B., Dias, G., MacLean, H.L., Dadfar, H., McDonald, I., Kludze, H., 2014.
 634 Implications of land class and environmental factors on life cycle GHG emissions of
 635 *Miscanthus* as a bioenergy feedstock. *GCB Bioenergy* 6, 401–413. doi:10.1111/gcbb.12062
- 636 Schlesinger, W.H., 2013. An estimate of the global sink for nitrous oxide in soils. *Glob. Chang.*
 637 *Biol.* 19, 2929–2931. doi:10.1111/gcb.12239
- 638 Soil Quality National, 2006. Crop Residue Removal For Biomass Energy Production: Effects on
 639 Soils and Recommendations. Tech. Note 19, 1–7.
- 640 Sokhansanj, S., Mani, S., Turhollow, A., Kumar, A., Bransby, D., Laser, L.L., Laser, M., 2009.
 641 Large-scale production, harvest and logistics of switchgrass (*Panicum virgatum* L.) –
 642 current technology and envisioning a mature technology. *Biofuels Bioprod. Bioprocess.*
 643 124–141. doi:10.1002/bbb
- 644 Stone, M.M., Kan, J., Plante, A.F., 2015. Parent material and vegetation influence bacterial
 645 community structure and nitrogen functional genes along deep tropical soil profiles at the
 646 Luquillo Critical Zone Observatory. *Soil Biol. Biochem.* 80, 273–282.
 647 doi:10.1016/j.soilbio.2014.10.019
- 648 Szukics, U., Abell, G.C.J., Hödl, V., Mitter, B., Sessitsch, A., Hackl, E., Zechmeister-
 649 Boltenstern, S., 2010. Nitrifiers and denitrifiers respond rapidly to changed moisture and
 650 increasing temperature in a pristine forest soil. *FEMS Microbiol. Ecol.* 72, 395–406.



- 651 doi:10.1111/j.1574-6941.2010.00853.x
- 652 Throbäck, I.N., Enwall, K., Jarvis, A., Hallin, S., 2004. Reassessing PCR primers targeting nirS,
653 nirK and nosZ genes for community surveys of denitrifying bacteria with DGGE. FEMS
654 Microbiol. Ecol. 49, 401–17. doi:10.1016/j.femsec.2004.04.011
- 655 Ver Hoef, J.M. and Boveng, P.L., 2007. Quasi-Poisson vs. Negative Binomial Regression: How
656 should we model overdispersed count data? Ecology 88, 2766–2772.
- 657 Watrud, L.S., Reichman, J.R., Bollman, M. a., Smith, B.M., Lee, E.H., Jastrow, J.D., Casler,
658 M.D., Collins, H.P., Fransen, S., Mitchell, R.B., Owens, V.N., Bean, B., Rooney, W.L.,
659 Tyler, D.D., King, G. a., 2013. Chemistry and Microbial Functional Diversity Differences
660 in Biofuel Crop and Grassland Soils in Multiple Geographies. BioEnergy Res. 6, 601–619.
661 doi:10.1007/s12155-012-9279-y
- 662 Wolsing, M., Priemé, A., 2004. Observation of high seasonal variation in community structure of
663 denitrifying bacteria in arable soil receiving artificial fertilizer and cattle manure by
664 determining T-RFLP of nir gene fragments. FEMS Microbiol. Ecol. 48, 261–71.
665 doi:10.1016/j.femsec.2004.02.002
- 666 Wu, L., Osmond, D.L., Graves, A.K., Burchell, M.R., Duckworth, O.W., 2012. Relationships
667 Between Nitrogen Transformation Rates and Gene Abundance in a Riparian Buffer Soil.
668 Environ. Manage. 50, 861–874. doi:10.1007/s00267-012-9929-z
- 669 Yin, C., Fan, F., Song, A., Li, Z., Yu, W., Liang, Y., 2014. Different denitrification potential of
670 aquic brown soil in Northeast China under inorganic and organic fertilization accompanied
671 by distinct changes of nirS - and nirK -denitrifying bacterial community. Eur. J. Soil Biol.
672 65, 47–56. doi:10.1016/j.ejsobi.2014.09.003
- 673 Zumft, W.G., 1997. Cell biology and molecular basis of denitrification. Microbiol. Mol. Biol.
674 Rev. 61, 533–616.
- 675

76 Table 1. Mean soil properties measured at the Elora Research Station.

Cropping System/ Harvest	N Rate (kg ha ⁻¹)	*Bulk Density (g cm ⁻³)	% Organic Carbon			% Total Carbon			Yield (dry tonnes ha ⁻¹)		†Mean
			0-15cm	15-30cm	0-15cm	15-30cm	2011	2012			
Corn-soybean	Fall	0	1.21 AB	1.88	1.06	2.22	1.86	5.341	2.912	E	
Corn-soybean	Fall	160	1.27 A	1.79	1.47	2.25	2.11	9.92	7.882	BC	
Miscanthus	Fall	0	1.10 B	2.06	1.44	2.27	1.72	17.62	12.77	A	
Miscanthus	Fall	160	1.10 B	2.13	1.63	2.36	1.84	17.43	18.32	A	
Miscanthus	Spring	0	1.13 AB	2.09	1.53	2.31	1.69	12.66	13.38	AB	
Miscanthus	Spring	160	1.13 AB	2.24	1.42	2.47	1.89	14.33	14.56	A	
Switchgrass	Fall	0	1.11 B	2.12	1.43	2.33	1.61	7.648	6.458	CD	
Switchgrass	Fall	160	1.09 B	2.12	1.34	2.32	1.73	11.1	10.45	AB	
Switchgrass	Spring	0	1.11 B	2.09	1.23	2.32	1.55	6.33	4.146	DE	
Switchgrass	Spring	160	1.21 AB	1.92	1.33	2.23	1.7	6.905	6.441	CD	

*Means of bulk density (n=6) followed by the same letter within one column are not significantly different according to a post-hoc Tukey's means comparison (p<0.05); carbon measurements (n=3) were not significantly different between treatments. †Mean yields (n=3) followed by the same letter are not significantly different according to a post-hoc Tukey's means comparison (p<0.05).

77

78

79

80

81

82

83

84

85

86

87





688 Table 2. Mean gene abundance responses to crop and harvest management, averaged over nitrogen application rate and time at the Elora Research
 689 Station.

Cropping System	Management	†Total soil bacteria (gene copy g ⁻¹ soil)		†Soil denitrifying bacteria (gene copy g ⁻¹ soil)	
		16S	<i>nirS</i>	<i>nosZ</i>	<i>nirS:nosZ</i> (x10 ⁻²)
Corn-Soybean	Fall Harvest	1.35 x 10 ⁹ b	1.95 x 10 ⁶ b	2.63 x 10 ⁵ b	7.42
	Fall Harvest	1.38 x 10 ⁹ b	2.30 x 10 ⁶ ab	4.47 x 10 ⁵ a	5.15
Miscanthus	Spring Harvest	1.91 x 10 ⁹ a	3.02 x 10 ⁶ a	5.25 x 10 ⁵ a	5.75
Switchgrass	Fall Harvest	1.41 x 10 ⁹ b	2.19 x 10 ⁶ b	3.55 x 10 ⁵ ab	6.17
Switchgrass	Spring Windrow	1.48 x 10 ⁹ ab	2.46 x 10 ⁶ ab	3.98 x 10 ⁵ ab	6.18

†Means followed by the same letter within one column are not significantly different according to post-hoc Tukey's means comparison at p < 0.05 (n = 24).



690 **Figure Captions**

691 **Figure 1.** Mean daily air temperature ($^{\circ}\text{C}$) and daily precipitation (mm) at the Elora Research Station from
692 January 2011 to November 2012. Soil gravimetric H_2O was measured on a per-sample basis and is shown as
693 crop means ($\pm\text{SE}$) for each sampling date (May 9th, 2011; October 30th, 2011; May 2nd, 2012 and October 20th,
694 2012) ($n=12$ in perennial grasses, $n=6$ in corn-soybean rotation).


695 **Figure 2.** Mean soil $\text{NH}_4\text{-N}$ and $\text{NO}_3\text{-N}$ (mg g^{-1} dry soil $\pm\text{SE}$) in annual and perennial biomass cropping
696 systems under varied harvest and N management at the Elora Research Station. CS = corn-soybean, SF = fall-
697 harvested switchgrass, SS = spring-harvested switchgrass, MF = fall-harvested miscanthus and MS = spring-
698 harvested miscanthus. Different letters within panels indicate significant differences according to a post-hoc
699 Tukey's test ($p<0.05$).

700 **Figure 3.** Mean log gene copies (g^{-1} dry soil $\pm\text{SE}$) in annual and perennial biomass cropping systems under
701 varied harvest management at the Elora Research Station ($n=6$) over time. Different letters within panels
702 indicate significant differences according to a post-hoc Tukey's test ($p<0.05$).

703 **Figure 4a.** Principal Component Analysis; factor 1 accounted for 43.89% variance and factor 2 accounted for
704 23.84% variance. **4b.** Loading plot for principal components of response variables (*nirS*, *nosZ* and 16S rRNA
705 gene copies, and soil $\text{NO}_3\text{-N}$, soil $\text{NH}_4\text{-N}$, gravimetric soil H_2O)..


706 Figure 1.

707






708 Figure 2.



709




710 Figure 3.



711




712 Figure 4a.



713

714 Figure 4b.



715