

Dear Editor of Soil,

Thank you for your message. We did our best to take into account every comment and suggestion and we hope that the new version of our manuscript has been improved in terms of quality.

Considering the points asked, here you find our improvements :

-In order to be clear to all readers, we have precised our objectives in the introduction and changed the subtitles « step 1 » and « step 2 » by « composting » and « soil incubation ». Moreover, the word « co-(vermi)composting » has been deleted and changed by « composting » followed by « in presence of worms » or « in absence of worms ».

- The term windrow will be defined as followed in the manuscript « Briefly the composting process was performed in windrow which are long narrow piles of green waste ».

-Figures have been revised according to the reviews. For the figure 6, the name of each treatment has been deleted because the figure was not readable if they were written, so we have changed them by symbols, according to the treatment.

You will find the latest version, taking into account the remarks and suggestions highlighted in yellow.

Kind regards,

Justine Barthod

# **The effects of worms, clay and biochar on CO<sub>2</sub> emissions during production and soil application of co-composts**

Justine Barthod <sup>1</sup>, Cornélia Rumpel <sup>1</sup>, Remigio Paradelo <sup>2</sup>, Marie-France Dignac <sup>1</sup>

5 <sup>1</sup> Institute of Ecology and Environmental Sciences- Paris (iEES-Paris) UMR CNRS, INRA, UPMC.  
78850 Thiverval-Grignon. France

<sup>2</sup> University of Vigo, Department of Plant Biology and Soil Science, Facultade de Ciencias, As  
Lagoas s/n, 32004 Ourense, Spain

10 *Correspondence: justine.barthod@grignon.inra.fr*

**Abstract :**

In this study we evaluated CO<sub>2</sub> emissions during **composting** of green wastes with clay and/or biochar **in the presence and absence of worms**. The stability of the final products as well as their effect on C mineralization in soil were evaluated. The aim of the study was to test the following  
15 hypotheses: (1) interactions between clay and biochar and organic wastes lead to reduced CO<sub>2</sub> emissions during the composting process, (2) these interactions are enhanced in the presence of worms, and (3) more carbon is sequestered in soil after the use of the resulting compost/vermicompost as amendments. We added two different doses of clay, biochar and their  
20 mixture to pre-composted green wastes and monitored carbon mineralisation during 21 days in presence or absence of worms (species of the *Eisenia* genus). The organic materials were then added to a loamy Cambisol and the CO<sub>2</sub> emissions were monitored during 30 days in a laboratory incubation.

Our results indicated that the addition of clay or clay/biochar mixture reduced carbon mineralization during co-composting without worms by up to 44%. However, in the presence of worms, CO<sub>2</sub>  
25 emissions increased for all treatments except for the low clay dose. The production conditions had more influence on carbon mineralization in soil for composts than for vermicomposts except for the low clay treatment, which showed a more reduced CO<sub>2</sub> emissions compared to regular compost.

**In summary, the addition of worms during co-composting with clay and biochar speeds up CO<sub>2</sub> emissions in most cases. Therefore,** the production of a low CO<sub>2</sub> emission amendment requires  
30 optimisation of OM source, co-composting agents and worm species. The effect of the resulting material on soil fertility has to be evaluated.

**Keywords:** carbon mineralization; worm; composting; biochar; clay; soil.

## 1.Introduction

Land use changes are responsible for the steady increase of CO<sub>2</sub> in the atmosphere, along with industrial activity and the use of fossil fuels. In this context, massive soil organic matter (OM) loss is observed, leading to the decline of many soil ecosystem services, such as fertility and carbon storage (Smith et al., 2015). These global changes of the earth's climate and (agro-)ecosystems have major environmental, agronomic but also social and economic consequences, which could be attenuated by the rebuilding of soil OM stocks (IPCC, 2014). Increasing soil carbon may be possible with the use of composted organic wastes as alternative fertilisers (Ngo et al, 2011, 2012), which could counterbalance the concentration of greenhouse gases in the atmosphere through soil carbon sequestration (Lashermes et al, 2009)

Two well-known aerobic processes based on microbial activity are able to transform organic wastes into valuable soil amendments: composting and vermicomposting. Composting has been traditionally used and leads to stabilized organic amendments with fertilization potential. During vermicomposting the presence of worms induces a continuous aeration resulting in a faster organic matter transformation (Lazcano et al, 2008; Paradelo et al, 2012). However, vermicomposting and composting both emit greenhouse gases such as CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O (Hobson et al, 2005; Chan et al, 2011; Thangarajan et al, 2013). In addition, the final products of these processes lead to greenhouse gas emissions after their application to soil (Cambardella et al, 2003; Bustamante et al, 2007). These emissions can originate from the mineralization of (vermi)compost OM itself or maybe due to the mineralization of native soil OM following increased microbial development and activity, a mechanism known as priming effect (Bustamante et al, 2010).

In order to optimize the recycling of waste carbon, there is a need to enhance OM stabilization during (vermi)composting. Stabilization mechanisms are poorly known for composting processes, while they have been widely studied in soils. Enhancing C stabilization in composts could thus benefit from an analogy with the mechanisms known to occur in soils (v. Lützow et al, 2006): spatial inaccessibility, selective preservation due to chemical recalcitrance, and formation of

organo-mineral associations. Among these processes, the association of OM with minerals is the most efficient for C stabilization on long time scales (Kleber *et al*, 2015). Therefore, a variety of minerals has been used to reduce gas emissions (CO<sub>2</sub>, CH<sub>4</sub>, NH<sub>3</sub> and N<sub>2</sub>O) during co-composting (Bolan *et al*, 2012; Wang *et al*, 2014; Chowdhury *et al*, 2015), e.g. clay minerals during composting of poultry manure (Bolan *et al*, 2012) or zeolite during (vermi)composting of wastes (Wang *et al*, 2014). However, to the best of our knowledge no studies have been carried out to evaluate the effect on carbon stability the resulting organic materials after their addition to soil.

In addition, many recent studies explored the potential benefits of biochar as soil amendment due to its physical and chemical properties, (Chan *et al*, 2007; Kookana *et al*, 2011). Biochar results from the incomplete combustion or pyrolysis of various feedstock materials. The biochar production process transforms OM into aromatic products, which are resistant against microbial decomposition and show increased adsorption properties compared to untransformed OM (Lehman *et al*, 2006). As a result, the use of biochar as co-composting agent leads to a reduction of carbon emissions due to adsorption of organic constituents on the biochar surface (Rogovska *et al*, 2011; Jindo *et al*, 2012; Vu *et al*, 2015).

To further enhance the protection of OM through the formation of organo-mineral or OM-biochar associations during co-composting, the addition of worms may be a promising avenue. In general, organo-mineral associations are enhanced by the presence of worms, due to the simultaneous ingestion of OM and minerals (Shipitalo and Protz, 1989). Micro-aggregates formed inside the worm guts improve physical protection of C (Bossuyt *et al*, 2005). However, these interactions have only been evidenced for soil earthworms and have never been evaluated as a strategy to reduce CO<sub>2</sub> emissions during co-composting. One study investigated the effect of biochar on worm activity during vermicomposting of a sludge biochar mixture (Malinska *et al*, 2016). However, to the best of our knowledge, no studies have investigated the effect on carbon emissions of biochar as a co-composting agent during vermicomposting.

In this study, we evaluated if the addition of clay, biochar and their mixture to organic wastes

influences CO<sub>2</sub> emissions (1) during (vermi)composting and (2) after the use of the final products as soil amendments. To do so, we carried out two laboratory experiments and evaluated CO<sub>2</sub> emissions during (1) production of organic amendments in presence or absence of worms and (2) soil incubation with the amendments. We hypothesized that carbon stabilization during composting may be increased by addition of (a) montmorillonite, a 2:1 clay, able to form organo-mineral associations; (b) biochar, able to protect OM by adsorption and (c) their mixture, which could create synergistic effects. In addition, we tested the effect of two different amounts of clay on the changes of CO<sub>2</sub> emissions during co-composting with and without worms and after addition to soil. The aim of the study was to investigate, if the presence of worms during composting with additives leads to enhanced C stabilisation.

## 2. Materials and methods

### 2.1 Compost, additives and worms

A pre-composted green waste was sampled in its maturation phase at BioYvelines service, a platform of green waste composting located 30 km West from Paris (France). The green wastes were a mix of shredded leaves, brushwood and grass cuttings collected from households or firms near the platform. Briefly, the composting process was performed in windrows, which are long narrow piles of green waste. Aerobic conditions and optimal humidity (approximately 45 %) were maintained through mechanical aeration and water sprinkling. The pre-composted material was sampled after 4 months, at the beginning of the maturation phase. Compost pH was 8.5 and the C:N ratio was 13.6 with 205.1 mg.g<sup>-1</sup> of organic carbon (OC) and 13.3 mg.g<sup>-1</sup> of nitrogen (N). After sampling, the compost was air-dried and sieved at 3 mm for homogenization.

Montmorillonite, a 2:1 clay, was purchased from Sigma-Aldrich. The clay's pH was between 2.5 and 3.5 and its specific surface area (SSA) was 250 m<sup>2</sup>/g. Montmorillonite was chosen because organo-mineral interactions depend on clay mineralogy (1:1 clay or 2:1 clay). In general, 2:1 minerals offer a bigger contact area for OM bonding and create stronger bonds with OM than the

1:1 minerals (Kleber *et al*, 2015). Thus numerous organo-mineral associations were expected due to  
115 this large SSA.

The biochar was provided by Advanced Gasification Technology (Italy). It was produced by gasification at 1200°C of a conifer feedstock and had a pH of 9.3 and a C:N ratio of 40:30, with 806 mg g<sup>-1</sup> of OC and 0.2 mg g<sup>-1</sup> of N (Wiedner *et al*, 2013).

*Eisenia andrei* and *Eisenia foetida* worms were purchased from La Ferme du Moutta, a worm farm  
120 in France. The two species were chosen because they present a high rate of consumption, digestion and assimilation of OM, can adapt to a wide range of environmental factors, have short life cycles, high reproductive rates and endurance and resistance to handling (Dominguez and Edwards, 2011).

### 2.3 Experimental setup

125 The present study was designed to evaluate and compare the CO<sub>2</sub> emissions of the different organic materials during the production phase and after their addition to soil (Fig.1)

### **Composting**

Composting was carried out at ambient temperature in the laboratory with 10 treatments and four  
130 replicates per treatment: (i) compost alone (control), (ii) compost with 25% (w/w) of montmorillonite (low clay treatment), (iii) compost with 50 % (w/w) of montmorillonite (high clay treatment), (iv) compost with 10% (w/w) of conifer biochar and (v) compost with a mixture of biochar (10% w/w) and montmorillonite (25% w/w). All treatments were established with and without worms (Table 1). Considering that a clay can retain 1 mg C per m<sup>2</sup> (Feng *et al*, 2011), 50%  
135 of clay and 25% of clay were chosen in order to theoretically retain 60% and 30% of the total carbon from the compost. In addition, biochar was moistened before addition to compost to avoid worm mortality due to desiccation (Li *et al*, 2011). The addition of 10% of biochar was chosen according to Weyers and Spokas (2011) to avoid negative effects on worms.

Worms were raised in the same compost as used in the experiment. Eight adult worms were chosen

140 and cleaned to remove adhering soil/compost before estimating their body mass and added to the organic material.

The experiments were carried out in 2L jars. A dry mass of 75 g of pre-composted material was used in each treatment. Water was sprinkled on jars at the beginning of the experiment to reach an optimal moisture level of 80-90% (water content by weight), which was maintained throughout the experimental period. Jars were placed in the dark at ambient temperature (24°C on average). The (vermi)composting was stopped after 21 days, when all the OM should have been ingested (75 g of compost for 8 worms). Indeed a worm can ingest its weight at maximum per day (0.5g).

At the end of the experiment, worms were counted and weighed again. The amount of cocoons and juveniles was recorded. The final (vermi)composts were air dried, sieved at 2 mm and an aliquot was ground for further analyses.

### **Soil incubation**

A loamy cambisol soil was collected for the laboratory experiment from the experimental site of a long-term observatory for environmental research (ORE-ACBB) of INRA, near Lusignan in the South-West of France. This soil was used for crop production for the last three years. The soil was collected at 0-10 cm depth, sieved at 4 mm, homogenized and kept at 4°C until the beginning of the experiment. The soil is carbonate-free and has the following characteristics: pH 6.4, N content 1.15 mgN g<sup>-1</sup>, carbon content 10.56 mgC g<sup>-1</sup>, sand 11%, clay 17% and silt 72% (Chabbi *et al*, 2009).

For all the treatments, 57 g of dry soil were weighed and placed into 2L glass jars. The mixtures were homogenized. All ten organic materials obtained during composting were applied to soil at a rate of 67g kg<sup>-1</sup> (dry weight). Amended and unamended soils were incubated in four replicates in the dark at ambient temperature. Soil moisture was adjusted to 18 % (dry weight) and maintained throughout the experiment by compensating weight losses with deionised water. The CO<sub>2</sub> emissions were measured during 30 days as described below.

165



### 2.3 Carbon mineralisation

CO<sub>2</sub> emissions were measured in the headspace of the jars according to Anderson (1982). All incubation jars contained a vial with 30 mL of 1M NaOH (composting) or 0.5M (soil incubation) to trap CO<sub>2</sub>. The NaOH vials were covered with a tissue to avoid contamination of the NaOH solution by worms. During co-composting step, NaOH traps were replaced at day 1, 2, 3, 4, 8, 11, 14, 16, 18 and 21. During the incubation with soil, vials were replaced at day 1, 2, 4, 7, 14 and 22. Phenolphalein and BaCl<sub>2</sub> solution in excess were added to a 10 mL aliquot of NaOH sampled from each vial. The solution was titrated with 1M HCl until neutrality to determine the CO<sub>2</sub>-C released. Three empty jars were used as control.

Results are expressed in mg CO<sub>2</sub>-C/ g compost (dry weight) or in mg CO<sub>2</sub>-C/ g total organic carbon (TOC) according to the formula:

$$\text{Released CO}_2 - \text{C} = \frac{(B - V) * M * E}{P}$$

where B is the volume of HCl used to titrate the control (mL); V the volume of HCl used to titrate the sample (mL); M the normality of HCL (1M); E (22) the molar mass of CO<sub>2</sub> divided by 2 (because 2 mol of OH<sup>-</sup> are consumed by one mol of CO<sub>2</sub>) and P the weight of the sample (grams).

### 2.4 Properties of the final products after composting

OC and N contents were measured using a CHN auto-analyzer (CHN NA 1500, Carlo Erba). A glass electrode (HANNA instruments) was used to measure pH in water extracts of (vermi)-composts (1:5). Dissolved organic carbon (DOC) contents were determined in 0.034 mol L<sup>-1</sup> K<sub>2</sub>SO<sub>4</sub> extracts (1:5 w/v) using a total organic carbon analyzer (TOC 5050A, Shimadzu).

### 2.5 Calculations and Statistical analysis

The amount of CO<sub>2</sub>-C mineralized was expressed as mgC per g of TOC. TOC includes for composting compost carbon and biochar carbon. For soil incubation, it includes soil carbon,

compost carbon and biochar carbon. Finally, a global carbon balance was done and calculated on the basis of the CO<sub>2</sub> emissions from the composting phase and the soil incubation after amendment. These results are expressed as mgC per g of TOC, including soil carbon, compost carbon and biochar carbon.

195 Additionally, for composting, the amount of CO<sub>2</sub>-C mineralized was expressed as mgC per g of compost in order to focus on the carbon from the pre-composted material (the amount of biochar and clay was not included). Biochar is not supposed to be mineralized during this step because it is produced at high temperatures and therefore its carbon is supposed to have a high chemical recalcitrance against biological decomposition (McBeath and Smernik, 2009). Biochar produced at high temperatures showed a very low carbon emissions during a 200 days incubation in soil (Naisse et al, 2014), so that we can hypothesize that its mineralization can be neglected compared to OM mineralization during 21 days.

A first-order model was applied to describe the rate of carbon mineralization during composting (step 1):

205 
$$C = C_0 (1 - e^{-(kt)}), \quad \text{equation 1}$$

where C is the cumulative amount of CO<sub>2</sub>-C mineralized after time t (mgC g<sup>-1</sup> compost), C<sub>0</sub> is the initial amount of organic carbon (mgC g<sup>-1</sup> compost), t is the incubation time (days), and k is the rate constant of CO<sub>2</sub>-C mineralization (day<sup>-1</sup>).

All reported data are the arithmetic means of four replicates. A Kruskal-Wallis test was performed to assess the significance of differences of CO<sub>2</sub> emissions from the different treatments. A Student t test was run to investigate the influence of the different substrates on the worm development. Significance was declared at the 0.05 level. Statistical analyses were carried out using the R 3.12 statistical package for Windows (<http://www.r-project.org>).

## 215 **3.Results**

### *3.1 Properties of the co-(vermi)composts*

After 21 days of composting, OM had been processed into a homogeneous and aerated material in the presence of worms whereas regular composts had a compact aspect, illustrating the positive effects of worms on the physical structure of the final product. Total N and OC contents, DOC and pH of initial material and the different composts are shown in Table 2. The pH of composts and vermicomposts ranged from 7.9 to 8.7. The lowest pH was observed for the high clay treatments due to the addition of acidic clay material (pH 2.5 to 3.5). Co-composting with biochars did not lead to any change in pH (Table 2). This may be due to the alkaline pH of the initial material and the low amount of biochar added. Presence of worms during the composting phase had no effect on pH.

Total OC in composts and vermicomposts ranged from 118.6 mg g<sup>-1</sup> to 241.9 mg g<sup>-1</sup> and total N from 8.5 mg g<sup>-1</sup> to 13.5 mg g<sup>-1</sup>. Compared to initial material, OC was decreased significantly after 21 days of (vermi)composting in both control treatments while N concentrations remained unchanged. Addition of clay produced lower OC and N concentrations due to dilution, whereas the addition of the C-containing biochar increased OC concentrations and decreased N concentrations.

Similarly to pH, the presence of worms had no effect on OC or N of the final product.

DOC in composts and vermicomposts ranged from 15.04 to 29.08 mg g<sup>-1</sup> TOC. DOC was similar to that of the initial material after 21 days of (vermi)composting for the two controls whereas the presence of additives significantly decreased the DOC in all other treatments. The lowest DOC concentrations were recorded for composts and vermicomposts produced with biochar/clay mixture.

The presence of worms had only an effect on DOC for compost produced with clay, decreasing its concentration by 12% (high clay treatment) and 16% (low clay treatment).

### *3.2 Worm growth and reproduction*

The number of worms and their total weight were measured before and after 28 days of composting. The number of worms did not vary (p-value > 0.07) and neither did their total weight (p-value = 0.34). Cocoons and juveniles were separated manually from the substrates and counted at the end of composting. The number of cocoons and juveniles in treatments ranged from none to 4: high and

low clay treatments did not differ significantly from the control treatments (p-value= 0.39). No cocoon and no juvenile were counted in the biochar treatment. Finally, in treatments with  
245 clay/biochar mixture, the number of cocoons and juveniles was significantly higher (p-value=0.003) compared to the treatment with biochar alone with an average of 3 cocoons and one juvenile.

### 3.3 Carbon mineralisation during composting

Cumulative carbon emissions at the end of the experiment ranged from 6.4 to 11.9 mg CO<sub>2</sub>-C g<sup>-1</sup> compost in treatments without worms and from 7.9 to 12.0 mg CO<sub>2</sub>-C g<sup>-1</sup> compost in treatments  
250 with worms (Fig. 3 and 4). The presence of worms during composting (Fig. 2) had contrasting effects on C mineralisation (mg g<sup>-1</sup> TOC): 1) no change in treatments free of additives; 2) decrease in the low clay treatments and 3) increase in the treatments with high clay and biochar/clay mixture. In both control treatments, the amounts of carbon mineralized after 21 days were similar, about 12 mg CO<sub>2</sub>-C g<sup>-1</sup> compost. Composting with clay led to a significant decrease of the carbon emissions  
255 compared to the controls: in the low clay treatment, emission decreased by 15% without worms and by 34% with worms (Fig. 3 and 4); in the high clay treatment emissions decreased by 43% without worms and by 24% with worms. With biochar, reduction of CO<sub>2</sub> emissions was only observed in treatments without worms: a 24% decreased with biochar alone and 46 % decrease with biochar/clay mixture (Fig. 2 and 3). The cumulative CO<sub>2</sub> emissions during composting did not reach  
260 a plateau for any treatment (Fig. 3 and 4), but the experimental period was limited by worms activity since worms had processed all organic material after 21 days.

Rate constants obtained with the first-order kinetic model (eq. 1) are listed in Table 2. Similar high rate constants suggest a rapid carbon mineralization from control treatments. In general, the presence of worms increased rate constants, except for the low clay and control treatments. Lowest  
265 rate constants were observed for the low clay treatment (with worms) and the high clay and clay/biochar mixture (without worms). Biochar alone decreased carbon mineralization only in treatments without worms.

### 3.4 Carbon mineralisation during incubation with soil

270 Carbon emissions from the soil amended with the organic materials are shown in Figure 5. Cumulative emissions at day 30 ranged from 8.95 to 18.20 mg g<sup>-1</sup> TOC. Generally, the application of organic materials to soil led to a larger amount of carbon mineralized compared to the soil without amendments. The carbon emissions were influenced by the compost production procedure (additives and worms). The highest emissions were recorded for soil amended with composts free  
275 of additives. Compared to soil amended with regular composts, vermicomposts decreased the carbon emissions from amended soil only when produced without additives or with low clay addition. Organic amendments produced with high clay addition induced similar carbon emissions from soil. Compost produced in the presence of biochar showed the lowest effect on mineralization in soil. When biochar was mixed with clay, the final product induced lower carbon emissions from  
280 soil when produced in the presence of worms compared to those produced without worms.

Figure 6 shows the correlation between the amount of carbon mineralized from the amended soil and the DOC of the respective organic material. The relationship was stronger for the soil amended with composts compared to the soil amended with vermicomposts (Fig.6, respectively R<sup>2</sup>=0.67 and R<sup>2</sup>=0.07).

285

## 4. Discussion

### 4.1 Effect of worms and additives on compost properties

The presence of worms had no effect on the pH (Table 2), all the treatments tending to a slightly alkaline pH. By contrast, some authors observed a decrease in pH during vermicomposting of  
290 household wastes (Frederickson *et al*, 2007) or cattle manure (Lazcano *et al*, 2008). The contrasting results may be explained by a lower production of CO<sub>2</sub> and organic acids by micro-organisms in our experiment due to the almost mature pre-composted material used compared to the fresh green wastes used in previous experiments.

The C:N ratio was significantly higher in (vermi)composts produced in the presence of biochar, due

295 to addition of carbon enriched material. No difference was observed in vermicompost treatments compared to compost treatments, concerning the OC and the total N. These results are in line with those obtained by Ngo *et al* (2013), who suggested that the elemental composition and the chemical structures present in different composts and vermicomposts could be similar.

#### 300 4.2 Effect of worms and additives on carbon mineralization during composting

Data recorded (Fig. 3 and 4) for control treatments indicated in contrast to what is generally observed (e.g. Chan *et al*, 2011), that the presence of worms did not lead to higher CO<sub>2</sub> emissions during composting. This is probably due to the OM used, which was almost mature compost and, may thus be characterised by lower degradability than the organic wastes originating from  
305 households usually used for (vermi)composting.

Addition of clay and biochar reduced carbon emissions during composting (Fig. 3). Similar results were obtained by other authors for co-composting in absence of worms with clay additives (Bolan *et al*, 2012) or biochar (Dias *et al*, 2010). These data may indicate carbon stabilization by physico-chemical protection of OM on clay and/or biochar surfaces. Carbon storage generally increases  
310 linearly with increasing clay concentration (Hassink, 1997). This is in line with our results, showing proportional CO<sub>2</sub> decrease, when clay content and thus **potentially available** surface area **increased**.

By contrast, in the presence of worms, C mineralization was more reduced for the low clay compared to the high clay treatment (Fig. 4). As we observed similar worm biomass in both treatments, we hypothesize that high clay contents may have negative effects on worm activity and  
315 therefore the formation of organo-mineral associations. This hypothesis is supported by the results of Klok *et al* (2007), who showed that *Lumbricus rubellus* worms can have their life cycle influenced by a high content of clay in soil leading to anaerobic conditions and soil compaction. Our results suggest that a 50% proportion of montmorillonite also impacts the activity of *Eisenia andrei* and *foetida* species. In contrast, in the low clay treatment, worm activity most probably  
320 increases the formation of organo-mineral associations, thus leading to higher reduction of CO<sub>2</sub>

emissions compared to regular composting (Fig. 3). These results indicate that the protective capacity of clay minerals may be enhanced by worm activity, up to a threshold of the clay:OM ratio, above which species of the *Eisenia* genus are no longer able to reduce CO<sub>2</sub> emissions. Species of the *Eisenia* genus (*Fetida and Andrei*) belong to the epigenic worm species living at the soil surface in leaf litter, one of the three ecological lifetypes described by Bouché (1977). Therefore, they are well adapted to process pure organic matter and may be less suited for co-composting with minerals. The optimal clay: OM ratio to allow for maximal reduction of CO<sub>2</sub> emissions remains to be assessed as well as the possibility to use other worm species more adapted to ingestion of minerals.

330

Without worms, biochar addition led to a reduction of CO<sub>2</sub> emissions up to 44% compared to the control (Fig. 3). This is in line with the capacity of biochar to adsorb and protect labile organic compounds from degradation (Augustenborg *et al*, 2012; Ngo *et al.*, 2013; Naisse *et al.*, 2015). However, other studies showed no significant reduction of CO<sub>2</sub> emissions when biochar was used for co-composting (Sánchez-García *et al*, 2015). These contrasting results may be explained by variable physico-chemical properties of biochar: the biochar used in this study was produced by gasification while Sánchez-García *et al*, 2015 used a biochar produced by pyrolysis.

In the presence of worms, the addition of biochar and biochar/clay mixture induced higher CO<sub>2</sub> emission (p-value > 0.1) compared to regular composting (Fig. 3 and 4). Three hypothesis might explain that worms drastically modify the complex interactions between clay, biochar and pre-composted OM: 1) the worms might increase in their gut the contact between clay and biochar, leading to the partial saturation of clay surfaces with carbon compounds originating from biochar and thus to a reduction of the available surface area; 2) the microbial colonization of biochar might be enhanced in the worm gut decreasing their long-term resistance to bio-degradation; 3) biochars might enhance worm activity, as suggested by Augustenborg *et al* (2012) to explain the increase of CO<sub>2</sub> emissions when biochar was added to soil in the presence of worms.

345

The incidence of these three hypotheses probably depends on the biochar quality, which influences the effects of biochar on worm activity. Indeed biochar addition had contradictory effects on worm reproduction. . Biochar alone reduced the number of juveniles and cocoons of *Eisenia* to zero, 350 contrary to what Malińska *et al.* (2016) observed during the vermicomposting of a sewage sludge-biochar mixture. These contrasting influences of biochar on worm activity may be explained by the different biochar chemical characteristics due to specific production processes (gasification in our study and pyrolysis in the Malinska *et al.* (2016) study)..

By contrast, we observed increased reproduction rates when biochar was used in combination with 355 clay, which is more representative of the soil matrix. For the development of soil earthworms, the presence of biochar has already been described as a potential risk (Liesch *et al.*, 2010). In soil, the negative effects of biochar on worm activity have been suggested to originate from a) a lack of nutrients following their adsorption on biochar, b) the presence of toxic compounds such as polycyclic aromatic hydrocarbons (PAH) mainly, or c) a lack of water (Li *et al.*, 2011). In our 360 experiment, the lack of nutrients was balanced by the presence of compost and the lack of water was avoided by a preliminary humidification of biochar before their addition. The presence of PAH or other potentially toxic substances might thus explain the effects that we observed. Further analyses and longer experiments should be carried out in order to investigate the reasons for these effects. Testing the influence of biochar of various origins (initial material and process) on 365 vermicomposting with clay compared with similar composting treatments would be necessary to elucidate the mechanisms responsible for their influence on carbon mineralization.

#### 4.3 Effect of co-(vermi)compost production conditions on carbon mineralization in soil and total carbon balance

370 The production conditions had more influence on carbon mineralization in soil for composts than vermicomposts (Table 3). Clay and biochar reduced the concentration of labile compounds in composts and vermicomposts leading to decreased DOC concentrations of the final amendments



(Table 1). But the CO<sub>2</sub> emissions after addition to soil were only reduced by clay and biochar addition when the compost was produced without worms (Fig 5). The rate of mineralization of organic amendments is generally linked to the labile carbon compounds (Chaoui *et al*, 2003) as was observed for the compost addition (Fig. 6). The lack of correlation between DOC and CO<sub>2</sub> emitted after addition to soil of vermicomposts suggests contrasted properties of DOC in composts and vermicomposts (Lazcano *et al*, 2008, Kalbitz *et al*, 2003).

CO<sub>2</sub> emitted from soil after the addition of amendments may originate from two sources: the mineralization of added carbon and the mineralization of native soil OM. Differences compared to the control (soil without amendments) may be explained by positive or negative priming effect, induced by microbial reaction to OM addition.

In the case of amendments produced with biochar alone, a negative priming effect could be observed, because the mineralization rate observed for this treatment was lower than for the control. This result is in line with many other studies reporting reduced mineralization of native soil OM after biochar amendment (Zimmerman *et al*, 2011). Our data evidenced that this phenomenon may also occur for composts, when biochar is used as co-composting agent. Negative priming was not observed for co-composts produced in the presence of worms.

In order to evaluate the positive or negative effect of each additive and process (with or without worms) on C mineralization, the CO<sub>2</sub> emissions during composting and during incubation of amended soil were summed up and expressed as mgC g<sup>-1</sup> TOC. The carbon emissions during both experiments were influenced differently by the (vermi)compost production procedure (additives and worms). The lowest total carbon emissions were recorded for compost and vermicompost produced in presence of biochar. Low clay treatment efficiently reduced carbon emissions in the presence of worms. In all other treatments, except the control, CO<sub>2</sub> emissions were higher for vermicompost due to higher emissions during the production step.

## 5. Conclusion

This study tested the influence of worm species of the *Eisenia* genus on CO<sub>2</sub> emissions during co-composting of green wastes with clay, biochar and their mixture. We established the complete carbon balance taking into account production of amendments as well as the effect after their addition to soil. Most additives were found to decrease CO<sub>2</sub> release during composting. In the presence of worms, 25% of clay led to greater OM protection than 50%. The opposite was observed in the absence of worms. Our results thus evidenced a threshold of clay concentrations for *Eisenia* worms, above which CO<sub>2</sub> emissions are no longer reduced. Biochar had a negative effect on carbon emissions for all treatments. Biochar/clay mixture resulted in synergistic effects for treatment without worms. We conclude that the use of additives may have the potential to greatly reduce CO<sub>2</sub> emissions during co-composting. Worms generally speed up carbon mineralization except in treatments with low clay dose. The effect of the amendments on C mineralization after addition to soil was small in the short-term. We suggest that production conditions during composting have to be optimized in terms of total CO<sub>2</sub> reduction by choosing the minerals, their optimal ratio with OM and testing different worm species. The effects of these amendments on soil fertility and plant growth remain to be investigated. Further work need to be done to assess the long-term effect of these amendments.

415

## 6.Acknowledgements

Authors are grateful to the Pierre and Marie Curie University for Doctoral fellowship. We also acknowledge Daniel Billiou and Valérie Pouteau for their technical assistance. Financial support was provided by CNRS under the framework EC2CO program (LOMBRICOM project), ADEME under the framework of the DOSTE program (VERMISOL project) and by the Australian Research Council by a discovery grant (C corundum project). Dr. R. Paradelo holds a *Juan de la Cierva* post-doctoral contract from the Spanish Government (JCI-2012-11778).

## 7. References

- Anderson J.P.E (1982) Soil Respiration. *Methods of Soil Analysis, part 2. Chemical and*  
425 *Microbiological Properties*, second ed. p. 331-871.
- Augustenborg C.A., Hepp S., Kammann C., Hagan D., Schmidt O. and Müller C. (2012) Biochar  
and earthworm effect on soil nitrous oxide and carbon dioxide emissions. *Journal of*  
*Environmental Quality* 41, 1203-1209.
- Barral M.T., Paradelo R., Moldes A.B., Domínguez M., Díaz-Fierros F. (2009) Utilization of MSW  
430 compost for organic matter conservation in agricultural soils of NW Spain. *Resources,*  
*Conservation and Recycling* 53, 529-534.
- Bolan N.S., Kunhikrishnan A., Choppala G.K, Thangarajan R. and Chung J.W (2012) Stabilization  
of carbon in composts and biochars in relation to carbon sequestration and soil fertility. *Science*  
*of the Total Environment* 424, 264-270.
- 435 Bossuyt H., Six J. and Hendrix P.F. (2005) Protection of soil carbon by microaggregates within  
earthworm casts. *Soil Biology & Biochemistry* 37, 251-258.
- Bouché, M.B. (1977) Stratégies lombriciennes. In : Lohm, U., Persson, T., Soil Organisms as  
Components of Ecosystems. *Ecological Bulletin* 25, 122-132.
- Bustamante M.A., Pérez-Murcia M.D., Paredes C., Moral R., Pérez-Espinosa A. and Moreno-  
440 Caselles J. (2007) Short-term carbon and nitrogen mineralization in soil amended with winery  
and distillery organic wastes. *Bioresource Technology* 98, 3269-3277.
- Bustamante M.A., Said-Pullicino D., Paredes C., Cecilia J.A. and Moral R. (2010) Influences of  
winery-distillery waste compost stability and soil type on soil carbon dynamics in amended  
soils. *Waste Management* 30, 1966-1975.
- 445 Cambardella C.A., Richard T.L and Russel A. (2003) Compost mineralization in soil as a function  
of composting process conditions. *European Journal of Soil Biology* 39, 117-127.
- Chabbi A., Kögel-Knabner I. And Rumpel C. (2009) Stabilised carbon in subsoil horizons is located  
in spatially distinct parts of the soil profile. *Soil Biology and Biochemistry* 41, 256-261.

- 450 Chan K.Y., Zwieten L. V., Meszaros I., Downie A. and Joseph S. (2007) Agronomic values of greenwaste biochar as a soil amendment. *Australian Journal of Soil Research*, 45, 629-634.
- Chan Y.C., Sinha R.K. and Wang W.J. (2011) Emission of greenhouse gases from home aerobic composting, anaerobic digestion and vermicomposting of household wastes in Brisbane (Australia). *Waste Management and Research* 29(5), 540-548.
- 455 Chaoui H.I., Zibilske L.M. and Ohno T. (2003) Effects of earthworms casts and compost on soil microbial activity and plant nutrient availability. *Soil Biology and Biochemistry* 35, 295-302
- Chowdhury S., Bolan N.S., Seshadri B., Kunhikrishnan A., Wijesekara H., Xu Y., Yang J., Kim G-H., Sparks D. and Rumpel C. (2015) Co-composting solid biowastes with alkaline materials to enhance carbon stabilization and revegetation potential. *Environmental Science and Pollution Research*, 1-12.
- 460 Dias B.O., Silva C.A., Higashikawa F.S., Roig A. and Sánchez-Monedero M.A. (2010) Use of biochar as bulking agent for the composting of poultry manure: effect on organic matter degradation and humification. *Bioresource Technology* 101, 1239-1246.
- Dominguez J. and Edwards C.A. (2011) *Biology and ecology of earthworm species used for vermicomposting*. Taylor and Francis Group.
- 465 Frederickson J., Howell G. and Hobson A.M. (2007) Effect of pre-composting and vermicomposting on compost characteristics. *European Journal of Soil Biology* 43, S320-S326.
- Hassink J. (1997) The capacity of soils to preserve organic C and N by their association with clay and silt particles. *Plant and Soil* 191: 77-87.
- 470 Hobson A.M., Frederickson J. and Dise N.B. (2005) CH<sub>4</sub> and N<sub>2</sub>O from mechanically turned windrow and vermicomposting systems following in-vessel pre-treatment. *Waste Management* 25, 345-352.
- IPCC (2014) *Climate change 2014. Synthesis Report*.
- Jindo K., Suto K., Matsumoto K., García C., Sonoki T. and Sánchez-Monedero M.A. (2012) Chemical and biochemical characterisation of biochar-blended composts prepared from poultry

- 475 manure. *Bioresource Technology* 110, 396-404.
- Kalbitz K., Schmerwitz J., Schwesig D. and Matzner E. (2003) Biodegradation of soil-derived dissolved organic matter as related to its properties. *Geoderma* 113, 273-291.
- Kleber M., Eusterhues K., Keiluweit M., Mikutta C., Mikutta R. and Nico P.S. (2015) Mineral-Organic Associations: formation, properties and relevance in soil environments. *Advances in*  
480 *Agronomy* 130, 1-140.
- Klok C., Faber J., Heijmans G. and Bodt J. (2007) Influence of clay content and acidity of soil on development of the earthworm *Lumbricus rubellus* and its population level consequences. *Biol. Fertil. Soils* 43: 549-556.
- Kookana R.S., Sarmah A.K., Zwieten L.V., Krull E. and Singh B. (2011) Biochar application to  
485 soil: agronomic and environmental benefits and unintended consequences. *Advances in Agronomy*, Volume 112, chapitre 3.
- Lahermes G., Nicolardot B., Parnaudeau V., Thuriès L., Chaussod R., Guillotin M.L., Linères M., Mary B., Metzger L., Morvan T., Tricaud A., Villette C. and Houot S. (2009) Indicator of potential residual carbon in soils after exogenous organic matter application. *European Journal*  
490 *of Soil Science* 60, 297-310.
- Lazcano C., Gomez-Brandon M. and Dominguez J. (2008) Comparison of the effectiveness of composting and vermicomposting for the biological stabilization of cattle manure. *Chemosphere* 72, 1013-1019.
- Lehmann J., Gaunt J. and Rondon M. (2006) Biochar sequestration in terrestrial ecosystems- a  
495 review. *Mitigation and adaptation strategies for global change* 11, 403-427.
- Li D., Hockaday W.C., Masiello C.A. and Alvarez P.J.J. (2011) Earthworm avoidance of biochar can be mitigated by wetting. *Soil Biology & Biochemistry* 43, 1732-1737.
- Liesch A. M., Weyers S.L., Gaskin J.W. and Das K.C (2010) Impact of two different biochars on earthworm growth and survival. *Annals of Environmental Science*, Vol 4, 1-9.
- 500 Malińska K., Zabochnicka-Świątek M., Cáceres R. and Marfà O. (2016) The effect of precomposted

sewage sludge mixture amended with biochar on the growth and reproduction of *Eisenia fetida* during laboratory vermicomposting. *Ecological Engineering* 90, 35-41.

- McBeath A.V. and Smernik R.J. (2009) Variation in the degree of aromatic condensation of chars. *Organic Geochemistry* 40, 1161-1168.
- 505 Naisse C., Girardin C., Davasse B., Chabbi A. and Rumpel C. (2015) Effect of biochar addition on C mineralisation and soil organic matter priming in two subsoil horizons. *J Soils Sediments* 15, 825-832.
- Ngo P-T, Rumpel C., Doan T.T. and Jouquet P. (2012) The effect of earthworms on carbon storage and soil matter composition in tropical soil amended with compost and vermicompost. *Soil*  
510 *Biology and Biochemistry* 50, 21-220.
- Ngo P-T, Rumpel C., Ngo Q-A., Alexix M., Vargas V.G., De la Luz Mora Gil M., Dang D-K and Jouquet P. (2013) Biological and chemical reactivity and phosphorous forms of buffalo manure compost, vermicompost and their mixture with biochar. *Bioresource Technology* 148, 401-407.
- Ngo P-T., Rumpel C., Dignac M-F., Billou D., Tran Duc T. and Jouquet P. (2011) Transformation of  
515 buffalo manure by composting or vermicomposting to rehabilitate degraded tropical soils. *Ecological Engineering* 37, 269-276.
- Owojori O.J., Reinecke A.J. and Rozanov A.B. (2010) Influence of clay content on bioavailability of copper in the earthworm *Eisenia fetida*. *Ecotoxicology and Environmental Safety* 73, 407-414.
- Paradelo R., Moldes A.B., González D., Barral M.T. (2012) Plant tests for determining the  
520 suitability of grape marc composts as components of plant growth media. *Waste Management & Research* 30, 1059-1065.
- Rogovska N., Laird D., Cruse R., Fleming P., Parkin T. and Meek D. (2011) Impact of biochar on manure carbon stabilization and greenhouse gas emissions. *Soil Biology and Biochemistry* 75, 871-879.
- 525 Sánchez-García M., Albuquerque J.A., Sánchez-Monedero M.A., Roig A. and Cayuela M.L. (2015) Biochar accelerates organic matter degradation and enhances N mineralisation during

- composting of poultry manure without a relevant impact on gas emissions. *Bioresource Technology* 192, 272-279.
- 530 Shipitalo M.J. and Protz R. (1989) Chemistry and micromorphology of aggregation in earthworm casts. *Geoderma* 45, 357-374.
- Smith P., House J.I., Bustamante M., Sobocka J., Harper R., Pan GG., West P.C., Clark J.M., Adhya T., Rumpel C., Paustian K., Kuikman P., Cotrufo M.F., Elliott J.A., McDowell R., Griffiths R.I., Asakawa S., Bondeau A., Jain A.K., Meersmans J. and Pugh T.A.M. (2015) Global change pressures on soils from land use and management. *Global Change Biology* 22, 1008-1028.
- 535 Thangarajan R., Bolan N.S., Tian G., Naidu R. and Kunhikrishnan A. (2013) Role of organic amendment application on greenhouse gas emission from soil. *Science of the Total Environment* 465, 72-96.
- v.Lützw M., Kögel-Knaber I., Ekschmitt K., Matzner E., Guggenberger G., Marschner B. and Flessa H. (2006) Stabilization of organic matter in temperate soils: mechanisms and their 540 relevance under different soil conditions – a review. *European Journal of Soil Science* 57, 426-445.
- Vu Q.D., de Neergaard A., Tran T. D., Hoang H. T., Vu V.T. and Jensen L.S. (2015) Greenhouse gas emissions from passive composting of manure and digestate with crop residues and biochar on small-scale livestock farms in Vietnam. *Environmental Technology* 36, 2924-2935.
- 545 Wang J., Hu Z., Xu X., Jiang X., Zheng B., Liu X., Pan X. and Kardol P. (2014) Emissions of ammonia and greenhouse gases during combined pre-composting and vermicomposting of duck manure. *Waste management* 34, 1546-1552.
- Weyers L.S. and Spokas A.K. (2011) Impact of biochar on earthworms populations: a review. *Applied and Environmental Soil Science*, Volume 2011, Article ID 541592, 12p.
- 550 Wiedner K., Rumpel C., Steiner C., Pozzi A., Maas R. and Glaser B. (2013) Chemical evaluation of chars produced by thermochemical conversion (gasification, pyrolysis and hydrothermal carbonization) of agro-industrial biomass on a commercial scale. *Biomass and Bioenergy* 59,

264-278.

Zimmerman A.R., Gao B. and Ahn M-Y. (2011) Positive and negative carbon mineralization  
555 priming effects among a variety of biochar-amended soils. *Soil Biology and Biochemistry* 43,  
1169-1179.



560 Table 1: Mean values of pH, DOC, content of total nitrogen and organic carbon after 21 days of co- (vermi)composting. Data are presented as means and standard error (n=4). Different small letters indicate significant differences between treatments (Kruskal-Wallis test,  $p < 0.005$ ).

	pH	C (mg g <sup>-1</sup> )	N (mg g <sup>-1</sup> )	DOC (mg g <sup>-1</sup> TOC)	C/N
Pre-composted material	8.5 ± 0.1 <sup>c</sup>	205.1 ± 3.0 <sup>b</sup>	13.3 ± 0.2 <sup>a</sup>	29.08 ± 0.86 <sup>a</sup>	15.4 ± 0.1 <sup>bc</sup>
<b>Organic materials after 21 days of co-composting</b>					
<b>Compost treatments</b>					
C	8.7 ± 0.1 <sup>ab</sup>	188.2 ± 9.1 <sup>c</sup>	13.5 ± 0.8 <sup>a</sup>	28.85 ± 0.38 <sup>a</sup>	13.5 ± 0.6 <sup>d</sup>
C + 25 % M	8.2 ± 0.1 <sup>d</sup>	153.1 ± 9.5 <sup>d</sup>	10.6 ± 0.5 <sup>c</sup>	21.77 ± 1.57 <sup>b</sup>	14.4 ± 0.8 <sup>d</sup>
C + 50 % M	7.9 ± 0.1 <sup>e</sup>	118.6 ± 2.9 <sup>e</sup>	8.5 ± 0.1 <sup>e</sup>	19.32 ± 0.94 <sup>c</sup>	14.0 ± 0.3 <sup>d</sup>
C + 10 % B	8.7 ± 0.1 <sup>a</sup>	241.9 ± 15.1 <sup>a</sup>	12.4 ± 0.5 <sup>b</sup>	21.26 ± 0.78 <sup>b</sup>	19.5 ± 0.8 <sup>ab</sup>
C + 10 % B+ 25 % M	8.2 ± 0.1 <sup>d</sup>	197.8 ± 5.9 <sup>b</sup>	10.0 ± 0.2 <sup>cd</sup>	15.04 ± 0.68 <sup>e</sup>	19.7 ± 0.3 <sup>a</sup>
<b>Vermicompost treatments</b>					
V	8.6 ± 0.1 <sup>b</sup>	185.0 ± 8.3 <sup>c</sup>	13.0 ± 0.6 <sup>ab</sup>	26.83 ± 0.49 <sup>a</sup>	14.3 ± 0.4 <sup>d</sup>
V + 25 % M	8.2 ± 0.1 <sup>d</sup>	150.2 ± 5.2 <sup>d</sup>	10.4 ± 0.5 <sup>cd</sup>	18.41 ± 0.66 <sup>cd</sup>	14.5 ± 0.3 <sup>d</sup>
V + 50 % M	7.9 ± 0.1 <sup>e</sup>	121.4 ± 6.0 <sup>e</sup>	8.6 ± 0.1 <sup>e</sup>	17.16 ± 0.7 <sup>d</sup>	14.1 ± 0.7 <sup>d</sup>
V + 10 % B	8.7 ± 0.1 <sup>ab</sup>	247.6 ± 12.3 <sup>a</sup>	12.5 ± 0.5 <sup>b</sup>	19.68 ± 0.49 <sup>bc</sup>	19.9 ± 0.9 <sup>a</sup>
V + 10 % B+ 25 % M	8.3 ± 0.1 <sup>d</sup>	206.0 ± 11.4 <sup>b</sup>	9.9 ± 0.3 <sup>d</sup>	15.18 ± 0.43 <sup>e</sup>	20.8 ± 1.4 <sup>a</sup>

Table 2: Effect of the addition of clay and/ or biochar on the rate constant  $k$  (day<sup>-1</sup>) during composting and vermicomposting.

	$k$ (10 <sup>-3</sup> day <sup>-1</sup> )	Std. Error (10 <sup>-5</sup> )
<b>Compost treatments</b>		
C	3.069 <sup>a</sup>	4.429
C + 25 % M	2.588 <sup>cd</sup>	4.539
C + 50 % M	1.699 <sup>g</sup>	2.776
C + 10 % B	2.313 <sup>ef</sup>	2.204
C + 10 % B+ 25 % M	1.762 <sup>g</sup>	5.265
<b>Vermicompost treatments</b>		
V	3.036 <sup>ab</sup>	4.089
V + 25 % M	1.973 <sup>fg</sup>	3.783
V + 50 % M	2.431 <sup>de</sup>	3.616
V + 10 % B	2.855 <sup>ab</sup>	4.869
V + 10 % B+ 25 % M	2.798 <sup>bc</sup>	4.251

565 Table 3: Carbon balance. Data are presented as means and standard error (n=4).

	<i>Composting phase</i> (mgC g <sup>-1</sup> TOC)	<i>Soil incubation phase</i> (mgC g <sup>-1</sup> TOC)	<i>Total carbon mineralized</i> (mgC g <sup>-1</sup> TOC)
<b>Compost treatments</b>			
C	17.11 <sup>a</sup>	18.20 <sup>a</sup>	35.31 <sup>a</sup>
C + 25 % M	13.55 <sup>b</sup>	15.68 <sup>ab</sup>	29.23 <sup>a</sup>
C + 50 % M	7.83 <sup>bc</sup>	14.03 <sup>bc</sup>	21.87 <sup>de</sup>
C + 10 % B	8.67 <sup>de</sup>	8.95 <sup>f</sup>	17.62 <sup>f</sup>
C + 10 % B+ 25 % M	6.36 <sup>e</sup>	13.58 <sup>c</sup>	19.94 <sup>ef</sup>
<b>Vermicompost treatments</b>			
V	15.75 <sup>a</sup>	13.11 <sup>cd</sup>	28.87 <sup>ab</sup>
V + 25 % M	10.59 <sup>c</sup>	13.72 <sup>c</sup>	24.31 <sup>cd</sup>
V + 50 % M	12.23 <sup>bc</sup>	13.73 <sup>c</sup>	25.96 <sup>bc</sup>
V + 10 % B	8.81 <sup>d</sup>	11.42 <sup>ef</sup>	20.22 <sup>ef</sup>
V + 10 % B+ 25 % M	10.59 <sup>c</sup>	12.67 <sup>de</sup>	23.27 <sup>cd</sup>






Figure 1. Experimental design to compare CO<sub>2</sub> emissions of different organic materials during composting and after their addition to soil.



570

Figure 2. Cumulative CO<sub>2</sub> emissions at day 21 from composts and vermicomposts. Different letters

(a, b, c, d, e and f) indicate statistically significant differences.



575 Figure 3: Cumulative CO<sub>2</sub> emissions during composting without worms of pre-composted material alone (C), with 25% of clay (C+25% M), with 50% of clay (C+ 50% M), with 10% of biochar (C+ 10%B) and, with 25% of clay and 10% of biochar (C+25%M + 10% B).





Figure 4: Cumulative CO<sub>2</sub> emissions during composting with worms of pre-composted material alone (V), with 25% of clay (V+25% M), with 50% of clay (V+ 50% M), with 10% of biochar (V+ 10%B) and, with 25% of clay and 10% of biochar (V+25%M + 10% B).



585 Figure 5. Cumulative CO<sub>2</sub> emissions at day 30 from composts and vermicomposts co-composting with 25% of clay (25% M), with 50% of clay (50% M), with 10% of biochar (10% B) and, with 25% of clay and 10% of biochar (25% M + 10% B) in soil. Different letters (a, b, c, d, e and f) indicate statistically significant differences.

590




Figure 6. Comparison between cumulative CO<sub>2</sub> emissions at day 30 from composts and vermicomposts in soil and DOC from these amendments co-composting with 25% of clay (25% M), with 50% of clay (50% M), with 10% of biochar (10% B) and, with 25% of clay and 10% of biochar (25% M + 10% B).