

1 **Thermal alteration of soil physico-chemical properties: A**  
2 **systematic study to infer response of Sierra Nevada**  
3 **climosequence soils to forest fires.**

4  
5 **Samuel N. Araya<sup>1</sup>, S. Mercer Meding<sup>2</sup> and Asmeret Asefaw Berhe<sup>1,3</sup>**

6 [1]Environmental Systems Graduate Group, University of California, Merced

7 [2]Soil, Water, and Environmental Science Dept., University of Arizona, Tucson

8 [3]Life and Environmental Sciences Unit, University of California, Merced

9 Corresponding author: S. N. Araya (e-mail: saraya@ucmerced.edu)

10  
11 **Abstract**

12 Fire is a common ecosystem perturbation that affects many soil properties. As global fire  
13 regimes continue to change with climate change, we investigated thermal alteration of soils'  
14 physical and chemical properties after they are exposed to a range of temperatures that are  
15 expected during prescribed and wildland fires. For this study, we used topsoils collected from  
16 a climosequence transect along the Western slope of the Sierra Nevada that spans from 210 to  
17 2865 m.a.s.l. All the soils we studied were formed on a granitic parent material and had  
18 significant differences in soil organic matter (SOM) concentration and mineralogy owing to the  
19 effects of climate on soil development. Topsoils (0 - 5 cm depth) from the Sierra Nevada  
20 climosequence were heated in a muffle furnace at six set temperatures that cover the range of  
21 major fire intensity classes (150, 250, 350, 450, 550 and 650 °C). We determined the effects of  
22 heating temperature on soil aggregate strength, aggregate size distribution, specific surface area  
23 (SSA), mineralogy, pH, cation exchange capacity (CEC), and carbon (C) and nitrogen (N)  
24 concentrations. With increasing temperature, we found significant reduction of total C, N and  
25 CEC. Aggregate strength also decreased with further implications for loss of C protected inside  
26 aggregates. Soil pH and SSA increased with temperature. Most of the statistically significant  
27 changes ( $p < 0.05$ ) occurred between 350 and 450 °C. We observed relatively smaller changes

1 at temperature ranges below 250 °C. This study identifies critical temperature thresholds for  
2 significant physico-chemical changes in soils that developed under different climate regimes.  
3 Our findings will be of interest to studies of inferences for how soils are likely to respond to  
4 different fire intensities under anticipated climate change scenarios.

## 5 **Keywords**

6 Chemical properties, Climate change, Climosequence, Fire intensity, Physical properties, Soil  
7 heating

8

## 9 **1 Introduction**

10 Fire is a common, widespread phenomenon in many ecosystems around the world (Bowman et  
11 al., 2009). Vegetation fires burn an estimated 300 to 400 million hectares of land globally every  
12 year (FAO, 2005). In the US alone, over 80,000 fires were reported in 2014—including about  
13 63,000 wildland fires, and 17,000 prescribed burns that burned over 1.5 million and 970,000 ha  
14 of land, respectively (National Interagency Fire Center, 2015). Climate and climatic variations  
15 exert strong control on the distribution, frequency, and severity of fires (Harrison et al., 2010).  
16 Significant changes in global fire regimes are anticipated because of climate change including  
17 an increase in frequency of fires in the coming decades (Pechony and Shindell, 2010; Westerling  
18 et al., 2006). However, our understanding of how climate change and changes in fire regimes  
19 will interact to influence topsoils' in fire affected ecosystems is limited.

20 Even though humans are responsible for causing a substantial proportion of vegetation fires  
21 (Caldararo, 2002), vegetation fires are also natural phenomena with an important role in  
22 maintaining the health of many ecosystems around the world (Harrison et al., 2010). In the  
23 Sierra Nevada, vegetation fires have a major influence on the landscapes (McKelvey et al.,  
24 1996).

25 In addition to alteration of vegetation, fire also significantly affects the physical, chemical and  
26 biological properties of soils (Certini, 2005; González-Pérez et al., 2004; Mataix-Solera et al.,  
27 2011). The degree of alteration caused by fires depends on the fire intensity and duration, which  
28 in turn depend on factors such as the amount and type of fuels, properties of above ground  
29 biomass, air temperature and humidity, wind, topography, and soil properties such as moisture

1 content, texture and soil organic matter (SOM) content (DeBano et al., 1998). The first-order  
2 effects of fire on soil are caused by the input of heat causing extreme soil temperatures in topsoil  
3 (Badía and Martí, 2003b; Neary et al., 1999) resulting in loss and transformation of SOM,  
4 changes in soil hydrophobicity, changes in soil aggregation, loss of soil mass, and addition of  
5 charred material and other combustion products (Albalasmeh et al., 2013; Mataix-Solera et al.,  
6 2011; Rein et al., 2008). Soil temperature thresholds for some important soil transformations  
7 are illustrated in Figure 1.

8 The duration of burning impacts soil properties because it determines the amount of energy  
9 transferred through the soil. Fires with longer durations typically have greater impact on soil  
10 properties and SOM than higher temperature fires if they are fast-moving (Frandsen and Ryan,  
11 1986; González-Pérez et al., 2004). Fires also impact soil by altering and removing above-  
12 ground vegetation and topsoil biomass, and increasing soil erodibility (Carroll et al., 2007;  
13 DeBano, 1991), subsequently leading to a shift in plant and microbial populations (Janzen and  
14 Tobin-Janzen, 2008).

15 The aim of this study is to investigate effects of heating temperatures on important soil  
16 properties. Here we aim to determine how the same input of energy from fires affects topsoils  
17 that vary significantly based on carbon content, mineralogy, and associated soil physical and  
18 chemical properties. The inferences derived from this work are essential for determining how  
19 changing climate regimes (and associated changes in vegetation dynamics and soil properties)  
20 are likely to influence the response of topsoil to wild- and prescribed-fires. We use a laboratory  
21 heating experiment on soils from a well-characterized climosequence in the western Sierra  
22 Nevada mountain range to determine: (1) magnitudes of change in soil physico-chemical  
23 properties associated with different fire heating temperatures; (2) identify critical thresholds for  
24 major changes in soil-physico-chemical properties for soils that significantly vary based on  
25 organic matter properties, texture, mineralogy, and other properties.; and (3) infer the  
26 implications of changing climate on topsoil physico-chemical properties that might experience  
27 changing fire regime. This study aims to contribute to the systematic evaluation and  
28 development of ability to predict the effect of different intensity fires on soil properties under  
29 changing climate scenarios.

## 1   **2   Materials and methods**

### 2   **2.1   Study site and soil description**

3   For this study, we collected soils from five sites across an elevation transect along the western  
4   slope of the central Sierra Nevada (Figure 2); the sites were previously characterized by  
5   Dahlgren et al. (1997). We selected four forested sites that are likely to experience forest fires  
6   and a fifth lower elevation grassland site for comparison.

7   All the sites have a Mediterranean climate characterized by warm to hot dry summers and cool  
8   to cold wet winters. Mean annual air temperature ranges from 16.7 °C at the lowest site located  
9   at 210 m to 3.9 °C at the highest elevation site which is at an elevation of 2865 m. Annual  
10   precipitation ranges from 33 cm at the lowest site to 127 cm at the highest site (Dahlgren et al.,  
11   1997; Rasmussen et al., 2007) (Table 1).

12   The lower elevation woodlands of Sierra Nevada experience less frequent fires than further  
13   upslope and the fires are often fast moving and lower severity (Skinner and Chang, 1996). At  
14   the middle-elevation zone of Sierran forest, the mixed conifer zones, frequent fires are low to  
15   moderate severity at lower altitudes but fire frequency generally increases with altitude towards  
16   the upper elevation of the mixed conifer forest (Caprio and Swetnam, 1995). Fires are infrequent  
17   and low severity within the high altitude, Subalpine, zone of Sierra (Skinner and Chang, 1996).

18   Soils from the lowest elevation site, Vista soils (210 masl), fall within the oak woodland zone  
19   (elevations < 1008 m). This is the only soil in our study that does not have an O-horizon, the  
20   soil has dense annual grass cover, however, and the A-horizon SOM originates mainly from  
21   root turnover. Musick soils (1384 masl) lie within oak/mixed-conifer forest (1008—1580 masl)  
22   and mixed-conifer forest (1580—2626 masl). These soils receive the highest biomass and litter  
23   fall. Shaver and Sirretta soils (2317 masl) fall within the mixed-conifer forest range zone while  
24   Chiquito soils (2865 masl) lies within the subalpine mixed-conifer forest range (2626—3200  
25   masl). These soils have lower biomass and litter fall compared to the lower elevation soils (van  
26   Wagtendonk and Fites-Kaufman, 2006).

27   The western slope of central Sierra Nevada presents a remarkable climosequence of soils that  
28   developed under similar granitic parent material and are located in landscapes of similar age,  
29   relief, slope and aspect (Trumbore et al., 1996) with significant developmental differences

1 attributed to climate. The soils at mid-elevation range (1000 to 2000 masl) tend to be highly  
2 weathered while soils at high and low elevations are relatively less developed (Dahlgren et al.,  
3 1997; Harradine and Jenny, 1958; Huntington, 1954; Jenny et al., 1949). Among the most  
4 important changes in soil properties along the climosequence include changes in soil organic  
5 carbon (SOC) concentration, base saturation, and mineral desilication and hydroxyl-Al  
6 interlayering of 2:1 layer silicates. Soil pH generally decreases with elevation and the  
7 concentrations of clay and secondary iron oxides show a step change at the elevation of present-  
8 day average effective winter snowline, i.e. 1600 m elevation (Tables 1 and 2) (California  
9 Department of Water Resources, 1952-1962; Dahlgren et al., 1997).

## 10 **2.2 Experimental design and sample collection**

11 To investigate the effect of heating temperature on physico-chemical properties of soils with  
12 significantly different carbon contents, mineralogy, and overall development, we collected top  
13 soils (0 to 5 cm depth) from five sites. Triplicate samples, approximately 10 m apart, were  
14 collected from each site. The soils were air-dried at room temperature and passed through 2 mm  
15 sieve. Prior to furnace heating, the soils were oven dried at 60 °C overnight. Soil bulk density  
16 and field soil moisture were determined from separate undisturbed core samples collected from  
17 each site (Table 2).

18 Sub-samples from each soil were heated in muffle furnace to one of six selected maximum  
19 temperatures (150, 250, 350, 450, 550 and 650 °C). To ensure uniform soil heating and reduce  
20 formation of heating gradient inside, the soils were packed 1 cm high in a 7 cm diameter  
21 porcelain flat capsule crucibles. Oxygen supply during the heating was only limited by the  
22 availability of space, the furnace was not sealed and the volume of soil sample to volume air in  
23 furnace was approximately 1:50. Furnace temperature was ramped a rate of 3 °C min<sup>-1</sup> and soils  
24 were exposed to the maximum temperature for 30 minutes. Once cooled to touch, soils were  
25 stored in air-tight polyethylene bags prior to analysis.

26 The six heating temperatures were selected to correspond with fire intensity categories that are  
27 based on maximum surface temperature (DeBano et al., 1977; Janzen and Tobin-Janzen, 2008;  
28 Neary et al., 1999), that is, low intensity (150 and 250 °C), medium intensity (350 and 450 °C),  
29 and high intensity (550 and 650 °C). These fire intensity classes generally correspond with

1 thresholds for important thermal reactions in soils observed by differential thermal analyses  
2 (Giovannini et al., 1988; Soto et al., 1991; Varela et al., 2010). Heating rate of  $3\text{ }^{\circ}\text{C min}^{-1}$  is  
3 preferred in laboratory fire simulation experiments (Giovannini et al., 1988; Terefe et al., 2008;  
4 Varela et al., 2010), the slow heating rate prevents sudden combustion when soil's ignition  
5 temperature is reached at about  $220\text{ }^{\circ}\text{C}$  (Fernández et al., 1997, 2001; Varela et al., 2010). The  
6 samples were exposed to the maximum set temperature for a period of 30 minutes. This length  
7 of time ensures that the entire sample is uniformly heated at the set temperature and is in keeping  
8 with wide majority of similar laboratory soil heating experiments (for example Badía and Martí,  
9 2003a; Fernández et al., 2001; Giovannini, 1994; Varela et al., 2010; Zavala et al., 2010). The  
10 duration of soil heating under vegetation fires is highly varied and not uniform across landscape  
11 (Parsons et al., 2010). The same heating procedure was used for all the soils so that it would be  
12 possible to compare how the soils from different climate regimes are likely to respond to the  
13 fires.

### 14 **2.3 Laboratory analysis**

15 Soil color was measured using the Munsell Color Charts. Dry color was measured from air-  
16 dried samples and moisture was added to same sample for moist color measurement. Dry-  
17 aggregate size distribution was measured by sieving. Samples were dry sieved into three  
18 aggregate size classes: 2–0.25 mm (macro-aggregates), 0.25–0.053 mm (micro-aggregates) and  
19  $<0.053$  mm (silt and clay sized particles). These aggregate size classes were selected to enable  
20 comparison with other studies that investigated the effect of different natural and anthropogenic  
21 properties on soil aggregate dynamics and aggregate protected organic matter (Six et al 2000).

22 Water-stable aggregate percent was measured by wet-sieving methods of Nimmo and Perkins  
23 (2002). For this procedure, a wet-sieving apparatus with 0.25mm mesh size sieve is used  
24 (Eijkelkamp Agrisearch Equipment, Giesbeek, The Netherlands). Four grams of soil sample  
25 was placed into a sieve and pre-wetted slowly from a moist tissue paper placed beneath the  
26 sieve. The soil sample was then wet-sieved with an up-down motion at a rate of 35 cycles per  
27 minute and a vertical distance of 1.3 cm. Mass of the soil passing through the sieve was recorded  
28 ( $M_I$ ) after evaporating the supernatant water in oven. The remaining soil sample was subjected  
29 to a second round of wet-sieving using a dispersing solution ( $2\text{ gL}^{-1}$  of sodium  
30 hexametaphosphate for the soils with  $\text{pH} >7$  and  $2\text{ gL}^{-1}$  NaOH for the soils with  $\text{pH} <7$ ) until

1 all particles smaller than the sieve screen opening pass. Mass of the soil that passed through the  
2 sieve in the second round of sieving ( $M_2$ ) was also determined by evaporating supernatant  
3 solution in oven and subtracting the weight of the dispersing-agent. The water-stable aggregate  
4 (WSA) fraction was calculated as:

$$5 \quad WSA = \frac{M_2}{M_1 + M_2} \times 100\%.$$

6 Specific surface area was measured using automated N<sub>2</sub>-BET analyzer (Micromeritics Tri-Star  
7 3000, Micromeritics Instrument Corporation, Norcross, GA, USA). For this procedure,  
8 approximately 1 g of soil was oven dried at 60 °C for 36 hours and out-gassed for another 30  
9 minutes using a flow of N<sub>2</sub> gas with outgassing station mantle set to a temperature of 105 °C.  
10 Measurement was done using ultra-high purity N<sub>2</sub> gas and the instrument was set to use seven-  
11 point N<sub>2</sub>-BET adsorption isotherm measurement from which the Micromeritics software  
12 calculated specific surface area.

13 Soil mineralogy was measured using X-ray diffraction analysis (XRD) using PANalytical Xpert  
14 Pro diffractometer (PANalytical Inc., Westborough, MA, USA). We used the PANalytical  
15 Xpert Pro software for identification of mineral phases and Rietveld refinement for  
16 quantification (Rietveld, 1969; Schulze and Dixon, 2002). Soil samples were ground to a fine  
17 powder consistency using a ball-mill (8000M MiXer/Mill, with 65 ml stainless steel grinding  
18 vial set, SPEX SamplePrep, LLC, Metuchen, NJ, USA) and oven dried at 60 °C for over 36  
19 hours. Samples were scanned at generator setting of 45 mA by 40 kV. Scan start position was  
20 set to 5° 2θ and end position was set to 120 2θ. Scan step time was set to 10 seconds with a step  
21 interval size of 0.0170° 2θ. At least two replicate measurements were run for each sample and  
22 samples were measured in random order.

23 Soil pH was measured using 1:2 solid:solution ratio mixtures in a deionized water and 0.01 M  
24 CaCl<sub>2</sub> solution. Five grams of soil was mixed by shaking with 10 ml of solution and allowed to  
25 stand for 30 minutes, stirring every 10 minutes. The pH reading was taken by placing electrodes  
26 directly in the sediment slurry immediately after stirring (Thomas, 1996).

27 Cation exchange capacity (CEC) was measured by the barium exchange method. Barium was  
28 used to quantitatively displace soil exchangeable cations, and excess barium was removed by  
29 four deionized water rinses. A known quantity of calcium is then exchanged for barium and

1 excess solution calcium is measured in order to determine CEC by the difference in the quantity  
2 of the calcium added and the amount left in the resulting solution. The method has a detection  
3 limit of 2.0 cmol<sub>c</sub>/kg (Rible and Quick, 1960).

4 Elemental concentrations of carbon (C) and nitrogen (N) were measured using an elemental  
5 combustion system (Costech ECS 4010 CHNSO Analyzer, Costech Analytical Technologies,  
6 Valencia, CA, USA) that is interfaced with a mass spectrometer (DELTA V Plus Isotope Ratio  
7 Mass Spectrometer, Thermo Fisher Scientific, Inc, Waltham, MA, USA). For the analyses, the  
8 soil samples were ground to a powder consistency using a ball-mill (8000M MiXer/Mill, with  
9 a 55 ml tungsten Carbide Vial, SPEX SamplePrep, LLC, Metuchen, NJ, USA) and oven-dried  
10 at 60 °C for over 36 hours. This lower temperature and longer duration oven-drying was used  
11 to avoid possible heating related C or N changes that might occur if drying was done 105 °C  
12 (Kaiser et al., 2015). The C and N concentration results were corrected for moisture by oven-  
13 drying subsamples at 105 °C overnight and to determine moisture content. The C and N  
14 concentration results were corrected for moisture by and adjusting for moisture as:  $W_{adj} =$   
15  $W \times (100 - W_m)$ . Where  $W_{adj}$  is the adjusted percent concentration,  $W$  is the concentration  
16 before moisture adjustment and  $W_m$  is the percent moisture content. All concentration changes  
17 resulting from moisture adjustment were a decrease of less than 1% of the value.

## 18 **2.4 Statistical Analysis**

19 All quantitative results are expressed as means of three replicates  $\pm$  standard error, unless  
20 otherwise indicated. Differences of means were tested by Analysis of Variance (ANOVA) and  
21 pairwise comparison of treatments done using Tukey's HSD test at  $p < 0.05$  significance level.  
22 The normality of the data and the homogeneity of variances was checked using Shapiro-Wilk's  
23 and Levene's tests respectively. All statistical analysis were performed using R statistical  
24 software (R Core Team, 2014). ANOVA was implemented using `aov()` function and Tukey's  
25 HSD test subsequently implemented on the `aov` fit by `TukeyHSD()` function. The ANOVA  
26 tables and Tukey's HSD comparison of means tables are given in supplementary material. The  
27 ordinary linear regression technique was used to examine relationships between soil properties.



## 1 **3 Results**

### 2 **3.1 Soil color**

3 We observed a marked soil color change, as inferred using the Munsell color system, with  
4 increasing heating temperature (Figure 3). As the heating temperatures increased, the soils  
5 initially got darker, reaching their darkest color at mid temperatures (250 - 350 °C when  
6 measured in dry soil, and 250 - 450 °C when soils were moist). At higher temperatures, the soils  
7 became markedly lighter and became increasingly reddish in color (with hue changing from  
8 10YR to 7.5YR at temperatures above 550 °C). Color change patterns were similar for both,  
9 dry and moist soil measurements except for the marked color change occurring at 450 °C when  
10 dry and at 550 °C when moist. Across the heating temperature range, Vista (210 masl) soils  
11 showed the least pronounced increase in darkness at 350 °C while Shaver (1737 masl) soils  
12 showed the most pronounced darkening at that temperature range (from dry color of 10YR 5/2  
13 unburned to 10YR 3/3 by 350 °C). At higher temperatures, Musick soils (1384 masl) showed  
14 the largest change in dry soil color going from 10YR 2/2 at 350 °C to 7.5YR 6/6 at 650 °C.

### 15 **3.2 Mass loss**

16 Mass loss was proportional to heating temperature in all the soils. For the high and low elevation  
17 soils, statistically significant mass loss, compared to unburned soils, was observed above 350  
18 °C. In contrast, significant mass loss was observed for the two mid elevation soils of Musick  
19 (1384 masl) and Shaver (1737 masl) starting 250 °C. There was no significant mass loss at  
20 temperatures above 450 °C for all soils. For all our soils, the steepest mass loss was observed  
21 between temperatures of 250 and 450 °C (Figure 4). Vista (210 masl) soils showed the lowest  
22 mass loss with heating while Musick soils (1384 masl) showed the highest mass loss with  
23 heating.

### 24 **3.3 Aggregate stability and size distribution**

25 Aggregate stability generally decreased with temperature for all soils. While aggregate stability  
26 decreased in an almost uniform manner with increase in temperature for the lower to mid  
27 elevation soils, the higher elevation Sirretta (2317 masl) and Chiquito (2865 masl) soils showed  
28 a stepwise decrease in aggregate stability at 250 °C and 350 °C respectively. At higher

1 temperature heating aggregate stability for the two soils showed only a small decrease from  
2 these two temperatures. Statistically significant decrease in aggregate strength, compared to  
3 unburned samples, was observed only at higher temperatures above 350 °C for Sirretta (2317  
4 masl) and Chiquito (2865 masl) soils, and above 450 °C for Musick (1384 masl), Vista (210  
5 masl) and Shaver (1737 masl) soils (Figure 4).

6 Although not statistically significant, all soils showed a decrease in macro-aggregate fraction  
7 accompanied by increase in micro-aggregate and silt-clay sized fractions (Figure 5). For the  
8 two lower elevation soils (Vista and Musick) the decrease in macro-aggregate fraction was over  
9 10% and less than 5% for all the other soils. Only Musick (1384 masl) soils showed a  
10 statistically significant decrease in macro-aggregate fraction between 150 and 350 °C  
11 temperatures.

### 12 **3.4 Specific surface area**

13 For all soils, we observed a stepwise increase in specific surface area (SSA) for samples were  
14 heated to between 250 to 450 °C (Figure 4). Changes in SSA between soils heated below 250  
15 °C and those heated above 450 °C were statistically significant at  $p < 0.05$  for all soils, except  
16 the high elevation soils Sirretta (2317 masl) and Chiquito (2865 masl). Sirretta soils showed a  
17 lot of variability and did not show any significant change in SSA throughout the temperature  
18 range while the Chiquito soil showed statistically significant increase between low temperature  
19 150 – 250 °C and higher temperature 350 – 550 °C range. The pattern of change in SSA with  
20 temperature was similar for all soils. The lowest SSA was recorded for all soils when soils were  
21 heated at 250 °C, and highest SSA was observed at 350 °C (for Musick and Chiquito soils) or  
22 450 °C (Vista and Shaver soils).

### 23 **3.5 Soil mineralogy**

24 The bulk soil XRD results of changes in soil mineralogy in response to heating are presented  
25 for basic mineral groups as: feldspar (microcline and orthoclase); plagioclase (albite and  
26 oligoclase); amphibole; mica/illite (biotite); kaolinite; gibbsite; and expandable phyllosilicate  
27 (montmorillonite and vermiculite). We identified vermiculite with low confidence, since we did  
28 not correct with oriented clay treatments, hence it is not certain if the identified peaks are indeed

1 representative of vermiculite, chlorite, or both. The XRD diagrams showed some significant  
2 transformations in soil mineralogy with heating, with shifted peaks at higher temperatures  
3 suggesting transformation of clay minerals. Layer silicates appeared to collapse structurally,  
4 possibly due to dehydration and the removal –OH (Figure 6). Summary mineral composition  
5 changes identified from XRD analysis using Rietveld method are presented in Figure 7. From  
6 the minerals identified, the concentration of kaolinite was the most affected by heating. In all  
7 soils, concentration of kaolinite showed a decrease when heated to 550 °C and above. Gibbsite  
8 was also not be detected in soils heated to > 450 °C. Furthermore, mica/illite, plagioclases and  
9 amphibole mineral groups changed consistently with increasing temperature. The largest  
10 change in soil mineralogy with heating was observed for the mid-elevation Musick (1384 masl)  
11 and Shaver (1737 masl) soils that are also among the most developed soils with highest  
12 proportions of the 1:1 clay minerals (kaolinite).

### 13 **3.6 Soil pH**

14 With increase in temperature, all soils showed a similar pattern of increase in pH (Figure 8).  
15 For all soils the largest increase in pH (2.5 – 5 units) occurred between 250 and 450 °C. All the  
16 soils started out with slightly to moderately acidic pH and with the exception of Chiquito (2865  
17 soils. Yet, all soils became alkaline at temperatures above 450 °C. The largest increase in pH  
18 was observed for the Musick (1384 masl) soils which reached a pH of 10 at temperatures above  
19 550 °C.

### 20 **3.7 Cation exchange capacity**

21 The CEC of studied soils ranged from an average of 6 cmol<sub>c</sub>/kg for Chiquito (2865 masl) soils  
22 to 25 cmol<sub>c</sub>/kg for Musick (1384 masl) soils. With increase of heating temperature all soils  
23 showed continued decrease of CEC. With the exception of Musick (1384 masl) soils, CEC  
24 eventually dropped to below our detection limit (2 cmol<sub>c</sub>/kg) at temperature above 550 °C  
25 (Figure 8). For the poorly weathered Chiquito (2865 masl) soils, CEC was below 2 cmol<sub>c</sub>/kg at  
26 temperatures 250 °C and above. For the rest of the soils, statistically significant changes in CEC  
27 ( $p < 0.05$ ) occurred at 450 °C with the exception of Musick (1384 masl) soils which showed  
28 statistically significant drop at 250 °C and again 350 °C. At 350 °C, all the soils except Musick

1 (1384 masl) showed a slightly higher CEC than at 250 °C thus interrupting continuous pattern  
2 of CEC decrease with increase in temperature.

### 3 **3.8 Carbon and nitrogen concentration**

4 The initial concentration of C ranged from 1.5% (Vista soil, 210 masl) to 7.7 % (Musick soils,  
5 1384 masl). Soil C concentration decreased with increase in temperature (Figure 9) with the  
6 largest decrease occurring between temperatures of 250 and 450 °C. At 450 °C, all soils had  
7 lost more than 95% of their initial C. At temperatures above 450 °C, C concentration changes  
8 were small and we did not find statistically significant changes at  $p < 0.05$ . The C:N ratio ranged  
9 from 10 (Vista soils, 210 masl) to 29 (Musick soils, 1384 masl). The C:N ratio decreased with  
10 increase in heating temperature in a similar pattern to the C concentration (Figure 9).

11 The loss of C and N from soils due to heating showed a similar response among all five soils.  
12 After 250 °C, all the soils lost more than 25% of their initial C (except Shaver soils that lost  
13 only about 10%). At 350 °C all soils lost 50 to 70% of C. Heating at 450 °C led to the loss of  
14 more than 95% of their initial C for all soils in this study. Loss of N was lower than that of C.  
15 At temperatures greater than 550 °C there was 5 to 15% of soil N still remaining. Consequently,  
16 we observed a decrease of C:N ratio with increase in heating temperature. All soils continued  
17 to lose about 15% soil N for every 100 °C increase and maintained more than 60% of their N at  
18 heating temperatures up to 350 °C. After heating at 450 °C, all soils lost more than 60% of the  
19 initial soil N and 85% by 550 °C.

## 20 **4 Discussion**

21 The topsoil layer is most affected by extreme temperature during vegetation fires. Our results  
22 show significant changes in soil properties as a result of temperature exposure. Our findings  
23 demonstrate that alterations and loss of SOM in topsoil because of heating, rather than  
24 alterations to soil minerals, was the most important driver for the observed changes in soil  
25 physico-chemical properties. Our XRD analysis shows notable changes in soil mineralogy only  
26 after the soils were heated to about 450 to 550 °C (Figure 7). In upland ecosystems, such as the  
27 Sierra Nevada Mountains, the soils typically have low clay content and low concentration of  
28 secondary minerals (Neary et al., 1999; Ubeda and Outeiro, 2009). In addition, these upland  
29 temperate ecosystems also tend to have relatively high concentration of SOM, including a fairly

1 well developed O-horizon. Consequently, strong relationships are observed between SOM  
2 concentrations and the soils' physical and chemical properties. Simple linear regression  
3 analyses between C concentration changes and other soil physical and chemical changes for our  
4 study soils shows that more than 80% of the variability in mass loss, aggregate strength, SSA,  
5 pH, CEC and N concentrations are associated with changes in C concentration at the different  
6 heating temperatures. Table 3 summarizes the correlation coefficients of soil property changes  
7 with changes in C concentration.

8 The changes in soil color observed were consistent with the charring of SOM which leads to  
9 darkening of the brownish color of the soils. At temperatures over 450 °C, the near complete  
10 removal of SOM by combustion and the addition of ash products likely explains the observed  
11 lighter color of soils. The increase in Munsell chromas and reddening of soils at these high  
12 temperatures has been noted in previous works (e.g. Giovannini et al., 1988; Ketterings and  
13 Bigham, 2000; Ulery and Graham, 1993) and is likely a result and the transformation of iron  
14 oxides that occurs during heating.

15 The extent of mass loss in topsoil layers due to vegetation fires is strongly correlated to SOM  
16 combustion (Rein et al., 2008). In all of our soils, statistically significant mass loss ( $p < 0.05$ )  
17 occurred within the temperature ranges of SOM combustion, between 250 °C to 450 °C.  
18 Proportion of soil mass loss with temperature was proportional to initial C concentration of the  
19 soils. For example, Musick (1384 masl) soils which had the highest initial C concentration  
20 (7.7%) had the steepest soil mass loss and lost 15% of mass at 550 °C, while Vista (210 masl)  
21 which had less than 2% C concentration showed the smallest mass loss losing less than 5%  
22 mass at even at the highest temperature. Mass loss at temperatures below 250 °C is likely driven  
23 by dehydration processes rather than by loss of SOM. On the other hand, mass loss at  
24 temperatures above which SOM combusts,  $>450$  °C, is likely driven by charring, ashing and  
25 volatilization processes.

26 Most studies report significant soil aggregate stability reduction with fire heating (e.g.  
27 Arcenegui et al. (2008); Zavala et al. (2010)), however, contrasting findings are also reported  
28 in other studies. Mataix-Solera et al. (2011) explains that increase in aggregate strength with  
29 heating is possible in clay-rich soils where the main cementing agents are inorganic minerals  
30 such as calcium carbonates and metallic oxides which would fuse under fire heating increasing

1 aggregate stability. Another possibility is, soils which initially had weak hydrophobicity may  
2 show increased aggregate stability due to increase in hydrophobicity, in such cases however  
3 aggregate strength would decline at higher severity fires as hydrophobicity is destroyed.

4 Aggregate stability in all our soils generally decreased with increase in heating temperature.  
5 Although we did not find a statistically significant difference, it is worth noting, that the lowest  
6 elevation soil (Vista, 210 masl) showed a trend of aggregate stability increase up to 350 °C  
7 while the high elevation soils, Sirretta (2317 masl) and Chiquito (2865 masl), showed a slight  
8 increase in aggregate stability at 150 °C. An increase in hydrophobicity at these temperatures is  
9 the most likely cause, substantial hydrophobicity was apparent with Chiquito soils (2865 masl)  
10 heated at 250 °C where resistance to slaking was remarkably evident during the aggregate  
11 stability test.

12 Soil specific surface area (SSA) is an important soil property that affects soil adsorption, ion  
13 exchange capacity, reactivity, aggregation and porosity (Feller et al., 1992). SSA of soil is  
14 largely dictated by clay-size particles and SOM (Carter et al., 1986). The increase in SSA with  
15 heating that we observed in this work is most likely the result of physical disintegration and  
16 charring of SOM, especially at temperatures below 500 °C. Changes in soil mineralogy are not  
17 likely to be responsible for the changes in SSA we observed. XRD analysis showed notable  
18 mineralogical changes only at temperatures above 450 °C where the kaolinite peak disappeared  
19 at 550 °C and the phyllosilicate peaks were diminished or disappeared at 450 – 550 °C (e.g.  
20 Figure 6). Because of combustion, pyrolysis and thermal dehydration, larger organic matter  
21 particles are likely to have fragmented and reduced in size with an increase in heating  
22 temperature leading to increase in bulk soil SSA. Furthermore, thermal alteration and the  
23 removal of organic matter from mineral surfaces may have increased surface area by reducing  
24 the size of particles. At higher temperatures (>450 °C), XRD spectra showed some collapse of  
25 mineral complexes through dehydration and de-hydroxylation of clay minerals that may reduce  
26 mineral particle size and increase surface area. These changes in mineralogy might have a  
27 significant effect on SOM. A study by Rosa et al. (2013) found that soils released more organic  
28 compounds during pyrolysis when the soils were treated with HF acid, suggesting that mineral  
29 complexes play a role in protecting organic compounds from pyrolysis. The collapse of mineral  
30 complexes we observed and the decrease in aggregate strength is likely to have enhanced

1 thermal oxidation of SOM. However mineralogical changes would play a more important role  
2 affecting soil properties at these high temperatures since SOM has been almost completely  
3 removed.

4 Soil pH generally increases with fire heating (Badía and Martí, 2003b; Chandler et al., 1983;  
5 Ubeda and Outeiro, 2009). In a soil heating experiment similar to our study, Fernández et al.  
6 (1997) observed a pH increase of 1.7 at 350 °C and 2.35 at 490 °C. Such increase in soil pH has  
7 been attributed to the denaturation of organic acids, release of base cations from combustion  
8 (K- and Na-hydroxides, Mg- and Ca-carbonates), deposition of ashes, and loss of hydroxyl  
9 groups from clays (Badía and Martí, 2003b; Certini, 2005; Ulery et al., 1993). In our soils, the  
10 higher elevation soils (Shaver, Sirretta and Chiquito) showed a statistically insignificant  
11 decrease of 0.3 to 0.5 pH units (measured in water) at 250 °C. The change in pH in our high  
12 elevation soils was consistent with previous results of Badía and Martí (2003b); Terefe et al.  
13 (2008) that found similar initial decrease. Terefe et al. (2008) hypothesized that this may be due  
14 to the combined effect of desiccation and heating effect which favor proton-reducing oxidation  
15 reactions. And the fact that this initial increase occurs below the temperature for the start of  
16 combustion of organic acids means contribution of SOM to this pH increase (organic acid  
17 denaturation and ash liming effect) was absent at this temperature. In a similar heating  
18 experiment Badía and Martí (2003b) found an increase in electric conductivity and soluble Ca  
19 along with decrease in pH at 250 °C. Such increase in soluble cations might explain our findings  
20 where we observed a decrease in pH when measured in water but not in CaCl<sub>2</sub> suggesting that  
21 the decrease in pH might have to do with increase of soluble salts with heating up to 250 °C.

22 The capacity of soil to exchange positively charged ions between soil and soil solution (CEC)  
23 decreased with increasing temperature. CEC of soils is a result of surface charges associated  
24 with secondary clay minerals and SOM (Sparks, 2005), and in our study soils, Dahlgren et al.  
25 (1997) had previously reported a strong relationship of CEC with soil organic carbon and clay  
26 concentrations. Different authors have attributed loss of CEC during heating mainly with loss  
27 of SOM (Fernández et al., 1997; Ubeda and Outeiro, 2009) partly because CEC loss starts to be  
28 observed at temperatures above 200 °C with little or no decrease at lower temperatures where  
29 SOM is not affected (Nishita and Haug, 1972; Soto and Diazfierros, 1993). The slight increase  
30 of CEC we observed at 350 °C may be due to the steep increase of specific surface area at that

1 temperature (Figure 4). The additional surface for cation adsorption might have to an extent  
2 compensated for the loss of SOM at that temperature. Furthermore, the contribution of surface  
3 oxidation of char products has been shown to increase CEC per unit C (Liang et al., 2006)  
4 because of the almost complete loss of C at temperatures above 450 °C and very little loss at  
5 temperatures below 250 °C. The soils most likely had highest concentration of charred SOM at  
6 350 °C temperature.

#### 7 **4.1 Importance of the 250 – 450 °C range**

8 Taking maximum surface temperature as a metric of fire intensity, low intensity fires reach  
9 surface temperatures of up to 250 °C, medium intensity fires reach surface temperatures of 400  
10 °C, and high intensity fires reach surface temperatures above 675 °C (Janzen and Tobin-Janzen,  
11 2008). In this study, the most significant changes of soil chemical properties occurred between  
12 250 and 450 °C. Figure 10 illustrates the changes between unburned and 650 °C burned soils  
13 and the amount of change that occurred within 250 to 450 °C heating temperature for a range  
14 of the variables discussed above. In all cases, the change in the 250 – 450 °C range accounts for  
15 most of the total change observed during our heating treatments. Among the variables we  
16 investigated in this study, we observed changes along two general lines: (1) mass loss, SSA and  
17 pH which showed a progressive increase with heating temperature, and (2) %C, %N, C:N ratio,  
18 CEC, and wet aggregate stability that showed a progressive decrease with increase in  
19 temperature (Figure 10), with the most significant changes in all cases being recorded in all  
20 soils between 250 – 450 °C.

21 Temperatures below 250 °C are very critical for many processes, water is lost at 95 °C and this  
22 has a significant effect on soil heat conduction and soil biota (Janzen and Tobin-Janzen, 2008).  
23 However, we found that temperatures below 200 °C have very little effect on quality or quantity  
24 of SOM. This implies that lower intensity fires, such as typical prescribed fires, where soil  
25 surface temperatures do not exceed below 250 °C temperature contribute little to soil C loss.  
26 Similarly, temperatures above 500 °C do little to SOM, which already has been lost or  
27 transformed into a pyrogenic product. The effect on soil inorganic particles starts at high  
28 temperature but the significance of change on minerals is not as large (Figure 1). Hence, we  
29 found that the most important soil changes occur the 250 – 450 °C range.



1 Important modifications of fire conditions that still allow for comparison of responses of  
2 different types of soils have to be adopted to conduct the type of heating experiments that we  
3 undertook in this work. A heating rate of 3 °C min<sup>-1</sup> is common in laboratory soil heating  
4 studies, often because of technical considerations and because such a slow rate prevents sudden  
5 combustion which otherwise would happen as soil's ignition temperature is reached at about  
6 220 °C (Fernández et al., 1997, 2001; Varela et al., 2010). However, it is important to recognize  
7 that during vegetation fires the rate of temperature increase experienced by the topmost layer  
8 of soil that is exposed to fire can be significantly higher. The rate of heating alone might have  
9 additional significant effects on soil properties beyond what we observe here. For example,  
10 Albalasmeh et al. (2013) found that slow rate of heating underestimates soil aggregate  
11 destruction of moist soils due to a slower buildup of pore-pressure.

## 12 **4.2 Climate Change Implications**

13 Investigation of the response of climosequence soils to different heating temperature in this  
14 study enables us to infer how changes in climate (and associated changes in soil properties) are  
15 likely to alter the effect of fires on topsoil physical and chemical properties. The overall pattern  
16 of soil property changes with increase in heating temperature were similar for all soils, however,  
17 the magnitudes of change were variable. Hence, these findings lead us to conclude that climate  
18 change is likely to alter the response of topsoil properties to different fire regimes. Along our  
19 study climosequence, we observed critical differences in response of topsoils based mostly on  
20 concentration OM in soil and soil development stages of each soil --both variables that are  
21 expected to respond to changes in climate (Berhe et al., 2012). Consequently, changes in soil C  
22 storage associated with climate change are expected to lead to different amounts of C loss due  
23 to fires. This is evidenced by the observed highest total mass of C loss from the mid-elevation  
24 Musick soil that had the highest carbon stock, compared to soils in either side of that elevation  
25 range. Anticipated changes in climate in the Sierra Nevada mountain ranges are expected to  
26 include upward movement of the rain-snow transition line exposing areas that now receive most  
27 of their precipitation as snow to rainfall and associated runoff (Arnold et al., 2015, 2014; Stacy  
28 et al., 2015). Moving of the rain-snow transition zone higher and promotion of more intense  
29 weathering at higher elevation zones then is likely to render more C to loss during fires. As we  
30 found in this study, more than 80% of the variability in mass loss, aggregate strength, SSA, pH,

1 CEC and N concentrations is associated with changes in C concentration at the different heating  
2 temperatures (Table 3). Improving our understanding of how topsoil properties are likely to  
3 respond to changes in climate becomes even more critical when we recognize that C  
4 concentration in soil is likely to respond quickly to changes in climate, compared to other soil  
5 physical and chemical properties (Berhe et al., 2012). Furthermore, the long-term fate of soil  
6 carbon in fire-affected ecosystems is also likely to be accompanied by changes in microbial  
7 community composition and OM decomposition kinetics (Holden et al., 2015; Tas et al., 2014)  
8 which are likely to have further implications for nutrient availability post-fire (Johnson et al.,  
9 2007b; Johnson et al., 1997).

10 The different responses of soil aggregation in our climosequence to the treatment temperatures  
11 also suggest potential loss and transformation of the physically protected C pool in topsoil.  
12 Degradation of aggregates during fire (Albalasmeh et al., 2013) is likely to render aggregate-  
13 protected C to potential losses through oxidative decomposition, leaching and erosion.  
14 Moreover, in systems such as the Sierra Nevada where steep slopes and organic matter-rich  
15 topsoils dominate, movement of the rain-snow transition zone upward is likely to increase  
16 proportion of precipitation that occurs as rain. The kinetic energy of raindrops and observed  
17 increase in hydrophobicity of soils post-fires (Johnson et al., 2007a; Johnson et al., 2004) can  
18 lead to higher rates of erosional redistribution of especially the free light fraction or particulate  
19 C that is not associated with soil minerals (Stacy et al., 2015). Moreover, the important  
20 differences in changes in pH, mineralogy, CEC in response to heating at different temperatures  
21 that we observed for soils along the climosequence suggest that changes in temperature are  
22 likely to lead to different effects on soil chemical properties in soils after fires.

23 Finally, with changes in climate it is anticipated that fires will increase in severity (Westerling  
24 et al., 2006). Our findings of important changes in soil physical and chemical properties  
25 occurring between 250-450 °C are important for recognizing that critical transformations of  
26 topsoil physical and chemical properties are likely to occur when, as a result of climate change,  
27 systems that are adapted to low severity fires experience medium to high severity fires.

## 28 **5 Conclusion**

29 The findings of this study showed that changes in soil properties during heating are closely  
30 related to changes in C concentrations in soil. The temperatures most critical to C loss and

1 alteration were found to be 250 °C, where charring of organic matter starts and 450 °C where  
2 most of the SOM is combusted. Most soil properties exhibited a steep change in this temperature  
3 range. Soil aggregate stability, CEC, and C and N concentrations significantly decreased with  
4 increase in heating temperature while soil pH and SSA significantly increased. The most  
5 important effect of heating on soil mineralogy as observed by XRD analysis was the  
6 disappearance of kaolinite, which was undetectable at temperatures above 500 °C.

7 This study presented the effects of heat input on topsoil properties. The study is necessary to  
8 understand thermally induced changes on soil properties in isolation from other variables that  
9 accompany vegetation fires such as the addition of pyrolysis products from plants and ash, and  
10 the fire induced soil moisture dynamics. Findings from this study will contribute towards  
11 estimating the amount and rate of change in carbon and nitrogen loss, and other essential soil  
12 properties that can be expected from topsoil exposure to different intensity fires under  
13 anticipated climate change scenarios.

#### 14 **Acknowledgements**

15 The authors would like to thank Prof. Randy A. Dahlgren for providing us with geo-references  
16 for the study sites, background data, and for his comments on an earlier version of this  
17 manuscript. We thank Drs. Marilyn Fogel and Christina Bradley for their help and expertise in  
18 elemental analysis of C and N; and Dr. Samuel Traina for his comments on an earlier version  
19 of this manuscript. The research was funded by a UC Merced Graduate Research Council grant  
20 and NSF grant (EAR-1352627) to AAB.

#### 21 **References**

22 Albalasmeh, A. A., Berli, M., Shafer, D. S., and Ghezzehei, T. A.: Degradation of moist soil  
23 aggregates by rapid temperature rise under low intensity fire, *Plant and Soil*, 362, 335-344,  
24 2013.

25 Arcenegui, V., Mataix-Solera, J., Guerrero, C., Zomoza, R., Matalx-Beneyto, J., and Garcia-  
26 Orenes, F.: Immediate effects of wildfires on water repellency and aggregate stability in  
27 Mediterranean calcareous soils, *Catena*, 74, 219-226, 2008.

1 Arnold, C., Ghezzehei, T. A., and Berhe, A. A.: Decomposition of distinct organic matter pools  
2 is regulated by moisture status in structured wetland soils, *Soil Biology and Biochemistry*, 81,  
3 28-37, 2015.

4 Arnold, C., Ghezzehei, T. A., and Berhe, A. A.: Early spring, severe frost events, and drought  
5 induce rapid carbon loss in high elevation meadows, *PloS one*, 9, e106058, 2014.

6 Badía, D. and Martí, C.: Effect of simulated fire on organic matter and selected microbiological  
7 properties of two contrasting soils, *Arid Land Research and Management*, 17, 55 — 69, 2003a.

8 Badía, D. and Martí, C.: Plant ash and heat intensity effects on chemical and physical properties  
9 of two contrasting soils, *Arid Land Research and Management*, 17, 23-41, 2003b.

10 Berhe, A. A., Suttle, K. B., Burton, S. D., and Banfield, J. F.: Contingency in the Direction and  
11 Mechanics of Soil Organic Matter Responses to Increased Rainfall, *Plant and Soil*, 358, 371-  
12 383. DOI 310.1007/s11104-11012-11156-11100, 2012.

13 Bowman, D. M. J. S., Balch, J. K., Artaxo, P., Bond, W. J., Carlson, J. M., Cochrane, M. A.,  
14 D'Antonio, C. M., DeFries, R. S., Doyle, J. C., Harrison, S. P., Johnston, F. H., Keeley, J. E.,  
15 Krawchuk, M. A., Kull, C. A., Marston, J. B., Moritz, M. A., Prentice, I. C., Roos, C. I., Scott,  
16 A. C., Swetnam, T. W., van der Werf, G. R., and Pyne, S. J.: Fire in the Earth System, *Science*,  
17 324, 481-484, 2009.

18 Caldararo, N.: Human ecological intervention and the role of forest fires in human ecology, *The*  
19 *Science of the Total Environment*, 292, 141-165, 2002.

20 California Department of Water Resources: Water Conditions in California. Bull. 120,  
21 Sacramento, 1952-1962.

1 Caprio, A. C. and Swetnam, T. W.: Historic fire regimes along an elevational gradient on the  
2 west slope of the Sierra Nevada, California, General Technical Report INT-GTR-320, 173-179  
3 pp., 1995.

4 Carroll, E. M., Miller, W. W., Johnson, D. W., Saito, L., Qualls, R. G., and Walker, R. F.:  
5 Spatial analysis of a large magnitude erosion event following a Sierran wildfire, *Journal of*  
6 *environmental quality*, 36, 1105-1111, 2007.

7 Carter, D. L., Mortland, M. M., and Kemper, W. D.: Specific surface area. In: *Methods of Soil*  
8 *Analysis. Part 1, Physical and Mineralogical Methods*, Klute, A. (Ed.), *Agronomy*, No.9, Part  
9 1, American Society of Agronomy/ Soil Science Society, Madison, 1986.

10 Certini, G.: Effects of fire on properties of forest soils: a review, *Oecologia*, 143, 1-10, 2005.

11 Chandler, C., Cheney, P., thomas, P., Trabaud, L., and William, D.: Fire effects on soil, water  
12 and air. In: *Fire in Forestry. Vol I: Forest fire behaviour and effects*, John Wiley Sons, New  
13 York, 1983.

14 Dahlgren, R. A., Boettinger, J. L., Huntington, G. L., and Amundson, R. G.: Soil development  
15 along an elevational transect in the western Sierra Nevada, California, *Geoderma*, 78, 207-236,  
16 1997.

17 DeBano, L. F.: The effect of fire on soil properties, Ogden, UT1991, 151-156.

18 DeBano, L. F.: The role of fire and soil heating on water repellency in wildland environments  
19 a review, *Journal of Hydrology*, 2000. 195-206, 2000.

20 DeBano, L. F., Dunn, P. H., and Conrad, C. E.: Fire's effect on physical and chemical properties  
21 of Chaparral soils, 1977.

- 1 DeBano, L. F., Neary, D. G., and Ffolliott, P. F.: Fire's effect on ecosystems John Wiley &  
2 Sons, Inc., New York, USA, 1998.
- 3 Diaz-Ravina, M., Prieto, A., Acea, M. J., and Carballas, T.: Fumigation extraction method to  
4 estimate microbial biomass in Heated Soils, *Soil Biol Biochem*, 24, 259-264, 1992.
- 5 FAO: State of the world's forest Food and Agricultural Organization of the United Nations,  
6 Rome, Italy, 2005.
- 7 Feller, C., Schouller, E., Thomas, F., Rouiller, J., and Herbillon, A. J.: N<sub>2</sub>-BET Specific surface  
8 areas of some low activity clay soils and their relationships with secondary constituents and  
9 organic matter contents, *Soil Sci*, 153, 1992.
- 10 Fernández, I., Cabaneiro, A., and Carballas, T.: Organic matter changes immediately after a  
11 wildfire in an Atlantic forest soil and comparison with laboratory soil heating, *Soil Biology and*  
12 *Biochemistry*, 29, 1-11, 1997.
- 13 Fernández, I., Cabaneiro, A., and Carballas, T.: Thermal resistance to high temperatures of  
14 different organic fractions from soils under pine forests, *Geoderma*, 104, 281-298, 2001.
- 15 Frandsen, W. H. and Ryan, K. C.: Soil-Moisture Reduces Belowground Heat-Flux and Soil  
16 Temperatures under a Burning Fuel Pile, *Can J Forest Res*, 16, 244-248, 1986.
- 17 Giovannini, G.: The effect of fire on soil quality. In: *Soil Erosion and Degradation as a*  
18 *Consequence of Forest Fires*, Sala, M. and Rubio, J. L. (Eds.), Geoforma Ediciones, Logrono,  
19 1994.
- 20 Giovannini, G., Lucchesi, S., and Giachetti, M.: Effect of heating on some physical and  
21 chemical parameters related to soil aggregation and erodibility, *Soil Sci*, 146, 1988.

1 González-Pérez, J., González-Vila, F., Almendros, G., and Knicker, H.: The effect of fire on  
2 soil organic matter—a review, *Environment international*, 30, 855-870, 2004.

3 Harradine, F. and Jenny, H.: Influence of parent material and climate on texture and nitrogen  
4 and carbon contents of virgin California soils: texture and nitrogen contents of soils, *Soil Sci*,  
5 85, 235-243, 1958.

6 Harrison, S. P., Marlon, J. R., and Bartlein, P. J.: Fire in the Earth System, *Int Year Planet Eart*,  
7 doi: Doi 10.1007/978-90-481-8716-4\_3, 2010. 21-48, 2010.

8 Holden, S. R., Berhe, A. A., and Treseder, K. K.: Decreases in soil moisture and organic matter  
9 quality suppress microbial decomposition following a boreal forest fire, *Soil Biology and*  
10 *Biochemistry*, 87, 1-9, 2015.

11 Huntington, G. L.: The effect of vertical zonality on clay content in residual granitic soils of the  
12 Sierra Nevada mountains, University of California, Berkeley, 1954.

13 Janzen, C. and Tobin-Janzen, T.: Microbial communities in fire-affected soils. In: *Microbiology*  
14 *of Extreme Soils*, Dion, P. and Nautiyal, C. S. (Eds.), *Soil Biology*, 13, Springer-Verlag Berlin  
15 Heidelberg, 2008.

16 Jenny, H., Gessel, S. P., and Bingham, F. T.: Comparative study of decomposition rates of  
17 organic matter in temperate and tropical regions, *Soil Sci*, 68, 419-432, 1949.

18 Johnson, D., Murphy, J., Walker, R., Glass, D., and MILLER, W.: Wildfire effects on forest  
19 carbon and nutrient budgets, *Ecological Engineering*, 31, 183-192, 2007a.

20 Johnson, D., Murphy, J. D., Walker, R. F., Glass, D. W., and Miller, W. W.: Wildfire effects  
21 on forest carbon and nutrient budgets, *Ecological Engineering*, 31, 183-192, 2007b.

1 Johnson, D., Susfalk, R., and Dahlgren, R.: Nutrient fluxes in forests of the eastern Sierra  
2 Nevada mountains, United States of America, *Global Biogeochemical Cycles*, 11, 673-681,  
3 1997.

4 Johnson, D. W., Susfalk, R. B., Caldwell, T. G., Murphy, J. D., Miller, W. W., and Walker, R.  
5 F.: Fire Effects on Carbon and Nitrogen Budgets in Forests, *Water, Air, & Soil Pollution:*  
6 *Focus*, 4, 263-275, 2004.

7 Kaiser, M., Kleber, M., and Berhe, A. A.: How air-drying and rewetting modify soil organic  
8 matter characteristics: An assessment to improve data interpretation and inference, *Soil Biology*  
9 *and Biochemistry*, 80, 324-340, 2015.

10 Ketterings, Q. M. and Bigham, J. M.: Soil color as an indicator of slash-and-burn fire severity  
11 and soil fertility in Sumatra, Indonesia, *Soil Science Society of America Journal*, 64, 1826-  
12 1833, 2000.

13 Knicker, H.: How does fire affect the nature and stability of soil organic nitrogen and carbon?  
14 A review, *Biogeochemistry*, 85, 91-118, 2007.

15 Liang, B., Lehmann, J., Solomon, D., Kinyangi, J., Grossman, J., O'Neill, B., Skjemstad, J. O.,  
16 Thies, J., Luizao, F. J., Petersen, J., and Neves, E. G.: Black carbon increases cation exchange  
17 capacity in soils, *Soil Science Society of America Journal*, 70, 1719-1730, 2006.

18 Massman, W. J., Frank, J. M., and Mooney, S. J.: Advancing investigation and physical  
19 modeling of first-order fire effects on soils, *Fire Ecology*, 6, 36-54, 2010.

20 Mataix-Solera, J., Cerda, A., Arcenegui, V., Jordan, A., and Zavala, L. M.: Fire effects on soil  
21 aggregation: a review, *Earth-Sci Rev*, 109, 44-60, 2011.



1 McKelvey, K. S., Skinner, C. N., Chang, C.-r., Erman, D. C., Husari, S. J., Parsons, D. J.,  
2 Wagtendonk, J. W. v., and Weatherspoon, C. P.: An overview of fire in the Sierra Nevada,  
3 Davis, California 37, 1996.

4 Munsell Color Science Laboratory: Munsell Color and CIExyY data. Laboratory, M. C. S.  
5 (Ed.).

6 National Interagency Fire Center: [https://www.nifc.gov/fireInfo/fireInfo\\_statistics.html](https://www.nifc.gov/fireInfo/fireInfo_statistics.html), last  
7 access: 6/5/2015 2015.

8 Neary, D. G., Klopatek, C. C., DeBano, L. F., and Ffolliott, P. F.: Fire effects on belowground  
9 sustainability a review and synthesis, *Forest Ecology and Management*, 122, 51-71, 1999.

10 Nimmo, J. R. and Perkins, K. S.: Aggregate stability and size distribution. In: *Methods of Soil*  
11 *Analysis. Part 4, Physical Methods*, Dane, J. H. and Topp, G. C. (Eds.), Soil Science Society of  
12 America, Madison, Wisconsin, 2002.

13 Nishita, H. and Haug, R. M.: Some physical and chemical characteristics of heated soils, *Soil*  
14 *Sci*, 113, 422-430, 1972.

15 Parsons, A., Robichaud, P. R., Lewis, S. A., Napper, C., and Clark, J.: Field guide for mapping  
16 post-fire soil burn severity, U.S. Department of Agriculture, Forest Service, Rocky Mountain  
17 Research Station, Fort Collins, CO, 2010.

18 Pechony, O. and Shindell, D. T.: Driving forces of global wildfires over the past millennium  
19 and the forthcoming century, *Proceedings of the National Academy of Sciences*, doi:  
20 10.1073/pnas.1003669107, 2010. 2010.

21 R Core Team: *R: A Language and Environment for Statistical Computing*. R Foundation for  
22 Statistical Computing, Vienna, Austria, 2014.

1 Rasmussen, C., Matsuyama, N., Dahlgren, R. A., Southard, R. J., and Brauer, N.: Soil Genesis  
2 and Mineral Transformation Across an Environmental Gradient on Andesitic Lahar, Soil  
3 Science Society of America Journal, 71, 225-237, 2007.

4 Rein, G., Cleaver, N., Ashton, C., Pironi, P., and Torero, J. L.: The severity of smouldering peat  
5 fires and damage to the forest soil, Catena, 74, 304-309, 2008.

6 Rible, J. M. and Quick, J.: Method S-19.0: Cation Exchange Capacity. In: Water, Soil, Plant  
7 Tissue: Tentative Methods of Analysis for Diagnostic Purposes, University of California  
8 Agricultural Experiment Service, Davis, CA, , 1960.

9 Rietveld, H. M.: A Profile Refinement Method for Nuclear and Magnetic Structures, J Appl  
10 Crystallogr, 2, 65-71, 1969.

11 Rosa, J. M. D. l., Gonzalez-Perez, J. A., Gonzalez-Vila, F. J., and Knicker, H.: Medium term  
12 effects of fire induced soil organic matter alterations on Andosols under Canarian pine (*Pinus*  
13 *canariensis*), Journal of Analytical and Applied Pyrolysis, 104, 269-279, 2013.

14 Rossel, R. A. V., Minasny, B., Roudier, P., and McBratney, A. B.: Colour space models for soil  
15 science, Geoderma, 133, 320-337, 2006.

16 Schulze, D. G. and Dixon, J. B.: An introduction to soil mineralogy. In: Soil mineralogy with  
17 environmental applications, Schulze, D. G. and Dixon, J. B. (Eds.), Soil Science Society of  
18 America, Inc. , Madison, Wisconsin, USA, 2002.

19 Skinner, C. N. and Chang, C.-R.: Fire Regimes, Past and Present. In: Sierra Nevada Ecosystem  
20 Project, Final Report to Congress, Vol II, Assessment and Scientific Basis for management  
21 Options, Davis: University of California, Center for Water and Wildland Resources, 1996.

1 Soto, B., Benito, E., Basanta, R., and Díaz-Fierros, F.: Influence of antecedent soil moisture on  
2 pedological effects of fire. In: Soil Erosion and Degradation as a Consequence of Forest Fires,  
3 Sala, M. and Rubio, J. L. (Eds.), Geoforma Ed., Logroño, Spain, 1991.

4 Soto, B. and Diazfierros, F.: Interactions between plant ash leachates and soil, *Int J Wildland*  
5 *Fire*, 3, 207-216, 1993.

6 Sparks, D. L.: *Environmental Soil Chemistry*, The Netherlands, 2005.

7 Stacy, E., Hart, S. C., Hunsaker, C. T., Johnson, D. W., and Berhe, A. A.: Soil carbon and  
8 nitrogen erosion in forested catchments: implications for erosion-induced terrestrial carbon  
9 sequestration, *Biogeosciences*, 12, 4861-4874, 2015.

10 Tas, N., Prestat, E., McFarland, J. W., Wickland, K. P., Knight, R., Berhe, A. A., Jorgenson, T.,  
11 Waldrop, M. P., and Jansson, J. K.: Impact of fire on active layer and permafrost microbial  
12 communities and metagenomes in an upland Alaskan boreal forest, *The ISME Journal.*, 2014.  
13 1-16, 2014.

14 Terefe, T., Mariscal-Sancho, I., Peregrina, F., and Espejo, R.: Influence of heating on various  
15 properties of six Mediterranean soils: a laboratory study, *Geoderma*, 143, 273-280, 2008.

16 Thomas, G. W.: Soil pH and soil acidity. In: *Methods of Soil Analysis. Part 3, Chemical*  
17 *Methods*, Sparks, D. L., Page, A. L., Helmke, P. A., and Loeppert, R. H. (Eds.), Soil Science  
18 Society of America/ American Society of Agronomy, Madison, Wisconsin, 1996.

19 Trumbore, S., Chadwick, O., and Amundson, R.: Rapid exchange between soil carbon and  
20 atmospheric carbon dioxide driven by temperature change, *Science*, 272, 393-396, 1996.

21 U.S. Geological Survey: *USGS NED and 1:1,000,000-Scale Contours*. USGS (Ed.), Reston,  
22 VA, 2015.

1 Ubeda, X. and Outeiro, L. R.: Physical and Chemical Effects of Fire on Soil. In: Fire Effects on  
2 Soils and Restoration Strategies, Cerda, A. and Robichaud, P. R. (Eds.), Land Reconstruction  
3 and Management, Science Publishers, Enfield, NH, USA, 2009.

4 UC Davis Soil Resource Laboratory: [http://casoilresource.lawr.ucdavis.edu/software/r-  
5 advanced-statistical-package/color-functions/convert-munsell-colors-computer-friendly-rgb-  
6 triplets/2015](http://casoilresource.lawr.ucdavis.edu/software/r-advanced-statistical-package/color-functions/convert-munsell-colors-computer-friendly-rgb-triplets/2015).

7 Ulery, A. L. and Graham, R. C.: Forest-fire effects on soil color and texture, Soil Science  
8 Society of America Journal, 57, 135-140, 1993.

9 Ulery, A. L., Graham, R. C., and Amrhein, C.: Wood-Ash Composition and Soil-pH Following  
10 Intense Burning, Soil Sci, 156, 358-364, 1993.

11 van Wagendonk, J. W. and Fites-Kaufman, J. A.: Sierra Nevada Bioregion. In: Fire in  
12 California's Ecosystems, Sugihara, N. G., van Wagendonk, J. W., Shaffer, K. E., and Thode,  
13 A. E. (Eds.), University of California Press, Berkeley, CA, USA, 2006.

14 Varela, M. E., Benito, E., and Keizer, J. J.: Effects of wildfire and laboratory heating on soil  
15 aggregate stability of pine forests in Galicia: the role of lithology, soil organic matter content  
16 and water repellency, Catena, 83, 127-134, 2010.

17 Westerling, A. L., Hidalgo, H. G., Cayan, D. R., and Swetnam, T. W.: Warming and earlier  
18 spring increase western U.S. forest wildfire activity, Science, 313, 940-943, 2006.

19 Zavala, L. M., Granged, A. J. P., Jordán, A., and Bárcenas-Moreno, G.: Effect of burning  
20 temperature on water repellency and aggregate stability in forest soils under laboratory  
21 conditions, Geoderma, 158, 366-374, 2010.

22



1 **Tables**

2 Table 1 Soil classification and site description for the five sites along elevational transect in the western slopes of  
 3 the Sierra Nevada (adapted from Dahlgren et al., 1997)

Soil Series	Elevation (m)	Ecosystem	MAT <sup>a</sup> (°C)	MAP <sup>b</sup> (cm)	Precip <sup>c</sup>	Dominant vegetation (listed in order of dominance)	Soil taxonomy (family)
Vista	210	Oak woodland	16.7	33	Rain	Annual grasses; <i>Quercus douglasii</i> ; <i>Quercus wislizeni</i>	Coarse-loamy, mixed, superactive,thermic; Typic Haploxerepts
Musick	1384	Oak/mixed-conifer forest	11.1	91	Rain	<i>Pinus ponderosa</i> ; <i>Calocedrus decurrens</i> ; <i>Quercus kelloggii</i> ; <i>Chamaebatia foliolosa</i>	Fine-loamy, mixed, semiactive, mesic; Ultic Haploxeralf
Shaver	1737	Mixed-conifer forest	9.1	101	Snow	<i>Abies concolor</i> ; <i>Pinus lambertiana</i> ; <i>Pinus ponderosa</i> ; <i>Calocedrus decurrens</i>	Coarse-loamy, mixed, superactive, mesic; Humic Dystroxerepts
Sirretta	2317	Mixed-conifer forest	7.2	108	Snow	<i>Pinus jeffreyi</i> ; <i>Abies magnifica</i> ; <i>Abies concolor</i>	Sandy-skeletal, mixed, frigid; Dystric Xerorthent
Chiquito <sup>d</sup>	2865	Subalpine mixed-conifer forest	3.9	127	Snow	<i>Pinus contorta murrayana</i> ; <i>Pinus monticola</i> ; <i>Lupinus</i> species	Sandy-skeletal, mixed; Entic Cryumbrept

4 <sup>a</sup> Mean annual air temperature, calculated from regression equation of Harradine and Jenny (1958)

5 <sup>b</sup> Mean annual precipitation

6 <sup>c</sup> Dominant form of precipitation

7 <sup>d</sup> Tentative soil series

8

1

2 Table 2 Bulk density, water content, pH, C concentration, cation exchange capacity (CEC), specific surface area  
3 (SSA) and particle size distribution for the five soils (mean  $\pm$ standard error, n=3)

Soil series and elevation (m)	Bulk density (g/cm <sup>3</sup> )	Gravimetric water content (%)	pH (CaCl <sub>2</sub> )	Carbon (%)	CEC (cmol/kg)	SSA (m <sup>2</sup> /g)	Particle size distribution <sup>a</sup> (%)		
							Sand	Silt	Clay
Vista (210)	1.26 $\pm$ 0.07	0.7 $\pm$ 0.0	5.53 $\pm$ 0.0	1.51 $\pm$ 0.2	8.40 $\pm$ 1.1	1.75 $\pm$ 0.2	79	11	10
Musick (1384)	0.90 $\pm$ 0.06	9.3 $\pm$ 1.6	4.67 $\pm$ 0.1	7.66 $\pm$ 0.8	25.20 $\pm$ 2.0	4.98 $\pm$ 0.3	60	27	15
Shaver (1737)	0.98 $\pm$ 0.06	8.3 $\pm$ 1.1	4.85 $\pm$ 0.3	2.84 $\pm$ 0.2	10.67 $\pm$ 2.1	3.08 $\pm$ 0.3	80	15	5
Sirretta (2317)	0.61 $\pm$ 0.09	9.9 $\pm$ 2.2	4.54 $\pm$ 0.1	4.74 $\pm$ 0.8	12.23 $\pm$ 2.6	6.63 $\pm$ 0.8	80	15	5
Chiquito (2865)	1.17 $\pm$ 0.03	6.1 $\pm$ 1.9	3.96 $\pm$ 0.1	4.10 $\pm$ 0.2	6.03 $\pm$ 1.8	1.00 $\pm$ 0.04	80	16	4

4 <sup>a</sup> Particle size distribution of topsoil profile from Dahlgren et al. (1997): Vista (0 – 14 cm),  
5 Musick (0 – 29 cm), Shaver (0 – 4 cm), Sirretta (0 – 6 cm) and Chiquito (0 – 6 cm)

6

7

1

2 Table 3 Linear correlation coefficients of changes in soil properties with changes in C concentration


Soil	Correlation coefficient ( $r^2$ ) values					
	Mass loss	SSA	Aggregate Stability	pH (CaCl <sub>2</sub> )	CEC	N concentration
Vista	0.74	0.73	0.21	0.77	0.78	0.89
Musick	0.89	0.58	0.77	0.89	0.96	0.83
Shaver	0.82	0.58	0.68	0.74	0.78	0.93
Sirretta	0.60	0.34	0.47	0.67	0.87	0.86
Chiquito	0.82	0.62	0.78	0.88	0.44	0.87

3




# 1 Figures

2




3


4 Figure 1: Temperature thresholds and ranges associated with fire heating. Figure adopted and  
5 expanded from Massman et al. (2010). (<sup>a</sup> DeBano et al. (1977), <sup>b</sup> Diaz-Ravina et al. (1992), <sup>c</sup>  
6 Giovannini et al. (1988), <sup>d</sup> DeBano (2000), and <sup>e</sup> Knicker (2007))




1  
 2 Figure 2: Map of the five sampling sites along elevational transect in the Western Sierra Nevada,  
 3 California (Base map from U.S. Geological Survey, 2015)



1  
 2 Figure 3: Soil color change across heating temperatures. Colors produced from CIExyY  
 3 colorspace equivalents to Munsell colors (Munsell Color Science Laboratory). CIExyY colors  
 4 were converted to RGB system (Rossel et al., 2006; UC Davis Soil Resource Laboratory) and  
 5 visually compared with Munsell Soil Color book for plotting.




1  
 2 Figure 4: Percent mass lost, water-stable aggregate percent and specific surface area changes  
 3 with increase in heating temperatures. Error bars represent standard error where n=3. Different  
 4 letters represent significantly different means ( $p < 0.05$ ) at temperature after Tukey's HSD  
 5 testing.




1

2 Figure 5: Weight fraction of aggregate sizes: macro (2-0.25 mm), micro (0.25-0.053 mm) and  
 3 silt-clay (<0.053 mm) sizes.



1


2 Figure 6: XRD diagram for Music series soils.



1


2 Figure 7: Relative amounts of minerals identified from powder XRD.

3




4

1 Figure 8: pH (geometric means) and cation exchange capacity (adjusted for mass loss) changes  
 2 with increase in heating temperature. CEC values below the 2 cmol<sub>c</sub>/kg are assigned a value of  
 3 1 for plotting. Error bars represent standard error where n=3. Different letters represent  
 4 significantly different means (p<0.05) at each temperature after Tukey's HSD testing.



5  
 6 Figure 9: Carbon concentration, Nitrogen concentration and C:N atomic ratio changes with  
 7 increase in heating temperature. Error bars represent standard error where n=3. Different letters  
 8 represent significantly different means (p<0.05) at each temperature after Tukey's HSD testing.





1  
 2 Figure 10: Total amount of change from unburned to 650 °C combusted soils showing amount  
 3 of change within the 250 to 450 °C range.