

1 **Thermal alteration of soil organic matter properties: a systematic study to**  
2 **infer response of Sierra Nevada climosequence soils to forest fires**

3

4 Samuel N. Araya<sup>1</sup>, Marilyn L. Fogel<sup>1,2</sup>, and Asmeret Asefaw Berhe<sup>1,2</sup>

5 <sup>1</sup>Environmental Systems Graduate Group, University of California, Merced, CA 95343

6 <sup>2</sup>Life and Environmental Sciences Unit, University of California, Merced, CA 95343

7

8

9 Corresponding author:

10 S. N. Araya

11 e-mail: saraya@ucmerced.edu

12

## 1 **Abstract**

2 Fire is a major driver of soil organic matter (SOM) dynamics, and contemporary global  
3 climate change is changing global fire regimes. We conducted laboratory heating experiment  
4 of soils from five locations across the western Sierra Nevada climosequence to investigate  
5 thermal alteration of SOM properties and determined temperature thresholds for major shifts  
6 in SOM properties. Topsoils (0 to 5 cm depth) were exposed to a range of temperatures that  
7 are expected during prescribed and wild fires (150, 250, 350, 450, 550 and 650 °C), with  
8 increase in temperature, we found that the concentrations of C and N decreased in a similar  
9 pattern among all five soils that varied considerably in their original SOM concentrations and  
10 mineralogies. Soils were separated into discrete size classes by dry sieving. The C and N  
11 concentrations in the larger aggregate size fractions (2-0.25mm) decreased with increase in  
12 temperature that at 450 °C temperature, the remaining C and N were almost entirely  
13 associated with the smaller aggregate size fractions (<0.25mm). We observed a general trend  
14 of <sup>13</sup>C enrichment with temperature increase. There was also <sup>15</sup>N enrichment with  
15 temperature increase followed by <sup>15</sup>N depletion when temperature increased beyond 350 °C.  
16 For all the measured variables, the largest physical, chemical, elemental, and isotopic changes  
17 occurred at the mid-intensity fire temperatures, i.e. 350 and 450 °C. The magnitude of the  
18 observed changes in SOM composition and distribution in three aggregate size classes, as  
19 well as the temperature thresholds for critical changes in physical and chemical properties of  
20 soils (such as specific surface area, pH, cation exchange capacity) suggest that transformation  
21 and loss of SOM are the principal responses in heated soils. Findings of this systematic  
22 investigation of soil and SOM response to heating are critical for predicting how soils are  
23 likely to be affected by future climate and fire regimes.

24 **Keywords :** Thermal alterations, Soil Organic Matter, Fire, Climosequence.

## 1 **1. Introduction**

2 Fire is a common, widespread phenomenon globally (Bowman et al., 2009) that controls  
3 dynamics of soil and soil organic matter (SOM). Vegetation fires burn an estimated 300 to  
4 400 million hectares of land globally every year (FAO, 2005). In the US alone, over 80,000  
5 fires were reported in 2014—including about 63,000 wildland fires, and 17,000 prescribed  
6 burns that burned over 1.5 million and 970,000 ha of land, respectively (National Interagency  
7 Fire Center, 2015). In the Sierra Nevada, vegetation fires have a major influence on the  
8 landscapes. Ecological functions such as plant regeneration, habitat revitalization, biomass  
9 accumulation and nutrient cycling, are influenced by fires (McKelvey et al., 1996).  
10 Historically most fires were caused by lightning fires and vegetation fires play important role  
11 in maintaining the health of many ecosystems around the world (Harrison et al., 2010). In  
12 recent decades, anthropogenic activities have become major causes of vegetation fires  
13 (Caldararo, 2002). Moreover, climate and climatic variations exert strong control on the  
14 distribution, frequency, and severity of fires (Harrison et al., 2010). Significant changes in  
15 global fire regimes are anticipated because of climate change including increased frequency  
16 of fires in the coming decades (Pechony and Shindell, 2010; Westerling et al., 2006).  
17 However, our understanding of how climate change and changes in fire regimes will interact  
18 to influence topsoils in fire affected ecosystems is limited.

19 In addition to combustion of aboveground biomass and alteration of vegetation dynamics,  
20 fires also affect the physical, chemical and biological properties of soils (Certini, 2005;  
21 González-Pérez et al., 2004; Mataix-Solera et al., 2011). The degree of alteration caused by  
22 fires depends on the fire intensity and duration, which in turn depend on factors such as the  
23 amount and type of fuels, properties of above ground biomass, air temperature and humidity,  
24 wind, topography, and soil properties such as moisture content, texture and soil organic  
25 matter (SOM) content (DeBano et al., 1998). The first-order effects of fire on soil are caused

1 by the input of heat causing extreme soil temperatures in topsoil (Badía and Martí, 2003b;  
2 Neary et al., 1999) resulting in loss and transformation of SOM, changes in soil  
3 hydrophobicity, changes in soil aggregation, loss of soil mass, and addition of charred  
4 material and other combustion products (Albalasmeh et al., 2013; Araya et al., 2016; Mataix-  
5 Solera et al., 2011; Rein et al., 2008; Santos et al., In press).

6 The duration of burning regulates the amount of energy transferred through the soil. Fires  
7 with longer residence time and lower temperature typically impact the soil and SOM more  
8 than fires with shorter residence time that burn at a higher temperature (Frandsen and Ryan,  
9 1986; González-Pérez et al., 2004). Penetration of heat down a soil profile depends on  
10 intensity and duration of fire as well as the thermal conductivity of the soil (Steward et al.,  
11 1990). Soil has a low thermal conductivity that in fires only the top few centimeters of soil  
12 experiences extreme temperature, For example, in short duration or low severity fires  
13 temperatures typically reach only 100 – 150 °C at 5 cm depth with no significant change of  
14 temperature at 30 cm depth (DeBano, 2000; Janzen and Tobin-Janzen, 2008).

15 Fire has multiple, complex effects on carbon (C) dynamics in soil. Wildfires alone lead to the  
16 release of up to 4.1 Pg C yr<sup>-1</sup> to the atmosphere in the form of carbon dioxide, with an  
17 additional 0.05 to 0.2 Pg C yr<sup>-1</sup> added to the soil as black or pyrogenic carbon ash (Singh et  
18 al., 2012). The changes in SOM characteristics due to combustion include reduced solubility  
19 of OM due to loss of external oxygen containing functional groups; reduced chain length of  
20 fatty acids, alcohols and other alkyl compounds; higher aromaticity due to transformation of  
21 carbohydrates and lipids; production of pyrogenic carbon; formation of heterocyclic nitrogen  
22 (N) compounds; and macromolecular condensation of humic substances (González-Pérez et  
23 al., 2004). In the long term, fires can affect soils by altering and removing vegetation and  
24 topsoil biomass, and increasing soil erodibility (Carroll et al., 2007; DeBano, 1991),

1 subsequently leading to a shift in plant and microbial populations (Janzen and Tobin-Janzen,  
2 2008).

3 The aim of this study is to determine the effects of heating temperatures on important SOM  
4 properties. We used a laboratory heating experiment on five soils from a well-characterized  
5 climosequence in the western Sierra Nevada mountain range (Dahlgren et al., 1997). We  
6 analyzed changes in SOM quantity and quality following heating treatment with the aim to:  
7 (1) determine magnitudes of change in SOM properties associated with different fire heating  
8 temperatures; (2) identify critical thresholds for these changes; and (3) infer the implications  
9 of changing climate on topsoil SOM properties that might experience changing fire regime.  
10 This study aims to contribute to the systematic evaluation and development of ability to  
11 predict the effect of different intensity fires on soil properties under changing climate and fire  
12 regimes.

## 13 **2. Materials and methods**

14 Following laboratory heating of five soils from western Sierra Nevada to temperatures  
15 ranging from 150 to 650 °C, we analyzed changes in SOM quality and quantity. We  
16 measured the changes in C and N concentration in the soil and changes in the distribution of  
17 C and N to different aggregate size classes. We also measured changes in isotopic  
18 composition of  $^{13}\text{C}$  and  $^{15}\text{N}$  in the soils and in the different aggregate size classes. Changes in  
19 SOM quality was analyzed using Fourier-transform infrared (FTIR) spectroscopy of soils.  
20 Description of the study site is given in Section 2.1 and details of the methods used are given  
21 in Sections 2.2 to 2.4.

### 22 **2.1. Study site and soil description**

23 For this study, we collected soils from five sites across an elevation transect along the  
24 western slope of the central Sierra Nevada, California (Figure 1); the sites were previously

1 characterized by Dahlgren et al. (1997). We selected four forested sites that are likely to  
2 experience forest fires and a fifth lower elevation grassland site. The thermal alterations of  
3 bulk soil physical and chemical properties from the same study soils was previously reported  
4 in Araya et al. (2016).

5 All the sites have a Mediterranean climate characterized by warm to hot dry summers and  
6 cool to cold wet winters. Mean annual air temperature ranges from 16.7 °C at the lowest site  
7 located at 210 m to 3.9 °C at the highest elevation site which is at an elevation of 2865 m.  
8 Annual precipitation ranges from 33 cm at the lowest site to 127 cm at the highest site  
9 (Dahlgren et al., 1997; Rasmussen et al., 2007) (Table 1).

10 The lower elevation woodlands of Sierra Nevada experience less frequent fires than further  
11 upslope and the fires are often fast moving and lower severity (Skinner and Chang, 1996). At  
12 the middle-elevation zone of Sierran forest, the mixed conifer zones, frequent fires are low to  
13 moderate severity at lower altitudes but fire frequency generally increases with altitude  
14 towards the upper elevation of the mixed conifer forest (Caprio and Swetnam, 1993). Fires  
15 are infrequent and low severity within the high altitude, Subalpine, zone of Sierra (Skinner  
16 and Chang, 1996).

17 Soils from the lowest elevation site, Vista Series soils (210 masl), fall within the oak  
18 woodland zone (elevations < 1008 m). This is the only soil in our study that does not have an  
19 O-horizon, the soil has dense annual grass cover, however, and the A-horizon SOM  
20 originates mainly from root turnover. The Musick Series soils (1384 masl) lie within  
21 oak/mixed-conifer forest (1008—1580 masl) and mixed-conifer forest (1580—2626 masl).  
22 These soils receive the highest litter fall biomass. The Shaver and Sirretta Series soils (1737  
23 and 2317 masl, respectively) fall within the mixed-conifer forest range zone while the  
24 Chiquito Series soils (2865 masl) lies within the subalpine mixed-conifer forest range

1 (2626—3200 masl). These soils have lower litter fall compared to the lower elevation soils  
2 (van Wagtenonk and Fites-Kaufman, 2006).

3 The western slope of central Sierra Nevada presents a remarkable climosequence of soils that  
4 developed under similar granitic parent material and are located in landscapes of similar age,  
5 relief, slope and aspect (Trumbore et al., 1996) with significant developmental differences  
6 attributed to climate. The soils at mid-elevation range (1000 to 2000 masl) tend to be highly  
7 weathered while soils at high and low elevations are relatively less developed (Dahlgren et  
8 al., 1997; Harradine and Jenny, 1958; Huntington, 1954; Jenny et al., 1949). Among the most  
9 important changes in soil properties along the climosequence include changes in soil organic  
10 carbon (SOC) concentration, base saturation, and mineral desilication and hydroxyl-Al  
11 interlayering of 2:1 layer silicates. Soil pH generally decreases with elevation and the  
12 concentrations of clay and secondary iron oxides show a step change at the elevation of  
13 present-day average effective winter snowline, i.e. 1600 m elevation (Tables 1 and 2)  
14 (California Department of Water Resources, 1952-1962; Dahlgren et al., 1997).

## 15 **2.2. Experimental design and sample collection**

16 Triplicate samples (0 to 5 cm depth) were collected at the five sites, approximately 10 m  
17 apart from each other. Any overlaying organic layer was removed prior to sampling so that  
18 only mineral soil was collected. The soils were air-dried at room temperature and passed  
19 through 2 mm sieve. Prior to furnace heating, the soils were oven dried at 60 °C overnight.  
20 Soil bulk density and field soil moisture were determined from separate undisturbed core  
21 samples collected from each site (Table 2).

22 Sub-samples from each soil were heated in muffle furnace to one of six selected maximum  
23 temperatures (150, 250, 350, 450, 550 and 650 °C). To ensure uniform soil heating and  
24 reduce formation of heating gradient inside, the soils were packed 1 cm high in a 7 cm

1 diameter porcelain flat capsule crucibles. Oxygen supply was not limited during the  
2 heating—the volume of soil sample to volume air in furnace was approximately 1:50.  
3 Furnace temperature was ramped a rate of 3 °C min<sup>-1</sup> and soils were exposed to the maximum  
4 temperature for 30 minutes. Once cooled to touch, soils were stored in in air-tight  
5 polyethylene bags prior to analysis.

6 The six heating temperatures were selected to correspond with fire intensity categories that  
7 are based on maximum surface temperature (DeBano et al., 1977; Janzen and Tobin-Janzen,  
8 2008; Neary et al., 1999), that is, low intensity (150 and 250 °C), medium intensity (350 and  
9 450 °C), and high intensity (550 and 650 °C). These fire intensity classes generally  
10 correspond with thresholds for important thermal reactions in soils observed by differential  
11 thermal analyses (Giovannini et al., 1988; Soto et al., 1991; Varela et al., 2010). Heating rate  
12 of 3 °C min<sup>-1</sup> is preferred in laboratory fire simulation experiments (Giovannini et al., 1988;  
13 Terefe et al., 2008; Varela et al., 2010), the slow heating rate prevents sudden combustion  
14 when soil's ignition temperature is reached at about 220 °C (Fernández et al., 1997, 2001;  
15 Varela et al., 2010). The samples were exposed to the maximum set temperature for a period  
16 of 30 minutes. This length of time ensures that the entire sample is uniformly heated at the set  
17 temperature and is in keeping with wide majority of similar laboratory soil heating  
18 experiments (for example Badía and Martí, 2003a; Fernández et al., 2001; Giovannini, 1994;  
19 Varela et al., 2010; Zavala et al., 2010). The duration of soil heating under vegetation fires is  
20 highly varied and not uniform across landscape (Parsons et al., 2010). The same heating  
21 procedure was used for all the soils so that it would be possible to compare how the soils  
22 from different climate regimes are likely to respond to the fires.



### 1 2.3. Laboratory analysis

2 Dry-aggregate size distribution was measured by sieving. Samples were dry sieved into three  
3 aggregate size classes: 2–0.25 mm (macro-aggregates), 0.25–0.053 mm (micro-aggregates)  
4 and <0.053 mm (silt and clay sized particles or composites). These aggregate size classes  
5 were selected to enable comparison with other studies that investigated the effect of different  
6 natural and anthropogenic properties on soil aggregate dynamics and aggregate protected  
7 organic matter (Six et al 2000).

8 C and N concentrations and stable isotope ratios were measured using an elemental  
9 combustion system (Costech ECS 4010 CHNSO Analyzer, Costech Analytical Technologies,  
10 Valencia, CA, USA) that is interfaced with a mass spectrometer (DELTA V Plus Isotope  
11 Ratio Mass Spectrometer, Thermo Fisher Scientific, Inc, Waltham, MA, USA). For the  
12 analyses, air-dried soil samples were ground to powder consistency on a ball-mill (8000M  
13 MiXer/Mill, with a 55 ml tungsten Carbide Vial, SPEX SamplePrep, LLC, Metuchen, NJ,  
14 USA) and oven dried at 60 °C for over 36 hours. This lower temperature and longer duration  
15 oven-drying was used to avoid possible heating related C or N changes that might occur if  
16 drying was done 105 °C (Kaiser et al., 2015). The C and N concentration results were  
17 corrected for moisture by oven-drying subsamples at 105 °C overnight. The C and N  
18 concentration results were corrected by adjusting for moisture as:  $W_{adj} = W \times (100 - W_m)$ .  
19 Where  $W_{adj}$  is the adjusted percent concentration,  $W$  is the concentration before moisture  
20 adjustment and  $W_m$  is the percent moisture content. All concentration changes resulting from  
21 moisture adjustment were a decrease of less than 1% of the value. The stable isotope ratios  
22 are presented using the  $\delta$  notation (per mill, ‰) as  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  calculated as:  $\delta =$   
23  $[(R_{sample} - R_{standard})/R_{standard}] \times 1000\text{‰}$ ; where  $R$  is ratio of  $^{13}\text{C}/^{12}\text{C}$  for  $\delta^{13}\text{C}$ , and  
24  $^{15}\text{N}/^{14}\text{N}$  for  $\delta^{15}\text{N}$ . The standards used for analyses are atmospheric  $\text{N}_2$   $\delta^{15}\text{N}$  and Vienna Pee  
25 Dee Belemnite (VPDB)  $\delta^{13}\text{C}$ .

1 Bulk soil organic matter composition was analyzed using Fourier-transform infrared (FTIR)  
2 spectroscopy on a Bruker IFS 66v/S vacuum FT-IR spectrometer (Bruker Biosciences  
3 Corporation, Billerica, MA, USA). We used diffuse reflectance infrared Fourier-transform  
4 (DRIFT) technique (Ellerbrock and Gerke, 2013; Parikh et al., 2014). Powder samples were  
5 dried overnight at 60 °C and scanned in mid-IR from 4000 to 400 cm<sup>-1</sup>. In this study, we used  
6 Non-KBr diluted samples were used after preliminary analyses revealed that dilution is not  
7 necessary. KBr dilution is not required for soils with low (<10%) organic matter  
8 concentrations (Ellerbrock and Gerke, 2013; Reeves III, 2003). The FTIR spectrum was  
9 collected using KBr background and was baseline corrected using to the Rubberband  
10 correction method with the default 64 baseline points that is part of the OPUS software  
11 (Bruker Corporation, 2009).

## 12 **2.4. Statistical Analysis**

13 All quantitative results are expressed as means of three replicates ± standard error, unless  
14 otherwise indicated. Differences of means were tested by Analysis of Variance (ANOVA)  
15 and pairwise comparison of treatments done using Tukey's HSD test at p<0.05 significance  
16 level. The normality of the data and the homogeneity of variances was checked using  
17 Shapiro-Wilk's and Levene's tests respectively. All statistical analysis were performed using  
18 R statistical software (R Core Team, 2014). The Pearson correlation coefficient was used to  
19 examine relationships between C concentration and changes in soil properties .

## 20 **3. Results**

### 21 **3.1. Carbon and nitrogen concentration**

22 The initial concentration of C ranged from 1.5% (Vista soil, 210 masl) to 7.7 % (Musick  
23 soils, 1384 masl). Soil C concentration continuously decreased with increasing temperature.  
24 The largest decrease occurred between temperatures of 250 and 450 °C. At 450 °C, all soils

1 lost more than 95% of their original C. C concentration changes with heating above 450 °C  
2 were small and not statistically significant at  $p < 0.05$ . The C:N ratio ranged from 10 (Vista  
3 soils, 210 masl) to 29 (Musick soils, 1384 masl). Following a similar pattern to C  
4 concentration changes, the C:N ratio decreased with an increase in heating temperature  
5 (Figure 2).

6 The loss of C and N from soils due to heating showed a similar response among all five soils  
7 (Figure 2). After 250 °C, all the soils lost more than 25% of their initial C (except Shaver  
8 soils that lost only about 10%). At 350 °C all soils lost 50 to 70% of C. Heating at 450 °C led  
9 to the loss of more than 95% of their initial C for all soils in this study. However, the rate of  
10 loss of N was lower than that of C. At temperatures greater than 550 °C there was 5 to 15%  
11 of soil N still remaining. Consequently, we observed a decrease of C:N ratio with increased  
12 heating temperature. All soils continued to lose about 15% soil N for every 100 °C increase  
13 and maintained more than 60% of their N at heating temperatures up to 350 °C. After heating  
14 at 450 °C, all soils lost more than 60% of their original soil N and 85% by 550 °C.

### 15 **3.2. Carbon and nitrogen stable isotopes**

16 The  $\delta^{13}\text{C}$  composition of all soils was indicative of C-3 vegetation. Soil  $\delta^{13}\text{C}$  composition  
17 was most negative at about -28‰ for the lowest elevation Vista site (210 m) and the value  
18 got consistently less negative with an increase in elevation reaching -24‰ for the highest two  
19 sites (i.e. >2317 m elevation). For all soils, there was a general trend of  $\delta^{13}\text{C}$  enrichment with  
20 temperature increase (Figure 2). The largest change (2.5 to 3.0‰) occurred at heating  
21 temperature between 250 and 450 °C for the lower elevation soils and between 150 and  
22 450 °C for the two highest elevation soils. For the two highest elevation soils, there was a  
23 significant ( $p < 0.05$ ) depletion above that temperature. For all soils, except Musick (1384 m)  
24 and Shaver (1737 m), the maximum enrichment occurred at 450 °C. All soils showed a

1 similar pattern  $\delta^{15}\text{N}$  composition change with temperature. The soils were increasingly  $\delta^{15}\text{N}$   
2 enriched with temperature increase up to 350 °C. At temperatures above 350 °C, the soils got  
3 more  $\delta^{15}\text{N}$  depleted with the most negative  $\delta^{15}\text{N}$  occurring at 650 °C (Figure 2).

### 4 **3.3. Carbon and nitrogen distribution in aggregate size fractions**

5 C and N concentrations, as well as  $^{13}\text{C}$  and  $^{15}\text{N}$  stable isotope ratios were measured for  
6 individual soil aggregate size class. The analysis was done on samples heated up to a  
7 temperature of 450°C. The concentration of C and N in samples heated above 450 °C was too  
8 low to measure significant changes in C distribution in the different aggregate size classes.

9 The distribution of C in the three aggregate sizes fractions followed the same general pattern  
10 with increase in the heating temperatures. The macro aggregate size fraction (2-0.25 mm) had  
11 the least C concentration and silt-clay size particles (<0.053 mm) had the largest  
12 concentration of C (Figure 3). N concentration for the macro size aggregates was below the  
13 detection limit at 450°C for Chiquito and Sirretta. The change in C and N concentration  
14 across heating temperature was similar for all soils

15 The distribution of C and N in different size aggregates did not change noticeably except at  
16 450 °C where concentration in all three fractions converged to zero. The distribution of N in  
17 the three aggregate sizes fractions was similar to that of C and followed a similar pattern  
18 across all the heating temperatures. Similarly, the macro aggregate size fraction (2-0.25 mm)  
19 had the least amount of N concentration and silt-clay size particles (<0.053 mm) had the  
20 largest concentration of N. For Shaver (1737 m), Sirretta (2317 m) and Chiquito (2865 m)  
21 soils, the macro size aggregate N concentration was too low and could not be detected  
22 (Figure 3). The atomic C:N ratio generally stayed the same for all soils through the  
23 temperatures. C:N ratio was highest in macro size aggregates, which had lowest C and N  
24 concentrations, followed by micro and by silt-clay sizes for all soils.

1 The stable isotope composition of  $^{13}\text{C}$  was very similar between aggregate sizes with silt-clay  
2 size aggregates being slightly more enriched except for Shaver (1737 m), which had slightly  
3 more enriched macro aggregates. On the other hand, the  $\delta^{15}\text{N}$  values showed clear differences  
4 among aggregate fractions even though the measured values of  $\delta^{15}\text{N}$  did not change notably  
5 with combustion temperatures.  $\delta^{15}\text{N}$  was highest in silt-clay size particles and lowest in  
6 macro size aggregates with the micro size aggregates showing intermediate values. The  
7 pattern of change in  $\delta^{15}\text{N}$  across combustion temperatures did not affect this order of  $\delta^{15}\text{N}$   
8 values among aggregate fractions. Most of the C and N in the soils was associated with the  
9 larger, macro and micro, aggregate size fractions. With the exception of Vista (210 m) soils,  
10 the distribution changed with increase in heating temperature where the concentrations in  
11 macro aggregates decreased markedly that the remaining C and N concentrations were  
12 distributed between the smaller aggregate fractions (Figure 4). At 450 C, most of the C and N  
13 of the higher altitude soils (Shaver, Sieretta and Chiquito) was now associated with the silt-  
14 clay sized fractions.

### 15 **3.4. FTIR spectroscopy**

16 Changes in chemical composition of SOM due to heating were analyzed by infrared  
17 spectroscopy using Diffuse reflectance infrared fourier transform (DRIFT) technique. The  
18 spectra and peaks after contrasting levels of thermal treatments exhibited qualitative  
19 similarities among the different soils. FTIR spectra for the soils are shown in Figure 5. One  
20 notable changes that occurred in the functional group composition of SOM with heating is the  
21 lowered absorbance intensity of aliphatic methylene groups (as represented by the aliphatic  
22 C–H stretching peak that appear at bands between  $2950 - 2850 \text{ cm}^{-1}$ ) at  $>250 \text{ }^\circ\text{C}$  in all soils.  
23 When comparing intensity of peaks at  $2910 - 2930$  and  $2853 \text{ cm}^{-1}$  wave numbers (from  
24 aliphatic methyl and methylene groups, band A) with those at  $1653$  and  $1400 \text{ cm}^{-1}$  (oxygen  
25 containing carboxyl and carbonyl groups, band B), the decrease in prominence in the

1 aliphatic C-H peak occurs early in the heating sequence while the C=O band shows little  
2 relative change. In addition, after heating at a temperature of 550 °C all soils lost the O-H  
3 stretching peaks (between 3700 – 3200 cm<sup>-1</sup>). In a pattern that is more prominent for the  
4 Musick soil that had the highest concentration of OM, the aromatic C=C stretch around 1600  
5 cm<sup>-1</sup> gets more resolved with increase in heating temperature. This pattern in the C=C is  
6 visible, but less well resolved in the rest of the soils, especially the Vista soil that showed the  
7 least resolved aromatic C=C stretch peak at this region.

## 8 **4. Discussion**

### 9 **4.1. Changes in SOM concentration, distribution and composition**

10 Our results show significant effects of combustion temperature on concentration, distribution,  
11 and composition of SOM on topsoils that experience the most intense heating during  
12 vegetation fires. Topsoils have relatively high OM and low clay content that render them  
13 more sensitive to heating as the SOM experiences significant changes during heating. In our  
14 study system, the effect of fire heating on SOM ranged from slight distillation (volatilization  
15 of minor constituents) typically at temperatures below 150°C, to charring which typically  
16 starts at temperatures above 350°C and complete combustion, consistent with findings of  
17 previous studies (Badía and Martí, 2003b; Certini, 2005). Our findings also confirmed that  
18 regardless of the differences our soils had in mineralogy and other soil physical and chemical  
19 properties, the heating treatments (as proxy for wild fires) led to consistent decrease in  
20 concentration of soil C. This was in agreement with previous studies that showed decrease in  
21 soil C concentration in topsoil after fires (for example Badía et al. (2014); Certini (2005)).  
22 However, this loss of C is expected to be restricted to topsoil while it is expected that the C  
23 concentration in subsoil is likely to remain unchanged or may even increase (for example

1 Dennis et al. (2013); Kavdir et al. (2005)) due to incorporation of necromass from surface  
2 biomass (Almendros et al., 1990; Knicker et al., 2005).

3 We observed significant changes in concentration, distribution, and composition of SOM  
4 with increasing heating temperature. The steep decline in concentration of C in soil that we  
5 observed between this study is consistent with decrease of about 25% C at 250°C and an  
6 almost 99% loss at 450°C (Figure 6). The magnitude of C loss with heating we observed is  
7 similar to the findings of (Terefe et al., 2008; Ulery and Graham, 1993) that investigated  
8 changes in soil C using artificial heating experiment. Similarly, Giovannini et al. (1988) also  
9 found OM decrease started at 220 °C with about 15% loss of OM and about 90% OM loss at  
10 460 °C; while Fernández et al. (1997) reported 37% of SOM loss at 220 °C and 90% at  
11 350 °C. Furthermore, along with the change in C concentration; between 150 °C and before  
12 almost total loss of C above 450 °C, the SOM went through significant qualitative changes  
13 that included decrease in C:N ratio, enrichment in  $\delta^{13}\text{C}$  isotope, changes in  $\delta^{15}\text{N}$  isotope, and  
14 changes in FTIR spectra. Loss in N after fire heating is the result of combustion and  
15 volatilization (Fisher and Binkley, 2000). In this study, we observed that N is not as  
16 significantly reduced until 350°C with about 75% N remaining as opposed to greater than  
17 50% loss of C concentration at the same temperature (Figure 6). Previously studies had  
18 showed that moderate to high intensity fires convert most organic-N into inorganic forms of  
19 N, such as Ammonium (Certini, 2005; Huber et al., 2013). Ammonium is the immediate  
20 combustion product that contributes to formation of nitrate ( $\text{NO}_3^-$ ) by nitrification reactions in  
21 weeks or months after fire. Other studies have showed that a considerable amount of N is  
22 transferred into pyrogenic OM products, to black N (de la Rosa and Knicker, 2011; Knicker,  
23 2010), which would also explain the decrease of the C:N. Decrease in C:N ratio with fire  
24 heating has previously been observed in both laboratory and field fire studies (Badía and  
25 Martí, 2003a; Certini, 2005; Fernández et al., 1997; González-Pérez et al., 2004) .

1 SOM has a C isotopic composition that reflects the  $\delta^{13}\text{C}$  signature of native vegetation. Plants  
2 are depleted in  $\delta^{13}\text{C}$  relative to atmosphere. The  $\delta^{13}\text{C}$  composition for our soils indicated that  
3 the dominant source of OM in all soils is C3 plant biomass that had average  $\delta^{13}\text{C}$  of -27%,  
4 with the higher elevation soils having more positive  $\delta^{13}\text{C}$  than the low elevation. Enrichment  
5 of  $^{13}\text{C}$  with heating is consistent with the loss of plant derived C. In addition, the fact that  
6 lipids (that have relatively more  $\delta^{13}\text{C}$  depleted than the woody materials) are combusted at  
7 lower temperatures than woody materials (such as cellulose and lignin), might contribute to  
8 the enrichment of  $\delta^{13}\text{C}$  with heating (Czimczik et al., 2002). The stable C and N isotope  
9 composition of our soils showed significant fractionation with temperature.  $\delta^{13}\text{C}$  values  
10 became more positive (enriched in  $\delta^{13}\text{C}$ ) up to 450 °C where up to 99% of C was lost (Figure  
11 6). At higher temperature there was a less uniform pattern among the soils. For the last <1%  
12 C, Sirretta and Chiquito soils continued to be more negative (depleted in  $\delta^{13}\text{C}$ ) at higher  
13 temperature while for the rest of the soils there was a slight depletion at 550°C followed by a  
14 slight enrichment at 650 °C (Figure 2). The depletion of  $\delta^{13}\text{C}$  at 550 and 650 °C we found in  
15 this study is likely a result of SOM charring as there was little or no decrease in C  
16 concentration between these temperatures. In a wood charring experiment (non-oxygen  
17 atmosphere) at 150, 340 and 480 °C, Czimczik et al. (2002), observed an enrichment of  $\delta^{13}\text{C}$   
18 at 150 °C where there was no C concentration change but a depletion of  $\delta^{13}\text{C}$  at 340 and  
19 480 °C with charring where the C concentration increased over 50% due to charring.

20 Fires tend to lead to enrichment of  $^{15}\text{N}$ , this is particularly observed in soils immediately in  
21 the aftermath of fires (Boeckx et al., 2005; Grogan et al., 2000; Herman and Rundel, 1989;  
22 Huber et al., 2013), but there is limited information available on the exact temperature ranges  
23 that cause specific levels of  $^{15}\text{N}$  enrichment. In this study, we observed enrichment of  $^{15}\text{N}$  up  
24 to 350 °C and depletion after 350 °C for all soils (Figure 2). It is likely that the continued  $^{15}\text{N}$   
25 enrichment with heating is the result of fractionation due to combustion and volatilization of



1 organic matter which discriminate against  $^{15}\text{N}$ . However, the exact mechanism behind  
2 continued depletion of  $^{15}\text{N}$  when heated above 350 °C remains unclear. One potential  
3 explanation for the  $^{15}\text{N}$  depletion at higher temperatures could be indiscriminate removal of N  
4 as higher temperatures cause the combustion and volatilization process to happen instantly,  
5 compared to charring of OM at lower temperatures. In a post fire-analysis of  $\delta^{15}\text{N}$  on a sub-  
6 alpine ecosystem in Australia, Huber et al. (2013), found that the  $^{15}\text{N}$  enrichment of bulk  
7 surface soil (from unburnt leaves) was higher than that of the charred OM, which was again  
8 higher than that of the ash. They attributed this difference in enrichment level to be the result  
9 of the lower heating intensity experienced by the bulk soil which provided slower processes  
10 for greater fractionation, while higher heat intensity experienced in by the ash result in full  
11 combustion of plant material providing little opportunity for isotopic discrimination. The  
12 temperature range where we observed the depletion of  $^{15}\text{N}$  in our experiment corresponds  
13 with the range where steep decline in N concentration happened (Figure 6), which would be  
14 consistent with the explanation.

#### 15 *4.1.1. Implication of SOM changes with heating*

16 The alterations and loss of SOM is likely more important cause of soil property changes  
17 rather than alterations to soil minerals. SOM is vulnerable to temperatures while soil minerals  
18 are only affected at much higher temperatures (Araya et al., 2016). In addition, all of the soils  
19 in our study are characterized by low clay content and low concentration of reactive minerals,  
20 but high concentration of SOM especially in topsoil leading to strong relationships between  
21 SOM concentrations and soils' physical properties.

22 Degradation of lignin and hemicellulose begins between 130 and 190 °C (Chandler et al.,  
23 1983); and carbohydrate signal is completely removed from  $^{13}\text{C}$  NMR spectra by 350 °C.  
24 Furthermore, Knicker observed loss of stable alkyl C and carboxyl C at 350 °C leading to

1 enrichment of aromatic functional groups in the remaining residue, consistent with what  
2 would expected from incomplete combustion of OM during fires, leading to transformation  
3 and production of charred products (Almendros et al., 2003; Knicker et al., 1996). FTIR  
4 analyses from our work showed that the aliphatic O–H stretch peak (bands 3700 – 3200 cm<sup>-</sup>  
5 <sup>1</sup>) disappeared at temperatures above 550 °C for all soils accompanied by nitriles or  
6 methanenitrile C≡N stretch (2300 – 2200 cm<sup>-1</sup>) at temperature above 450 suggesting  
7 condensation of aromatic functional groups.

8 Loss of OM from soil due to combustion has multiple implications on soil physical and  
9 chemical properties. Simple linear correlation between C concentration changes and other  
10 soil physical and chemical changes that we observed with heating (reported here and in Araya  
11 et al. (2016)) show that more than 80% of the variability in mass loss, aggregate strength,  
12 SSA, pH, CEC and N concentrations is associated with changes in C concentration at the  
13 different heating temperatures. Table3 summarizes the correlation coefficients of soil  
14 property changes with change in C concentration. Analyses of associations between C  
15 concentration and several soil properties showed linear association between: C and N  
16 ( $R^2 > 0.8$ ), mass loss ( $R^2 > 0.8$ , except for Vista and Sirretta soils), pH ( $R^2 > 0.8$ , except for  
17 Shaver and Sirretta), CEC ( $R^2 > 0.7$ , except for Chiquito). Linear association between C  
18 concentration and aggregate strength ( $R^2 > 0.7$ , except for Musick and Chiquito which had  
19  $R^2 \sim 0.7$ ). Specific surface area showed relation with C ( $R^2 > 0.7$  except for Vista and Musick).

20 In this study, the greatest changes in SOM occurred between temperatures 250 and 450 °C  
21 and we found that temperatures below 250 °C had little effect on the quality and quantity of  
22 SOM. This implies that lower intensity fires, such as typical prescribed fires, where soil  
23 surface temperatures do not exceed 250 °C (Janzen and Tobin-Janzen, 2008) have minimum  
24 impact on SOM.

## 1 **4.2. Climate Change Implications**

2 Investigation of the response of climosequence soils to different heating temperature in this  
3 study enables us to infer how, in the long-term, changes in climate are likely to alter the  
4 effect of fires on topsoil physical and chemical properties. Along our study climosequence,  
5 we observed critical differences in response of topsoils based mostly on concentration OM in  
6 soil and soil development stages of each soil. Soil OM concentration and composition in  
7 particular has been shown to respond to changes in precipitation amount and distribution, as  
8 is expected in the Sierra Nevada (Berhe et al., 2012b). Consequently, changes in soil C  
9 storage associated with climate change are expected to lead to different amounts of C loss  
10 due to fires. This is evidenced by the observed highest total mass of C loss from the mid-  
11 elevation Musick soil that had the highest carbon stock, compared to soils in either side of  
12 that elevation range. Anticipated changes in climate in the Sierra Nevada mountain ranges are  
13 expected to include upward movement of the rain-snow transition line exposing areas that  
14 now receive most of their precipitation as snow to rainfall and associated runoff (Arnold et  
15 al., 2015, 2014; Stacy et al., 2015). Upward moving of the rain-snow transition zone under  
16 anticipated climate change scenarios and associated more intense weathering at higher  
17 elevation zones can render more C to loss during fires. More than 80% of the variability in  
18 mass loss, aggregate strength, SSA, pH, CEC and N concentrations is associated with  
19 changes in C concentration (Table 3). Hence, as the vulnerability of these ecosystems to  
20 increased fire frequency increases, due to climate change (Westerling et al., 2006), we can  
21 expect more soil C loss with fires, along with associated changes in soil chemical and  
22 physical properties. In particular, our findings of important changes in soil physical and  
23 chemical properties occurring between 250-450 °C are important for recognizing that critical  
24 transformations of topsoil SOM are likely to occur when, as a result of climate change,  
25 systems that are adapted to low severity fires experience medium to high severity fires.

1 The different responses of soil aggregation in our climosequence to the treatment  
2 temperatures also suggest potential loss and transformation of the physically protected C pool  
3 in topsoil. Degradation of aggregates during fire (Albalasmeh et al., 2013) is likely to render  
4 aggregate-protected C to potential losses through oxidative decomposition, leaching and  
5 erosion. Moreover, in systems such as the Sierra Nevada which are dominated by steep  
6 slopes, movement of the rain-snow transition zone upward is likely to increase proportion of  
7 precipitation that occurs as rain. The kinetic energy of raindrops and the observed increase in  
8 hydrophobicity of soils after fires (Johnson et al., 2007; Johnson et al., 2004) can lead to  
9 higher rates of erosional redistribution of especially the free light fraction or particulate C that  
10 is not associated with soil minerals (Berhe et al., 2012a; Berhe and Kleber, 2013; McCorkle  
11 et al., 2016; Stacy et al., 2015).

## 12 **5. Conclusion**

13 Previously, considerable amount of work had been published to demonstrate how fires affect  
14 OM concentration and composition in biomass. This study fills critical gaps by determining  
15 how and to what extent OM in soil experiences changes due to heating. The findings of this  
16 study also showed that changes in soil properties during heating are closely related to changes  
17 in C concentrations in soil. The temperatures most critical to C loss and alteration were found  
18 to be 250 °C, where charring of organic matter starts and 450 °C where most of the SOM is  
19 combusted. Most soil properties exhibited a steep change in this temperature range. SOM  
20 exhibited largest change, i.e. soils became enriched in <sup>13</sup>C and <sup>15</sup>N isotopic composition until  
21 approximately 90% of C and N was lost, at higher temperatures slight depletion of <sup>13</sup>C and  
22 steep depletion of <sup>15</sup>N is observed. FTIR spectroscopy showed the reduction and  
23 disappearance of aliphatic OH functional groups with temperature increase and accumulation  
24 of aromatic carbon groups.

1 This study presented the effects of heat input on topsoil properties. The study is necessary to  
2 understand thermally induced changes on soil properties in isolation from other variables that  
3 accompany vegetation fires such as the addition of pyrolysis products from plants and ash,  
4 and the fire induced soil moisture dynamics. Findings from this study will contribute towards  
5 estimating the amount and rate of change in carbon and nitrogen loss, and other essential soil  
6 properties that can be expected from topsoil exposure to different intensity fires under  
7 anticipated climate change scenarios.

## 8 **6. Acknowledgements**

9 The authors would like to thank Prof. Randy A. Dahlgren for providing us with geo-  
10 references for the study sites, background data, and for his comments on an earlier version of  
11 this manuscript. We thank Dr. Christina Bradley for her help and expertise in analysis of C  
12 and N; and Dr. Samuel Traina for his comments on an earlier version of this manuscript.  
13 Funding for this work was provided by UC Merced Graduate Research Council grant  
14 and<sup>[1]</sup><sub>SEP</sub> National Science Foundation (CAREER EAR - 1352627) award to A. A. Berhe.

## 1 7. References

- 2 Albalasmeh, A. A., Berli, M., Shafer, D. S., and Ghezzehei, T. A.: Degradation of moist soil aggregates by rapid  
3 temperature rise under low intensity fire, *Plant Soil*, 362, 335-344, 2013.
- 4 Almendros, G., Gonzalez-Vila, F. J., and Martin, F.: Fire-induced transformation of soil organic matter from an  
5 oak forest: an experimental approach to the effects of fire on humic substances, *Soil Sci*, 149, 158-168, 1990.
- 6 Almendros, G., Knicker, H., and González-Vila, F. J.: Rearrangement of carbon and nitrogen forms in peat after  
7 progressive thermal oxidation as determined by solid-state <sup>13</sup>C-and <sup>15</sup>N-NMR spectroscopy, *Organic*  
8 *Geochemistry*, 34, 1559-1568, 2003.
- 9 Araya, S. N., Meding, M., and Berhe, A. A.: Thermal alteration of soil physico-chemical properties: a systematic  
10 study to infer response of Sierra Nevada climosequence soils to forest fires, *Soil*, 2, 351-366, 2016.
- 11 Arnold, C., Ghezzehei, T. A., and Berhe, A. A.: Decomposition of distinct organic matter pools is regulated by  
12 moisture status in structured wetland soils, *Soil Biology and Biochemistry*, 81, 28-37, 2015.
- 13 Arnold, C., Ghezzehei, T. A., and Berhe, A. A.: Early spring, severe frost events, and drought induce rapid carbon  
14 loss in high elevation meadows, *PloS one*, 9, e106058, 2014.
- 15 Badía, D. and Martí, C.: Effect of simulated fire on organic matter and selected microbiological properties of two  
16 contrasting soils, *Arid Land Research and Management*, 17, 55-69, 2003a.
- 17 Badía, D. and Martí, C.: Plant ash and heat intensity effects on chemical and physical properties of two contrasting  
18 soils, *Arid Land Research and Management*, 17, 23-41, 2003b.
- 19 Badía, D., Martí, C., Aguirre, A. J., Aznar, J. M., González-Pérez, J. A., De la Rosa, J. M., León, J., Ibarra, P.,  
20 and Echeverría, T.: Wildfire effects on nutrients and organic carbon of a Rendzic Phaeozem in NE Spain: Changes  
21 at cm-scale topsoil, *Catena*, 113, 267-275, 2014.
- 22 Berhe, A. A., Harden, J. W., Torn, M. S., Kleber, M., Burton, S. D., and Harte, J.: Persistence of soil organic  
23 matter in eroding versus depositional landform positions, *J Geophys Res-Bioge*, 117, 2012a.
- 24 Berhe, A. A. and Kleber, M.: Erosion, deposition, and the persistence of soil organic matter: mechanistic  
25 considerations and problems with terminology, *Earth Surface Processes and Landforms*, 38, 908-912, 2013.
- 26 Berhe, A. A., Suttle, K. B., Burton, S. D., and Banfield, J. F.: Contingency in the Direction and Mechanics of Soil  
27 Organic Matter Responses to Increased Rainfall, *Plant Soil*, 358, 371-383. DOI 310.1007/s11104-11012-11156-  
28 11100, 2012b.
- 29 Boeckx, P., Paulino, L., Oyarzun, C., van Cleemput, O., and Godoy, R.: Soil  $\delta^{15}\text{N}$  patterns in old-growth forests  
30 of southern Chile as integrator for N-cycling, *Isot Environ Healt S*, 41, 249-259, 2005.
- 31 Bowman, D. M. J. S., Balch, J. K., Artaxo, P., Bond, W. J., Carlson, J. M., Cochrane, M. A., D'Antonio, C. M.,  
32 DeFries, R. S., Doyle, J. C., Harrison, S. P., Johnston, F. H., Keeley, J. E., Krawchuk, M. A., Kull, C. A., Marston,  
33 J. B., Moritz, M. A., Prentice, I. C., Roos, C. I., Scott, A. C., Swetnam, T. W., van der Werf, G. R., and Pyne, S.  
34 J.: Fire in the Earth System, *Science*, 324, 481-484, 2009.
- 35 Bruker Corporation: OPUS Spectroscopy Software (Version 6.5). 2009.
- 36 Caldararo, N.: Human ecological intervention and the role of forest fires in human ecology, *The Science of the*  
37 *Total Environment*, 292, 141-165, 2002.
- 38 California Department of Water Resources: Water Conditions in California. Resources, C. D. o. W. (Ed.), Bull.  
39 120, Sacramento, 1952-1962.
- 40 Caprio, A. C. and Swetnam, T. W.: Historic fire regimes along an elevational gradient on the west slope of the  
41 Sierra Nevada, California, Missoula, MT, March 30 - April 1 1993, 173-179, 1995.

- 1 Carroll, E. M., Miller, W. W., Johnson, D. W., Saito, L., Qualls, R. G., and Walker, R. F.: Spatial analysis of a  
2 large magnitude erosion event following a Sierran Wildfire, *Journal of environmental quality*, 36, 1105-1105,  
3 2007.
- 4 Certini, G.: Effects of fire on properties of forest soils: a review, *Oecologia*, 143, 1-10, 2005.
- 5 Chandler, C., Cheney, P., thomas, P., Trabaud, L., and William, D.: Fire effects on soil, water and air. In: *Fire in*  
6 *Forestry. Vol I: Forest fire behaviour and effects*, John Wiley Sons, New York, 1983.
- 7 Czimeczik, C. I., Preston, C. M., Schmidt, M. W. I., Werner, R. A., and Schulze, E.-D.: Effects of charring on  
8 mass, organic carbon, and stable carbon isotope composition of wood, *Organic Geochemistry*, 33, 1207-1223,  
9 2002.
- 10 Dahlgren, R. A., Boettinger, J. L., Huntington, G. L., and Amundson, R. G.: Soil development along an elevational  
11 transect in the western Sierra Nevada, California, *Geoderma*, 78, 207-236, 1997.
- 12 de la Rosa, J. M. and Knicker, H.: Bioavailability of N released from N-rich pyrogenic organic matter: An  
13 incubation study, *Soil Biol Biochem*, 43, 2368-2373, 2011.
- 14 DeBano, L. F.: The effect of fire on soil properties, Boise, ID, April 10-12, 1990 1991, 151-156.
- 15 DeBano, L. F.: The role of fire and soil heating on water repellency in wildland environments: a review, *Journal*  
16 *of Hydrology*, 231, 195-206, 2000.
- 17 DeBano, L. F., Dunn, P. H., and Conrad, C. E.: Fire's effect on physical and chemical properties of Chaparral  
18 soils, 1977.
- 19 DeBano, L. F., Neary, D. G., and Ffolliott, P. F.: Fire's effect on ecosystems John Wiley & Sons, Inc., New York,  
20 USA, 1998.
- 21 Dennis, E. I., Usoroh, A. D., and Ijah, C. J.: Soil properties dynamics induced by passage of fire during agricultural  
22 burning, *International Journal of Advance Agricultural Research*, 1, 43-52, 2013.
- 23 Ellerbrock, R. H. and Gerke, H. H.: Characterization of organic matter composition of soil and flow path surfaces  
24 based on physicochemical principles—a review. In: *Advances in Agronomy*, Sparks, D. L. (Ed.), Academic Press,  
25 Burlington, 2013.
- 26 FAO: State of the world's forest Food and Agricultural Organization of the United Nations, Rome, Italy, 2005.
- 27 Fernández, I., Cabaneiro, A., and Carballas, T.: Organic matter changes immediately after a wildfire in an Atlantic  
28 forest soil and comparison with laboratory soil heating, *Soil Biology and Biochemistry*, 29, 1-11, 1997.
- 29 Fernández, I., Cabaneiro, A., and Carballas, T.: Thermal resistance to high temperatures of different organic  
30 fractions from soils under pine forests, *Geoderma*, 104, 281-298, 2001.
- 31 Fisher, R. F. and Binkley, D.: Ecology and management of forest soils, John Wiley and Sons, Inc., , New York,  
32 USA., 2000.
- 33 Frandsen, W. H. and Ryan, K. C.: Soil moisture reduces belowground heat flux and soil temperatures under a  
34 burning fuel pile, *Canadian Journal of Forest Research*, 16, 244-248, 1986.
- 35 Giovannini, G.: The effect of fire on soil quality. In: *Soil Erosion and Degradation as a Consequence of Forest*  
36 *Fires*, Sala, M. and Rubio, J. L. (Eds.), Geofoma Ediciones, Logrono, 1994.
- 37 Giovannini, G., Lucchesi, S., and Giachetti, M.: Effect of heating on some physical and chemical parameters  
38 related to soil aggregation and erodibility, *Soil Sci*, 146, 255-261, 1988.
- 39 González-Pérez, J., González-Vila, F., Almendros, G., and Knicker, H.: The effect of fire on soil organic matter—  
40 a review, *Environment International*, 30, 855-870, 2004.

- 1 Grogan, P., Burns, T. D., and Chapin III, F. S.: Fire effects on ecosystem nitrogen cycling in a Californian bishop  
2 pine forest, *Oecologia*, 122, 537-544, 2000.
- 3 Harradine, F. and Jenny, H.: Influence of parent material and climate on texture and nitrogen and carbon contents  
4 of virgin California soils: texture and nitrogen contents of soils, *Soil Science*, 85, 235-243, 1958.
- 5 Harrison, S. P., Marlon, J. R., and Bartlein, P. J.: Fire in the Earth System. In: *Changing Climates, Earth Systems  
6 and Society.*, Dodson, J. (Ed.), International Year of Planet Earth, Springer, New York, 2010.
- 7 Herman, D. J. and Rundel, P. W.: Nitrogen isotope fractionation in burned and unburned Chaparral soils, *Soil  
8 Science Society of America Journal*, 53, 1229-1236, 1989.
- 9 Huber, E., Bell, T. L., and Adams, M. A.: Combustion influences on natural abundance nitrogen isotope ratio in  
10 soil and plants following a wildfire in a sub-alpine ecosystem, *Oecologia*, 173, 1063-1074, 2013.
- 11 Huntington, G. L.: The effect of vertical zonality on clay content in residual granitic soils of the Sierra Nevada  
12 mountains, University of California, Berkeley, 1954.
- 13 Janzen, C. and Tobin-Janzen, T.: Microbial communities in fire-affected soils. In: *Microbiology of Extreme Soils*,  
14 Dion, P. and Nautiyal, C. S. (Eds.), *Soil Biology*, 13, Springer-Verlag Berlin Heidelberg, 2008.
- 15 Jenny, H., Gessel, S. P., and Bingham, F. T.: Comparative study of decomposition rates of organic matter in  
16 temperate and tropical regions, *Soil Science*, 68, 419-432, 1949.
- 17 Johnson, D., Murphy, J., Walker, R., Glass, D., and MILLER, W.: Wildfire effects on forest carbon and nutrient  
18 budgets, *Ecological Engineering*, 31, 183-192, 2007.
- 19 Johnson, D. W., Susfalk, R. B., Caldwell, T. G., Murphy, J. D., Miller, W. W., and Walker, R. F.: Fire Effects on  
20 Carbon and Nitrogen Budgets in Forests, *Water, Air, & Soil Pollution: Focus*, 4, 263-275, 2004.
- 21 Kaiser, M., Kleber, M., and Berhe, A. A.: How air-drying and rewetting modify soil organic matter characteristics:  
22 An assessment to improve data interpretation and inference, *Soil Biol Biochem*, 80, 324-340, 2015.
- 23 Kavdır, Y., Ekinci, H., Yüksel, O., and Mermut, A. R.: Soil aggregate stability and <sup>13</sup>C CP/MAS-NMR assessment  
24 of organic matter in soils influenced by forest wildfires in Çanakkale, Turkey, *Geoderma*, 129, 219-229, 2005.
- 25 Knicker, H.: "Black nitrogen" - an important fraction in determining the recalcitrance of charcoal, *Organic  
26 Geochemistry*, 41, 947-950, 2010.
- 27 Knicker, H., Almendros, G., González-Vila, F. J., Martín, F., and Lüdemann, H.-D.: <sup>13</sup>C- and <sup>15</sup>N-NMR  
28 spectroscopic examination of the transformation of organic nitrogen in plant biomass during thermal treatment,  
29 *Soil Biology and Biochemistry*, 28, 1053-1060, 1996.
- 30 Knicker, H., Gonzalezvila, F., Polvillo, O., Gonzalez, J., and Almendros, G.: Fire-induced transformation of C-  
31 and N- forms in different organic soil fractions from a Dystric Cambisol under a Mediterranean pine forest, *Soil  
32 Biology and Biochemistry*, 37, 701-718, 2005.
- 33 Mataix-Solera, J., Cerda, A., Arcenegui, V., Jordan, A., and Zavala, L. M.: Fire effects on soil aggregation: a  
34 review, *Earth-Science Reviews*, 109, 44-60, 2011.
- 35 McCorkle, E. P., Berhe, A. A., Hunsaker, C. T., McFarlane, K. J., Johnson, D., Fogel, M. L., and Hart, S. C.:  
36 Tracing the source of soil organic matter eroded from temperate forested catchments using carbon and nitrogen  
37 isotopes *Chemical Geology*, 2016. 2016.
- 38 McKelvey, K. S., Skinner, C. N., Chang, C.-r., Erman, D. C., Husari, S. J., Parsons, D. J., Wagtendonk, J. W. v.,  
39 and Weatherspoon, C. P.: *An overview of fire in the Sierra Nevada*, Davis, California, 1996.
- 40 National Interagency Fire Center: [https://www.nifc.gov/fireInfo/fireInfo\\_statistics.html](https://www.nifc.gov/fireInfo/fireInfo_statistics.html), last access: 5  
41 June 2015.




- 1 Neary, D. G., Klopatek, C. C., DeBano, L. F., and Ffolliott, P. F.: Fire effects on belowground sustainability: a  
2 review and synthesis, *Forest Ecology and Management*, 122, 51-71, 1999.
- 3 Parikh, S. J., Goyne, K. W., Margenot, A. J., Mukome, F. N. D., and Calderón, F. J.: Soil chemical insights  
4 provided through vibrational spectroscopy. In: *Advances in Agronomy*, Donald, L. S. (Ed.), Academic Press,  
5 2014.
- 6 Parsons, A., Robichaud, P. R., Lewis, S. A., Napper, C., and Clark, J.: Field guide for mapping post-fire soil burn  
7 severity, U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fort Collins, CO,  
8 2010.
- 9 Pechony, O. and Shindell, D. T.: Driving forces of global wildfires over the past millennium and the forthcoming  
10 century, *Proceedings of the National Academy of Sciences of the United States of America*, 107, 19167-19170,  
11 2010.
- 12 R Core Team: *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical  
13 Computing, Vienna, Austria, 2014.
- 14 Rasmussen, C., Matsuyama, N., Dahlgren, R. A., Southard, R. J., and Brauer, N.: Soil Genesis and Mineral  
15 Transformation Across an Environmental Gradient on Andesitic Lahar, *Soil Science Society of America Journal*,  
16 71, 225-237, 2007.
- 17 Reeves III, J. B.: Mid-infrared diffuse reflectance spectroscopy: is sample dilution with KBr necessary, and if so,  
18 when?, *American Laboratory*, 35, 24-28, 2003.
- 19 Rein, G., Cleaver, N., Ashton, C., Pironi, P., and Torero, J. L.: The severity of smouldering peat fires and damage  
20 to the forest soil, *Catena*, 74, 304-309, 2008.
- 21 Santos, F., Russel, D., and Berhe, A. A.: Thermal alteration of water extractable organic matter in climosequence  
22 soils from the Sierra Nevada, California, *Journal of Geophysical Research - Biogeosciences*, In press. In press.
- 23 Singh, N., Abiven, S., Torn, M. S., and Schmidt, M. W. I.: Fire-derived organic carbon in soil turns over on a  
24 centennial scale, *Biogeosciences*, 9, 2847-2857, 2012.
- 25 Skinner, C. N. and Chang, C.-R.: *Fire Regimes, Past and Present*. In: *Sierra Nevada Ecosystem Project, Final  
26 Report to Congress, Vol II, Assessment and Scientific Basis for management Options*, Davis: University of  
27 California, Center for Water and Wildland Resources, 1996.
- 28 Soto, B., Benito, E., Basanta, R., and Díaz-Fierros, F.: Influence of antecedent soil moisture on pedological effects  
29 of fire. In: *Soil Erosion and Degradation as a Consequence of Forest Fires*, Sala, M. and Rubio, J. L. (Eds.),  
30 Geofoma Ed., Logroño, Spain, 1991.
- 31 Stacy, E., Hart, S. C., Hunsaker, C. T., Johnson, D. W., and Berhe, A. A.: Soil carbon and nitrogen erosion in  
32 forested catchments: implications for erosion-induced terrestrial carbon sequestration, *Biogeosciences*, 12, 4861-  
33 4874, 2015.
- 34 Steward, F. R., Peter, S., and Richon, J. B.: A Method for Predicting the Depth of Lethal Heat Penetration into  
35 Mineral Soils Exposed to Fires of Various Intensities, *Can J Forest Res*, 20, 919-926, 1990.
- 36 Terefe, T., Mariscal-Sancho, I., Peregrina, F., and Espejo, R.: Influence of heating on various properties of six  
37 Mediterranean soils: a laboratory study, *Geoderma*, 143, 273-280, 2008.
- 38 Trumbore, S. E., Chadwick, O. A., and Amundson, R.: Rapid exchange between soil carbon and atmospheric  
39 carbon dioxide driven by temperature change, *Science*, 272, 393-396, 1996.
- 40 U.S. Geological Survey: *NLCD 2011 Land Cover (2011 Edition, amended 2014)*. 2014.
- 41 Ulery, A. L. and Graham, R. C.: Forest-fire effects on soil color and texture, *Soil Science Society of America  
42 Journal*, 57, 135-140, 1993.

- 1 van Wagtendonk, J. W. and Fites-Kaufman, J. A.: Sierra Nevada Bioregion. In: Fire in California's Ecosystems,  
2 Sugihara, N. G., van Wagtendonk, J. W., Shaffer, K. E., and Thode, A. E. (Eds.), University of California Press,  
3 Berkeley, CA, USA, 2006.
- 4 Varela, M. E., Benito, E., and Keizer, J. J.: Effects of wildfire and laboratory heating on soil aggregate stability  
5 of pine forests in Galicia: the role of lithology, soil organic matter content and water repellency, *Catena*, 83, 127-  
6 134, 2010.
- 7 Westerling, A. L., Hidalgo, H. G., Cayan, D. R., and Swetnam, T. W.: Warming and earlier spring increase western  
8 U.S. forest wildfire activity, *Science*, 313, 940-943, 2006.
- 9 Zavala, L. M., Granged, A. J. P., Jordán, A., and Bárcenas-Moreno, G.: Effect of burning temperature on water  
10 repellency and aggregate stability in forest soils under laboratory conditions, *Geoderma*, 158, 366-374, 2010.

11

1 **Figures**




2

3 Figure 1: (a) Location of the sampling site on the western slopes of the Sierra Nevada,  
4 California, and (b) map of the five sampling locations and percent tree canopy cover (U.S.  
5 Geological Survey, 2014).


6

7




1

- 1 Figure 2: Bulk soil Carbon and Nitrogen concentrations, C:N atomic ratio, and  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$
- 2 isotope (‰) changes with increase in heating temperature. Error bars represent standard error
- 3 where  $n=3$ . Different letters represent significantly different means ( $p<0.05$ ) at each
- 4 temperature after Tukey's HSD testing.



1

2 Figure 3. C and N concentrations, C:N atomic ratio, and  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  isotope (‰) changes  
 3 in macro (2-0.25 mm), micro (0.25-0.053 mm) and silt-clay sized (<0.053 mm) aggregates  
 4 with increase in heating temperature. Error bars represent standard error where n=3.








1

2 Figure 4: C and N distributions in macro (2-0.25 mm), micro (0.25-0.053 mm) and silt-clay

3 sized (<0.053 mm) aggregates.


4



Absorption peaks (cm <sup>-1</sup> )	Peak assignment
3700, 3625	O-H
3270	O-H, N-H (H bonded)
2940	C-H (Aliphatic)
2345	C≡N
1870	C=O
1600	C=C, C-O, C=O
1310	C-H
1159, 995	Ester, Phenol, C-O-C, C-OH
800	C-H (Aromatic)



1 Figure 5: FTIR spectra of the five soils at the different heating temperatures. Heating  
 2 temperatures, in Celsius, are shown to the right of each spectrum.



3  
 4 Figure 6: (a) Percentage of C and (b) N loss with heating; and (c) change in  $\delta^{13}\text{C}$  and (d)  $\delta^{15}\text{N}$   
 5 versus percent of total C and N lost from soils (error bars represent standard error where  
 6  $n=3$ ).

7

1

2 **Tables**3 Table 1 Soil classification and site description for the five sites along elevational transect in  
4 the western slopes of the Sierra Nevada (adapted from Dahlgren et al., 1997)

Soil Series	Elevation (m)	Ecosystem	MAT <sup>a</sup> (°C)	MAP <sup>b</sup> (cm)	Precip <sup>c</sup>	Dominant vegetation (listed in order of dominance)	Soil taxonomy (family)
Vista	210	Oak woodland	16.7	33	Rain	Annual grasses; <i>Quercus douglasii</i> ; <i>Quercus wislizeni</i>	Coarse-loamy, mixed, superactive,thermic; Typic Haploxerepts
Musick	1384	Oak/mixed-conifer forest	11.1	91	Rain	<i>Pinus ponderosa</i> ; <i>Calocedrus decurrens</i> ; <i>Quercus kelloggii</i> ; <i>Chamaebatia foliolosa</i>	Fine-loamy, mixed, semiactive, mesic; Ultic Haploxeralf
Shaver	1737	Mixed-conifer forest	9.1	101	Snow	<i>Abies concolor</i> ; <i>Pinus lambertiana</i> ; <i>Pinus ponderosa</i> ; <i>Calocedrus decurrens</i>	Coarse-loamy, mixed, superactive, mesic; Humic Dystroxerepts
Sirretta	2317	Mixed-conifer forest	7.2	108	Snow	<i>Pinus jeffreyi</i> ; <i>Abies magnifica</i> ; <i>Abies concolor</i>	Sandy-skeletal, mixed, frigid; Dystric Xerorthent
Chiquito <sup>d</sup>	2865	Subalpine mixed-conifer forest	3.9	127	Snow	<i>Pinus contorta murrayana</i> ; <i>Pinus monticola</i> ; <i>Lupinus</i> species	Sandy-skeletal, mixed; Entic Cryumbrept

5 <sup>a</sup> Mean annual air temperature, calculated from regression equation of Harradine and Jenny  
6 (1958); <sup>b</sup> Mean annual precipitation; <sup>c</sup> Dominant form of precipitation; <sup>d</sup> Tentative soil series  
7

1

2 Table 2 Bulk density, water content, pH, C concentration, cation exchange capacity (CEC),  
 3 specific surface area (SSA) and particle size distribution for the five soils (mean  $\pm$ standard  
 4 error, n=3)

Soil series and elevation (m)	Bulk density (g/cm <sup>3</sup> )	Gravimetric water content (%)	pH (CaCl <sub>2</sub> )	Carbon (%)	CEC (cmol <sub>c</sub> /kg)	SSA (m <sup>2</sup> /g)	Particle size distribution <sup>a</sup> (%)		
							Sand	Silt	Clay
Vista (210)	1.26 $\pm$ 0.07	0.7 $\pm$ 0.0	5.53 $\pm$ 0.0	1.51 $\pm$ 0.2	8.40 $\pm$ 1.1	1.75 $\pm$ 0.2	79	11	10
Musick (1384)	0.90 $\pm$ 0.06	9.3 $\pm$ 1.6	4.67 $\pm$ 0.1	7.66 $\pm$ 0.8	25.20 $\pm$ 2.0	4.98 $\pm$ 0.3	60	27	15
Shaver (1737)	0.98 $\pm$ 0.06	8.3 $\pm$ 1.1	4.85 $\pm$ 0.3	2.84 $\pm$ 0.2	10.67 $\pm$ 2.1	3.08 $\pm$ 0.3	80	15	5
Sirretta (2317)	0.61 $\pm$ 0.09	9.9 $\pm$ 2.2	4.54 $\pm$ 0.1	4.74 $\pm$ 0.8	12.23 $\pm$ 2.6	6.63 $\pm$ 0.8	80	15	5
Chiquito (2865)	1.17 $\pm$ 0.03	6.1 $\pm$ 1.9	3.96 $\pm$ 0.1	4.10 $\pm$ 0.2	6.03 $\pm$ 1.8	1.00 $\pm$ 0.04	80	16	4

5 <sup>a</sup> Particle size distribution of top soil profile from Dahlgren et al. (1997): Vista (0 – 14 cm),  
 6 Musick (0 – 29 cm), Shaver (0 – 4 cm), Sirretta (0 – 6 cm) and Chiquito (0 – 6 cm)

7

8

1

2 Table 3 Linear correlation coefficients of changes in soil properties with changes in C  
3 concentration. All correlation coefficients have p-values < 0.01 unless otherwise indicated.

Correlation coefficient ( $r^2$ ) values						
Soil	Mass loss	SSA	Aggregate Stability	pH (CaCl <sub>2</sub> )	CEC	N concentration
Vista	0.74	0.73	0.21 <sup>a</sup>	0.77	0.78	0.89
Musick	0.89	0.58	0.77	0.89	0.96	0.83
Shaver	0.82	0.58	0.68	0.74	0.78	0.93
Sirretta	0.60	0.34 <sup>b</sup>	0.47	0.67	0.87	0.86
Chiquito	0.82	0.62	0.78	0.88	0.44	0.87

4 <sup>a</sup> p = 0.078; <sup>b</sup> p = 0.035