

# Decision support for the selection of reference sites using $^{137}\text{Cs}$ as a soil erosion tracer

Laura Arata<sup>1</sup>, Katrin Meusburger<sup>1\*</sup>, Alexandra Bürge<sup>1</sup>, Markus Zehringer<sup>2</sup>, Michael E. Ketterer<sup>3</sup>, Lionel Mabit<sup>4</sup> and Christine Alewell<sup>1</sup>

<sup>1</sup> Institute of Environmental Geosciences, Department of Environmental Sciences, University of Basel, Switzerland

<sup>2</sup> State Laboratory Basel-City, Basel, Switzerland

<sup>3</sup> Chemistry Department, Metropolitan State University of Denver, Colorado, USA

<sup>4</sup> Soil and Water Management & Crop Nutrition Laboratory, FAO/IAEA Agriculture & Biotechnology Laboratory, Austria.

*Correspondence to:* Katrin Meusburger (katrin.meusburger@unibas.ch)

**Abstract.** The classical approach to use  $^{137}\text{Cs}$  as soil erosion tracer is based on the comparison between stable reference sites and sites affected by soil redistribution processes, and enables to derive soil erosion and deposition rates. The method is associated with potentially large sources of uncertainty with major parts of this uncertainty being associated with the selection of the reference sites. We propose a decision support tool to Check the Suitability of reference Sites (CheSS). Commonly the variation among  $^{137}\text{Cs}$  inventories of spatial replicate reference samples are taken as sole criteria to decide on the suitability of a reference inventory. Here we propose an extension of this procedure using a repeated sampling approach, where the reference sites are resampled after a certain time period. Suitable reference sites are expected to present no significant temporal variation in their decay-corrected  $^{137}\text{Cs}$  depth profiles. Possible causes of variation are assessed by a decision tree. More specifically, the decision tree tests for (i) uncertainty connected to small scale variability of  $^{137}\text{Cs}$  due to its heterogeneous initial fallout (such as in areas affected by the Chernobyl fallout), (ii) signs of erosion/deposition processes, (iii) artefacts due to the collection, preparation and measurement of the samples and (iv) finally, if none of the above can be assigned, this variation might be attributed to “turbation” processes (e.g. bioturbation, cryoturbation and mechanical turbation such as avalanches or rock falls). CheSS was applied in a case study site in a Swiss alpine valley, where the apparent temporal variability was questioning the suitability of selected reference sites. In general we suggest the application of CheSS to implement first steps towards a comprehensible approach to test for the suitability of reference sites.


Keywords: FRN, fallout radionuclides, soil degradation,  $^{210}\text{Pb}_{\text{ex}}$ ,  $^{239+240}\text{Pu}$ , comparability of gamma spectrometers, Cesium-137

## 1 Introduction

Soil erosion is a global threat (Lal, 2003). Recent estimated erosion rates range from low rates of  $0.001\text{--}2\text{ t ha}^{-1}\text{ yr}^{-1}$  on flat relatively undisturbed lands (Patric, 2002) to high rates under intensive agricultural use of  $> 50\text{ t ha}^{-1}\text{ yr}^{-1}$ . In mountainous

regions, rates ranging from 1–30 t ha<sup>-1</sup> yr<sup>-1</sup> have been reported (e.g. Descroix et al. 2003, Frankenberg et al. 1995, Konz et al., 2012) where they often exceed the natural process of soil formation (Alewell et al., 2015). The use of the artificial radionuclide <sup>137</sup>Cs as a soil erosion tracer has been increasing during the last decades, and the method has been applied all over the world with success (e.g. Mabit et al., 2013; Zapata, 2002). The use of <sup>137</sup>Cs as soil erosion tracer allows an integrated temporal estimate of the total net soil redistribution rate per year since the time of the main fallout, including all erosion processes by water, wind and snow during summer and winter seasons (Meusbürger et al., 2014).

<sup>137</sup>Cs was released into the atmosphere during nuclear bomb tests and as a consequence of nuclear power plant (NPP) accidents such as Chernobyl in April 1986. It reached the land surface by dry and wet fallouts and once deposited on the ground, it is strongly bound to fine particles at the soil surface. Due to its low vertical migration rates, it moves predominantly in association with fine soil particles through physical processes, and provides an effective track of soil and sediment redistribution processes (Mabit et al., 2008). The traditional approach in using the <sup>137</sup>Cs method is based on the comparison between the inventory (total radionuclide activity per unit area) at a given sampling site and that of a so-called reference site, located in a flat and undisturbed/stable area. The method indicates the occurrence of erosion processes at sites with lower <sup>137</sup>Cs inventory as compared to the reference site, and sediment deposition processes at sites with a greater <sup>137</sup>Cs inventory (Figure 1, A). Specific mathematical conversion models allow then to derive from the latter comparison quantitative estimates of soil erosion and deposition rates (IAEA, 2014).



50 **Figure 1: Concept of the fallout radionuclide (FRN) traditional method, in which the FRN content of a reference site located in a flat and undisturbed area (R) is compared to the FRN content of disturbed sites (E and D). If the FRN at the site under**

**investigation is lower than at the reference site, the site has experienced erosion processes (E), while if the FRN content is greater than at the reference site, the site has experienced deposition processes (D).**

55 The efficacy of the method relies on an accurate selection of representative reference sites (Mabit et al., 2008; Owens and Walling, 1996, Sutherland,1996). The measured total  $^{137}\text{Cs}$  inventory at the reference sites represents the baseline fallout (i.e. reference inventory), a fundamental parameter for the qualitative and quantitative assessment of soil redistribution rates (Loughran et al., 2002). It is used for the comparison with the total  $^{137}\text{Cs}$  inventories of the sampling sites, and it therefore determines if and how strongly a site is eroding or accumulating sediments. Moreover, the depth profile of the  $^{137}\text{Cs}$  distribution in the soil at the reference site plays a very important role, as the shape of this profile is used in the conversion models to convert changes in  $^{137}\text{Cs}$  inventory changes into quantitative estimates of soil erosion rates (Walling et al., 2002).  
60 Recent studies demonstrated the sensitivity of conversion models to uncertainties or even biases in the reference inventory (e.g. Arata et al., 2016; Iurian et al., 2014; Kirchner, 2013).

A close proximity of a reference site to the area under investigation is required to meet the assumption that both experienced similar initial fallout. The latter is particularly important if the study area was strongly affected by Chernobyl fallout, which is, besides global fallout from nuclear weapons testing, the major input of  $^{137}\text{Cs}$  in many regions of Europe. Because of different geographical situations and meteorological conditions at the time of passage of the radioactive cloud, the contamination associated with Chernobyl fallout was very heterogeneous (Chawla et al., 2010, Alewell et al., 2014). Therefore, in some areas a significant small scale variability of  $^{137}\text{Cs}$  distribution may be expected and, as already pointed out by Lettner et al. (1999) and Owens and Walling (1996), might impede the comparison between reference and sampling sites. To consider adequately the spatial variability of the FRN fallout, multiple reference sites should be selected and the variability within the sites properly tackled (Kirchner, 2013, Mabit et al., 2013, Pennock and Appleby, 2002). In addition, the reference site should not have experienced any soil erosion or deposition processes since the main  $^{137}\text{Cs}$  fallout (which generally requires that it was under continuous vegetation cover such as perennial grass). Different forms of turbation, including animal-, anthropogenic- and cryoturbation or snow processes may also affect the  $^{137}\text{Cs}$  soil depth distribution at the reference site. Finally, the collection of the samples, the preparation process and the gamma analysis might introduce a certain level of uncertainty, which should be carefully considered. For instance, Lettner et al. (1999) estimated that the preparation and measuring processes contribute 12.2% to the overall variability of the reference inventory. Guidance in form of independent indicators (e.g. stable isotopes as suggested by Meusburger et al., 2013) for the suitability of reference sites might help to assist with the selection of these sites.  
75

80 All in all the suitability or unsuitability of references site is crucial, maybe even the most crucial step, in all FRN based erosion assessments. The general suitability of  $^{137}\text{Cs}$  based erosion assessment has been recently discussed very controversially (Parsons and Forster 2011, 2013; Mabit et al., 2013). We would like to propose that the FRN community needs to agree on general concepts and sampling strategies to test the suitability of reference sites in order to improve the method as well as increase the confidence in this useful erosion assessment method. Until now, the variability among spatial replicate samples at reference sites are commonly the sole criteria to decide on the suitability of a reference value. We  
85

propose an extended method to Check the Suitability of reference Sites (CheSS) using a repeated sampling strategy and, as such, an assessment of the temporal variability of reference sites. The suitability of reference sites for an accurate application of  $^{137}\text{Cs}$  as soil erosion tracer is tested at Urseren Valley (Canton Uri, Swiss Central Alps).

## 90 **2 CheSS (Check the Suitability of reference Sites): a concept to assess the suitability of reference sites for application of $^{137}\text{Cs}$ as soil erosion tracer**

### **2.1 Repeated sampling strategy and calculation of inventories**

The time period for the repeated sampling of reference sites needed for the application of  $^{137}\text{Cs}$  as soil erosion tracer will be site- and case-specific and depends on the initial small scale spatial variability and the depth distribution of the reference inventory. The time span should be of sufficient length to cause an inventory change that is larger than the uncertainty related to the inventory assessment (e.g. > 35%). In our study site being effected by anthropogenic disturbance and snow erosion of several mm per winter already 2 years can be considered sufficient (Meusburger et al., 2014). Several spatial repetitions following the suggestion of Sutherland et al. (1996) are necessary and should be analysed separately to investigate the small scale variability of  $^{137}\text{Cs}$  in the area. As we detected measurement differences between different detectors (see section 3.4), all samples should ideally be measured for  $^{137}\text{Cs}$  activity using the same analytical facilities. Finally,  $^{137}\text{Cs}$  activity needs to be decay-corrected to the same date (either the period of the first sampling campaign or the second one), considering the half-life of  $^{137}\text{Cs}$  (30.17 years).

The decay-corrected  $^{137}\text{Cs}$  activities (*act*, Bq kg<sup>-1</sup>), of each soil layer of the depth profile are converted into inventories (*inv*, Bq m<sup>-2</sup>) with the following equation:

$$Inv = act \times xm \quad (1)$$

105 where *xm* is the measured mass depth of fine soil material (<2 mm fraction) (kg m<sup>-2</sup>) of each soil sample. The depth profile of each reference site is then displayed as inventory (Bq m<sup>-2</sup>) against the depth of each layer (cm). The repeated-sampling inventory change (*Inv<sub>change</sub>*) can then be defined as:

$$Inv_{change} = \frac{Inv_{t_0} - Inv_{t_1}}{Inv_{t_0}} \times 100 \quad (2)$$

110 where *t<sub>0</sub>* and *t<sub>1</sub>* are the dates of the first and the second sampling campaigns respectively, *Inv<sub>t<sub>1</sub></sub>* is the  $^{137}\text{Cs}$  inventory (Bq m<sup>-2</sup>) at *t<sub>1</sub>*, and *Inv<sub>t<sub>0</sub></sub>* is the  $^{137}\text{Cs}$  inventory at *t<sub>0</sub>*. Positive values of *Inv<sub>change</sub>* indicate erosion, whereas negative values correspond to deposition.

### **2.2 A decision tree to assess the suitability of reference sites**

We evaluated the suitability of the reference sites by analyzing both the spatial and the temporal variability of the  $^{137}\text{Cs}$  inventory. Given the assumption that no additional deposition of  $^{137}\text{Cs}$  occurred at the sites during the investigated time

115 window (which is valid worldwide except for the areas affected by the Fukushima-Daiichi fallout, Winiarek et al., 2014, Oh et al., 2014), any temporal variation of the  $^{137}\text{Cs}$  content should be attributable to different forms of soil disturbance or to artefacts in the preparation/measurement of the samples. The potential causes of the spatial and temporal variation in the  $^{137}\text{Cs}$  total inventories and depth profiles are examined through a decision tree which includes three main nodes (Figure 2).

### **Node 1: Spatial variation of FRN total inventory**

120 First, the spatial variation of the  $^{137}\text{Cs}$  total inventory at each reference site is tested. Ideally, several replicates have been collected. If the coefficient of variation (CV) exceeds 35% as suggested by Sutherland (1996), this could be a sign of unsuitability of the reference site, although it leaves the possibility of i) increasing sampling numbers, ii) analysing the causes for the spatial variation (see CheSS A to D) and iii) moving to node 2 and 3 in CheSS.

### **Node 2: Variation of the $^{137}\text{Cs}$ depth profile**

125 Second, it is tested whether there is a significant variation between the  $^{137}\text{Cs}$  depth profiles measured as spatial or temporal (in  $t_0$  and  $t_1$ ) replicates. In theory, at a stable site, the shape of the depth profile should not change between replicates. Consequently, a regression between the FRN activity depth profiles collected as spatial or temporal replicates should follow a 1:1 line and the variability should lie within the range of the observed spatial uncertainty (node 1). A deviation of the linear regression coefficient from the 1:1 line in combination with high residues and low  $R^2$  values ( $<0.5 R^2$ ) indicates an  
130 immediate and significant change of the profile, which is typically caused by anthropogenic disturbance. For the FRN application at ploughed sites, the reference site might still be considered appropriate if the total inventory is not affected, because conversion models used for ploughed sites are less sensitive to the shape of the FRN depth distribution. For unploughed soils, again the analysis of other causes (see the causes labelled A to D in Figure 2) might help to identify the sources of this variability. Alternative options would be to take temporal replicates to evaluate the stability and thus  
135 suitability of the reference site (node 3).

### **Node 3: Temporal variation of FRN total inventory**

If the CV of all replicates taken in  $t_0$  and  $t_1$  is  $<35\%$ , the reference site might be used for the FRN method. Further a suitable test for significant differences should confirm or reject the hypothesis of  $^{137}\text{Cs}$  total inventory stability over time. If the potential causes for variation (A to D) do not apply, the site is not suitable for the traditional FRN approach. Still a repeated  
140 sampling approach could be used to assess soil redistribution rates based on FRN methods (Porto et al., 2014; Kachanoski & de Jong, 1984).




Figure 2: The CheSS decision tree to evaluate the suitability of a reference site for using <sup>137</sup>Cs as a soil erosion tracer.

145 **A: Signs of disturbance associated with erosion and deposition processes**

A variation in the <sup>137</sup>Cs depth profile may have been caused by soil movement processes affecting the site (Figure 2, A). If the site experienced a loss of soil due to erosion, we expect to observe a removal of the top soil layers of the profile

measured for instance during the second sampling campaign (Figure 3, red values below the reference profile). Further, the regression coefficient of the reference site that was affected by erosion will tend to be  $<0.9$  when plotted against a suitable reference profile or (for node 3) the reference profile before the disturbance (Figure 3). In case of deposition, a sedimentation layer should be found on the top of the reference depth profile, assuming that no ploughing operations affected the site (Figure 3, red values above the reference profile). In this case, the regression coefficient will be  $>1.1$ . Information on the depth distribution of another FRN might provide additional reliable confirmation. If redistribution processes are confirmed, the site is not suitable as a reference site and other location or a repeated FRN sampling approach to estimate erosion rates between the two sampling campaign should be considered (Kachanoski & de Jong, 1984).

reference	deposition	erosion
34	27	17.6
30	33	19.4
17.6	34	10
19.4	30	9.3
10	17.6	10.2
9.3	19.4	9.3
10.2	10	5
9.3	9.3	4.2
5	10.2	1
4.2	9.3	1





Figure 3: Signs of sheet erosion (A) and deposition (B) on a depth profile of a hypothetical reference site.

### B: Sampling or preparation artefacts

One very common artefact which might bias the comparison between the samples collected at different sites or at  $t_0$  and  $t_1$  is the difference in the skeleton content (the percentage of soil fractions  $> 2\text{mm}$ ) (Figure 2, B). The presence of stones might create preferential pathways for water as well as for very fine particles and solutes through the soil and thus influences the accumulation/migration of  $^{137}\text{Cs}$  through the soil layers. As  $^{137}\text{Cs}$  reaches the soil by the fallout from the atmosphere, the common shape of the  $^{137}\text{Cs}$  distribution along the undisturbed depth profile can be described by an exponential function, with

the highest  $^{137}\text{Cs}$  concentrations located in the uppermost soil layers (Mabit *et al.*, 2008; Walling *et al.*, 2002). This is particularly the case for soils with low skeleton content (Figure 4, A) since the presence of stones may affect  $^{137}\text{Cs}$  depth distribution either through (i) impeding the  $^{137}\text{Cs}$  downward migration ( $^{137}\text{Cs}$  activity could then be concentrated in the layer above the stone (Figure 4, B) or (ii) creating macro- and micro-pores favouring  $^{137}\text{Cs}$  associated with fine particles to “migrate” to deeper layers (Figure 4, C) or causing lateral movement, which will induce a lower  $^{137}\text{Cs}$  content in our samples.



**Figure 4: Possible influence of stones on the FRN depth distribution.**

As such, a spatial or of a temporal variation in the depth profile within the soil samples might be caused by a different skeleton content and/or by a different bulk densities among the samples. Higher bulk densities will result in higher increment inventories even if  $^{137}\text{Cs}$  activities in the layers are comparable. Thus, a thorough control (eventually through a statistical test such as a paired t test) if skeleton content and bulk densities are comparable between replicates is suggested. Finally, sampling, preparation artefacts and measurement procedures may produce various sources of errors between different sites and years. The latter is especially the case, when different persons prepared the samples. An estimation of possible errors might be considered, for example through a simulation of different increment assignments along the profile. If different detectors or different calibration sources and/or geometries are used in the two sampling campaigns, a comparability check of the measurements is advisable. For instance, a subset of samples could be measured with the two different detectors and any potential discrepancy of the results should be properly reported.

### **C: Signs of soil disturbance**

Different forms of disturbance, such as bio-, cryoturbation or even human induced soil perturbation (e.g. tillage, seed bed preparation, digging etc.) might have influenced the  $^{137}\text{Cs}$  depth distribution between different sites and  $t_0$  and  $t_1$  (Figure 2, C). Occurrence of turbation is often difficult to identify prior to sampling although it may eventually be detected by using

other tracing approaches, such as the  $\delta^{13}\text{C}$  depth distribution (Meusbürger *et al.*, 2013; Schaub and Alewell, 2009). In case of  
turbation the shape of the depth profile will be highly variable and should not be considered in the estimation of soil  
redistribution rates for unploughed soils. Nonetheless, the total inventory of  $^{137}\text{Cs}$  at a ploughed site could still be used in  
combination with simple and basic mathematical conversion models, such as the proportional model (Ritchie and McHenry,  
1990, IAEA, 2014), which require information only about the total reference inventory of  $^{137}\text{Cs}$ , and do not need detailed  
information about the  $^{137}\text{Cs}$  depth distribution.

#### **D: Signs of a heterogeneous initial fallout of $^{137}\text{Cs}$ over the area**

Finally, a significant difference between reference replicates may be caused by a high small scale spatial variability of  $^{137}\text{Cs}$   
distribution at the site, due to heterogeneous initial fallout over the study area (Figure 2, D). In Europe, significant small  
scale variability of  $^{137}\text{Cs}$  distribution is known to be due to the Chernobyl fallout, which was characterized by a high  $^{137}\text{Cs}$   
deposition associated with few rain events. Compared to the nuclear bomb tests fallout, the Chernobyl fallout was  
significantly more heterogeneous (e.g. Alewell *et al.*, 2014). Therefore, in the areas affected by the Chernobyl fallout, sites  
sampled close to each other may present very different  $^{137}\text{Cs}$  contents. It is therefore necessary to investigate the small scale  
spatial variability (e.g. the same scale as distance between reference site replicates) measured at both or at least one sampling  
campaign, looking at the CV again, as presented in the previous sections, or through a statistical test (for example the  
Analysis of the Variance, ANOVA). If the spatial variability is highly significant, the site should not be envisaged as a  
reference site for the application of the  $^{137}\text{Cs}$  method unless the number of samples collected for the determination of the  
reference baseline is large enough (at least 10) to counterweight the small scale variability within the site (Mabit *et al.*, 2012;  
Sutherland, 1996, Kirchner, 2013). A possible validation of this cause of heterogeneity might be a comparison with the  
spatial distribution of another FRN such as  $^{239+240}\text{Pu}$  or  $^{210}\text{Pb}_{\text{ex}}$  (Porto *et al.*, 2013). (Figure 2, D). As the fallout deposition of  
 $^{239+240}\text{Pu}$  after the Chernobyl accident was confined to a restricted area in the vicinity of the Nuclear Power Plant (Ketterer *et al.*,  
2004), the origin of Plutonium fallout in the rest of Europe is linked to the past nuclear bomb tests only. Consequently,  
Pu fallout distribution was more homogeneous (Alewell *et al.*, 2014; Ketterer *et al.*, 2004; Zollinger *et al.*, 2015). If the  
 $^{239+240}\text{Pu}$  depth profiles do not vary significantly between the two sampling years, there should be no disturbance (e.g.  
turbation, erosion) or measurement artefacts. As such, it might be concluded that the heterogeneous deposition of  $^{137}\text{Cs}$  at the  
time of the fallout may prevent the use of Cs at this site.

### **3 The application of the CheSS decision tree**

#### **3.1 Study area**

To test the methodology described above, we used a dataset from an alpine study area, the Urseren Valley (30 km<sup>2</sup>) in  
Central Switzerland (Canton Uri), which has an elevation ranging from 1440 to 3200 m a.s.l. At the valley bottom (1442 m  
a.s.l.), average annual air temperature for the period 1980–2012 is around  $4.1 \pm 0.7$  °C and the mean annual precipitation is

1457 ± 290 mm, with 30% falling as snow (MeteoSwiss, 2013). The U-formed valley is snow-covered from November to April. On the slopes, pasture is the dominant land use, whereas hayfields are dominant near the valley bottom.

### 3.2 Sampling design

220 Supportive information was provided by the local landowners to select the reference sites in both valleys. Sites used for ploughing and grazing activities were excluded. A first sampling campaign was undertaken in autumn 2010 for  $^{239+240}\text{Pu}$  and 2013 for  $^{137}\text{Cs}$  measurements. Six reference sites (REF1 to REF6) were identified in flat and undisturbed areas along the valley. At each site 3 cores (40 cm depth), collected 1 m apart from each other, were sampled. The cores were cut into 3 cm increments, to derive the  $^{137}\text{Cs}$  depth profile. The three cores from each site were bulked to provide one composite sample  
225 per site. During the second sampling campaign in spring 2015, all six reference sites were resampled. Considering the typical and high soil redistribution dynamics of the valley of >1cm per year caused by snow-induced soil removal (Meusburger et al., 2014), the period considered is sufficiently long to ensure the possibility to observe changes in the depth profiles if soil erosion and deposition processes affected the area. At each site, we collected three replicates, which were analyzed separately, to investigate the small scale variability of the FRN content. All cores were air-dried (40°C for 72h), sieved (<2  
230 mm) to remove coarse particles and to determine the skeleton content. The bulk density (BD) was also determined.

### 3.3 Measurement of anthropogenic FRN activities and inventories

The  $^{137}\text{Cs}$  activity measurements ( $\text{Bq kg}^{-1}$ ) were performed with high resolution HPGe detectors. The  $^{137}\text{Cs}$  activity ( $\text{Bq kg}^{-1}$ ) from the samples collected in 2013 were analysed at the Institute of Physics of the University of Basel using a coaxial, high resolution germanium lithium detector (Princeton Gammatech) with a relative efficiency of 19% (at 1.33 MeV,  $^{60}\text{Co}$ ).  
235 Counting time was set to 24 hours per sample. Samples collected in 2015 were analysed at the state laboratory Basel-City using coaxial high resolution germanium detectors having 25% to 50% relative efficiencies (at 1.33 MeV,  $^{60}\text{Co}$ ). Counting times were set to provide a precision of less than ±10% for  $^{137}\text{Cs}$  at the 95% level of confidence.

All soil samples were counted in sealed discs (65 mm diameter, 12 mm height, 32 cm<sup>3</sup>) and the measurements were corrected for sample density and potential radioactivity background. The detectors located at the state laboratory Basel-City  
240 were calibrated with a reference solution of the same geometry. The reference contained  $^{152}\text{Eu}$  and  $^{241}\text{Am}$  (2.6 kBq resp. 7.7 kBq) to calibrate the detectors from 60 to 1765 keV. It was obtained from the Czech Metrology Institute, Prague. This solution was bound in a silicon resin with a density of 1.0. The efficiency functions were corrected for coincidence summing of the  $^{152}\text{Eu}$  lines using a Monte Carlo simulation program (Gespecor). The  $^{137}\text{Cs}$  was counted at 662 keV with an emission probability of 0.85 and a (detector) resolution of 1.3 to 1.6 keV (FWHM). All measurements and calculations were  
245 performed with the gamma software Interwinner 7. The  $^{137}\text{Cs}$  activity measurements were all decay-corrected to the year 2015.

To compare the  $^{137}\text{Cs}$  results with those obtained with another artificial FRN, all samples were also analyzed for  $^{239+240}\text{Pu}$  activity. The determination of Plutonium isotopes from both valleys and for both sampling years were performed using a

Thermo X Series II quadrupole ICP-MS at the Northern Arizona University, USA. Detailed description of the ICP-MS specifications and sample preparation procedure can be found in Alewell *et al.*, 2014. The activities of  $^{137}\text{Cs}$  and  $^{239+240}\text{Pu}$  ( $\text{act}$ ,  $\text{Bq kg}^{-1}$ ) were converted into inventories ( $\text{Bq m}^{-2}$ ) according to equation (1).

### 3.4 Application of the CheSS decision support tool to the reference sites

Because the  $^{137}\text{Cs}$  activity of the samples was measured with different detectors for the two sampling years, we investigated the potential variability between the results obtained from these two detectors. A selected subset of samples ( $n= 24$ ) was analysed using both detectors (i.e. the one located at the Institute of Physics of the University of Basel and the other located at the State Laboratory Basel-City). The results highlight a high correspondence of the measurements obtained with the two analytical systems ( $R^2 = 0.97$ ;  $p < 0.005$ ), although the detector of the State Laboratory Basel-City returns slightly lower  $^{137}\text{Cs}$  activities (Figure 5). Thus, the  $^{137}\text{Cs}$  activities of the samples measured in 2013 were corrected to be comparable to those obtained with the detector of State Laboratory Basel-City (higher efficiency) to allow comparability between the different data sets.





Figure 5: The comparison between the  $^{137}\text{Cs}$  measurements of a subset of samples ( $n=16$ ) performed with two different HpGe detectors, where detector 1: detector hosted at the Physics department of the University of Basel (CH) and detector 2: detector hosted at the State-Laboratory of Basel (CH).

Total  $^{137}\text{Cs}$  inventories (decay-corrected to 2015) of the six reference sites collected in the Urseren Valley in 2013 ranged from 3858 to 5057  $\text{Bq m}^{-2}$ , with a mean value of 4515  $\text{Bq m}^{-2}$  and a standard deviation (SD) of 468  $\text{Bq m}^{-2}$ . Data from 2015 ranged between 3925 to 8619  $\text{Bq m}^{-2}$ , with a mean value of 5701  $\text{Bq m}^{-2}$  and a SD of 1730  $\text{Bq m}^{-2}$  (Figure 6).




**Figure 6: Temporal variation between the total <sup>137</sup>Cs inventories measured at the reference sites in the Urseren Valley, where Time 0 = 2013 and Time 1 = 2015. The errors bars indicate the standard deviations of the inventories among the replicates collected at each reference site in 2015.**

When following the CheSS decision tree, we investigated the variation in the <sup>137</sup>Cs total inventories at each reference site (node 1). The replicate samples were analyzed separately only during the second sampling campaign (t1), while during the first sampling campaign (t0) only composite samples were analysed. Reference sites 3, 5 and 6 presented signs of high small scale variability, as expressed by a CV of 48 % (Table 1). Such variability excluded them from any further use as reference sites without subsequent additional sampling. For reference sites 1, 2 and 4, the CV was between 19 – 31%.


Passing to node 2 of the CheSS decision tree, the analysis focused on the variation of the shape of the <sup>137</sup>Cs depth profile (Figure 7). Here, we examined the regression between the reference depth profiles in t0 and t1. For the three sites with acceptable spatial variability (i.e. reference site 1, 2 and 4) the site REF4 showed sign of deposition with a regression coefficient between t0 and t1 = 1.34. The deposition was confirmed by field observation of construction works that were conducted between the two sampling campaigns. Thus, after this disturbance, the site is not a suitable reference site anymore. Among the sites with high spatial variability REF6 showed signs of erosion with a regression coefficient between t0 and t1 = 0.79.

In node 3, the temporal differences in total inventories between t0 and t1 were assessed. Here, only REF4 showed a significant difference of the total <sup>137</sup>Cs inventories between t0 and t1. This confirmed the unsuitability of the site to provide a reference value after the construction works.



290 **Figure 7: The  $^{137}\text{Cs}$  depth profiles of the six investigated reference sites in the Urseren Valley for the two different sampling campaigns. The errors bars indicate the standard deviations of the inventories among the replicates collected at each reference site in 2015. The regression equation between the depth profile at t0 and t1 is displayed together with the  $R^2$ .**

To further investigate the causes for the spatial variation,  $^{239+240}\text{Pu}$  inventories measured at the three replicates of each site were analysed for t0 = 2010 and t1 = 2015 (Figure 8). Clearly, deposition for REF4 and erosion processes for REF6 were confirmed with an increase of 46% and a decrease of 27% in the total  $^{239+240}\text{Pu}$  inventory between t0 and t1, respectively.



**Figure 8: Temporal variation between the total <sup>239+240</sup>Pu inventories measured at the reference sites in the Urseren Valley, where Time 0 = 2010 and Time 1 = 2015. The errors bars indicate the standard deviations of the inventories among the replicates collected at each reference site in 2015.**

300

Further, the depth profiles of the three replicates at reference site 1 presented also significant differences. We then looked at the differences in the skeleton content of the three replicates (Figure 2, B). An ANOVA test showed a significant difference (p-value of 0.025), which indicates that a difference in the presence of stones in the three soil cores might have affected the FRN depth distribution. In particular, a Tukey's HSD (Honest Significant Difference) Post-hoc pairwise comparison identified the replicate number 3 at REF1 as a potential outlier. To validate the suitability of REF1 as a reference site, more replicates should be collected and measured, in order to compare their <sup>137</sup>Cs depth profiles to those results obtained during the first sampling campaign. In summary, REF2, REF4 (before the construction works) appeared to be most suitable for <sup>137</sup>Cs-based studies. As for sites REF3 and REF5, a visual inspection of their soil profiles excluded that any soil disturbance

affected the sites (Figure 2, C). Consequently, at those sites, the variation in their depth profiles is due to a heterogeneous  
315 fallout with high spatial variability (Figure 2, D).. These sites may be suitable for studies based on other FRNs or even on  
 $^{137}\text{Cs}$  if more samples are collected to constrain the spatial heterogeneity that was introduced by the  $^{137}\text{Cs}$  Chernobyl fallout.

#### 4 Conclusions

With the decision tree CheSS, a support tool was proposed to verify the suitability of reference sites for a  $^{137}\text{Cs}$  based soil  
315 erosion assessment. Great attention has to be given to the analysis of the small scale variability of  $^{137}\text{Cs}$  distribution in the  
reference areas, especially in those regions affected by heterogeneous post-accidental nuclear fallout. To cope with a small  
scale variability, sampling numbers might be increased, or the temporal variation of  $^{137}\text{Cs}$  or another radionuclide, such as  
 $^{239+240}\text{Pu}$ , might be analysed. The CheSS test in the Urseren Valley indicated that the heterogeneity and disturbance of  $^{137}\text{Cs}$   
distribution prejudiced the suitability of some reference sites. In addition, the presence of stones affected the shapes of the  
320 depth profile in at least one replicate sample. Including unsuitable reference sites, the application of the traditional  $^{137}\text{Cs}$   
approach, based on a spatial comparison between reference and sampling sites, is compromised. To derive soil redistribution  
rates, a  $^{137}\text{Cs}$  repeated sampling approach should be preferred. This approach is based on a temporal comparison of the FRN  
inventories measured at the same site in different times (Kachanoski & de Jong, 1984). It doesn't require the selection of  
reference sites, because the inventory documented by the initial sampling campaign is used as the reference inventory for  
325 that point (Porto *et al.*, 2014).

Accurate soil erosion assessments are crucially needed to validate soil erosion modelling, which can help prevent and  
mitigate soil losses on larger spatial scales. In this context, FRN could play a decisive role, if we are able to overcome its  
potential pitfalls, especially those related to the selection of suitable reference sites. The decision tree CheSS provides a new  
concept for an objective and comparable reference site testing, which enables to exclude those sites which present signs of  
330 uncertainty and should therefore not be used as reference sites. With this approach we are convinced to contribute improving  
the reliability of the FRN-based soil erosion assessments.

#### Authors contributions.

L. Arata, K. Meusburger, L. Mabit and C. Alewell designed the concept of the method and analysed the data. A. Bürge  
contributed to the collection and preparation of the soil samples, and to the analysis of the data. M. Zehringer measured the  
335  $^{137}\text{Cs}$  activity of the soil samples and analysed the results. M. E. Ketterer measured the  $^{239+240}\text{Pu}$  activity of the soil samples.  
L. Arata prepared the manuscript with contributions from all co-authors.

## Acknowledgements

The authors would like to thank Annette Ramp, Gregor Juretzko, Simon Tresch, Carmelo La Spada and Axel Birkholz for support during field work. This work was financially supported by the Swiss National Science Foundation (SNF), project no. 200021-146018, and has been finalized in the framework of the IAEA Coordinated Research Project (CRP) on “*Nuclear techniques for a better understanding of the impact of climate change on soil erosion in upland agro-ecosystems*” (D1.50.17).

## References

- 345 Alewell, C., Egli, M., & Meusbürger, K. 2015. An attempt to estimate tolerable soil erosion rates by matching soil formation with denudation in Alpine grasslands. *Journal of Soils and Sediments*, 15(6), 1383-1399.
- Alewell, C., Meusbürger, K., Juretzko, G., Mabit, L., & Ketterer, M. E. (2014). Suitability of  $^{239+240}\text{Pu}$  and  $^{137}\text{Cs}$  as tracers for soil erosion assessment in mountain grasslands. *Chemosphere*, 103, 274-280.
- Arata, L., Meusbürger, K., Frenkel, E., A'Campo-Neuen, A., Iuran, A.R., Ketterer, M.E., Mabit, L., Alewell, C., 2016b.
- 350 Modelling Deposition and Erosion rates with RadioNuclides (MODERN) - Part 2: A comparison of different models to convert  $^{239+240}\text{Pu}$  inventories into soil redistribution rates at unploughed sites. *Journal of environmental radioactivity*, 162, 97-106.
- Chawla, F., Steinmann, P., Pfeifer, H.R., Froidevaux, P., 2010. Atmospheric deposition and migration of artificial radionuclides in Alpine soils (Val Piora, Switzerland) compared to the distribution of selected major and trace elements.
- 355 *Science of the Total Environment*, 408(16), 3292–3302.
- Descroix, L., & Mathys, N., 2003. Processes, spatio-temporal factors and measurements of current erosion in the French southern Alps: a review. *Earth Surface Processes and Landforms*, 28(9), 993-1011.
- Frankenberg, P., Geier, B., Proswitz, E., Schütz, J., & Seeling, S., 1995. Untersuchungen zu Bodenerosion und Massenbewegungen im Gunzesrieder Tal/Oberallgäu. *Forstwissenschaftliches Centralblatt vereinigt mit Tharandter*
- 360 *forstliches Jahrbuch*, 114(1), 214-231.
- IAEA (International Atomic Energy Agency), 2014. Guidelines for using Fallout radionuclides to assess erosion and effectiveness of soil conservation strategies. IAEA-TECDOC-1741. IAEA publication. Vienna, Austria. 213 p.
- Iurian, A. R., Mabit, L., Cosma, C., 2014. Uncertainty related to input parameters of  $^{137}\text{Cs}$  soil redistribution model for undisturbed fields. *Journal of environmental radioactivity*, 136, 112-120.
- 365 Kachanoski, R. G., & De Jong, E., 1984. Predicting the temporal relationship between soil cesium-137 and erosion rate. *Journal of Environmental Quality*, 13(2), 301-304.
- Ketterer, M.E., Hafer, K.M., Jones, V.J., Appleby, P.G., 2004. Rapid dating of recent sediments in Loch Ness: ICPMS measurements of global fallout Pu. *Science of the Total Environment*, 322, 221–229.

- Kirchner, G., 2013. Establishing reference inventories of  $^{137}\text{Cs}$  for soil erosion studies: methodological aspects. *Geoderma* 211, 107–115.
- Konz, N., Prasuhn, V., Alewell, C., 2012. On the measurement of alpine soil erosion. *Catena*, 91, 63-71.
- Lal, R., 2003. Soil erosion and the global carbon budget. *Environ. Int.* 29, 437–450.
- Lettner, H., Bossew, P., Hubmer, A.K., 1999. Spatial variability of fallout Caesium-137 in Austrian alpine regions. *J. Environ. Radioact.* 47, 71–82.
- 375 Loughran, R.J., Pennock, D.J., Walling, D.E., 2002. Spatial distribution of caesium-137, in: *Handbook for the Assessment of Soil Erosion and Sedimentation Using Environmental Radionuclides*. Springer, pp. 97–109.
- Mabit, L., Benmansour, M., Walling, D. E., 2008. Comparative advantages and limitations of the fallout radionuclides  $^{137}\text{Cs}$ ,  $^{210}\text{Pb}_{\text{ex}}$  and  $^7\text{Be}$  for assessing soil erosion and sedimentation. *Journal of Environmental Radioactivity*, 99(12), 1799-1807.
- Mabit, L., Chhem-Kieth, S., Toloza, A., Vanwalleghem, T., Bernard, C., Amate, J.I., de Molina, M.G., Gómez, J.A., 2012.
- 380 Radioisotopic and physicochemical background indicators to assess soil degradation affecting olive orchards in southern Spain. *Agric. Ecosyst. Environ.* 159, 70–80.
- Mabit, L., Meusburger, K., Fulajtar, E., Alewell, C., 2013. The usefulness of  $^{137}\text{Cs}$  as a tracer for soil erosion assessment: A critical reply to Parsons and Foster (2011). *Earth-Science Reviews*, 127, 300–307.
- Meusburger, K., Leitinger, G., Mabit, L., Mueller, M.H., Walter, A., Alewell, C., 2014. Soil erosion by snow gliding—a first
- 385 quantification attempt in a subalpine area in Switzerland. *Hydrology and Earth System Sciences*, 18(9), 3763–3775.
- Meusburger, K., Mabit, L., Park, J.H., Sandor, T., Alewell, C., 2013. Combined use of stable isotopes and fallout radionuclides as soil erosion indicators in a forested mountain site, South Korea. *Biogeosciences* 10, 5627–5638.
- Oh, J. S., Lee, S. H., Choi, J. K., Lee, J. M., Lee, K. B., & Park, T. S. (2014). Atmospheric input of  $^{137}\text{Cs}$  and  $^{239,240}\text{Pu}$  isotopes in Korea after the Fukushima nuclear power plant accident. *Applied Radiation and Isotopes*, 87, 53-56.
- 390 Owens, P.N., Walling, D.E., 1996. Spatial variability of caesium-137 inventories at reference sites: an example from two contrasting sites in England and Zimbabwe. *Appl. Radiat. Isot.* 47, 699–707.
- Parsons, A. J. and I. D. L. Foster, 2011. "What can we learn about soil erosion from the use of  $^{137}\text{Cs}$ ?" *Earth Science Reviews* 108: 13.
- Pennock, D.J., Appleby, P.G., 2002. Site selection and sampling design, in: *Handbook for the Assessment of Soil Erosion and Sedimentation Using Environmental Radionuclides*. Springer, pp. 15–40.
- 395 Porto, P., Walling, D.E., Alewell, C., Callegari, G., Mabit, L., Mallimo, N., Meusburger, K., Zehring, M., 2014. Use of a  $^{137}\text{Cs}$  re-sampling technique to investigate temporal changes in soil erosion and sediment mobilisation for a small forested catchment in southern Italy. *J. Environ. Radioact.* 138, 137–148.
- Porto P, Walling DE, Callegari G., 2013. Using  $^{137}\text{Cs}$  and  $^{210}\text{Pb}_{\text{ex}}$  measurements to investigate the sediment budget of a small
- 400 forested catchment in southern Italy. *Hydrological Processes*; 27: 795-806.

- Ritchie, J. C., Ritchie, C. A., 2001. Bibliography of publications of Cesium-137 studies related to erosion and sediment deposition. USDA-ARS Hydrology and Remote Sensing Laboratory.
- Schaub, M., Alewell, C., 2009. Stable carbon isotopes as an indicator for soil degradation in an alpine environment (Urseren Valley, Switzerland). *Rapid Commun. Mass Spectrom.* 23, 1499–1507.
- 405 Sutherland, R.A., 1996. Caesium-137 soil sampling and inventory variability in reference locations: A literature survey. *Hydrol. Process.* 10, 43–53.
- Walling, D.E., He, Q., Appleby, P.G., 2002. Conversion models for use in soil-erosion, soil-redistribution and sedimentation investigations. In: Zapata, F. (Ed.) *Handbook for the assessment of soil erosion and sedimentation using environmental radionuclides*. Kluwer, Dordrecht. Netherlands. pp. 111–164.
- 410 Winiarek, V., Bocquet, M., Duhanyan, N., Roustan, Y., Saunier, O., & Mathieu, A. (2014). Estimation of the caesium-137 source term from the Fukushima Daiichi nuclear power plant using a consistent joint assimilation of air concentration and deposition observations. *Atmospheric environment*, 82, 268-279.
- Zapata, F. (Ed.), 2002. *Handbook for the assessment of soil erosion and sedimentation using environmental radionuclides* (Vol. 219). Dordrecht: Kluwer Academic Publishers.
- Zollinger, B., Alewell, C., Kneisel, C., Meusbürger, K., Brandová, D., Kubik, P., Schaller M., Ketterer M., Egli, M., 2014.
- 415 The effect of permafrost on time-split soil erosion using radionuclides ( $^{137}\text{Cs}$ ,  $^{239+240}\text{Pu}$ , meteoric  $^{10}\text{Be}$ ) and stable isotopes ( $\delta^{13}\text{C}$ ) in the eastern Swiss Alps. *Journal of Soils and Sediments*, 15(6), 1400–1419.