



1 **Soil organic carbon stocks are systematically overestimated by**
2 **misuse of the parameters bulk density and stone content**

3

4 Christopher Poeplau, Cora Vos, Axel Don

5 Thünen Institute of Climate-Smart Agriculture, Bundesallee 50, 38116 Braunschweig, Germany

6

7 *Correspondence to:* Christopher Poeplau, christopher.poeplau@thuenen.de

8

9 Key words: Fine soil stock, soil skeleton, rock content, bulk density estimation

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25



26

27 **Abstract** Estimation of soil organic carbon (SOC) stocks requires estimates of the carbon content, bulk density,
28 stone content and depth of a respective soil layer. However, different application of these parameters could
29 introduce a considerable bias. Here, we explain why three out of four frequently applied methods overestimate
30 SOC stocks. In stone rich soils (>30 Vol. %), SOC stocks could be overestimated by more than 100%, as
31 revealed by using German Agricultural Soil Inventory data. Due to relatively low stone content, the mean
32 systematic overestimation for German agricultural soils was 2.1-10.1% for three different commonly used
33 equations. The equation ensemble as re-formulated here might help to unify SOC stock determination and avoid
34 overestimation in future studies.

35

36 1 Introduction

37 Size and changes in the soil organic carbon (SOC) pool are major uncertainties in global earth system models
38 used for climate predictions. Accurate estimation of SOC stocks is vital to understanding the links between
39 atmospheric and terrestrial carbon (Friedlingstein et al., 2014). Estimates of global SOC stocks are based on soil
40 inventories from regional to continental scale, involving multiplication of measured carbon content by soil bulk
41 density (BD, oven-dry mass of soil per unit volume) and the depth of the respective soil layer (Batjes, 1996). The
42 content of elements such as carbon and nitrogen in soils is usually determined in an aliquot sample of the fine
43 soil, which is defined as the part of the soil that passes through a 2 mm sieve (Corti et al., 1998). Mineral
44 fragments >2mm (gravel and stones) are considered free of SOC (Perruchoud et al., 2000), although this may not
45 be completely true as shown by Corti et al. (2002). Furthermore, living root fragments >2mm are not considered
46 part of SOC, but usually as part of plant biomass. It is thus widely accepted that accurate estimates of SOC
47 stocks should account in some way for the presence of gravel, stones (coarse soil, in the following only referred
48 to as stones) and roots (Rytter, 2012; Throop et al., 2012).

49 The accuracy of SOC estimates depends in the first instance on the available data and their quality. Soil organic
50 carbon content of the fine soil is usually measured with high throughput and precision in elemental analyzers,
51 while BD and stone content are often only assessed in plot scale studies due to much more elaborate sampling
52 requirements (Don et al., 2007). In regional scale studies or national soil inventories, BD is therefore often
53 approximated using pedotransfer functions and the fraction of stones is often ignored (Wiesmeier et al., 2012).
54 Stoniness is therefore regarded as the greatest uncertainty in SOC stock estimates (IPCC, 2003). However, even
55 when all parameters are recorded, considerable difference in SOC stocks can arise from varying use of the
56 parameters in equations. Apart from the methodological bias caused by using different methods for determining
57 BD and stone content (Beem-Miller et al., 2016; Blake, 1965), the different calculation approaches could lead to
58 systematically different SOC stock estimates if soils contain stones. Several of the approaches commonly used to
59 calculate SOC stocks are not correct and inflate SOC stocks. The aim of this study was i) to reveal the
60 conceptual differences in widely used methods for SOC stock calculation, ii) to quantify the methodological bias
61 in SOC stocks in a regional scale soil inventory, iii) to identify the most affected soil layers and finally iv) to
62 suggest the most adequate method for unified and unbiased SOC stock calculation.



63

64

65 2 Materials and Methods

66 In a preliminary literature review we selected a total of 100 publications for which the method used to calculate
67 SOC stocks was recorded. The search was restricted to publications listed in ISI Web of Knowledge, where ‘soil
68 carbon stocks’ was used as the search term. We ordered the 4915 search results by ‘relevance’, excluded reviews
69 and modeling studies and avoided redundant senior authors (Tab. S1). In the literature we identified four
70 different methods where identified, which vary in use of the parameters BD and stone content (Henkner et al.,
71 2016; Lozano-García and Parras-Alcántara, 2013; Poeplau and Don, 2013; Wang and Dalal, 2006):

72 In method one (M1), a certain volume of soil is sampled, dried and weighed to determine BD. Thereby, no
73 separation into fine soil and coarse soil (gravel, stones, roots) fraction is made, while C concentration is
74 determined in a sieved fine soil sample (usually <2 mm). Soil organic carbon stocks are then calculated as
75 follows:

76 M1:

$$77 \quad BD_{sample} = \frac{mass_{sample}}{volume_{sample}} \quad (\text{Eq. 1})$$

$$78 \quad SOCstock_i = SOCcon_{fine\ soil} \times BD_{sample} \times depth_i \quad (\text{Eq. 2})$$

79 where BD_{sample} is the bulk density of the total sample, $SOCstock_i$ is the SOC stock of the investigated soil layer
80 (i) [Mg ha^{-1}], $SOCcon_{fine\ soil}$ is the content of SOC in the fine soil [%] and $depth_i$ is the depth of the respective
81 soil layer [cm]. This method leads to biased SOC stocks estimates due to inadequate representation of the stones
82 as almost SOC free mass. In method two (M2), a certain volume of soil is sampled, dried and weighed.
83 However, after sieving, the mass and volume of stones and coarse roots are determined. In the following, we
84 simplify the equations by omitting coarse roots, which is also ‘common practice’, although the volume occupied
85 by roots can be considerably high. This source of error is not further discussed in this study. By approximating a
86 stone density (ρ_{stones}) of 2.6 g cm^{-3} (root density is usually assumed to be close to 1 g cm^{-3}), BD of the fine soil
87 is subsequently calculated as:

88 M2:

$$89 \quad BD_{fine\ soil} = \frac{mass_{sample} - mass_{stones}}{volume_{sample} - \frac{mass_{stones}}{\rho_{stones}}} \quad (\text{Eq. 3})$$

$$90 \quad SOCstock_i = SOCcon_{fine\ soil} \times BD_{fine\ soil} \times depth_i \quad (\text{Eq. 4})$$

91 Thus in M2, coarse soil content is accounted for in equation (3), not in equation (4). The opposite is true for the
92 next method (M3), in which the *stone fraction* [dimensionless] is determined, but only applied to reduce the soil
93 volume (Eq. 6), and not to determine $BD_{fine\ soil}$:

94 M3:



95 $BD_{sample} = \frac{mass_{sample}}{volume_{sample}}$ (Eq. 1)

96 $SOCstock_i = SOCcon_{fine\ soil} \times BD_{sample} \times depth_i \times (1 - stone\ fraction)$ (Eq. 5)

97 In method four (M4), the coarse soil fraction is accounted for in both equations, i.e. to calculate $BD_{fine\ soil}$ (Eq.
98 3) and the volume of the fine soil (Eq. 3)

99 M4:

100 $BD_{fine\ soil} = \frac{mass_{sample} - mass_{stones}}{volume_{sample} - \frac{mass_{stones}}{\rho_{stones}}}$ (Eq. 3)

101 $SOCstock_i = SOCcon_{fine\ soil} \times BD_{sample} \times depth_i \times (1 - stone\ fraction)$ (Eq. 5)

102 In the German Agricultural Soil Inventory, more than 3000 agricultural soils (cropland and grassland) have been
103 sampled as described by Grüneberg et al. (2014). To date, a total of 2515 sites were sampled and analysed for all
104 relevant parameters (stone content, fine soil mass, carbon content of the fine soil) in five different depth
105 increments: 0-10, 10-30, 30-50, 50-70 and 70-100 cm. Here, we considered only mineral soils with a SOC
106 content <8.7% giving a total of 2350 sites and 11,514 soil samples. We expected the strongest effects in soils
107 with high stoniness and therefore stratified the dataset by stone content [vol. %]. Statistical analyses were not
108 conducted, since a systematic deviation implied significant difference between all methods. Data analysis and
109 plotting was performed in the R 3.1.2 environment (R Development Core Team, 2010).

110

111 3 Results and Discussion

112 3.1 Bias of three calculation methods to estimate SOC stocks

113 Three out of the four SOC calculation methods produced systematically overestimated SOC stocks. These
114 deviations are systematic errors (bias) that cannot be reduced with optimised methods to determine the
115 parameters SOC content, BD and stone content but reduce the accuracy of SOC stock estimates. As expected,
116 the differences in SOC stocks between calculation methods increased with stone content (Fig. 1). This is in line
117 with findings by Rytter (2012) that the method of BD estimation is most important in very stony soils. While
118 differences between methods for soils with a stone content of less than 5 vol. % were small to almost negligible,
119 M1-M3 deviated strongly from M4 in soils with >30% stones (Fig. 1, Tab. 1). Since M4 is the closest
120 approximation to reality, the systematic bias was expressed as relative deviation from M4 (Tab. 1). In soils with
121 >30% stones, M1 caused the highest bias of all three calculation methods, overestimating SOC stocks by on
122 average 144%, i.e. more than doubling the real SOC stocks. Methods M2 and M3 also produced biased SOC
123 stocks with 98% and 21% overestimations for the highest stone content class (>30% stone content). Using an
124 average $BD_{fine\ soil}$ of 1.2, we plotted the deviation from M4 as a function of volumetric stone content for M1-
125 M3 (Fig. 2). Thereby, M1 and M2 showed exponential responses, while M3 showed a linear response. These
126 responses would increase with decreasing bulk density of the fine soil. The literature review revealed that M1,
127 M2, M3 and M4 were used by 52, 5, 36, and 7 studies respectively. Thus, in 93% of all studies reviewed, SOC



128 stocks were systematically overestimated assuming a stone fraction >0 . More than half of the studies reviewed
129 did not account for the stone fraction at all.

130 The number of soils with high stone contents in the German dataset is limited due to the dominance of parent
131 material from glacio-fluvial deposits (Tab. 1). Thus, the majority of soils (67-78%, depending on soil depth
132 increment) had a volumetric stone content of $<5\%$. As a consequence, the average SOC stocks were only
133 moderately influenced by the calculation method (2.1-10.1% deviation, Tab. 1). For forests, which are usually
134 found on soils less suitable for agriculture, e.g. due to high stoniness, the bias would be stronger. Overall, the
135 results highlight the importance of a correct use of the parameters BD and stone fraction when calculating SOC
136 stocks.

137 3.2 Evaluation of the four different calculation methods

138 Since all four methods use the same $SOC_{con_{fine\ soil}}$ due to equal preparation of the fine soil, differences
139 between the calculation methods arise from differences in use of the parameters BD and stone content. The
140 individual bias of each method is visualized in Figure 3. In M1, BD of the soil containing SOC (fine soil) is
141 overestimated due to inclusion of stones in the BD estimate. The mass of fine soil present in the respective soil
142 layer is also overestimated, since the stone fraction is not subtracted from the soil volume with which BD_{sample}
143 is multiplied (Eq. 2). Thus, M1 ‘fills’ the space occupied by stones with fine soil with an overestimated BD. In
144 the German Agricultural Soil Inventory, only 9% of all sampled layers were found to be free of stones. Thus, for
145 most soils M1 is not the correct way to calculate SOC stocks. Similarly, M2 overestimates SOC_{stock_i} by filling
146 the volume of stones with fine soil. However, BD is calculated and used correctly leading to a smaller systemic
147 overestimation of SOC compared to M1. Finally, M3 correctly accounts for the stone fraction that can be
148 assumed to be SOC free. However, in M3 an overestimated BD is applied as in M1, i.e. BD_{sample} and not the
149 $BD_{fine\ soil}$. Methods to estimate BD and stone content vary, primarily owing to size and abundance of the latter
150 and may have large uncertainty (Blake, 1965; Parfitt et al., 2010; Rytter, 2012). However, the presented
151 difference between calculation methods is independent of the method of determination of these parameters
152 with one exception: If the sampled soil layer contains no gravel, but only fine soil and stones that exceed the
153 diameter of a soil ring used to determine BD_{sample} , and this ring is placed at a position (in the profile wall)
154 which is completely free of stones, while the stone content is estimated with a different method and accounted
155 for, then M3 does resemble M4. Bulk density is often determined with soil rings with a volume between 100 and
156 500 cm³ or soil probes (Walter et al., 2016). In the German Agricultural Soil Inventory, 250 cm³ soil rings are
157 used to determine BD. In 89% of all soils inventoried, small stones were detected which end up in the soil ring
158 and have to be corrected for. Thus, method M3 is rarely a correct method to estimate SC stocks. It is erroneously
159 often cited as the IPCC default method. However, while the equations given in IPCC resemble M3, IPCC
160 provides a footnote that is most likely often overlooked, which states that BD estimates should be corrected for
161 the proportion of ‘coarse fragments’ (IPCC, 2003). Even if the stone fraction might store a certain amount of
162 organic carbon (Corti et al., 2002), which might lead to slight underestimation of SOC stocks in M4, we suggest
163 use of this method in future studies.

164 3.3 Proposed equations to calculate SOC stocks



165 Bulk density might be of interest as an important soil property. However, for the calculation of SOC stocks alone
166 it is not needed, while it is the fine soil stock of the investigated soil layer (FSS_i , Mg ha^{-1}) that is of interest since
167 it contains the SOC. Thus, the equations in M4 could be reformulated as:

$$168 \quad FSS_i = \frac{\text{mass}_{\text{fine soil}}}{\text{volume}_{\text{sample}}} \times \text{depth}_i \quad (\text{Eq. 6})$$

$$169 \quad \text{SOC}_{\text{stock}_i} = \text{SOC}_{\text{con}_{\text{fine soil}}} \times FSS_i \quad (\text{Eq. 7})$$

170 This has implications for sample preparation: For $BD_{\text{fine soil}}$ the volume of coarse fragments has to be estimated
171 by weighing stones and coarse roots separately, while FSS_i would only need the total mass of the fine soil
172 contained in the known volume of sample. When using the probe method, SOC stock calculation can thus be
173 simplified, while FSS_i is slightly more complicated to calculate when the profile pit method (determination of
174 stones within and outside the soil ring) is used (Grüneberg et al., 2014).

175

176 **4 Conclusions**

177 We show here that substantially different methods are used for the calculation of SOC stocks. These methods
178 differ in use of the parameters bulk density and coarse soil fraction, which causes systematic overestimation of
179 SOC stocks in three out of four, more or less frequently applied methods, or in 93 of 100 publications reviewed.
180 We showed that this overestimation can exceed 100% in stone-rich soils. For future studies, we suggest to
181 calculate the fine soil stock of a certain soil layer which is to be multiplied with its SOC content to derive
182 unbiased SOC stock estimates.

183

184 **Acknowledgements**

185 This study was funded by the German Federal Ministry of Food and Agriculture in the framework of the German
186 Agricultural Soil Inventory.

187

188

189

190

191

192

193

194

195 **References**

- 196 Batjes, N.: Total carbon and nitrogen in the soils of the world, *Eur. J. Soil Sci.*, 47, 151-163, 1996.
- 197 Beem-Miller, J. P., Kong, A. Y. Y., Ogle, S., and Wolfe, D.: Sampling for Soil Carbon Stock Assessment in
198 Rocky Agricultural Soils, *Soil Sci. Soc. Am. J.*, 80, 1411-1423, 2016.
- 199 Blake, G.: Bulk density, *Methods of Soil Analysis. Part 1. Physical and Mineralogical Properties, Including*
200 *Statistics of Measurement and Sampling*, 1965. 374-390, 1965.
- 201 Corti, G., Ugolini, F., Agnelli, A., Certini, G., Cuniglio, R., Berna, F., and Fernández Sanjurjo, M.: The soil
202 skeleton, a forgotten pool of carbon and nitrogen in soil, *Eur. J. Soil Sci.*, 53, 283-298, 2002.
- 203 Corti, G., Ugolini, F. C., and Agnelli, A.: Classing the Soil Skeleton (Greater than Two Millimeters): Proposed
204 Approach and Procedure, *Soil Sci. Soc. Am. J.*, 62, 1620-1629, 1998.
- 205 Don, A., Schumacher, J., Scherer-Lorenzen, M., Scholten, T., and Schulze, E. D.: Spatial and vertical variation
206 of soil carbon at two grassland sites - Implications for measuring soil carbon stocks, *Geoderma*, 141, 272-282,
207 2007.
- 208 Friedlingstein, P., Meinshausen, M., Arora, V. K., Jones, C. D., Anav, A., Liddicoat, S. K., and Knutti, R.:
209 Uncertainties in CMIP5 Climate Projections due to Carbon Cycle Feedbacks, *Journal of Climate*, 27, 511-526,
210 2014.
- 211 Grüneberg, E., Ziche, D., and Wellbrock, N.: Organic carbon stocks and sequestration rates of forest soils in
212 Germany, *Glob. Change Biol.*, 20, 2644-2662, 2014.
- 213 Henkner, J., Scholten, T., and Kühn, P.: Soil organic carbon stocks in permafrost-affected soils in West
214 Greenland, *Geoderma*, 282, 147-159, 2016.
- 215 IPCC: Good practice guidance for land use, land-use change and forestry, *Good practice guidance for land use,*
216 *land-use change and forestry*, 2003. 2003.
- 217 Lozano-García, B. and Parras-Alcántara, L.: Land use and management effects on carbon and nitrogen in
218 Mediterranean Cambisols, *Agriculture, ecosystems & environment*, 179, 208-214, 2013.
- 219 Parfitt, R. L., Ross, C., Schipper, L. A., Claydon, J. J., Baisden, W. T., and Arnold, G.: Correcting bulk density
220 measurements made with driving hammer equipment, *Geoderma*, 157, 46-50, 2010.
- 221 Perruchoud, D., Walthert, L., Zimmermann, S., and Lüscher, P.: Contemporary carbon stocks of mineral forest
222 soils in the Swiss Alps, *Biogeochemistry*, 50, 111-136, 2000.
- 223 Poeplau, C. and Don, A.: Sensitivity of soil organic carbon stocks and fractions to different land-use changes
224 across Europe, *Geoderma*, 192, 189-201, 2013.
- 225 R Development Core Team: R: A language and environment for statistical computing. , R Foundation for
226 Statistical Computing, Vienna, Austria, 2010. 2010.
- 227 Rytter, R.-M.: Stone and gravel contents of arable soils influence estimates of C and N stocks, *CATENA*, 95,
228 153-159, 2012.
- 229 Throop, H. L., Archer, S. R., Monger, H. C., and Waltman, S.: When bulk density methods matter: Implications
230 for estimating soil organic carbon pools in rocky soils, *Journal of Arid Environments*, 77, 66-71, 2012.
- 231 Walter, K., Don, A., Tiemeyer, B., and Freibauer, A.: Determining Soil Bulk Density for Carbon Stock
232 Calculations: A Systematic Method Comparison, *Soil Sci. Soc. Am. J.*, 2016. 2016.
- 233 Wang, W. and Dalal, R.: Carbon inventory for a cereal cropping system under contrasting tillage, nitrogen
234 fertilisation and stubble management practices, *Soil and Tillage Research*, 91, 68-74, 2006.



235 Wiesmeier, M., Sporlein, P., Geuss, U., Hangen, E., Haug, S., Reischl, A., Schilling, B., von Lutzow, M., and
 236 Kogel-Knabner, I.: Soil organic carbon stocks in southeast Germany (Bavaria) as affected by land use, soil type
 237 and sampling depth, *Glob. Change Biol.*, 18, 2233-2245, 2012.

238

239 List of Tables

240 Table 1: Fraction of total observations for different volumetric stone content classes in the German Agricultural
 241 Soil Inventory and average soil organic carbon stock deviations [%] from M4 for the calculation methods M1 -
 242 M3 in different depth increments.

Depth	Fraction of total observations					Average relative deviation from M4		
	<5%	5-10%	10-20%	20-30%	>30%	M1	M2	M3
0-10	78.4	12.9	5.7	1.8	1.2	6.1	3.6	2.2
10-30	72.4	14.0	6.4	3.1	4.2	7.3	4.3	2.5
30-50	68.4	10.3	6.4	4.1	10.7	8.4	5.3	2.2
50-70	67.5	9.4	6.4	4.1	12.6	8.8	5.8	2.1
70-100	68.4	9.3	5.7	3.3	13.3	10.1	6.5	2.3

243

244

245

246

247

248

249

250

251

252

253

254


255

256

257




258 **Figures**



259

260 Figure 1: Soil organic carbon stocks of the German Agricultural Soil Inventory in different depth increments
 261 calculated by different calculation methods (M1-M4) for five volumetric stone content classes. Error bars
 262 indicate standard errors.




263

264 Figure 2: Systematic deviations in SOC stock from calculation method M4 for methods M1-M3 as a function of
 265 volumetric stone content. Bulk density of the fine soil was set to 1.2 [g cm⁻³] in this example.

266



267



268

269 Figure 3: Schematic overview on the four methods applied to estimate the mass of soil
270 organic carbon stocks. Different shades of brown are used to indicate different densities. Thereby the stone
271 fraction (ellipsoids) has the darkest brown and the fine soil fraction the lightest brown.