

1 **Response to Editor:**

2

3 Dear Editor,

4

5 First, thank you for your effort and detail in reviewing our manuscript. We have incorporated all of the  
6 line to line comments from the referees and have addressed all of the main concerns in this revised  
7 version of the manuscript. You will also find a response to your individual concerns below as well as in  
8 the submitted "mark-up" version of the manuscript found below.

9

10 1. Why the variation of soil properties in depth is consider in this research to be less than the expected  
11 variations due to anthropogenic impacts?

12 A: I must apologize that I do not fully understand the question. However, we have detailed the  
13 differences in the native soil properties compared to the ones that are impacted primarily by golf course  
14 management in the paragraph starting at line 164 in the manuscript. Our results show that human  
15 alteration is significantly greater than the natural variability.

16

17 2. Why variation within each location (golf course) is assumed to be less relevant than variations  
18 between locations? It is necessary to explain in the manuscript, particularly in the methods section all  
19 the evidences that the authors have in relation to this issue.

20 A: Thank you for bringing this important aspect forward. We have taken the opportunity to include an  
21 additional paragraph in the Methods section that details the decision to analyze between courses rather  
22 than within courses (lines 110-119 in the revised manuscript).

23

24 3. Please could you go deeper in the discussion on the possible ecological relevance of your results? As  
25 well as in the implications on the differences on sand content?

26 A: We have included information on ecological implications in each section of the discussion. These are  
27 mostly contained in lines: 195-201, 230-234, and 253-259.

28

29 4. Could you be a bit more conceptual and less descriptive in your conclusion? Which relevance for  
30 sustainability on medium and long term, for soil health has that the soils of golf courses are affected by  
31 management? You can introduce also this issue in the discussion.

32 A: We have added a conceptual portion to the conclusion that includes the relevance and implications of  
33 this preliminary research, supporting the need for a more detailed study. Similar changes have been  
34 included in the revised abstract.

35

36

37

38

39 **Effects of golf course management on subsurface soil properties in Iowa**

40 Matthew T. Streeter\* and Keith E. Schilling

41 Iowa Geological Survey, 300 Trowbridge Hall, University of Iowa, Iowa City, IA 52242

42 \*corresponding author: [matthew-streeter@uiowa.edu](mailto:matthew-streeter@uiowa.edu); (319) 335-1593

43 **Abstract**

44 Currently, in the USA and especially in the Midwest region, urban expansion is developing  
45 turfgrass landscapes surrounding commercial sites, homes, and recreational areas on soils that  
46 have been agriculturally managed for decades. Often, golf courses are at the forefront of  
47 conversations concerning anthropogenic environmental impacts since they account for some of  
48 the most intensively managed soils in the world. Iowa golf courses provide an ideal location to  
49 evaluate whether golf course management is affecting the quality of soils at depth. Our study  
50 evaluated how soil properties relating to soil health and resiliency varied with depth at golf  
51 courses across Iowa and interpreted relationships of these properties to current golf course  
52 management, previous landuse, and inherent soil properties. Systematic variation in soil  
53 properties including sand content, NO<sub>3</sub>, and SOM were observed with depth at six Iowa golf  
54 courses among three landform regions. Variability in sand content was identified between the  
55 20 and 50 cm depth classes at all courses, where sand content decreased by as much as 37%.  
56 Highest concentrations of SOM and NO<sub>3</sub> were found in the shallowest soils, whereas total C and  
57 P variability was not related to golf course management. Sand content and NO<sub>3</sub> were found to  
58 be directly related to golf course management, particularly at shallow depths. The effects of  
59 golf course management dissipated with depth and deeper soil variations were primarily due to  
60 natural geologic conditions. These two soil properties that were most noticeably altered by golf  
61 course management may directly impact crop productivity, soil health, and water quality and

62 while NO<sub>3</sub> may be altered relatively quickly in soil through natural processes, particle size of the  
63 soil may not be altered without extensive mitigation. Iowa golf courses continue to be  
64 developed in areas of landuse change from historically native prairies and more recently  
65 agriculture to urban landscapes. As soils are continually altered by human impacts, it is  
66 imperative that we monitor the changes, both physical and chemical, in order to establish  
67 management practices that maintain environmental sustainability and productivity.

68

## 69 **1. Introduction**

70 Critical evaluation of resilience and health of soil resources has been fueled by a recent  
71 urgency to understand anthropogenic impacts on environmental resources relating to local,  
72 regional, and international environmental sustainability and quality (Doran and Zeiss,  
73 2000;Glanz, 1995). Urban expansion is a primary cause for these recent concerns. Currently, in  
74 the USA and especially in the Midwest region, urban expansion is developing turfgrass  
75 landscapes surrounding commercial sites, homes, and recreational areas on soils that have  
76 been agriculturally managed for decades (Qian and Follett, 2002). Often, golf courses are at the  
77 forefront of conversations concerning anthropogenic environmental impacts since they account  
78 for some of the most intensively managed soils in the world (Balogh and Walker, 1992).

79 Limited information is available documenting changes in soils due to golf course  
80 management, especially at depth, since most studies focus on the rooting depth of turfgrass  
81 and near the surface where soils are engineered to ideal texture classes (Bauters et al., 2000).  
82 Qian and Follett (2002) conducted a study on the effects of landuse change to turfgrass on soil  
83 organic matter (SOM) and found that previous landuse imparted a strong baseline control.

84 However, while this study analyzed data from nearly 700 data sets, the results were limited to  
85 the top 15 cm of soil. In contrast, deep soil quality affects via landuse management has been  
86 observed in agricultural systems. Tomer and Burkart (2003) observed a significant increase in  
87 soil nutrient concentrations at a depth of 17 m below ground surface that was associated with  
88 historical fertilizer applications that occurred 20 years prior. Furthermore, they estimated  $\text{NO}_3^-$   
89 N percolation rates in loess soils to be approximately 0.67 m/yr. With the apparent decades-  
90 long translocation of soil nutrients after agricultural soil management, it is of interest to  
91 determine whether golf course management is similarly affecting the resiliency and long-term  
92 sustainability of soil resources.

93 There are over 400 golf courses in Iowa and many of these have been in operation for  
94 decades. With some of the most productive agricultural soils in the world, Iowa golf courses  
95 provide an ideal location to evaluate whether golf course management is affecting the quality  
96 of soils at depth. Specifically, our study objectives were to: 1) evaluate how soil physical and  
97 chemical properties relating to soil health and resiliency varied with depth at golf courses  
98 across Iowa; and 2) interpret relationships of these properties to current golf course  
99 management, previous landuse, and inherent soil properties.

## 100 **2. Materials and methods**

101 Our study sites consisted of 6 golf courses across Iowa, which were selected from  
102 approximately 421 golf courses in the state. A stratified random design was used to select  
103 courses by alphabetically grouping courses based on their location in eastern, central, and  
104 western Iowa and separating them into 18-hole and 9-hole classes. Courses were then selected  
105 using a random number generator. If permission for sampling was not obtained from the

106 course, the next randomly selected course was selected. Ultimately, three 18-hole and three 9-  
107 hole courses were selected (Figure 1). Iowa is comprised of several landform regions that are  
108 associated with different glacial advances and post-glacial erosion (Prior, 1991), and courses  
109 were selected in a variety of landform regions. One course was located on the lowan Surface  
110 landform region in Eastern Iowa which consists of a significantly dissected glacial landscape and  
111 loamy pedisediment. Two courses were located on the Southern Iowa Drift Plain, which is a  
112 highly weathered pre-Illinoian glacial landscape capped in loess. Finally, three courses were  
113 located on the Des Moines Lobe which consists of recent Wisconsinan age glacial deposits and  
114 hummocky topography (Table 1).

115 Three of the courses (Central 9, West 9, and West 18) were combined public and private  
116 use in towns of less than 5,000 people. The other three courses (East 9, East 18, and Central 18)  
117 were privately managed country clubs positioned in communities of 10-60,000 people. The  
118 smaller 9-hole courses ranged from 20 to 33 ha in size compared to the larger 18-hole courses  
119 which ranged from 37 to 68 ha. Tee boxes, greens, and fairways at the courses consisted of  
120 bentgrass (Argostis) except at the West 9 and Central 9 courses where fairways consisted of a  
121 variety of bentgrass, bluegrass (Poa Pratensis), and ryegrass (Lolium Perenne). Grass varieties in  
122 roughs varied widely but consisted mainly of bentgrass, bluegrass, fescue (Festuca), ryegrass,  
123 and annual bluegrass (Poa Annua) in varying proportions. For each course, superintendents  
124 recorded and made available annual fertilizer application rates (typically four applications per  
125 year) for tee boxes, fairways, and roughs (Table 2). Generally, the roughs received the least  
126 fertilizer while the tee boxes received the most. Four out of six roughs received no fertilizer.  
127 Only one course (West 9) applied P fertilizer, whereas all courses applied N and K fertilizer. Each

128 course was fertilized between 15 and 30 days prior to soil sampling. Prior land use at the  
129 courses was identified from historical air photography which was available starting in the 1930s  
130 (Iowa Department of Natural Resources, 2017). Golf course opening dates were verified with  
131 the superintendents (Table 1).

132 At each course, soil samples were collected in conjunction with shallow monitoring well  
133 installations being performed for a water quality study (Figure 2; Schilling and Streeter (2017)).  
134 Based on conversation with course superintendents, sampling locations were selected that  
135 could be easily accessed multiple times with minimal course disruption. Soil samples were  
136 collected according to the stratigraphy encountered at each site. Sampling depths were  
137 consistent within each course, but varied among courses based on the depth required to  
138 breach the water table for the water quality assessment (Table 1).

139 Continuous core was collected from each borehole by which the soil was described  
140 according to Schoeneberger (2012). Soil samples were collected, air dried and ground to pass a  
141 2 mm sieve and lab analyzed according to Brown (1998) for P by strong Bray extraction.  
142 Furthermore, total soil carbon (TC) and total soil nitrogen (TN) were determined by elemental  
143 analysis via dry combustion and chromatography (Costech Analytical Technologies, 2015). Soil  
144 organic Matter (SOM) was determined by weight loss on ignition (Walkley and Black, 1934) as  
145 described by Schulte (1995). Nitrate nitrogen (NO<sub>3</sub>) was measured by segmental flow analysis.  
146 Particle size distribution was determined by x-ray absorption (Micromeritics Instrument  
147 Corporation, 2015). A total of 127 soil samples were collected and analyzed (Table 1).

148 While specific management strategies at each golf course varied slightly between the  
149 tee, rough, and fairway areas, we found through interviews with golf course managers that

150 management strategy varied much more significantly between courses than between locations  
151 on the course. For example, the two central courses were located on the same landform region,  
152 yet course average N fertilizer was more than doubled at Central 18 compared to Central 9.  
153 When looking at these same two courses, N fertilization was the same for Tees and Fairways at  
154 the Central 9 course and was only 25% less on the fairway compared to the Tee on the Central  
155 18 course (Table 2). Based on these types of observations, the decision was made to combine  
156 samples within golf courses (3 sites per course) and compare between courses where  
157 differences in management were most pronounced.

### 158 **3. Results**

159 Site stratigraphy was expectedly variable across the range of courses. The courses were  
160 located on varying landscape positions and soils formed in a variety of different parent  
161 materials including alluvium, glacial till, and loess. Depth of soil development ranged from 26  
162 cm in glacial till at the central 18 course to 172 cm in loess at the west 18 course. Soils were less  
163 variable within courses, but variations were observed due to golf course management,  
164 primarily in terms of soil texture. In many locations, natural sand layers were identified at  
165 varying depths as well as evidence of human alteration to soil texture near the soil surface. On  
166 average, sand content was 25% higher in the upper 20 cm compared to the next lower depth  
167 class (50 cm depth) for all courses (Table 3 and Figure 3). Decreasing sand content with depth  
168 was most noticeable in courses on the Southern Iowa Drift Plain where average sand content  
169 decreased from 45 to 8%. However, since background sand content in soils from different  
170 geologic regions ranged widely among the courses (11-74%), we could not quantify the  
171 statistical significance of the sand content differences due to the limited sample size. In

172 essence, testing for changes in sand content in courses could be done within the same landform  
173 region because soils have similar background conditions, but in our study we have only 1-3  
174 courses per landform region.

175 In contrast to physical properties, varying concentrations of soil chemical properties  
176 were identified near the soil surface at multiple courses (Table 3). Course average TC and SOM  
177 ranged from 0.45-4.11% and 0.76-1.89%, respectively and ranged from 0.11-4.82% and 0.21-  
178 5.50% for individual samples. Likewise, SOM ranged from 2.9 to 5.5% in samples collected at 20  
179 cm, and was weakly correlated to TC ( $r=0.27$ ,  $p<0.05$ ). SOM decreased with depth at all courses  
180 ( $r=0.47$ ,  $p<0.05$ ) regardless of landform region, whereas TC was not correlated to depth.

181 Additional sampling is required to test the statistical significance of changes in soil chemical  
182 properties with depth, but these results do help to highlight the variability in the naturally  
183 calcareous parent material. NO<sub>3</sub> and TN concentrations also decreased with increasing depth  
184 ( $r=0.41$  and  $r=0.40$ , respectively,  $p<0.05$ ). NO<sub>3</sub> concentrations were observed to decrease at the  
185 greatest rates between the first two depth classes (20 to 50 cm), whereas TN decreased more  
186 gradually. NO<sub>3</sub> concentrations decreased substantially from 15 to 2 mg/kg between 20 and 50  
187 cm at East 18 located on the lowan surface while concentrations dropped by more than 50% at  
188 all other locations by 100 cm. C/N ratio ranged from 13 at the East 9 course (floodplain) to 57 at  
189 the Central 18 course (glacial upland). C/N ratios of more than 20 generally occurred at depths  
190 greater than 20 cm. However, C/N ratios were not significantly correlated with depth. Mean P  
191 concentrations ranged from 18-67 mg/kg and were not correlated with any other soil property  
192 measured.

#### 193 4. Discussion



#### 194 4.1. Soil Physical Properties

195 In this study, we evaluated soil conditions at six randomly selected Iowa golf courses  
196 located in three different landform regions of the state. Despite inherent variation associated  
197 with parent materials, study results indicated systematic variation in several soil properties with  
198 depth at the Iowa golf course sites that may be directly related to landuse. These systematic  
199 variations by depth are intriguing, but require further study to expand sample sizes in order to  
200 test statistical significance. Even so, preliminary conclusions may be drawn regarding the  
201 significance of landuse on soil properties.

202 At a gross scale, soil particle size distributions were mainly associated with the parent  
203 material of the region. Typical sand contents of loess in western Iowa consist of less than 10%  
204 sand, whereas soils formed in glacial till in central Iowa often exceed 50% sand and may be  
205 highly variable (Soil Survey Staff, 2016). Our study found less than 10% sand in the unaltered  
206 soil formed in loess from 20-100 cm at the West 18 course. However, the surface soil horizon  
207 was much higher (45%) at this course. Unlike the deeper depth classes, the elevated sand  
208 content near the soil surface was not likely an inherent soil property due to the nature of the  
209 loess parent material, but rather caused by golf course landuse. Our study found sand content  
210 was much higher at the four courses located on glacial till derived soils (West 9, Central 9 and  
211 18, and East 18) compared to the courses located on loess (West 18) or alluvium (East 9) (Table  
212 4). With the exception of the Central 9 course, the glacial derived soils exhibited a sharp  
213 decrease in sand content between the 20 and 50 cm depths (as much as 37%) that may not be  
214 explained primarily by inherent soil properties. Although natural variability in sand content

215 varied widely between landform regions, average sand content for all courses was 25% higher  
216 in the 0-20 cm interval compared to the 20 to 50 cm depth.

217 Golf course soils are regularly aerated and top-dressed by adding new sand to the  
218 surface of the established soil profile (Bandaranayake et al., 2003;Bauters et al., 2000). Golf  
219 course superintendents at our study sites confirmed that topdressing was performed on an  
220 annual basis at all but the Central 9 course. This topdressing management approach explains  
221 the elevated sand content near the surface at our study sites. Increased sand content of the  
222 surface soil horizons due to topdressing may have several implications for soil health and  
223 development. Altering the particle size distribution of the soil may in-turn alter several soil  
224 properties including bulk density and porosity (Arya and Paris, 1981;Gupta and Larson, 1979),  
225 soil compaction (Bodman and Constantin, 1965), hydraulic conductivity and water holding  
226 capacity (Campbell and Shiozawa, 1992;Jabro, 1992), as well as SOM content and distribution  
227 (Anderson et al., 1981;Puget et al., 2000;Tiessen and Stewart, 1983), and NO<sub>3</sub> concentrations  
228 via CEC (Anderson et al., 1981;Arya and Paris, 1981;Ersahin et al., 2006). Bandaranayake et al.  
229 (2003) explained that topdressing soil with sand prolongs the time required for SOM content to  
230 reach equilibrium in the soil profile. Since SOM affects several other soil properties, it is likely  
231 that the soil conditions at our study sites are much less stable than those at surrounding non-  
232 golf course sites.

233 Implications of these results may be considered long-term. Soil physical properties,  
234 especially soil particle size, develop naturally over geologic time-scales and even then, are  
235 highly dependent upon the physical properties of the parent material. In the case of our study,  
236 we have shown that surficial particle size content has been significantly altered by golf course

237 management over a relatively short time period. While the effects of topdressing may benefit  
238 golf course management in the short-term, these soils have become less stable and their ability  
239 to process soil nutrients and water have in effect been permanently altered.

#### 240 **4.2. Soil chemical properties**

241 Similar to the inherent nature of soil texture, TC content may also be primarily  
242 dependent on the soil parent material and may be highly spatially variable (Huang et al., 2007).  
243 Courses on unaltered glacial deposits (West 9, Central 9 and 18) exceeded 4% TC due to high  
244 inorganic carbon concentrations, whereas courses on reworked Iowan Surface glacial sediments  
245 (East 18) as well as weathered Southern Iowa Drift Plain deposits had TC concentrations less  
246 than 2%. These reworked and weathered soils have likely experienced inorganic carbon  
247 leaching which has left the soils void of almost all inorganic carbon. This was quite noticeable  
248 when comparing TC between landform regions at all depths and it also explains the variability  
249 in C/N ratio at depths greater than 100 cm in our study where C/N ratio is 4 to 10 times greater  
250 in Des Moines Lobe soils. The variability in TC (and C/N ratio below 100 cm) that we have  
251 identified through this study may be entirely affected by parent material and natural  
252 weathering patterns and serve to highlight geologic variation across landform regions.-

253 Soil chemical properties including SOM at the Iowa golf course were altered from their  
254 long-term equilibrium conditions established under native perennial vegetation in response to  
255 modern agricultural management and urban landuse. For example, agricultural drainage in  
256 conjunction with row crop agriculture can decrease SOM between 24 and 89% compared to  
257 that of perennially managed soils (Knops and Tilman, 2000;Kucharik et al., 2001;VandenBygaart  
258 et al., 2003). In urban areas, Qian and Follett (2002) conducted a study on the effects of landuse

259 change to turfgrass and found that previous landuse imparted a strong baseline control on SOM  
260 concentrations. Bandaranayake et al. (2003) estimated that turfgrass systems (e.g. golf courses)  
261 could sequester up to 32 Mg/ha of soil organic carbon (approximately 64Mg/ha of SOM) in the  
262 top 20 cm of soil within 30 years of establishment. At the Iowa golf courses, we observed  
263 substantially lower SOM contents ~~in the upper 20 cm (2.9 to 5.5%)~~ in samples collected ~~at~~  
264 below 20 cm. ~~but~~ C concentrations were highest in the uppermost layer and decreased with  
265 depth at all courses. Although accumulation of SOM in the upper 20 cm is consistent with  
266 Bandaranayake et al. (2003), attributing specific SOM changes to golf course management is  
267 not possible with our study. The SOM profiles could define native perennial, agricultural, or  
268 urban turfgrass landuse. Still, the implications of SOM alteration must be considered. SOM  
269 content may directly impact the soils ability to retain water and improve its quality via filtration  
270 (Rawls et al., 2003) and buffer the effects of climate change (Lal, 2004). Further sampling and  
271 critical analysis is necessary to draw definitive conclusions about the effects of golf course  
272 management on SOM.

273 In agricultural systems, application of N fertilizer has been found to migrate through the  
274 soil profile (De Haan et al., 2017; Tomer and Burkart, 2003). In deep loess soils of western Iowa,  
275 Tomer and Burkart (2003) documented a zone of high soil NO<sub>3</sub> located 17 m below land surface  
276 due to over-application of commercial fertilizer 20 years earlier. De Haan et al. (2017) observed  
277 that cropping systems has a large impact on residual soil NO<sub>3</sub> in the upper 1.8 m of the soil  
278 profile. Residual soil NO<sub>3</sub> in the soil profile under continuous corn with a ryegrass cover crop  
279 was more than seven times higher than a perennial grass system. Others have similarly  
280 observed residual NO<sub>3</sub> concentrations in soil under agricultural land use in Iowa as high as 60

281 mg/kg (Blackmer et al., 1989). However, the golf courses in our study applied varying rates of  
282 slow release N fertilizer to Tee boxes and fairways, and only two courses (East 9 and 18) applied  
283 N fertilizer to the roughs (Table 2). We estimated (based on typical bulk density values for  
284 surface soils in Iowa) that soil NO<sub>3</sub> concentrations ranged from 15-35 kg/ha for our study sites.  
285 Our study shows evidence of higher NO<sub>3</sub> levels in golf course soils than that of typical  
286 perennially managed soils (less than 10 kg/ha), but not necessarily as high as typical agricultural  
287 management (40-70 kg/ha) (De Haan et al., 2017). Recent studies provide evidence that once  
288 application of N fertilizer ceases, the top 1 m of the soil may return to native concentrations of  
289 NO<sub>3</sub> within 10 years (Streeter and Schilling, 2017). Since the courses for our study have been  
290 established for over 50 years, it is likely that the current NO<sub>3</sub> concentrations near the soil  
291 surface reflect golf course management rather than historical agricultural practices. Overall,  
292 limited field-scale research suggests that nutrients like NO<sub>3</sub> that are discharged from turf  
293 managed soils at golf course facilities are typically less than levels of major concern (King et al.,  
294 2007). However, a recent study in Iowa has suggested that a decade-long delay may be expected  
295 between changes in golf course management and groundwater quality (Schilling and Streeter,  
296 2018, 2017). Therefore, continued monitoring is necessary to assess changes in soil nutrients via  
297 golf course management and their impacts on environmental quality over time.

298 P concentrations at the courses were not correlated with any other soil property  
299 measured during our study. Furthermore, P fertilizer was only applied at 1 course (West 9) on  
300 the rough. Typically, background P concentrations in Iowa soils vary from 0-100 mg/kg  
301 depending on the mineralogy of the soil parent material (Fenton, 1999). Loess derived soils  
302 generally have the highest P, while glacial till derived soils will have some of the lowest P

303 concentrations (Fenton, 1999). Similarly, our study identified the highest P concentrations at  
304 the West 18 course (loess derived soil) and the lowest P concentrations at the Central 18 course  
305 (glacial till derived soil). Our study results suggest no significant alterations to background soil P  
306 concentrations by golf course management.

## 307 5. Conclusions

308 In this study, systematic variation in soil properties including sand content, NO<sub>3</sub>, and  
309 SOM were observed with depth at six Iowa golf courses among three landform regions.  
310 Variability in sand content was identified between the 20 and 50 cm depth classes at all  
311 courses, where sand content decreased by as much as 37%. Highest concentrations of SOM and  
312 NO<sub>3</sub> were found in the shallowest soils, whereas TC and P variability was not related to golf  
313 course management and did not correlate with depth. Although many of the soil properties  
314 measured for this study may be influenced by parent material and native vegetation, sand  
315 content and NO<sub>3</sub> were found to be directly related to golf course management, particularly at  
316 shallow depths. The effects of golf course management dissipated with depth and deeper soil  
317 variations were primarily due to natural geologic conditions. These two soil properties that  
318 were most noticeably altered by golf course management may directly impact crop  
319 productivity, soil health, and water quality and while NO<sub>3</sub> may be altered relatively quickly in  
320 soil through natural processes, particle size of the soil may not be altered without extensive  
321 mitigation. ~~The effects of golf course management dissipated with depth and deeper soil~~  
322 ~~variations were primarily due to natural geologic conditions.~~ Iowa golf courses continue to be  
323 developed in areas of landuse change from historically native prairies and more recently  
324 agriculture to urban landscapes. As soils are continually altered by human impacts, it is

325 imperative that we monitor the changes, both physical and chemical, in order to establish  
326 management practices that maintain environmental sustainability and productivity. Further  
327 work is recommended to increase sample size of course sites within different landform regions  
328 to better quantify the variability of soil properties with depth. Likewise, additional investigation  
329 of spatial patterns within individual courses would improve characterization of soil quality  
330 patterns among managed and unmanaged areas. Despite limitations, our study indicates that  
331 golf course management is affecting-impacting both short-term (soil N) and long-term (physical)  
332 quality ofthe surface and subsurface quality of soils.

### 333 **References**

- 334 Anderson, D., Saggar, S., Bettany, J., and Stewart, J.: Particle size fractions and their use in studies of soil  
335 organic matter: I. The nature and distribution of forms of carbon, nitrogen, and sulfur, Soil Science  
336 Society of America Journal, 45, 767-772, 1981.
- 337 Arya, L. M., and Paris, J. F.: A physicoempirical model to predict the soil moisture characteristic from  
338 particle-size distribution and bulk density data, Soil Science Society of America Journal, 45, 1023-1030,  
339 1981.
- 340 Balogh, J. C., and Walker, W. J.: Golf course management & construction: Environmental issues, CRC  
341 Press, 1992.
- 342 Bandaranayake, W., Qian, Y., Parton, W., Ojima, D., and Follett, R.: Estimation of soil organic carbon  
343 changes in turfgrass systems using the CENTURY model, Agronomy Journal, 95, 558-563, 2003.
- 344 Bauters, T., Steenhuis, T., DiCarlo, D., Nieber, J., Dekker, L., Ritsema, C., Parlange, J.-Y., and Haverkamp,  
345 R.: Physics of water repellent soils, Journal of Hydrology, 231, 233-243, 2000.
- 346 Blackmer, A., Pottker, D., Cerrato, M., and Webb, J.: Correlations between soil nitrate concentrations in  
347 late spring and corn yields in Iowa, Journal of Production Agriculture, 2, 103-109, 1989.
- 348 Bodman, G., and Constantin, G.: Influence of particle size distribution in soil compaction, California  
349 Agriculture, 36, 567-591, 1965.
- 350 Brown, J. R.: Recommended chemical soil test procedures for the North Central Region, Missouri  
351 Agricultural Experiment Station, University of Missouri--Columbia, 1998.
- 352 Campbell, G., and Shiozawa, S.: Prediction of hydraulic properties of soils using particle-size distribution  
353 and bulk density data, Indirect methods for estimating the hydraulic properties of unsaturated soils.  
354 University of California, Riverside, 317-328, 1992.
- 355 De Haan, R. L., Schuiteman, M. A., and Vos, R. J.: Residual soil nitrate content and profitability of five  
356 cropping systems in northwest Iowa, PloS one, 12, e0171994, 2017.
- 357 Doran, J. W., and Zeiss, M. R.: Soil health and sustainability: managing the biotic component of soil  
358 quality, Applied soil ecology, 15, 3-11, 2000.
- 359 Ersahin, S., Gunal, H., Kutlu, T., Yetgin, B., and Coban, S.: Estimating specific surface area and cation  
360 exchange capacity in soils using fractal dimension of particle-size distribution, Geoderma, 136, 588-597,  
361 2006.

362 Fenton, T.: Phosphorus in Iowa soils, Agronomy Department, Iowa State University, 1999.

363 Glanz, J.: Saving our soil: solutions for sustaining earth's vital resource, Johnson Books, 1995.

364 Gupta, S., and Larson, W.: Estimating soil water retention characteristics from particle size distribution,  
365 organic matter percent, and bulk density, Water resources research, 15, 1633-1635, 1979.

366 Huang, X., Senthilkumar, S., Kravchenko, A., Thelen, K., and Qi, J.: Total carbon mapping in glacial till soils  
367 using near-infrared spectroscopy, Landsat imagery and topographical information, Geoderma, 141, 34-  
368 42, 2007.

369 Natural Resources Geographic Information Systems Library: <https://programs.iowadnr.gov/nrgislib/>,  
370 access: April, 2017.

371 Jabro, J.: Estimation of saturated hydraulic conductivity of soils from particle size distribution and bulk  
372 density data, Transactions of the ASAE, 35, 557-560, 1992.

373 King, K., Balogh, J., Hughes, K., and Harmel, R.: Nutrient load generated by storm event runoff from a  
374 golf course watershed, Journal of environmental quality, 36, 1021-1030, 2007.

375 Knops, J. M., and Tilman, D.: Dynamics of soil nitrogen and carbon accumulation for 61 years after  
376 agricultural abandonment, Ecology, 81, 88-98, 2000.

377 Kucharik, C. J., Brye, K. R., Norman, J. M., Foley, J. A., Gower, S. T., and Bundy, L. G.: Measurements and  
378 modeling of carbon and nitrogen cycling in agroecosystems of southern Wisconsin: potential for SOC  
379 sequestration during the next 50 years, Ecosystems, 4, 237-258, 2001.

380 Lal, R.: Soil carbon sequestration impacts on global climate change and food security, science, 304, 1623-  
381 1627, 2004.

382 Prior, J. C.: Landforms of Iowa, University of Iowa Press, 1991.

383 Puget, P., Chenu, C., and Balesdent, J.: Dynamics of soil organic matter associated with particle-size  
384 fractions of water-stable aggregates, European Journal of Soil Science, 51, 595-605, 2000.

385 Qian, Y., and Follett, R. F.: Assessing soil carbon sequestration in turfgrass systems using long-term soil  
386 testing data, Agronomy Journal, 94, 930-935, 2002.

387 Rawls, W., Pachepsky, Y. A., Ritchie, J., Sobecki, T., and Bloodworth, H.: Effect of soil organic carbon on  
388 soil water retention, Geoderma, 116, 61-76, 2003.

389 Schilling, K. E., and Streeter, M. T.: Groundwater Nutrient Concentrations and Mass Loading Rates at  
390 Iowa Golf Courses, JAWRA Journal of the American Water Resources Association, 2017.

391 Schilling, K. E., and Streeter, M. T.: Groundwater Nutrient Concentrations and Mass Loading Rates at  
392 Iowa Golf Courses, JAWRA Journal of the American Water Resources Association, 54, 211-224, 2018.

393 Schoeneberger, P. J., Wysocki, D.A., Benham, E.C., Soil Survey Staff.: Field Book for Describing and  
394 Sampling Soils, Version 3. 0, Government Printing Office, 2012.

395 Schulte, E.: Recommended soil organic matter tests, Recommended Soil Testing Procedures for the  
396 North Eastern USA. Northeastern Regional Publication, 52-60, 1995.

397 Web Soil Survey: <http://websoilsurvey.nrcs.usda.gov>, access: December, 2016.

398 Streeter, M. T., and Schilling, K. E.: Soil Properties in Native, Reconstructed, and Farmed Prairie Potholes:  
399 A Chronosequence Study of Restoration Timeframes, Ecological Restoration, 35, 6-12, 2017.

400 Tiessen, H., and Stewart, J.: Particle-size fractions and their use in studies of soil organic matter: II.  
401 Cultivation effects on organic matter composition in size fractions, Soil Science Society of America  
402 Journal, 47, 509-514, 1983.

403 Tomer, M., and Burkart, M.: Long-term effects of nitrogen fertilizer use on ground water nitrate in two  
404 small watersheds, Journal of Environmental Quality, 32, 2158-2171, 2003.

405 VandenBygaart, A., Gregorich, E., and Angers, D.: Influence of agricultural management on soil organic  
406 carbon: A compendium and assessment of Canadian studies, Canadian Journal of Soil Science, 83, 363-  
407 380, 2003.

408 Walkley, A., and Black, I. A.: An examination of the Degtjareff method for determining soil organic  
409 matter, and a proposed modification of the chromic acid titration method, Soil science, 37, 29-38, 1934.



411 **Figure Captions:**

412 Figure 1. Location map showing the six golf courses chosen for this study.

413 Figure 2. Soil sampling and monitoring well installation at East 18 course.

414 Figure 3. Profile of selected soil constituents with depth at the six Iowa golf courses.

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430 Table 1. Golf course site information.

<b>Course</b>	<b>Year Opened</b>	<b>Previous Landuse</b>	<b>Landform</b>	<b>Parent Material</b>	<b>Max. Depth (cm)</b>	<b>Number of Samples</b>
West 18	1963	row crop	Southern Iowa Drift Plain	loess	767	19
West 9	1938	row crop	Des Moines Lobe	glacial till	380	20
Central 9	1965	row crop	Des Moines Lobe	glacial till	457	18
Central 18	1915	no prior history	Des Moines Lobe	glacial till	506	26
East 18	1965	row crop	Iowan Surface	glacial sediments	546	22
East 9	1920	no prior history	Southern Iowa Drift Plain	alluvium	405	22

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445 Table 2. Summary of annual fertilizer rates at selected Iowa golf courses.

Course	Position	N kg/ha	Weighted Course Average N		
			kg/ha	P kg/ha	K kg/ha
West 18	Tee	90		0	44
West 18	Rough	0		0	0
West 18	Fairway	77	28.5	0	11
West 9	Tee	73		5.4	22
West 9	Rough	0		0	0
West 9	Fairway	37	15.4	0	6
Central 18	Tee	146		0	26
Central 18	Rough	0		0	0
Central 18	Fairway	110	41.7	0	0
Central 9	Tee	51		0	17
Central 9	Rough	0		0	0
Central 9	Fairway	51	18.5	0	17
East 18	Tee	87		0	132
East 18	Rough	112		0	17
East 18	Fairway	63	95.8	0	107
East 9	Tee	169		0	20
East 9	Rough	22		0	7
East 9	Fairway	115	59.2	0	19

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461 Table 3. Soil conditions (mean and standard deviation) by depth at Iowa golf courses.

Course	n	Depth Class(cm)	Sand %	P mg/kg	SOM %	NO3 mg/kg	TC %	TN %	C/N
Central 18	2	20	53 +/-42	18 +/-2	5.50 +/-4.24	5.00 +/-2.82	4.63 +/-0.17	0.33 +/-0.28	22 +/-19
Central 18	5	50	40 +/-23	18 +/-9	4.46 +/-3.69	8.20 +/-6.22	4.82 +/-2.35	0.29 +/-0.21	47 +/-75
Central 18	6	100	40 +/-21	03 +/-3	1.65 +/-1.83	3.83 +/-2.31	3.08 +/-2.18	0.10 +/-0.08	30 +/-33
Central 18	12	500	26 +/-25	02 +/-2	0.48 +/-0.27	1.25 +/-0.45	4.21 +/-2.14	0.05 +/-0.05	84 +/-37
Central 18	1	1000	05 +/-	02 +/-	0.30 +/-	1.00 +/-	4.69 +/-	0.07 +/-	67 +/-
Central 9	2	20	35 +/-2	14 +/-1	5.50 +/-0.14	15.50 +/-	2.58 +/-0.48	0.18 +/-0.07	14 +/-4
Central 9	2	50	50 +/-1	10 +/-0	2.35 +/-1.76	4.00 +/-4.24	1.81 +/-0.43	0.08 +/-0.01	22 +/-12
Central 9	5	100	46 +/-10	04 +/-9	1.52 +/-1.13	1.60 +/-0.89	1.15 +/-0.84	0.06 +/-0.05	19 +/-1
Central 9	9	500	61 +/-10	04 +/-3	0.48 +/-0.30	1.11 +/-0.33	1.51 +/-1.25	0.01 +/-0.02	15 +/-5
East 18	1	20	81 +/-	31 +/-	2.60 +/-	15.00 +/-	1.54 +/-	0.11 +/-	14 +/-
East 18	6	50	74 +/-9	27 +/-1	1.53 +/-0.95	2.83 +/-2.31	0.77 +/-0.54	0.04 +/-0.03	19 +/-6
East 18	4	100	78 +/-8	17 +/-1	0.40 +/-0.21	1.25 +/-0.50	0.18 +/-0.12	0.01 +/-0.00	18 +/-9
East 18	9	500	79 +/-25	15 +/-3	0.21 +/-0.16	1.11 +/-0.33	0.11 +/-0.08		
East 18	2	1000	42 +/-0	02 +/-2	0.80 +/-0	1.00 +/-0	1.04 +/-0.10		
East 9	4	20	11 +/-5	16 +/-1	3.15 +/-1.96	4.50 +/-3.31	1.74 +/-1.77	0.10 +/-1.12	17 +/-22
East 9	6	50	07 +/-2	23 +/-5	1.48 +/-0.42	1.00 +/-0	0.94 +/-0.36	0.09 +/-0.02	10 +/-2
East 9	12	500	28 +/-33	27 +/-2	0.57 +/-0.26	1.00 +/-0	0.22 +/-0.08	0.02 +/-0.02	11 +/-1
West 18	1	20	45 +/-	97 +/-	2.90 +/-	9.00 +/-	1.72 +/-	0.14 +/-	12 +/-
West 18	6	50	06 +/-11	28 +/-4	2.92 +/-0.95	5.83 +/-3.43	0.90 +/-0.75	0.04 +/-0.05	22 +/-22
West 18	3	100	01 +/-0	46 +/-2	2.20 +/-0.85	3.67 +/-2.08	1.07 +/-0.58	0.08 +/-0.01	13 +/-5
West 18	6	500	16 +/-36	98 +/-3	0.63 +/-0.35	1.17 +/-0.40	0.24 +/-0.12	0.01 +/-0.01	31 +/-36
West 18	3	1000	10 +/-6	48 +/-3	0.67 +/-0.11	1.00 +/-0	0.20 +/-0.05	0.01 +/-0.01	20 +/-
West 9	1	20	26 +/-	19 +/-	2.90 +/-	3.00 +/-	3.12 +/-	0.12 +/-	26 +/-
West 9	3	50	17 +/-4	28 +/-9	4.97 +/-1.54	5.33 +/-2.08	3.04 +/-0.84	0.2 +/-0.07	15 +/-4
West 9	5	100	33 +/-15	21 +/-2	1.56 +/-0.92	3.60 +/-2.61	2.27 +/-0.84	3.6 +/-0.05	23 +/-9
West 9	11	500	33 +/-24	38 +/-4	0.76 +/-0.56	1.27 +/-0.64	1.87 +/-0.86	0.02 +/-0.03	19 +/-16

462  
463  
464  
465  
466  
467  
468  
469  
470  
471

472

473 Table 4. ~~Mean s~~Soil conditions (mean and standard deviation) by course at Iowa golf courses.

Course	Sand %	P mg/kg	SOM %	NO3 mg/kg	TC %	TN %	C/N
Central 18	33 +/-25	18 +/-25	1.89 +/-2.65	3.46 +/-3.87	4.11 +/-2.06	0.13 +/-0.15	57 +/-48
Central 9	52 +/-12	50 +/-26	1.53 +/-1.74	3.16 +/-5.54	1.56 +/-1.05	0.04 +/-0.06	18 +/-6
East 18	74 +/-19	37 +/-24	0.76 +/-0.84	2.22 +/-3.17	0.45 +/-0.50	0.01 +/-0.03	22 +/-7
East 9	19 +/-26	57 +/-22	1.29 +/-1.26	1.63 +/-1.86	0.69 +/-0.91	0.05 +/-0.06	13 +/-13
West 18	11 +/-22	67 +/-44	1.72 +/-1.24	3.42 +/-3.18	0.64 +/-0.63	0.03 +/-0.04	23 +/-22
West 9	30 +/-20	31 +/-35	1.7 +/-1.69	2.55 +/-2.13	2.20 +/-0.91	0.06 +/-0.07	20 +/-10

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492


493

494

495

496

497 Figure 1.



498

499


500 Figure 2.



501

502

503 Figure 3.



504

505

506