

Interactive comment on “Spatial assessments of soil organic carbon for stakeholder decision-making. A case study from Kenya” by Tor-Gunnar Vågen et al.

Tor-Gunnar Vågen et al.

t.vagen@cgiar.org

Received and published: 2 July 2018

Abstract: We have edited the abstract as suggested, with objectives now earlier in the abstract. Also, some edits were made to the final part of the abstract to improve the flow of the language after moving these sections from the beginning of the abstract.

References in introduction: Thank you for this suggestion. We have added references to the SDGs as well as additional references related to our section on indicators in the introduction.

Units: We use metric units throughout the manuscript. The reason for using m^2 , and

km² is simply that expressing for example 100 km² in m² (100,000,000 m²) is not very intuitive for readers. The same also applies to smaller areas, where expressing these in km² would lead to numbers with a lot of decimals. Hence, we have not changed these units in the manuscript.

Description of study site: We have added a brief description of Turkana County.

Interactive comment on SOIL Discuss., <https://doi.org/10.5194/soil-2017-37>, 2018.

Interactive comment

[Printer-friendly version](#)

[Discussion paper](#)



Interactive comment on “Spatial assessments of soil organic carbon for stakeholder decision-making. A case study from Kenya” by Tor-Gunnar Vågen et al.

Tor-Gunnar Vågen et al.

t.vagen@cgiar.org

Received and published: 2 July 2018

Thank you for your various suggestions and corrections.

We expanded the methods section around the mapping of soil and land health, while attempting to not make this too technical as the other papers that are cited provide a more technical description and discussion.

The SHARED approach has been elaborated on somewhat, including the co-design elements of it, tying these to indicator frameworks and use of evidence in decision making. Some of these edits were done in response to reviewer 1.

Printer-friendly version

Discussion paper



Figure 1: Thanks for noticing the typo - we have fixed this (including in the text of the manuscript)

Figure 5 missing: This has been rectified. We also noticed a couple of cross-references that had not rendered correctly and have fixed these.

Interactive comment on SOIL Discuss., <https://doi.org/10.5194/soil-2017-37>, 2018.

SOILD

Interactive
comment

Printer-friendly version

Discussion paper



Spatial assessments of soil organic carbon for stakeholder decision-making. A case study from Kenya

Tor-Gunnar Vågen^{*1}, Leigh Ann Winowiecki¹, Constance Neely¹, Sabrina Chesterman¹ and Mieke Bourne¹

¹World Agroforestry Centre (ICRAF), P.O. Box 30677 - 00100, Nairobi, Kenya

5 *Corresponding author: t.vagen@cgiar.org

Abstract

Land degradation impacts the health and livelihoods of about 1.5 billion people worldwide. Given that the state of the environment and food security are strongly interlinked in tropical landscapes, the increasing need for land for food

10 production, urbanization and other uses pose several threats to sustainability in the long term. This paper demonstrates the integration of land and soil health maps with socioeconomic datasets into an online, open-access platform called the Resilience Diagnostic and Decision Support Tool for Turkana County in Kenya, using the Stakeholder Approach to Risk Informed and Evidence Based Decision Making (SHARED) methodology. The paper highlights the utility of spatial assessments of soil organic carbon (SOC) for monitoring of land degradation neutrality (LDN) compliance, understanding
15 the drivers of SOC dynamics, and inclusion in stakeholder decision-making. The main objectives of this paper were to: 1) demonstrate the application of a systematic approach for land health assessments, including spatial mapping of soil organic carbon; 2) show an operational interdisciplinary framework for assessing ecosystem health; and 3) showcase the application
20 of evidence-based tools for stakeholder engagement using the SHARED approach. Through the approaches and tools presented, the paper addresses the increasing need for more integrated approaches when assessing and managing ecosystem health to meet the targets of the 2030 Agenda, including SDG 15.3. In addition to systematic and reliable biophysical and socio-economic assessments, stakeholder engagement with evidence is crucial to support such integrated approaches.

Keywords: soil organic carbon, stakeholder engagement, remote sensing, evidence-based decision making, resilience

Formatted: Font: Italic

1. Introduction

The 2030 Agenda, including multi-lateral environmental agreements and the Sustainable Development Goals (SDGs), has set the stage for greater appreciation and understanding of the complex nature and interactions among environmental challenges facing society ([United Nations General Assembly, 2015](#)).

SDG 15 calls for the protection, restoration and promotion of sustainable use of terrestrial ecosystems, sustainably managed forests, combating desertification, halting and reversing land degradation and halting biodiversity loss. At their core, initiatives such as Land Degradation Neutrality (LDN) and SDG 15 emphasize healthy ecosystem function and resilient landscapes, which underpin healthy economies and societal well-being. Because these issues are intrinsically inter-related, decisions around soil and land cannot be taken in isolation. Achieving the associated targets, requires a robust evidence base for measuring and monitoring land health indicators and associated land management practices, coupled with local and policy level awareness of the importance of land health in supporting multiple sectors. Also, local capacity to implement, monitor and assess these indicators is needed, including mechanisms for integrated, cross-sectoral coordination and inclusive, multi-stakeholder collaboration to achieve impact.

Many assessments of land health suffer from (i) disagreements about the definition of land degradation/land health; (ii) an abundance of indicators ([Heink and Kowarik, 2010](#)) that are often not feasible to measure and hence operationalize, and (iii) a lack of rigorous science-based [indicators \(Niemeijer and Groot, 2008\)](#) and analytical frameworks. Comparable indicators are critical when assessing environmental conditions and progress made towards the mitigation or avoidance of land degradation, but they are also important to effectively communicate information both to stakeholders such as farmers or advisory services, and to policy makers. In addition to being comparable, indicators for assessment and monitoring of land health should be a) science based; b) readily measurable (quantifiable); c) rapid; d) based on field assessment across multiple scales (plot, field, landscape, region) and e) representative of the complex processes of land degradation in landscapes ([Dale and Beyeler, 2001](#)). The methods, tools and approaches presented in this paper offer realistic and cost-effective options both for baseline assessments of LDN indicators and for monitoring of progress towards LDN targets. Several assessments of methodologies for LDN compliance (Aynekulu et al., 2017; Caspari et al., 2015; Stavi and Lal, 2015) also highlight the need for consistent, spatially explicit data over time.

The Land Degradation Surveillance Framework (LDSF) is an example of a method that has been applied in a number of projects in the global tropics to provide more rigorous, science-based, assessments of land degradation risk and status, as well as soil health (Vågen et al., 2013a). An important indicator of soil health is soil organic carbon (SOC), and field datasets collected using the LDSF have been used to provide spatially explicit assessments of SOC and other indicators of land degradation processes such as soil erosion, in addition to other soil functional properties (Vågen et al., 2012, 2016; Winowiecki et al., 2016a, 2016b). These assessments can be used to inform spatially explicit land and soil health monitoring systems, which are critical in order for countries to avoid land degradation, or to restore already degraded ecosystems.

Combining systematic data collection efforts with rigorous analytical frameworks, evidence-based approaches can be used to engage stakeholders with interactive online user-friendly platforms to inform national and international policy makers.

Within this context, the engagement of stakeholders becomes an important element to prioritize investment strategies for accelerating the achievement of the SDGs. The Stakeholder Approach to Risk-informed and Evidence-based Decision-making (SHARED)¹ emerged in response to this need and was developed around a number of key factors, steps and principles, including: i) advancing a holistic or systems view to raise awareness on the integrated nature of environmental, social, cultural and economic dimensions and causal relationships; ii) establishing a clear understanding of the influencing factors of human and group decision making including stakeholder analysis; iii) facilitating different government sectors and multi-stakeholder platforms of diverse societal sectors; iv) collectively articulating mutually agreed, desired sustainable development outcomes and indicators building upon fundamental ecosystem services and nested within national and global goals, v) generating evidence and experience and tailoring tools in a readily consumable way for problem solving and options identification, vi) testing options based on collectively defined criteria, including risks and potential synergies and vii) designing option implementation with monitoring and evaluation and co-learning feedback into the process.

This paper demonstrates the integration of land and soil health maps, using spatial assessments of SOC as an example, with socio-economic datasets into an online tool, the Resilience Diagnostic and Decision Support Tool² (RDDST) in Turkana County, Kenya. The main objectives of the efforts discussed in this paper were to: 1) demonstrate the application of a systematic approach for land health assessments, including spatial mapping of soil organic carbon; 2) demonstrate the operationalization of interdisciplinary framework for assessing ecosystem health; and 3) showcase the application of evidence-based tools for stakeholder engagement using the SHARED approach. The approaches presented can also be applied as part of LDN efforts to assist countries in more effectively restoring degraded lands and increasing agricultural productivity, including opportunities for sustainable development and biodiversity conservation (IUCN, 2015). A recent assessment of land degradation in Kenya estimates that the annual cost of land degradation is at approximately 1.5 billion USD, which represents about 5% of the country's GDP (Munoz, 2016). Kenya has endorsed the SDG 15.3 targets and discussions on baseline assessments and monitoring are underway. Given the large extent of drylands in Kenya, their management will be critical to achieving these targets.

1 <http://www.worldagroforestry.org/shared>

2 <http://landscapeportal.org:3838/sharedApp>

2. Methodology

This paper utilizes spatial assessments of SOC that were conducted using data from a network of Land Degradation Surveillance Framework (LDSF) sites in the global tropics. The LDSF was designed for practical and cost-effective soil and ecosystem health surveillance, including for mapping of SOC (Vågen et al., 2013a, 2016; Winowiecki et al., 2016a, 2016b).

5 The framework is also designed to monitor changes over time, and provides opportunities for targeting improved soil management and land restoration activities (Lohbeck et al., 2018). Specifically, the LDSF systematically assesses several ecological metrics simultaneously at four different spatial scales (100 m^2 , 1000 m^2 , 1 km^2 and 100 km^2), using a spatially stratified, hierarchical sampling design (Vågen et al., 2013b). The LDSF also applies the latest soil infrared (IR) spectroscopy technologies in analysis of SOC and other soil properties, which are cost effective and hence allow for the 10 scaling of soil measurements. The spatial assessments of SOC included in this study were developed based on an archive of 10,000 georeferenced LDSF plots with soil samples that were analyzed for SOC and mid-infrared analysis at the ICRAF Soil and Plant Diagnostics Lab in Nairobi, Kenya and used to conduct spatial predictions of SOC concentrations and stocks using reflectance data from the Moderate Resolution Imaging Spectroradiometer (MODIS) satellite sensor (Vågen et al., 2016). In
15 brief, annual MODIS reflectance image composites were created based on the date of maximum fractional vegetation cover in each MODIS pixel and these reflectance data were used to train models using the Random Forest algorithm (Breiman, 2001) to predict soil properties such as SOC and pH, as well as land degradation processes such as soil erosion. Modeling of soil and land health variables was conducted in R Statistics (R Core Team, 2018) using the *randomForest* (Liaw and Wiener, 2002) R library. The spatial assessments of SOC published in Vågen et al. (2016) were clipped for Turkana County and applied as part of the interactive tools described below. In brief, the accuracy of the spatial estimates of SOC based on
20 MODIS as measured using the root-mean squared error of prediction (RMSEP) for three independent model test runs ranged from 9.6 to 11.5 g C kg⁻¹.

An interactive dashboard was developed for Turkana County, which is located in north-western Kenya, bordering Uganda, Sudan and Ethiopia. Turkana County is part of the Arid and Semi-Arid Lands (ASALs) of Kenya with an annual average rainfall of about 250 mm, a population of almost 1 million people, the majority of which as pastoralists. We used the Shiny 25 web framework for R statistics (Chang et al., 2017; R Core Team, 2018) to develop an interactive dashboard integrating existing and new data and providing robust data management and graphical tools to allow users to interact with these data in a meaningful way. The RDDST dashboard is hosted on the ICRAF Landscape Portal³, which provides a platform and various tools for spatial data storage and visualization.

3 <http://landscapeportal.org/sharedApp/>

5 The SHARED approach was applied to foster the innovation required to shape and embed land health assessments into inclusive negotiation and decision-making processes through a co-design process with stakeholders in Turkana County. By utilizing a comprehensive framework tailored to specific decision needs, coupled with science-based evidence, processes, evidence and tools were brought together to shift the decision paradigm towards more inclusive, inter-sectoral and inter-institutional integration to tackle complex decisions and achieve desired outcomes. The approach emphasises the importance of rigorous indicator frameworks that are simple and readily measurable, but that allow reflect the complexity of both ecological systems and decision making processes.

10 Through robust data management and advanced data visualization, the RDDST was co-developed with stakeholders in Turkana County to help facilitate communication of data and analysis between scientists and stakeholders. This allows for the interrogation of evidence, an increased rate of discovery and helps contextualize the data used. Third-party data sources for the RDDST included the Armed Conflict Location & Event Data Project (ACLED) for security data, Kenya government statistics for education and health data, and data from the Hunger Safety Net Programme (HSNP) for information on livestock populations and energy use within the county.

15 The targeted facilitation The targeted facilitation and co-design approach employed ensures local ownership and cohesive communication across multiple institutions, including different political levels and knowledge systems to develop capacity and the evidence-base as a continuously linked process. At the national level, the SHARED approach has been used in collaboration with Kenya's Ministry of Environment and Natural Resources and the Ministry of Agriculture, Livestock and Fisheries to synthesise the evidence of 44 integrated crop-livestock-tree system projects to develop recommendations for climate-resilient approaches (Chesterman and Neely, 2015) and to directly inform the drafting of Kenya's National Climate

20 Change Policy Framework (Neely, 2014).

3. Results

25 Spatial estimates of SOC stocks for Kenya, show that Kenyan soils store approximately 2.4 Gt carbon (C) in the upper 30 cm of the soil profile (Minasny et al., 2017). Average SOC stocks for Kenya were estimated to 42 Mg C ha^{-1} (Figure 1) for 0 to 30cm depth. The lowest estimated C stocks were found in arid and semi-arid regions of the country ($<20 \text{ Mg C ha}^{-1}$ on average), such as Turkana County. These estimates are consistent with those reported by Batjes and Batjes (2004) for arid- and semiarid parts of the country. Higher SOC stocks were found in sub-humid and humid areas, including the central highlands and western parts of the country (Figure 1). The highest SOC stocks are found in forest systems, such as around Mt Kenya ($>100 \text{ Mg C ha}^{-1}$), the Aberdares, the Mau Forest Complex and Kakamega forest. These forest systems, while making up a small proportion of the country's area, are critical carbon pools and also represent important water towers that 30 millions of Kenyans rely on for their water supply (Mogaka et al., 2005). Also, wetland ecosystems such as inland riverine and palustrine areas are critical SOC pools in Kenya and in much of East Africa. In the drylands, such wetland systems are

particularly critical for SOC storage and land health, frequently reaching between 80 and 100 $Mg\ C\ ha^{-1}$ at 0 to 30cm depth. Other wetland systems, many of them under threat, such as around the Rift Valley lakes and lacustrine wetlands along Kenya's coast are also examples of ecosystems that are critical for C storage (Minasny et al., 2017; Saunders et al., 2007) and other ecosystem services (Zedler and Kercher, 2005) in the country.

5 [\[Figure 1 here\]](#)

of Kenyans rely on for their water supply (Mogaka et al., 2005). Also, wetland ecosystems such as inland riverine and palustrine areas are critical SOC pools in Kenya and in much of East Africa. In the drylands, such wetland systems are particularly critical for SOC storage and land health, frequently reaching between 80 and 100 $Mg\ C\ ha^{-1}$ at 0 to 30cm depth. Other wetland systems, many of them under threat, such as around the Rift Valley lakes and lacustrine wetlands along Kenya's coast are also examples of ecosystems that are critical for C storage (Minasny et al., 2017; Saunders et al., 2007) and other ecosystem services (Zedler and Kercher, 2005) in the country.

10 [\[Figure 1 here\]](#)

15 Soil organic carbon (SOC) concentrations were mapped for Africa at 500m spatial resolution (Vågen et al., 2016) and clipped for Kenya (Figure 2) for use in stakeholder engagement processes within Turkana County (see county outlined in Figure 2). Large parts of the drylands have SOC concentrations that are lower than $15\ g\ C\ kg^{-1}$, which is generally considered very low, as shown in Figure 3. Given the emphasis on management of SOC as a critical resource for both agriculture and rangeland management in Turkana County, SOC concentrations were applied in the RDDST rather than estimates of SOC stocks.

20 [\[Figure 2 here\]](#)

25 Concentrations of SOC follow a similar pattern as the SOC stocks, as shown in Figure 2 and Figure 3, with very low SOC in parts of the drier northern and north-eastern parts of the country. However, it is worth noting that SOC is relatively heterogenous within these areas and that there are pockets of high SOC in parts of the drylands such as the Matthews Range and mountains such as Ndoto, Marsabit and Kulal, as well as Loima Hills in Turkana County. These areas represent critical SOC pools, are important resources for pastoralists in the drylands for dry-season grazing (Oba et al., 2000), and are hotspots for biodiversity in the region.

30 ~~of SOC shown in Figure 2. Figure 2 shows very low SOC in northern and north-eastern parts of the country. However, it is worth noting that SOC is relatively heterogenous within these areas and that there are pockets of high SOC in parts of the drylands such as the Matthews Range and mountains such as Ndoto, Marsabit and Kulal, as well as Loima Hills in Turkana County. These areas represent critical SOC pools, are important resources for pastoralists in the drylands for dry-season grazing (Oba et al., 2000), and are hotspots for biodiversity in the region.~~

[Figure 3 here]

In Turkana County, maps of SOC concentrations were included in the RDDST tool to assess resilience within the county and support decision making processes (Figure 4), integrating data from across numerous sectors, from security and education to human health and land health, support decisionmaking processes (Figure 4), integrating data from across numerous sectors, security education human health and land health. Through a series of workshops facilitated using the

5 SHARED process, the RDDST was co-developed with representatives from the Turkana County government and other stakeholders such as representatives from UN organizations and Non-Governmental Organizations (NGOs) in the county.

Furthermore, the outcomes of these workshops and the RDDST was applied in the development of the Turkana County Integrated Development Plan for the period 2018 to 2022. The land health part of the RDDST was used to interactively 10 visualize and explore SOC (Figure 5) along with other key land health indicators across the county, including soil erosion prevalence, root-depth restrictions and soil pH. In addition, a module for exploring vegetation cover status and trends was applied for assessing vegetation performance and trends. These tools are being used to demonstrate the critical role of the underpinning natural resource base and the importance of taking this into account in decision making across multiple sectors. Visualization of local land health parameters in conjunction with data from other sectors shifted the discourse in the County 15 Government and resulted in the identification of integrated flagships for multiple sectors working together on development outcomes with a base in land restoration.

In addition, a module for exploring vegetation cover status and trends was applied for assessing vegetation performance and trends. These tools are being used to demonstrate the critical role of the underpinning natural resource base and the importance of taking this into account in decision making across multiple sectors. Visualization of local land health parameters in conjunction with data from other sectors shifted the discourse in the County Government and resulted in the 20 identification of integrated flagships for multiple sectors working together on development outcomes with a base in land restoration.

[Figure 4 here]

[Figure 5 here]

25 4. Discussion

Consistently developed indicator frameworks such as the LDSF, combined with systematic field and lab measurement protocols, allow for spatial assessments of SOC and other land health indicators at multiple spatial scales and with unprecedented accuracy and spatial coverage. Such spatial assessments can be applied at continental and regional scales (Vågen et al., 2016), as well as at country and local scales (Figure 2 and Figure 3), with results integrated directly into 30 decision making processes at local level, as demonstrated for Turkana County (Figure 5). In Kenyan drylands, which make

up more than 80% of the area of the country and where the primary land use is livestock production, SOC concentrations and stocks are low due to a combination of climatic conditions and widespread land degradation. Land degradation in these drylands is the result of several factors, including substantial migration into these areas, which has resulted in a large growth in their population starting as early as over 30 years ago (Bernard, 1985) and soils that are alkaline and prone to soil erosion. 5 This, along with other factors, has in turn resulted in overgrazing or inappropriate agricultural practices that have again resulted in accelerated land degradation. As a result, Counties in the drylands of Kenya face serious challenges in terms of restoring degraded lands, including implementing policies and incentives that can help target restoration efforts both in terms of space and context. By applying spatial assessments of land health indicators, SOC being a key indicator in this regard, areas that are degraded can be identified for further assessments of the drivers and processes that are leading to degradation, 10 while at the same time making more realistic assessments of the types of restoration efforts that are likely to be effective given conditions in a certain area or location. of Kenyans rely on for their water supply (Mogaka et al., 2005). Also, wetland ecosystems such as inland riverine and palustrine areas are critical SOC pools in Kenya and in much of East Africa. In the drylands, such wetland systems are particularly critical for SOC storage and land health, frequently reaching between 80 and 100 Mg C ha⁻² at 0 to 30cm depth. Other wetland systems, many of them under threat, such as around the Rift Valley lakes and lacustrine wetlands along Kenya's coast are also examples of ecosystems that are critical for C storage (Minasny et al., 15 2017; Saunders et al., 2007) and other ecosystem services (Zedler and Kercher, 2005) in the country.

Land degradation in drylands is the result of several factors, including substantial migration into these areas, which has resulted in a large growth in their population starting as early as over 30 years ago (Bernard, 1985) and soils that are alkaline and prone to soil erosion. This, along with other factors, has in turn resulted in overgrazing or inappropriate agricultural practices that have again resulted in accelerated land degradation. As a result, counties in the drylands of Kenya face serious challenges in terms of restoring degraded lands, including implementing policies and incentives that can help target restoration efforts both in terms of space and context. By applying spatial assessments of land health indicators, SOC being a key indicator in this regard, areas that are degraded can be identified for further assessments of the drivers and processes that are leading to degradation, while at the same time making more realistic assessments of the types of restoration efforts that are likely to be effective given conditions in a certain area or location. 25

There is great value in using the soil health maps depicting SOC and other attributes for awareness raising and to inform decisions of stakeholders and decision makers. Building these data into the RDDST and displaying them in conjunction with data and statistics from other sectors including human health, education, pastoral economies, environment, agriculture, trade, infrastructure and energy among others, supported the facilitation of a new-found attention to the socio-ecological system 30 and associated inter-relationships and an understanding of causal dynamics among the sectors. The Ministry of Economic Planning, working with all of the government sectors and several development actors in Turkana County chose to use the SHARED methodology and the RDDST to guide the new County Integrated Development Plan (CIDP), a 5-year program (2017-2022). Based on the knowledge of the prevalence and location of land degradation in the county, there were multiple

county integrated flagships aimed at restoring land health while meeting the priorities of social and economic sectors. Further, the county government chose to cancel an earlier planned infrastructural effort that would have opened up relatively healthy land for degradation. ~~Presenting readily accessible soil health maps initiated discussion on the fundamental importance of land restoration to achieving development outcomes and lead to a commitment to bring land health management into integrated planning processes within the county. Presenting readily accessible soil health maps initiated discussion on the fundamental importance of land restoration to achieving development outcomes and lead to a commitment to bring land health management into integrated planning processes within the county.~~

5. Conclusions

Recent developments in the use of earth observation data to model and map SOC concentrations and stocks have resulted in 10 the availability of spatial estimates with high accuracy and moderate to fine spatial resolution. Decision support tools increase the utility of spatial assessments of SOC, along with other indicators of soil and land health, by providing users with both interactive dashboards that allow them to map and interact with data and analytical tools for land health diagnostics and targeting of interventions. Through the application of open source platforms and tools, the use of evidence in land health management can also be effectively mainstreamed. In the context of the SHARED approach, this process is enhanced 15 through structured stakeholder engagement, co-learning and co-design of tools. These approaches and tools have the potential to greatly enhance the uptake of management interventions to increase SOC and avoid or reverse land degradation, hence also contributing towards the achievement of LDN and SDG targets.

6. References

Aynekulu, E., Lohbeck, M., Nijbroek, R., Ordóñez, J., Turner, K., Vågen, T.-G. and Winowiecki, L. A.: Review of 20 methodologies for land degradation neutrality baselines. Sub-national case studies from Costa Rica and Namibia, 58 [online] Available from: <http://hdl.handle.net/10568/80563>, 2017.

Batjes, N. and Batjes, N.: Soil carbon stocks and projected changes according to land use and management: a case study for Kenya, Soil Use and Management, 20(3), 350–356, doi:[10.1079/SUM2004269](https://doi.org/10.1079/SUM2004269), 2004.

Bernard, F. E.: Planning and Environmental Risk in Kenyan Drylands, Geographical Review, 75(1), 58, 25 doi:[10.2307/214578](https://doi.org/10.2307/214578), 1985.

Breiman, L.: Random forests, Machine learning, 45, 35, doi:10.1023/A:1010933404324, 2001.

Caspari, T., Lynden, G. van and Bai, Z.: Land Degradation Neutrality: An Evaluation of Methods, ISRIC-World Soil Information, Wageningen., 2015.

Chang, W., Cheng, J., Allaire, J. J., Xie, Y. and McPherson, J.: shiny: Web Application Framework for R. [online] Available from: <https://cran.r-project.org/package=shiny>, 2017.

5 Chesterman, S. and Neely, C.: Evidence [into decision making for resilience planning and policy implications of Climate-Smart Agriculture in Turkana County: Stakeholder Approach for Risk Informed and Evidence Based Decision Making \(SHARED\)](#).Kenya, 112, 2015.

Dale, V. H. and Beyeler, S. C.: Challenges in the development and use of ecological indicators, *Ecological Indicators*, 1(1), 3–10, doi:10.1016/S1470-160X(01)00003-6, 2001.

Heink, U. and Kowarik, I.: What are indicators? On the definition of indicators in ecology and environmental planning, *Ecological Indicators*, 10(3), 584–593, doi:10.1016/j.ecolind.2009.09.009, 2010.

10 IUCN: Land Degradation Neutrality. Implications and opportunities for conservation, Nairobi, Kenya., 2015.

Liaw, A. and Wiener, M.: Classification and Regression by randomForest, R news [online] Available from: [ftp://131.252.97.79/Transfer/WFRE\(_ \)Articles/Liaw\(_ \)02\(_ \)Classification_and_regression_by_randomForest.pdf](ftp://131.252.97.79/Transfer/WFRE(_)Articles/Liaw(_)02(_)Classification_and_regression_by_randomForest.pdf), 2002.

15 Lohbeck, M., Winowiecki, L., Aynekulu, E., Okia, C. and Vågen, T.-G.: Trait-based approaches for guiding the restoration of degraded agricultural landscapes in East Africa, edited by M. Isaac, *Journal of Applied Ecology*, 55(1), 59–68, doi:10.1111/1365-2664.13017, 2018.

Minasny, B., Malone, B. P., McBratney, A. B., Angers, D. A., Arrouays, D., Chambers, A., Chaplot, V., Chen, Z.-S., Cheng, K., Das, B. S., Field, D. J., Gimona, A., Hedley, C. B., Hong, S. Y., Mandal, B., Marchant, B. P., Martin, M., McConkey, B. G., Mulder, V. L., O'Rourke, S., Richer-de-Forges, A. C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I.,

20 Stolbovoy, V., Stockmann, U., Sulaeman, Y., Tsui, C.-C., Vågen, T.-G., Wesemael, B. van and Winowiecki, L.: Soil carbon 4 per mille, *Geoderma*, 292, 59–86, doi:10.1016/j.geoderma.2017.01.002, 2017.

Mogaka, H., Gichere, S., Davis, R. and Hirji, R.: Climate Variability and Water Resources Degradation in Kenya, The World Bank., 2005.

Munoz, P.: Land Degradation Neutrality (LDN) in Kenya. The Economic Case, UNCCD, Nairobi, Kenya., 2016.

25 Neely, C.: Transitioning towards Climate-Smart Agriculture in Kenya. *Linking Research, Practice and Policy*, 6, 2014.

Niemeijer, D. and Groot, R. S. de: A conceptual framework for selecting environmental indicator sets, *Ecological Indicators*, 8(1), 14–25, doi:10.1016/j.ecolind.2006.11.012, 2008.

Oba, G., Stenseth, N., BioScience, W. L. and 2000, U.: New perspectives on sustainable grazing management in arid zones of sub-Saharan Africa, *BioScience*, 50(1), 35–51, doi:10.1641/0006-3568(2000)050[0035:NPOSGM]2.3.CO;2, 2000.

[R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria. \[online\] Available from: <https://www.r-project.org/>, 2018.](https://www.r-project.org/)

Saunders, M. J., Jones, M. B. and Kansiime, F.: Carbon and water cycles in tropical papyrus wetlands, *Wetlands Ecology and Management*, 15(6), 489–498, doi:[10.1007/s11273-007-9051-9](https://doi.org/10.1007/s11273-007-9051-9), 2007.

5 Stavi, I. and Lal, R.: Achieving Zero Net Land Degradation: Challenges and opportunities, *Journal of Arid Environments*, 112(PA), 44–51, doi:[10.1016/j.jaridenv.2014.01.016](https://doi.org/10.1016/j.jaridenv.2014.01.016), 2015.

[United Nations General Assembly: Transforming our world: The 2030 agenda for sustainable development, <https://sustainabledevelopment.un.org/content/documents/7891Transforming%20Our%20World.pdf>, \(1\), 1–5, doi:\[10.1007/s13398-014-0173-7\]\(https://doi.org/10.1007/s13398-014-0173-7\), 2015.](https://sustainabledevelopment.un.org/content/documents/7891Transforming%20Our%20World.pdf)

10 Vågen, T.-G., Davey, F. and Shepherd, K. D.: Land health surveillance: Mapping soil carbon in Kenyan rangelands, in *Agroforestry - the future of global land use*, edited by P. R. Nair and D. Garrity, pp. 455–462, Springer., 2012.

Vågen, T.-G., Winowiecki, L. A., Abegaz, A. and Hadgu, K. M.: Landsat-based approaches for mapping of land degradation prevalence and soil functional properties in Ethiopia, *Remote Sensing of Environment*, 134, 266–275, doi:[10.1016/j.rse.2013.03.006](https://doi.org/10.1016/j.rse.2013.03.006), 2013a.

15 Vågen, T.-G., Winowiecki, L. A., Tamene Desta, L. and Tondoh, J. E.: the Land Degradation Surveillance Framework (LDSF) - Field Guide, 14 [online] Available from: <http://landscapeportal.org/documents/705>, 2013b.

[Vågen, T.-G., Winowiecki, L.A.: Mapping of soil organic carbon stocks for spatially explicit assessments of climate change mitigation potential. *Environ. Res. Lett.* 8, 15011. doi:\[10.1088/1748-9326/8/1/015011\]\(https://doi.org/10.1088/1748-9326/8/1/015011\), 2013.](https://doi.org/10.1088/1748-9326/8/1/015011)

20 Vågen, T.-G., Winowiecki, L. A., Tondoh, J. E., Desta, L. T. and Gumbrecht, T.: Mapping of soil properties and land degradation risk in Africa using MODIS reflectance, *Geoderma*, 263, 216–225, doi:[10.1016/j.geoderma.2015.06.023](https://doi.org/10.1016/j.geoderma.2015.06.023), 2016.

Winowiecki, L., Vågen, T.-G. and Huisng, J.: Effects of land cover on ecosystem services in Tanzania: A spatial assessment of soil organic carbon, *Geoderma*, 263, 274–283, doi:[10.1016/j.geoderma.2015.03.010](https://doi.org/10.1016/j.geoderma.2015.03.010), 2016a.

25 Winowiecki, L., Vågen, T.-G., Massawe, B., Jelinski, N. A., Lyamchai, C., Sayula, G. and Msoka, E.: Landscape-scale variability of soil health indicators: effects of cultivation on soil organic carbon in the Usambara Mountains of Tanzania, *Nutrient Cycling in Agroecosystems*, 105(3), 263–274, doi:[10.1007/s10705-015-9750-1](https://doi.org/10.1007/s10705-015-9750-1), 2016b.

Zedler, J. B. and Kercher, S.: WETLAND RESOURCES: Status, Trends, Ecosystem Services, and Restorability, *Annual Review of Environment and Resources*, 30(1), 39–74, doi:[10.1146/annurev.energy.30.050504.144248](https://doi.org/10.1146/annurev.energy.30.050504.144248), 2005.

Figures




Figure 1 The distribution of SOC stocks to 30cm depth in Kenya. The vertical dashed line shows the mean (42 Mg ha⁻¹) for the country.




Figure 2. Soil organic carbon (SOC) map of Kenya at 500m resolution (for 2012). Turkana County is located in the north-western corner of the country, and is outlined on the map. [Adapted from Vågen et al. \(2016\)](#).




Figure 3. Histograms showing the distribution of SOC in Turkana County (green) versus the rest of Kenya (red). The vertical dashed line represents a SOC concentration of 15 g kg^{-1}

TURKANA COUNTY

HOME

EDUCATION

SECURITY

LIVESTOCK

LAND HEALTH

ENERGY

IRRIGATION

HEALTH

NUTRITION

WASH

TOURISM



Resilience diagnostic and decision support tool

We welcome you to explore the Turkana Resilience and Decision Support Tool. The tool is structured with **thematic modules** such as nutrition, education, land health, irrigation and education, allowing for easy visualization of data and for information on these themes to assist with decision making and resilience planning. The Turkana County Government is in the process of developing an information and data management system to allow for easy access and visualization of County data for all stakeholders and the tool is continually being updated with new data and information.

If you have data or information, please get in touch with the Ministry of Finance and Economic Planning



The tool process

1. Data capture, digitising and cleaning

2. Data synthesis/analysis

3. Interactive analysis and data visualization

4. Evidence outputs

SHARED Methodology for Evidence Based Decision Making

Our partners

In partnership with the World Agroforestry Centre (ICRAF), UNICEF and with support from USAID AHADI, Turkana County Government is committed to evidence based decision making and planning. Using the SHARED methodology, the Ministry of Economic Planning and Development has the resources to allow for easy access to evidence, to assist County decision makers and stakeholders.



Figure 4. The front page of the Turkana Resilience diagnostic and decision support tool.




Figure 5. The SOC assessment part of the Turkana Resilience diagnostic and decision support tool, which is part of the land health section of the tool.