

Challenges of soil carbon sequestration in NENA Region

Darwish^{1,3*}, Talal², Atallah², ThereseThérèse²; and Ali Fadel¹

1. National Council for Scientific Research, Beirut, Lebanon
 2. Faculty of Agricultural and Veterinary Sciences, Lebanese University
 3. Intergovernmental Technical Panel on Soil (ITPS)
- * Corresponding author: tdarwich@cnrs.edu.lb

Abstract

Near East North Africa (NENA) region spans over 14% of the total surface of the Earth and hosts 10% of its population. Soils of the NENA region are mostly highly vulnerable to degradation, and food security will depend much on sustainable agricultural measures. Weather variability, drought and depleting vegetation are dominant causes of the decline in soil organic carbon (SOC). In this work the ~~situation-status~~ of SOC was studied, using a land capability model and soil mapping. The land capability model showed that most NENA countries (17 out of 20), suffer from low productive lands (>80%). Stocks of SOC were mapped (1:5 Million) in topsoils (0-0.30 m) and subsoils (0.30-1.00 m). The maps showed that 69% of soil resources present a stock of SOC below the threshold of 30 tons ha⁻¹. The stocks varied between ≈ 10 tons ha⁻¹ in shrublands and 60 tons ha⁻¹ for evergreen forests. Highest stocks were found in forests, irrigated crops, mixed orchards and saline flooded vegetation. The stocks of soil inorganic carbon (SIC) were higher than those of SOC. -In subsoils, the SIC ranged between 25 and 450 tons ha⁻¹, against 20 to 45 tons ha⁻¹ for SOC. ~~This paper also highlights~~Results highlight the ~~modest~~ contribution of NENA region to global SOC stock in the topsoil ~~not exceeding~~(<4.1%). The paper also discusses agricultural practices that are favorable to carbon sequestration such as: Practices of conservation agricultureOrganic amendment, no till or minimum tillage, crop rotation, and mulching and the constraints caused by geomorphological and climatic conditions. ~~Further, could be effective, as the presence of soil cover reduces the evaporation, water and wind erosions. In semi arid east Mediterranean~~Further, the introduction of legumes, as part of a cereal legume rotation, and the application of nitrogen fertilizers to the cereal, caused a notable increase of SOC after 10 years. The effects of crop rotations on SOC are related to the amounts of above and belowground biomass produced and retained in the system. Some knowledge gaps exist, especially in aspects related to the impact of climate change and effect of irrigation on SOC, and on SIC at the level of soil profile and soil landscape. Still, major constraints facing soil carbon sequestration are policy relevant and socio-economic in nature, rather than scientific.

Keywords: Drylands, soil organic carbon, soil inorganic carbon, land capability, C stock, conservation practices.

1. ~~1.~~ Introduction

The Near East North Africa (NENA) region spans over 14% of the total surface of the Earth and hosts 10% of its population (Elhadi, 2005). The largest importer of wheat in the world, this region is also one of the poorer (FAO, 2015). A recent assessment of global

Formatted: Font: (Default) Times New Roman, Complex Script Font: Times New Roman

Formatted: Font: (Default) Times New Roman, 14 pt, Complex Script Font: Times New Roman, 14 pt

Formatted: List Paragraph, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"

Formatted: Indent: Before: 0.25"

hunger index (GHI), based on four indicators -undernourishment, child wasting, child stunting, and child mortality- showed that most of the NENA countries present low to moderate GHI. Countries suffering from armed conflicts, Syria, Iraq and Yemen, are at a serious risk (von Grebmer et al., 2017). With the scarce natural resources and difficult socio-economic conditions, it is questionable whether food security will be reached by 2030, unless a significant change in agricultural practices and governance occurs (FAO, 2017).

Formatted: Indent: First line: 0"

Most of the land area of the NENA region falls in the hyper-arid, arid and semi-arid climatic zones. Climate change is expected to exacerbate the scarcity of water, ~~and drought-and drought effect~~. Weather variability, drought and depleting vegetation are major concerns in the loss of soil productivity and agricultural sustainability. ~~Instabilities~~ Changes in soil organic carbon (SOC) can affect the ~~density-emission~~ of greenhouse gases ~~in-to~~ the atmosphere and negatively ~~affect-influence~~ the global climate ~~change~~ (Lal, 2003). In fact, destructive land management practices are impacting soil functions. Land use change, mono-cropping and frequent tillage are considered to cause a rapid loss of SOC (Guo et al., 2016). These agricultural practices ~~disrupt-disturb~~ the stability of ~~inherited~~ soil characteristics, built under local land cover and climate (Bhogal et al. 2008). Thus, most NENA lands contain ~1 % of SOC, and frequently less than 0.5%.

Formatted: Indent: First line: 0"

Formatted: Indent: First line: 0.5"

~~Despite the constraints of~~ NENA pedo-climatic conditions present major constraints to carbon sequestration, increasing SOC levels is critical and challenging (Atallah et al., 2015). To maintain soil productivity and land quality, several technical and socio-economic measures need to be adopted. Additional efforts oriented to maintaining and increasing ~~-SOC, SOC~~ can contribute to poverty reduction and achieve food security (Plaza-Bonilla et al., 2015). Good agricultural practices, based on low tillage or no tillage, may result in the reduction of SOC breakdown and the enhancement of the soil carbon pool (Atallah et al., 2012; Cerdá et al., 2012; Boukhoudoud et al., 2016).

Formatted: Indent: First line: 0"

Formatted: Indent: First line: 0.5"

Formatted: Indent: First line: 0"

In order to compare situations and problems, global soil organic carbon maps are a priority. As recently as December 2017, the GSP-FAO, ITPS launched the version 1 of the

global soil organic carbon map, showing the SOC stock in topsoil (<http://www.fao.org/3/a-i8195e.pdf>). A preliminary assessment of regional SOC stocks, using unified background information, is needed to analyze the challenges facing C sequestration. Quantifying SOC content in the NENA countries using available soil data is crucial, even at a small scale, to assess the nature and potential of available soil resources and analyze the associated threats. Mapping the spatial distributions of national and regional OC stocks can be used to monitor and model regional and global C cycles under different scenarios of soil degradation and climate change. Accurately quantifying SOC stocks in soils and monitoring their changes are considered essential to assessing the state of land degradation. At the same time, the predominantly calcareous soils of NENA region are rich in soil inorganic carbon (SIC). The dynamics of SIC and its potential in sequestering soil carbon in soils are remain largely unknown and as such deserves thorough investigation. This paper analyzes the state of SOC and SIC in NENA countries and outlines challenges and barriers for devising organic carbon sequestration in NENA's impoverished and depleted soils. It also highlights several questions which scientists need to resolve. Finally, it discusses practical agricultural measures to promote SOC sequestration.

2. Materials and Methods

Data on SOC and soil inorganic carbon (SIC) contents in soils of the NENA region were retrieved from the soil database of the FAO-UNESCO digital soil map of the world (DSMW) at 1:5 Million. Within The the database, contains large number of 1700 geo-referenced soil profiles, collected and harmonized from each all member states of the NENA region, are. These were excavated, sampled by horizon, down to the rock, and analyzed in the laboratory according to the standard world accepted methods included (FAO, 2007).

In terms of area, the largest soil units are Yermosols (4670.6 km²), Lithosols (2914.3 km²) and Regosols (1193.2 km²). The great majority of soil classes presents very low to low resistance to erosion and degradation (Table 1). To the contrary, Cambisols, Fluvisols and

Formatted: Font: Bold, Complex Script Font: Bold

Formatted: Font: (Default) Times New Roman, 12 pt, Bold, Complex Script Font: Times New Roman, 12 pt, Bold

Formatted: List Paragraph, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"

Formatted: Font: Bold, Complex Script Font: Bold

Formatted: Space After: 10 pt, Line spacing: Multiple 1.15 li, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Formatted: Font: (Default) Times New Roman, 12 pt, Bold, Complex Script Font: Times New Roman, 12 pt, Bold

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Regosols are highly resistant to erosion. ~~For the SIC, soil classes dominated by calcareous rocks (Solonchaks, Rendzinas and Aridisols) have the highest SIC contents (Table 1), while Lithosols and Xerosols, subject to regular water and wind erosion, show the smallest SIC.~~

Table 1. Soil inorganic carbon level and the resistance to land degradation in the major soil units of Near East North Africa region (Source: DSMW, FAO, 2007).

Soil Classes	Area, 1000 km ²	Resistance to Land Degradation	Average SIC content (%)	
			topsoil	subsoil
Cambisols	178.9	High	0.25	0.64
Fluvisols	232.7	High	1.12	1.40
Kastanozems	26.0	High	1.69	3.96
Regosols	1193.2	High	1.18	0.23
Luvisols	121.6	Moderate	0.02	0.11
Phaeozems	3.8	Moderate	0.40	0.70
Rendzinas	25.6	Low	2.80	4.80
Lithosols	2914.3	Low	0.01	0.06
Vertisols	45.4	Low	0.45	0.72
Xerosols	498.5	Low	0.25	0.45
Yermasols (Aridisols)	4670.6	Low	2.50	2.30
Solonchaks	230.1	Very low	3.60	3.90
Solonetz	31.2	Very low	0.06	0.36
Arenosols	384.0	Very low	0.00	0.00

~~The soil map was prepared using the topographic map series of the American Geographical Society of New York, as a base, at a nominal scale of 1:5.000.000. Country boundaries were checked and adjusted using the FAO UNESCO Soil Map of the World, on the basis of FAO and UN conventions. Soil classification was based on horizon designation, depth, texture, slope gradient and soil physico-chemical and chemical properties. Statistical (weighted) average was calculated for the topsoil (0-30 cm) and for the subsoil (30-100 cm) for the full series of chemical and physical parameters sufficient to assess main agricultural soil properties. To fill the gap in some attributes and complete the fields for which no data were available, an expert opinion internationally known soil scientists was used.~~

Formatted: Right

Formatted: Right

Formatted: Right

Formatted: Right

Formatted: Right

Formatted: Right

Formatted: Right

Formatted: Right

Formatted: Right

Formatted: Right

Formatted: Right

Formatted: Right

Formatted: Right

Formatted: Right


Formatted: Space After: 10 pt, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Formatted: Indent: First line: 0.5"

Using the DSMW and its updated attribute database, the maps of the SOC and SIC stock and distribution in 20 NENA states were produced. The scale used in the DSMW is 1:5 Million (FAO, 2007). The soil map was prepared using the topographic map series of the American Geographical Society of New York, as a base, at a nominal scale of 1:5,000,000. Country boundaries were checked and adjusted using the FAO UNESCO Soil Map of the World, on the basis of FAO and UN conventions. To produce the maps representing the spatial distribution of SOC and SIC, ArcMap 10.3 was used to join the symbology of the C stocks and density with quantities classified into five 0-14, 14-22, 22-31, 31-61 and 61-236 tons/ha for SOC density and 0, 0.1-30, 30-45, 46-74 and 74-200 tons/ha for SIC density numerical categories with natural breaks. The global LC maps at 300-m spatial resolution on an annual basis from 1992 to 2015 was produced by ESA. The Coordinate Reference System used is a geographic coordinate system based on the World Geodetic System 84 (WGS84) reference ellipsoid. The legend assigned to the global LC map has been defined using the UN Land Cover Classification System.

The SOC content in studied soils varied between values as low as 0.13% and 0.16% and as high as 1.74% and 0.9% in topsoils and subsoils of Yermosols (Aridisols) and Rendzinas (Mollisols) respectively (Table 1). Worth noting that less than 20% of soil resources in NENA region have sequestered and accumulated SOC to an extent exceeding 1.0%. The SOC content can vary depending on soil type, topography, land cover, erosion-sedimentation and soil management. Within the topsoils (0-0.30 m), the SOC contents are between 0.13% and 1.74%, while in the subsoils (0.30-1 m) values range between 0.16 and 0.9% (Figure 1). Two out of the three predominant soil classes (Xerosols and Aridisols) have SOC contents below 0.5%. Overall, the NENA soils are poor in SOC, as less than 20% of soil resources have SOC contents above 1.0%.

Formatted: Indent: First line: 0"



Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Figure -1. SOC content (%) in topsoil and subsoil in the major soil groups of NENA region with standard deviation related to soil class.

The NENA soil map was prepared using the topographic map series of the American Geographical Society of New York, as a base, at a nominal scale of 1:5,000,000. Country boundaries were checked and adjusted using the FAO-UNESCO Soil Map of the World. Soil classification was based on horizon designation, depth, texture, slope, and soil physico-chemical properties. Main agricultural soil properties were assessed using the statistical (weighted) average in the topsoil and subsoil. For the production of the maps of C stocks and distribution (FAO, 2007), ArcMap 10.3 was used to join the geometric database with the C stocks. These were ranked into five categories for SOC density (0-14, 14-22, 22-31, 31-61 and 61-236 tons/ha) and five others for SIC (0, 0.1-30, 30-45, 46-74 and 74-200 tons/ha).

The land cover map was that of ESA, at 300 m spatial resolution. The Reference Coordinate System used was a geographic coordinate system based on the World

Geodetic System 84 (WGS84) reference ellipsoid. The legend assigned to the global LC map has been defined using the UN-Land Cover Classification System.

To assess the potential soil productivity in the NENA region, a land capability model proposed by USDA (1999), which includes the soil geomorphological features (geology and topography), other soil physico-chemical parameters conditioning soil fertility like soil depth, texture, organic matter content, salinity and sodicity hazards was adopted. The soils of the area were classified into four classes of arable soils: class I (highly productive), class II (medium productive), class III (low productivity) and class IV (very low productivity) and one non-arable soil class V, where lands suitable for wild vegetation and recreation and lands with rock outcrops were grouped.

Map 10.1 was used for the mapping of soil types and OC stock and density of each soil unit based on the geographic or spatial distribution of the soil type. Total SOC and SIC stocks and the stock of SOC were calculated separately for the topsoil (0-0.3m) and subsoil (0.3-1.0m) using the following equations:

$$\text{Total National OC Stock (ton)} = [\text{Area (m}^2\text{)} * \text{Depth (m)} * \text{Bulk Density (ton m}^{-3}\text{)} * \text{OC content (\%)}] / 100 \quad \text{— equation (1)}$$

$$\text{Stock SOC or SIC density (ton ha}^{-1}\text{)} = \text{Stock in soil unit (ton)} / \text{Soil Unit Area (ha)} \quad \text{(equation-2)}$$

The stocks of SOC under different land cover/land use were evaluated, as well. Since 1990, the European Space Agency, (Climate Change Initiative project), started to produce land cover (LC) maps of the NENA region. The version used in the study (Website 1) corresponds to the second phase of the 2015 global LC map (<http://maps.elie.ucl.ac.be/CCI/viewer/index.php>) with a spatial resolution of 300 m of spatial resolution, using the Coordinate Reference System (CRS) in a geographic coordinate system (GCS) based on the World Geodetic System 84 (WGS84) reference ellipsoid.

The legend assigned to the global LC map has been defined using the UN-LCCS.

3. Results and Discussion

3.1. Land capability and SOC in NENA region

Formatted: Indent: First line: 0"

Formatted: Indent: First line: 0"

Formatted: Centered

Formatted: Line spacing: 1.5 lines

Field Code Changed

Formatted: Default Paragraph Font, Font: (Default) +Body (Calibri), 11 pt, Complex Script Font: 11 pt

Formatted: Indent: First line: 0"

Formatted: Line spacing: 1.5 lines

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: List Paragraph, Indent: Hanging: 0.5", Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 3 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.5"

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: List Paragraph

The most abundant soil classes in the NENA region are Arenosols, Xerosols and Aridisols representing together more than 80% of available soil resources (FAO, 2007). All three soil classes have low resilience resistance to degradation (Table 1). According to the model results based on the land capability model, showing the proportion (%) of different soil productivity classes in each country of NENA region, the majority of soils (40 to 100% of soils in the) region belongs to fall within the low, very low and non-arable classes (Figure 42). Thus, the proportion of highly and medium productive soils varies between 0% (Bahrain, Qatar, Oman and UAE) and 60% (Jordan). Countries like Iraq, Lebanon, Morocco, Palestine, Somalia, Syria and Tunisia, present between 9 and 20% of highly to medium productive soils. The remaining NENA countries have less than 5% of their lands as high and medium productive. Some of these countries belong to the food insecure nations.




Figure 42. Distribution of land capability classes (% of total national area) Land-capability classification for the 20 countries of the NENA region, based on the USDA M-model (1999) and Digital Soil Map of the WorldSMW₁ (FAO, 2007).

Thus, the proportion of highly and medium productive soils varies between 0% (Bahrain, Qatar, Oman and UAE) and 60% (Jordan). Countries like Iraq, Lebanon, Morocco, Palestine, Somalia, Syria and Tunisia present between 9 and 20% of highly to medium

productive soils. The rest of NENA countries have less than 5% of their lands as high and medium productive soils. Some of these countries belong to the seriously endangered and food unsecure nations.

Formatted: Strikethrough

The ~~potential medium soil~~ productivity concept is based strictly on soil properties. But, with the lack of water in drylands and the prevalence of rainfed agriculture, the soil cannot show its full potential for food production. Similarly, irrigation with brackish and saline water restricts crop productivity due to the development of secondary soil salinity. With ~~hen properly managed irrigated~~, the medium productive lands can provide moderately good harvests. For instance, our field observation in Jordan showed that ~~d that due to climate change and climate variability, a large area of good productive lands was cropped with barley, not because of land suitability, but due to low rainfall (< 200 mm). In drought affected years, the land is converted into grazing area for small ruminants following crop failure to make the minimal profit from the exploitation. The presence of kaolinite in red soils of Jordan developed from hard limestone under semi arid climate points to the inheritance of material formed under more aggressive climate (Kusus and Ryan, 1985). The same was confirmed by Lucke et al., 2013 for Red Mediterranean Soils of Jordan, which require new insights in their origin, genesis and role as a source of information on paleoenvironment.~~

The low productivity of the soil is reflected in the SOC contents. ~~Two out of the three predominant soil classes (Xerosols and Aridisols) have SOC contents below 0.5% (Table 1). Overall, the NENA soils are poor in SOC, as less than 20% of soil resources have SOC contents exceeding 1.0%.~~ The accumulation of SOC in NENA region is refrained by the high mineralization rate (Bosco et al., 2012). Climate change and recurrent drought events affect SOC sequestration in the soil. It is estimated that a rise in temperature of 3 °C would increase the emission of carbon dioxide by 8% (Sharma et al., 2012).

Formatted: Indent: First line: 0"

Among the soil properties affecting SOC, the clay and calcium carbonate contents are most relevant. ~~High Clay clay fraction content~~ tends to counteract the decomposition of SOC, as found in clay soils of Morocco and in Vertisols of northern Syria (FAO and ITPS, 2015). But, the dominant soil classes, ~~Xerosols, Aridisols or Arenosols~~ (Table 1) characterized by sandy and sandy loam textures, are subject to fast decomposition.

Formatted: Indent: First line: 0"

Next to the clay texture, the presence of calcium carbonate decreased the decomposition of composted organic material in sub-humid coastal Lebanon (Al Chami et al., 2016). This slower turnover of organic matter was explained by the low porosity and prevalence of micropores in soil macro-aggregates (Fernández-Ugalde et al., 2014). ~~For the SIC, the highest values are found in soil classes dominated by calcareous rocks, that is the Solonchaks, Rendzinas and Aridisols (Table 1). The lowest stocks were detected in Lithosols and Xerosols, subject to water and wind erosions that remove the surface layer of eroded lands.~~

3.2. Mapping of soil carbon stocks ~~in the soils of NENA region~~

~~In order to compare situations and problems, global soil organic carbon maps are a priority. As recently as A preliminary assessment of regional SOC stocks, using unified background information, needed to analyze the challenges facing C sequestration.~~

~~Thus, the need for more detailed and harmonized soil mapping and coding of available national information arises to downscale to national and local soil assessment and mapping, which is currently on the agenda of the Global Soil Partnership (GSP).~~

~~To obtain an idea of the status of the soil carbons, assessed under different mapping, the spatial distributions of SOC and SIC stocks for the NENA region assessed in this study at 1:5 Million was compared with a larger scale mapping (1:50,000) for Lebanon. A moderate discrepancy (11% to 14%) was found between the two scales for carbon stock in topsoil and subsoil respectively~~

~~Based on the mapping of SOC density (ton ha^{-1}), 69% of the regional soils have a stock density inferior below to 30 tons ha^{-1} (Figure 34), value considered as a threshold for C deficient soils (Batjes and Sombroek, 1997). This could be linked to the arid conditions prevailing in the region with flat lands and limited humid mountain areas. Consequently,~~

~~The majority of the countries of NENA region presents moderate to relatively low total stocks of SOC.~~

Formatted: Indent: First line: 0"

Formatted: Not Highlight

Formatted: Indent: First line: 0", Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Indent: First line: 0"




Figure 3. Spatial distribution of SOC density (ton ha⁻¹) in topsoils (a) and subsoils (b) of the NENA region.

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Indent: First line: 0"

Formatted: Indent: First line: 0"

Field Code Changed

Formatted: Hyperlink, Font: (Default) +Body (Calibri), 11 pt, Complex Script Font: +Body CS (Arial), 11 pt, Not Highlight

Formatted: Indent: First line: 0"

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted Table

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

This is especially relevant to the Gulf countries, Iran, Tunisia and Morocco, with values below 221 Mega tons (Figure 243). Such low OC sequestration potential can be explained by the prevalence of arid climate and rare sparse natural vegetation and the reliance on irrigation to produce food and feed crops. A regional implementation plan for sustainable management of NENA soils appeared in 2017. (Website 2, <http://www.fao.org/3/a-bl105e.pdf>). In fact, the total stocks, in the topsoils (0-0.3 m) of NENA countries, represent only 4.1% of the world global stock (Website 3) https://drive.google.com/drive/folders/1454FhX_p2_GxjZT-fsi0-Rncl2QpSN.




Figure 4. Spatial view of total soil organic carbon stock (Megatons) across the countries of the NENA region. Mega-ton (a-topsoil; b-subsoil).

A comparison has been undertaken between the FAO methodology, adopted in the present work (two soil layers: 0-0.3 m; 0.3-1 m), and a previous study where the soil profile was divided into six depths down to 2 m (Hengl et al., 2014). Both approaches agree about the NENA region (SOC content: 1-2% and SOC stock: 20-204 ton ha⁻¹), as confirmed by the Global Soil Organic Carbon Map (<http://www.fao.org/global-soil-partnership/pillars-action/4-information-and-data/global-soil-organic-carbon-gsoc-map/en/>). The global gridded soil information based on machine learning (SoilGrid250m) contains 1.6 billion pixels also predicted SOC density but the arid and semi-arid zones are under measured and thus pseudo observations based on expert knowledge were introduced to predict the SOC content in the soil at seven standard soil depths (Hengl et al., 2017).

~~Mapping was done on a small coarse scale, which could be a source of a loss of information. The choice of scale when using or producing soil maps may lead to uncertainty in small countries and fragmented land use (Darwish et al., -2009). The coarse scale adopted in this work (1:5 Million) could be a source of uncertainty. -Practically, this corresponds to a polygon -of -0.5 cm x 0.5 cm (area < 0.25 cm²), -the equivalent of an area of 6.25 km² on the ground. -To test this, the results of the current estimation (1:5 Million) of SOC stocks in Lebanon, were compared with the large scale mapping undertaken recently to produce the unified soil map of Lebanon at 1:50,000 for Lebanon (Darwish et al., 2006+50,000). The FAO scale gave an overestimation of 11.2% in the topsoil, underestimation of 16.4% in subsoil and a 14.4% underestimation in the whole profile. This comparison showed discrepancies between 11% for the topsoil and 14% for the subsoil (Darwish and Fadel, 2017). Therefore, the level of mapping uncertainty falls within the reported admitted diagnostic power of soil maps, with a matching close to map purity of 65-70%. -Therefore, the level of uncertainty falls within the admitted diagnostic power of soil mapping, estimated to be close enough to the reported range of map units' purity in reference areas, i.e., a matching between 65% and 70% (Finke et al., 2000). In our study, the matching reached 83.6-88.2%. Any Loss-loss of information related to small, non-mappable soil units in small coarse scale mapping (1:5 Million or 1:1 Million)~~

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Centered, Space After: 0 pt

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Indent: First line: 0"

Formatted: Indent: First line: 0"

Formatted: Font: (Default) +Headings CS (Times New Roman), Complex Script Font: +Headings CS (Times New Roman)

Formatted: Font: (Default) +Headings CS (Times New Roman), Complex Script Font: +Headings CS (Times New Roman)

could be ~~corrected~~rectified by national and sub-regional large scale soil mapping (1:50,000 and 1:20,000).

~~The SOC content in studied soils varied between values as low as 0.13% and 0.16% and as high as 1.74% and 0.9% in topsoils and subsoils of Yermosols (Aridisols) and Rendzinas (Mollisols) respectively (Table 1). Another source of uncertainty-error can be associated with the used average SOC content in soil classes or major groups from the DSMW (FAO, 2007), containing several soil types. The level of uncertainty in the assessment of the SOC density in NENA countries depends on the variability le content of of SOC, as suggested by them a given soil unit or type that can represent large standard deviations of the means (Figure 41). Soil classification considers several soil forming factors including soil depth, horizon designation and evolution and topographic location. The SOC content can vary depending on soil type location on slopping or level lands, land cover, erosion-sedimentation and soil management. Therefore, the SOC content is the major source of variability forfor the SOC density when assessing at the at higher classification levels (soil class level.); with Cambisols and Fluvisols showpresenting the largest standard deviation, caused by a long land use history and large anthropologicanthropogenic impact. Subsoil is less subject to pedo-turbation and direct human influence, thus SOC content has lower variability.~~

3.3. Land cover mapping and ~~stocks of~~effect on SOC stock


~~Based on the land cover map of ESA, the bare lands correspond to Land cover map of NENA region shows nearly 80% of the whole NENA regionarea is covered with bare lands (Figure 45a, b). Grassland, sparse vegetation cover and rainfed agriculture representare close by area varying between 4.39% and 5.27%. The irrigated crops do not exceed 0.66% of the total area, distributed in limited usefulcultivated area (Figure 65). The NENA region possesses a land area about 15 million km², with a total population exceeding 400 million inhabitants (about 6% of world population) but with only 1% of the world's renewable water resources (<https://www.slideshare.net/FAOoftheUN/plenary1-keynote-speech-16dec2013az>). Apparently for this reason, TheIne fact that factthe NENA region, represents anthe irrigated area equivalentcorresponds to to 247.5 m² per capita. This iscould be one of the reasons for the high dependency on imported food, exacerbated by-region is becoming increasingly dependent on food imports, because of demographic pressure, rapid urbanization, water scarcity and climate change (FAO, 2015).~~

Formatted: Indent: First line: 0"

Formatted: Indent: First line: 0"

Formatted: Superscript

Formatted: Superscript



*Others: Mixed trees, Needle-leaved, Evergreen trees and Urban

Figure 5. Proportion of main land cover (% total land area) and land use in NENA region. (Source: <http://maps.elie.ucl.ac.be/CCI/viewer/index.php>, ESA, 2015)




Figure 6. Land cover map of NENA region (Source: ESA, 2015; <http://maps.elie.ucl.ac.be/CCI/viewer/index.php>)

Formatted: Indent: First line: 0.5", Space After: 0 pt, Line spacing: single

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Centered, Space After: 0 pt, Line spacing: single

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Text 1, Complex Script Font: Times New Roman, 12 pt

Formatted: Font: 12 pt, Font color: Text 1, Complex Script Font: 12 pt

Formatted: Font: 12 pt, Complex Script Font: 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Line spacing: single

392 ~~Comparing our results with the global SOC map produced by Hengl et al., (2014, based on~~
 393 ~~soilGrids1km layers showing the soil organic carbon content in permille in 0-5 cm and the~~
 394 ~~predicted global distribution of the soil organic carbon stock in tonnes per ha for 0-200 cm~~
 395 ~~to be beyond the followed by FAO methodology of SOC stock estimation and~~
 396 ~~presentation. In this paper the standard methodology of the measured SOC stock and~~
 397 ~~density in topsoil (0-30 cm) and subsoil (30-100 cm) was followed. The first Global SOC~~
 398 ~~Map was launched on December 5, 2017. However, a comparison of values of SOC~~
 399 ~~content (%) and SOC stock revealed comparable trends values for the C content and stock~~
 400 ~~(1-2% and 20-204 ton/ha), with higher upper density in Hengl et al approach. Organic~~
 401 ~~Carbon ([http://www.fao.org/global-soil-partnership/pillars-action/4-information-and-data/global-](http://www.fao.org/global-soil-partnership/pillars-action/4-information-and-data/global-soil-organic-carbon-gsoc-map/en/)~~
 402 ~~[soil-organic-carbon-gsoc-map/en/](http://www.fao.org/global-soil-partnership/pillars-action/4-information-and-data/global-soil-organic-carbon-gsoc-map/en/)).~~

Formatted: Indent: First line: 0"

Formatted: Font: 11 pt, Complex Script Font: 11 pt

403
 404 ~~FurtherIn our studyT,~~ the combination of SOC stock map with the land cover map
 405 ~~showed, a the significant effect of land cover on -SOC stocks in NENA region. of SOC~~
 406 ~~were studied in relation to land cover/land use. As can be expected, Shrublands shrublands,~~
 407 sparse vegetation and bare lands gave the smallest values, between 14 and 26 ton ha⁻¹
 408 (Figure ~~76~~). In a mixture of shrublands and herbaceous vegetation, the SOC increases to
 409 40 ton ha⁻¹. The highest density (30 and 60 ton ha⁻¹) is found under forest stands.

Formatted: Indent: First line: 0"

410
 411 Despite the expected impact of frequent plowing, the soils under mixed trees and irrigated
 412 crops have ~~ve~~ higher SOC density than rainfed crops. ~~This could be linked to the higher~~
 413 ~~biomass produced under irrigated conditions in these water-limited areas. The highest~~
 414 ~~SOC stock was~~ observed under evergreen forests ~~land~~ whose area is very limited (3380
 415 ~~km² corresponding to 0.02% of from the total area~~). Surprisingly, the stock found under
 416 urban soils (≈ 30 tons ha⁻¹) ~~was is~~ moderate. This could be related to the urban
 417 encroachment on prime soils. Between 1995 and 2015, rapid urban growth caused the loss
 418 of over 53 Million tons of soils, 16% of which correspond to prime soils (Darwish and
 419 Fadel, 2017). ~~The assessment of SOC content in NENA region showed a decline of OC~~
 420 ~~time and space in relation to land cover showed a decline of OC content in topsoil by up to~~
 421 ~~1% between 2001 and 2009 (Stockman et al., 2015). L~~Overtime, ~~L~~and cover change
 422 ~~was~~ considered as the primary agent of change ~~of that influenced SOC change overtime,~~
 423 ~~followed by temperature and precipitation.~~

Formatted: Left, Indent: Before: 0", First line: 0"

Formatted: Font: (Default) +Headings CS (Times New Roman), 12 pt, Complex Script Font: +Headings CS (Times New Roman), 12 pt

Formatted: Font: (Default) +Headings CS (Times New Roman), 12 pt, Complex Script Font: +Headings CS (Times New Roman), 12 pt

Formatted: Superscript

Formatted: Font: (Default) +Body (Calibri), 11 pt, Complex Script Font: +Body CS (Arial), 11 pt




Figure 76. SOC stock density (tons ha⁻¹) in the topsoils (0-0.3 m) and subsoils (0.3-1.0 m), calculated from the FAO DSMW (FAO, 2007), under corresponding land cover (<http://maps.elie.ucl.ac.be/CCI/viewer/index.php>) as related.

In addition to the stocks of SOC in relation to land cover and use, the stocks of SOC and SIC were established per country (Figure 875). The stock of SIC was compared to that of SOC (Figure 5). The range of SIC stocks is very wide, from less than 25 tons ha⁻¹ (Gaza subsoil) to 450 tons ha⁻¹ (Bahrain subsoil), while that of SOC varied between ≈ 20 tons ha⁻¹ (Bahrain subsoil) and 45 tons ha⁻¹ (Sudan subsoil). Based on the stocks of SIC in the subsoils, the countries were separated into three groups. The first, represented by six countries (Bahrain, Oman, Egypt, Saudi Arabia, UAE and Yemen) was dominated by calcareous parent materials, with values in the subsoil exceeding 200 tons ha⁻¹ (Figure 57). The second group, with eight countries (Kuwait, Libya, Iran, Iraq, Algeria, Qatar, Morocco and Tunisia), presents a SIC density between 100 and 200 tons ha⁻¹. Finally, the third group (Gaza, Jordan, Lebanon, Mauritania, Syria and Sudan) has less than 100 tons ha⁻¹.

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Centered

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Complex Script Font: Times New Roman, 12 pt, Line spacing: Multiple 1.15 li, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers

Formatted: Indent: First line: 0"





Figure 87. Stocks of soil organic carbon and soil inorganic carbon density (tons ha⁻¹) in the topsoils (0-0.3 m) and subsoils (0.3-1.0 m) of the 20 countries in the NENA region. The red line represents the threshold for organic carbon, and the yellow lines correspond to the limits of classes for inorganic carbon.

3.3. Challenges of carbon sequestration in NENA agro-ecosystems

Climatic conditions characterized by wetting/drying cycles, a long dry and hot season (Boukhoudoud et al., 2016) promote the decomposition of SOC. Further, frequent cultivation, irrigation with saline water, and soil salinity rise in coastal areas the prevailing agriculture practices (Boukhoudoud et al., 2017) exert significant effects on soil

Formatted: Indent: First line: 0"

microbial functional properties. For instance three months after the application of glyphosate-based herbicide under olive in coastal Lebanon, lipase activities significantly decreased (Boukhoudoud et al., 2017). Soil classification and SOC mapping help identifying hot spots that need to be improved or require special management measures and bright spots with satisfactory C accumulation levels that need to be protected. -In this section, ~~there will be a presentation of~~ major practices affecting SOC will be presented followed by a discussion of preventive and remediation measures.

3.3.1. Tillage and SOC

Tillage practices contribute to the vulnerability of soils to water erosion. If not properly managed, some 41 million hectares in NENA region would be affected by water erosion (FAO and ITPS, 2015). The erosion of soil surface layers can affect the soil carbon in two possible ways. The greater exposure of carbonates to climatic elements could increase the loss of SIC to the atmosphere and ground water. Compared to stable soils, ~~Also,~~ the higher decomposition of SOC in eroded soils decreases the productivity of cultivated crops and can reduce SOC stock, if not properly managed (Plaza-Bonilla et al., 2015).

A possible measure to reduce the risk of erosion is the no-tillage practice. No-tillage coupled with mulching, to reduce weed development and omit herbicide application, as part of conservation agriculture (CA), aims to return more plant residues to the soil, enhance C sequestration, increase soil aggregates, improve water infiltration and ~~to provide a protection to~~ soil carbon from decomposers (Palm et al., 2014).- Through a modification of common practices, such as the frequency and depth of tillage, changes in the SOC could be promoted in most soils. Experiments conducted by ICARDA, Syria, showed that no-tillage performed well in terms of energy and soil conservation (Plaza-Bonilla et al., 2015). Elsewhere, in Palestine soil conservation was found to pay, with a net profit 3.5 to 6 times higher than without conservation measures (FAO and ITPS, 2015). In dryland regions, agricultural activities based on CA practices are beneficial as crop residues are left on the soil surface (Plaza-Bonilla et al., 2015). The presence of residues would protect the soils from high evaporation, water and wind erosions. This is especially relevant to soils that are sensitive to degradation, such as the very shallow Lithosols, the easily-periodically wetted (swelling) and dry (shrinking) Vertisols, Gypsic Yermasols (Aridisols), the poorly-structured Solonchaks and Solonetz, the sandy-textured Arenosols, and the desert soils (Xerosols).

Formatted: Indent: First line: 0"

Formatted: Indent: First line: 0"

Major constraints facing soil conservation measures, in East Mediterranean, were due to knowledge and perception, prevailing practices of complete removal as hay or forage and some-times burning of residues after harvest, land tenure and type of landscape (FAO, 2012; FAO and ITPS, 2015). These ~~major~~ factors are socio-economic in nature, rather than scientific. They are related to the ability of growers to accept new techniques and adopt them. In many situations, the transfer from the research stations to the farmers was not smooth. For instance, CA was successfully tested in experimental stations in Morocco and Lebanon, but several social and technical barriers prevented it from reaching farmers (Mrabet et al., 2012; FAO, 2012).

A debate has been taking place about the effect of no-tillage on SOC. ~~Most~~ Many authors agree that under CA, SOC increases near the soil surface, but not necessarily throughout the profile. A study compared 100 pairs, where no-tillage has been practiced for over 5 years. The absence of tillage lead to higher C stocks (0-~~0.3~~ 30-cm soil depth) in 54% of pairs, while 39% showed no difference in stocks (Palm et al., 2014). In the absence of tillage, the slower decomposition of residues would result in higher ~~belowground~~ C accumulation on the soil surface. Over a period of 5 years, zero tillage promoted an increase in SOC equal to 1.38 Mg ha⁻¹ as compared to the conventional tillage in northern Syria (Sommer et al., 2014).

3.3.2. Agricultural practices and SOC

Practices, such as the application of N fertilizers, ~~of the~~ organic amendments, ~~the~~ incorporation of residues and crop rotations, influence the levels of SOC. In soil mining practices without minimal input of fertilizers, ~~the~~ lack of ~~available~~ accessible nutrients ~~and soil mining~~ makes most crops entirely reliant on the mineralization of accumulated SOC, ~~causing soil mining~~ (Plaza-Bonilla et al., 2015). In East Africa, 14-years of continuous cultivation without any inputs, decreased SOC from 2% to 1% (Sharma et al., 2012). The application of N fertilizers was associated with increased levels of soil C, as compared to the absence of N fertilizers (Palm et al., 2014). In a 10-year rotation of wheat-~~grain~~ grain legume (~~vetch~~) in northern Syria, the application of nitrogen fertilizers to the cereal caused a notable increase of SOC, in the top 1m of soil, equal to 0.29 Mg ha⁻¹ year⁻¹ (Sommer et al., 2014). Similarly, in semi-arid Lebanese area (Anti-Lebanon mountains), the growth of intercropped legumes as winter cover crop ~~crop legumes (Vicia sp., Lathyrus sp.)~~

Formatted: Indent: First line: 0"

Formatted: Indent: First line: 0"

Formatted: Font: Italic, Complex Script Font: Italic

Formatted: Font: Italic, Complex Script Font: Italic

Formatted: Font: Italic, Complex Script Font: Italic

~~intercropped alone or with barley (*Hordeum vulgare*) significantly; between cherry trees in semi arid Lebanese area (Jourd Aarsal, eastern Lebanese mountains), increased SOC in cherry orchards significantly notably when legumes were mixed with barley (Darwish et al., 2012). Roots of cover crops contributed some Results showed that the sites were supplemented with OM varying between 140 and 250 kg ha⁻¹season⁻¹ of organic matter (OM) against resulting from the decomposition of plant root residues. The above ground plants provided the orchards with 95-665.7 kg ha⁻¹season⁻¹ of OM for the aboveground parts.~~

~~Plant residues provided additional feedstuff for small ruminants; the soils were enriched with OM and fixed nitrogen with more efficient use of surface soil moisture.~~

The effects of crop rotations on SOC are related to the amounts of above and belowground biomass produced and retained in the system. In a study conducted in semi-arid northern Syria, a 12-year rotation gave higher SOC in wheat-medic (12.5 g SOC kg⁻¹ soil) and wheat-vetch (13.8 g SOC kg⁻¹ soil) rotations, as compared to continuous wheat (10.9 g SOC kg⁻¹ soil) or wheat-fallow (Masri and Ryan, 2006). In this rainfed system, the introduction of a forage legume (vetch/medic) with wheat, over a decade, was able to significantly raise the level of SOC. Further, the combination of crop rotations and no-tillage was found to sequester more C than monocultures (Palm et al., 2014). One means of building up biomass is through ~~cover-winter cover~~ crops. Their beneficial impact on C sequestration and water infiltration has been demonstrated. The presence of a cover crop on the soil surface protects the soil against erosion. ~~In the NENA region, their cultivation is restricted to sub humid to humid areas (> 600 mm of rainfall).~~ Still, more research is needed about the best species to be used, the optimum termination strategies of the cover crop as well as the best date and density of planting and best management practices of consequent crops (Plaza-Bonilla et al., 2015). The choice of cover crops in NENA region is crucial as these can compete with the main crop for the limited water resources.

In poor dryland regions ~~Poverty~~, especially in the rainfed agricultural systems, ~~prevents~~ some practices hinder the accumulation of all such as the incorporation of SOC residues. Overall, crop residues serve as fodder or for household cooking or heating, leaving little plant material remains on the soil surface. Even animal dung is used as cooking fuel in many regions. The low SOC content could be improved by increasing the crop residues

Formatted: Font: Italic, Complex Script Font: Italic

Formatted: Indent: First line: 0"

Formatted: Font: (Default) +Headings CS (Times New Roman), 12 pt, Complex Script Font: +Headings CS (Times New Roman), 12 pt, Not Highlight

Formatted: Font: (Default) +Headings CS (Times New Roman), Complex Script Font: +Headings CS (Times New Roman)

Formatted: Indent: First line: 0"

Formatted: Indent: First line: 0"

produced and incorporated. – Such an approach requires the application of fertilizers in order to avoid the depletion of soil nutrients (Plaza-Bonilla et al., 2015). By removing residues, animal dungs and crops, no residues are left in the soil except roots. In the absence of fertilizers, these practices can mine the soil N and over the years the pool of nutrients in the soil can be imbalanced and depleted.

Some authors question the validity of remediation measures to ~~build-up~~promote SOC accumulation in most of the NENA region. –Results from research stations in Egypt and Syria provide ~~proof~~evidence~~evidences~~ to the contrary. In a trial in north-east Cairo, Egypt, the irrigation of a sandy soil with sewage water, for 40 years, changed its texture to loamy sand (Abd el-Naim et al., 1987). This modification of the soil texture leads to a significant improvement of the soil physical properties. Further, within the same long-term trial, the irrigation with sewage water, for 47 years, increased SOC to 2.79%, against 0.26% in the control (Pescod and Arar, 2013). This rather slow accumulation could be related to the sandy soil texture and to the input of the organic matter in labile, soluble forms.

–The addition of more stable composted materials was tested in semi-arid north Syria. The amount of compost, 10 Mg ha⁻¹ every two years, needed to raise the SOC, was too large in these rainfed systems. This amount is larger than the compost available in these conditions. Rather than relying on composts, the authors found that a combination of reduced tillage and a partial retention of crop residues moderately increased SOC (Sommer et al., 2014). The quality of residues seems to affect the SOC on the short-term but on the medium-term it is the quantity that matters (Palm et al., 2014).

3.3.3. Impact of irrigation on agricultural soils

The irrigated land ~~might~~represents a minor fraction of agriculture in NENA region, but irrigated crops are essentially found on prime soils (Figure 4). Frequent wetting of irrigated soils make them more likely to lose C as compared to dry soils. But, this partial loss is compensated by higher biomass production and greater OM inputs from roots, even if residues are removed. Lack of moisture limits soil mineralization (Sharma et al., 2012). Irrigated soils promote intense microbial activity and a rapid decomposition of SOC. In the fertile region of Doukkala, Morocco, known for producing wheat and sugar beet, a

Formatted: Indent: First line: 0"

Formatted: Font: (Default) +Headings CS (Times New Roman), Complex Script Font: +Headings CS (Times New Roman)

Formatted: Font: (Default) +Headings CS (Times New Roman), 12 pt, Complex Script Font: +Headings CS (Times New Roman), 12 pt

decade of irrigated farming decreased SOM-soil organic matter by 0.09% per year (FAO and ITPS, 2015). This loss could have been reduced through the incorporation of crop residues. But, in these mixed farming systems, aboveground residues are consumed by farm animals.

The irrigation of soils in NENA region is expected to affect the SIC. Dryland soils were considered to contain, ~~at least, as much~~ equivalent stock of SIC as SOC (Sharma et al., 2012). ~~But~~ However, this study showed much higher SIC than SOC were found in this study, notably in the subsoils. ~~Despite~~ Despite this large stock, there is a major knowledge gap regarding the effects of land use and management on the dynamics of SIC. This is especially relevant to the irrigation with calcium or sodium-enriched groundwater (Plaza-Bonilla et al., 2015). ~~In~~ In these conditions, the formation of calcium carbonate could be accompanied by some release of carbon dioxide while the development of sodicity can cause irreversible SOC loss.

4. Conclusions

NENA area consisting of 14% of the earth surface contributes only 4.1% of total SOC stocks in topsoil. The soil resources of NENA region are developed under dry conditions with prevailing of rainfed agriculture. The majority of lands in NENA countries are of low productivity. The current mapping of SOC density showed that 69% of soil resources represent a SOC stock below 30 tons ha⁻¹, indicating the soils of NENA region are not enriched with OC. Highest stocks (60 tons ha⁻¹) were found in forests, irrigated crops, mixed orchards and saline flooded vegetation. This means that SOC can be increased in the soils of the NENA region under appropriate and sustainable soil management practices. The moderate density (≈30 tons ha⁻¹) in urban areas indicates land take by urban growth and expansion on prime lands. The stocks of SIC were higher than SOC density, due to the calcareous nature of soils. In subsoil, the SIC stock ranged between 25 and 450 tons ha⁻¹, against 20 to 45 tons ha⁻¹ for SOC. The OC sequestration in the NENA region is a possible task to mitigate climate change and sustain food security despite the hostile climatic conditions and poor land stewardship and governance. Practices of conservation agriculture (no-tillage, intercropping and agro-pastoral system, winter cover crops, proper rotation) could be effective in reducing evaporation, water and wind erosion and promoting aboveground and belowground biomass production. Land cover and land use affected the amounts of SOC retained in the soil ecosystem. A good result was achieved in

Formatted: Indent: First line: 0"

Lebanon through winter cover crop consisting of fruit trees-legume-barley intercropping system. Knowledge gaps exist with respect to the effect of irrigation on SOC and SIC. Constraints facing soil conservation measures and carbon sequestration in the NENA region can be faced with awareness raising and capacity building at the level of stakeholders and decision-makers. Sustainable soil management can contribute to alleviate the pressure on soil resources, improve SOC sequestration and maintain soil resistance to degradation.

The total stocks of SOC (0-0.3m) showed a small contribution to global SOC stock in the topsoil (4.1%), against 14% of the earth surface area. Soils of the NENA region are mostly highly vulnerable to degradation, and food security will depend much on sustainable agricultural measures. The land capability model showed that most NENA countries (17 out of 20), suffer from low productive lands (>80%). To obtain an idea of the status of the soil carbons, the spatial distributions of SOC and SIC stocks for the NENA region were mapped (1:5 Million). This small scale mapping was compared with a larger scale mapping (1:50,000) for Lebanon. A moderate discrepancy (11% to 14%) was found between the two scales. The results of the mapping Mapping of the stocks of SOC and SIC showed that 69% of soil resources present a stock of SOC below the threshold of 30 tons ha^{-1} , considered as limit value for carbon deficient soils. The stocks varied between ~ 10 tons ha^{-1} in shrublands and 60 tons ha^{-1} for evergreen forests. Highest stocks were found in forests, irrigated crops, mixed orchards and saline flooded vegetation. The moderate stock (~ 30 tons ha^{-1}) in urban areas indicates that some urban growth took place on was at the expenses of prime soils. The stocks of SIC were higher than those of SOC due to. In subsoils, the SIC ranged between 25 and 450 tons ha^{-1} , against 20 to 45 tons ha^{-1} for SOC.

Decomposition of SOC is accelerated by climatic conditions, high temperatures, wetting/drying cycles, and by sandy soil textures. OC sequestration in the NENA region is problematic due to specific geomorphological and climatic factors, requiring the protection of the topsoils. Practices of conservation agriculture (no tillage, presence of soil cover...) could be effective in as the presence of residues reduces the evaporation, as well as water and wind erosions. This is especially relevant to soil classes that are susceptible to degradation. Further, the combination of crop rotations and no tillage was found to sequester more C than monocultures. The introduction of legumes, as part of a cereal legume rotation, and the application of nitrogen fertilizers to the cereal caused a

Formatted: Indent: First line: 0"

Formatted: Indent: First line: 0.5"

Formatted: Strikethrough

notable increase of SOC, after 10 years. The effects of crop rotations on SOC are related to the amounts of above and belowground biomass produced and retained in the system.

Some knowledge gaps exist, especially in aspects related to the effect of irrigation on SOC, as well as on SIC at the level of soil profile and soil landscape. Still, major constraints facing soil conservation measures and carbon sequestration are socio-economic in nature, rather than scientific. They are related to the ability of growers to accept new techniques and adopt them. Awareness raising and capacity building at the level of stakeholders and decision makers can contribute to alleviate the pressure on vulnerable soil resources, improve SOC sequestration, and maintain soil resilience to degradation and strengthen food security.

5. Acknowledgements

This paper was supported by the FAO, GSP-ITPS, UNESCWA and CNRS Lebanon within a land degradation assessment and SOC mapping project related to food security.

6. References

Abd Elnaim ~~E. M.~~, ~~M.S.~~ Omran, ~~M. S. T.M.~~ Waly, ~~T. M.~~ and ~~El Nashar, B.M.B. El Nashar~~ ~~B. M. B.~~ ~~El Nashar~~ (1987) Effects of prolonged sewage irrigation on some physical properties of sandy soil. *Biological Wastes*, 22, 269-274, 1987.

Al Chami, Z., ~~S.~~ Bou Zein Eldeen, ~~S. L.~~ Al Bitar ~~L.~~ and ~~T.~~ Atallah ~~T.~~ (2016) Decomposition of olive-mill waste compost, goat manure and Medicago sativa in Lebanese soils as measured using the litterbag technique. *Soil Research*, 54, 191-199, 2016.

Atallah, T., ~~C.~~ Jamous, ~~C. P.~~ Debs, ~~P.~~ and ~~Darwish, T.~~ ~~Darwish~~ (2012) Biosolid recycling to enhance carbon sequestration in mountainous Lebanese conditions. *Lebanese Science Journal*, 13, (2), 69-79, 2012.

Batjes, N. H., and Sombroek, W. G. (1997) Possibilities for carbon sequestration in tropical and subtropical soils. *Global Change Biology*, (3), 161-173, 1997.

Bosco S., ~~C.~~ Di Bene, ~~C. E.~~ Bonari ~~E.~~ (2012) The effect of crop management on soil organic matter in the carbon footprint of agricultural products. 8th Int. Conf. on Life

Formatted: Indent: First line: 0"

Formatted: Indent: Before: 0", First line: 0"

688 Cycle assessment in the agri-food sector. Saint-Malo, France, Oct-1-4 October 2012,
689 Saint-Malo, France, 2012.

690 Bosco, S., ~~C.~~ Di Bene, ~~C.~~ M. Galli, ~~M.~~ D. Remorini, ~~D.~~ R. Massai, ~~R.~~ and ~~E.~~ Bonari,
691 ~~E.~~ (2013) Soil organic matter accounting in the carbon footprint analysis of the wine
692 chain. ~~International Int. Journal Life Cycle Assess., essment~~ 18, ~~:973-989.~~ DOI
693 10.1007/s11367-013-0567-3, 2013.

694 Boukhoudoud, N., Gros, R., Darwish, T., and Farnet Da Silva, A.M.: A.M. 2016.
695 Agriculture practice and coastal constraint effects on microbial functional properties of
696 soil in Mediterranean olive orchards. European Journal of Soil Science., 67. DOI:
697 10.1111/ejss.1234. July 2016, 67: 470-477. 2016.

698 Boukhoudoud, N., Farnet Da Silva, A. M., Darwish, T., and Gros, R.: R. 2017. Olive mill
699 waste and glyphosate-based herbicide addition to olive grove soils: effects on microbial
700 activities and their responses to drying/rewetting cycles. Soil Use and Management.,
701 DOI: 10.1111/sum.12367, 2017.

702 Cerdà A., Gonzalez Penaloza F., Santin C., and ~~S.H.~~ Doerr S.H. (2012) Land
703 abandonment, fire recurrence and soil carbon content in the Macizo del Caroig, Eastern
704 Spain. Geophysical Research Res. Abstracts 14, EGU2012-14331, 2012.

705 Darwish, T., Faour, Gh., and ~~M.~~ Khawlie, M.: (2004) Assessing Soil Degradation by
706 Landuse-Cover Change in Coastal Lebanon. Lebanese Science Journal 5, ~~: 45-59.~~ 2004.

707

708 Darwish, T., Khawlie, M., Jomaa, I., Abou Daher, M., Awad, M., Masri, T., Shaban, A.,
709 Faour, Gh., Bou Kheir, R., Abdallah, C. and Haddad, T. (Eds 2006). Soil Map of Lebanon
710 1/50000. Publications CNRS-Lebanon, Monograph Series 4, 1 pp-367, 2006.

711 Darwish, T., Abou Daher, M., Jomaa, I., and Atallah, T.: Soil organic carbon stock
712 estimation in Lebanese territories. 10th International Meeting on Soils with Mediterranean
713 Type of Climate. CNRS-Lebanon, Book of Extended Abstracts, 113-118, 2009.

714 Darwish, T., Atallah, T., Francis, R., Saab, C., Jomaa, I., Shaaban, A., Sakka, H., and ~~P.~~
715 Zdruli P.: (2011) Observations on soil and groundwater contamination with nitrate, a case
716 study from Lebanon-East. Mediterranean Agr. Water Management, DOI
717 10.1016/j.agwat.2011.07.016, 2011.

Formatted: French (France)

Formatted: Indent: Before: 0", First line: 0",
Space After: 0 pt

718 Darwish, T.M.; Jomaa, I.; Atallah, T.; Hajj, S.; Shaban, A.; Zougheib, R.; and F. Sibai
 719 Ouayda, F.: (2012) An agropastoral system as a practice to enhance organic matter in
 720 Lebanese inland mountainous soils. *Lebanese Lib. Science Journal*, vol. 13 (1), 1-14,
 721 2012.

722 Darwish, T., and A. Fadel, A.: (2017) Mapping of Soil Organic Carbon Stock in the Arab
 723 Countries to Mitigate Land Degradation. *Arab J. Geosci.* (2017) 10, 474.
 724 <https://doi.org/10.1007/s12517-017-3267-7>, 2017.

725 Elhadi, M.Y.: 2005. Postharvest technology of food crops in the Near East and North
 726 Africa (NENA) region, i. In: *Ramdane Dris Ph.D. (ed.), 'Crops: Growth, Quality and
 727 Biotechnology'*, pp. 643-664, WFL Publisher, Meri-Rastilan tie 3 C, 00980 Helsinki,
 728 Finland, 2005.

729 FAO: 2007. The digital soil map of the world. Food and Agriculture Organization of the
 730 United Nations, Version 3.6, completed January 2003 and updated 2007
 731 (<http://www.fao.org/geonetwork/srv/en/metadata.show?id=14116>), accessed 20
 732 December 2017.

733 FAO: 2012. Country Study on Status of Land Tenure, Planning and Management in
 734 Oriental Near East Countries Case of Lebanon, FAO, RNE, SNO, Cairo, Egypt, 161 p.,
 735 2012.

736 FAO: 2015. Regional Overview of Food Insecurity - Near East and North Africa:
 737 Strengthening Regional Collaboration to Build Resilience for Food Security and Nutrition,
 738 Cairo, Egypt, FAO, 2015.

739 FAO: 2017. Voluntary Guidelines for Sustainable Soil Management, Food and
 740 Agriculture Organization of the United Nations Rome, Italy, <http://www.fao.org/3/a-bl813e.pdf>, accessed November 17, 2017.

742 FAO and ITPS: 2015. Status of the World's Soil Resources (SWSR) – Main Report,
 743 Food and Agriculture Organization of the United Nations and Intergovernmental
 744 Technical Panel on Soils, Rome, Italy, 2015, www.fao.org/3/a-i5199e.pdf accessed 13
 745 December, 2017.

746 Fernandez-Ugalde, O.; Virto, I.; Barre, P.; Apesteguia, M.; Enrique, A.; Imaz, M. J. and P.
 747 Bescansa, P.: (2014). Mechanisms of macroaggregate stabilisation by carbonates:

Formatted: French (France)

748 implications for organic matter protection in semi-arid calcareous soils. *Soil Resear*
749 *ch* 52: 180-192, 2014.

750 Finke, P., Hartwich, R., Dudal, R., Ibanez, J., Jamagne, M., King, D., Montanarella, L., and
751 N. Yassoglou N.: (2000) Georeferenced Soil Database for Europe. Manual of Procedures,
752 Version 1.1. European Soil Bureau, 2000.

753 Guo, L.J., Lin, S., Liu, T., Q., Cao, C., G., Li, C., F.: (2016) Effects of conservation tillage
754 on topsoil microbial metabolic characteristics and organic carbon within aggregates under
755 a rice (*Oryza sativa* L.)-wheat (*Triticum aestivum* L.) cropping system in Central China.
756 *PLoS One* 11:e0146145, 2016.

757 Hengl, T., de Jesus, J. M., MacMillan, R. A., Batjes, N. H., Heuvelink, G. B. M., Ribeiro,
758 E., Rosa, A. S., Kempen, B., Leenaars, J. G. B., Walsh, M. G., Gonzalez, M. R.: (2014)
759 *SoilGrids1km-Global Soil Information Based on Automated Mapping. PLoS ONE* 9,(8):
760 e105992, DOI:doi:10.1371/journal.pone.0105992, 2014.

761 Hengl, T., Mendes de Jesus, J., Heuvelink, G. B. M., Ruiperez Gonzalez, M., Kilibarda,
762 M., Blagotić, A., et al. : *SoilGrids250m: Global gridded soil information based on machine*
763 *learning. PLoS ONE* 12(2): e0169748. <https://doi.org/10.1371/journal.pone.0169748>,
764 2017, accessed 20 February 2017.

765 Lal, R.: (2003) Soil erosion and the global carbon budget. *Environment International*,
766 29,29 (4): 437-450, 2003.

767 Masri, Z. and Ryan, J.: Ryan (2006) Soil organic matter and related physical properties in
768 a Mediterranean wheat-based rotation trial. *Soil Tillage Resear* 87: 146-154, 2006.

769 Mrabet, R., Moussadek, R., Fadlaoui, A., and van Ranst, E.: van Ranst (2012)
770 Conservation agriculture in dry areas of Morocco. *Field Crops Resear* 132: 84-94,
771 2012.

772 Pescod, M. B., and Arar, A.: Arar (2013) Treatment and use of sewage effluent for
773 irrigation. Butterworths. <https://books.google.com.lb/books>, ISBN: 1483162257, Isbn:
774 1483162257, Pages 211-212, 2013.

Formatted: English (U.S.)

Formatted: French (France)

Formatted: French (France)

Formatted: English (U.S.)

Formatted: text, Complex Script Font: Times
New Roman

Formatted: text, Complex Script Font: Times
New Roman

775 Plaza-Bonilla, D., Arrúe, J. L., Cantero-Martínez, C., Fanlo, R., Iglesias, A., and Álvaro-
 776 Fuentes, J.: ~~-(2015)~~ Carbon management in dryland agricultural systems. A review.
 777 Agronomy for Sustainable Development 35, ~~-(4):~~ 1319-1334, 2015.

778 Sharma, P., Abrol, V., Abrol, S., and Kumar, R.: ~~Kumar (2012)~~ Climate change and
 779 carbon sequestration in dryland soils. ~~i-~~In Tech open access book, chapter 6. 26 pages.
 780 <http://dx.doi.org/10.5772/52103>, 2012, accessed last time 31 July 2018.

781 Sommer, R., Piggitt, C., Feindt, D., Ansar, M., van Delden, L., Shimonaka, K., Abdalla,
 782 J., Douba, O., Estefan, G., Haddad, A., Haj-Abdo, R., Hajdibo, A., Hayek, P., Khalil, Y.,
 783 Khoder, A., and Ryan, J.: ~~Ryan (2014)~~ Effects of zero tillage and residue retention on soil
 784 quality in the Mediterranean region of northern Syria. Open Journal of Soil Science 4 ~~-(3),~~
 785 Article ID: 44383. 17 pages. DOI:10.4236/ojss.2014.43015, 2014.

786 [Stockmann, U., Padarian, J., McBratney, A., Minasny, B., deBrogniez, D., Montanarella,](#)
 787 [L., Young Hong, S., Rawlins, B. G., Damien, J.: \(2015\) Field-Global soil organic carbon](#)
 788 [assessment. Global Food Security, 4\(6\), 9-16, 2015.](#)

789 USDA: ~~(1999)~~ Land capability classification, ~~-~~NRCS-USDA, 1999.

790 von Grebmer, K., Bernstein, J., Hossain, N., Brown, T., Prasai, N., Yohannes, Y.,
 791 Patterson, F., Sonntag, A., Zimmermann, S. M., Towey, O., and Foley, C.: ~~Foley (2017)~~
 792 Global Hunger Index: The Inequalities of Hunger, Washington, DC, International Food
 793 Policy Research Institute, Bonn, Welthungerhilfe, and Dublin, Concern Worldwide,
 794 2017.

795 Yu, L., Dang, Z.-Q., Tian, F.-P., Wang, D., and Wu, G.-L.: ~~Wu (2017)~~ Soil organic
 796 carbon and inorganic carbon accumulation along a 30-year grassland restoration
 797 chronosequence in semi-arid regions (China). Land Degrad. Develop., 28, 189–198,
 798 2017.

799 [Webography](#)

800 [Website 1. http://maps.elie.ucl.ac.be/CCI/viewer/index.php](http://maps.elie.ucl.ac.be/CCI/viewer/index.php)

801 [Website 2. http://www.fao.org/3/a-bl105e.pdf](http://www.fao.org/3/a-bl105e.pdf)

802 [Website 3. https://drive.google.com/drive/folders/1454FhX_p2_GxjZT_fsi0_Rncl2QpSN](https://drive.google.com/drive/folders/1454FhX_p2_GxjZT_fsi0_Rncl2QpSN)

Formatted: French (France)

Formatted: Indent: Before: 0"

Formatted: Indent: Before: 0", Space After: 10 pt, Adjust space between Latin and Asian text, Adjust space between Asian text and numbers