

Authors response: Comments on the manuscript “Dynamic modelling of weathering rates – Is there any benefit over steady-state modelling?”

Authors: Veronika Kronnäs, Cecilia Akselsson and Salim Belyazid

veronika.kronnas@nateko.lu.se

5 MS No.: soil-2018-25

2018-11-27

Contents

	Authors response: Comments on the manuscript “Dynamic modelling of weathering rates – Is there any benefit over steady-state modelling?”	1
10	Comments from reviewer 1	2
	GENERAL COMMENTS.....	2
	SPECIFIC COMMENTS	4
	Comments from reviewer 2	10
	GENERAL COMMENTS.....	10
15	SPECIFIC COMMENTS	10
	Manuscript: Dynamic modelling of weathering rates – the benefit over steady-state modelling	1
	1 Introduction.....	1
	2 Methods	3
	2.1 PROFILE	3
20	2.2 ForSAFE	3
	2.3 Site descriptions	4
	2.4 Scenarios and time series of driver parameters	4
	3 Results	5
	3.1 Weathering rates from PROFILE and ForSAFE	5
25	3.2 Seasonal, yearly and decadal variation in weathering rates from ForSAFE	6
	3.3 Effect of forestry on weathering.....	6
	3.4 Effect of acidification on weathering.....	7
	3.5 Effect of climate change on weathering.....	7

3.6 Overall effect of forestry, acidification and climate change	7
4 Discussion.....	7
4.1 Implications of model differences	7
4.2 Weathering dynamics in a changing environment.....	8
5 4.3 Model limitations and development	9
5 Conclusions.....	10
References	11
Tables	15
Figures.....	17
10	

We would like to thank both reviewers so much for your comments. They were very helpful and we think that they have helped us improve the manuscript and make it more clear and understandable. Below we give our answers to the questions the reviewers had.

15

Comments from reviewer 1

GENERAL COMMENTS

Comment: “A neat and generally well written and structured paper, the subject of which (weathering) falls within the scope of SOIL and is of broad international interest. However, several related and somewhat similar applications of the models have already been published, as indicated by the strong Sweden related references.”

Answer: Thank you! It is true that other outputs from ForSAFE (such as soil water chemistry) have been analysed and evaluated in a number of papers, but the actual weathering rates themselves have not been compared with any other methods. This study shows that the weathering calculations in ForSAFE give weathering rates of the same size as the much more tested and evaluated model PROFILE.

Comment: “And - as stated by the authors - as both models are based on the same weathering equations, it is hardly surprising that the results (long-term) from both models are similar and suggests little benefit is gained in using the more complex model when looking at long-term climate change and forest management impacts.”

Answer: Both models are based on the same weathering equations, but in ForSAFE the equations are dynamic, while they are not in PROFILE. There are processes in ForSAFE, which are important for the weathering and which are not modelled

at all in *PROFILE* – soil hydrology for example. Also, the timing of different processes relative to each other can be important for the weathering and affect the weathering rates; also the long term average rates. This cannot be captured in *PROFILE*. For example the timing of the relatively high uptake to vegetation in the beginning of the vegetation period vs the timing of the high weathering during warmer months, and high leaching during wetter months – these processes affect soil water chemistry which in its turn affect weathering in the next time step. Also, *PROFILE* assumes steady state and cannot take into consideration how the system got there, if the changes were slow and gradual or abrupt. Reality is never in steady state, and climate in the future is less in steady state now than in a long time, which makes a dynamic model more important than in earlier calculations of what loads of acidity the ecosystem in climatic steady state could tolerate. We tried to describe this in the discussion, P7L26-P8L8. See also our answer to the second reviewer’s second question.

10 **Comment:** “Furthermore, the dynamic ForSAFE model, by definition is bound to provide more detailed and seasonal results and information, making the answers to the question posed by the title and to the hypotheses some- what self-evident.”

Answer: We have changed the title to “Dynamic modelling of weathering rates – the benefit over steady-state modelling” (P1L2), as suggested by the second reviewer. We have also added a bit to the second objective (P2L31-32): “...scenarios, representing important ecological issues: acidification, climate change and nutrient removal through land use.”, since a large part of the paper describes the weathering response to environmental drivers (i.e. describes in detail some benefits that can be gained by dynamic modelling). By doing those changes, the title and the second objective fits better to what we actually do in the paper –describing the benefits and the dynamics in weathering. For example, we show that even though ForSAFE weathering rates vary a lot, averages over the same period represented in the *PROFILE* modelling, fall so close to *PROFILE* values, which gives credit to the ForSAFE weathering. We didn’t know this before we performed the study.

20 **Comment:** “I think the paper could be substantially improved if there were more focus on and calibration of the ForSAFE model with empirical field data; for example, with measured soil temperature, soil moisture, forest growth (base cation uptake) and leaching data.”

Answer: ForSAFE is not calibrated with empirical field data except base saturation and soil carbon and nitrogen. Other data can be used for comparisons with model results. Unfortunately we don’t have measurements of soil moisture or soil temperature for these sites. Since soil moisture and temperature are very important parameters, a very good next step would be to compare ForSAFE modelled soil moisture and soil temperature to measured data, but that is outside the scope of this paper, as that would have to be on other sites. We include this suggestion in the discussion (P9L28-30) as a suggested future study. There are measured forest growth and soil water concentrations for these sites, but there are already papers comparing ForSAFE modelled values with these kinds of measurements (Yu et al in the reference list and others).

30

SPECIFIC COMMENTS

Comment: “p. 1, l. 20 (and p. 4, l. 31). Annual precipitation may remain similar but what about the seasonal distribution? How will temperature change affect snowfall and snowmelt, surely a very important feature of climate change in such latitudes, the water cycle and weathering?”

5 **Answer:** *In this part of Sweden, there is already very little snow. Otherwise it would probably be an important feature. The seasonal distribution of precipitation does change in this scenario, with less precipitation in summer in the second half of the century. The last decades the yearly precipitation increases somewhat too, about 8%. We corrected the text in the abstract with regards to seasonal distribution of precipitation and did the same in the chapter about scenarios (2.4) and in the discussion (P1L21-23, P3L5-8, P8L20-23).*

10 **Comment:** “p. 3, l. 19 (and p. 8, l.11). Why/how are base cations, Al³⁺ and organic acids inhibitors of weathering? Do you mean that if the concentrations of weathering products in the soil solution increase, the weathering reaction slows down as equilibrium concentrations are reached? But then aren't the weathering products being continually taken away through uptake, leaching or adsorption by the soil allowing weathering to proceed?”

Answer: *Yes, ions in the soil solution slow the weathering down, they don't stop it altogether, since the weathering products are leached, adsorbed or taken up by trees. Nonetheless, in acidified conditions the concentration of Al in the soil increases substantially compared to non-acidified conditions, and slows weathering of silicates down. Also, in the C-horizon, where water flows are slower and uptake to trees are lower, concentrations of weathering products increase and should slow weathering down to a minimum – that is probably why the C-horizon consists of less weathered parent material even though there are so much weatherable material. We corrected the text; the inhibitors base cations and aluminium are products of the weathering (P3L19).*

Comment: “p. 4, l.16. There is little description of the soil type at the two sites. From the horizon abbreviations listed in Table 2, it would appear the soil are not Podzols?”

25 **Answer:** *The two soils are assessed in the SWETHRO database as transition types (developing towards podzols). O is the thin uppermost layer consisting of mostly organic material, A is greyish, but not deemed a true E zone, AB is a light brown, B is a reddish brown and C is the parent material, which is till. We added information about the soil type in chapter 2.3 (P4L17-18).*

Comment: “p. 5, l.16. Furthermore, thickness of the mineral soil horizons at Hissmossa is 55 cm (and not 50 cm; Table 2) and how/why is the organic layer included where surely it is a question of organic matter decomposition rather than mineral weathering? It is not stated how stoniness was derived and if the hydraulic parameter values (Table 3) have been corrected for stone content. Given the concluded importance of soil moisture, the fixed value used in PROFILE and same value for all layers (0.2) seems rather crude. The field capacity values in Table 3 also seem somewhat low – is this because of stone

content correction? How have any time related changes in organic layer thickness (and therefore soil moisture content) been taken into account?”

Answer: *The modelled soil horizon at Hissmossa is 55 cm+organic layer, but the root zone is still 50 cm+organic layer. We included the C-horizon in the modelling, even though it lies below the root zone, but we did not include it in calculations representing the weathering in the root zone. We clarified this in the text to figure 1. The organic layers do contain some mineral soil and the modelled weathering in those layers is from the minerals, not from the organic matter. They are included in the modelling since they are important for the modelled ecosystem and thus for the soil water chemistry for the rest of the layers – for example much of the nutrient uptake takes place here, especially of nitrogen. Stoniness is estimated at the soil sampling. We added this in chapter 2.3 (P4L22-26). PROFILE, being a steady state model, can only handle fixed (in time) values. The same soil moisture value for all horizons may be crude, but without measurements over long time periods in all soil layers, there has not been much of an alternative –and in Västra Torup’s case it is in accordance with the modelling (see new figure 2). The field capacity values are calculated according to Balland (2008). Dynamic changes in layer thicknesses are not modelled by any of the models.*

Comment: “P. 4, l. 31: by “rainfall” you mean annual precipitation?”

Answer: *Yes, thanks. I have changed the text (P5L5).*

Comment: “p. 5, l. 6-11. The description of the scenarios is unclear, at least to me. For example, does the base scenario mean there are two thinnings and a clear cut every 70 years (and starting from the year of planting – Table 1), deposition loads constant from “today’s” (which year?) levels into the future plus climate change temperature (but no change in precipitation)? What is the whole tree harvesting treatment: stems + branches or stems + branches + stumps? Is it carried out every 70 years during the 1900-2100 period? It would be useful to number or letter the scenarios and refer to them in the text and, tables and figures.”

Answer: *We have given the scenarios three letter abbreviations and use those throughout the text and figures, as suggested. We have clarified the scenario descriptions, both the description of the base scenario (BSC) and how the others differ from the BSC scenario (P4L31-P5L20). Yes, the BSC has forestry with a clear cut approximately every 70 years (Västra Torup: 1940, 2010, 2080. Hissmossa: 1972, 2040, 2100) and thinnings at about 25 and 45 years of plant age (the thinnings are more “on time” in the future scenario than in actual history of the sites). Deposition of SO_x peaks in 1970 and decrease sharply afterwards and is in the future scenarios kept constant at 13% of the peak deposition after 2020. Deposition of N peaks in 1985 and decrease more slowly. It is kept constant at 50% of the peak deposition after 2020. Climate: Both temperature and precipitation changes in the BSC scenario, (as in the SRESA2 scenario) – the temperature increase almost exponentially from 1900-1910 to 2090-2100 with on average about 5.9°C in winter and 3.7° in summer. Yearly precipitation changes less and without trend, but summer precipitation is lower in 2050-2100 than before and winter and spring precipitation is higher. Whole tree harvest is, in the WTH scenario of this study, stems and 60% of branches, treetops and*

needles, but stumps are not removed. Whole tree harvest in the WTH scenario is carried out in thinning and clear cutting from the clear cutting in 2010 (Västra Torup) or 2040 (Hissmossa) and forward.

Comment: “Which scenario is used for Figure 1?”

5 **Answer:** *Base scenario (BSC). We gave the scenarios abbreviations and clarified in the running text and in the figure texts which scenario was used.*

Comment: “Doesn’t the 70 year rotation period cover a different set of years between the two sites (20112080 vs. 2041-2100), when the climate change has changed the climate. Doesn’t this explain the differences in weathering between the two sites rather than differences in soil texture (p. 5, l. 21), which anyway would also affect the soil hydraulic properties besides surface area?”

10 **Answer:** *Yes, the rotation periods are different for the two sites. As can be seen in figure 3 (in the previous version of the manuscript figure number 2, since we added a new figure number 2 showing soil moisture), this does not explain the difference in weathering rates between the sites, as the increase in weathering because of climate change in 20 to 30 years is a lot smaller than the difference in weathering rates between the sites. Also, if the increase in temperature in the time between the clear cuts of the two sites would have been the reason for the difference in weathering between the sites, then*
15 *Hissmossa would have had the higher weathering rates and not the lower. Also, the difference between the sites exist also in the scenarios with no climate change (NCC and BGR). See also P6L2-5. According to our calculations of soil hydraulic properties following Balland (2008), field capacity is slightly lower in Hissmossa than in Västra Torup, which means that soil moisture also is slightly lower, which should affect the weathering rates as calculated by ForSAFE, but not those calculated by PROFILE, since we used the same soil moisture value for both sites in PROFILE. And the differences in*
20 *weathering rates between the sites are as large in the PROFILE calculations as in the ForSAFE ones (see figure 1).*

Comment: “Why is only Mg weathering presented in Figure 2 and in Figure 4 to represent silicate mineral weathering? Wouldn’t the sum of base cations be a more appropriate measure of overall silicate weathering?”

25 **Answer:** *We want to show the effect of the acidification on silicate mineral weathering rather than the total size of it, in figure 5 (previously figure 4). One of the base cations, Ca, comes in about equal amounts from weathering of apatite and of silicates. The weathering of Ca is also relatively large. Thus, to examine the dynamics of release of base cations from only silicate weathering in our models, we chose to show one of the base cations that is only from silicate weathering, not apatite, Mg. We clarify this somewhat in the figure caption for figure 5 (previously figure 4). Figure 3 (previously figure 2) could as well show the sum of base cations, since the focus of the figure isn’t difference between silicates and apatite, so we changed it to sum of base cations instead.*

30 **Comment:** “As weathering largely takes place by acid (proton) attack, why is silicate weathering decreased by acidified conditions (p. 6, l. 17) and why would apatite weathering be increased?”

Answer: *Apatite weathering is increased by more acid conditions because there are more protons in solution, and not at the same time high concentrations of something that inhibits weathering. Silicate weathering was not increased by the more acid conditions on those two sites, according to ForSAFE. Those two sites had high concentration of Al in soil solution during the acidified conditions, as Al is more soluble in acid conditions, and Al inhibits weathering of silicates in ForSAFE, but not of apatite since there is no Al in apatite. We added a little text in the discussion (P9L11-12) to clarify this.*

Comment: “Doesn’t the base scenario include harvesting effects besides climate change effects (p. 6, l. 29)? See also p. 8, l. 21-24.”

Answer: *Yes, but so does the NCC scenario. Forestry and acidification are equal in the NCC and the BSC scenario. Only climate differs. We clarified the scenario descriptions a bit (chapter 2.4).*

10 **Comment:** “p. 7. l. 19: by “more detailed forestry plans” do you mean timing of thinning and timing and intensity of harvesting?”

Answer: *Yes (P8L13-14). And with additional soil data, at least on soil texture, differences within a stand or a couple of nearby stands could also be modelled, so that more sensitive areas within the stand could get a less intensive forestry.*

15 **Comment:** “p. 8, l. 13- Isn’t a matter of litter decomposition and not weathering? And I think you need to give a reference that supports the statement about harvesting intensity effects of soil solution base cation concentrations. Concentrations and leaching loads may increase with whole-tree harvesting as a result of increased drainage (percolation) and disturbance of the site.”

20 **Answer:** *We didn’t explain this clearly enough, thank you for the question. With standard clear cutting where branches and treetops and litter are left on the site, litter decomposing after harvest lead to increases in base cation concentration, especially of K. This can be seen in data from SWETHRO sites (krondroppsstatet.ivl.se) and in Piirainen et al. (2004). After whole tree harvesting, where branches and treetops, with most of the needles, are removed, there will be less decomposable litter on the site, which should mean that concentrations in soil water do not increase as much after WTH as after stem only harvest (as discussed in Ågren et al., 2010). The concentrations in the soil water in its turn affect the weathering rates. See P8L31-P9L2.*

25 **Comment:** “p. 8. L. 25-. The contribution of your C-horizon is small (Table 2). And if the material is less weathered, then there would be more weatherable minerals and therefore potential or weathering? Is the explanation for taking silica concentrations into account the same as mentioned above, i.e. equilibrium concentrations reached?”

30 **Answer:** *The contribution of the C-horizon is small because we only model a small portion of it. But the rates of weathering per volume in the C-horizon are equivalent to the rates in the horizons above, in the modelling. The material in the C-horizon is less weathered since the weathering in the C-horizon is smaller than in the B-horizon, even though there is more weatherable material. The environment in the C-horizon inhibits weathering (or the C horizon would already be weathered*

and not be a C horizon), probably because of slow water movements, so that some reactants are depleted and some products of the weathering build up and inhibit further weathering. And Si is one of the products that are building up, one that we do not model with ForSAFE and PROFILE yet. We added a little text in the discussion to clarify this, P9L33-P10L4, P10L16-21.

5 **Comment:** “p. 9. l. 6. A paper by Starr & Lindroos (Geoderma (2006) 133: 269–280) shows this.”

Answer: *Thank you! We included it in the text, P10L17.*

Comment: “Conclusions: I appreciate the recognition of the importance of soil moisture to weathering (and decomposition) and for the reasons stated (time step). However, wouldn’t a model with a daily time step be more suited to forest stand nutrient/biogeochemical cycling studies rather than modelling long-term climate change effects.”

10 **Answer:** *A daily time step is more suited for modelling of hydrology, which in its turn affects all other processes both on short term and possibly also their long term averages. The effects on long term averages might be large, if average soil moisture is affected significantly or if the dynamic shows that the timing of different processes produce for example lack of soil moisture at high temperatures or lack of nutrients when vegetation needs them. In any case, studying if the daily time step that has been implemented in another version of ForSAFE has an impact seems like a reasonable next step. We clarified*
15 *the text on model development (4.3) somewhat (P9L24).*

Comment: “Units: Wouldn’t it be more correct to present weathering in units of moles charge rather than equivalents? And why sometimes meq m⁻² yr⁻¹ and sometimes meq m⁻³ y⁻¹ (Fig. 2, 3, 4, 5). Is it somehow because the latter refer to a specific layer (of differing thicknesses) rather than to the fixed organic layer + 50 cm layer or simply a typo?”

Answer: *To our understanding, moles of charge and equivalents are equivalent in this case. We used weathering per area in*
20 *figure 1 to make it easy to sum up the weathering from the different layers, even though they have different thicknesses, and to show how much each layer contributes to the total weathering. But since the layers have different thicknesses, the total weathering per forest floor area for a layer is not as easily comparable to other soils in other studies as weathering per soil volume, and therefore we converted the weathering rates into meq m⁻³ y⁻¹ for the text and the rest of the figures. We clarified the use of different unit in figure 1 in the caption to figure 1.*

25 **Comment:** “Table 1. Coordinates are decimal degree latitude (N) and longitude (E).”

Answer: *Yes, thank you. We added unit in the table.*

Comment: “Figure 1. Title: Using which scenario and for which period of time?”

Answer: *BSC scenario and the “future forest rotation”: 2011-2080 in Västra Torup and 2041-2100 in Hissmossa. We added this in the figure caption.*

Comment: “Figures 2 & 6. Is it necessary to include the ForSAFE monthly values? See Fig.3.

Answer: *Since one of the objectives of the paper is to describe the dynamics of the weathering in ForSAFE it seems necessary to show the dynamics of the weathering and not only averages over time periods of different length at least in figure 2 (now figure 3) where ForSAFE weathering (=the monthly weathering values) is compared to PROFILE weathering.*

5 *We removed the monthly values from figure 6 (now figure 7).*

Comment: “Fig. 3. Use a circle around the years of clear cutting on the whole tree harvesting scenario and base scenario lines rather than the vertical line that intersects all scenarios. Are the years 1941, 2010 and 2080 for Västra Torup and 1973 and 2043 for Hissmossa?”

10 **Answer:** *A circle would still seem to indicate all scenarios, since the average weathering is the same before the clear cut for all three scenarios. We think that the vertical lines are more clearly visible and divides the time into the different rotation periods, but we have clarified the figure caption somewhat. The years of clear cut are 1940, 2010 and 2080 in Västra Torup and 1972, 2040 and 2101 in Hissmossa.*

Comment: “Fig. 4. Is each dot is a year of the 70 year rotation period? Why only layer 4 (B-horizon)?”

15 **Answer:** *Each dot represents a month. L4 is the thickest layer of the root zone in Hissmossa and one of the two thickest layers of the root zone in Västra Torup and it lies at the depth of the soil water chemistry measurements. Therefore it is the best layer for comparing modelled and measured soil water chemistry and thus the layer we look at most. Figure 4 (now figure 5) is intended to show the modelled different reaction to acidification between silicate weathering and apatite weathering and for this any layer could be used, but we chose to use L4.*

20 **Comment:** “Fig. 5. The base scenario includes the climate change temperature scenario and does the “constant climate change scenario” refer to the last one in the list on p. 5?”

Answer: *We have now given the scenarios names and refer to them in the text and in the figures with these. The base scenario (BSC) include climate change, yes, and the other scenario used in figure 5 (now figure 6), the NCC, is the same as BSC with regards to forestry and acidifying deposition – but it has no climate change. The last scenario of the list (the BGR) has no forestry, no acidification and no climate change and is used in figure 6 (now figure 7).*

25

Comments from reviewer 2

GENERAL COMMENTS

Comment: “The study presents soil weathering rates for two sites in Sweden. It is concise and well written, and the subject is within the scope of SOIL.”

5 **Answer:** *Thank you!*

Comment: “However, I would urge the authors to rethink the focus of the manuscript; the comparison of output from a weathering sub-model within a steady-state model to output from the same weathering sub-model in a dynamic model is somewhat trivial, and the results are unsurprising.”

10 **Answer:** *ForSAFE consists of many more parts than just the weathering part and they interact with one another. Soil moisture, for example, are not modelled at all in PROFILE and is a very important parameter for the weathering. The weathering sub model is also not exactly the same, since it is dynamic in ForSAFE. For us it was not obvious that the results would be so similar to each other. The models are different enough that one cannot just be switched to the other without studying if the newer one gives reasonable results, which is what we have done by comparing with PROFILE weathering. See also our answer to the first reviewer’s general questions. We included text in the discussion clarifying how*
15 *much the models differ (P7L27-P8L7). We also expanded the second objective to refocus the paper somewhat (P2L32-33).*

Comment: “Similarly, it is also unsurprising that a dynamic model provides more temporal data compared to a steady-state model. The current aims of the manuscript are trivial. There is merit in presenting the dynamics of weathering; the Title and presentation could be refocused to ‘Dynamic modelling of weathering rates – the benefit over steady-state modelling’.”

20 **Answer:** *We agree with the change of title. We think there is merit also in comparing the results of a new (with regards to weathering calculations) model with a previously used model, to examine if the new model would make very different critical load or acidification sensitivity assessment than the old assessment – and if the new assessment would have been very different – how this can be explained and if it is to be believed.*

Comment: “Secondly, the discussion somewhat repeats the results, there are few references to literature, and overall it
25 feels more like a report.”

Answer: *We have restructured the discussion somewhat, expanded it and put in more references (P7L25-P10L21).*

SPECIFIC COMMENTS

Comment: “P1L10. Do not mention SWETHRO in the abstract.”

Answer: *We have removed it. See P1L10.*

Comment: “P1L20. The results / discussion of scenario’s does not fit with the objectives or title of the manuscript.”

Answer: *We changed the title and expanded the description of objective number two somewhat (P2L31-32).*

Comment: “P1L23. This is not a result of this study.”

5 **Answer:** *What is not a result of the study? ForSAFE has not been used for weathering calculations before this study, but we show that weathering rates calculated by ForSAFE are similar to weathering rates from the previously used model PROFILE and thus that ForSAFE can be used for weathering calculations too.*

Comment: “P2L23. Given the importance of weathering (Title / objectives), it is surprising that weathering is given limited attention in the introduction.”

10 **Answer:** *We have focused on why knowing the rate of weathering is important, rather than the weathering chemistry. The process understanding of chemical weathering, which is used in the models, has been described thoroughly in previous papers, some of which we refer to.*

Comment: “P2L27. I recommend that you remove the first objective and expand (refocus) the second.”

15 **Answer:** *We don’t agree on removing the first objective. Weathering from the ForSAFE model has never before been investigated and compared to other, well used, ways of calculating weathering. The fact that the process is in the model and the model gives reasonable other output does not necessarily mean that the weathering estimates are robust and useful. This study shows that the weathering estimates from ForSAFE are of the same size as estimates from the previously often used model PROFILE, and this we did not know before.*

20 *We expanded the description of second objective to explain the scenarios: “...scenarios, representing important ecological issues: acidification, climate change and nutrient removal through land use.” (P2L31-32) The scenarios are included to illustrate what kinds of questions dynamic modelling can help answer, using scenarios with relevant questions such as how weathering responds to climate change, changes in forestry practices and changes in acidifying deposition.*

Comment: “P2L30. What is the objective of the scenarios?”

Answer: *See answer to the comment above.*

Comment: “P3L2. What were the models applied to two forests?”

25 **Answer:** *The steady state model PROFILE and the dynamic model ForSAFE, described in the text in chapter 2.*

Comment: “P3L3. Please provide more background. For the external reader SWETHRO has no meaning or context.”

Answer: *We provided some more background in chapter 2.3 (P4L3-4). There is also a reference to a paper describing SWETHRO for the reader who wishes to know details about the monitoring (on P3L2).*

Comment: “P3L17. Does ‘factors affecting’ mean sensitive parameters? Can you site previous sensitivity studies?”

Answer: *We meant parameters used to model the weathering, not only the sensitive parameters. We changed the word factors to parameters, to clarify. We cited two previous sensitivity studies in chapter 4.3. (P3L17, P9L16-17)*

Comment: “P4L14. There is more sand at Hissmossa (Table 2) but more less Quartz (Table 3). How is this?”

5 **Answer:** *Texture and total chemistry have been analysed separately and show that there are more material in the sand fraction in Hissmossa than in Västra Torup, and more Si in Västra Torup than in Hissmossa. Quartz is not only found in sand and sand does not only consist of quarts, especially not in relatively young till soils, which can explain the observed pattern. We added in the description of the sites that texture has been measured (P4L7).*

10 **Comment:** “P4L20. It is surprising that a fixed value is used for soil moisture given (a) that soil moisture is an ‘important’ parameter (as noted by the authors), and that (b) the determinants of soil moisture (texture, bulk density and organic matter) are very different between both sites (as noted by the authors).”

15 **Answer:** *A steady state model cannot use a value that varies with time, so in that sense it has to be a fixed value in PROFILE. It does not have to be the same value for all sites though, but there are no measurements of soil moisture for the sites, nor are there usually any for sites that are used for PROFILE modelling. The values used are based on observation of the vegetation on the sites, which gives a “soil moisture class”, which is translated into a soil moisture value that is supposed to represent average soil moisture over a whole forest rotation, for all layers. In this study it happens to be the same for both sites. Yes, it is very crude. Modelling it, as in ForSAFE, should be a lot better. We clarified in the text in chapter 2.3 that it is a site specific value (P4L24).*

Comment: “P4L21. The authors need to provide better context (justification) for the scenarios.”

20 **Answer:** *We rewrote the text explaining the scenarios in chapter 2.4 and expanded the second objective to include the scenarios better (P2L31-32, P4L32-P5L20).*

Comment: “P5L5. The list of scenarios suggests a study objective different than that presented.”

Answer: *We expanded the second objective somewhat to include the scenarios better (P2L31-32).*

Comment: “P5L25. Given the importance of soil moisture, why are these data not shown (Figure or Table)”

25 **Answer:** *Good idea, data on modelled soil moisture is included as a new figure 2.*

Comment: “P5 Figure 1. The difference between PROFILE and ForSAFE in L4 (and L5) at Hissmossa needs more quantitative explanation / support. It might be soil moisture but this is not clearly shown.”

Answer: *In Hissmossa, there is a very strong relationship between difference in soil moisture in the two models and*

5 difference in weathering in the two models. This can be more easily seen now when the soil moisture is shown in a new figure 2. In Västra Torup the differences between soil moisture in the two models are small, except in the organic layer, and differences between weathering rates between the two models are also small, except in the organic layer. There are other differences between the models that affect the difference in weathering rates too, but soil moisture has a large role. See figure below.




Figure 1. Difference in calculated weathering between ForSAFE and PROFILE vs difference in soil moisture between ForSAFE and PROFILE, in percent. The largest relative differences are in the organic layers of both sites, where weathering is very low.

10 **Comment:** "P6 Figures 2 to 6. Many of the figures forces on magnesium or calcium but the sum of base cations is the focus of the text (primarily)."

Answer: We changed figure 2 (now figure 3) to two diagrams that show the sum of base cations instead of Mg. Figure 3 (now figure 4) shows sum of base cations. Figure 4 (now 5) and 6 (now 7) show Mg and P, since the rates of weathering of these two react in the opposite way to acidification, which was the focus of figure 4 (now 5) and the reason behind some of the difference between the base scenario and the background scenario shown in figure 6 (now 7). Figure 5 (now 6) shows Ca as an example of one of the base cations.

Comment: "P7L10. The discussion has notably few citations. . . it is a discussion?"

Answer: We have restructured the discussion and put in some more references. See answer to the general question regarding the discussion.

Comment: “P9L15–L19. These are not surprising conclusions (and more-or-less were previously known).”

Answer: *That two different models give comparable results needs to be verified before using the new, previously untested model (with regards to this process), which is what we have done. For us, the results were not at all self-evident. We changed the wording of the conclusions somewhat, since yes, it is quite self-evident that the ForSAFE model provides more results, but not necessarily that they are useful (P10L26).*

Comment: “Table 3. It appears that surface area is estimated for Clay, Silt and Sand. How are areas for the O horizon estimated?”

Answer: *For the mineral part of the soils in the organic layers, mineralogies and texture analyses from the second layers are used, since there are no texture analyses for the organic layers and the total chemistry analysis of them include the ash of the organic part and thus aren't useful for calculating mineralogy. The surface areas of the mineral part of the organic layers are calculated from the texture in the second layers, but adjusted for how much less mineral matter there is in the organic layers (about 10% of the matter). We clarified this in the text in chapter 2.3 (P4L22-24).*

Manuscript:

Dynamic modelling of weathering rates – the benefit over steady-state modelling

Veronika Kronnäs¹, Cecilia Akselsson¹, Salim Belyazid²

¹Department of Physical Geography and Ecosystem Science, Lund University, 223 62 Lund, Sweden

5 ²Department of Physical Geography, Stockholm University, 106 91 Stockholm, Sweden

Correspondence to: Veronika Kronnäs (veronika.kronnas@nateko.lu.se)

Abstract. Weathering rates are of considerable importance in estimating the acidification sensitivity and recovery capacity of soil, and are thus important in the assessment of the sustainability of forestry in a time of changing climate and growing demands for forestry products. In this study, we modelled rates of weathering in mineral soil at two forested sites in southern Sweden included in a monitoring network, using two models. The aims were to determine whether the dynamic model ForSAFE gives comparable weathering rates as the steady-state model PROFILE, and whether the ForSAFE model provided believable and useful extra information on the response of weathering to changes in acidification load, climate change and land use.

The average weathering rates calculated with ForSAFE were very similar to those calculated with PROFILE for the two modelled sites. The differences between the models regarding the weathering of certain soil layers seemed to be due mainly to differences in calculated soil moisture. The weathering rates provided by ForSAFE vary seasonally with temperature and soil moisture, as well as on longer time scales, depending on environmental changes. Long-term variations due to environmental changes can be seen in the ForSAFE results, for example: the weathering of silicate minerals is suppressed under acidified conditions due to elevated aluminium concentration in the soil, whereas the weathering of apatite is accelerated by acidification. The weathering of both silicates and apatite is predicted to be enhanced by increasing temperature during the 21st century. In this part of southern Sweden, yearly precipitation is assumed to be similar to today's level during the next forest rotation, but with more precipitation in winter and spring and less in summer, which leads to somewhat drier soils in summer, but still increased weathering. In parts of Sweden with bigger projected decrease in soil moisture, weathering might not increase despite increasing temperature.

25 These results show that the dynamic ForSAFE model can be used for weathering rate calculations and that it gives average results comparable to those from the PROFILE model. However, dynamic modelling provides extra information on the variation in weathering rates with time, and offers much better possibilities for scenario modelling.

1 Introduction

Most parts of Sweden are covered with glacial till, composed largely of slowly weathering minerals of granitic origin, the type of rugged landscape with mostly very shallow soil depth described in Krabbendam and Bradwell (2014). This makes

both soils and lakes sensitive to acidification. Two thirds of Sweden is covered by boreal and northern temperate forests, mostly consisting of Norway spruce and Scots pine, together with birch and a few other deciduous trees. Forests are one of Sweden's most important natural resources, and are used for timber (32 million $\text{m}^3 \text{y}^{-1}$ in 2013, Christiansen, 2014), pulp wood (31 million $\text{m}^3 \text{y}^{-1}$ in 2013) and biomass for energy production (6 million $\text{m}^3 \text{y}^{-1}$ in 2013). The last is especially
5 important due to the need to replace fossil fuels with renewable sources of energy (Chu and Majumdar, 2012). Forests and their soil also determine the water quality of most lakes and streams, since the catchments of most lakes are forested and surface water is filtered through forest soils.

During the 1960s, lakes in Scandinavia became increasingly acidified (Odén, 1968). The cause of this was found to be air pollution in the form of atmospheric sulphur and nitrogen (Overrein, 1972), much of it from fossil fuel combustion. The
10 regions most severely affected were those with high deposition of acidifying substances on shallow soils containing base cation poor minerals with low weathering rates and release of base cations (i.e. calcium, magnesium, potassium and sodium) (Galloway et al., 1983). In 1979, the Convention on Long-Range Transboundary Air Pollution (CLRTAP) was formulated by the United Nations Economic Commission for Europe (UNECE). CLRTAP was extended by the addition of several
15 protocols for the mitigation of air pollutants, where participating countries were urged to submit data on emissions of pollutants and ecosystem sensitivity. A need thus emerged for ways of assessing ecosystem sensitivity, and different methods of estimating critical loads of acidity for sulphur and nitrogen for forest and lake ecosystems were developed (Sverdrup and Warfvinge, 1995). One of these was the PROFILE model, developed by researchers at Lund University during the 1990s (Sverdrup and Warfvinge, 1993; Sverdrup et al., 2005).

CLRTAP led to a considerable reduction in the emission of acidifying pollution, and lakes and soils in large parts of
20 acidified areas in Europe slowly started to recover (Engardt et al., 2017; Garmo et al., 2014; Johnson et al., 2018). However, acidifying pollution is still a large and increasing problem in some parts of the world, for example, Southeast Asia (Cho et al., 2016). Forestry is also a potentially acidifying practice, as buffering base cations are removed during harvest (Farley and Werritty, 1989; Akselsson et al., 2016; Zetterberg et al., 2013). Furthermore, as the demand for forest products is growing, while both climate conditions and atmospheric deposition are changing, there is an increasing need to evaluate the sensitivity
25 of forest soils and the weathering of base cations in greater detail, as an aid in forestry planning and regulation. The dynamic ecosystem model ForSAFE (Wallman et al., 2005; Belyazid et al., 2006), which consists of a dynamic development of the PROFILE model, together with models for tree growth and decomposition, has the potential to do this.

The aims of this study were:

- to investigate whether ForSAFE gives comparable weathering rates to those estimated with the PROFILE model, and to explain the results based on differences in the formulation of the models, and
30
- to investigate the seasonal, inter-annual and decadal weathering dynamics provided by ForSAFE for different scenarios, representing important ecological issues: acidification, climate change and nutrient removal through land use.

2 Methods

The PROFILE and ForSAFE models were applied to two spruce forest sites in southernmost Sweden, Västra Torup and Hissmossa, included in the Swedish Throughfall Monitoring Network (SWETHRO) (Pihl Karlsson et al., 2011). Different scenarios for the input parameters were modelled with ForSAFE. ForSAFE-modelled weathering for the base scenario was averaged over the 21st century forest rotation and compared with PROFILE-modelled weathering. The weathering rates from the different scenarios from the ForSAFE model were examined in detail.

2.1 PROFILE

The PROFILE model is a steady-state mechanistic biogeochemistry model, developed at Lund University in the 1990s (Sverdrup and Warfvinge, 1993; Warfvinge and Sverdrup, 1995). It has been widely used for calculations of critical loads of acidification, weathering as an aid to improving the sustainability of forestry in Europe (including Iceland with its very different mineralogy), North America and East Asia, and has even been applied to agricultural land (Akselsson et al., 2016; Erlandsson et al., 2016; Phelan et al., 2014; Fumoto et al., 2001; Holmqvist et al., 2003; Stendahl et al., 2013). The ecosystem in PROFILE is represented by a soil profile divided into layers, each with its own chemical and physical properties, to which water, nutrients and pollutants are added via atmospheric deposition and litterfall from trees, and from which water, nutrients and pollutants are removed via uptake by trees and downward leaching. Chemical equilibrium reactions and weathering take place in the soil profile. Weathering is modelled using transition state theory, and the parameters affecting it are soil temperature, soil moisture, mineralogy, soil texture, expressed as the exposed mineral surface area, soil density, and the concentrations of H^+ , organic ligands and carbon dioxide, as well as the concentrations of inhibitors: base cations (Ca, Mg, K and Na), Al^{3+} (products of the weathering reaction) and organic acids.

2.2 ForSAFE

The ForSAFE model consists of a dynamic development, SAFE, of the PROFILE model (Alveteg et al., 1995; Martinsson et al., 2005), together with the DECOMP model of the decomposition of soil organic matter (Wallman et al., 2006; Walse et al., 1998), the PnET model of tree growth (Aber and Federer, 1992) and the hydrological PULSE model (Lindström and Gardelin, 1992). ForSAFE was developed to better model the process of recovery from acidification and the effects on ecosystems of forestry and climate change, with dynamic feedbacks between soil chemistry and forest growth. Many parameters used as input data in the PROFILE model are modelled by the ForSAFE model. These include runoff, soil moisture, decomposition of litter and the uptake of nutrients by trees. The model is being continuously developed (Belyazid et al., 2011; Phelan et al., 2016; Zanchi et al., 2014; Yu et al., 2016; Rizzetto et al., 2016; Gaudio et al., 2015). In this study, a ForSAFE version with monthly time steps was used.

2.3 Site descriptions

The characteristics of the two SWETHRO sites, at Västra Torup and Hissmossa, are presented in Table 1 and Table 2. SWETHRO is a Swedish network started in the 1980's to monitor deposition of acidifying substances to Swedish managed forest and how the forest and forest soil is affected by the deposition. Each site consists of a 30 m x 30 m square plot in a forest stand, where throughfall deposition is measured every month, and soil water chemistry parameters are measured with lysimeters at a depth of 50 cm three times per year; at Västra Torup since 1996, and at Hissmossa since 2010. Open field deposition is measured near the stands. Soil chemistry, texture and other properties as well as forest parameters have been measured previously (Tables 1 and 2).

Västra Torup has previously been modelled by Belyazid et al. (2006) with an earlier version of the ForSAFE model, using less detailed input data. Zanchi et al. (2014) have also modelled this site using the same version of ForSAFE as in the present study, as well as most of the input data, with the aim of describing changes in forest ecosystem services in a changing climate.

The forest at Västra Torup was clear cut in 2010, and the site at Hissmossa, 5 km to the north, was introduced into SWETHRO as a replacement site. Hissmossa has previously been modelled with ForSAFE, with the aim of explaining why this site shows continuously elevated concentrations of nitrate in soil water, while Västra Torup did not, prior to clear cutting (Olofsson et al., manuscript). Hissmossa has courser, very sandy soil. Both soils are high in quartz and feldspars. Both sites are highly productive sites for Norway spruce, but was probably grazing lands up to the beginning of the 20th century. The soils are assessed as transition types.

The soil parameters used in the modelling are given in Table 3. Values of the field capacity and wilting point were calculated using the equations given by Balland et al. (2008). Mineral content was calculated from total soil chemistry data using A2M, a mathematical model that uses total chemistry of the soil samples to come up with possible mineral compositions (Posch and Kurz, 2007). For the uppermost, organic layers, mineralogy and texture from the second layers were used, since there are no texture analyses for the organic layers and the total chemistry analyses of the organic layers include the ash of the organic matter. The soil moisture input value for PROFILE is an estimated site specific value based on observations at the sites. In this case the soil moisture value is equal at both sites: $0.2 \text{ m}^3_{\text{soil water volume}} \text{ m}^{-3}_{\text{soil volume}}$ for all layers. The fraction of stones in the soils is also estimated at the time of the soil sampling.

2.4 Scenarios and time series of driver parameters

ForSAFE uses time series of climate parameters, forest management and the deposition of atmospheric pollutants and base cations to the site. A set of these time series, from 1900 to 2100, is here called a scenario. The purpose of the different scenarios used in this study is to investigate how ForSAFE-modelled weathering rates responded to changes in the driving parameters. Thus, the scenarios used consist of a base scenario (BSC), four scenarios in which one aspect of the environment differs from the BSC scenario and a background scenario (BGR) without forestry, acidification and climate change.

The **BSC** scenario represents the actual drivers at the sites from 1900 to today, followed by a reasonably realistic future to the year 2100 with regards to forestry management, climate and deposition. This scenario has been used by Zanchi et al. (2014), and Olofsson et al. (manuscript). The future climate is based on a high-CO₂ emission scenario (SRESA2, modelled with ECHAM5: Nakićenović et al., 2000; Roeckner et al., 2006), with an approximately exponentially increasing temperature during the 21st century. Annual precipitation is almost unaffected by the climate change in this scenario for this part of Sweden up to 2080, after which it increases, but only by about 8%. The distribution of precipitation during the year changes after 2050, with more precipitation during winter and spring and less during summer. Past and future forest management of the sites in the **BSC** scenario is based on normal, but not intensively, managed forestry in Sweden today, with two thinnings (at approximately 30 and 45 years after planting) and clear cuttings approximately every 70 years, where only stem wood is removed. The deposition of pollutants and base cations is based on data from the EMEP programme (Simpson et al., 2012), with SO_x-deposition peaking in 1970 and decreasing sharply after that and nitrogen deposition peaking in 1985 with a smaller decrease after that. Future deposition is assumed to be constant after 2020.

Five scenarios were compared with the **BSC** scenario, where climate, deposition or forest management were changed (for the whole or part of the period 1900 - 2100), while the other input parameters were as in the **BSC** scenario. The scenarios were:

- **BSC**: Base scenario, described above.
- **NFO**: No forestry: no thinning or clear cutting between 1900 and 2100. Deposition and climate change as in **BSC**.
- **WTH**: Whole-tree harvest at clear cutting and thinning from 2010. Deposition and climate change as in **BSC**.
- **NAC**: No acidification: no increase in acidifying deposition after 1900. Forestry and climate change as in **BSC**.
- **NCC**: No climate change: no increase in temperature between 1900 and 2100. Forestry and deposition as in **BSC**.
- **BGR**: Background: no clear cutting or thinning, no increase in acidifying deposition and no climate change.

3 Results

3.1 Weathering rates from PROFILE and ForSAFE

The total weathering rates obtained with ForSAFE with the **BSC** scenario, averaged over a forest rotation, were similar to the weathering rates obtained with PROFILE for all soil layers and modelled elements, and almost equal for many of them (Figure 1). At Västra Torup, the total annual weathering rate of the base cations (Ca, Mg, K and Na) in the root zone (organic layer plus the 50 uppermost cm of the mineral soil, L1-L5) was 115 meq m⁻³ y⁻¹ on average, according to ForSAFE (varying for different months between 51 meq m⁻³ y⁻¹ and 260 meq m⁻³ y⁻¹), and 106 meq m⁻³ y⁻¹ according to PROFILE. At Hissmossa, the total weathering rate of base cations in the root zone (L1-L4) estimated with ForSAFE was 38 meq m⁻³ y⁻¹ (varying from 16 meq m⁻³ y⁻¹ to 86 meq m⁻³ y⁻¹) and 45 meq m⁻³ y⁻¹ according to PROFILE.

The estimated weathering rate of base cations is lower at Hissmossa than that at Västra Torup according to both models. This is due to the coarser soil texture at Hissmossa, leading to a significantly lower exposed mineral surface area. Also, according to field measurements, Hissmossa has a more acid soil solution than Västra Torup, with twice the concentration of inorganic

aluminium at Västra Torup. Dissolved inorganic aluminium, a product of the weathering of silicate minerals, inhibits the weathering of silicate minerals. The rotation periods at Västra Torup and Hissmossa are not the same, so average rates for the forest rotation are not directly comparable since climate changes during the period. The differences in weathering rates between the sites are much larger than the changes in rates because of climate change in the two to three decades differing in rotation period.

Differences in the weathering rates predicted by the two models are greater for soil layers where the differences between the values of soil moisture are higher between the two models (Figure 2), i.e. in the organic layers (where weathering is very small, due to very small mineral mass) and in L4 in Hissmossa. The input value for PROFILE was $0.2 \text{ m}^3_{\text{soil water volume}} \text{ m}^{-3}_{\text{soil volume}}$ for all layers at both these sites. The soil moisture is dynamically modelled in ForSAFE, with average values close to the defined field capacity for the respective layers (Table 3). The average soil moisture at Västra Torup, for the forest rotation 2011 - 2080, was 0.18 - 0.21 in the mineral layers and 0.29 in the thin organic upper layer. In the sandy soil at Hissmossa the average soil moisture in ForSAFE (for the forest rotation 2041 - 2100) was 0.13 - 0.18 in the mineral soil layers and 0.4 in the organic soil layer. The difference between the value of soil moisture used in PROFILE and that calculated by ForSAFE is thus greater at Hissmossa, and the differences in weathering rates between the two models are thus also greater at Hissmossa than at Västra Torup.

3.2 Seasonal, yearly and decadal variation in weathering rates from ForSAFE

The weathering rates obtained with ForSAFE vary seasonally with temperature and soil moisture, as well as on longer time scales, depending, for example, on forest stage, the acidification status of the soil and the climate (Figure 3). On the seasonal scale, weathering is lowest in winter and highest in the warmest period of summer, unless the soil is too dry. Weathering rates during the warmest month of the year are typically 3 to 4 times higher than during the coldest month, except for Ca and P, where weathering in the warmest month is 5 to 8 times higher than in the coldest month. On longer time scales, the yearly average weathering rates can vary by a factor of two during a forest rotation.

3.3 Effect of forestry on weathering

Thinning and clear cutting at Västra Torup increased the weathering of base cations by 9 % in the future forest rotation (2011 - 2080) in the BSC scenario, compared to the NFO scenario with no clear cutting or thinning (Figure 4). Whole-tree harvesting in the WTH scenario increased the weathering by a further one percent. At Hissmossa the increase in weathering between the NFO scenario and the BSC scenario was 14 % for the forest rotation between 2041 and 2100, with a further increase of 2 % for the WTH scenario. The difference in weathering between scenarios occurs during the first half of the forest rotation.

3.4 Effect of acidification on weathering

In ForSAFE, the weathering of silicate minerals is decreased by the acidified conditions in the soils during the second half of the 20th century in the BSC scenario, whereas the weathering of the only P-containing mineral, apatite, is enhanced (Figure 5). The effect of acidification on weathering is smaller than the effects of temperature and soil moisture. For the forest rotation 1941 - 2010 in Västra Torup, the weathering of base cations was 11 % lower in the BSC scenario than in the non-acidification NAC scenario, while the P weathering was 11 % higher. At Hissmossa, for the forest rotation 1973 - 2040 (i.e., mostly after the most acidified period), the weathering of base cations was 6 % lower and the weathering of P 17 % higher in the BSC scenario than in the NAC scenario.

3.5 Effect of climate change on weathering

Temperature has a considerable effect on weathering rates. In the BSC scenario, the yearly average temperature increased from 7°C in the 1990s to 11°C in the 2090s. This leads to an increase in ForSAFE weathering rates of the base cations of 7 % per degree increase in temperature. The increase in temperature is greatest in winter (6°C difference between 1900 - 1930 and 2080 - 2100) and smallest in summer (4°C difference between 1900 - 1930 and 2080 - 2100). In Hissmossa, the weathering rates of Ca in L4 are 44 % to 49 % higher in 2080 - 2100 in the BSC scenario than in the constant climate scenario, NCC, for all seasons (Figure 6).

3.6 Overall effect of forestry, acidification and climate change

The overall effect of human practices on weathering rates, as in the BSC scenario: forestry, historical acidification and climate change, is positive, compared to the background scenario, BGR. Climate change and forestry have a positive effect on silicate weathering, while acidification has a negative effect, but not of such a magnitude that it cancels out the first two. For apatite weathering, the combined effect of climate change, forestry and decreasing acidification is an increase of the weathering in the future, especially for newly planted forest. The weathering-enhancing effect of forestry is also seen in the first part of a forest rotation for silicate weathering, whereas an aging forest has slightly decreasing weathering rates. Increasing temperatures combined with the forestry induced weathering dynamic with higher weathering in young forest, produces a step-like increase in weathering rates of silicates in the BSC scenario (Figure 7).

4 Discussion

4.1 Implications of model differences

The weathering calculations in PROFILE and ForSAFE are based on the same equations, but in ForSAFE they are dynamic, while PROFILE has no time dimension. The models also differ in that several processes are only given as input data into PROFILE while they are modelled dynamically with ForSAFE and that feedbacks between these processes affect the system

in ForSAFE. In the PROFILE model, lack of nutrients because of low weathering can never affect tree growth, since uptake of nutrients to trees are input data. Low soil moisture during summers can also never affect weathering rates in PROFILE, because there are neither seasons nor modelled soil moisture values. PROFILE was developed at a time when climate change was usually not considered, to answer the question of what long term loads of acidity the ecosystem could tolerate (under the premise of unchanging forestry and climate), and for this it was sufficient. As acidification loads decreased, the role of forestry intensity for recovery from acidification increased (Iwald et al., 2013). A more complex model was needed and ForSAFE was developed, which include these processes and feedbacks. We have shown that despite their differences, the two models produce comparable estimates of weathering rates on these two sites.

The PROFILE model has often been used for critical load assessments and weathering estimates. This study shows that the more advanced model ForSAFE is as reliable as the PROFILE model and can be used to gain more information on the variation in weathering rates due to forestry practices, climate changes and temperature change, which could increase our understanding of the dynamics of ecosystem sensitivity. General conclusions regarding acid sensitivity, critical loads and the sustainability of forestry would not change significantly, but our ability to make customised or more detailed forestry plans with regards to intensity of harvest or to take acidification countermeasures would be improved.

4.2 Weathering dynamics in a changing environment

Another parameter that has a significant influence on weathering rates is the temperature. The climate is becoming warmer, and in some regions in Sweden, as elsewhere, it is possibly also becoming drier in the summer (Kjellström et al., 2018). Higher temperatures increase weathering, as shown in our simulations. However, drier conditions inhibit weathering, and dry periods in the summer, when weathering otherwise would be much higher than in the rest of the year, might affect the yearly weathering considerably. These two sites, although having lower soil moisture in the summer on average (Figure 2), does not seem to experience really dry summers more often in these future scenarios than during the 20th century, for the same forest stand age. Future studies, on regions that are believed to become much drier in summer in the future may help elucidate this. Akselsson et al. (2016) calculated the increase in weathering rate due to climate change in the 21st century in Sweden, using the PROFILE model. They found that the increase in weathering rates due to temperature increase up to 2050 varied at different locations in Sweden. The median increase in base cation weathering rate was 20 % for the ECHAM projection and 33 % for the HADLEY projection, which are both equivalent to about 10 % °C⁻¹. This is slightly higher than our result of a 7 % increase per degree increase in temperature. The difference is due to the fact that ForSAFE is a more complex model, with dynamic feedbacks between the uptake by trees, soil solution chemistry, soil moisture and weathering. Forestry also affects weathering. After clear cutting, both soil moisture and soil temperature increase, leading to an increase in weathering rate. As uptake of nutrients to trees are halted and as the remaining litter starts to decompose, concentrations of base cations start to increase (Pirainen et al. 2004). Base cations in soil solution inhibit weathering of base cations in the model (like inorganic aluminium inhibit weathering of aluminium), but the increase in base cations is not sufficient to reduce the rate of weathering, since the soil moisture is still high. With whole-tree harvesting, much of the litter is removed, so that

there are less base cations to be released to soil water through decomposing, and the concentrations of base cations should not increase as much as with stem only harvesting (Ågren et al. 2010). This might be the reason for the very slight increase in weathering following whole-tree harvesting compared to stem only harvesting, found in this study. If base cation concentrations do not increase as much after whole-tree harvesting as after stem only harvesting, this also leads to less leaching of base cations after whole-tree harvest than after stem only harvest. The slightly increased weathering rate and the decreased leaching may explain the diminishing difference in soil conditions with time between whole-tree harvesting and stem harvesting that has been seen in field experiments, despite the fact that a large quantity of base cations is removed from the ecosystem by whole-tree harvesting (Zetterberg et al., 2013).

According to ForSAFE, the weathering of silicate minerals is considerably suppressed by the atmospheric deposition of acidifying substances, whereas the weathering of apatite (P and some of the Ca) was enhanced. The reason for this is the combined effects of H^+ as a driver of weathering and Al^{3+} as an inhibitor of silicate weathering, but not of apatite weathering, since apatite does not contain Al. The solubility of Al increases with lower pH, thus inhibiting the weathering of silicates as the soil acidifies.

4.3 Model limitations and development

The results of this study demonstrate the importance of soil moisture on weathering rates. In PROFILE the soil moisture is an input, previously known from uncertainty studies to be of great importance for the weathering rates (Jönsson et al. 1995, Barkman and Alveteg, 2001), but often based on observation of the site and rough assumptions, whereas it is modelled in ForSAFE with soil texture, precipitation and temperature as inputs. For these two sites, average soil moisture modelled by ForSAFE is similar to the rough estimates of moisture used as input for PROFILE for most of the soil layers. The soil moisture modelled by ForSAFE is also close to the calculated field capacity most of the time. Average soil moisture being close to field capacity could partly be an effect of the monthly time step, which evens out precipitation and gives enough time for draining of excess water each time step. A new version of ForSAFE with a daily time step is under development. A daily time step, with a more realistic time distribution of precipitation, with rainfall events and dry periods in between, affects the calculations of soil moisture on the short term, might affect the seasonal average soil moisture values and might thus affect the predicted weathering rates; giving a greater variability in weathering between drier and wetter periods and potentially shifting the average. A shorter time step would potentially give more accurate results, given that soil moisture is an important parameter for weathering and soil moisture is highly variable on a shorter time scale than monthly.

In the SWETHRO sites, soil moisture and soil temperature are not measured and thus modelled soil moisture can't be compared to measured values. Another future study could model sites where such measurements are made and compare these, for the weathering important parameters, with measurements.

Both the PROFILE model and the ForSAFE model are known to overestimate weathering in the lower soil layers (Stendahl et al., 2013; Zanchi, 2016). The soil horizon C consists of the less weathered parent material at the bottom of the soil profile, where weathering rates are low because the conditions in the soil inhibits weathering, despite the relative abundance of

weatherable minerals. Both PROFILE and ForSAFE currently calculate rather high weathering rates in the C horizon, if this soil layer is included in the calculations. In the modelling presented in this paper only a few centimetres of the C horizon are included, thus the total contribution of weathering from horizon C is small, but the rates per soil volume are equivalent to the layers above. Most of the C-horizon is usually located below the root zone, usually defined as the uppermost 50 cm of mineral soil for spruce forests in Sweden, where more than 90 % of the spruce roots can be located (Rosengren and Stjernquist, 2004) and therefore not included in the modelling. The overestimation of weathering in the lower soil layers by these two models is likely to be, at least partly, due to the lack of calculation of the concentrations of dissolved silica in the soil water in both models. The dissolved silica, being a product of weathering of silicate minerals, acts as an inhibitor on the weathering of these minerals, i.e. all the minerals modelled in this study except apatite. The concentration of dissolved silica in the soil water is currently being included in the ForSAFE model.

When the PROFILE and ForSAFE weathering profiles at Västra Torup and Hissmossa are compared to weathering rates at a nearby site, Skånes Värnsjö, calculated with the depletion method (Stendahl et al., 2013), PROFILE and ForSAFE predict substantially higher weathering rates in the lower soil horizons, in line with the above discussion on overestimation in the lower layers. The weathering rates modelled in the upper horizons by PROFILE and ForSAFE are, on the other hand, lower than the rates obtained with the depletion method. However, the depletion method does not calculate present-day weathering, but average weathering in the soil layer since deglaciation. The weathering rates have varied with time, both because new soils have more easily weatherable material and weathers much faster than older soils (Starr and Lindroos, 2006) and because environmental conditions have varied since the end of the last glaciation. This means that weathering rates calculated with methods that calculate average weathering since the deglaciation, such as the depletion method, should generally be higher than PROFILE and ForSAFE weathering rates, except for the lower soil layers, since the weathering front moves down .

5 Conclusions

We have shown that despite the differences between PROFILE and ForSAFE, the two models give comparable estimates of annual weathering rates.

The PROFILE model has often been used for critical load assessments and weathering estimates. This study shows that the more advanced model, ForSAFE, can be used to gain much more information on the variation in weathering rates in response to forestry and climate change.

The results from ForSAFE presented in this paper demonstrate that weathering rates vary considerably; between seasons, between years and on longer time scales. This dynamic behaviour can be of importance in nutrient leaching and nutrient availability to the trees: during seasons with high nutrient demand there might be risk of nutrient deficiency, even though there might be higher availability of nutrients than demand and nutrient losses through leaching during other seasons.

References

- Aber, J.D. and Federer, C.A.: A Generalized, Lumped-Parameter Model of Photosynthesis, Evapotranspiration and Net Primary Production in Temperate and Boreal Forest Ecosystems. *Oecologia* 92: 463 – 474, 1992.
- Akselsson, C., Olsson, J., Belyazid, S., Capell, R.: Can increased weathering rates due to future warming compensate for base cation losses following whole-tree harvesting in spruce forests? *Biogeochemistry* 128(1 - 2): 89 – 105, 2016.
- Ågren, A., Buffam, I., Bishop, K. and Laudon, H. Sensitivity of pH in a boreal stream network to a potential decrease in base cations caused by forest harvest. *Can. J. Fish. Aquat. Sci.* 67: 1116–1125, 2010.
- Alveteg, M., Sverdrup, H. and Warfvinge, P.: Regional assessment of the temporal trends in soil acidification in southern Sweden, using the SAFE model. *Water, Air, Soil Pollut.* 85, 2509 – 2514, 1995.
- Balland, V., Pollacco, J.A.P., Arp, P.A.: Modeling soil hydraulic properties for a wide range of soil conditions. *Ecological modelling* 219, 300–316, 2008.
- Barkman, A., Alveteg, M. Identifying potentials for reducing uncertainty in critical load calculations using the PROFILE model. *Water, Air, and Soil Pollution* 125: 33–54, 2001.
- Belyazid, S., Westling, O., Sverdrup, H.: Modelling changes in forest soil chemistry at 16 Swedish coniferous forest sites following deposition reduction. *Environmental Pollution* 144, 596 – 609, 2006.
- Belyazid, S., Sverdrup, H., Kurz, D., Braun, S.: Exploring Ground Vegetation Change for Different Deposition Scenarios and Methods for Estimating Critical Loads for Biodiversity Using the ForSAFE-VEG Model in Switzerland and Sweden. *Water, Air and Soil Pollution* 216:289–317, 2011.
- Christiansen, L. (red.): Swedish Statistical Yearbook of Forestry 2014. Swedish Forestry Agency. ISBN 978-91-87535-05-5, 2014.
- Cho, Y., Kang, H., Jeon, E.-C.: The Effects of Reactive Nitrogen (Nr) Compounds on the Acidification in Soil and Water Environment Ecosystems and the Mitigation Strategy. *KJEE* 49(1): 1 - 10 (in Korean), 2016.
- Chu, S., Majumdar, A.: Opportunities and challenges for a sustainable energy future. *Nature*. 8/16/2012, Vol. 488 Issue 7411, p 294 – 303, 2012.
- Engardt, M., Simpson, D., Schwikowski, M., Granat, L.: Deposition of sulphur and nitrogen in Europe 1900–2050. Model calculations and comparison to historical observations. *Tellus B: Chemical and Physical Meteorology*, 69:1, 1328945, DOI: 10.1080/16000889.2017.1328945, 2017.
- Erlandsson, M., Oelkers, E.H., Bishop, K., Sverdrup, H., Belyazid, S., Ledesma, J.L.J., Köhler, S.J.: Spatial and temporal variations of base cation release from chemical weathering on a hillslope scale. *Chemical Geology* 441: 1 – 13, 2016.
- Farley, D.A., Werritty, A.: Hydrochemical budgets for the Loch Dee experimental catchments, southwest Scotland (1981 - 1985). *Journal of Hydrology*, 109, 351 – 368, 1989.

- Fumoto, T., Shindo, J., Oura, N., Sverdrup, H.: Adapting the PROFILE model to calculate the critical loads for east Asian soils by including volcanic glass weathering and alternative aluminium solubility system. *Water, Air and Soil Pollution* 130: 1247–1252, 2001.
- Galloway, J.N., Norton, S.A., Robbins Church, M.: Freshwater acidification from atmospheric deposition of sulfuric acid: A conceptual model. *Environ. Sci. Technol.*, Vol. 17, No. 11, 1983.
- 5 Garmo, Ø. A., Skjelkvåle, B.L., de Wit, H.A., Colombo, L., Curtis, C., Fölster, J., Hoffmann, A., Hruška, J., Høgåsen, T., Jeffries, D.S., Keller, W.B., Krám, P., Majer, V., Monteith, D.T., Paterson, A.M., Rogora, M., Rzychon, D., Steingruber, S., Stoddard, J.L., Vuorenmaa, J., Worsztynowicz, A.: Trends in Surface Water Chemistry in Acidified Areas in Europe and North America from 1990 to 2008. *Water, Air, & Soil Pollution*, March 2014, 225:1880, 2014.
- 10 Gaudio, N., Belyazid, S., Gendre, X., Mansat, A., Nicolas, M., Rizzetto, S., Sverdrup, H., Probst, A.: Combined effect of atmospheric nitrogen deposition and climate change on temperate forest soil biogeochemistry: A modeling approach. *Ecological Modelling* 306: 24 – 34, 2015.
- Hellsten, S., Stadmark, J., Pihl Karlsson, G., Karlsson, P.-E., Akselsson, C.: Increased concentrations of nitrate in forest soil water after windthrow in southern Sweden. *Forest Ecology and Management* 356, 234–242, 2015.
- 15 Holmqvist, J., Øgaard, A.F., Öborn, I., Edwards, A.C., Mattsson, L., Sverdrup, H.: Application of the PROFILE model to estimate potassium release from mineral weathering in Northern European agricultural soils. *European Journal of Agronomy* 20 149 – 163, 2003.
- Iwald, J., Löfgren, S., Stendahl, J., Karlton, E. Acidifying effect of removal of tree stumps and logging residues as compared to atmospheric deposition. *For. Ecol. Manage.* 290, 49–58. <http://dx.doi.org/10.1016/j.foreco.2012.06.022>. 2013.
- 20 Johnson, J., Graf Pannatier, E., Carnicelli, S., Cecchini, G., Clarke, N., Cools, N., Hansen, K., Meesenburg, H., Nieminen, T.M., Pihl-Karlsson, G., Titeux, H., Vanguelova, E., Verstraeten, A., Vesterdal, L., Waldner, P., Jonard, M.: The response of soil solution chemistry in European forests to decreasing acid deposition. *Global Change Biology*. England, Mar. 31. ISSN: 1365 - 2486. 2018.
- Jönsson, C., Warfvinge, P. and Sverdrup, H. Uncertainty in predicting weathering rate and environmental stress factors with the PROFILE model. *Water, Air, and Soil Pollut.* 81, 1. 1995.
- 25 Kjellström, E., Nikulin, G., Strandberg, G., Bøssing Christensen, O., Jacob, D., Keuler, K., Lenderink, G., van Meijgaard E., Schär, C., Somot, S., Lund Sørland, S., Teichmann, C., Vautard, R.: European climate change at global mean temperature increases of 1.5 and 2 °C above pre-industrial conditions as simulated by the EURO-CORDEX regional climate models. *Earth Syst. Dynam.*, 9, 459–478, 2018.
- 30 Krabbendam, M., Bradwell, T. Quaternary evolution of glaciated gneiss terrains: pre-glacial weathering vs. glacial erosion. *Quaternary Science Reviews* 95: 20-42. 2014.
- Lindström, G. and Gardelin, M.: Modelling groundwater response to acidification, Report from the Swedish integrated groundwater acidification project. 1992.

- Martinsson, L., Alveteg, M., Kronnäs, V., Sverdrup, H., Westling, O., Warfvinge, P.: A regional perspective on present and future soil chemistry at 16 Swedish forest sites. *Water, Air, and Soil Pollution* 162: 89–105, 2005.
- Nakićenović, N., Alcamo, J., Davis, G., de Vries, B., Fenhann, J., Gaffin, S., Gregory, K., Grubler, A., Jung, T.Y., Kram, T., La Rovere, E.L., Michaelis, L., Mori, S., Morita, T., Pepper, W., Pitcher, H.M., Price, L., Riahi, K., Roehrl, A., Rogner, H.-H., Sankovski, A., Schlesinger, M., Shukla, P., Smith, S.J., Swart, R., van Rooijen, S., Victor, N., Dadi, Z. (eds.): IPCC Special Report on Emissions Scenarios: a special report of Working Group III of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge. 2000.
- Odén, S.: Nederbördens och luftens försurning: dess orsaker, förlopp och verkan i olika miljöer. Ecology Committee Bulletin No.1. Swedish National Science Research Council, Stockholm ISSN: 0375-2208 (in Swedish with English summary). 1968.
- 10 Olofsson, J., Akselsson, C., Zanchi, G., Belyazid, S. Identifying soil parameters decisive for nitrogen leaching in forests through a combined empirical and modelling approach. Manuscript.
- Overrein, L.N.: Sulphur Pollution Patterns Observed; Leaching of Calcium in Forest Soil Determined *Ambio*, Vol. 1, No. 4 (Sep., 1972), pp. 145 – 147, 1972.
- 15 Piirainen, S., Finer, L., Mannerkoski, H. and Starr, M. Effects of forest clear-cutting on the sulphur, phosphorus and base cations fluxes through podzolic soil horizons. *Biogeochemistry* 69: 405–424, 2004.
- Phelan, J., Belyazid, S., Kurz, D., Guthrie, S., Cajka, J., Sverdrup, H., Waite, R.: Estimation of soil base cation weathering rates with the PROFILE model to determine critical loads of acidity for forested ecosystems in Pennsylvania, USA: Pilot Application of a Potential National Methodology. *Water Air and Soil Pollution* 225(9), 2014.
- Phelan, J., Belyazid, S., Jones, P., Cajka, J., Buckley, J., Clark, C.: Assessing the Effects of Climate Change and Air 20 Pollution on Soil Properties and Plant Diversity in Northeastern U.S. hardwood forests: Model simulations from 1900 - 2100. *Water, Air and Soil Pollution*, 227:84, 2016.
- Pihl Karlsson, G., Akselsson, C., Hellsten, S., Karlsson, P.-E.: Reduced European emissions of S and N - Effects on air concentrations, deposition and soil water chemistry in Swedish forests. *Environmental Pollution* 159, 3571 – 3582, 2011.
- Posch, M., Kurz, D.: A2M - A program to compute all possible mineral modes from geochemical analyses. *Computers & 25 Geosciences* 33, 563–572, 2007.
- Rizzetto, S., Belyazid, S., Gegout, J.C., Nicolas, M., Alard, D., Corcket, E., Gaudio, N., Sverdrup, H., Probst, A.: Modelling the impact of climate change and atmospheric N deposition on French forests biodiversity. *Environmental Pollution* 213: 1016 – 1027, 2016.
- Roeckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kornbluh, L., Manzini, E., Schlese, U., Schulzweida, 30 U.: Sensitivity of simulated climate to horizontal and vertical resolution in the ECHAM5 atmosphere model. *Journal of Climate* 19, 3771 – 3791, 2006.
- Rosengren, U. and Stjernquist, I.: *Gå på djupet! Om rottdjup och rotproduktion i olika skogstyper*. Alnarp : SUFOR, ISBN 91-576-6617-2. 2004.

- Simpson, D., Benedictow, A., Berge, H., Bergström, R., Emberson, L. D., Fagerli, H., Flechard, C. R., Hayman, G. D., Gauss, M., Jonson, J. E., Jenkin, M. E., Nyíri, A., Richter, C., Semeena, V. S., Tsyro, S., Tuovinen, J.-P., Valdebenito, Á., and Wind, P.: The EMEP MSC-W chemical transport model – technical description, *Atmos. Chem. Phys.*, 12, 7825 – 7865, 2012.
- 5 Starr, M., Lindroos, A.-J. Changes in the rate of release of Ca and Mg and normative mineralogy due to weathering along a 5300-year chronosequence of boreal forest soils. *Geoderma* 133: 269–280, 2006.
- Stendahl, J., Akselsson, C., Melkerud, P.A., Belyazid, S.: Pedon-scale silicate weathering: comparison of the PROFILE model and the depletion method at 16 forest sites in Sweden. *Geoderma* 211: 65 – 74, 2013.
- Sverdrup, H. and Warfvinge, P.: Calculating field weathering rates using a mechanistic geochemical model PROFILE. *Applied Geochemistry* 8: 273 – 283, 1993.
- 10 Sverdrup, H. and Warfvinge, P.: Critical loads of acidity for Swedish forest ecosystems. - *Ecol. Bull. (Copenhagen)* 44: 75 – 89, 1995.
- Sverdrup, H., Martinson, L., Alveteg, M., Moldan, F., Kronnäs, V., Munthe, J.: Modeling Recovery of Swedish Ecosystems from Acidification. *Ambio*, Vol. 34, No. 1, pp. 25 – 31, 2005.
- 15 Wallman, P., Svensson, M.G.E., Sverdrup, H., Belyazid, S.: ForSAFE—an integrated process-oriented forest model for long-term sustainability assessments. *Forest Ecology and Management* 207, 19 – 36, 2005.
- Wallman, P., Belyazid, S., Svensson, M.G.E., Sverdrup, H.: DECOMP – a semi-mechanistic model of litter decomposition. *Environmental Modelling & Software* 21, 33–44, 2006.
- Walse, C., Berg, B., Sverdrup, H.: Review and synthesis of experimental data on organic matter decomposition with respect to the effects of temperature, moisture and acidity. *Environmental Review* 6, 25 – 40, 1998.
- 20 Warfvinge, P., Sverdrup, H.: Critical loads of acidity to Swedish forest soils. Methods, data and results. Reports in ecology and environmental engineering 1995:5, Lund University. 1995.
- Yu L., Belyazid S., Akselsson C., van der Heijden G., Zanchi G.: Storm disturbances in a Swedish forest-A case study comparing monitoring and modelling. *Ecological Modelling* 320: 102 – 113, 2016.
- 25 Zanchi, G.: Modelling nutrient transport from forest ecosystems to surface waters: The model ForSAFE2D. Lund, Sweden: Lund University, Faculty of Science, Department of Physical Geography and Ecosystem Science. ISBN: 9789185793686. 2016.
- Zanchi, G., Belyazid, S., Akselsson, C., Yu, L.: Modelling the effects of management intensification on multiple forest services: a Swedish case study. *Ecological Modelling* 284: 48 – 59, 2014.
- 30 Zetterberg, T., Olsson, B.A., Löfgren, S., von Brömssen, C., Brandtberg, P.-O.: The effect of harvest intensity on long-term calcium dynamics in soil and soil solution at three coniferous sites in Sweden. *Forest Ecology and Management* 302, 280–294, 2013.

Tables

Table 1. Characteristics of the two sites.

	Västra Torup	Hissmossa
Coordinates (°N, °E)	56.135, 13.510	56.181, 13.515
Active years	1988 - 2010	2010 -
Year of planting	1941	1973
Year of clear cutting	2010	-
Standing stem biomass (g m ⁻²) (year in parenthesis)	18841 (2010)	10559 (2011)
Measured throughfall		
Precipitation (mm)	430 - 780	460 - 730
S deposition (kg ha ⁻¹ y ⁻¹)	4.5 - 27 *	3.6 - 6.9
N deposition (kg ha ⁻¹ y ⁻¹)	6.2 - 12	6.8 - 11
Cl deposition (kg ha ⁻¹ y ⁻¹)	21 - 50	33 - 87
Ca+Mg+Na+K deposition (kg ha ⁻¹ y ⁻¹)	31 - 57	39 - 80
Measured soil water chemistry		
pH	4.4 - 4.9	4.2 - 4.5
SO ₄ -S (mg l ⁻¹)	0.8 - 7.3	2.1 - 4.4
Cl (mg l ⁻¹)	3.2 - 20	17 - 51
NO ₃ -N (mg l ⁻¹)	0 - 0.1	0.5 - 3.3
NH ₄ -N (mg l ⁻¹)	0 - 0.2	0 - 0.1
Ca (mg l ⁻¹)	0.2 - 1.0	0.2 - 1.7
Mg (mg l ⁻¹)	0.2 - 1.0	0.6 - 1.9
Na (mg l ⁻¹)	2.8 - 8.4	12 - 23
K (mg l ⁻¹)	0.1 - 1.1	0.2 - 1.0
Inorganic Al (mg l ⁻¹)	0.2 - 3.4	0.6 - 5.3
Organic Al (mg l ⁻¹)	0 - 0.4	0.6 - 1.1
Al-tot (mg l ⁻¹)	0.4 - 3.7	1.5 - 6.2
TOC (mg l ⁻¹)	3.5 - 15	8.2 - 21

* Decreasing steeply with time

Table 2. Measured soil parameters for the five soil layers (O, A, AB, B and C) at the two sites. Above: thickness of layer, bulk density, percentage organic matter, estimated percentage stones, measured size fractions, pH, exchangeable ions, cation exchange capacity, base saturation, fraction of carbon and nitrogen. Below: total chemistry of all dry soil matter.


Horizon	Thickness (m)	Bulk density (kg m ⁻³)	OM (% of DW)	Stoniness (%)	Clay Silt Sand			pH H ₂ O	Exchangeable ions (µeq g ⁻¹)					CEC (µeq g ⁻¹)	BS (%)	Tot-C (g (kg DW) ⁻¹)	Tot-N (g (kg DW) ⁻¹)	
					(% of mineral soil)				Al	H	Na	K	Mg					Ca
Västra Torup																		
O	0.05	181	87	0				4.0	29	84.5	<0.1	13.0	27.9	50.1	205	43.0	543	20.9
A	0.06	959	6	20	5	27	68	4.1	31	16.5	<0.1	1.0	0.7	0.8	50	4.9	34	2.0
AB	0.20	1062	5	20	5	31	64	4.6	27	6.1	<0.1	0.6	0.4	1.2	36	6.4	25	1.7
B	0.20	1279	4	20	3	21	76	4.8	16	1.3	<0.1	0.4	0.1	0.6	18	6.5	18	1.3
C	0.04	1446	2	20	0	17	83	4.9	13	4.8	<0.1	0.4	0.1	0.5	19	5.0	8	0.6
Hissmossa																		
O	0.05	394	65	0				3.5	45	63.5	3.7	8.1	21.4	26.4	164	36.8	391	19.8
A	0.13	909	8	10	0	5	91	3.8	36	15.3	0.6	1.4	3.8	2.9	60	17.6	46	3.0
AB	0.10	1075	8	10	1	8	89	4.6	27	4.6	0.4	0.8	2.5	2.3	37	17.8	38	3.0
B	0.28	1276	3	10	0	9	88	4.5	12	0.3	0.4	0.7	2.3	2.3	17	34.2	17	2.2
C	0.04	1316	3	10	0	8	88	4.7	11	0.7	0.4	0.7	2.4	2.3	17	36.1	14	2.0

Horizon	Total chemistry of mineral and organic matter (mg (kg DW) ⁻¹)									
	Si	Al	Ca	Fe	K	Mg	Mn	Na	P	Ti
Västra Torup										
O	33100	5060	1760	2540	2540	623	167	1440	789	337
A	349000	50000	7110	20000	26300	1540	386	15900	253	4280
AB	335000	56400	8190	28600	26600	2610	469	16300	401	4510
B	341000	58600	8920	22600	27700	2950	436	17100	540	3700
C	348000	59500	9820	24400	28700	3240	493	17800	602	4240
Hissmossa										
O	77300	13500	1970	4370	5710	580	103	3540	787	844
A	322000	51500	5040	19200	28400	822	277	14600	166	3540
AB	322000	65700	7030	29100	29600	2070	459	16900	217	3340
B	340000	66000	7490	27600	30700	2140	419	18900	231	3290
C	329000	70500	8840	28800	33600	2670	822	19000	451	3030

5


Table 3. Soil input data to the models, standard values (partial pressure of CO₂ and gibbsite constant) or calculated from measured soil parameters at the two sites (mineral area, field capacity, wilting point, field saturation and percentage of minerals). The modelled layers L1 - L5 correspond to soil layers O, A, AB, B and C in the two soils. Hissmossa L5 is below the modelled root zone of 50 cm.

Layer	Mineral area (10 ⁶ m ² m ⁻³)	pCO ₂	Kgibb	FC	WP	FS	Quartz	K-feldspar	Albite	Anorthite	Muscovite	Epidote	Hornblende	Apatite	Illite	Vermiculite1	Vermiculite2	Chlorite1	Chlorite2
							SiO ₂	KAlSi ₃ O ₈	NaAlSi ₃ O ₈	CaAl ₂ Si ₂ O ₈	a	b	c	d	e	f	g	h	i
Västra Torup																			
L1	214161	10	6.5	0.31	0.11	0.87	50	17	19	2.2	2.4	2.0	0.4	0.2	1.3	0.6	0.2	0.4	0.2
L2	1131959	20	7.6	0.21	0.06	0.68	50	17	19	2.2	2.4	2.0	0.4	0.2	1.3	0.6	0.2	0.4	0.2
L3	1334007	20	8.6	0.24	0.06	0.64	46	16	19	2.4	4.4	2.1	0.7	0.2	2.3	1.0	0.4	0.6	0.4
L4	1167398	20	9.2	0.22	0.06	0.54	45	16	20	2.5	4.1	2.2	0.8	0.3	2.2	1.1	0.5	0.7	0.5
L5	909226	20	9.2	0.18	0.03	0.47	44	17	20	2.6	3.3	2.3	0.9	0.3	1.9	1.3	0.5	0.8	0.5
Hissmossa																			
L1	143372	10	6.5	0.42	0.17	0.80	42	18	17	1.5	1.2	1.4	0.2	0.1	2.9	0.3	0.1	0.2	0.1
L2	330775	20	7.6	0.18	0.05	0.65	42	18	17	1.5	1.2	1.4	0.2	0.1	2.9	0.3	0.1	0.2	0.1
L3	491284	20	8.6	0.20	0.06	0.58	39	17	19	2.1	2.9	1.8	0.6	0.1	5.2	0.7	0.3	0.4	0.3
L4	534872	20	9.2	0.14	0.03	0.51	41	17	21	2.5	4.8	1.7	0.5	0.1	3.5	0.7	0.3	0.4	0.3
L5	538935	20	9.2	0.15	0.03	0.50	37	18	21	2.7	5.5	2.0	0.7	0.2	3.7	0.9	0.4	0.5	0.4



10

Figures



5 **Figure 1.** Weathering rates ($\text{meq m}^{-2} \text{y}^{-1}$) calculated with the PROFILE model and the ForSAFE model (averages over a forest rotation, BSC scenario), for the sites at Västra Torup and Hissmossa, for soil layers L1 (top layer) to L5 (bottom layer at ~50 cm depth). The time period is from one clear cut to the next and is different for the two sites: 2011-2080 for Västra Torup and 2041-2100 for Hissmossa. Note that the rates are shown here per layer, so that the bars show directly how much of the total weathering each soil layer contributes. For Hissmossa, L5 is shown, even though it lies below the root zone and is not included in calculations of weathering rates in the root zone.




Figure 2. Soil moisture in all soil layers, BSC scenario, forest rotation 2010-2080 in Västra Torup and 2040-2100 in Hissmossa, compared to wilting point and PROFILE input soil moisture.






Figure 3. Modelled Ca+Mg+K+Na weathering in Västra Torup (above) and Hissmossa (below) from 1950 to 2100 (note the difference in scale for the two sites). PROFILE calculates the average weathering rates for the time period represented by the input values, while monthly weathering values were calculated with ForSAFE, using the BSC scenario.



5 **Figure 4.** Yearly average weathering of base cations in the whole soil profile, for the BSC scenario, the whole-tree harvest WTH scenario, and the NFO scenario without any clear cutting or thinning. The years of clear cuts in the BSC and WTH scenarios are marked with vertical lines. In Västra Torup clear cuts are in the years 1940, 2010 and 2080 and in Hissmossa in 1972, 2040 and 2101.




Figure 5. Comparison of weathering rates of Mg and P in soil layer L4 in the non-acidification scenario NAC and the base scenario BSC ($\text{meq m}^{-3} \text{y}^{-1}$). With the mineralogy of these sites, Mg is only weathered from silicate minerals and P is only weathered from apatite. One dot represent one month.




Figure 6. The effect of the increased temperature of the BSC scenario on Ca weathering in L4 at Hissmossa, compared to the NCC scenario with no climate change, shown as averages for seasons over periods of 30 years. Winter = December, January and February, spring = March, April and May, summer = June, July and August and autumn = September, October and November.





Figure 7. Weathering of Mg (from silicates) and P (from apatite) at Västra Torup, under the **BSC** scenario and the **BGR** scenario with neither acidification, climate change nor forestry.