

1 **Refining physical aspects of soil quality and soil health when** 2 **exploring the effects of soil degradation and climate change on** 3 **biomass production: an Italian case study.**

4 Antonello Bonfante^{1,3}, Fabio Terribile^{2,3}, Johan Bouma⁴

5 ¹Institute for Mediterranean Agricultural and Forest Systems - CNR-ISAFOM, Ercolano, Italy

6 ²University of Naples Federico II, Department of Agriculture, Portici, (NA), Italy

7 ³University of Naples Federico II, CRISP Interdepartmental Research Centre

8 ⁴Em. Prof. Soils Science, Wageningen University, The Netherlands

9 *Correspondence to:* Antonello Bonfante (antonello.bonfante@cnr.it)

10 **ABSTRACT**

11 This study focuses on soil physical aspects of soil quality and - health with the objective to define
12 procedures with worldwide rather than only regional applicability, reflecting modern developments in
13 soil physical and agronomic research and addressing important questions regarding possible effects of
14 soil degradation and climate change. In contrast to water and air, soils cannot, even after much research,
15 be characterized by a universally accepted quality definition and this hampers the internal and external
16 communication process. Soil quality expresses the capacity of the soil to function. Biomass production
17 is a primary function, next to filtering and organic matter accumulation, and can be modeled with soil-
18 water-atmosphere-plant simulation models, as used in the agronomic yield-gap program that defines
19 potential yields (Y_p) for any location on earth determined by radiation, temperature and standardized
20 crop characteristics, assuming adequate water and nutrient supply and lack of pests and diseases. The
21 water-limited yield (Y_w) reflects, in addition, the often limited water availability at a particular
22 location. Real yields (Y_a) can be considered in relation to Y_w to indicate yield gaps, to be expressed
23 in terms of the indicator: $(Y_a/Y_w) \times 100$. Soil data to calculate Y_w for a given soil type (the genoform)
24 should consist of a range of soil properties as a function of past management (various phenoforms)
25 rather than as a single “representative” dataset. This way a Y_w -based characteristic soil quality range
26 for every soil type is defined, based on semi-permanent soil properties. In this study effects of subsoil
27 compaction, overland flow following surface compaction and erosion were simulated for six soil series
28 in the Destre Sele area in Italy, including effects of climate change. Recent proposals consider soil

29 health, which appeals more to people than soil quality and is now defined by separate soil physical,
30 chemical and –biological indicators. Focusing on the soil function biomass production, physical soil
31 health at a given time of a given type of soil can be expressed as a point (defined by a measured Ya)
32 on the defined soil quality range for that particular type of soil, thereby defining the seriousness of the
33 problem and the scope for improvement. The six soils showed different behavior following the three
34 types of land degradation and projected climate change up to the year 2100. Effects are expected to be
35 major as reductions of biomass production of up to 50% appear likely under the scenarios. Rather than
36 consider soil physical, chemical and biological indicators separately, as proposed now elsewhere for
37 soil health, a sequential procedure is discussed logically linking the separate procedures.

38 **Keywords:** soil quality, soil health, climate change, simulation modeling, water-limited crop yield.

39

40

41 1. INTRODUCTION

42 The concept of Soil Health has been proposed to communicate the importance of soils to stakeholders
43 and policy makers (Moebius-Clune et al., 2016). This follows a large body of research on soil quality,
44 recently reviewed by Bünemann et al., (2018). The latter conclude that research so far has hardly
45 involved farmers and other stakeholders, consultants and agricultural advisors. This may explain why
46 there are as yet no widely accepted, operational soil quality indicators in contrast to quality indicators
47 for water and air which are even formalised into specific laws (e.g. EU Water Framework Directive).
48 This severely hampers effective communication of the importance of soils which is increasingly
49 important to create broad awareness about the devastating effects of widespread soil degradation. New
50 soil health initiatives, expanding the existing soil quality discourse, deserve therefore to be supported.
51 A National Soil Health Institute has been established in the USA (www.soilhealthinstitute.org) and
52 Cornell University has published a guide for its comprehensive assessment after several years of
53 experimentation (Moebius-Clune et al, 2016). Soil health is defined as:”*the continued capacity of the*
54 *soil to function as a vital living ecosystem that sustains plants, animals and humans*”(NRCS, 2012).
55 Focusing attention in this paper to soil physical conditions, the Cornell assessment scheme (Moebius-
56 Clune et.al, 2016) distinguishes three soil physical parameters: wet aggregate stability, surface and

57 subsurface hardness to be characterized by penetrometers and the available water capacity (AWC:
58 water held between 1/3 and 15 bar). The National Soil Health Institute reports 19 soil health parameters,
59 including 5 soil physical ones: water-stable aggregation, penetration resistance, bulk density, AWC
60 and infiltration rate.

61 Techniques to determine aggregate stability and penetrometer resistance have been introduced many
62 years ago (e.g. Kemper and Chepil, 1965; Lowery, 1986; Shaw et al., 1943). Aggregate stability is a
63 relatively static feature as compared with dynamic soil temperature and moisture content with
64 drawbacks in terms of (1) lack of uniform applied methodology (e.g. Almajmaie et al., 2017), (2) the
65 inability of dry and wet sieving protocols to discriminate between management practices and soil
66 properties (Le Bissonnais, 1996; Pulido Moncada et al., 2013) and above all: (3) the mechanical work
67 applied during dry sieving is basically not experienced in real field conditions (Díaz-Zorita et al., 2002).
68 Measured penetrometer resistances are known to be quite variable because of different modes of
69 handling in practice and seasonal variation. Finally, the AWC is a static characteristic based on fixed
70 values as expressed by laboratory measurements of the pressure head for “field capacity” and “wilting
71 point” that don’t correspond with field conditions (e.g. Bouma, 2018).

72 These drawbacks must be considered when suggesting the introduction for general use as physical soil
73 health indicators. More recent developments in soil physics may offer alternative approaches, to be
74 explored in this paper, that are more in line with the dynamic behavior of soils.

75 The definition of soil health is close to the soil quality concept introduced in the 1990’s: “*the capacity*
76 *of the soil to function within ecosystem and land-use boundaries to sustain productivity, maintain*
77 *environmental quality and promote plant and animal health*” (Bouma, 2002; Bünemann et al., 2018;
78 Doran and Parkin, 1994; Karlen et al., 1997). Discussions in the early 2000’s have resulted in a
79 distinction between *inherent* and *dynamic* soil quality. The former would be based on relatively stable
80 soil properties as expressed in soil types that reflect the long-term effect of the soil forming factors
81 corresponding with the basic and justified assumption of soil classification that soil management
82 should not change a given classification. Still, soil functioning of a given soil type can vary
83 significantly as a result of the effects of past and current soil management, even though the name of
84 the soil type does not change (this can be the soil series as defined in USDA Soil Taxonomy (Soil
85 Survey Staff, 2014 as expressed in Table 1) but the lowest level in other soil classification systems
86 would also apply. In any case, the classification should be unambiguous. *Dynamic* soil quality would
87 reflect possible changes as a result of soil use and management over a human time scale, which can
88 have a semi-permanent character when considering , for example, subsoil plowpans (e.g. Mobius-
89 Clune et al, 2016). This was also recognized by Droogers and Bouma, (1997) and Rossiter and Bouma

90 (2018) when defining different soil phenoforms reflecting effects of land use for a given genoform as
91 distinguished in soil classification. Distinction of different soil phenoforms was next translated into a
92 range of characteristically different soil qualities by using simulation techniques (Bouma and Droogers,
93 1998). The term soil health appears to have a higher appeal for land users and citizens at large than the
94 more academic term soil quality, possibly because the term “health” has a direct connotation with
95 human wellbeing in contrast to the more distant and abstract term: “quality”. Humans differ and so do
96 soils; some soils are genetically more healthy than others and a given soil can have different degrees
97 of health at any given time, which depends not only on soil properties but also on past and current
98 management and weather conditions. Mobius-Clune et al. (2016) have recognized the importance of
99 climate variation by stating that their proposed system only applies to the North-East of the USA and
100 its particular climate and soil conditions. This represents a clear limitation and could in time lead to a
101 wide variety of local systems with different parameters that would inhibit effective communication to
102 the outside world. This paper will therefore explore possibilities for a science-based systems approach
103 with general applicability. To apply the soil health concept to a wider range of soils in other parts of
104 the world, the attractive analogy with human health not only implies that “health” has to be associated
105 with particular soil individuals, but also to climate zones. In addition, current questions about soil
106 behavior often deal with possible effects of climate change. In this paper, the proposed systems analysis
107 can – in contrast to the procedures presented so far- also deal with this issue. Using soils as a basis for
108 the analysis is only realistic when soil types can be unambiguously defined, as was demonstrated by
109 Bonfante and Bouma (2015) for five soil series in the Italian Destre Sele area that will also be the focus
110 of this study. In most developed countries where soil surveys have been completed, soil databases
111 provide extensive information on the various soil series, including parameters needed to define soil
112 quality and soil health in a systems-analysis as shown, for example, for clay soils in the Netherlands
113 (Bouma and Wösten, 2016). The recent report of the National Academy of Sciences, Engineering and
114 Medicine (2018) emphasizes the need for the type of systems approaches as followed in this study.
115 The basic premise of the Soil Health concept, as advocated by Moebius-Clune et.al. 2016 and others,
116 is convincing. Soil characterization programs since the early part of the last century have been
117 exclusively focused on soil chemistry and soil chemical fertility and this has resulted in not only
118 effective recommendations for the application of chemical fertilizers but also in successful pedological
119 soil characterization research. But soils are living bodies in a landscape context and not only chemical
120 but also physical and biological processes govern soil functions. The Soil Health concept considers
121 therefore not only soil chemical characteristics, that largely correspond with the ones already present
122 in existing soil fertility protocols, but also with physical and biological characteristics that are
123 determined with well defined methods, with particular emphasis on soil biological parameters

124 (Moebius-Clune et al, 2016). However, the proposed soil physical methods by Moebius-Clune et al (
125 2016) don't reflect modern soil physical expertise and procedures need to have a universal rather than
126 a regional character, while pressing questions about the effects of soil degradation and future climate
127 change need to be addressed as well. The proposed procedures do not allow this. Explorative simulation
128 studies can be used to express possible effects of climate change as, obviously, measurements in future
129 are not feasible. Also, only simulation models can provide a quantitative, interdisciplinary integration
130 of soil-water-atmosphere-plant processes that are key to both the soil quality and soil health definitions,
131 as mentioned above.

132 In summary, the objectives of this paper are to: (i) explore alternative procedures to characterize: "soil
133 physical quality and health" applying a systems analysis by modeling the soil-water-atmosphere-plant
134 system, an analysis that is valid anywhere on earth; (ii) apply the procedure to develop quantitative
135 expressions for the effects of different forms of soil degradation and (iii) explore effects of climate
136 change for different soils also considering different forms of soil degradation.

137

138 **2. MATERIALS AND METHODS.**

139 **2.1. Soil functions as a starting point**

140 The soil quality and - health definitions both mention: "*the continued capacity of a soil to function*".
141 Soil functions have therefore a central role in the quality and health debate. EC (2006) defined the
142 following soil functions: (1) Biomass production, including agriculture and forestry; (2) Storing,
143 filtering and transforming nutrients, substances and water; (3) Biodiversity pool, such as habitats,
144 species and genes; (4) Physical and cultural environment for humans and human activities; (5) Source
145 of raw material; (6) Acting as carbon pool, and (7) Archive of geological and archaeological heritage.
146 Functions 4, 5 and 7 are not covered in this contribution since, if considered relevant, specific measures
147 have to be taken to set soils apart by legislative measures. The other functions are directly and indirectly
148 related to function 1, biomass production. Of course, soil processes not only offer contributions to
149 biomass production, but also to filtering, biodiversity preservation and carbon storage. Inter- and
150 transdisciplinary approaches are needed to obtain a complete characterization, requiring interaction
151 with other disciplines, such as agronomy, hydrology, ecology and climatology and, last but not least,
152 with stakeholders and policy makers. Soil functions thus contribute to ecosystem services and,
153 ultimately, to all seventeen UN Sustainable Development Goals (e.g. Bouma, 2016, 2014; Keesstra et
154 al., 2016). However, in the context of this paper, attention will be focused on function 1, biomass
155 production.

156 Soil physical aspects play a crucial role when considering the role of soil in biomass production, as
157 expressed by Function 1, which is governed by the dynamics of the soil-water-atmosphere-plant system
158 in three ways:

159 (1) Roots provide the link between soil and plant. Rooting patterns as a function of time are key factors
160 for crop uptake of water and nutrients. Deep rooting patterns imply less susceptibility to moisture stress.
161 Soil structure, the associated bulk densities, and the soil water content determine whether or not roots
162 can penetrate the soil. When water contents are too high, either because of the presence of a water table
163 or of a dense, slowly permeable soil horizon impeding vertical flow, roots will not grow because of
164 lack of oxygen. For example, compact plow-pans, resulting from applying pressure on wet soil by
165 agricultural machinery, can strongly reduce rooting depth. In fact, soil compaction is a major form of
166 soil degradation that may affect up to 30% of soils in some areas. (e.g. FAO & ITPS, 2015).

167 (2) Availability of water during the growing season is another important factor that requires, for a start,
168 infiltration of all rainwater into the soil and its containment in the unsaturated zone, constituting “green-
169 water” (e.g. Falkenmark and Rockström, 2006). When precipitation rates are higher than the infiltrative
170 capacity of soils water will flow laterally away over the soil surface, possibly leading to erosion and
171 reducing the amount of water available for plant growth, and:

172 (3) the climate and varying weather conditions among the years govern biomass production. Rainfall
173 varies in terms of quantities, intensities and patterns. Radiation and temperature regimes vary as well.
174 In this context, definitions of location-specific potential yield (Y_p), water-limited yield (Y_w) and actual
175 yield (Y_a) are important, as will be discussed later .

176 Soil Function 2 requires first soil infiltration of water followed by good contact between percolating
177 water and the soil matrix, where clay minerals and organic matter can adsorb cations and organic
178 compounds, involving chemical processes that will be considered when defining soil chemical quality.
179 However, not only the adsorptive character of the soil is important but also the flow rate of applied
180 water that can be affected by climatic conditions or by management when irrigating. Rapid flow rates
181 generally result in poor filtration as was demonstrated for viruses and fecal bacteria in sands and silt
182 loam soils (Bouma, 1979).

183 Soil Functions 3 and 6 are a function of the organic matter content of the soil (or %C) ,the quantity of
184 which is routinely measured in chemical soil characterization programs (also in the soil health protocols
185 mentioned earlier that also define methods to measure soil respiration). The organic matter content of
186 soils is highly affected by soil temperature and moisture regimes and soil chemical conditions. Optimal

187 conditions for rootgrowth in terms of water, air and temperature regimes will also be favorable for soil
 188 biological organisms, linking soil functions 1, 3 and 6.

189 When defining soil physical aspects of soil quality and soil health, focused on soil function 1,
 190 parameters will have to be defined that integrate various aspects, such as: (1) weather data, (2) the
 191 infiltrative capacity of the soil surface, considering rainfall intensities and quantities, (3) rootability as
 192 a function of soil structure, defining thresholds beyond which rooting is not possible, and: (4) hydraulic
 193 and root extraction parameters that allow a dynamic characterization of the soil-water-plant-
 194 atmosphere system that can only be realized by process modeling, that requires these five parameters
 195 and modeling is therefore an ideal vehicle to realize interdisciplinary cooperation. Simulation models
 196 of the soil-water-atmosphere-plant system are ideal to integrate these various aspects.

197 **2.2. The role of dynamic modeling of the soil-water-plant-atmosphere system**

198 When analysing soil quality and soil health, emphasis must be on the dynamics of *vital, living*
 199 *ecosystems* requiring a dynamic approach that is difficult to characterize with static soil characteristics
 200 (such as bulk density, organic matter content and texture) except when these characteristics are used
 201 as input data into dynamic simulation models of the soil-water-plant-climate system. Restricting
 202 attention to soil physical characteristics, hydraulic conductivity (K) and moisture retention properties
 203 ($\Theta(h)$) of soils are applied in such dynamic models. Measurement procedures are complex and can only
 204 be made by specialists, making them unsuitable for general application in the context of soil quality
 205 and health. They can, however, be easily derived from *pedotransferfunctions* that relate static soil
 206 characteristics such bulk density, texture and %C to these two properties, as recently summarized by
 207 Van Looy et al. (2017). The latter soil characteristics are available in existing soil databases and are
 208 required information for the dynamic models predicting biomass production.

209 Simulation models of the soil-water-plant-atmosphere system, such as the Soil Water Atmosphere,
 210 Plant model (SWAP) (Kroes et al., 2008) to be discussed later in more detail, integrate weather
 211 conditions, infiltration rates, rooting patterns and soil hydrological conditions in a dynamic systems
 212 approach that also allows exploration of future conditions following climate change. The worldwide
 213 agronomic Yield-Gap program (www.yieldgap.org) can be quite helpful when formulating a soil
 214 quality and – health program with a global significance. So-called water-limited yields (Yw) can be
 215 calculated, **assuming optimal soil fertility and lack of pests and diseases** (e.g Gobbett et al., 2017;
 216 van Ittersum et al., 2013; Van Oort et al., 2017). Yw reflects climate conditions at any given location
 217 in the world as it is derived from potential production (Yp) that reflects radiation, temperature and
 218 basic plant properties, **assuming that water and nutrients are available and pests and diseases**
 219 **don't occur**. Yw reflects local availability of water. Yw is usually, but not always, lower than Yp. Yw

220 can therefore act as a proxy value for physical soil quality, focusing on function 1. Note that Y_p and
221 Y_w , while providing absolute science-based points of reference, include assumptions on soil fertility
222 and crop health.

223 Actual yields (Y_a) are often, again, lower than Y_w (e.g. Van Ittersum et al, 2013). The ratio Y_a/Y_w is
224 an indicator of the so-called “yield-gap” showing how much potential there is at a given site to improve
225 production (www.yieldgap.org) (Bouma, 2002). When multiplied with 100, a number between 1 and
226 100 is obtained as a quantitative measure of the “yield gap” for a given type of soil . Y_w can be
227 calculated for a non-degraded soil. Y_a should ideally be measured but can also be calculated as was
228 done in this exploratory study (in terms of Y_w values) on the basis of the assumed effects of different
229 forms of soil degradation, such as subsoil soil compaction, poor water infiltration at the soil surface
230 due to surface compaction or crusting and erosion. This requires introduction of a compact layer
231 (plowpan) in the soil, a reduction of rainfall amounts with the volume of estimated overland flow and
232 by removing topsoil. Each variant of the analyzed soil series represents a Phenoform. In this exploratory
233 study Y_a values were simulated but, ideally, field observations should be made in a given soil type to
234 define effects of management as explored, for example, by Pulleman et al., (2000) for clay soils and
235 Sonneveld et al., (2002) for sandy soils. They developed Phenoforms based on different %C of surface
236 soil and such Phenoforms could also have been included here to provide a link with soil biology but
237 field data were not available to do so, Field work identifying phenoforms includes important
238 interaction with farmers as also mentioned by Moebius-Clune et al, (2016). Sometimes, soil
239 degradation processes, such as erosion, may be so severe that the soil classification (the soil genoform)
240 changes. Then, the soil quality and soil health discussion shifts to a different soil type.

241 This approach will now be explored with a particular focus on the Mediterranean environment. Physical
242 soil quality is defined by Y_w for each soil, considering a soil without assumed degradation phenomena
243 (the reference) and for three variants (hypothetical Y_a , expressed in terms of Y_w) with: (1) a compacted
244 plowlayer, (2) a compacted soil surface resulting in overland flow, and (3) removal of topsoil following
245 erosion, without a resulting change in the soil classification. This way a characteristic range of Y_w
246 values is obtained for each of the six soil series, reflecting positive and negative effects of soil
247 management and representing a range of soil physical quality values of the particular soil series
248 considered. Within this range an actual value of Y_a will indicate the soil physical health of the particular
249 soil at a given time and its position within the range of values will indicate the severity of the problem
250 and potential for possible improvement.

251 The ratio $(Y_a/Y_w) \times 100$ is calculated to obtain a numerical value that represents “soil health” as a point
252 value, representing actual conditions. Health is relatively low when real conditions occur in the lower

253 part of the soil quality range for that particular soil and relatively high when it occurs in the upper
254 range. Again, in this exploratory study measured values (at current climate conditions) for Ya have not
255 been made, so Ya only applies to the three degraded soil forms being distinguished here where
256 hypothetical effects of soil degradation have been simulated as related to the corresponding calculated
257 Yw values. Of course, actual measured Ya values can't be determined at all when considering future
258 climate scenario's and simulation is the only method allowing exploratory studies. We assume that
259 climate change will not significantly affect soil formation processes until the year 2100. Soil properties
260 will therefore stay the same.

261 To allow estimates of the possible effects of climate change RCP 8.5- IPCC scenario will be applied.
262 Obviously, only computer simulations can be used when exploring future conditions, another important
263 reason to use dynamic simulation modeling in the context of characterizing soil quality and soil health.
264 The approach in this paper extends earlier studies on soil quality for some major soil types in the world
265 that did not consider aspects of soil health nor effects of climate change (Bouma, 2002; Bouma et al.,
266 1998).

267

268 **2.3. Simulation modeling**

269 The Soil–Water–Atmosphere–Plant (SWAP) model (Kroes et al., 2008) was applied to solve the soil
270 water balance. SWAP is an integrated physically-based simulation model of water, solute and heat
271 transport in the saturated–unsaturated zone in relation to crop growth. In this study only the water flow
272 module was used; it assumes unidimensional vertical flow processes and calculates the soil water flow
273 through the Richards equation. Soil water retention $\theta(h)$ and hydraulic conductivity $K(\theta)$ relationships
274 as proposed by van Genuchten (1980) were applied. The unit gradient was set as the condition at the
275 bottom boundary. The upper boundary conditions of SWAP in agricultural crops are generally
276 described by the potential evapotranspiration ET_p , irrigation and daily precipitation. Potential
277 evapotranspiration was then partitioned into potential evaporation and potential transpiration according
278 to the LAI (Leaf Area Index) evolution, following the approach of Ritchie (1972). The water uptake
279 and actual transpiration were modeled according to Feddes *et al.* (1978), where the actual transpiration
280 declines from its potential value through the parameter α , varying between 0 and 1 according to the
281 soil water potential.

282 The model was calibrated and validated by measured soil water content data at different depths for
283 Italian conditions (Bonfante et al., 2010; Crescimanno and Garofalo, 2005) and in the same study area
284 by (Bonfante et al., 2011, 2017). In particular, the model was evaluated in two farms inside of Destra
285 Sele area, on three different soils (Udic Calciustert, Fluventic Haplustept and Typic Calciustoll), under

286 maize crop (two cropping seasons) during a Regional project “Campania Nitrati” (Regione Campania,
287 2008) (Table.2).

288 Soil hydraulic properties of soil horizons in the area were estimated by the pedotransfer function (PTF)
289 HYPRES (Wösten et al., 1999). A reliability test of this PTF was performed on $\theta(h)$ and $k(\theta)$ measured
290 in the laboratory by the evaporation method (Basile et al., 2006) on 10 undisturbed soil samples
291 collected in the Destra Sele area. The data obtained were compared with estimates by HYPRES and
292 were considered to be acceptable ($RMSE = 0.02 \text{ m}^3 \text{ m}^{-3}$) (Bonfante et al., 2015).

293 Simulations were run considering a soil without assumed degradation phenomena (the reference) and
294 for three variants with a compacted plowlayer, surface runoff and erosion, as discussed above:

295 (i) The compacted plowlayer was applied at -30cm (10 cm of thickness) with following physical
296 characteristics: 0.30 WC at saturation, 1.12 n, 0.004 "a" and Ks of 2 cm/day. Roots were restricted to
297 the upper 30 cm of the soil. (ii) Runoff from the soil surface was simulated removing ponded water
298 resulting from intensive rainfall events. Rooting depth was assumed to be 80 cm. (iii) Erosion was
299 simulated for the Ap horizon, reducing the upper soil layer to 20 cm. The maximum rooting depth was
300 assumed to be 60 cm (A+B horizon) with a higher root density in the Ap horizon.

301 Variants were theoretical but based on local knowledge of the Sele Plain. Compaction is relevant
302 considering the highly specialized and intensive horticulture land use of the Sele plain which typically
303 involves repetitive soil tillage at similar depth. Runoff and erosion easily occur at higher altitude plain
304 areas especially where the LON0, CIF0/RAG0, GIU0 soil types occur (Figure 1).

305

306 **2.4. Soils in the Destra Sele area in Italy.**

307 The “Destra Sele” study area, the plain of the River Sele (22,000 ha, of which 18,500 ha is farmed) is
308 situated in the south of Campania, southern Italy (Figure 1). The main agricultural production consists
309 of irrigated crops (maize, vegetables and fruit orchards), greenhouse-grown vegetables and mozzarella
310 cheese from water buffalo herds. The area can be divided into four different landform classes
311 (hills/footslopes, alluvial fans, fluvial terraces and dunes) with heterogeneous parent materials in which
312 twenty different soil series were distinguished (within Inceptisol, Alfisol, Mollisol, Entisol and Vertisol
313 soil orders) (Regione Campania, 1996), according to Soil Taxonomy (Soil Survey Staff, 1999). Six soil
314 series were selected in the area to test application of the soil quality and soil health concepts.
315 Representative data for the soils are presented in Table 1.

316 Decision trees were developed to test whether the selection process of the soil series was based on
317 stable criteria, allowing extrapolation of results from measured to unmeasured locations when
318 considering effects of climate change. While extrapolation in space of soil series data has been a

319 common procedure in soil survey (e.g. Soil Survey Staff, 2014; Bouma et al., 2012), extrapolation in
320 time has not received as much attention. A basic principle of many taxonomic soil classification
321 systems is a focus on stable soil characteristics when selecting diagnostic criteria for soil types. Also,
322 emphasis on morphological features allows, in principle, a soil classification without requiring
323 elaborate laboratory analyses (e.g. Soil Survey Staff, 2014). A given soil classification should, in order
324 to obtain permanent names, not change following traditional management measures, such as plowing.
325 This does, however, not apply to all soils and then a different name will have to be assigned.
326 This way, soil classification results in an assessment of the (semi)-permanent physical constitution of
327 a given soil in terms of its horizons and textures. That is why soil quality is defined for each soil type
328 as a characteristic range of Yw values, representing different effects of soil management that have not
329 changed the soil classification.

330

331 **2.5. Climate information**

332 Future climate scenarios were obtained by using the high resolution regional climate model (RCM)
333 COSMO-CLM (Rockel et al., 2008), with a configuration employing a spatial resolution of
334 0.0715°(about 8 km), which was optimised over the Italian area. The validations performed showed
335 that these model data agree closely with different regional high-resolution observational datasets, in
336 terms of both average temperature and precipitation in Bucchignani et al. (2015) and in terms of
337 extreme events in Zollo et al. (2015).

338 In particular, the Representative Concentration Pathway (RCP) 8.5 scenario was applied, based on the
339 IPCC (Intergovernmental Panel on Climate Change) modelling approach to generate greenhouse gas
340 (GHG) concentrations (Meinshausen et al., 2011). Initial and boundary conditions for running RCM
341 simulations with COSMO-CLM were provided by the general circulation model CMCC-CM
342 (Scoccimarro et al., 2011), whose atmospheric component (ECHAM5) has a horizontal resolution of
343 about 85 km. The simulations covered the period from 1971 to 2100; more specifically, the CMIP5
344 historical experiment (based on historical greenhouse gas concentrations) was used for the period
345 1976–2005 (Reference Climate scenario - RC), while, for the period 2006–2100, a simulation was
346 performed using the IPCC scenario mentioned. The analysis of results was made on RC (1971–2005)
347 and RCP 8.5 divided into three different time periods (2010–2040, 2040–2070 and 2070–2100). Daily
348 reference evapotranspiration (ET_0) was evaluated according to Hargreaves and Samani, (1985)
349 equation (HS). The reliability of this equation in the study area was performed by Fagnano et al.,
350 (2001) comparing the HS equation with the Penman–Monteith (PM) equation (Allen et al., 1998).

351 Under the RCP 8.5 scenario the temperature in Destra Sele is expected to increase approximately two
352 degrees celsius respectively every 30 years to 2100 starting from the RC. The differences in
353 temperature between RC and the period 2070–2100 showed an average increase of minimum and
354 maximum temperatures of about 6.2°C (for both min and max). The projected increase of temperatures
355 produces an increase of the expected ET_0 . In particular, during the maize growing season, an average
356 increase of ET_0 of about 18% is expected until 2100.

357

358 **3. RESULTS**

359 **3.1. Soil physical quality of the soil series, as expressed by Y_w , under current and future climates.**

360 Soil physical quality of the six soil series, expressed as calculated Y_w values for the reference climate
361 and for future climate scenario RCP 8.5, expressed for three time periods are shown in Figure 2.
362 Considering current climate conditions, the Longobarda and Cifariello soils with loamy textures have the
363 highest values, while the sandy soil Lazzaretto is lower. This can be explained by higher water retention
364 of loamy soils (180 and 152 mm of AWC in the first 80 cm for Longobarda and Cifariello respectively)
365 compared to the sandy soil (53 mm of AWC in the first 80 cm for Lazzaretto). The effects of climate
366 change are most pronounced and quite clear for the two periods after 2040. Reductions compared with
367 the period up to 2040 range from 20-40%, the highest values associated with sandier soil textures. This
368 follows from the important reduction of projected rainfall during the cropping season (Figure 3) ranging
369 from an average value of 235 (± 30) mm in the 2010-2040 period to 185 (± 26) mm (-21%) and to 142
370 (± 24) mm (-40%) in the 2040-2070 and 2070-2100 periods, respectively (significant at $p < 0.01$). The
371 figure also includes a value for Y_p , potential production (under RC with optimal irrigation), which is 18
372 $t\ ha^{-1}$, well above the Y_w values. Only a Y_p value is presented for current conditions because estimates
373 for future climates involve too many unknown factors.

374

375 **3.2. Projected effects of soil degradation processes**

376 *3.2.1. Projected effects of subsoil compaction.*

377 The projected effects of soil compaction are shown in Figure 4. The effects of compaction are very strong
378 in all soils, demonstrating that restricting the rooting depth has major effects on biomass production.
379 Compared with the reference, reductions in Y_w do not occur in the first time window (2010-2040), while
380 the projected lower precipitation rates are expected to have a significant effect on all soils, strongly
381 reducing Y_w values by 44-55% with, again, highest values in the sandy soils. Clearly, any effort to
382 increase effective rooting patterns of crops should be a key element when considering attempts to combat
383 effects of climate change. Data indicate that reactions are soil specific.

384 *3.2.2. Projected effects of overland flow.*

385 Results, presented in Figure 5, show relatively small differences (5% or less) with results presented in
386 Figure 2 that was based on complete infiltration of rainwater. This implies that surface crusting or
387 compaction of surface soil, leading to lower infiltration rates and more surface runoff, does not seem to
388 have played a major role here in the assumed scenario's. Real field measurements may well produce
389 different results. Even though projected future climate scenario's predict rains with higher intensities,
390 that were reflected in the climate scenario's being run, the effects of lower precipitation, as shown in
391 Figure 3, appear to dominate.

392 *3.2.3. Projected effects of erosion.*

393 Results, presented in Figure 6, show significant differences with results presented in Figure 2. Yw values
394 are lower in all soils as compared with reference climate conditions, but loamy and clayey subsoils still
395 can still provide moisture to plant roots, leading to relatively low reductions of Yw (e.g 10%-20% for the
396 Longobarda and Cifariello soils, with an AWC of the remaining 60 cm depth of 150 mm and 120 mm,
397 respectively) even though topsoils with a relatively high organic matter content have been removed. Next
398 are the Picciola, Giuliarossa and San Vito soils with reductions between 35 and 45%, all with an AWC
399 of appr. 107 mm. Effects of erosion are strongest in the sandy Lazzaretto soil, where loss of the A horizon
400 has a relatively strong effect on the moisture supply capacity of the remaining soil with an AWC of 33
401 mm up to the new 60 cm depth. The reduction with the reference level is 30%, which is relatively low
402 because the reference level was already low as well. Projected effects of climate change are again strong
403 for all soils, leading to additional reductions of Yw of appr. 30%.

404 *3.2.4. Indicators for the soil quality range.*

405 Figure 7 presents the physical soil quality ranges for all the soil series, expressed separately as bars for
406 each of the climate periods. The $(Y_a/Y_w) \times 100$ index illustrates that ranges are significantly different.
407 The upper limit is theoretically 100%. But Van Ittersum et al (2013) have suggested that an 80% limit
408 would perhaps be more realistic. Figure 7, ranging to 100%, shows the lower limits for the ranges to vary
409 from e.g. 35 (Longobarda) to 55 (Lazaretto) for the reference climate with values for the three
410 phenofoms in between. $(Y_a/Y_w) \times 100$ decreases as a projected reaction to climate change (e.g. 20 for
411 Longobarda and 40 for Lazaretto). This provides important signals for the future.

412 As discussed, the presented ranges are soil specific and are based on hypothetical conditions associated
413 with different forms of land degradation. Field research may well result in different ranges also possibly
414 considering different soil degradation factors beyond compaction, surface runoff and erosion. Still,
415 principles involved are identical. Ranges presented in Figure 7 represent a physical soil quality range that
416 is characteristic for that particular type of soil. Actual values (Y_a) will fit somewhere in this range and
417 will thus indicate how far they are removed from the maximum and minimum value, thereby presenting
418 a quantitative measure for soil physical health. This can not only be important for communication

419 purposes but it also allows a judgment of the effects of different forms of degradation in different soils
420 as well as potential for improvement.

421

422 **4. DISCUSSION**

423 Linking the soil quality and soil health discussion with the international research program on the *yield*
424 *gap* allows direct and well researched expressions for crop yields, defining soil function 1, as discussed
425 above. The potential yield (Y_p) and water-limited yield (Y_w) concepts apply worldwide and provide,
426 therefore, a sound theoretical basis for a general soil quality/health classification, avoiding many local
427 and highly diverse activities as reviewed by Büneman et al, (2017). Of course, different indicator crops
428 will have to be defined for different areas in the world.

429 Linking soil quality and health to specific and well defined soil types is essential because soil types, such
430 as the soil series presented in this paper, uniquely reflect soil forming processes in a landscape context.
431 They provide much more information than just a collection of soil characteristics, such as texture, organic
432 matter content and bulk density. They are well known to stakeholders and policy makers in many
433 countries. A good example is the USA where State Soils have been defined.

434 Defining (semi-permanent) soil quality for specific soil types, in terms of a characteristic range of Y_w
435 values reflecting effects of different forms of land management, represents a quantification of the more
436 traditional Soil Survey interpretations or land evaluations where soil performance was judged by
437 qualitative, empirical criteria. (e.g. FAO, 2007, Bouma et al 2012).

438 In this exploratory study, hypothetical effects of three forms of soil degradation were tested. In reality,
439 soil researchers should go to the field and assemble data for a given soil series as shown on soil maps,
440 establishing a characteristic range of properties, following the example of Pulleman et al (2000) for a
441 clay soil and Sonneveld et al, (2002) for a sand soil, but not restricting attention to %C, as in these two
442 studies, but including at least bulk density measurements. This way, a characteristic series of Phenofoms
443 can be established. Physical soil quality (for a given soil type=Genoform) has a characteristic range of
444 Y_w values, as shown in Figure 7. Soil physical health at any given time is reflected by the position of
445 real Y_a values within that range and can be expressed by a number $(Y_a/Y_w) \times 100$.

446 One could argue that this “range” acts as a “thermometer” for a particular type of soil allowing
447 determination of the physical “health” of a given soil by the placement of Y_a .

448 But calculating Y_w has implications beyond defining physical soil quality and health. As discussed, Y_w
449 not only reflects the effects of soil moisture regimes but also assumes that chemical conditions for crop
450 growth are optimal and that pests and diseases don't occur. Defining Y_w can thus function as a starting
451 point of a general soil quality/soil health discussion. If Y_a is lower than Y_w the reasons must be found.
452 Is it lack of water, nutrients or occurrence of pests and diseases? Irrigation may be difficult to realize but

453 fertility can be restored rather easily and many methods, biological or chemical, are available to combat
 454 pests and diseases. If Phenofoms would be included that consider different %C of surface soil (as
 455 discussed above), also low %C contents could be a reason for relatively low Yw values. This would cover
 456 soil biological quality with %C acting as proxy value. This way, the Yw analysis can be a logical starting
 457 point for follow-up discussions defining appropriate forms of future soil management.

458 This paper has focused on physical aspects but the proposed procedure has potential to extend the
 459 discussion to chemical and biological aspects, to be further explored in future. Rather than consider the
 460 physical, chemical and biological aspects separately, each with their own indicators as proposed by
 461 Moebius-Clune et al, (2016), following a logical and interconnected sequence considering first
 462 pedological (soil types), and soil physical (Yw) characterizations, to be followed by analysing chemical
 463 and biological aspects, that can possibly explain relatively low Ya values, could be more effective. This
 464 is the more relevant because definition of reproducible biological soil health parameters are still object
 465 of study (Wade et al., 2018) and %C might be an acceptable proxy for soil biology for the time being.
 466 Recent tests of current soil-health protocols have not resulted in adequately expressing soil conditions in
 467 North Caolina (Roper et al, 2017), indicating the need for further research as suggested in this paper.

468

469 5. CONCLUSIONS

470

- 471 1. Lack of widely accepted, operational criteria to express soil quality and soil health is a barrier
 472 for effective external communication of the importance of soil science
- 473 2. Using well defined soil types as “carriers” of information on soil quality and soil health can
 474 improve communication to stakeholders and the policy arena.
- 475 3. A universal system defining soil quality and soil health is needed based on reproducible
 476 scientific principles that can be applied all over the world, avoiding a multitude of different
 477 local systems. Models of the soil-water-plant-atmosphere system can fulfil this role.
- 478 4. Connecting with the international *yield gap* program, applying soil-water-plant-atmosphere
 479 simulation models, will facilitate cooperation with agronomists which is essential to quantify
 480 the important soil function 1: biomass production.
- 481 5. The proposed system allows an extension of classical soil classification schemes, defining
 482 genoforms, by allowing estimates of effects of various forms of past and present soil
 483 management (phenofoms) within a given genoform that often strongly affects soil
 484 performance. Quantitative information thus obtained can improve current empirical and
 485 qualitative soil survey interpretations and land evaluation.
- 486 6. Rather than consider physical, chemical and biological aspects of soil quality and - health

- 487 separately, a combined approach starting with pedological and soil physical aspects followed
488 by chemical and biological aspects, all to be manipulated by management, is to be preferred.
- 489 7. Only the proposed modeling approach allows exploration of possible effects of climate change
490 on future soil behaviour which is a necessity considering societal concerns and questions.
- 491 8. Field work, based on existing soil maps to select sampling locations for a given genoform, is
492 needed to identify a characteristic range of phenoforms for a given genoform, which, in turn,
493 can define a characteristic soil quality range by calculating Y_w values.

494

495

496 **6 ACKNOWLEDGMENTS**

497 We acknowledge Dr. Eugenia Monaco and Dr. Langella Giuliano for the supporting in the analysis
498 of climate scenario. The “Regional Models and Geo-Hydrogeological Impacts Division”, Centro
499 Euro-Mediterraneo sui Cambiamenti Climatici (CMCC), Capua (CE) – Italy, and the Dr. Paola
500 Mercogliano and Edoardo Bucchignani for the climate information applied in this work.

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520 **7.REFERENCES**

- 521
- 522 Almajmaie, A., Hardie, M., Acuna, T., Birch, C.: Evaluation of methods for determining soil aggregate
523 stability. *Soil Tillage Res.* 167, 39–45. <http://dx.doi.org/10.1016/j.still.2016.11.003>, 2017
- 524 Allen, R. G., Pereira, L. S., Raes, D., Smith, M. and W, a B.: Crop evapotranspiration - Guidelines for
525 computing crop water requirements - FAO Irrigation and drainage paper 56, *Irrig. Drain.*, 1–15,
526 doi:10.1016/j.eja.2010.12.001, 1998.
- 527 Basile, A., Coppola, A., De Mascellis, R. and Randazzo, L.: Scaling approach to deduce field
528 unsaturated hydraulic properties and behavior from laboratory measurements on small cores, *Vadose*
529 *Zo. J.*, 5(3), 1005–1016, doi:10.2136/vzj2005.0128, 2006.
- 530 Bonfante, A. and Bouma, J.: The role of soil series in quantitative land evaluation when expressing
531 effects of climate change and crop breeding on future land use, *Geoderma*, 259–260, 187–195, 2015.
- 532 Bonfante, A., Basile, A., Acutis, M., De Mascellis, R., Manna, P., Perego, A. and Terribile, F.: SWAP,
533 CropSyst and MACRO comparison in two contrasting soils cropped with maize in Northern Italy,
534 *Agric. Water Manag.*, 97(7), 1051–1062, doi:10.1016/j.agwat.2010.02.010, 2010.
- 535 Bonfante, A., Basile, A., Manna, P. and Terribile, F.: Use of Physically Based Models to Evaluate
536 USDA Soil Moisture Classes, *Soil Sci. Soc. Am. J.*, 75(1), 181, doi:10.2136/sssaj2009.0403, 2011.
- 537 Bonfante, A., Monaco, E., Alfieri, S. M., De Lorenzi, F., Manna, P., Basile, A. and Bouma, J.: Climate
538 change effects on the suitability of an agricultural area to maize cultivation: Application of a new
539 hybrid land evaluation system, *Adv. Agron.*, 133, 33–69, doi:10.1016/bs.agron.2015.05.001, 2015.
- 540 Bonfante, A., Impagliazzo, A., Fiorentino, N., Langella, G., Mori, M. and Fagnano, M.: Supporting
541 local farming communities and crop production resilience to climate change through giant reed
542 (*Arundo donax* L.) cultivation: An Italian case study, *Sci. Total Environ.*, 601–602,
543 doi:10.1016/j.scitotenv.2017.05.214, 2017.
- 544 Bouma, J.: Subsurface applications of sewage effluent, *Plan. uses Manag. L.*, (planningtheuses), 665–
545 703, 1979.
- 546 Bouma, J.: Land quality indicators of sustainable land management across scales, *Agric. Ecosyst.*
547 *Environ.*, 88(2), 129–136, doi:10.1016/S0167-8809(01)00248-1, 2002.
- 548 Bouma, J.: Soil science contributions towards sustainable development goals and their implementation:
549 linking soil functions with ecosystem services, *J. plant Nutr. soil Sci.*, 177(2), 111–120, 2014.
- 550 Bouma, J.: Hydropedology and the societal challenge of realizing the 2015 United Nations Sustainable
551 Development Goals, *Vadose Zo. J.*, 15(12), 2016.
- 552 Bouma, J. and Droogers, P.: A procedure to derive land quality indicators for sustainable agricultural
553 production, *World Bank Discuss. Pap.*, 103–110 [online] Available from:
554 <http://www.archive.org/details/plantrelationsfi00coul>, 1998.
- 555 Bouma, J. and Wösten, J. H. M.: How to characterize good and greening in the EU Common
556 Agricultural Policy (CAP): The case of clay soils in the Netherlands, *Soil Use Manag.*, 32(4), 546–
557 552, 2016.
- 558 Bouma, J., Batjes, N. H. and Groot, J. J. R.: Exploring land quality effects on world food supply1,
559 *Geoderma*, 86(1–2), 43–59, 1998.
- 560 Bucchignani, E., Montesarchio, M., Zollo, A. L. and Mercogliano, P.: High-resolution climate
561 simulations with COSMO-CLM over Italy: performance evaluation and climate projections for the 21st
562 century, *Int. J. Climatol.*, 36(2), 735–756, 2015.
- 563 Bünemann, E. K., Bongiorno, G., Bai, Z., Creamer, R. E., De Deyn, G., de Goede, R., Fleskens, L.,
564 Geissen, V., Kuyper, T. W., Mäder, P. and others: Soil quality--A critical review, *Soil Biol. Biochem.*,
565 120, 105–125, 2018.
- 566 Crescimanno, G. and Garofalo, P.: Application and evaluation of the SWAP model for simulating water
567 and solute transport in a cracking clay soil, *Soil Sci. Soc. Am. J.*, 69(6), 1943–1954, 2005.
- 568 Doran, J. W. and Parkin, T. B.: Defining and assessing soil quality, *Defin. soil Qual. a Sustain.*
569 *Environ.*, (definingsoilqua), 1–21, 1994.

- 570 Droogers, P. and Bouma, J.: Soil survey input in exploratory modeling of sustainable soil management
571 practices, *Soil Sci. Soc. Am. J.*, 61(6), 1704–1710, 1997.
- 572 Fagnano, M., Acutis, M. and Postiglione, L.: Valutazione di un metodo semplificato per il calcolo
573 dell'ET₀ in Campania, *Model. di Agric. sostenibile per la pianura meridionale Gest. delle risorse idriche*
574 *nelle pianure irrigue*. Gutenberg, Salerno, ISBN, 88–900475, 2001.
- 575 Falkenmark, M. and Rockström, J.: The new blue and green water paradigm: Breaking new ground for
576 water resources planning and management, 2006.
- 577 Feddes, R. A., Kowalik, P. J., Zaradny, H. and others: Simulation of field water use and crop yield.,
578 Centre for Agricultural Publishing and Documentation., 1978.
- 579 Van Genuchten, M. T.: A closed-form equation for predicting the hydraulic conductivity of unsaturated
580 soils, *Soil Sci. Soc. Am. J.*, 44(5), 892–898, 1980.
- 581 Gobbett, D. L., Hochman, Z., Horan, H., Garcia, J. N., Grassini, P. and Cassman, K. G.: Yield gap
582 analysis of rainfed wheat demonstrates local to global relevance, *J. Agric. Sci.*, 155(2), 282–299, 2017.
- 583 Hargreaves, G. H. and Samani, Z. A.: Reference crop evapotranspiration from temperature, *Appl. Eng.*
584 *Agric.*, 1(2), 96–99, 1985.
- 585 van Ittersum, M. K., Cassman, K. G., Grassini, P., Wolf, J., Tittonell, P. and Hochman, Z.: Yield gap
586 analysis with local to global relevance a review, *F. Crop. Res.*, 143, 4–17, 2013.
- 587 Karlen, D. L., Mausbach, M. J., Doran, J. W., Cline, R. G., Harris, R. F. and Schuman, G. E.: Soil
588 quality: a concept, definition, and framework for evaluation (a guest editorial), *Soil Sci. Soc. Am. J.*,
589 61(1), 4–10, 1997.
- 590 Keesstra, S. D., Bouma, J., Wallinga, J., Tittonell, P., Smith, P., Cerdà, A., Montanarella, L., Quinton,
591 J. N., Pachepsky, Y., van der Putten, W. H., Bardgett, R. D., Moolenaar, S., Mol, G., Jansen, B. and
592 Fresco, L. O.: The significance of soils and soil science towards realization of the United Nations
593 Sustainable Development Goals, *Soil*, 2(2), 111–128, doi:10.5194/soil-2-111-2016, 2016.
- 594 Kemper, W. D. and Chepil, W. S.: Size distribution of aggregates. p. 499--509. CA Black (ed.) *Methods*
595 *of soil analysis. Part I. Agron. Monogr. 9. ASA and SSSA, Madison, WI., Size Distrib. aggregates. p.*
596 *499--509. CA Black Methods soil Anal. Part I. Agron. Monogr. No. 9. ASA, SSSA, Madison, WI.,*
597 *1965.*
- 598 Kroes, J. G., Van Dam, J. C., Groenendijk, P., Hendriks, R. F. A. and Jacobs, C. M. J.: SWAP version
599 3.2. Theory description and user manual, Alterra Rep., 1649, 2008.
- 600 Van Looy, K., Bouma, J., Herbst, M., Koestel, J., Minasny, B., Mishra, U., Montzka, C., Nemes, A.,
601 Pachepsky, Y., Padarian, J. and others: Pedotransfer functions in Earth system science: challenges and
602 perspectives, *Rev. Geophys.*, 2017.
- 603 Lowery, B.: A Portable Constant-rate Cone Penetrometer 1, *Soil Sci. Soc. Am. J.*, 50(2), 412–414,
604 1986.
- 605 Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L. T., Lamarque, J. F.,
606 Matsumoto, K., Montzka, S. A., Raper, S. C. B., Riahi, K. and others: The RCP greenhouse gas
607 concentrations and their extensions from 1765 to 2300, *Clim. Change*, 109(1–2), 213, 2011.
- 608 Moebius-Clune, B. N., Moebius-Clune, D. J., Gugino, B. K., Idowu, O. J., Schindelbeck, R. R., Ristow,
609 A. J. and others: Comprehensive assessment of soil health: The Cornell Framework Manual, Edition
610 3.1, Cornell Univ., Ithaca, NY, 2016.
- 611 Van Oort, P. A. J., Saito, K., Dieng, I., Grassini, P., Cassman, K. G. and Van Ittersum, M. K.: Can
612 yield gap analysis be used to inform R&D prioritisation?, *Glob. Food Sec.*, 12, 109–118, 2017.
- 613 Pulleman, M. M., Bouma, J., Van Essen, E. A. and Meijles, E. W.: Soil organic matter content as a
614 function of different land use history, *Soil Sci. Soc. Am. J.*, 64(2), 689–693, 2000.
- 615 Regione_Campania: I Suoli della Piana in Destra Sele. Progetto carta dei Suoli della Regione Campania
616 in scala 1:50.000 e lotto CP1 e Piana destra Sele (Salerno)., 1996.
- 617 Ritchie, J. T.: Model for predicting evaporation from a row crop with incomplete cover, *Water Resour.*
618 *Res.*, 8(5), 1204–1213, 1972.
- 619 Rockel, B., Will, A. and Hense, A.: The regional climate model COSMO-CLM (CCLM), Meteorol.

- 620 Zeitschrift, 17(4), 347–348, 2008.
- 621 Rossiter, D. G. and Bouma, J.: A new look at soil phenoforms--Definition, identification, mapping,
622 Geoderma, 314, 113–121, 2018.
- 623 Scoccimarro, E., Gualdi, S., Bellucci, A., Sanna, A., Fogli, P. G., Manzini, E., Vichi, M., Oddo, P. and
624 Navarra, A.: Effects of Tropical Cyclones on Ocean Heat Transport in a High-Resolution Coupled
625 General Circulation Model, *J. Clim.*, 24(16), 4368–4384, doi:Doi 10.1175/2011jcli4104.1, 2011.
- 626 Shaw, B. T., Haise, H. R. and Farnsworth, R. B.: Four Years' Experience with a Soil Penetrometer 1,
627 *Soil Sci. Soc. Am. J.*, 7(C), 48–55, 1943.
- 628 Soil Survey Staff: Keys to soil taxonomy, 1999.
- 629 Soil Survey Staff: Keys to soil taxonomy, *Soil Conserv. Serv.*, 12, 410, doi:10.1109/TIP.2005.854494,
630 2014.
- 631 Sonneveld, M. P. W., Bouma, J. and Veldkamp, A.: Refining soil survey information for a Dutch soil
632 series using land use history, *Soil Use Manag.*, 18(3), 157–163, 2002.
- 633 Wade, J., Culman, S. W., Hurisso, T. T., Miller, R. O., Baker, L. and Horwath, W. R.: Sources of
634 variability that compromise mineralizable carbon as a soil health indicator, *Soil Sci. Soc. Am. J.*, 82(1),
635 243–252, 2018.
- 636 Wösten, J. H. ., Lilly, A., Nemes, A. and Le Bas, C.: Development and use of a database of hydraulic
637 properties of European soils, *Geoderma*, 90(3–4), 169–185, doi:10.1016/S0016-7061(98)00132-3,
638 1999.
- 639 Zollo, A. L., Turco, M. and Mercogliano, P.: Assessment of hybrid downscaling techniques for
640 precipitation over the Po river basin, in *Engineering Geology for Society and Territory-Volume 1*, pp.
641 193–197, Springer., 2015.
- 642
- 643

644
645
646
647
648
649
650
651

Tab. 1. Main soil features of selected soil series.

Env. Systems	SMU	STU	Soil family	Soil description		Texture			Hydrological properties				
				Horiz.	Depth (m)	sand (g 100g ⁻¹)	silty	clay	Θ _s (m ³ m ⁻³)	K ₀ (cm d ⁻¹)	α (l cm ⁻¹)	l	n
Hills/foothills	LON0	Longobarda	Pachic Haploxerolls, fine loamy, mixed, thermic	Ap	0-0.5	33.0	40.6	26.4	0.46	27	0.04	-3.44	1.15
				Bw	0.5-1.5	21.7	48.9	29.4	0.61	69	0.02	-1.79	1.18
Alluvial fans	CIF0/ RAGO	Cifariello	Typic Haploxerepts, coarse loamy, mixed, thermic	Ap	0-0.6	33.0	49.5	17.5	0.42	18	0.03	-2.52	1.21
				Bw1	0.6-0.95	33.2	50.2	16.6	0.47	37	0.03	-2.14	1.20
				Bw2	0.95-1.6	29.8	52.2	18.0	0.50	49	0.03	-2.02	1.20
Fluvial Terraces	GIU0	Giuliarossa	Mollic Haploxeralf, fine, mixed, thermic	Ap	0-0.4	27.1	31.9	41.0	0.47	39	0.04	-3.72	1.13
				Bw	0.4-0.85	19.8	28.9	51.3	0.49	7	0.02	-1.28	1.10
				Bss	0.85-1.6	46.3	28.8	24.9	0.40	18	0.05	-2.75	1.16
	SVI0	San Vito	Typic Haploxererts fine, mixed, thermic	Ap	0-0.5	17.3	39.4	43.3	0.44	31	0.03	-3.58	1.15
				Bw	0.5-0.9	16.1	39.6	44.3	0.49	11	0.02	-3.35	1.09
				Bk	0.9-1.3	11.2	40.5	48.3	0.49	10	0.02	-2.52	1.10
	LAZ0	Lazzaretto	Typic Xeropsammets, mixed, thermic	Ap	0-0.45	75.3	12.8	11.9	0.38	77	0.07	-2.26	1.30
C				0.45- >0.65	100.0	0.0	0.0	0.34	123	0.08	2.04	1.85	
Dunes	PET0/ PICO	Picciola	Typic Haploxerepts, coarse loamy, mixed, thermic	Ap	0-0.6	33.3	34.7	32.0	0.48	36	0.04	-3.60	1.13
				Bw	0.6-0.95	30.5	41.2	28.3	0.44	18	0.03	-3.61	1.13
				2Bw	0.95-1.35	28.6	50.0	21.4	0.42	21	0.03	-2.77	1.17

652

653
654
655
656
657
658
659
660
661
662
663
664
665
666
667


668
669
670
671
672
673
674
675

Tab. 2. Main performance indexes of SWAP application in the three soils (Udic Calciustert, Fluventic Haplustept and Typic Calciustoll) under maize cultivation (data from "Nitrati Campania" regional project, Regione Campania, 2008).

Soil	RMSE*	Pearson's R*	n° of soil depths meas.	number of data
Udic Calciustert	0.043 (\pm 0.03)	0.716 (\pm 0.11)	7	1964
Typic Calciustoll	0.044 (\pm 0.03)	0.72 (\pm 0.13)	6	190
Fluventic Haplustept	0.031 (\pm 0.02)	0.821 (\pm 0.09)	6	318

* (average value \pm standard deviation)


676
677
678
679
680
681
682
683
684
685
686
687
688
689
690



691
692

Figure 1: The four landform classes of the “Destra Sele” area and the Soil Map Units (SMU) of selected six Soil Typological Units (STUs, which are similar to the USDA soil series) (CIF0/RAG0= Cifariello; GIU0= Giuliarossa; LAZ0= Lazzaretto; LON0= Longobarda; PET0/PIC0= Picciola; SVI= San Vito).


693
694



695
696
697
698
699
700
701


Figure 2: Simulated Y_w values for all soil series, considering the reference climate (RC; 1971-2005) and future climate scenarios (RCP 8.5) expressed for three time periods (2010-2040; 2040-2070; 2070-2100). The Y_p (potential yield) is the maize production for the Destre Sele area assuming optimal irrigation and fertilization and no pests and diseases. Y_p is only calculated for the reference climate.

702
703




704
705
706
707
708

Figure 3: Cumulated rainfall during the maize growing season (April–August) in the four climate periods.




709
710
711
712
713

Figure 4: The projected effects of simulated soil compaction on Y_w for all the soil series assuming the presence of a compacted plowlayer at 30 cm depth. Other terms are explained in Figure 2.




714
 715
 716 Figure 5: The projected effects of simulated surface runoff of water on Yw for all the soil series. Runoff
 717 occurs when rainfall intensity is higher than the assumed infoltrative capacity of the soil. Other terms
 718 are explained in Figure 2.

719
 720
 721
 722



723
 724 Figure 6: The projected effects of erosion on Yw for all the soil series. Other terms are explained in
 725 Figure 2.

726
 727
 728
 729



730
 731
 732
 733
 734
 735
 736
 737
 738
 739
 740
 741
 742
 743

Figure 7: Range of soil physical quality indexes (Y_a/Y_w) x 100) for all the soil series, expressing the effects of different forms of soil degradation and climate change. The vertical bars for each type of soil (the Genofom) represent a “Thermometer” indicating a characteristic range of values obtained by establishing a series of Phenofoms, represented by their Y_w values. Soil Quality for a given soil is thus represented by a characteristic range of values. Soil Health is indicated by the particular location of an actual Y_a within this range.