

The authors have duly considered and addressed the various queries from the editor and reviewers and provided adequate replies, in so far as possible. For example, evaluation statistics in Table 4 are with respect to profiles from WoSIS, quite a number of which were considered in the compilation of the IGBP, HWSD and GSDE products. Although this is duly mentioned in the text, this aspect should also be indicated in a footnote to Table 4, as tables are often considered without reference to the text.

Reply:

Thanks for the editor and reviewers for providing queries and comment for this manuscript. This helped a lot in improving it. We hope this review will provide some value to the soil data development and Earth system modeling. We added a footnote to Table 4 as suggested.

Modification:

We added a sentence in the text: though quite a number of WoSIS soil profiles were considered in the compilation of these datasets which means that this evaluation is not independent validation.

A footnote to Table 4 was added: Quite a number of WoSIS soil profiles were considered in the compilation of the four products.

Overall, this is a useful and timely review. Unfortunately, the manuscript is still not very well written. There are still various flaws in the English used; these should be corrected by a professional editor.

Reply: We have taken a language service for English correction by two professional editors. Modifications can be seen in the marked manuscript.

Minor remarks (line numbers according to annotated manuscript)

line 69: usually single point observations at a given moment in time, instead of 'usually time-independent'.

Reply: modified.

line 135-142: Check numbering/naming of sections against text.

Reply: we checked this and modified the text.

Modification: In section 2, we first introduce soil datasets produced by the linkage method and digital soil mapping technology at global and national scales, and then, we introduce the soil datasets that have already been incorporated into ESMs, and we also present PTFs that are used in ESMs to estimate soil hydraulic and thermal parameters. In section 3, several global soil datasets are compared and evaluated with a global soil profile database. In section 4, two issues regarding the model use of soil data are described and existing challenges related to the application of soil datasets in ESMs are discussed. In Section 5, a summary and the outlook of further improvements are provided.

line 158, 321-322: remove bold case

Reply: modified.

line 484: the proportion of coarse fragments is also considered in HWSD and WISE30sec.

Reply: Here we made a small mistake. Only IGBP does not contain coarse fragments.

Modification: Notably, only the IGBP does not contain coarse fragments.

line 1628-1630, 1714: rephrase for clarity. PTFs are empirical, predictive functions of certain soil properties (e.g. hydraulic conductivity) from more easily obtained soil properties (e.g. sand, silt, clay and organic carbon content).

Reply: modified.

Modification: PTFs are the empirical, predictive functions that account for the relationships between certain soil properties (e.g., hydraulic conductivity) and more easily obtainable soil properties (e.g. sand, silt, clay and organic carbon content).

line 1931-1933: the soil map is the base map here, not a co-variate.

Reply: modified. However, we argue that from the perspective of 'scorpan' framework soil map can be considered as the major covariate (sometimes the only one) for the linkage method.

Modification: the linkage method usually considers the soil map to be a base map

line 1967: ... countries, most of these are not yet freely available to the international community.

Reply: modified.

Modification: countries, although most of these are not yet freely available to the international community.

line 1969: Should refer to NCSS as the data provider here:  
<https://ncsslabsdatamart.sc.egov.usda.gov/>

Reply: modified.

line 1337: Fifth column, GSDE. 0 is given as depth of bottom layer. Delete this.

Reply: deleted.

# A review on the global soil property maps for Earth system models

Yongjiu Dai<sup>1\*</sup>, Wei Shangguan<sup>1\*</sup>, Dagang Wang<sup>2</sup>, Nan Wei<sup>1</sup>, Qinchuan Xin<sup>2</sup>, Hua Yuan<sup>1</sup>, Shupeng Zhang<sup>1</sup>, Shaofeng Liu<sup>1</sup>, Xingji Lu<sup>1</sup>, Fapeng Yan<sup>3</sup>

<sup>1</sup> Guangdong Province Key Laboratory for Climate Change and Natural Disaster Studies, School of Atmospheric Sciences, Sun Yat-sen University, Guangzhou, China.

<sup>2</sup>School of Geography and Planning, Sun Yat-sen University, Guangzhou, China.

<sup>3</sup>College of Global Change and Earth System Science, Beijing Normal University, Beijing, China

Correspondence to: Yongjiu Dai ([daiyj6@mail.sysu.edu.cn](mailto:daiyj6@mail.sysu.edu.cn)) and Wei Shangguan ([shgwei@mail.sysu.edu.cn](mailto:shgwei@mail.sysu.edu.cn))

**Abstract.** Soil is an important regulator of Earth system processes, but remains one of the least well-described data layers in Earth System Models (ESMs). We reviewed global soil property maps from the perspective of ESMs, including soil physical and, chemical and biological properties, which can also offer insights to soil data developers. These soil datasets provide model inputs, initial variables and benchmark datasets. For modelling use, the dataset should be geographically continuous, scalable and have with uncertainty estimates. The popular soil datasets used in ESMs are often based on limited soil profiles and coarse resolution soil type maps with various uncertainty sources. Updated and comprehensive soil information needs to be incorporated in ESMs. New generation soil datasets derived throughby digital soil mapping with abundant, harmonized and quality controlled soil observations and environmental covariates are preferred to those derived throughby the linkage method (i.e., taxotransfer rule-based method) for ESMs. Soilgrids SoilGrids has the highest accuracy and resolution among the global soil datasets at the time, while other recently developed datasets offerare useful compensationeompliments. Because there is no universal pedotransfer function, an ensemble of them may be more suitable to provide derived soil properties to ESMs. Aggregation and upscaling of soil data are needed for model use but can be avoided ed by taking-using a subgrid method in ESMs at the expenseeost of increases in model complexity. Producing soil property maps in a time series remainsis still challenging. The un Uncertainty iesy ofin soil data needs to be estimated and incorporated into ESMs.

## 1 Introduction

Soil or ~~the~~ pedosphere is a key component of ~~the~~ Earth system, and plays an important role in ~~the~~ water, energy and carbon balances and other biogeochemical processes. An accurate description of soil properties is essential in ~~advancing the modeling/modelling~~ capability~~yies~~ of Earth System Models (ESMs) to predict land surface processes at the global and regional scales (Luo et al., 2016). Soil information is required by ~~the~~ land surface models (LSMs), which ~~are~~is a component of ESMs. With the ~~aid/help~~ of computer-based geographic systems, many researchers have produced geographical databases to organize and harmonize large amounts of soil information generated from soil surveys during ~~recent/the last~~ decades (Batjes, 2017; Hengl et al., 2017). However, soil datasets used in ESMs ~~are~~is not ~~yet~~ well updated ~~nor~~ well utilized ~~yet~~ (Sanchez et al., 2009; FAO/IIASA/ISRIC/ISS-CAS/JRC, 2012). The popular soil datasets used in ESMs are outdated and ~~have~~with limited accuracy~~iesy~~. Some soil properties, such as gravel (or coarse fragment) and depth to bedrock, are not utilized in most ESMs. ~~Meanwhile, it is needed to change The~~ ESMs' schemes and structures ~~must be changed~~ to represent soil processes ~~in a~~ more realistic ~~manner when~~ utilizing new soil information (Brunke et al., 2016; Luo et al., 2016; Oleson et al., 2010). For example, Brunke et al., (2016) incorporated the depth to bedrock data in a land surface model using variable soil layers ~~and~~ instead of the previous constant depth. Better soil information with ~~a~~ high resolution and better representation of soil in models ~~has~~have improved and will improve the performance ~~of~~in simulating the Earth system (eg., Livneh et al., 2015; Dy and Fung, 2016; Kearney and Maino, 2018).

ESMs require detailed information on ~~the the soil~~ physical, chemical and biological properties ~~of the soil~~. Site observations (called soil profiles) from soil surveys include soil properties such as soil depth, soil texture (sand, silt and clay fractions), organic matter, coarse fragments, bulk density, soil colour, soil nutrients (carbon (C), nitrogen (N), phosphorus (P), potassium (K) and sulph~~ph~~ur (S)), amount of roots ~~and so on, etc~~. The range of soil data collected during a soil survey, varies with scale, ~~specifications of a country or a regional specifications~~, and projected applications of the data (i.e., type of soil surveys, routine versus specifically designed surveys). As a result, the availability of soil properties differs in different soil databases. However, soil hydraulic and thermal parameters as well as biogeochemical parameters are usually not observed in soil surveys, which need to be estimated by pedotransfer functions (PTFs) (Looy et al., 2017). This review focuses ~~on the~~ soil data (usually ~~single point observations at a given moment in timetime-invariant~~) from soil surveys, while variables such as soil temperature and soil moisture are beyond ~~the this scope of this paper's scope~~.

Soil properties ~~are~~functioned in three aspects in ESMs:

- 1) Model inputs to estimate parameters. The soil thermal (soil heat capacity and ~~the~~ thermal conductivity) and hydraulic characteristics (empirical parameters of ~~the~~ soil water retention curve and hydraulic conductivity) are usually obtained by fitting equations (PTFs) to easily measured and widely available soil properties, such as sand, silt and clay fractions, organic matter content, rock fragments and bulk density

(Clapp and Hornberger, 1978; Farouki, 1981; Vereecken et al., 2010; Dai et al., 2013). Soil albedos are significantly correlated with ~~the~~ Munsell soil ~~coloureolor~~ value (Post et al., 2000). For some ESMs, the ~~parameters~~ derived ~~parameters~~-by PTFs are used as direct input instead of ~~being ealeulating-calculatedthem~~ in the models.

2) Initial variables. The nutrient (C, N, P, K, S ~~and so on,-ete.~~) amounts and the nutrients associated parameters (pH, cation-exchange capacity, etc.) in soils can be used to initialize the simulations. Generally, their initial values are assumed to be at steady state by running ~~the~~model over thousands of model years (i.e., spin-up) until ~~there is no change~~ trend ~~of change~~ in pool sizes (McGuire et al., 1997; Thornton and Rosenbloom, 2005; Doney et al., 2006; Luo et al., 2016). To initialize nutrient amounts using soil data derived from observations as background ~~field-fields~~ could largely reduce the times of model spin-up, and ~~also~~-could avoid the possibility of ~~athe~~ non-linear singularity evolution of the ~~model,modeling~~ which means ~~that~~ that ~~the~~ models may have multiple equilibria, and then provide ~~a~~ better estimate of the true terrestrial nutrient state. The ~~setting-of~~ initial nutrient stocks ~~settings are is-a~~ major factors leading to model-to-model variation in ~~the~~ simulation (Todd-Brown et al., 2014).

3) Benchmark data. Soil data, as measurements, could serve as a reference for ~~modeling~~ calibration, validation and comparison. Soil carbon stock is one of the ~~sol~~ ~~properties that is~~ most frequently used ~~soil-properties~~ as benchmark data (Todd-Brown et al., 2013). Other nutrient stocks, such as nitrogen stock, can also be used as benchmark data if an ESM simulated ~~these propertiesthem~~.

Soil properties ~~haveare-of~~ great spatial heterogeneity both horizontally and vertically. As a result, ESMs usually incorporate soil property maps (i.e., horizontal spatial distribution) for multiply layers rather than a global constant or a single layer. ESMs, especially LSMs, are evolving towards hyper-resolutions of 1 km or finer with more detailed parameterization schemes to accommodate the land surface heterogeneity (Singh et al., 2015; Ji et al., 2017). ~~Therefore,So~~ spatially explicit soil data at high resolutions are necessary to improve land surface representations and simulations. Because soil properties are observed at individual locations, soil mapping or spatial prediction models ~~are-is~~ needed to derive ~~athe~~ 3D representation of ~~the~~ soil distribution. The traditional ~~methodway~~ (i.e., the linkage method, also called ~~the~~ taxotransfer rule-based method) ~~involvesis-to~~ linking soil profiles and soil mapping units on soil type maps, sometimes with ancillary maps such as topography and land use (Batjes, 2003; FAO/IIASA/ISRIC/ISS-CAS/JRC, 2012). In ~~recentthe past~~ decades, various digital soil mapping technologies ~~have beenwere~~ proposed by finding the relationships between soil and environmental covariates (usually remote sensing data), such as climate, topography, land use, geology and so on (McBratney et al., 2003).

There are many challenges related to ~~the~~ application of soil datasets in ESMs. First, soil datasets are usually not appropriately ~~lyd~~ scaled or formatted for the use of ESMs and some upscaling issues, which ~~areis~~ the most frequently encountered, need to be addressed. The soil datasets produced by the linkage methods are polygon-based and need to be converted to fit the grid-based ESMs. This conversion can be

performed by either the subgrid method or spatial aggregation. The up-to-date soil data are provided at a resolution of 1 km or finer, while the LSMs are mostly ran at a coarser resolution. Therefore, soil data upscaling is necessary before it can be used by ESMs. Proper upscaling methods need to be chosen carefully to minimize the uncertainty introduced by these methods in the modelling results introduced by them (Hoffmann and Christian Biernath, 2016; Kuhnert et al., 2017). Second, all the current global soil datasets represent the average state of the last decades, and the production of soil property maps in a time series is still challenging. Soil landscape and pedogenic models are developed to simulate soil formation processes and soil property changes, which can be incorporated into ESMs. The prediction of changing soil properties can also be performed by digital soil mapping using the changing climate and land use as covariates. Third, the uncertainty in the soil properties can be estimated, and adaptive surrogate modelling based on statistical regression and machine learning may be used to assess the uncertainty effects of the uncertainty of soil properties on ESMs (Gong et al., 2015; Li et al. 2018). Last but not the least, the layer schemes of soil data sets need to be converted for model use, and missing values for deeper soil layers need to be filled.

This paper is organized in the following sections. In section 2, we first introduce soil datasets at global and national scales produced by the linkage method and digital soil mapping technology at global and national scales, and then, we introduce the soil datasets that have already been incorporated into ESMs. Section 3 and we also presents PTFs that are used in ESMs to estimate soil hydraulic and thermal parameters. In section 3, several global soil datasets are compared and evaluated with a global soil profile database. In section 4, two issues regarding the model use of soil data are described and existing challenges related to the application of soil datasets in ESMs are discussed. Section 4 describes how to deal with soil data derived by the linkage methods. Section 5 introduces the upscaling of high-resolution soil data to the coarse resolution of ESMs. In Section 5, 6 gives the a summary and an the outlook of further improvements are provided.

## 2 General methodology of deriving soil datasets for ESMs

### 2.1 Global and national soil datasets

Two kinds of soil data are generated from soil surveys: a-maps (usually in the form of polygon maps) representing the main soil types in a landscape units and soil profiles with observations of soil properties measurements which are considered to be representative of the main component soils of the respective mapping units. ESMs usually require the spatial distribution of soil properties (i.e., soil property maps) rather than information about soil types. Two kinds of methods, i.e., the linkage method and the digital soil mapping method, are used to derive the soil property maps.

Soil maps (the term soil map refers to soil type map in this paper) show the geographical distribution of soil types, which are compiled under a certain soil



classification system. There are many soil mapping units (SMUs) in a soil map and ~~ana~~ SMU is composed of more than one component (i.e. soil type) in most cases. At the global level, there is only one generally accepted global soil map, i.e., the FAO-UNESCO Soil Map of the World (SMW) (FAO, 1971-1981). ~~It~~The SMW was made based on soil surveys conducted between the 1930s and ~~the~~ 1970s, and technology ~~that was~~ available in ~~the~~ 1960s. Several versions exist in the digital format (FAO, 1995, 2003b; Zöbler, 1986) and these products are known to be outdated. The information on the initial SMW and DSMW has since been updated for large sections of the world in the ~~Harmonized World Soil Database (HWSD)~~HWSD product (FAO/IIASA/ISRIC/ISS-CAS/JRC, 2012), which has recently been revised in WISE30sec (Batjes, 2016).–

At the regional and national ~~levels~~level, there are many soil maps based on either national or international soil classifications. ~~Here are–~~ ~~Somesome~~ examples of major soil maps available in digital formats ~~are as follows~~: the Soil and Terrain Database (SOTER) databases (Van Engelen and Dijkshoorn, 2012) for different regions, the European Soil Database (ESB, 2004), the 1: 1 million Soil Map of China (National Soil Survey Office, 1995), the U.S. General Soil Map (GSM), the 1:1 million Soil Map of Canada (Soil Landscapes of Canada Working Group, 2010) and the Australian Soil Resource Information System (ASRIS) (Johnston et al., 2003).–

Soil profiles are composed of multiple layers called soil horizons. For each horizon, soil properties are observed (e.g., site data) or measured (e.g., pH, sand, silt, ~~and~~ clay content). At the global level, several soil profile databases exist. Here, we ~~only~~ discuss ~~only~~ the two most comprehensive ~~ones~~databases. The World Inventory of Soil Emission Potentials (WISE) database was developed as a homogenized set of soil profiles (Batjes, 2008). The newest version (WISE 3.1) contains 10,253 soil profiles and 26 physical and chemical properties. The soil profiles database of ~~the~~ World Soil Information Service (WoSIS) contains the most abundant profiles (about 118,400) from national and global databases including most of the databases mentioned below (Batjes et al., 2017), ~~although~~ only a selection of important soil properties (12) are included (Ribeiro et al., 2018). Data ~~from~~served through WoSIS have been standardized, with special attention ~~to for~~ the description and comparability of soil analytical methods worldwide. However, many countries, although having a large collection of soil profile data, are not yet sharing such data (Arrouays et al., 2017).–

At the regional and national ~~levels~~, there are many soil profile databases, usually with soil classifications corresponding to the local soil maps. ~~and~~ ~~H~~here are some examples: the USA National Cooperative Soil Survey Soil Characterization database (<http://ncsslabsdatamart.sc.egov.usda.gov/>), profiles from the USA National Soil Information System (<http://soils.usda.gov/technical/nasis/>), Africa Soil Profiles database (Leenaars, 2012), the ~~ASRIS~~Australian Soil Resource Information System (Karssies, 2011), the Chinese National Soil Profile database (Shangguan et al., 2013), soil profile archive from the Canadian Soil Information System (MacDonald and Valentine, 1992), soil profiles from SOTER (Van Engelen and Dijkshoorn, 2012), the soil profile analytical database for Europe (Hannam et al., 2009), the Mexico soil profile database ( Instituto Nacional de Estadística y Geografía, 2016), and the

212 Brazilian national soil profile database (Cooper et al., 2005).

213 The linkage method (called the taxotransfer rule-based method) ~~is to involve~~  
214 linking soil maps (with ~~SMUs soil mapping units~~ or soil polygons) and soil profiles  
215 (with soil properties) according to taxonomy-based pedotransfer (taxotransfer in short,  
216 note that ~~here~~, pedotransfer here does ~~not~~ mean ~~pedotransfer functions~~ PTFs, which  
217 ~~are~~ is a different thing) rules (Batjes, 2003). The criteria used in the linkage could be  
218 one or many factors, ~~such~~ as following: soil class, soil texture class, depth zone,  
219 topographic class, distance between soil polygons and soil profiles ~~and so on~~  
220 (Shangguan et al., 2012). Each soil type is represented by one or a group of soil  
221 profiles that meet the criteria, and usually, the median or mean value of a soil property  
222 is assigned to the soil type. Because the linkage method assigned only one value or a  
223 statistical distribution to a soil type in ~~the~~ soil polygons (usually a polygon contains  
224 multiple soil types with their fractions), the intra-polygonal spatial variation is not  
225 ~~considered taken into account~~. At the global level, many databases were derived by the  
226 linkage method: the FAO ~~SMW Soil Map of the World~~ with derived soil properties  
227 (FAO, 2003a), the Data and Information System of International Geosphere-  
228 Biosphere Programme (IGBP-DIS) database (Global Soil Data Task, 2000), the Soil  
229 and Terrain Database (Van Engelen and Dijkshoorn, 2012) for multiply regions and  
230 countries, the ISRIC-WISE derived soil property maps (Batjes, 2006), the  
231 ~~Harmonized World Soil Database (HWSD)~~ (FAO/IIASA/ISRIC/ISS-CAS/JRC,  
232 2012), the Global Soil Dataset for Earth System Model (GSDE) (Shangguan et al.,  
233 2014) and WISE30sec (Batjes, 2016). ~~The T~~ three most recent ~~databases ones~~ are  
234 HWSD, GSDE and WISE30sec. HWSD was built ~~by via~~ combining the existing  
235 regional and national ~~soil information~~ updates ~~of soil information~~. GSDE, as an  
236 improvement of HWSD, incorporated more soil maps and more soil profiles related to  
237 the soil maps, with more soil properties. GSDE accomplished the linkage based on the  
238 local soil classification, which required no correlation between classification systems  
239 and avoided the error brought by ~~the~~ taxonomy reference. In addition, GSDE provides  
240 ~~an~~ estimation of eight layers to ~~at the~~ depth of 2.3 m, while HWSD provides ~~an~~  
241 estimation of two layers to the depth of 1 m. WISE30sec is another improvement of  
242 HWSD ~~that~~ incorporates ~~sd~~ more soil profiles with seven layers up to 200 cm depth and  
243 with uncertainty estimated by ~~the~~ mean  $\pm$  standard deviation. WISE30sec used the soil  
244 map from HWSD with minor corrections and climate zone maps as categorical  
245 covariates. Many national and regional agencies around the world have organized  
246 their soil surveys by linking soil maps and soil profiles, including the USA State Soil  
247 Geographic Database (STATSGO2) (Soil Survey Staff, 2017), Soil Landscapes of  
248 Canada (Soil Landscapes of Canada Working Group, 2010), the ASRIS (Johnston et  
249 al., 2003), the Soil-Geographic Database of Russia (Shoba et al., 2008), the European  
250 Soil Database (ESB, 2004), ~~and~~ the China dataset of soil properties (Shangguan et al.,  
251 2013) ~~and so on~~.

252 Digital soil mapping (McBratney et al., 2003) is the creation and ~~the~~ population  
253 of a geographically referenced soil database, generated at a given resolution by using  
254 field and laboratory observation methods coupled with environmental data through  
255 quantitative relationships (<http://digitalsoilmapping.org/>). Usually, the soil datasets



derived by digital soil mapping provide grid-based spatially continuous estimation while the soil datasets derived by the linkage method provide estimations with abrupt changes at the boundaries of soil polygons. The GlobalSoilMap is a global consortium that aims to create global digital maps for key soil properties (Sanchez et al., 2009). This global effort takes a bottom-up framework and will produce the best available soil map of soil at a resolution of 3 arc sec (about 100 m) along with the 90% confidence of the predictions. Soil properties will be provided for six soil layers (i.e., 0–5, 5–15, 15–30, 30–60, 60–100, and 100–200 cm). Many countries have produced soil maps following the GlobalSoilMap specifications (Odgers et al., 2012; Viscarra Rossel et al., 2015; Mulder et al., 2016; Ballabio et al., 2016; Ramcharan et al., 2018; Arrouays, 2018). The SoilGrids system (<https://www.soilgrids.org>) is another global soil mapping project (Hengl et al., 2014; Hengl et al., 2015; Hengl et al., 2017). The newest version (Hengl et al., 2017) at a resolution of 250 m was produced by fitting an ensemble of machine learning methods based on about 150,000 soil profiles and 158 soil covariates, which is currently the most detailed estimation of global soil distribution. A third global soil mapping project is the Global SOC (soil organic carbon) Map of the Global Soil Partnership, which focuses on country-specific soil organic carbon estimates (Guevara et al., 2018).

Because soil property maps are derived products that are derived based on soil measurements of soil profiles (point observations) and spatial continuous covariates (including soil maps), it is necessary to discuss the sources of uncertainty sources, spatial uncertainty estimation and accuracy assessment of these derived data (the last two are different aspects of uncertainty estimation). More attention should be given to this issue in ESM applications instead of taking soil property maps as observations without error. There are various uncertainty sources in the derivation of deriving soil property maps, including uncertainty from soil maps, soil measurements, soil-related covariates and the linkage method itself (Shangguan et al., 2012; Batjes, 2016; Stoorvogel et al., 2017). The following uncertainties may not be the are not a complete list of uncertainties, but the major uncertainties are listed below. The uncertainties in soil maps are a major source of global datasets derived by the linkage methods. For these datasets, large sections of the world are incorporated into the coarse FAO SMW map, and the purity of soil maps (referring to the following website for the definition: [https://esdac.jrc.ec.europa.eu/ESDB\\_Archive/ESDBv2/esdb/sgdbe/metadata/purity\\_maps/purity.htm](https://esdac.jrc.ec.europa.eu/ESDB_Archive/ESDBv2/esdb/sgdbe/metadata/purity_maps/purity.htm)) is likely to be around 50 to 65% (Landon, 1991). Another important source of uncertainty is the limited comparability of different analytical methods for a given soil property when using soil profiles coming from various sources. A weak correlation or even a negative correlation was found between different analytical methods, although a strong positive correlation was revealed in most cases (McLellan et al. 2013). Both datasets of the linkage method and those by digital soil mapping are subject to suffer this uncertainty. Although there are no straightforward mechanisms to harmonize the data, efforts have been undertaken to address this issue and provide quality assessment (Batjes, 2017; Pillar 5 Working Group, 2017).

Another source of uncertainty comes from the geographic and taxonomic distribution of soil profiles, especially for the under-represented areas and soils (Batjes, 2016). The fourth source of uncertainty is from the linkage method itself. The linkage method does not represent the intra-polygon spatial variation and usually does not explicitly consider soil-related covariates explicitly like digital soil mapping, although there are cases where climate and topography are considered; and Stoorvogel et al. (2017) proposed a methodology to incorporate landscape properties in the linkage method. Finally, uncertainty from the covariates is minor because spatial prediction models such as machine learning in digital soil mapping can reduce its influences (Hengl et al., 2014), although a more comprehensive list of covariates with higher resolution and accuracy will improve the predicted soil property maps. Spatial uncertainty is estimated by different methods for the linkage method and digital soil mapping methods. For the linkage method, statistics such as standard deviation and percentiles can be used for the spatial uncertainty estimation, while these statistics are calculated for the population of soil profiles linked to a soil type or a land unit (Batjes, 2016). This estimation has some limitations because soil profiles are not taken probabilistically but based on their availability, especially for the global soil datasets. Uncertainty will be underestimated when the sample size is not large enough to represent a soil type. For digital soil mapping, spatial uncertainty could be estimated by methods such as geostatistical methods and quantile regression forest (Vaysse and Lagacherie, 2017), which make sense of the statistics. The accuracy of the soil datasets derived by digital soil mapping is estimated by independent validation or cross-validation. But it However, this estimation is not trivial for those data derived by the linkage method due to the global scale, the support of the data and independent data (Stoorvogel et al., 2017), and most of these maps are validated by statistics such as the mean error and coefficient of determination. Instead, some datasets, including WISE and GSDE, use some indicators such as the linkage level of soil class and sample size to offer quality control information (Shangguan et al. 2014; Batjes, 2016). A simple way to compare the accuracy of using datasets with both methods may be to use a global soil profile database as a validation dataset, though some quite a number of these profiles were used when deriving these datasets and questions will be raised. We evaluated several global soil property maps in section 3.—

## 2.2 Soil dataset incorporated in ESMs

Table 1 shows ESMs (specifically, their LSMs land surface models) and their input soil datasets. The ESMs in Table 1 cover the list of CMIP5 (Coupled Model Intercomparison Project) list except those without information about the input soil datasets inputs. Land surface models (LSMs) are key tools to predict the dynamics of land surfaces under climate change and land use. Five datasets are widely used, i.e., the datasets by Wilson and Henderson-Sellers (1985), Zöbner (1986), Webb et al. (1993), Reynolds et al. (2000), Global Soil Data Task (2000), and Miller and White (1998). Except for GSDE, HWSD and STATSGO (Miller and White, 1998) for the USA in Table 1, these datasets were derived from the SMW Soil Map of the World

(note that large sections of GSDE and HWSD still used this map as a base map because there are no available regional or national maps) (FAO, 1971-1981) and limited soil profile data (no more than 5,800 profiles), which gained popularity ~~becausebecause of~~ its simplicity and ease of use. ~~However,But~~ these ~~datasets~~ are outdated and should no longer be used because much better soil information, as introduced in Section 2.1, can be incorporated (Sanchez et al., 2009; FAO/IIASA/ISRIC/ISS-CAS/JRC, 2012).

In recent years, efforts ~~have been made~~~~were taken~~ to improve the soil data condition in ESMs. The Land-Atmosphere Interaction Research Group at Beijing Normal University (BNU, now at Sun Yat-sen University) has put much efforts ~~into~~~~on~~ this topic. Shangguan et al. (2012, 2013) developed a China ~~dataset of~~ soil property~~ies~~ ~~dataset~~ for land surface ~~modeling~~~~modelling~~ based on 8,979 soil profiles and the Soil Map of China using the linkage method. Dai et al. (2013) derived soil hydraulic parameters using ~~PTF~~~~pedotransfer functions~~ based on the soil properties by Shangguan et al. (2013). Shangguan et al. (2014) further developed a comprehensive global dataset for ESMs. The above soil datasets were widely used in the ESMs. Soil properties from these soil datasets, including soil texture fraction, organic carbon, bulk density and derived soil hydraulic parameters, were implemented in the Common Land Model Version 2014 (CoLM2014, <http://land.sysu.edu.cn/>). Li et al. (2017) show~~eds~~ that CoLM2014 was more stable than the previous version and had comparable performance to that of CLM4.5, which may be ~~partially~~ attributed ~~to in-~~~~part to~~ the new soil parameters ~~being used~~ as input. Wu et al. (2014) show~~eds~~ that soil moisture values are closer to the observations when simulated by CLM3.5 with the China dataset than those simulated with FAO. Zheng and Yang (2016) estimated ~~the~~ effects of soil texture datasets from FAO and BNU ~~based~~ on regional terrestrial water cycle simulations with the Noah-MP land surface model. Tian et al. (2012) used the China soil texture data in a land surface model (GWSiB) coupled with a groundwater model. Lei et al. (2014) used the China soil texture data in CLM to estimate the impacts of climate change and vegetation dynamics on runoff in the mountainous region of the Haihe River basin. Zhou et al. (2015) estimated age-dependent forest carbon ~~sink-sinks~~ with a terrestrial ecosystem model utilizing ~~the~~China soil carbon data ~~of China~~. Dy and Fung (2016) updated the soil data for the Weather Research and Forecasting model (WRF).—

Researchers have also put efforts ~~into~~~~to~~ ~~updating~~~~update~~ ESMs with other soil data. Lawrence and Chase (2007) used MODIS data to derive soil reflectance, which was used as a soil colour parameter in ~~the~~ Community Land Model 3.0 (CLM). De Lannoy et al. (2014) updated the ~~NASA~~ Catchment land surface model ~~of the NASA~~ with soil texture and organic matter data from HWSD and STATSGO2. Livneh et al. (2015) evaluated the influence of soil textural properties on hydrologic fluxes by comparing the FAO data and STATSGO2. Folberth et al. (2016) evaluated the impact of soil input data on yield estimates in a ~~globally~~~~global~~ gridded crop model. Slevin et al. (2017) utilized the HWSD to simulate global gross primary productivity in the JULES land surface model. Trinh et al. (2018) proposed an approach that can assimilate coarse global soil data by finer land use and coverage ~~datasets, dataset~~

which improved the performance of hydrologic ~~modeling~~modelling at the watershed scale. Kearney and Maino (2018) incorporated the new generation of soil data produced by the digital soil mapping method into a climate model and found that, compared to the old soil information, ~~this improved the simulation of soil moisture the soil moisture simulation was improved at a~~ at fine spatial and temporal resolution over Australia. A dataset of globally A-global gridded hydrologic soil groups (HYSOGs250m) ~~werewas~~ developed based on soil texture and depth to bedrock of ~~Soilgrids~~SoilGrids (Hengl et al., 2017) and groundwater table depth (Fan et al., 2013) for curve-number based runoff ~~modeling~~modelling of the U.S. Department of Agriculture (Ross et al., 2018).

Except for soil properties, the estimation of underground boundaries, including the groundwater table depth, the depth to bedrock (DTB) and depth to regolith and its implementation in ESMs is also a new focus. Fan et al. (2013) compiled global observations of water table depth and inferred the global patterns using a groundwater model. Pelletier et al. (2016) developed a global DTB dataset ~~by~~ using process-based models for upland and an empirical model for lowland. This dataset was implemented in ~~the~~ CLM4.5, and ~~found that~~ there were significant influences on the water and energy simulations compared to the default constant depth (Brunke et al., 2015). Shangguan et al. (2017~~8~~) developed a global DTB by digital soil mapping based on about 1.7 million observations from soil profiles and water wells, which has a much higher accuracy than the dataset by Pelletier et al. (2016). Vrettas and Fung (2016) ~~showed~~shows that ~~the~~ weathered bedrock stores a significant fraction (more than 30%) of the total water despite its low porosity. Jordan et al. (2018) estimated the global permeability of the unconsolidated and consolidated earth for groundwater modelling. However, due to the lack of data, an accurate global estimation of depth to regolith is not feasible. Caution should be ~~paid to use of the products of~~ used when employing the so-called soil depth products in ESMs so-called soil depth in ESMs. Soil depth maps are usually estimated based on observations from soil surveys, and soil depth (or depth to the R horizon) is assumed to be equal to DTB. However, these observations are usually less than 2 ~~metres~~meters and usually do not reach~~meet~~ the ~~DTB-depth to bedrock~~ (Shangguan et al., 2017). Thus, soil depth maps based on only soil profiles ~~only~~ are significantly underestimated (one order of magnitude lower) compared to the actual ~~DTBdepth to bedrock~~ and should not be taken as the lower boundary of ESMs.–

### 2.3 Estimating secondary parameters using PTFs~~pedotransfer functions~~

Earth system modellers have employed different ~~pedotransfer functions~~ ~~(PTFs)~~PTFs to estimate soil hydraulic parameters (SHP), soil thermal parameters (STP), and biogeochemical parameters (Looy et al., 2017; Dai et al., 2013) or used these parameters as model inputs. ~~Almost~~ Nearly all ESMs incorporated SHPs and STPs estimated by PTFs but not biogeochemical parameters. PTFs are the empirical, predictive functions that account for the relationships between certain soil properties (e.g., hydraulic conductivity) ~~these secondary parameters (i.e., derived soil properties)~~ and more easily obtainable soil properties (e.g. sand, silt, clay and organic carbon content)~~data~~. Direct measurement of these parameters is difficult, expensive and in

most cases impractical ~~to take for obtaining~~ sufficient samples to reflect ~~the~~ spatial variation. Thus, most soil databases do not contain these ~~secondary~~ parameters. PTFs provide ~~the an~~ alternative ~~means of estimating these parametersto estimate them~~. In ESMs, SHPs and STPs are usually derived using simple PTFs, ~~taking using~~ only soil texture data as the input. As more soil properties become ~~globally~~ available ~~globally~~, including gravel, soil organic matter and bulk density, more sophisticated PTFs ~~using~~ ~~that use~~ additional soil properties can be ~~employedutilized~~ in ESMs.—

PTFs can be expressed as either numerical equations or by machine learning methodology which is more flexible ~~to for simulatingsimulate~~ the highly nonlinear relationship in analysed data. PTFs can also be developed based on soil processes. Most researches ~~did have~~ not indicated ~~d~~ where the PTFs can potentially be used, and the accuracy of a PTF outside of its development dataset is essentially unknown (McBratney et al., 2011) ~~McBratney et al. (2011)~~. PTFs ~~are~~ generally ~~are~~ not portable from one region to ~~the otheranother~~ (i.e. locally or regionally validated). Therefore, ~~PTFs they~~ should never be considered as an ultimate source of parameters in soil modelling. Looy et al. (2017) reviewed PTFs extensively in earth system science and emphasized that PTF development ~~has to must~~ go hand in hand with suitable extrapolation and upscaling techniques such that the PTFs correctly represent the spatial heterogeneity of soils in ESMs. ~~Although Though~~ the PTFs were evaluated, it is ~~not clear which are the best set of PTFsunclear which set of PTFs are the best~~ for global applications. Due to these limitations, a better way to estimate these ~~secondary~~ parameters may be to use an ensemble of PTFs, which can ~~providegive~~ the ~~variability~~ ~~of~~ parameters ~~variability~~. Dai et al. (2013) derived a global soil hydraulic parameter database using the ensemble method. Selection of PTFs was carried out based on the following rules, including ~~the a~~ consistent physical definition, ~~large enoughadequately~~ ~~large~~ training sample and positive evaluations ~~in comparisonthat are comparable~~ with other PTFs. The ~~selected~~ PTFs ~~selected included~~ not only ~~included~~ those in equations but also ~~PTFs of~~ machine learning ~~PTFs~~. As a result, the modellers could use these parameters as inputs instead of calculating them in ESMs every time ~~running~~ the model ~~was run~~.

~~The new~~ New generation soil information has already been utilized to derive SHPs and STPs in some ~~studies-researches~~. Montzka et al. (2017) produced a global map of SHPs at a ~~0.25°~~ resolution ~~of 0.25°~~ based on the ~~SoilGridsSoilGrids~~ 1 km dataset. Tóth et al. (2017) calculated SHPs for Europe with ~~the~~ EU-HYDI PTFs (Tóth et al., 2015) based on ~~the SoilGridsSoilGrids~~ 250 m. Wu et al. (2018) used an integrated approach that ensembles PTFs to map ~~the~~ field capacity of China based on multi-source soil datasets.

The ~~PTF~~ performance ~~of PTF~~ in ESMs ~~has beenis~~ evaluated in many ~~researchesstudies~~, ~~althoughthough~~ PTFs ~~havehas~~ not been fully exploited and integrated into ESMs (Looy et al., 2017). ~~Here are s~~Some examples ~~are as follows~~. Chen et al. (2012) incorporated soil organic matter to estimate soil porosity and thermal parameters for ~~the use of land surface modelsin LSMs~~. Zhao et al. (2018a) evaluated PTFs performance to estimate SHPs and STPs for land surface modelling over the Tibetan Plateau. Zheng et al. (2018) developed PTFs to estimate the soil



optical parameters to derive soil albedo for the Tibetan Plateau, and the PTFs ~~that~~ were incorporated into an eco-hydrological model ~~which~~ improved the model simulation of a surface energy budget. Looy et al. (2017) envisaged two possible approaches to improve parameterization of ~~ESMsEarth system models~~ by PTFs. One approach is to replace constant coefficients in ~~the~~ current ESMs ~~that have~~ with spatially distributed values ~~with~~ by PTFs. The other approach is to develop spatially exploitable PTFs to parameterize specific processes using knowledge of environmental controls and variations ~~in~~ of soil properties.

### 3 Comparison of available global soil datasets

For the convenience of ESMs' application, we compared several available soil datasets and evaluated them with soil profiles from WoSIS for some of the key variables (~~Sandsand~~, clay content, organic carbon, coarse fragment and bulk density) used in ESMs. In addition to the most recent developed soil datasets, we also included one old data set (i.e., IGBP) used in ESMs for the evaluation. It is not necessary to compare all the old data sets because they are based on similar, limited and outdated source data as described in section 2.2. ~~They-These datasets~~ have coarser resolutions (Table 1) than the newly developed soil datasets (Table 2).

We present basic descriptions ~~about~~ of the new soil datasets in Table 2 and 3. As described in section 2.1, four available global soil datasets, i.e., HWSD, GSDE, WISE30sec and ~~SoilgridsSoilGrids~~, have been developed in the last several years (Table 2). These soil datasets are selected to be shown here because they have ~~a~~ global coverage with key variables used by ESMs and were developed with relatively good data sources in recent years; ~~and these data~~ are also freely available. Old versions of these datasets are not shown here. Table 3 shows the available soil properties of these soil datasets. Except for WISE30sec, ~~all~~ none of these databases ~~do not~~ contain spatial uncertainty estimations. The explained ~~variance of~~ soil properties variance in ~~SoilgridsSoilGrids~~ is between 56% and 83%, while the other datasets do not offer quantitative accuracy assessments. GSDE has the largest number of soil properties, while ~~SoilgridsSoilGrids~~ currently contains ten primary soil properties defined by the GlobalSoilMap consortium.

The accuracy of the newly developed soil datasets (~~SoilgridsSoilGrids~~, GSDE and HWSD) and an old dataset (IGBP) are evaluated for five key variables using 94,441 soil profiles from WoSIS (Table 4), though quite a number of the WoSIS soil profiles were considered in the complication of these datasets which means that this evaluation is not independent validation. We used four statistics in the evaluation, including mean error (ME), root mean squared error (RMSE), coefficient of variation (CV) and coefficient of determination ( $R^2$ ). All soil datasets are evaluated for topsoil (0-30 cm) and subsoil (30-100 cm). The layer schemes of soil datasets are different (Table 1) and ~~they~~ were converted to the two layers. Soil datasets are ~~in~~ high in resolution and were converted to the resolution of 10 km by averaging. All datasets have relatively small ME. In general, ~~SoilgridsSoilGrids~~ has have much better accuracy than the other three due to RMSE, CV and  $R^2$ , and GSDE ranks the second, followed by IGBP and HWSD. However, IGBP is slightly better than GSDE for bulk density and organic carbon



content of topsoil. ~~Note~~ Notably, that only the IGBP does not contain coarse fragments, which is needed ~~when~~ calculating soil carbon stocks. We did not evaluate the WISE30sec here to save ~~some~~ time in data processing, because previous evaluation using WoSIS showed that WISE30sec had slightly better accuracy than HWSD (<https://github.com/thengl/SoilGrids250m/tree/master/grids/HWSD>). This evaluation has some limitations. First, ~~because~~ the datasets developed by the linkage method, which give the mean value of a SMU, resulted in an abrupt change between the boundaries of soil polygons ~~while~~ whereas the datasets developed by digital soil mapping simulated the soil as a continuum with a spatial continuous change ~~in~~ soil properties; ~~thus, by these datasets~~ may not be ~~so~~ comparable. Second, the original resolutions of soil datasets are different, which means that maps with higher resolutions provides more spatial details, and we should judge the map quality ~~by due to~~ not only the accuracy assessment but also by the resolution. As a result, datasets with higher resolutions (i.e. HWSD, WISE30sec and GSDE) are preferred ~~to those than that~~ with lower resolutions (i.e., IGBP) ~~as because they the higher resolution datasets~~ have similar accuracy, especially when the LSMs are run at a high resolution, such as 1 km. Third, the vertical variation ~~is~~ better represented by ~~Soilgrids~~ SoilGrids, GSDE and WISE30sec with more than 2 layers and ~~to a depth of~~ over 2m (Table 2), ~~which~~ This will provide more useful information for ESMs, especially when they model deeper soils with multiple layers.

The new generation soil dataset produced by the digital soil mapping method gave a quite-very different distribution of soil properties from those produced by the linkage method. Figure 1 shows the soil sand and clay ~~fraction~~ fractions at the surface 0-30 cm layer from ~~Soilgrids~~ SoilGrids, IGBP and GSDE. Figure 2 shows the SOC ~~soil organic carbon~~ and bulk density at the surface 0-30 cm layer from ~~Soilgrids~~ SoilGrids, IGBP and GSDE. Significant differences are visible in these datasets. This difference will lead to different modelling results in ESMs. Tifafi et al. (2018) found that the global ~~soil organic carbon~~ SOC stocks down to a depth of 1 m is 3,400 Pg when estimated by SoilGrids and estimated by Soilgrids while it is 2500 Pg by according to HWSD, and the estimates by ~~Soilgrids~~ SoilGrids are closer to the actual observations, ~~although~~ though they all datasets underestimated the soil carbon stocks. Figure 1 of Tifafi et al. (2018) ~~showed~~ shows the global distribution of soil carbon stocks by ~~Soilgrids~~ SoilGrids and HWSD.

In general, ~~Soilgrids~~ SoilGrids is preferred for ESMs' application ~~as because~~ it currently has the highest accuracy and resolution ~~at the time~~. When soil properties are not available in ~~Soilgrids~~ SoilGrids, WISE30sec and GSDE offers ~~the~~ alternative options. However, model sensitivity simulations need to be ~~performed~~ done to investigate the effects of different soil datasets on ESMs in future studies.

## 4 Soil data usage in ESMs and existing challenges

### 4.1 Model use of soil data derived by the linkage method

Soil data by the linkage method are derived for each ~~soil mapping unit~~ SMU or land unit and thus ~~are~~ is polygon-based, while ESMs are usually grid-based. However, soil data derived by digital soil mapping are grid-based. ~~So~~ Therefore, the

compatibility between soil data derived by the linkage method and ESMs ~~must needs~~  
~~to~~ be addressed. In the soil map, a ~~soil mapping unit~~ (SMU) is composed of more than  
one component soil unit in most cases, and thus, a one-to-many relationship exists  
between the SMU and ~~the~~ profile attributes of the respective soil units. This condition  
makes representing ~~the~~ attributes characterizing ~~ana~~ SMU a non-trivial task. To keep  
the whole ~~soil~~ variation of ~~soil~~ in ~~ana~~ SMU, ~~the best way it is~~ ~~best to use~~ using the  
subgrid method in ESMs (Oleson et al., 2010), i.e. aggregate values of soil properties,  
and provide the area percentage of each value. This will bring ~~about~~ the problem of  
~~how to map mapping~~ the soil subgrids with land cover (or plant function type)  
subgrids. A possible solution is to: classify ~~the~~ soil according to ~~the~~ soil properties  
and ~~get obtain~~ a number of defined soil classes (~~SC~~, n classes) ~~like such as~~ land cover  
types (~~LCT~~, m classes);  
overlay the defined soil classes with land cover types and ~~get~~  
~~obtain~~ n by m combinations assuming ~~the~~ soil classes and land cover types are  
independent. However, this will increase the computing time and ~~the~~ complexity of  
~~the~~ ESMs' structures, which ~~needs require~~ ~~sto~~ implementation the soil processes over  
each subgrid soil column within a grid instead of the entire model grid.

Usually, the compatibility issue is addressed by converting the SMU-based soil  
data to grid data using spatial aggregation. The ESMs uses grid data as input, and  
each grid cell has one unique value of a soil property. Three spatial aggregation  
methods were proposed to aggregate compositional attributes in ~~a-an~~ SMU to a  
representative value (Batjes, 2006; Shangguan et al., 2014). The area-weighting  
method (method A) ~~obtain~~ ~~stakes the~~ area-weighting of soil attributes. The dominant  
type method (method D) ~~takes obtains~~ the soil attribute of the dominant type. The  
dominant binned method (method B) classifies the soil attributes into several  
preselected classes and ~~obtain~~ ~~stakes~~ the dominant class. All three methods can be  
applied to quantitative data, while ~~the~~ method D and ~~the~~ method B can be applied to  
categorical data. The advantages and disadvantages of these methods ~~were have been~~  
discussed (Batjes, 2006; Shangguan et al., 2014). The choice should be made  
according to the specific applications (Hoffmann and Christian Biernath, 2016). ~~The-~~  
~~m~~Method B provides binned classes, which are not convenient for modelling, ~~though-~~  
~~although~~ method B is considered more appropriate to represent a grid cell. ~~The-~~  
~~m~~Method A ~~keeps maintains~~ mass conservation, which can meet most ~~demands of-~~  
model application ~~demandss~~. However, ~~the~~ method A may be misleading in cases  
~~when where~~ extreme values appeared in ~~ana~~ SMU. For the linkage method, the  
uncertainty is usually estimated by ~~obtaining giving~~ the 5 and 95 percentile soil  
properties (or other statistics) of the soil profiles that ~~are~~ linked to ~~ana~~ SMU. Because  
the frequency distribution of ~~the~~ soil properties within a SMU is usually not a normal  
distribution or any other typical ~~statistical statistic~~ distribution, the application of  
statistics such as standard deviation ~~to in~~ model use is not proper. This means that the  
uncertainty ~~in the of~~ soil dataset derived by the linkage method ~~can not cannot~~ be  
incorporated into ESMs in a ~~straightforward~~ ~~straight forward~~ way, and technology  
such as bootstrap may be more suitable than methods ~~that make making~~ assumptions  
on ~~regarding~~ the distribution.

The basic soil properties are often used to derive ~~the~~ secondary parameters,

including SHPs and STPs by PTFs and soil carbon stock or other nutrient stocks by certain equations (Shangguan et al., 2014). This procedure could be ~~done-performed~~ either before or after the aggregation (~~here~~-referred to here as “aggregating after” and “aggregating first”). Because the relationship between the soil basic properties and the derived soil parameters is usually nonlinear, the “aggregating first” method should be ~~used~~taken. This was also ~~proven~~proved by case studies (Romanowicz et al., 2005; Shangguan et al., 2014). However, some ~~researches~~researchers have used the “aggregating after” method ~~to produce~~producing misleading results (Hiederer and Köchy, 2012).

The aggregation smooths the variation ~~in the of~~ soil properties between soil components within a given SMU (Odgers et al., 2012). To avoid ~~the~~ aggregation, the spatial disaggregation of soil type maps can be used to determine the location of the SMU components, ~~though-although~~ the location error may be high in some cases (Thompson et al., 2010; Stoorvogel et al., 2017). This method depends on the high density of soil profiles to establish soil and landscape relationships. Folberth et al. (2016) ~~shows~~showed that the correct spatial allocation of the soil type to the present cropland was very important in global crop yield simulations. Currently, aggregation is still the practical method to use ~~pragmatic-way~~ at the global scale due to lack of data.

#### 4.2 Upscaling detailed soil data for model use

The updated soil datasets derived by both the linkage method and digital soil mapping are usually at a resolution from 1 km to 100 m, and upscaling or aggregation is required to derive lower resolution datasets for model use. The aggregation methods mentioned above can be used. Moreover, there are ~~plenty-of~~many upscaling methods such as the window median, variability-weighted methods (Wang et al., 2004), variogram method (Oz et al., 2002), fractal theory (Quattrochi et al., 2001) and the Miller–Miller scaling approach (Montzka et al., 2017). However, few studies have been devoted to ~~test-out-which~~determining the upscaling methods that are suitable for soil data. A preliminary effort was ~~done-made~~ by (Shangguan, 2014). Five upscaling methods ~~were~~ compared, ~~including-were~~ the window average, ~~widow-window~~ median, ~~widow-window~~ modal, arithmetic average variability-weighted method and bilinear interpolation method. Differences between aggregation methods varied from 10% to 100% for different parameters. The upscaling methods affected the data derived by the linkage method more than the data derived by digital soil mapping. The window average, window median and arithmetic average variability-weighted method performed similar in upscaling. The ~~root-mean-square-error~~RMSE increased rapidly when the window size was less than 40 pixels. Similar to the aggregation of SMUs, the “aggregating first” method is recommended when secondary soil parameters are derived. Again, an alternative to avoid the aggregation into one single value for a grid cell is to use the subgrid methods in ESMs.–

The upscaling effect of soil data on the model simulation has been investigated in previous studies with controversial conclusions. For example, Melton et al. (2017) used two linked algorithms to provide tiles of representative soil textures for subgrids in a

terrestrial ecosystem model and found that the model is relatively insensitive to subgrid soil textures compared to a simple grid-mean soil texture at a global scale. However, the treatment without soil subgrid structure in JULES resulted in soil-moisture dependent anomalies in simulated carbon flux (Park et al., 2018). Further researches are necessary to investigate the upscaling effect on models.

#### 4.3 The changing soil properties

There ~~is are no not any~~ global soil property maps in the time-series because we do not have enough available data. In all ~~the~~ global soil property maps, all ~~the~~ available soil observations in ~~the last decades~~ recent decades have been are used in the development of soil property maps without considering the changing environment. ~~So~~ Therefore, these datasets should be considered as an average state. The critical issue for mapping global soil properties in ~~a time-series~~ time series is to establish a soil profile database with time stamps and then divide them into two or more groups of different periods such as the 1950s-1970s. This is still quite challenging at the global scale because the spatial coverage of soil profiles is quite uneven for different periods and the sample size may not be ~~big enough~~ adequately large to derive maps with ~~satisfied~~ satisfactory accuracy.

Soil properties are changing, but we are now usually considering them to be taking it as static in ESMs. As some ESMs already simulate the soil carbon, this may be considered in PTFs used to estimate soil hydraulic and thermal parameters. Other soil properties affecting soil hydraulic and thermal parameters include soil texture, bulk density, and soil structure ~~and so on~~, but the change is relatively slow. The effect of environmental change on soil properties is the topic of the quantitative ~~modeling~~ modelling of soil forming processes, i.e., soil landscape and pedogenic models (Gessler et al., 1995; Minasny et al., 2008). If we need to simulate the change ~~of in~~ soil properties, a coupling of ESMs and soil landscape and pedogenic models will be needed. Otherwise, we need to predict the soil properties in the future using soil landscape and pedogenic models, which are small scale ~~models and has with~~ high uncertainty. The prediction of changing soil properties may also be ~~done performed~~ by digital soil mapping taken the changing (especially for the future) climate and land use as covariates, which may be easier and ~~the~~ more feasible than dynamic models.

#### 4.4 Incorporating the uncertainty of soil data in ESMs

Incorporating the uncertainty of soil data in ESMs is ~~a rising challenge~~ increasing challenging. Except for WISE30sec, all the current global soil ~~datasets~~ data-sets do not have a corresponding uncertainty map for a soil property. ~~But~~ However, the spatial uncertainty can be estimated by the methods mentioned in section 2.1, and soil ~~datasets~~ data-sets with uncertainty ~~map~~ maps will be made available sooner or later. It is too expensive to run multiply ESM simulations that combine the ~~combining~~ upper and lower bounds in all possible combinations to quantify the effect of soil data uncertainty on ESMs. Instead, adaptive surrogate ~~modeling~~ modelling based on statistical regression and machine learning can be used to emulate the responses of ESMs to the variation of soil properties at each location, which ~~costs much lower~~ uses

~~much less~~ computing time and proves to be effective and efficient (Gong et al., 2015; Li et al. 2018). ~~Surrogate models are used to emulate the responses of ESMs to the variation of soil properties at each location.~~

#### 4.5 Layer schemes and lack of deep layer soil data

The layer scheme of a soil data set needs to be ~~converted~~~~devoted~~ to that of ESMs for model use. A simple ~~methodway~~ for this conversion is the depth weighting method. When a more accurate conversion is needed, the equal-area quadratic smoothing spline functions can be used, which is ~~proved to be~~ advantageous in predicting the depth function of soil properties (Bishop et al., 1999). Mass conservation for a soil property of a layer is guaranteed by this method under the assumption of ~~a~~ continuous vertical variation ~~of in~~ soil properties. This method may produce some negative values ~~thatwhich~~ should be set to zero.

The depth of soil observations in ~~the~~ soil survey ~~is~~~~are~~ usually less than 2 m and thus ~~results~~~~resulted~~ in missing values for the deep layers of ESMs. For the lack of deep soil data, there is ~~nonot any~~ good solution other than ~~extrapolating~~~~extrapolate~~ the values based on the observations of shallower layers, which will lead to higher uncertainty of soil properties for deep layers. The extrapolation can be ~~done~~~~performed~~ by the ~~abovementioned~~~~above-mentioned~~ spline method or simply by assigning ~~the~~ soil properties of the last layer to the rest of ~~the~~ deeper soil layers. ~~The DTB~~~~Depth to bedrock~~ map (Shangguan et al., 2017~~8~~) can be utilized ~~in defining~~~~to define~~ the low boundary of soil layers, and a default set of thermal and hydraulic ~~characteristic~~~~characteristic~~ can be assigned for bedrocks.

#### 5 Summary and outlook

~~In T~~his paper, ~~reviews~~ the status of soil datasets and their usage in ESMs ~~is~~ ~~reviewed~~. Soil physical and chemical properties served as model parameters, initial variables or benchmark datasets in ESMs. Soil profiles, soil maps and soil datasets derived by the linkage method and digital soil mapping are reviewed at national, regional and global levels. The soil datasets derived by digital soil mapping are considered to provide ~~a~~ more realistic estimation of soils than those derived by the linkage method, because digital soil mapping ~~provide~~~~provides~~ ~~spatial~~~~spatially~~ continuous estimations of soil properties using spatial prediction models with various soil-related covariates. Due to the evaluation of soil datasets by WoSIS, ~~Soilgrids~~~~SoilGrids~~ have the most accurate estimation of soil properties. However, other soil datasets, including GSDE and WISE30sec, can be considered as ~~compensation and~~ they provide more soil properties.

The popular soil datasets used in ESMs are outdated and there are ~~updated~~ soil datasets available ~~for the updates~~. In ~~the~~ recent ~~several~~ years, efforts ~~were taken~~~~have been made~~ to update the soil data in ESMs. The effects of updated soil properties which are used to estimate soil hydraulic and thermal parameters, were evaluated. Other major updates include soil reflectance, ground water tables and ~~DTB~~~~depth to bedrock~~.

~~Pedotransfer functions~~ (PTFs) are employed to estimate secondary soil parameters, including soil hydraulic and thermal parameters, and biogeochemical parameters. PTFs



can take more soil properties (i.e., ~~soil-organic-carbon~~SOC, bulk density-~~etc~~ and so on.) as input in addition to soil texture data. An ensemble of PTFs may be more suitable to provide secondary soil parameters as direct input to ESMs, because the ensemble method has a number of benefits and potential over a single PTF (Looy et al., 2017).

Soil data derived by the linkage methods and high-resolution data can be aggregated by different methods to ~~fit-the~~be use in ESMs. The aggregation should be ~~done-performed~~ after the secondary parameters are estimated. However, the aggregation will omit the ~~variation-of~~ soil properties variation. To avoid ~~the~~ aggregation, the subgrid method in ESMs is an alternative ~~that~~which increases the model complexity. The effect of different upscaling methods on the performance of ESMs needs to be further investigated ~~further~~.

Because digital soil mapping has many advantages compared to the traditional linkage method, especially in representing spatial heterogeneity and quantifying uncertainty in the predictions, the new generation soil datasets derived by digital soil mapping need to be tested in ESMs, and some regional studies have shown that these datasets provided ~~d~~ better modelling results than products by the linkage method (Kearney and Maino, 2018; Trinh et al., 2018). Moreover, many studies from digital soil mapping have identified that soil maps are not very important ~~for predicting~~to predict soil properties and are usually not used as a covariate in most studies (eg., Hengl et al., 2014; Viscarra Rossel et al., 2015; Arrouays et al., 2018). However, the linkage method usually ~~takes-considers the~~ soil map as to be a base map~~the major covariate~~, which essentially ~~affect-affects~~ the accuracy of the derived soil property maps, especially for areas without detailed soil maps. As a data-driven method, digital soil mapping requires soil profiles ~~observations-measurements~~ and environmental covariates (in which the importance of soil maps is low), and by including more of these data in mapping will improve the global predictions (Hengl et al., 2017). More quality assessed data, analysed according to comparable analytical methods, are needed to support such efforts. The ~~harmonization-of~~ soil data harmonization ~~is undertaking~~ undertaken by the work of GSP Pillar 5 (Pillar 5 Working Group, 2017) and WoSIS (Batjes et al., 2017). Data derived from proximal sensing, although with higher uncertainty than traditional soil measurements, can be used in soil mapping (England and Viscarra Rossel, 2018). To avoid spatial extrapolation, soil profiles should have ~~a~~ good geographical coverage. The temporal variation ~~of-in~~ global soil is quite challenging due to a lack of data. Soil image fusion is also needed to merge the local and global soil maps, ~~which-and this fusion consider~~consider these maps~~them~~ as ~~components-of~~ soil variation components for ensemble predictions (Hengl et al., 2017). ~~It may take years before Aa~~ system for automated soil image fusion ~~might take years before~~is fully functional in an operational system for global soil data fusion ~~is fully functional~~. Mapping the soil depth and ~~depth-to-bedrock~~DTB separately at the global level ~~is-also~~ remains~~still~~ challenging due to a lack of data and the understanding of relevant processes. Uncertainty estimation, especially spatial uncertainty estimation should be included in the soil datasets developed in the future. However, incorporating the spatial uncertainty of the soil properties in ESMs is still challenging due to the cost, and an alternative may be to use adaptive surrogate ~~modeling~~modelling.



The gap is large between the amount of data that has been ~~taken-obtained~~ in  
 surveys and the amount of data freely available ~~is large~~. The soil profiles included ~~by~~  
~~in~~ global soil databases such as WoSIS ~~make-up~~comprise a very small fraction of the  
 soil pits dug by human beings. For example, there are more than 100,000 soil profiles  
 from the second national soil survey of China (Zhang et al., 2010) and no more than  
 9,000 were used to produce the national soil property maps that are freely available  
 (Shangguan et al., 2013). In the last century, national soil surveys ~~was~~have been  
~~accomplished~~widely accomplished, ~~majorly~~primarily for agriculture purpose.  
 However, most of these legacy data are not digitalized and they are usually not made  
 available to the science community even if digitalized. ~~How to flush out~~Obtaining these  
 hidden soil data will requires some mechanism such as government  
~~mandated~~mandatory regulations and ~~investing~~money investments on making them to  
make these data available (Pillar four Working Group, 2014; Pillar 5 Working Group,  
 2017). Arrouays et al. (2017) reported that about 800,000 soil profiles have been  
~~obtained~~reseued from ~~in~~ the selected countries, although most of these are not yet freely  
available to the international community. In addition, investments ~~in~~on new soil  
 samplings should be made, especially in the under-represented areas. A good example  
 is the USU.S., which has the most abundant soil data freely available  
 (<http://ncsslabsdatamart.sc.egov.usda.gov/>~~Batjes et al., 2017~~) ~~like~~similar to many other  
 data. Censored information produces censored maps and ~~things~~so on. If the hidden data  
 could be made available in any way, science and the whole human being will be  
 promoted. A true big data era is waiting for us. ~~The D~~data compatibility of different  
 analysis methods and different description protocols including soil classifications is  
 also an important issue and data harmonization is necessary when the data are made  
 available to the public.

**Acknowledgements.** This work was supported by the National Key Research and  
 Development Program of China under grants 2017YFA0604303 and  
 2016YFB0200801 and the Natural Science Foundation of China (under grants  
 41575072, 41730962 and U1811464).

## References

- Arora, V.K., Boer, G.J., Christian, J.R., Curry, C.L., Denman, K.L., Zahariev, K., Flato, G.M., Scinocca, J.F., Merryfield, W.J. and Lee, W.G.: The Effect of Terrestrial Photosynthesis Down Regulation on the Twentieth-Century Carbon Budget Simulated with the CCCma Earth System Model, *Journal of Climate* 22(22), 6066-6088, 2009.
- Arrouays, D., Leenaars, J. G. B., Richer-de-Forges, A. C., Adhikari, K., Ballabio, C., Greve, M., Grundy, M., Guerrero, E., Hempel, J., Hengl, T., Heuvelink, G., Batjes, N., Carvalho, E., Hartemink, A., Hewitt, A., Hong, S.-Y., Krasilnikov, P., Lagacherie, P., Lelyk, G., Libohova, Z., Lilly, A., McBratney, A., McKenzie, N., Vasquez, G. M., Mulder, V. L., Minasny, B., Montanarella, L., Odeh, I., Padarian, J., Poggio, L., Roudier, P., Saby, N., Savin, I., Searle, R., Solbovoy, V., Thompson, J., Smith, S., Sulaeman, Y., Vintila, R., Rossel, R. V., Wilson, P., Zhang, G.-L., Swerts, M., Oorts, K., Karklins, A., Feng, L., Ibelle Navarro, A. R., Levin, A., Laktionova, T., Dell'Acqua, M., Suvannang, N., Ruam, W., Prasad, J., Patil, N., Husnjak, S., Pásztor, L., Okx, J., Hallett, S., Keay, C., Farewell, T., Lilja, H., Juilleret, J., Marx, S., Takata, Y., Kazuyuki, Y., Mansuy, N., Panagos, P., Van Liedekerke, M., Skalsky, R., Sobocka, J., Kobza, J., Eftekhari, K., Alavipanah, S. K., Moussadek, R., Badraoui, M., Da Silva, M., Paterson, G., Gonçalves, M. d. C., Theocharopoulos, S., Yemefack, M., Tedou, S., Vrscaj, B., Grob, U., Kozák, J., Boruvka, L., Dobos, E., Taboada, M., Moretti, L., and Rodriguez, D.: Soil legacy data rescue via GlobalSoilMap and other international and national initiatives, *GeoResJ*, 14, 1-19, <https://doi.org/10.1016/j.grj.2017.06.001>, 2017.
- Arrouays, D., Savin, I., Leenaars, J. , McBratney, A.: *GlobalSoilMap - Digital Soil Mapping from Country to Globe*, CRC Press, London, 2018.
- Ballabio, C., Panagos, P., and Montanarella, L.: Mapping topsoil physical properties at European scale using the LUCAS database, *Geoderma*, 261, 110-123, 2016.
- Batjes, N. H.: A taxotransfer rule-based approach for filling gaps in measured soil data in primary SOTER databases, International Soil Reference and Information Centre, Wageningen, 2003.
- Batjes, N. H.: ISRIC-WISE derived soil properties on a 5 by 5 arc-minutes global grid. Report 2006/02, ISRIC- World Soil Information, Wageningen (with data set), 2006.
- Batjes, N. H.: ISRIC-WISE harmonized global soil profile dataset (ver. 3.1). Report 2008/02, ISRIC - World Soil Information, Wageningen, 2008.
- Batjes, N. H.: Harmonized soil property values for broad-scale modelling (WISE30sec) with estimates of global soil carbon stocks, *Geoderma*, 269, 61-68, <https://doi.org/10.1016/j.geoderma.2016.01.034>, 2016.
- Batjes, N. H., Ribeiro, E., van Oostrum, A., Leenaars, J., Hengl, T., Mendes de Jesus, J.: WoSIS: Serving standardised soil profile data for the world, *Earth Syst. Sci. Data*, 9, 1-14, 2017.
- Best, M. J., Pryor, M., Clark, D. B., Rooney, G. G., Essery, R. L. H., Ménard, C. B., Edwards, J. M., Hendry, M. A., Porson, A., Gedney, N., Mercado, L. M., Sitch, S., Blyth, E., Boucher, O., Cox, P. M., Grimmond, C. S. B., and Harding, R. J.: The Joint UK Land Environment Simulator (JULES), model description– Part 1: Energy and water fluxes, *Geosci. Model Dev.*, 4, 677-699, [10.5194/gmd-4-677-2011](https://doi.org/10.5194/gmd-4-677-2011), 2011.
- Bishop, T. F. A., McBratney, A. B., and Laslett, G. M.: Modelling soil attribute depth

858 functions with equal-area quadratic smoothing splines, *Geoderma*, 91, 27–45, 1999.  
 859 Blyth, E. M. a. C.: JULES: A new community land surface mode. *Global Change*  
 860 *Newsletter*, NO. 66, IGBP, Stockholm, Sweden, 9-11, 2006.  
 861 Brunke, M. A., Tucson, A., Broxton, P. D., Pelletier, J., Gochis, D. J., Hazenberg, P.,  
 862 Lawrence, D. M., Niu, G. Y., Troch, P. A., and Zeng, X.: Implementation and testing of  
 863 variable soil depth in the global land surface model CLM4.5, 27th Conference on  
 864 Climate Variability and Change, Phoenix, 2015,  
 865 Brunke, M. A., Broxton, P., Pelletier, J., Gochis, D., Hazenberg, P., Lawrence, D. M.,  
 866 Leung, L. R., Niu, G.-Y., Troch, P. A., and Zeng, X.: Implementing and evaluating  
 867 variable soil thickness in the Community Land Model version 4.5 (CLM4.5), *Journal of*  
 868 *Climate*, 29, 3441–3461, doi:10.1175/JCLI-D-15-0307.1, 2016.  
 869 Chen, F., and Dudhia, J.: Coupling an advanced land surface-hydrology model with  
 870 the Penn State-NCAR MM5 modeling system. Part I: Model implementation and  
 871 sensitivity, *Monthly Weather Review*, 129, 569-585, 2001.  
 872 Chen, Y., Yang, K., Tang, W., Qin, J., and Zhao, L.: Parameterizing soil organic carbon's  
 873 impacts on soil porosity and thermal parameters for Eastern Tibet grasslands, *Science*  
 874 *China Earth Sciences*, 55, 1001-1011, 10.1007/s11430-012-4433-0, 2012.  
 875 Clapp, R. W., and Hornberger, G. M.: Empirical equations for some soil hydraulic  
 876 properties, *Water Resources Res.*, 14, 601-604, 1978.  
 877 Clark, D. B., Mercado, L. M., Sitch, S., Jones, C. D., Gedney, N., Best, M. J., Pryor, M.,  
 878 Rooney, G. G., Essery, R. L. H., Blyth, E., Boucher, O., Harding, R. J., Huntingford, C.,  
 879 and Cox, P. M.: The Joint UK Land Environment Simulator (JULES), model description  
 880 – Part 2: Carbon fluxes and vegetation dynamics, *Geosci. Model Dev.*, 4, 701-722,  
 881 10.5194/gmd-4-701-2011, 2011.  
 882 Cooper, M., Mendes, L. M. S., Silva, W. L. C., and Sparovek, G.: A national soil profile  
 883 database for brazil available to international scientists, *Soil Science Society of*  
 884 *America Journal*, 69, 649–652, 2005.  
 885 Cox, P. M., Betts, R. A., Bunton, C. B., Essery, R. L. H., Rowntree, P. R., and Smith, J.:  
 886 The impact of new land surface physics on the GCM sensitivity of climate and climate  
 887 sensitivity, *Climate Dynamics*, 15, 183-203, 1999.  
 888 Dai, Y., Zeng, X., Dickinson, R. E., Baker, I., Bonan, G. B., Bosilovich, M. G., Denning, A.  
 889 S., Dirmeyer, P. A., Houser, P. R., Niu, G., Oleson, K. W., Schlosser, C. A., and Yang, Z.:  
 890 The Common Land Model, *Bull. Amer. Meteor. Soc.*, 84, 1013-1023, 2003.  
 891 Dai, Y., Shangguan, W., Duan, Q., Liu, B., Fu, S., and Niu, G.: Development of a China  
 892 Dataset of Soil Hydraulic Parameters Using Pedotransfer Functions for Land Surface  
 893 Modeling, *Journal of Hydrometeorology*, 14, 869–887, 2013.  
 894 De Lannoy, G. J. M., Koster, R. D., Reichle, R. H., Mahanama, S. P. P., and Liu, Q.: An  
 895 updated treatment of soil texture and associated hydraulic properties in a global land  
 896 modeling system, *Journal of Advances in Modeling Earth Systems*, 6, 957-979,  
 897 10.1002/2014ms000330, 2014.  
 898 Dickinson, R. E., Henderson-Sellers, A., and Kennedy, P. J.: Biosphere-Atmosphere  
 899 Transfer Scheme (BATS) Version 1e as Coupled to the NCAR Community Climate  
 900 Model. NCAR-TN-387+STR, National Center for Atmospheric Research, Boulder,  
 901 Colorado, 88, 1993.

902 Doney, S. C., Lindsay, K., Fung, I., and John, J.: Natural variability in a stable, 1000-yr  
 903 global coupled climate-carbon cycle simulation, *Journal of Climate*, 19, 3033-3054,  
 904 2006.  
 905 Dy, C. Y., and Fung, J. C. H. C. J.: Updated global soil map for the Weather Research  
 906 and Forecasting model and soil moisture initialization for the Noah land surface  
 907 model, *Journal of Geophysical Research: Atmospheres*, 121, 8777-8800,  
 908 10.1002/2015jd024558, 2016.  
 909 Elguindi, N., Bi, X., Giorgi, F., Nagarajan, B., Pal, J., Solmon, F., Rauscher, S., Zaakey, A.,  
 910 O'Brien, T., Nogherotto, R., and Giuliani, G.: Regional climatic model RegCM  
 911 Reference Manual version 4.6, ITCP, Trieste, 33, 2014.  
 912 England, J. R., and Viscarra Rossel, R. A.: Proximal sensing for soil carbon accounting,  
 913 *SOIL*, 4, 101-122, 10.5194/soil-4-101-2018, 2018.  
 914 Fan, Y., Li, H., and Miguez-Macho, G.: Global Patterns of Groundwater Table Depth,  
 915 *Science*, 339, 940-943, 10.1126/science.1229881, 2013.  
 916 Guevara, M., Olmedo, G. F., Stell, E., Yigini, Y., Aguilar Duarte, Y., Arellano Hernández,  
 917 C., Arévalo, G. E., Arroyo-Cruz, C. E., Bolivar, A., Bunning, S., Bustamante Cañas, N.,  
 918 Cruz-Gaistardo, C. O., Davila, F., Dell Acqua, M., Encina, A., Figueredo Tacona, H.,  
 919 Fontes, F., Hernández Herrera, J. A., Ibelle Navarro, A. R., Loayza, V., Manueles, A.  
 920 M., Mendoza Jara, F., Olivera, C., Osorio Hermosilla, R., Pereira, G., Prieto, P., Ramos,  
 921 I. A., Rey Brina, J. C., Rivera, R., Rodríguez-Rodríguez, J., Roopnarine, R., Rosales  
 922 Ibarra, A., Rosales Riveiro, K. A., Schulz, G. A., Spence, A., Vasques, G. M., Vargas, R.  
 923 R., and Vargas, R.: No silver bullet for digital soil mapping: country-specific soil  
 924 organic carbon estimates across Latin America, *SOIL*, 4, 173-193, 10.5194/soil-4-173-  
 925 2018, 2018.  
 926 FAO: Soil Map of the World, UNESCO, Paris. Vol. 110, 1971-1981.  
 927 FAO: Digitized Soil Map of the World and Derived Soil Properties, FAO, Rome, 1995.  
 928 FAO: Digital soil map of the world and derived soil properties, Food and Agriculture  
 929 Organization of the United Nations, Land and Water Digital Media Series, CD-ROM,  
 930 2003a.  
 931 FAO: The Digitized Soil Map of the World Including Derived Soil Properties (version  
 932 3.6), FAO, Rome, 2003b.  
 933 FAO/IIASA/ISRIC/ISS-CAS/JRC: Harmonized World Soil Database (version1.2), FAO,  
 934 Rome, Italy and IIASA, Laxenburg, Austria, 2012.  
 935 Farouki, O. T.: Thermal Properties of Soils. Monograph, No. 81-1, U.S. Army Cold  
 936 Regions Research and Engineering Laboratory, 1981.  
 937 Folberth, C., Skalský, R., Moltchanova, E., Balkovič, J., Azevedo, L. B., Obersteiner, M.,  
 938 and van der Velde, M.: Uncertainty in soil data can outweigh climate impact signals in  
 939 global crop yield simulations, *Nature Communications*, 7, 11872,  
 940 10.1038/ncomms11872, 2016.  
 941 Gessler, P.E., Moore, I.D., McKenzie, N.J. and Ryan, P.J.: Soil-landscape modelling and  
 942 spatial prediction of soil attributes. *International journal of geographical information*  
 943 *systems*, 9, 421-432, 1995.  
 944 Global Soil DataTask: Global Soil Data Products CD-ROM (IGBP-DIS). International  
 945 Geosphere-Biosphere Programme - Data and Information Services, Available online

at from the ORNL Distributed Active Archive Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee, U.S.A., 2000.

Gong, W., Duan, Q., Li, J., Wang, C., Di, Z., Dai, Y., Ye, A., and Miao, C.: Multi-objective parameter optimization of common land model using adaptive surrogate modeling, *Hydrol. Earth Syst. Sci.*, 19, 2409-2425, 10.5194/hess-19-2409-2015, 2015.

Gurney, K. R., Baker, D., Rayner, P., and Denning, S.: Interannual variations in continental-scale net carbon exchange and sensitivity to observing networks estimated from atmospheric CO<sub>2</sub> inversions for the period 1980 to 2005, *Global Biogeochemical Cycles*, 22, doi:10.1029/2007GB003082, 2008.

Hagemann, S., Botzet, M., Dümenil, L., and Machenhauer, B.: Derivation of global GCM boundary conditions from 1 km land use satellite data. MPI Report No. 289, 34, 1999.

Hagemann, S.: An Improved Land Surface Parameter Dataset for Global and Regional Climate Models. MPI Report No. 336, 28, 2002.

Hannam, J. A., Hollis, J. M., Jones, R. J. A., Bellamy, P. H., Hayes, S. E., Holden, A., Van Liedekerke, M. H., and Montanarella, L.: SPADE-2: The soil profile analytical database for Europe, Version 2.0 Beta Version March 2009, unpublished Report, 27pp, 2009.

Hengl, T., de Jesus, J. M., MacMillan, R. A., Batjes, N. H., Heuvelink, G. B. M., Ribeiro, E., Samuel-Rosa, A., Kempen, B., Leenaars, J. G. B., Walsh, M. G., and Gonzalez, M. R.: SoilGrids1km — Global Soil Information Based on Automated Mapping, *PLoS ONE*, 9, e105992, 10.1371/journal.pone.0105992, 2014.

Hengl, T., Heuvelink, G. B. M., Kempen, B., Leenaars, J. G. B., Walsh, M. G., Shepherd, K. D., Sila, A., MacMillan, R. A., Jesus, J. M. d., Tamene, L., and Tondoh, J. E.: Mapping Soil Properties of Africa at 250 m Resolution: Random Forests Significantly Improve Current Predictions, *PLOS ONE*, 10, e0125814, 2015.

Hengl, T., J., M. d. J., Heuvelink, G. B. M., Gonzalez, R., M., K., M., Blagotic, A., Shangguan, W., Wright, M. N., Geng, X., Bauer-Marschallinger, B., Guevara, M. A., Vargas, R., MacMillan, R. A., Batjes, N. H., Leenaars, J. G. B., Ribeiro, E., Wheeler, I., Mantel, S., and Kempen, B.: SoilGrids250m: global gridded soil information based on Machine Learning, *PLOS One*, 12, 2017.

Hiederer, R., and Köchy, M.: Global Soil Organic Carbon Estimates and the Harmonized World Soil Database, Publications Office of the European Union, Luxembourg, 79, 2012.

Hoffmann, H., G. Zhao, S. Asseng, M. Bindi, and Christian Biernath, J. C., Elsa Coucheney, Rene Dechow, Luca Doro, Henrik Eckersten, Thomas Gaiser, Balázs Grosz, Florian Heinlein, Belay T. Kassie, Kurt-Christian Kersebaum, Christian Klein, Matthias Kuhnert, Elisabet Lewan, Marco Moriondo, Claas Nendel, Eckart Priesack, Helene Raynal, Pier P. Roggero, Reimund P. Rötter, Stefan Siebert, Xenia Specka, Fulu Tao, Edmar Teixeira, Giacomo Trombi, Daniel Wallach, Lutz Weihermüller, Jagadeesh Yeluripati, Frank Ewert: Impact of Spatial Soil and Climate Input Data Aggregation on Regional Yield Simulations, *Plos One*, 11, e0151782, 2016.

Hugelius, G., Tarnocai, C., Broll, G., Canadell, J. G., Kuhry, P., and Swanson, D. K.: The Northern Circumpolar Soil Carbon Database: spatially distributed datasets of soil coverage and soil carbon storage in the northern permafrost regions, *Earth Syst. Sci.*

990 Data, 5, 3-13, 10.5194/essd-5-3-2013, 2013.  
 991 Ji, P., Yuan, X., and Liang, X.-Z.: Do Lateral Flows Matter for the Hyperresolution Land  
 992 Surface Modeling?, *Journal of Geophysical Research: Atmospheres*, 122, 12,077-  
 993 012,092, doi:10.1002/2017JD027366, 2017.  
 994 Johnston, R. M., Barry, S. J., Bleys, E., Bui, E. N., Moran, C. J., Simon, D. A. P., Carlile,  
 995 P., McKenzie, N. J., Henderson, B. L., Chapman, G., Imhoff, M., Maschmedt, D., Howe,  
 996 D., Grose, C., and Schoknecht, N.: ASRIS: the database, *Australian Journal of Soil*  
 997 *Research*, 416, 1021-1036, 2003.  
 998 Instituto Nacional de Estadística y Geografía: Conjunto de Datos de Perfiles de Suelos  
 999 Escala 1: 250 000 Serie II (Continuo Nacional), INEGI, Aguascalientes, Ags. Mexico,  
 1000 2016.  
 1001 Jordan, H., Tom, G., Jens, H., and Janine, B.: Compiling and Mapping Global  
 1002 Permeability of the Unconsolidated and Consolidated Earth: GLobal HYdrogeology  
 1003 MaPS 2.0 (GLHYMPS 2.0), *Geophysical Research Letters*, 45, 1897-1904,  
 1004 doi:10.1002/2017GL075860, 2018.  
 1005 Karssies, L.: CSIRO National Soil Archive and the National Soil Database (NatSoil). No.  
 1006 v1 in Data Collection, CSIRO, Canberra, 2011.  
 1007 Kearney, M. R., and Maino, J. L.: Can next-generation soil data products improve soil  
 1008 moisture modelling at the continental scale? An assessment using a new  
 1009 microclimate package for the R programming environment, *Journal of Hydrology*,  
 1010 561, 662-673, <https://doi.org/10.1016/j.jhydrol.2018.04.040>, 2018.  
 1011 Koster, R. D., and Suarez, M. J.: Modeling the land surface boundary in climate  
 1012 models as a composite of independent vegetation stands, *Journal of Geophysical*  
 1013 *Research: Atmospheres*, 97, 2697-2715, doi:10.1029/91JD01696, 1992.  
 1014 Kowalczyk, E., Stevens, L., Law, R., Dix, M., Wang, Y., Harman, I., Haynes, K.,  
 1015 Srbinovsky, J., Pak, B. and Ziehn, T: The land surface model component of ACCESS:  
 1016 description and impact on the simulated surface climatology, *Australian*  
 1017 *Meteorological and Oceanographic Journal*, 63, 65–82, 2013.  
 1018 Krinner, G., N. Viovy, N. de Noblet-Ducoudré, J. Ogée, J. Polcher, P. Friedlingstein, P.  
 1019 Ciais, S. Sitch, and I. C. Prentice: A dynamic global vegetation model for studies of the  
 1020 coupled atmosphere-biosphere system, *Global Biogeochemical Cycles*, 19, GB1015,  
 1021 2005.  
 1022 Kuhnert, M., Yeluripati, J., Smith, P., Hoffmann, H., van Oijen, M., Constantin, J.,  
 1023 Coucheney, E., Dechow, R., Eckersten, H., Gaiser, T., Grosz, B., Haas, E., Kersebaum, K.-  
 1024 C., Kiese, R., Klatt, S., Lewan, E., Nendel, C., Raynal, H., Sosa, C., Specka, X., Teixeira,  
 1025 E., Wang, E., Weihermüller, L., Zhao, G., Zhao, Z., Ogle, S., and Ewert, F.: Impact  
 1026 analysis of climate data aggregation at different spatial scales on simulated net  
 1027 primary productivity for croplands, *European Journal of Agronomy*, 88, 41-52,  
 1028 <https://doi.org/10.1016/j.eja.2016.06.005>, 2017.  
 1029 Landon, J.R., 1991. *Booker Tropical Soil Manual*. Longman Scientific & Technical,  
 1030 New York.  
 1031 Lawrence, P. J., and Chase, T. N.: Representing a new MODIS consistent land surface  
 1032 in the Community Land Model (CLM 3.0), *Journal of Geophysical Research*, 112,  
 1033 10.1029/2006JG000168, 2007.



1034 Leenaars, J. G. B.: Africa Soil Profiles Database, Version 1.0. A compilation of geo-  
 1035 referenced and standardized legacy soil profile data for Sub Saharan Africa (with  
 1036 dataset). ISRIC report 2012/03, Africa Soil Information Service (AfSIS) project and  
 1037 ISRIC - World Soil Information, Wageningen, the Netherlands, 2012.  
 1038 Lei, H., Yang, D., and Huang, M.: Impacts of climate change and vegetation dynamics  
 1039 on runoff in the mountainous region of the Haihe River basin in the past five  
 1040 decades, *Journal of Hydrology*, 511, 786-799,  
 1041 <http://dx.doi.org/10.1016/j.jhydrol.2014.02.029>, 2014.  
 1042 Li, C., Lu, H., Yang, K., Wright, J. S., Yu, L., Chen, Y., Huang, X., and Xu, S.: Evaluation of  
 1043 the Common Land Model (CoLM) from the Perspective of Water and Energy Budget  
 1044 Simulation: Towards Inclusion in CMIP6, *Atmosphere*, 8, 141, 2017.  
 1045 Li, J., Duan, Q., Wang, Y.-P., Gong, W., Gan, Y., and Wang, C.: Parameter optimization  
 1046 for carbon and water fluxes in two global land surface models based on surrogate  
 1047 modelling, *International Journal of Climatology*, 38, e1016-e1031,  
 1048 [doi:10.1002/joc.5428](https://doi.org/10.1002/joc.5428), 2018.  
 1049 Liang, X., Lettenmaier, D. P., Wood, E. F., and Burges, S. J.: A simple hydrologically  
 1050 based model of land surface water and energy fluxes for general circulation models,  
 1051 *Journal of Geophysical Research: Atmospheres*, 99, 14415-14428,  
 1052 [doi:10.1029/94JD00483](https://doi.org/10.1029/94JD00483), 1994.  
 1053 Livneh, B., Kumar, R., and Samaniego, L.: Influence of soil textural properties on  
 1054 hydrologic fluxes in the Mississippi river basin, *Hydrological Processes*, 29, 4638-  
 1055 4655, [dx.doi.org/10.1002/hyp.10601](https://doi.org/10.1002/hyp.10601), 2015.  
 1056 Looy, K. V., Bouma, J., Herbst, M., Koestel, J., Minasny, B., Mishra, U., Montzka, C.,  
 1057 Nemes, A., Pachepsky, Y. A., Padarian, J., Schaap, M. G., Tóth, B., Verhoef, A.,  
 1058 Vanderborght, J., Ploeg, M. J., Weihermüller, L., Zacharias, S., Zhang, Y., and  
 1059 Vereecken, H.: Pedotransfer Functions in Earth System Science: Challenges and  
 1060 Perspectives, *Reviews of Geophysics*, 55, 1199-1256, [doi:10.1002/2017RG000581](https://doi.org/10.1002/2017RG000581),  
 1061 2017.  
 1062 Luo, Y., Ahlström, A., Allison, S. D., Batjes, N. H., Brovkin, V., Carvalhais, N., Chappell,  
 1063 A., Ciais, P., Davidson, E. A., Finzi, A., Georgiou, K., Guenet, B., Hararuk, O., Harden, J.  
 1064 W., He, Y., Hopkins, F., Jiang, L., Koven, C., Jackson, R. B., Jones, C. D., Lara, M. J.,  
 1065 Liang, J., McGuire, A. D., Parton, W., Peng, C., Randerson, J. T., Salazar, A., Sierra, C.  
 1066 A., Smith, M. J., Tian, H., Todd-Brown, K. E. O., Torn, M., van Groenigen, K. J., Wang, Y.  
 1067 P., West, T. O., Wei, Y., Wieder, W. R., Xia, J., Xu, X., Xu, X., and Zhou, T. C. G. B.:  
 1068 Toward more realistic projections of soil carbon dynamics by Earth system models,  
 1069 *Global Biogeochemical Cycles*, 30, 40-56, [10.1002/2015gb005239](https://doi.org/10.1002/2015gb005239), 2016.  
 1070 MacDonald, K. B., and Valentine, K. W. G.: CanSIS/NSDB. A general description  
 1071 (Centre for Land and Biological Resources Research), Research Branch, Agriculture  
 1072 Canada, Ottawa, 1992.  
 1073 Mauritsen, Thorsten, Jürgen Bader, Tobias Becker, Jörg Behrens, Matthias Bittner,  
 1074 Renate Brokopf, Victor Brovkin, Martin Claussen, Traute Crueger, Monika Esch, Irina  
 1075 Fast, Stephanie Fiedler, Dagmar Fläschner, Veronika Gayler, Marco Giorgetta, Daniel  
 1076 S. Goll, Helmuth Haak, Stefan Hagemann, Christopher Hedemann, Cathy Hohenegger,  
 1077 Tatiana Ilyina, Thomas Jahns, Diego Jimenez de la Cuesta Otero, Johann Jungclaus,

1078 Thomas Kleinen, Silvia Kloster, Daniela Kracher, Stefan Kinne, Deike Kleberg, Gitta  
 1079 Lasslop, Luis Kornblueh, Jochem Marotzke, Daniela Matei, Katharina Meraner, Uwe  
 1080 Mikolajewicz, Kameswarrao Modali, Benjamin Möbis, Wolfgang A. Müller, Julia E. M.  
 1081 S. Nabel, Christine C. W. Nam, Dirk Notz, Sarah-Sylvia Nyawira, Hanna Paulsen,  
 1082 Karsten Peters, Robert Pincus, Holger Pohlmann, Julia Pongratz, Max Popp, Thomas  
 1083 Raddatz, Sebastian Rast, Rene Redler, Christian H. Reick, Tim Rohrschneider, Vera  
 1084 Schemann, Hauke Schmidt, Reiner Schnur, Uwe Schulzweida, Katharina D. Six, Lukas  
 1085 Stein, Irene Stemmler, Bjorn Stevens, Jin-Song von Storch, Fangxing Tian, Aiko Voigt,  
 1086 Philipp de Vrese, Karl-Hermann Wieners, Stiig Wilkenskjeld, Alexander Winkler, and  
 1087 Erich Roeckner: Developments in the MPI-M Earth System Model version 1.2 (MPI-  
 1088 ESM 1.2) and its response to increasing CO<sub>2</sub>, *Journal of Advances in Modeling Earth*  
 1089 *Systems*, 2019.  
 1090 McBratney, A. B., Santos, M. L. M., and Minasny, B.: On digital soil mapping,  
 1091 *Geoderma*, 117, 3-52, 10.1016/s0016-7061(03)00223-4, 2003.  
 1092 McBratney, A. B., Minasny, B., and Tranter, G.: Necessary meta-data for pedotransfer  
 1093 functions, *Geoderma*, 160, 627-629, 2011.  
 1094 McGuire, A. D., Melillo, J. M., Kicklighter, D. W., Pan, Y. D., Xiao, X. M., Helfrich, J.,  
 1095 Moore, B., Vorosmarty, C. J., and Schloss, A. L.: Equilibrium responses of global net  
 1096 primary production and carbon storage to doubled atmospheric carbon dioxide:  
 1097 sensitivity to changes in vegetation nitrogen concentration, *Global Biogeochem.*  
 1098 *Cycles*, 11, 173-189, 1997.  
 1099 McLellan, I., Varela, A., Blahgen, M., Fumi, M. D., Hassen, A., Hechminet, N., Jaouani,  
 1100 A., Khessairi, A., Lyamlouli, K., Ouzari, H.-I., Mazzoleni, V., Novelli, E., Pintus, A.,  
 1101 Rodrigues, C., Ruiu, P. A., Pereira, C. S., and Hursthouse, A.: Harmonisation of physical  
 1102 and chemical methods for soil management in Cork Oak forests - Lessons from  
 1103 collaborative investigations, *African Journal of Environmental Science and*  
 1104 *Technology*, 7, 386-401, 2013.  
 1105 Melton, J. R., Sospedra-Alfonso, R., and McCusker, K. E.: Tiling soil textures for  
 1106 terrestrial ecosystem modelling via clustering analysis: a case study with CLASS-CTEM  
 1107 (version 2.1), *Geosci. Model Dev.*, 10, 2761-2783, 10.5194/gmd-10-2761-2017, 2017.  
 1108 Miller, D. A., and White, R. A.: A conterminous United States multilayer soil  
 1109 characteristics dataset for regional climate and hydrology modeling, *Earth*  
 1110 *Interactions*, 2, 1-26, 10.1175/1087-3562(1998)002<0001:ACUSMS>2.3.CO;2, 1998.  
 1111 Minasny, B., McBratney, A.B. and Salvador-Blanes, S.: Quantitative models for  
 1112 pedogenesis—A review. *Geoderma*, 144, 140-157, 2008.  
 1113 Moigne, P.: SURFEX scientific documentation, Centre National de Recherches  
 1114 Meteorologiques, 2018  
 1115 Montzka, C., Herbst, M., Weihermüller, L., Verhoef, A., and Vereecken, H.: A global  
 1116 data set of soil hydraulic properties and sub-grid variability of soil water retention  
 1117 and hydraulic conductivity curves, *Earth Syst. Sci. Data*, 9, 529-543, 10.5194/essd-9-  
 1118 529-2017, 2017.  
 1119 Mulder, V. L., Lacoste, M., Richer-de-Forges, A. C., and Arrouays, D.: GlobalSoilMap  
 1120 France: High-resolution spatial modelling the soils of France up to two meter depth,  
 1121 *Science of The Total Environment*, 573, 1352-1369,

<http://dx.doi.org/10.1016/j.scitotenv.2016.07.066>, 2016.  
 National Soil Survey Office: Soil Map of China (in Chinese), China Map Press, Beijing, 1995.  
 Niu, G.-Y., Yang, Z.-L., Mitchell, K. E., Chen, F., Ek, M. B., Barlage, M., Kumar, A., Manning, K., Niyogi, D., Rosero, E., Tewari, M., and Xia, Y.: The community Noah land surface model with multiparameterization options (Noah-MP): 1. Model description and evaluation with local-scale measurements, *Journal of Geophysical Research: Atmospheres*, 116, doi:10.1029/2010JD015139, 2011.  
 Odgers, N. P., Libohova, Z., and Thompson, J. A.: Equal-area spline functions applied to a legacy soil database to create weighted-means maps of soil organic carbon at a continental scale, *Geoderma*, 189-190, 153-163, 2012.  
 Oleson, K. W., Lawrence, D. M., B, G., Flanner, M. G., Kluzek, E., J., P., Levis, S., Swenson, S. C., Thornton, E., Feddes, J., Heald, C. L., Lamarque, J.-f., Niu, G.-y., Qian, T., Running, S., Sakaguchi, K., Yang, L., Zeng, X., and Zeng, X.: Technical Description of version 4.0 of the Community Land Model (CLM). NCAR Technical Note NCAR/TN-478+STR, National Center for Atmospheric Research, Boulder, 257, 2010.  
 Oleson, K. W., D.M. Lawrence, G.B. Bonan, B. Drewniak, M. Huang, C.D. Koven, S. Levis, F. Li, W.J. Riley, Z.M. Subin, S.C. Swenson, P.E. Thornton, A. Bozbiyik, R. Fisher, E. Kluzek, J.-F. Lamarque, P.J. Lawrence, L.R. Leung, W. Lipscomb, S. Muszala, D.M. Ricciuto, W. Sacks, Y. Sun, J. Tang, Z.-L. Yang: Technical Description of version 4.5 of the Community Land Model (CLM). Ncar Technical Note NCAR/TN-503+STR, National Center for Atmospheric Research, Boulder, CO, 422, 2013.  
 Orth, R., Dutra, E. and Pappenberger, F.: Improving Weather Predictability by Including Land Surface Model Parameter Uncertainty. *Monthly Weather Review* 144(4), 1551-1569, 2016.  
 Oz, B., V. Deutsch, C., and Frykman, P.: A visualbasic program for histogram and variogram scaling, *Computers & Geosciences*, 28, 21-31, [http://dx.doi.org/10.1016/S0098-3004\(01\)00011-5](http://dx.doi.org/10.1016/S0098-3004(01)00011-5), 2002.  
 Park, J., Kim, H.-S., Lee, S.-J., and Ha, T.: Numerical Evaluation of JULES Surface Tiling Scheme with High-Resolution Atmospheric Forcing and Land Cover Data, *SOLA*, 14, 19-24, 10.2151/sola.2018-004, 2018.  
 Patterson, K. A.: Global distributions of total and total-available soil water-holding capacities, Master, University of Delaware, Newark, DE, 1990.  
 Pelletier, J. D., P. D. Broxton, P. Hazenberg, X. Zeng, P. A. Troch, G.-Y. Niu, Z. Williams, M. A. Brunke, and D. Gochis: A gridded global data set of soil, immobile regolith, and sedimentary deposit thicknesses for regional and global land surface modeling, *Journal of Advances in Modeling Earth Systems*, 8, 10.1002/2015MS000526, 2016.  
 Pillar 5 Working Group: Implementation Plan for Pillar Five of the Global Soil Partnership, FAO, Rome, 2017.  
 Pillar four Working Group: Plan of Action for Pillar Four of the Global Soil Partnership, FAO, Rome, 2014.  
 Post, D. F., Fimbres, A., Matthias, A. D., Sano, E. E., Accioly, L., Batchily, A. K., and Ferreira, L. G.: Predicting Soil Albedo from Soil Color and Spectral Reflectance Data, *Soil Science Society of America Journal* 64, 1027-1034, 2000.

1166 Quattrochi, D. A., Emerson, C. W., Lam, N. S.-N., and Qiu, H.-I.: Fractal  
 1167 Characterization of Multitemporal Remote Sensing Data, in: Modelling Scale in  
 1168 Geographical Information System, edited by: Tate, N., and Atkinson, P., John Wiley &  
 1169 Sons, London, 13-34, 2001.

1170 Ramcharan, A., Hengl, T., Nauman, T., Brungard, C., Waltman, S., Wills, S., and  
 1171 Thompson, J.: Soil Property and Class Maps of the Conterminous United States at  
 1172 100-Meter Spatial Resolution, *Soil Science Society of America Journal*, 82, 186-201,  
 1173 10.2136/sssaj2017.04.0122, 2018.

1174 Ribeiro, E., Batjes, N. H., and Oostrom, A. v.: World Soil Information Service (WoSIS) -  
 1175 Towards the standardization and harmonization of world soil data, ISRIC - World Soil  
 1176 Information, Wageningen, 2018.

1177 Reynolds, C. A., Jackson, T. J., and Rawls, W. J.: Estimating soil water-holding  
 1178 capacities by linking the Food and Agriculture Organization Soil map of the world  
 1179 with global pedon databases and continuous pedotransfer functions, *Water Resour.*  
 1180 *Res.*, 36, 3653-3662, 2000.

1181 Romanowicz, A. A., Vanclooster, M., Rounsevell, M., and Junesse, I. L.: Sensitivity of  
 1182 the SWAT model to the soil and land use data parametrisation: a case study in the  
 1183 Thyle catchment, Belgium, *Ecological Modelling*, 187, 27-39, 2005.

1184 Rosenzweig, C., and Abramopoulos, F.: Land surface model development for the GISS  
 1185 GCM, *J. Climate*, 10, 2040-2054, 1997.

1186 Ross, C. W., Prihodko, L., Anchang, J., Kumar, S., Ji, W., and Hanan, N. P.:  
 1187 HYSOGs250m, global gridded hydrologic soil groups for curve-number-based runoff  
 1188 modeling, *Scientific Data*, 5, 180091, 10.1038/sdata.2018.91, 2018.

1189 Rotstayn, L. D., S. J. Jeffrey, M. A. Collier, S. M. Dravitzki, A. C. Hirst, J. I. Syktus, and K.  
 1190 K. Wong: Aerosol- and greenhouse gas-induced changes in summer rainfall and  
 1191 circulation in the Australasian region: a study using single-forcing climate simulations,  
 1192 *Atmos. Chem. Phys.*, 12, 6377–6404, 2012.

1193 Saha, S., Moorthi, S., Wu, X., Wang, J., Nadiga, S., Tripp, P., Behringer, D., Hou, Y.-T.,  
 1194 Chuang, H.-y., Iredell, M., Ek, M., Meng, J., Yang, R., Mendez, M.P., Dool, H.v.d.,  
 1195 Zhang, Q., Wang, W., Chen, M. and Becker, E.: The NCEP Climate Forecast System  
 1196 Version 2. *Journal of Climate* 27(6), 2185-2208, 2014.

1197 Sanchez, P. A., Ahamed, S., Carré, F., Hartemink, A. E., Hempel, J., Huising, J.,  
 1198 Lagacherie, P., McBratney, A. B., McKenzie, N. J., Mendonça-Santos, M. d. L.,  
 1199 Budiman Minasny, L. M., Okoth, P., Palm, C. A., Sachs, J. D., Shepherd, K. D., Vågen, T.-  
 1200 G., Vanlauwe, B., Walsh, M. G., Winowiecki, L. A., and Zhang, G.-L.: Digital soil map of  
 1201 the world, *Science*, 325, 680-681, 2009.

1202 Sellers, P. J., Randall, D. A., Collatz, G. J., Berry, J. A., Field, C. B., Dazlich, D. A., Zhang,  
 1203 C., Collelo, G. D., and Bounoua, L.: A revised land surface parameterization (SiB2) for  
 1204 atmospheric GCMs. Part I: model formulation, *Journal of Climate*, 9, 676-705, 1996.

1205 Shangguan, W., Dai, Y., Liu, B., Ye, A., and Yuan, H.: A soil particle-size distribution  
 1206 dataset for regional land and climate modelling in China, *Geoderma*, 171-172, 85-91,  
 1207 2012.

1208 Shangguan, W., Dai, Y., Liu, B., Zhu, A., Duan, Q., Wu, L., Ji, D., Ye, A., Yuan, H., Zhang,  
 1209 Q., Chen, D., Chen, M., Chu, J., Dou, Y., Guo, J., Li, H., Li, J., Liang, L., Liang, X., Liu, H.,

1210 Liu, S., Miao, C., and Zhang, Y.: A China dataset of soil properties for land surface  
 1211 modeling, *Journal of Advances in Modeling Earth Systems*, 5, 212-224,  
 1212 10.1002/jame.20026, 2013.  
 1213 Shangguan, W.: Comparison of aggregation ways on soil property maps, 20th World  
 1214 Congress of Soil Science, Jeju, Korea, 2014,  
 1215 Shangguan, W., Dai, Y., Duan, Q., Liu, B., and Yuan, H.: A global soil data set for earth  
 1216 system modeling, *Journal of Advances in Modeling Earth Systems*, 6, 249-263, 2014.  
 1217 Shangguan, W., Hengl, T., Mendes de Jesus, J., Yuan, H., and Dai, Y.: Mapping the  
 1218 global depth to bedrock for land surface modeling, *Journal of Advances in Modeling*  
 1219 *Earth Systems*, 9, 65-88, 10.1002/2016ms000686, 2017<sup>8</sup>.  
 1220 Shoba, S. A., Stolbovoi, V. S., Alyabina, I. O., and Molchanov, E. N.: Soil-geographic  
 1221 database of Russia, *Eurasian Soil Science*, 41, 907-913, 10.1134/s1064229308090019,  
 1222 2008.  
 1223 Singh, R. S., Reager, J. T., Miller, N. L., and Famiglietti, J. S.: Toward hyper-resolution  
 1224 land-surface modeling: The effects of fine-scale topography and soil texture on  
 1225 CLM4.0 simulations over the Southwestern U.S, *Water Resources Research*, 51, 2648-  
 1226 2667, doi:10.1002/2014WR015686, 2015.  
 1227 Slevin, D., Tett, S. F. B., Exbrayat, J. F., Bloom, A. A., and Williams, M.: Global  
 1228 evaluation of gross primary productivity in the JULES land surface model v3.4.1,  
 1229 *Geosci. Model Dev.*, 10, 2651-2670, 10.5194/gmd-10-2651-2017, 2017.  
 1230 Soil Survey Staff, N. R. C. S., United States Department of Agriculture: Web Soil  
 1231 Survey. Available online at <http://websoilsurvey.nrcs.usda.gov/>. Accessed 1/1/2017,  
 1232 2017.  
 1233 Soil Landscapes of Canada Working Group: Soil Landscapes of Canada version 3.2.,  
 1234 Agriculture and Agri-Food Canada, Ottawa, Ontario, 2010.  
 1235 Stoorvogel, J. J., Bakkenes, M., Temme, A. J. A. M., Batjes, N. H., and Brink, B. J. E.: S-  
 1236 World: A Global Soil Map for Environmental Modelling, *Land Degradation &*  
 1237 *Development*, 28, 22-33, doi:10.1002/ldr.2656, 2017.  
 1238 Takata, K., Emori, S., and Watanabe, T.: Development of the minimal advanced  
 1239 treatments of surface interaction and runoff. *Global Planet. Change*, 38, 209–222,  
 1240 2003.  
 1241 Thompson, J. A., Prescott, T., Moore, A. C., Bell, J., Kautz, D. R., Hempel, J. W.,  
 1242 Waltman, S. W., and Perry, C. H.: Regional approach to soil property mapping using  
 1243 legacy data and spatial disaggregation techniques, 19th World Congress of Soil  
 1244 Science, Brisbane, Queensland, 2010,  
 1245 Thornton, P. E., and Rosenbloom, N. A.: Ecosystem model spin-up: estimating steady  
 1246 state conditions in a coupled terrestrial carbon and nitrogen cycle model, *Ecological*  
 1247 *Modelling*, 189, 25-48, 2005.  
 1248 Tian, W., Li, X., Wang, X. S., and Hu, B. X.: Coupling a groundwater model with a land  
 1249 surface model to improve water and energy cycle simulation, *Hydrol. Earth Syst. Sci.*  
 1250 *Discuss.*, 2012, 1163-1205, 10.5194/hessd-9-1163-2012, 2012.  
 1251 Tifafi, M., Guenet, B., and Hatté, C.: Large Differences in Global and Regional Total  
 1252 Soil Carbon Stock Estimates Based on SoilGrids, HWSD, and NCSCD: Intercomparison  
 1253 and Evaluation Based on Field Data From USA, England, Wales, and France, *Global*

1254 Biogeochemical Cycles, 32, 42-56, doi:10.1002/2017GB005678, 2018. Todd-Brown, K.  
 1255 E. O., Randerson, J. T., Post, W. M., Hoffman, F. M., Tarnocai, C., Schuur, E. A. G., and  
 1256 Allison, S. D.: Causes of variation in soil carbon simulations from CMIP5 Earth system  
 1257 models and comparison with observations, Biogeosciences, 10, 1717-1736,  
 1258 10.5194/bg-10-1717-2013, 2013.  
 1259 Todd-Brown, K. E. O., Randerson, J. T., Hopkins, F., Arora, V., Hajima, T., Jones, C.,  
 1260 Shevliakova, E., Tjiputra, J., Volodin, E., Wu, T., Zhang, Q., and Allison, S. D.: Changes  
 1261 in soil organic carbon storage predicted by Earth system models during the 21st  
 1262 century, Biogeosciences, 11, 2341-2356, 10.5194/bg-11-2341-2014, 2014.  
 1263 Tóth, B., Weynants, M., Nemes, A., Makó, A., Bilas, G., and Tóth, G.: New generation  
 1264 of hydraulic pedotransfer functions for Europe, European Journal of Soil Science, 66,  
 1265 226-238, doi:10.1111/ejss.12192, 2015.  
 1266 Tóth, B., Weynants, M., Pásztor, L., and Hengl, T.: 3D soil hydraulic database of Europe  
 1267 at 250 m resolution, Hydrological Processes, 31, 2662-2666, doi:10.1002/hyp.11203,  
 1268 2017.  
 1269 Trinh, T., Kavvas, M. L., Ishida, K., Ercan, A., Chen, Z. Q., Anderson, M. L., Ho, C., and  
 1270 Nguyen, T.: Integrating global land-cover and soil datasets to update saturated  
 1271 hydraulic conductivity parameterization in hydrologic modeling, Science of The Total  
 1272 Environment, 631-632, 279-288, <https://doi.org/10.1016/j.scitotenv.2018.02.267>,  
 1273 2018.  
 1274 Van Engelen, V., and Dijkshoorn, J.: Global and National Soils and Terrain Digital  
 1275 Databases (SOTER), Procedures Manual, version 2.0. ISRIC Report 2012/04, ISRIC -  
 1276 World Soil Information, Wageningen, the Netherlands, 2012.  
 1277 Vaysse, K., and Lagacherie, P.: Using quantile regression forest to estimate  
 1278 uncertainty of digital soil mapping products, Geoderma, 291, 55-64,  
 1279 <https://doi.org/10.1016/j.geoderma.2016.12.017>, 2017.  
 1280 Vereecken, H., Weynants, M., Javaux, M., Pachepsky, Y., Schaap, M. G., and  
 1281 Genuchten, M. T. v.: Using pedotransfer functions to estimate the van Genuchten-  
 1282 Mualem soil hydraulic properties: a review, Vadose Zone Journal, 9, 795-820, 2010.  
 1283 Viscarra Rossel, R., Chen, C., Grundy, M., Searle, R., Clifford, D., and Campbell, P.: The  
 1284 Australian three-dimensional soil grid: Australia's contribution to the GlobalSoilMap  
 1285 project, Soil Research, 53, 845-864, 2015.  
 1286 Versegny, D.: The Canadian land surface scheme (CLASS): Its history and future,  
 1287 Atmosphere-Ocean, 38:1, 1-13, 2000.  
 1288 Vrettas, M. D., and Fung, I. Y.: Toward a new parameterization of hydraulic  
 1289 conductivity in climate models: Simulation of rapid groundwater fluctuations in  
 1290 Northern California, Journal of Advances in Modeling Earth Systems, 7, 2105-2135,  
 1291 10.1002/2015ms000516, 2016.  
 1292 Wang, G., Gertner, G., and Anderson, A. B.: Up-scaling methods based on variability-  
 1293 weighting and simulation for inferring spatial information across scales, International  
 1294 Journal of Remote Sensing, 25, 4961- 4979, 2004.  
 1295 Webb, R. S., Rosenzweig, C. E., and Levine, E. R.: Specifying land surface  
 1296 characteristics in general circulation models: Soil profile data set and derived water-  
 1297 holding capacities, Global Biogeo. Cyc., 7, 97-108, 1993.



1298 Wilson, M. F., and Henderson-Sellers, A.: A global archive of land cover and soils data  
 1299 for use in general circulation climate models, *Journal of Climatology*, 5, 119-143,  
 1300 1985.  
 1301 Wu, L., Wang, A., and Sheng, Y.: Impact of Soil Texture on the Simulation of Land  
 1302 Surface Processes in China, *Climatic and Environmental Research (in Chinese)*, 19,  
 1303 559-571, doi:10.3878/j.issn.1006-9585.2013.13055, 2014.  
 1304 Wu, T., Song, L., Li, W., Wang, Z., Zhang, H., Xin, X., Zhang, Y., Zhang, L., Li, J., Wu, F.,  
 1305 Liu, Y., Zhang, F., Shi, X., Chu, M., Zhang, J., Fang, Y., Wang, F., Lu, Y., Liu, X., Wei, M.,  
 1306 Liu, Q., Zhou, W., Dong, M., Zhao, Q., Ji, J., Li, L. and Zhou, M.: An overview of BCC  
 1307 climate system model development and application for climate change studies.  
 1308 *Journal of Meteorological Research*, 28(1), 34-56, 2014. Wu, X., Lu, G., Wu, Z., He, H.,  
 1309 Zhou, J., and Liu, Z.: An Integration Approach for Mapping Field Capacity of China  
 1310 Based on Multi-Source Soil Datasets, *Water*, 10, 728, 2018.  
 1311 Zhang, W. L., Xu, A. G., Ji, H. J., Zhang, R. L., Lei, Q. L., Zhang, H. Z., Zhao, L. P., and  
 1312 Long, H. Y.: Development of China digital soil map at 1:50,000 scale, 19th World  
 1313 Congress of Soil Science, *Soil Solutions for a Changing World*, Brisbane, Australia,  
 1314 2010,  
 1315 Zhao, H., Zeng, Y., Lv, S., and Su, Z.: Analysis of soil hydraulic and thermal properties  
 1316 for land surface modeling over the Tibetan Plateau, *Earth Syst. Sci. Data*, 10, 1031-  
 1317 1061, 10.5194/essd-10-1031-2018, 2018a.  
 1318 Zhao, M., Golaz, J.-C., Held, I. M., Guo, H., Balaji, V., Benson, R., Chen, J.-H., Chen, X.,  
 1319 Donner, L. J., Dunne, J. P., Dunne, K., Durachta, J., Fan, S.-M., Freidenreich, S. M.,  
 1320 Garner, S. T., Ginoux, P., Harris, L. M., Horowitz, L. W., Krasting, J. P., Langenhorst, A.  
 1321 R., Liang, Z., Lin, P., Lin, S.-J., Malyshev, S. L., Mason, E., Milly, P. C. D., Ming, Y., Naik,  
 1322 V., Paulot, F., Paynter, D., Philipps, P., Radhakrishnan, A., Ramaswamy, V., Robinson,  
 1323 T., Schwarzkopf, D., Seman, C. J., Shevliakova, E., Shen, Z., Shin, H., Silvers, L. G.,  
 1324 Wilson, J. R., Winton, M., Wittenberg, A. T., Wyman, B., and Xiang, B.: The GFDL  
 1325 Global Atmosphere and Land Model AM4.0/LM4.0: 2. Model Description, Sensitivity  
 1326 Studies, and Tuning Strategies, *Journal of Advances in Modeling Earth Systems*, 10,  
 1327 735-769, doi:10.1002/2017MS001209, 2018b.  
 1328 Zheng, G., Yang, H., Lei, H., Yang, D., Wang, T., and Qin, Y.: Development of a  
 1329 Physically Based Soil Albedo Parameterization for the Tibetan Plateau, *Vadose Zone*  
 1330 *Journal*, 17, 10.2136/vzj2017.05.0102, 2018.  
 1331 Zheng, H., and Yang, Z. L.: Effects of soil type datasets on regional terrestrial water  
 1332 cycle simulations under different climatic regimes, *Journal of Geophysical Research:*  
 1333 *Atmospheres*, Accepted, 10.1002/2016jd025187, 2016.  
 1334 Zhou, T., Shi, P. J., Jia, G. S., Dai, Y. J., Zhao, X., Shangguan, W., Du, L., Wu, H., and Luo,  
 1335 Y. Q.: Age-dependent forest carbon sink: Estimation via inverse modeling, *Journal of*  
 1336 *Geophysical Research-Biogeosciences*, 120, 2473-2492, 10.1002/2015jg002943,  
 1337 2015.  
 1338 Zöbner, L.: A world soil file for global climate modeling, NASA Tech. Memo. 87802,  
 1339 NASA, New York, 33, 1986.

1340 Table 1. Lists of the soil dataset used by land surface models (LSM) of Earth System Models (ESM) or climate models (CM).  
1341

Dataset	Resolution	ESM or CM	LSM	Input soil data
Elguindi et al. (2014)		RegCM	BATS1e (Dickinson et al., 1993) or CLM3.5	Soil texture classes and Soil color classes prescribed for BATS vegetation/land cover type
FAO (2003 a,b)	5'	CanESM2	CTEM (Arora et al., 2009) CLASS3.4 (Verseghy, 2000)	Soil texture
FAO (2003 a,b)	5'	EC-EARTH	HTESSEL (Orth et al., 2016)	Soil texture classes
FAO (2003 a,b; outside Conterminous US) STATSGO (Miller and White, 1998)	5' 30"	WRF CWRF	Noah (Chen and Dudhia, 2001) Noah-MP (Niu et al., 2011) CLM4 Other LSMs	Soil texture
GSDE (Shangguan et al., 2014)	30"	CAS_ESM BNU_ESM GRAPES	CoLM 2014(Dai et al., 2003)	Soil texture, gravel, soil organic carbon, bulk density
GSDE (Shangguan et al., 2014)	30"	WRF CWRF	Noah (Chen and Dudhia, 2001) Noah-MP (Niu et al., 2011) CLM4 Other LSMs	Soil texture
GSDE (Shangguan et al., 2014)	30"	BCC_CSM 1.1 BCC_CSM 1.1(m)	BCC_AVIM 1.1 (Wu et al., 2014)	Soil texture
Hagemann (2002)	0.5° (8km over Africa)	MPI-ESM ICON-ESM	JSBACH4 (Mauritsen et al. (2019)	Soil albedo

Hagemann (2002)	0.5°	MPI-ESM ICON-ESM	JSBACH4 (Mauritsen et al. (2019)	Field capacity, Plant-available soil water holding capacity and wilting point prescribed for ecosystem type
Hagemann et al. (1999)	0.5°	MPI-ESM ICON-ESM	JSBACH4 (Mauritsen et al. (2019)	Volumetric heat capacity and thermal diffusivity prescribed for 5 soil types of FAO soil map
HWSD (FAO/IIASA/ISRIC/ISS -CAS/JRC, 2012)	30"	GFDL ESM	GFDL LM4 (Zhao et al., 2018b)	Soil texture classes
HWSD (FAO/IIASA/ISRIC/ISS -CAS/JRC, 2012)	30"	HadCM3 HadGEM2 QUEST	JULES/MOSESvn 5.4 (Best et al., 2011; Clark et al., 2011)	Soil texture
HWSD (FAO/IIASA/ISRIC/ISS -CAS/JRC, 2012)	30"	CNRM- CM5	SURFEX8.1 (Moigne, 2018)	Soil texture, soil organic matter
IGBP-DIS (Global Soil DataTask, 2000)	5'	CESM CCSM CMCC– CESM FIO-ESM FGOALS (s2,gl,g2)	CLM 3.0 or CLM 4.0 or CLM 5.0 (Oleson, 2013)	Soil texture (sand, clay)
		NorESM1		
ISRIC-WISE (Batjes, 2006) combined with NCSD (Hugelius et al., 2013)	5'; 0.25°	CESM CCSM CMCC– CESM FIO-ESM FGOALS (s2,gl,g2) NorESM1	CLM 3.0 or CLM 4.0 or CLM 5.0 (Oleson, 2013)	Soil organic matter

Lawrence and Chase (2007)	0.05°	CESM CCSM CMCC– CESM FIO-ESM FGOALS (s2,gl,g2) NorESM1	CLM 3.0 or CLM 4.0 or CLM 5.0 (Oleson, 2013)	Soil color class
Reynolds et al. (2000)	5'	GLDAS	Mosaic (Koster and Suarez, 1992) CLM2 Noah (Chen and Dudhia, 2001) VIC (Liang et al., 1994)	Soil texture classes
Webb et al. (1993) and Zöbler (1986)	1°	GISS-E2	GISS-LSM (Rosenzweig and Abramopoulos, 1997)	Soil texture
Wilson and Henderson- Sellers (1985)	1°	HadCM3 HadGEM2 QUEST	JULES/MOSESvn 5.4 (Best et al., 2011;Clark et al., 2011)	Soil texture
Zöbler (1986)	1°	ACCESS- ESM	CABLE2.0 (Kowalczyk et al, 2013)	Soil texture classes
Zöbler (1986)	1°		SiB (Sellers et al., 1996; Gurney et al., 2008)	Soil texture classes
Zöbler (1986)	1°	CFSv2	CFSv2/Noah(Saha et al., 2014)	Soil texture
Zöbler (1986)	1°	CSIRO- Mk3.6.0	CSIRO-Mk3.6.0 (Rotstayn et al., 2012)	Soil texture classes
Zöbler (1986)	1°	MIROC (4h,5) MIROC- ESM	MATSIRO (Takata et al., 2003)	Soil texture classes

	Zöbler (1986); Reynolds et al. (2000)	1°; 5'	IPSL-CM6	ORCHIDEE [rev 3977] (Krinner, 2005)	Soil texture classes
1342					
1343	ACCESS = Australia Community Climate and Earth System Simulator				
1344	BATS = Biosphere-Atmosphere Transfer Scheme				
1345	BCC_CSM = Beijing Climate Center Climate System Model				
1346	BCC_AVIM = Beijing Climate Center Atmosphere and Vegetation Interaction Model				
1347	BNU_ESM = Beijing Normal University Earth System Model				
1348	CABLE = Community Atmosphere Biosphere Land Exchange				
1349	CanESM = Canadian Earth System Model				
1350	CAS_ESM = Chinese Academy of Sciences Earth System Model				
1351	CCSM = Community Climate System Model.				
1352	CESM = Community Earth System Model				
1353	CFS = Climate Forecast System				
1354	CLASS = Canadian Land Surface Scheme				
1355	CLM = Community Land Model				
1356	CMCC-CESM = Euro-Mediterranean Centre on Climate Change Community Earth System Model				
1357	CNRM-CM = Centre National de Recherches Meteorologiques Climate Model				
1358	CoLM = Common Land Model				
1359	CSIRO-Mk = Commonwealth Scientific and Industrial Research Organization climate system model				
1360	CTEM = Canadian Terrestrial Ecosystem Model				
1361	EC-EARTH = European community Earth-System Model				
1362	FAO = Food and Agriculture Organization (FAO-UNESCO) digital Soil Map of the World (SMW) at <a href="#">a</a> 1:5 million scale				
1363	FGOALS = Flexible Global Ocean-Atmosphere-Land System Model				
1364	FIO-ESM = First Institute of Oceanography Earth System Model				
1365	GRAPES = Global/Regional Assimilation Prediction System				
1366	GFDL = Geophysical Fluid Dynamics Laboratory				
1367	GISS = Goddard Institute for Space Studies				
1368	GLDAS = Global Land Data Assimilation System				
1369	GSDE = Global Soil Dataset for Earth System Model				
1370	HadCM = Hadley Centre Coupled Model				

1371 HadGEM2-ES = Hadley Global Environment Model 2 - Earth System  
 1372 HTESSEL = Tiled ECMWF Scheme for Surface Exchanges over Land  
 1373 HWSO = Harmonized World Soil Database  
 1374 ICON-ESM = Icosahedral non-hydrostatic Earth System Model  
 1375 IGBP-IDS = Data and Information System of International Geosphere-Biosphere Programme  
 1376 IPSL-CM = ~~Institut~~Institute Pierre Simon Laplace Climate Model  
 1377 ISRIC-WISE = World Inventory of Soil Emission Potentials of International Soil Reference and Information Centre  
 1378 JSBACH = Jena Scheme of Atmosphere Biosphere Coupling in Hamburg  
 1379 JULES/MOSES= Joint UK Land Environment Simulator/Met Office Surface Exchange Scheme  
 1380 MATSIRO = Minimal Advanced Treatments of Surface Interaction and Runoff  
 1381 MIROC = Model for Interdisciplinary Research on Climate  
 1382 MPI-ESM = Max Planck Institute for Meteorology Earth System Model  
 1383 Noah-MP = Noah-multiparameterization  
 1384 NorESM1 = Norwegian Earth System Model  
 1385 NCSD = Northern Circumpolar Soil Carbon Database  
 1386 ORCHIDEE = Organising Carbon and Hydrology In Dynamic Ecosystems  
 1387 QUEST = Quantifying and Understanding the Earth System  
 1388 RegCM = Regional Climate Model  
 1389 SiB = Simple ~~Biosphere~~Biosphere Model  
 1390 STATSGO = State Soil Geographic Database  
 1391 SURFEX = Surface Externalisée  
 1392 WRF = Weather Research and Forecasting Model

Table 2 Four new global soil datasets for ~~the~~ESM updates ~~of~~ ESMs.

Dataset*	Resolution	Number of layers	Number of properties	depth to the bottom of a layer (cm)	Mapping method
HWSD	1km	2	22	30, 100	Linkage method
GSDE	1km	8	39	0, 4.5, 9.1, 16.6, 28.9, 49.3, 82.9, 138.3, 229.6	Linkage method
WISE30sec	1km	7	20	20,40,60,80,100,150,200	Linkage method
<del>Soilgrids</del> <u>SoilGrids</u>	250m	6	7	5, 15, 30, 60, 100, 200	Digital soil mapping

\*HWSD, GSDE, WISE30sec and ~~Soilgrids~~SoilGrids are freely available at <http://www.iiasa.ac.at/web/home/research/researchPrograms/water/HWSD.html>, <http://globalchange.bnu.edu.cn/research/data>, <https://www.isric.org/explore/wise-databases>, and <http://www.soilgrids.org/>, respectively.

Table 3 Derived soil properties considered in ~~f~~Four global soil datasets.

Soil property*	HWSD	GSDE	WISE30sec	<del>Soilgrids</del> <u>SoilGrid</u> <u>s</u>	Soil property*	HWSD	GSDE	WISE30sec	<del>Soilgrids</del> <u>SoilGrid</u> <u>s</u>
Drainage class	√	√	√		Total carbon		√		
AWC class	√	√			Total nitrogen		√	√	
Soil phase	√	√			Total sulfur		√		
Impermeable layer	√	√			pH(KCL)		√		√
Obstacle to roots	√	√			pH(CaCl <sub>2</sub> )		√		
Additional property	√	√			Exchangeable Ca		√		
Soil water regime	√	√			Exchangeable Mg		√		
Reference soil depth	√	√			Exchangeable K		√		
Depth to bedrock				√	Exchangeable Na		√		
Gravel	√	√	√	√	Exchangeable Al		√		
Sand, Silt, Clay	√	√	√	√	Exchangeable H		√		
Texture class**	√				VWC at -10 kPa		√		
Bulk density	√	√	√	√	VWC at -33 kPa		√	√	
Organic Carbon	√	√	√	√	VWC at -1500 kPa		√	√	
pH(H <sub>2</sub> O)	√	√	√	√	Phosphorous by Bray method		√		
CEC (clay)	√		√		Phosphorous by Olsen method		√		
CEC (soil)	√	√	√		Phosphorous by New Zealand method		√		
Effective CEC			√		Water soluble phosphorous		√		



Base saturation	✓	✓	✓	Phosphorous by Mechlich method	✓		
TEB	✓		✓	Total phosphorous	✓		
Calcium Carbonate	✓	✓	✓	Total Potassium	✓		
Gypsum	✓	✓	✓	Salinity (ECE)	✓	✓	✓
Sodicity (ESP)	✓		✓	Aluminium saturation			✓
C/N ratio			✓				

1400 \*CEC is cation exchange capacity. The base saturation measures the sum of exchangeable cations (nutrients) Na, Ca, Mg and K as a  
1401 percentage of the overall exchange capacity of the soil (including the same cations plus H and Al). TEB is the total exchangeable base  
1402 including Na, Ca, Mg and K. ESP is the exchangeable sodium percentage, which is calculated as  $Na * 100 / CEC_{soil}$ . ECE is electrical  
1403 conductivity. AWC is the available water storage capacity. The first 9 soil properties on the left, including the drainage class and,—  
1404 AWC class ~~and so on~~ are available for each soil type, while the other properties are available for each layer. ~~It should be noted—~~  
1405 ~~that~~ Notably, many different analytical methods have been used to derive a given soil property, which is a major source of uncertainty.  
1406 \*\*texture class can be calculated using sand, silt and clay content.

Table 4 Evaluation statistics of soil datasets using ~~WoSIS~~ soil profiles from World Soil Information Service (WoSIS). ~~ME is mean error. RMSE is root mean squared error. CV is coefficient of variation. R<sup>2</sup> is coefficient of determination.~~

Soil property	Dataset	Topsoil (0-30 cm)*				Subsoil (30-100 cm)			
		ME	RMSE	CV	R <sup>2</sup>	ME	RMSE	CV	R <sup>2</sup>
Sand content (% in weight)	<del>Soilgrids</del> SoilGrids	-0.906	18.6	0.457	0.518	-0.27	19.1	0.501	0.492
	GSDE	-0.443	23.2	0.571	0.247	-1.31	23.8	0.625	0.211
	HWSD	6.64	27.4	0.673	0.014	2.08	27.6	0.725	-0.058
	IGBP	3.74	26.3	0.647	0.051	4.06	26.3	0.691	0.055
Clay content (% in weight)	<del>Soilgrids</del> SoilGrids	1.34	12.5	0.554	0.339	0.39	13.6	0.485	0.382
	GSDE	-0.949	14.6	0.643	0.104	-0.79	16.4	0.584	0.105
	HWSD	0.77	16.2	0.718	-0.119	1.42	18.9	0.672	-0.182
	IGBP	3.27	15.4	0.678	0.044	2.44	16.8	0.597	0.084
Bulk density (kg/m <sup>3</sup> )	<del>Soilgrids</del> SoilGrids	-79.7	237	0.164	0.338	-33.5	212	0.136	0.327
	GSDE	-68.4	279	0.193	0.030	-65.5	269	0.173	-0.043
	HWSD	-105	298	0.206	-0.033	-168	317	0.204	-0.107
	IGBP	-55.6	273	0.189	0.050	-112	294	0.189	-0.130
Coarse fragment (% in volume)	<del>Soilgrids</del> SoilGrids	1.53	10.1	1.68	0.319	1.23	12.8	1.47	0.335
	GSDE	3.2	13.5	2.24	-0.165	3.18	16.8	1.93	-0.115
	HWSD	1.8	13.2	2.2	-0.164	-0.40	16.2	1.87	-0.081
Organic carbon (g/kg)	<del>Soilgrids</del> SoilGrids	6.21	29.8	1.69	0.218	0.99	23.5	3.32	0.134
	GSDE	-0.354	34.5	1.95	-0.095	0.45	27.4	3.87	-0.174
	HWSD	-3.67	36.2	2.05	-0.194	-1.38	27.4	3.87	-0.172
	IGBP	0.61	33.4	1.89	-0.026	1.67	28.5	4.02	-0.268

\*Quite a number of WoSIS soil profiles were considered in the compilation of the four products.  
ME is the mean error. RMSE is the root mean squared error. CV is the coefficient of variation. R<sup>2</sup> is the coefficient of determination.




Figure 1 Soil sand and clay fraction at the surface 0-30 cm layer from ~~Soilgrids~~SoilGrids, IGBP-DIS and GSDE. The difference among them will lead to different ~~modeling~~modelling results for ESMs. IGBP-DIS is Data and Information System of International Geosphere-Biosphere Programme, and GSDE is Global Soil Dataset for Earth System Model.




Figure 2 Soil organic carbon and bulk density at the surface 0-30 cm layer from Soilgrids, GSDE and IGBP.