

Interactive comment on “Beneath the arctic greening: Will soils lose or gain carbon or perhaps a little of both?” by Jennifer W. Harden et al.

Jennifer W. Harden et al.

jaodonnell@nps.gov

Received and published: 5 April 2019

Response to Anonymous Referee #2 General Comments

1. Harden et al. used the space for time approach in order to get insights on the critical question about the fate of permafrost soil carbon under climate change. The authors combined physico-chemical fractionation of soil C pools with radio carbon dating and exponential equation fitting with soil depth. The results showed depth distributions of organic C were related mainly to depths of rooting and changes in bulk density. According to the study, thawing of PF will cause changes in specific C pools. The first period until the year 2100 will result in net C loss of unprotected pools, while mineral protected pools will gain C. Further warming beyond 2100 will cause losses

C1

from the mineral protected C pools, while deeper rooting stimulate the gain of light fraction materials. These results are of strong importance for the scientific community. Not only for permafrost research, but also for general information about changes in stabilization mechanisms of soil C under a future climate. The authors did a great job evaluating ^{14}C in different SOM pools, which is crucial in order to understand SOM stabilization mechanisms. The study is written in excellent scientific English and well organized.

We appreciated the positive feedback.

2. I found, however, some drawbacks, which need to be considered further in the review. The authors provide only little information about parent material, except that there is some loess underlain. It would be good to have a map of the sites or some information about the depth of the loess sediments and the material below. Texture and mineralogical composition are crucial parameters for the storing capacity of OC in mineral soil layers. OC contents strongly correlate positively with mineral parameters such as clay, silt, Fe-Al- hydroxides in soil. Already small changes in these parameters have strong impact on the overall OC storage capacity. The space for time approach assumes that these parameters are similar between the sites. Unfortunately, no information on texture are presented. An increase of clay content by only 5% can result, for example, in up to 2% higher OC concentrations in temperate arable soils. If for example, the Inceptisols would have the highest clay content, than the gain in the mineral-associated fraction could be explained by that. Similar, the loss toward Mollisols could be explained by a slightly lower clay or Fe content. I'm afraid that the message could be biased without considering these very important parameters. Incorporating these parameters in mixed effects models, or at least showing that clay is not a principle driver for OC stock change between the sites should solve the problem. Further, the gradient of sampling sited not only reflects a temperature gradient but also a precipitation gradient, from 270 mm in Gelisols to 850mm in Iova. How does the climate scenarios reflect changes in precipitation in the arctic? Precipitation and thus, soil moisture are next to

C2

temperature, the main drivers for OC mineralization. Therefore, it would be good to read how this moisture gradient reflects the model results.

We included more information about particle size and citations to mineralogy AND we entered those data into the online repository for these profiles (International Soil Radiocarbon Database). It is true that the parent materials are not exactly comparable among sites (e.g., clay contents, Fe oxides, etc vary) but their variation is likely far less than that of climate and biotic systems which is the basis for the comparison. As for precipitation, we don't know. We included this "unknown" in the caveats paragraph on page 16.

3. I found no information's about how many profiles or soil samples have been analyses. Also the data in supplement were not very helpful. How many samples have been fractionated?

We have added text to show the number of profiles analyzed in the Methods section. We also note that only one profile of soil samples were fractionated for each soil type.

Specific Comments

1. Specific comments: P2L3: "Fitting an exponential equation to depth trends in soil C: : ." please explain if specific pools are fitted or the bulk soil. The same for the depths of rooting and changes in bulk density. Pools or bulk?

While we fit the exponential equation to both bulk and fractionated soils, this statement specifically refers to depth trends and controls on bulk soil samples. We revised text in the Abstract to reflect this.

2. P3L3-12: the paragraph described that SOC stocks and MRT depend from environmental and substrate-specific factors. In terms of substrates, the authors refer mainly to the quality and quantity of plant residue inputs. One crucial factor for the SOC storing quantity is the parent material or the substrate for soil formation. Clay, silt and Fe-Al-(oxy)hydroxide content effecting the overall storage capacity of SOC (Kleber et

C3

al., 2015; von Lützow et al.,2006). This should be mentioned here, because mineral-organic interactions are part of the manuscript. There are also some latest works on organo-mineral stabilization in permafrost soils (Gentsch et al., 2015, 2018; Mueller et al., 2017).

Excellent point. We added text and some citations recommended by the reviewer to this paragraph.

3. P4L16 following: Please describe how the samples were taken. How was bulk density measured, which is used in calculating the C density?

We added text to this section of the Methods to this section to better describe how soils were sampled and bulk density was determined.

4. P5L12: "dramatic differences" sounds a bit fishy. Please chance the phrase.

We changed the wording here to state "considerable differences".

5. P8L15: I found it pretty hard to understand what Zmin means. Would be nice to have a quick excess explanation

We added a sentence here to better define Zmin and also Cmin.

6. P9L14: please delete relatively before modern. Everything $F > 1$ is per definition modern. So the whole profile LF is modern.

Agreed. We deleted "relatively" from this sentence.

7. P12-13 Fig 4: there is probably a mistake in units by description of the model results from Fig 4. In the text the changes are given in g C m^{-2} which is reasonable for me. Figure 4 reported values in $\text{kg C m}^{-2} \text{y}^{-1}$, which resulted in incredible amounts of C when scaling them up to a larger area or over 200years.

We revised the Results text in this section to better describe fluxes (and not stocks) in Figure 4. Based on these flux estimates, which are constrained by our radiocarbon

C4

measurements, we provide total stock changes over relevant time scales.

8. P13L17: correct Zmin lower case. Also in later sentences.

We corrected the formatting of Zmin throughout the manuscript.

9. Figure 3: please specify how many profiles were involved in the model

We added text to the Figure legend to note that model-data fits are based on fractionated soils for one profile from each soil type.

Interactive comment on SOIL Discuss., <https://doi.org/10.5194/soil-2018-41>, 2019.