

Suggestions for revision or reasons for rejection (will be published if the paper is accepted for final publication)

The manuscript SOIL-2019-15 “Impacts of land use and topography on soil organic carbon in a Mediterranean landscape (north-western Tunisia)” has been improved by the authors after having into account most of the comments made by the referees. In this new version, provided information concerning the study area and methodology and sampling design is much clear and understandable. However, still some aspects should be considered before being publishing. Above **all discussion should be improved, in the present form** is wordy and it is difficult to extract the meaning. **Why Authors did not include a brief discussion about the 3.1 soil spectral library as an integrative indicator of soil quality.** This is one of the novel aspects of the article.

Please **avoid the repetition of the information.** Be careful to keep verb tense consistent within sections of a paper. For example, **the Results section is usually in the past tense** (because the experiments have already been done).

- Please keep the same terminology in the whole manuscript; **choose among topographic characteristics or features (instead of topographic units or factors).** **Similarly SOC levels or contents.**
- The acronyms should be **defined for the first time.** Revise in line 8 soil organic carbon (SOC) and line 51 Soil organic matter (OM); Line 84 SOC instead of soil organic carbon
- Please some concepts should be rewritten to be more accurate Lines 29-30 change people by human **activities ... exacerbates soil erosion**”. Lines 31- 32 **Long-term anthropogenic pressure from agricultural use (Kosmas et al., 2015) together with abiotic factors such as climatic and topographical variability.**
- A deeper English proofreading **can improve the readability of the text.** Please remove to avoid the repetition of information Line 35 **“Soil degradation is a key component of land degradation** in the Mediterranean region” a similar sentence was included in line 28. In addition, revise to avoid the repetition of the same word e.g. “soil quality **deterioration contributes to the deterioration**”. See also lines 89-90
- It is **not clear if Authors refers to north-western Tunisia (lines 67-73, line 101)**
- Lines 77, 360 **Include a reference**
- Line 83 soil **organic carbon levels?**
- Lines 128-129 **ranges of slopes corresponding to high, steep and moderate** can be included. These categories are the same as in table 2?
- Line 185 **Soil type should be included of the study soils**
- Figure 2 The 24 **different units can be delineating**

- Table 2 include aspect for steep category. “in” can be removed (in %). The exact number of soil samples per unit should be included in Table 2 (see lines 155 – 156) by including land use

- Lines 219 and 251 31%

- Line 203 This sentence from the previous version of the manuscript “Soil samples were air dried (to 30 °C) and sieved to pass through a 2 mm mesh.” Should be included.

- Please avoid the repetition of information by deleting lines 206-207 (info repeated in lines 199-200); lines 207-213 (repeated in lines 200-201 and figure 2); line 245 (info is provided in lines 232-233); line 268 (info is included in line 271); Lines 418 – 423 (info is repeated in lines 394-409)

- Line 253 (vario MICRO tube, Elementar)

- Line 258 This study is focused on SOC then this sentence can be omitted.

Did Authors check that TC measurements and SOC measurements are coherent? TC >> SOC

Authors measured SIC by other method? To control / verify their SOC estimates?

- Line 342 – 350 Please revise the readability for a fluent English style. Similarly in the abstract. See the repetition of decline in two continuous sentences (lines 344-345)

- Lines 366 -372 This is not a discussion of the results. This paragraph presents materials and methods and objectives. Please remove because this information was included previously.

- Line 387 “enhanced the SOC content” instead of improved

- Line 388 appropriate land management instead of good land management

- Line 389 “as it reduced both the area covered only by very old cereal monocultures and the soil degradation.” This sentence is unclear in the present form.

- Line 405 assuming that forest soils are not affected by soil erosion because their higher SOC contents. I do not agree with this statement.

- Line 409 Not only low erosion rates please include other key factors for high SOC content in forest soils: higher carbon input, low litter decomposition ...

- Lines 410-417 This general information about Mediterranean soils should be move to the beginning of the discussion

- Lines 420 – 421 Please revise and rewrite, the sentence in the present form is unclear

- “In lands where field crops once were, if they have been interplanted with permanent crops, the SOC content has improved”

- Line 421 “Overgrazing and bad management of grazing lands has led to SOC decreases.” Please move to line 412 and link with inappropriate land management
- Line 422 Remove “Finally, forest land use has the highest SOC content, as it is protected by forest regulation and less disturbed.” This info is included in line 409 “which is highly related to the lower disturbance in the forests”
- Line 428 “Different soil properties encountered among” Which soil properties have been determined by authors apart from SOC. This information should be very useful to include a description in materials and methods about soil characteristics and soil type
- To say “undoubtedly have effects on SOC content” these key soil properties should be included
- Line 436 in soils
- Line 462 I doubt about this statement: “This can be explained by the fact that soils on flat slopes tend to be thicker as a result of deposition”. Have authors any evidence from the results?
- Line 476 nutrients loss instead of nutrition loss
- Lines 478-483 This information is repeated and discussed previously in Discussion section
- Lines 392 and Lines 547 – 552 Since the work is not focused on C sequestration and it is not introduced previously. The main topic was SOC variability as soil quality parameter and its controlling factors (topography and land use). I recommend remove this paragraph.

1 **Impacts of land use and topography on soil organic carbon in a Mediterranean landscape (north-**
2 **western Tunisia)**

3 Donia Jendoubi^{1, 2}, Hanspeter Liniger¹ and Chinwe Ifejika Speranza²

4 ¹ Centre for Development and Environment (CDE), University of Bern, Bern, 3012, Switzerland

5 ² Institute of Geography, University of Bern, Bern, 3012, Switzerland

6 *Correspondence to:* Donia Jendoubi (Donia.jendoubi@cde.unibe.ch)

7 **Abstract:**

8 This study evaluates the impact of land use and topographic features (slope and aspect) on soil organic
9 carbon (SOC) within the Wadi Beja watershed in north-western Tunisia. A soil spectral library was set
10 to assess the variation in the SOC of 1440 soil samples from four land use types (field crops, permanent
11 crops, forest, and grazing land), three slope categories (flat, moderate, and steep) and two aspects
12 (north- and south-facing). For field crops, only one factor – slope – significantly affected SOC, with SOC
13 content in north-facing areas appearing to be higher in flat areas (0.75%) than in hilly areas (0.51%).
14 However, in south-facing areas, SOC content was also higher in flat areas (0.74%) than in hilly areas
15 (0.50%). For permanent crops, which were inter-planted with field crops, the slope significantly affected
16 SOC content, which improved to 0.97% in flat north-facing and 0.96% in flat south-facing areas, scoring
17 higher than hilly south- and north-facing areas (0.79%). In the grazing land use system, both of the
18 investigated factors – aspect and slope – significantly affected the SOC content, which was significantly
19 higher in flat areas (north-facing: 0.84%, south-facing: 0.77%) than in hilly areas (north-facing: 0.61%,
20 south-facing: 0.56%). For the forest, none of the factors had a significant effect on SOC content, which
21 was higher in flat areas (north-facing: 1.15%, south-facing: 1.14%) than in steep areas (1.09% in north-
22 facing and 1.07% in south-facing). This study highlights the ability of visible and near-infrared (VNIR)
23 spectroscopy to quantify C in diverse soils collected over a large diverse geographic area to indicate
24 that calibrations are feasible, and therefore, assessing the variation of SOC content under land use and
25 topographic features (slope and aspect) will result in better sustainable land management planning.

26 **Keywords:** soil organic carbon – land use – spectroscopy – topography – north-western Tunisia

27 **1. Introduction:**

28 Land degradation is a major challenge for Mediterranean arid and semi-arid ecosystems (Hill et al.,
29 2008). In Tunisia, human activities are responsible for land degradation through deforestation,
30 overgrazing, removal of natural vegetation, and agricultural practices that exacerbate soil erosion
31 (Sarraf et al., 2004). Long-term anthropogenic pressure from agricultural use (Kosmas et al., 2015),
32 together with abiotic factors such as climatic and topographical variability (Scarascia-Mugnozza et al.,

Deleted: units

Deleted: units

Deleted: people

Deleted: erode soils

Deleted: in addition to

Deleted: variability

39 2000), create diverse situations for which it is difficult to draw generally valid assumptions concerning
40 SOC distribution and its determinant factors (Jobbagy and Jackson, 2000).

41 Further, soil quality degradation contributes to the deterioration of other land resources (e.g. water and
42 vegetation) (Karamesouti et al., 2015). Soil degradation processes include biological degradation (e.g.
43 soil fertility and soil fauna decline), physical degradation (e.g. compaction, soil erosion, and
44 waterlogging), and chemical degradation (e.g. acidification nutrient and depletion (Diodato and
45 Ceccarelli, 2004; Post and Kwon, 2000), which are caused by agricultural practices.

46 The soil quality concept has been proposed for application in studies on sustainable land management
47 (Doran, 2002). When using the term “soil quality”, it must be linked to a specific function. In this study,
48 soil quality is seen in relation to soil conservation in agricultural systems, which aims to maintain the
49 capacity of soil to function as a vital living system for sustaining biological productivity, promoting
50 environmental quality, and maintaining plant and animal health (Doran and Zeiss, 2000).

51 To measure soil quality, minimum data sets have been suggested that allow a detailed description by
52 including soil chemical and physical indicators (Lal 1998). However, integrative indicators are more
53 appropriate for preliminary studies, as they efficiently provide insights into general soil quality. Soil
54 organic matter (OM) is one such integrative measure of soil quality, influencing soil stability, soil
55 fertility, and hydrological soil properties. OM plays a crucial role in soil erosion: when the erosion
56 removes surface soil, the OM and clay vanish, resulting in a decline of soil fertility and biological
57 activity, and to soil aggregation (Wolfgramm et al., 2007). In soils with high calcareous silty amounts
58 and in the absence of clay, OM is particularly important with regard to the soil's physical properties
59 (e.g. soil structure, porosity, and bulk density), which again determine erodibility (Hill and Schütt,
60 2000).

61 Mediterranean soils are characterized by low amounts of OM, which results in a soil fertility decline
62 and structure loss (Van-Camp et al., 2004). Furthermore, SOC is variable across land use (Brahim et al.,
63 2010), and most agricultural soils are poor in OM, often comprising less than 1% (Achiba et al., 2009;
64 Parras-Alcántara et al., 2016; Muñoz-Rojas et al., 2012). In Mediterranean soils, loss of OM leads to a
65 reduction in root penetration, soil moisture, and soil permeability, which in turn reduces vegetation
66 cover and biological activity, and increases runoff and risk of erosion (Stanners and Bourdeau, 1995).

67 Tunisia has one the highest SOC depletion rates among Mediterranean countries (Brahim et al., 2010).
68 Its low soil fertility is considered a sign of its predominant inappropriate land management systems
69 (Hassine et al., 2008; Achiba et al., 2009). In north-western Tunisia, soils are mostly derived from an
70 alteration of carbonate sedimentary parent material (marl, limestone, clay), cultivated under rainfed
71 conditions to produce cereal crops (wheat and barley), (Hassine et al., 2008). This form of cultivation
72 decelerates the mineralization of OM through a series of unsustainable practices including deep
73 ploughing in spring and summer, stubble ploughing in autumn to protect wheat against Fusarium, and
74 various tillage operations preceding sowing (Hassine et al., 2008). This relatively intensive soil

Deleted: soil organic carbon (

Deleted:)

Deleted: Soil degradation is a key component of land degradation in the Mediterranean region (Hartemink 2003), and s

Deleted: deterioration

Moved (insertion) [2]

Deleted: ¶

Moved up [2]: When using the term “soil quality”, it must be linked to a specific function. In this study, soil quality is seen in relation to soil conservation in agricultural systems, which aims to maintain the capacity of soil to function as a vital living system for sustaining biological productivity, promoting environmental quality, and maintaining plant and animal health (Doran and Zeiss, 2000). ¶

Formatted: Not Highlight

Deleted: resulting in fertility decline, biological activity, and aggregation

Deleted: T

Deleted: he

Deleted: from the study area

Formatted: English (United Kingdom)

95 cultivation, accompanied by the practice of an annual application of phosphate and nitrogen fertilizers,
96 is at the root of the decrease in OM content following stimulation of microbial activity (Álvaro-Fuentes
97 et al., 2008).

98
99 Understanding the above dynamics and SOC distribution as influenced by land use systems and
100 topographic features is critical for assessing land use management planning (Kosmas et al., 2000). SOC
101 contents are influenced by topographic features and climate variation, specifically temperature and
102 water (García Ruiz et al., 2012).

103 Currently, the north-west region of Tunisia is enduring extensive field crop monoculture and land
104 degradation owing to population increase, inappropriate land management, and rough topographic
105 features. Much of the cropped land is unsuitable for agriculture and degrades quickly. The impacts of
106 agricultural practices and topography on nutrient cycling and ecological health, however, have not
107 been studied extensively in the Tunisian northwest region.

108 Due to this dispute, SOC contents measured through time can establish the long-term productivity and
109 possible sustainability of a land use system. In a nutrient-poor system, SOC can play an important role
110 in the stability, quality, and fertility of the soil. Farmers and land use planners are therefore interested
111 in land use management that will enhance soil carbon contents.

112 In this study, we explore SOC distribution according to land use across topography (slopes and aspects)
113 in north-western Tunisia. The aim of this study is to quantify SOC content and evaluate the factors that
114 affect SOC variation, specifically the mechanisms affecting differences in SOC distribution patterns
115 along different land use systems and topographic features (slope and aspect) in a Mediterranean
116 ecosystem dominated by agricultural activities.

117 Information on soil quality is crucial for improving decision-making around efficient support of
118 sustainable land management. Thus, methods are needed to allow fast and inexpensive prediction of
119 important soil quality indicators such as SOC. The potential of diffuse reflectance spectroscopy in the
120 visible and near infrared (VNIR) range for fast prediction of soil properties in a non-destructive and
121 efficient way has been demonstrated by a number of studies (Amare et al., 2013; Shiferaw and
122 Hergarten, 2014; Shepherd and Walsh, 2002).

123 We are not aware of any study evaluating the impacts of topographic features (slope and aspect) or
124 existing land use systems on SOC dynamics in Mediterranean agricultural soils, specifically in Tunisia,
125 based on an accurate and consistent database such as a soil spectral library.

126 Most soils in north-western Tunisia are exposed to water erosion, which is provoked by poor cover,
127 cultivation practices and hilly topography. The Wadi Beja watershed was selected because it comprises
128 a variety of degraded areas and areas where soil and water conservation practices (SWC) are applied.
129 It is the most productive and extended cereal area in Tunisia, and faces serious risks associated with
130 monoculture production of field crops under inappropriate land management practices. Some new

Deleted: the

Formatted: English (United Kingdom)

Deleted: distribution

Deleted: is

Deleted: factors

Deleted: formations

Deleted: soil carbon

Deleted: levels

Deleted: soil organic carbon (

Deleted:)

Deleted: levels

Deleted: units

Deleted: factors

143 practices, such as agroforestry, were introduced into the region in the 1980s, along with permanent
144 crops such as olive and almond trees.

145 The first research objective was to build a soil spectral library in order to apply it in the Wadi Beja
146 watershed, as there was no accurate or valid soil database for the studied region or even for the whole
147 country. The second objective was to examine the distribution of SOC under the different slopes,
148 aspects, and land use systems. The third objective was to investigate, specifically, three research
149 questions: (1) How does SOC vary under cereal monoculture and then after inter-planting with
150 permanent crops? (2) How and why are ecosystems more sensitive to soil degradation (SOC loss) on
151 steep and south-facing slopes than on gentle and north-facing slopes? (3) How can land management
152 practices under different abiotic factors (e.g. topography) influence soil SOC variation, and what
153 practices are recommended in this case study?
154

Deleted: interplantation

Deleted: organic carbon (


Deleted:)

155 2. Materials and Methods:

156 2.1. Study area

157 The study area, the Wadi Beja watershed, lies at 36°37'60" N and 9°13'60" E in north-western Tunisia.
158 Upstream of Wadi Beja is the Amdoun region, and downstream the junction with Wadi Medjerdah in
159 the Mastutah region. Wadi Beja is a tributary of the Wadi Majerdah, the most important river in Tunisia
160 (figure 1).

161



165

166 **Figure 1.** Characterization of household income, location and land uses of the study area, Wadi Beja
 167 watershed, north-western Tunisia. Source: Jendoubi et al., 2019

168 The watershed (about 338 km²) covers diverse topographic features, with an elevation ranging from
 169 110 m a.s.l to nearly 750 m a.s.l; slopes ranges from flat, moderate to steep surfaces, with 64% having a
 170 high to steep slope and 36% a moderate slope. Annual rainfall is irregular and varies from 200 mm to
 171 800 mm. Early October to the end of April (late autumn to early spring) are considered the rainy season,
 172 (AVFA, 2016). During the summer it is very dry and hot. The maximum temperatures are recorded at
 173 the end of July and range from 38°C to 44°C. Minimum temperatures are recorded at the end of
 174 December and fall between 6°C and 8°C (AVFA, 2016). In the Beja region, the population is mainly
 175 rural (56%), with 48.5% active in the agricultural sector. Agriculture remains the main source of
 176 household income (55%, including livestock) (figure 1). Nearly 78% of rural households live entirely
 177 off their farms (AVFA, 2016). There are three types of farming systems: extensive (83%), intensive (6%),
 178 and mixed (11%). Five different land use systems (LUS) have been defined: field crops (71%), grazing
 179 lands (10%), forest (9%), permanent crops (7%), and built-up areas (3%).

Deleted: environments

Deleted: an

Deleted: d

Deleted: which,

Deleted:

Deleted: of the surface has

Deleted: has

Deleted: s

188 The current soil types in the study area are vertisols, which cover 46% of the total area, isohumic soils
189 (23%), brown calcareous soils (12%) and regosols (10%). Rendzinas soils, lithosoils, hydromorphic soils
190 and fersiallitic soils exist, covering small areas that add up to less than 9% according to the agricultural
191 map of Tunisia.

Deleted: database

192 Land management in the study area is similar in relation to land preparation, organic amendments,
193 crop rotation, and mulching (stubble, roots). Mineral fertilizers have been applied for several decades,
194 and cropland – the major land use – has been used for monoculture of cereal crops such as wheat and
195 barley.

196

197 2.2. Methods

198 2.2.1. Land use change history

199 A land use system (LUS) is defined as the sequence of goods and services obtained from land, but can
200 involve particular management interventions undertaken by the land users as well. It is generally
201 determined by socio-economic market forces, as well as the biophysical constraints and potentials
202 imposed by the ecosystems in which they occur (Nachtergaele et al., 2010).

203 This study investigated four land use systems – field crops, permanent crops, forest plantation, and
204 grazing land – in order to assess their effects on the variation of SOC (table 1). Built-up areas and roads
205 were excluded. We used atmospherically corrected Landsat Surface Reflectance data images (Bands 4-
206 5, 7 and 8) from 1985, 2002, and 2016 to derive the land use maps, in order to evaluate the changes over
207 that time period (Jendoubi et al., 2019).

208 The Landsat scenes were selected from among all those available in the green season (out of harvesting)
209 for the corresponding years; we considered only those with less than 20% of cloud cover overall and
210 without clouds on the study site area. Unsupervised classification was carried out for the images in
211 order to define the major land use systems. Following this, a validation based on ground truth data was
212 made in order to confirm the generated land use maps and assess their accuracies.

213 Table 1 illustrates substantial land use and land cover change (LULC) in the Wadi Beja watershed after
214 1980.

215 **Table 1.** The five major land use and management classes studied in the Wadi Beja watershed,
216 Tunisia

Aggregated land use classes	1985		2002		2016	
	%	km ²	%	km ²	%	km ²
Field crops	82.1	272.7	76.4	254.0	71.0	236.2
Grazing lands	9.3	30.9	10.2	33.7	9.7	32.2
Forests	3.9	13.1	7.7	25.6	8.9	29.6
Permanent crops	3.4	11.2	4.2	14.1	7.3	24.4
Built-up areas	1.3	4.5	1.5	4.9	3.1	10.0
Total	100	332.4	100	332.4	100	332.4

Formatted: Left

Formatted Table

219
220
221
222

Source: Jendoubi et al., 2019

223 Field crops constituted the predominant land use type, accounting for approximately 82% in 1985 and
224 71% in 2016. Plantation forest also increased from 3.9% in 1985 to 9% of the watershed in 2016. In 1980,
225 to remedy the degrading effects of monoculture of annual cropping, deforestation, and overgrazing on
226 the pastures and the forests, a programme developed by ODESYFANO (Office Development Sylvo-
227 Pastoral Nord Ouest) and financed by the World Bank implemented some conservation activities
228 including development of permanent vegetative cover using olive trees and sylvo-pastoral
229 management. An agroforestry (agro-sylvo-pastoral) system was introduced in 1982 as an alternative
230 programme for development and conservation in the region. This system included converting annual
231 cropping into a combination of annual crops inter-planted with olive trees (in this study classified as
232 “permanent crops”). This area increased from 3.4% in 1985, when it was introduced for the first time in
233 the region, to 7.3% in 2016. The local farmers took this alternative as they believed that their soils had
234 become poor and no longer gainful for annual crop production. Grazing land remained almost
235 unchanged in terms of area, as it is spread over badlands, barren lands, and riverbanks with a high
236 concentration of eroded and poor soils.

237 2.2.2. Soil sampling

238

239 We selected four land use systems (LUS) (excluding built-up areas), three slope classes, and two aspect
240 classes to study their interrelations and their effects on SOC. The LUS were forests, field crops,
241 permanent crops, and grazing land (table 1). Aspect and slope units were derived from Lidar DTM,
242 aligned, and resampled to 30m. Slope categorization was based on the FAO soil description guidelines
243 (Barham et al., 1997). The slope categories were grouped into three: flat, moderate and steep. Aspect
244 was categorized into two classes: north and south. Details about slope and aspect categories are
245 presented in table 2.

246

Table 2. Slope and aspect

247

248

Slope (%)	Aspect (azimuth degrees)
0 to 8 (Flat)	0 to 90, 270 to 360 (North)
8 to 16 (Moderate)	90 to 270 (South)
> 16 (Steep)	


249

250

251 From all slope, aspect classes, and different land use systems (LUS), soil samples were collected
252 randomly from the topsoil (0-20 cm). In a factorial randomized design considering the four land use
253 types, the three slopes, and two aspects, a total of 24 different sampling units ($n=4 \times 3 \times 2$) were

Deleted: in

255 considered. In total, 1440 soil samples were collected from all the sampling units in the topsoil layer (0-
256 20 cm) using a soil auger (10 cm diameter) with an average of 60 samples per sampling unit.



257
258

Figure 2. Location of the soil samples and the sampling design.

259 The sampling design shown in figure 2 summarizes the strategy of the sampling. Sampling units are
260 listed as shown in table 3.

261 **Table 3. Sampling units with their corresponding number of soil samples**

Sampling units	Number of soil samples
Field crops - flat - north	65
Field crops - flat - south	66
Field crops - moderate - north	60
Field crops - moderate - south	62
Field crops - steep - north	57
Field crops - steep - south	59
Permanent crops - flat - north	60
Permanent crops - flat - south	62
Permanent crops - moderate - north	55
Permanent crops - moderate - south	57
Permanent crops - steep - north	63
Permanent crops - steep - south	65
Forests - flat - north	60
Forests - flat - south	54
Forests - moderate - north	57
Forests - moderate - south	60
Forests - steep - north	61
Forests - steep - south	59
Grazing lands - flat - north	60
Grazing lands - flat - south	63
Grazing lands - moderate - north	55
Grazing lands - moderate - south	59
Grazing lands - steep - north	62
Grazing lands - steep - south	59

262

263

264 2.2.3. Soil analysis and spectral library

265 The soil spectral library was set according to protocols cited by Shepherd and Walsh (2002), and
266 includes the following steps: (1) representative sampling of soil variability in the study area; (2)
267 establishing the soil reflectance spectral dataset using VNIR spectrometry; (3) selecting a reference
268 dataset to be analysed using traditional soil chemical methods required as reference values (450

Deleted: , where each soil sample can be taken in a randomized way from any specific sampling unit

Deleted: follows

Formatted: Font: (Default) +Body (Calibri), Not Bold, Complex Script Font: 11 pt, Not Bold

272 samples, or 31% of the total, were selected according to their spectral variability); (4) determination of
273 SOC by means of soil chemical analysis (CNS elemental analysis); (5) calibrating soil property data to
274 soil reflectance spectra by applying multivariate calibration models; and finally (6) prediction of new
275 samples using the spectral library.

Deleted: 0

276 The soil spectral library for prediction of SOC was adjusted using a mug-light for illumination as
277 described by Mutuo et al. (2006). Soil spectral reflectance was measured under standard conditions in
278 the laboratory. Soil samples were air dried (to 30°C) and sieved to pass through a 2 mm mesh. Soil
279 samples of 2 mm thickness were filled into borosilicate Duran glass Petri dishes with optimal optical
280 characteristics. The Petri dishes were placed on a mug-light equipped with a Tungsten Quartz Halogen
281 light source (Analytical Spectral Devices, Boulder, CO). Spectral reflectance readings were collected
282 through the bottom of the Petri dishes using a FieldSpec PRO FR spectro-radiometer (Analytical
283 Spectral Devices, Boulder, CO). Every sample was measured twice, with the sample rotated by 90
284 degrees for the second measurement. The two measurements were averaged, which minimized light
285 scatter effects from uneven particle size distribution on the Petri dish floor. The instrument works with
286 three spectro-radiometers to cover the wavelengths from 350 to 2500 nm at an interval of 1 nm. The
287 fore-optic view was set to 8 degrees. For dark current readings, 25 scans were averaged, while for white
288 reference and soil spectral readings 10 scans were averaged by the spectro-radiometer. Before each
289 sample reading, white reference readings were taken from a spectralon (Labsphere) that was placed on
290 a trimmed Petri dish bottom.

Moved up [1]: Soil samples were air dried (to 30°C)

Moved (insertion) [1]

Deleted: Soil samples were air dried (to 30°C)

Deleted: Air-dried ground soil

291 Pre-processing of soil reflectance data to decrease the noise present in the data and thus increase
292 robustness of reflectance spectral data is common in VNIR spectrometry, and is especially
293 important in the case of measuring setups that are difficult to control (e.g. due to power fluctuations
294 or different operators during different measuring sessions). The main pre-processing steps conducted
295 were as follows: Spectra were compressed by selection of every 10th nm. Spectral bands in the lowest
296 (350-430 nm) and highest (2440-2500 nm) measurement ranges were omitted due to a low signal to
297 noise ratio (lower than 90). The final number of wavelengths used as model input was 205. Information
298 for these 205 wavelengths was further processed. Steps in the spectral reflectance curves were observed
299 at the spectrometer changeovers. Most likely, this effect resulted from the Petri dishes used as sample
300 holders and their specific index of refraction.

Deleted: : the instrument covers the full wavelength range with three spectro-radiometers.

301 When choosing the validation set, care was taken to assure that validation samples were representative
302 for the whole study area. Thus, samples were systematically chosen by selecting from every land use
303 system and under the different (slope and aspect) sampling units. These samples, which constituted
304 31% of the total samples, were selected for chemical analysis, which was used to validate SOC model
305 prediction.

Deleted: 0

313 Regarding the chemical method, the elemental CNS analyser (vario ~~MICRO~~ tube, ~~Elementar~~) was used
314 for SOC estimates. For SOC measurement, 1 g soil is pre-treated with 10 drips of H₃PO₄ in order to
315 remove carbonate. The sample is combusted at 1150°C with constant helium flow, carrying pure oxygen
316 to ensure complete oxidation of organic materials. The CO₂ gas is produced and detected by a thermal
317 conductivity detector.

Deleted: micro

Deleted: c

318 A calibration and validation with Partial Least Square Regression were used based on cross-validation
319 (“leave one out”) in order to ensure simultaneous reduction/correlation of both the spectral
320 information and the concentration data obtained from the chemical analysis.
321 After prediction of the remaining SOC sample values, a set of statistical parameters was applied in
322 order to assess the accuracy of results such as: the coefficient of determination (R²), which measures
323 how well a regression line estimates real data points; the Residual Prediction Deviation (RPD), which
324 evaluates the quality of a validation; and the Root Mean Square Error of the prediction (RMSEP), which
325 assesses the accuracy of the model. These parameters evaluate the performance quality of the soil
326 spectroscopy model (Rossel et al., 2006).

Deleted: Total soil carbon is measured, using the same procedure without pre-treatment with H₃PO₄. Soil inorganic carbon is calculated as the difference between total soil carbon and SOC.

327 **2.2.4. Statistical analysis**

328 Regarding the soil spectral library analysis, the partial least squares regression (PLS regression) was
329 used in RStudio to validate the spectral prediction model while assessing the coefficient of
330 determination (R²), residual prediction deviation (RPD), and root mean square error of the prediction
331 (RMSEP).

Deleted: A script for running various steps was made in RStudio in order to generate the soil spectral library. ¶

332 After generating the soil spectral library, a test of normality based on Sullivan and Verhoosel (2013)
333 was carried out in order to check the normality distribution of the data.


334 The Statistical Package for the Social Sciences (SPSS 20.0) software was used in order to compare the
335 averages obtained under the different factors. Variance analyses and multiple comparisons (MANOVA
336 test) were carried out to determine the effect of the different factors (land use, slope, and aspect) on the
337 variation of the SOC. Results were significant when p < 0.05. The interaction effect between the factors
338 was tested using the technique of split file. The results were grouped according to the land use factor
339 and the effect of the slope and aspect were tested in each land use value.

340 Results were presented in histograms using Excel XLSTAT. We then assessed the variation of SOC
341 under the different selected factors.

342 **3. Results**

343 **3.1. Soil spectral library as an integrative indicator of soil quality**

344 SOC content was plotted against SOC content predictions as displayed in figure 3.



353

354

Figure 3: SOC values from chemical analysis plotted against SOC prediction.

355 The obtained spectral prediction model has $R^2=0.85$, RPD= 2.11, and RMSEP= 0.35%, which was rated
 356 excellent for prediction because RPD>2 (Viscarra Rossel et al. 2006). This means that the model is able
 357 to determine accurately the SOC content of 85% of the samples. The RPD (2.11>2) also showed that the
 358 model developed is of good quality and can be used to predict the remaining spectra and for further
 359 development of the spectral library.

Deleted: shows

360 Regarding the normality of the data, the test showed a high correlation of 0.95 between the overall data
 361 and their corresponding z-scores. Therefore, this means that the data are approximately normally
 362 distributed.

Deleted: s

365 **3.2. Significance effects of all the variables**

366 A multivariate MANOVA analysis revealed which variables had statistically significant differences in
 367 SOC related to land use systems, slopes, and aspects. Table 4 **shows** the results of the significance
 368 analysis for each of the three variables. The highest significance was reported for land use, followed by
 369 slope and aspect.

Deleted: reveals

370 **Table 4.** MANOVA results showing the significance of the impact of land use, slope, and aspect for
 371 SOC (n= 1440)

	F	Sig.
LUS	395.263	0.000
slope	76.505	0.000
aspect	11.093	0.001

Deleted: soil organic carbon (

Deleted:)

Formatted Table

372 Sig. < 0.05 (statistically significant difference), in bold.

373 Sig. > 0.05 (no statistically significant difference)

374

375 The analysis of the significance of the different variables for each land use type is presented in table 5.

376 **Table 5.** MANOVA results regarding significance of all the variables under different LUS.

LUS	Variables	F	Sig.
Forests	slope	1.806	0.176
	aspect	2.931	0.094
Field crops	slope	51.429	0.000
	aspect	1.028	0.312
Permanent crops	slope	36.474	0.000
	aspect	0.068	0.795
Grazing lands	slope	8.242	0.001
	aspect	5.971	0.017

Formatted: Left

Formatted Table

377 Sig. < 0.05 (statistically significant difference), in bold.

378 Sig. > 0.05 (no statistically significant difference)

379


380 For forest land use, no variables were significant, indicating that the variation of the SOC with high
 381 **contents** in those components was not related to slope or aspect. For field crops and permanent crops,
 382 only slope had a significant effect on SOC. For grazing lands, both variables (slope and aspect) revealed
 383 significant effects on SOC content.

Deleted: amounts

388 **3.3. SOC according to land use systems**

389 SOC content for different land use systems is shown in figure 4. The forest LUS had the highest SOC
390 content, with 1.09%. Permanent crops had the second highest values with 0.87% of SOC. The lowest
391 SOC content was found for field crops (0.70%) and grazing soils (0.74%).

392



393


394 **Figure 4.** SOC contents according to land use systems in the Wadi Beja watershed, Tunisia.

Deleted: rates

395 According to the MANOVA results, land use systems significantly affect SOC content. In the study
396 area, the lowest SOC content was found in field cropping soils (0.70%), and the highest SOC content in
397 the forests (1.09%).

398 **3.4. Impact of slope and aspect on SOC**

399



401

402

Figure 5. SOC rates according to slope and aspect in the Wadi Beja watershed, Tunisia.

403

Figure 5 shows the highest SOC content (0.81%-0.83%) on flat slopes and slightly reduced SOC on moderate slopes (0.78%-0.79%). Both flat and moderate slopes revealed no significant difference between northern and southern slopes (difference <0.02%). The lowest SOC was on steep southern slopes with 0.63%, followed by steep northern slopes with 0.69% SOC.

404

405

406

407

408

3.5. Impact of land use, slope, and aspect on SOC

Deleted: s

Deleted: show




Figure 6. SOC contents according to slope and aspect for the different land use systems.

When evaluating the impact of slopes on SOC variations under the different LUS, the results presented in figure 6 revealed that in forest plantations, the highest SOC contents were observed in flat (1.15%) and north-facing areas. 1.14% SOC was found on moderate slopes in north-facing areas, and 1.09% on steep north-facing areas. As previously shown, statistically, the slope has no significant effect on SOC variation under the forest LUS.

For field crops, the highest SOC content was found in flat north-facing areas (0.75%), followed by 0.69% on moderate slopes in north-facing areas and then very low figures of 0.51% on steep slopes in north-facing areas. Figure 6 clearly shows a marked decline in SOC with increased slopes under field crops. For permanent crops, the decrease with increasing slopes is less than that of the field crops. The highest SOC content was found in north-facing areas, first on flat slopes (0.97%), second on moderate (0.84%) and third on steep slopes (0.79%). Finally, on grazing lands, the different slopes showed marked differences. In flat north-facing areas, SOC was 0.84% and 0.77% in flat south-facing areas, in moderate north-facing areas it was 0.76% and in moderate south-facing areas, 0.70%, in steep north-facing areas it was 0.61% and 0.56% in steep south-facing areas.

The MANOVA test showed that aspect has no significant effect on SOC variation for forests, field crops, or permanent crops. Only for grazing land does aspect have a significant effect on SOC variation, with north-facing soils having a greater SOC content than south-facing areas. See figure 6 and table 5.

- Deleted: rates
- Deleted: amounts
- Deleted: and 0.50% in south-facing areas.
- Deleted: s
- Deleted: increasing
- Deleted: decline
- Deleted: , and the SOC for all slopes are increased.
- Deleted: flat
- Deleted: followed by
- Deleted: 0.84% in
- Deleted: areas
- Deleted: 0.79% in
- Deleted:
- Deleted: areas
- Deleted: ,
- Deleted: with
- Deleted: of SOC in flat north-facing areas,
- Deleted: in flat
- Deleted: in moderate north-facing areas
- Deleted: ,
- Deleted: in moderate south-facing areas
- Deleted: in steep north-facing areas,
- Deleted: s
- Deleted: level

454 4. Discussion

455 The Mediterranean region is generally characterized by poor soils with low SOC content (around 2 %) due to their nature and to being overused by agriculture, which means that they have low carbon
456 inputs from plant residues and low canopied density, and are subjected to inappropriate management
457 practices (Verheye and De la Rosa, 2005; Cerdà et al., 2015).

Deleted: OM

Deleted: 1

Deleted: C

459 Land management is shown to be a key indicator affecting SOC distribution, influencing topsoil in
460 particular (Ferreira et al., 2012). In Mediterranean areas in particular, land management is a significant
461 factor given the limitations to SOC accumulation. Moreover, high SOC reflects undisturbed soil and
462 high soil quality, as is the case in forest land use (Corral-Fernández et al., 2013).

463 Regarding the soil spectral library, the obtained spectral responses were well correlated, which means
464 that the prediction model is excellent. Some outliers were detected and their corresponding soil samples
465 were assessed to check the reasons for their reflectance. This differences can be explained by the fact
466 that some spectral responses were influenced by the colour of the soils. Wolfgramm (2008) also found
467 that the spectral responses of dark soils give over-predicted values and those of light soils give under-
468 predicted values. In this case, if samples are identified as outliers, the existing spectral library needs to
469 be extended to include all the variable soils. Thus additional reference values from soil chemical
470 analysis have to be obtained and calibration models for an extended reference sample set need to be
471 developed.

Deleted: herefore, t

Deleted: Thus, as

Deleted: s

Deleted: an

Deleted: s

Deleted: an

Deleted: , as is the case in a study made by Wolfgramm (2008)

Deleted: c

Deleted: ing the results

Deleted: from

Deleted: carried out

Deleted: OM

Deleted: have

Deleted: have been

Deleted: We thus focused on the SOC content, which we calibrated from soil reflectance spectra. Chemical analysis of SOC made it possible to calibrate and validate a model using soil spectra to predict a wider range of soil samples. The soil spectral library of 1440 soil samples was used to investigate SOC under the various combinations of land use, slopes, and aspects.

472 Compared with the study by Hassine et al. (2008), which concluded that SOC content does not exceed
473 2% in north-western Tunisia, our prediction model falls within this amount with a maximum organic
474 carbon percentage of 1.2%. This state of low SOC in soils used for agriculture, compared to forests with
475 little indication of soil degradation, has been confirmed by various authors (Arrouays et al., 1994; Cerri,
476 1988; Robert, 2002). This low content has negative impacts on the soil structure, which is built mainly
477 by means of mineral colloids and whose stability is affected, leading to numerous deficiencies in
478 production and susceptibility to degradation factors. Cereal soils may have acquired a balance between
479 SOC inputs and losses, but at a very low equilibrium level compared to forests; with the latter having
480 less decline of SOC and being protected against erosion, which is the main type of land degradation in
481 the study area (Hassine et al., 2008).

482 Previous studies show that SOC can play a significant role in monitoring soil quality related to land
483 use and reduction of soil degradation (Shukla et al., 2006; Hassine et al., 2008). The soil spectral library
484 made it feasible to gain some interpretations and therefore to generate some recommendations for land
485 use planners regarding assessing SOC variability, as an integrative soil quality measure.

Deleted: and assess SOC variation in the study area,

Deleted: .

514 The results on the impacts of land use on SOC indicate that field crops have the lowest SOC content.
515 This could be the result of land degradation due to inappropriate agricultural management such as
516 intensive tillage, the removal of crop residues, reduced vegetation cover, deteriorated soil aggregation
517 and erosion, and a continuous monoculture system. This finding is coherent with the results of several
518 researchers (Lemenih and Itanna, 2004; Lal, 2005; Muñoz-Rojas et al., 2015; Hamza and Anderson, 2005)
519 who have revealed a significant decline in SOC content in cropland compared to natural forests. Herrick
520 and Wander (1997) found that in annual cropping systems, the distribution of SOC is highly influenced
521 by land management practices such as reduced tillage, rotation, fertilization, and shifting cultivation.
522 Consistent with the study by Hassine et al. (2008) in north-western Tunisia, the reduced OM
523 decomposition rates are a result of intensive agricultural practices; monoculture, tillage on steep slopes,
524 and tillage in wet seasons, in addition to other topographic features, which may lead to a decrease in
525 SOC.

Field Code Changed

Deleted: OM

Deleted: factors

526 Changing annual field crops by inter-planting them with permanent tree crops has increased the SOC
527 of soils under previous annual field crops almost halfway to the level of SOC in forests (0.87%).
528 Intercropping previously mono-cropped fields with tree crops (olive, almond, and pomegranate trees)
529 between 1982 and 1985 significantly enhanced the SOC within 30-35 years. Creating agroforestry
530 systems in this way is considered to have been an appropriate land management intervention in north-
531 western Tunisia. However, some farmers made no changes to their land management, as they did not
532 perceive the advantages of the agroforestry system (Jendoubi and Khemiri, 2018). Yet agroforestry
533 systems are globally recognized, since they are more accomplished at capturing and utilizing resources
534 than grassland systems or single-species cropping (Nair et al., 2011).

Deleted: improved

Deleted: a good

Deleted: , as it reduced both the area covered only by very old cereal monocultures and the soil degradation.

Deleted: as having a high potential to sequester C

535 Grazing lands, even though they are not tilled, have a low content of SOC (0.74%), only slightly higher
536 than annual crops (Figure 3). Continued overgrazing and reduction of vegetation cover seem to
537 degrade the soils and their SOC. A low SOC content can continue due to a lack of appropriate grassland
538 management. Open pasture without canopies and weak grass-vegetation cover increase the
539 vulnerability of this land use system to soil degradation and SOC decline. Various studies have shown
540 that the way grazing land is managed affects SOC (Wu et al., 2003; Soussana et al., 2004): overused
541 grazing lands with less vegetation cover are more affected by soil erosion and soil exposure to wind
542 and rain, leading to greater SOC loss. Notably, grassland management strongly affects SOC contents,
543 which decrease as grazing intensities increase (Neff et al., 2005).

Deleted: level

Deleted: stocks

544 The highest SOC contents were found in the forests. The explanation for this is that forest has a dense
545 cover that protects soil from being exposed to any other factors such as erosion, hence the SOC contents
546 are less affected. This finding has been confirmed by many authors who have shown that in
547 Mediterranean areas, many forest soils are rich in OM; as a consequence, these soils supply a large

Deleted: amounts

Deleted: and

Deleted: cannot be

Deleted: It can be assumed that the soil under forest has no degradation caused by soil erosion from water, as observed in some surrounding fields.

563 input of carbon, and have a low litter decomposition, which means that they are distinguished by high
564 SOC (Lal, 2005; FAO, 2010), which is highly related to the lower disturbance in the forests.

565 Regarding the impact of slope on SOC variation, our results show that the higher the slope, the lower
566 the SOC content. Irvin (1996) specified that generally, with increasing slope, OM lixiviation is reduced,
567 mineral is weathered, clay is translocated, and horizons are differentiated.

568 Moreover, topography has a significant impact on soil temperature, soil erosion, runoff, drainage, and
569 soil depth - and hence soil formation. The accumulation of SOC variation on hillslopes is explained by
570 the decomposition rates of OM and litter input differences (Yimer et al., 2006).

571 When assessing the results of the impact of aspects on SOC variation, south-facing terrain has lower
572 SOC content than north-facing terrain, which is explained by its exposure to the highest solar radiation
573 and, in particular, the highest temperature during the vegetation period and the long hot summers.
574 This implies high evaporation and a high burn down of OM due to high temperature, less moisture in
575 soils, and consequently a slow-down of the decomposition of OM.

576 In addition, according to our findings, the impact of both slope and aspect on SOC content was very
577 distinct, as indicated statistically by a significant effect on SOC content in the MANOVA. The issue is
578 that steep and south-facing slopes are more sensitive to degradation than other areas, which is
579 explained by the fact that steepness increases runoff and soil erosion, and southern exposure increases
580 evapotranspiration and temperatures, thus decreasing the availability of nutrients, water, and SOC to
581 plants. Apart from differences in land use management, SOC variation is mainly affected by
582 environmental factors in soil along with topographic features (slope and aspect).

583 The literature links temperature and moisture to OM decomposition in soils (García Ruiz et al., 2012;
584 Griffiths et al., 2009). As shown by Garcia-Pausas (2007), in the Mediterranean area, shaded areas such
585 as northern-facing or colder southern areas sustain regularly high moisture content for longer and
586 consequently become more fertile and productive, in contrast to the southern-facing areas that are
587 exposed to high radiation and thus occasional water deficits.

588 With regard to steepness and aspect, the higher the slope, the more exposed to the south, and the more
589 affected by erosion and different climatic conditions, the lower the SOC content (Yimer et al., 2007;
590 Yimer et al., 2006). Different topographic features are considered to have different microclimatic and
591 vegetation community types and thus significant variations in SOC. Topography (slope and aspect)
592 hence plays a crucial role in relation to temperature and moisture regimes. The temperature is highly
593 influenced by solar radiation, which has a role in soil chemical and biological processes and vegetation
594 distribution (Bale et al., 1998). Hence, the temperature of the soil plays a key role in monitoring the
595 biomass decomposition rate, and thus affects the SOC distribution, either by delaying or accelerating
596 its decomposition (Scowcroft et al., 2008).

597 From the results of assessing the impact of slope combined with land use, we can see that the highest
598 SOC content was observed in the flat area under all land use systems, and it tended to decrease in steep
599 positions. In general, under all land use systems, we can observe the same tendency of SOC variation,
600 ranging from highest SOC content in flatter positions to lowest in steep positions.

Deleted: quantity

Deleted: The Mediterranean region is generally characterized by poor soils with low OM content (around 1%) due to their nature and to being overused by agriculture, which means that they have low C inputs from plant residues and low canopied density, and are subjected to inappropriate management practices (Verheye and De la Rosa, 2005; Cerdà et al., 2015). ¶

Land management is shown to be a key indicator affecting SOC distribution, influencing topsoil in particular (Ferreira et al., 2012). In Mediterranean areas in particular, land management is a significant factor given the limitations to SOC accumulation. Moreover, high SOC reflects undisturbed soil and high soil quality, as is the case in forest land use (Corral-Fernández et al., 2013). ¶

The interpretations emphasize that the impacts of land use on SOC variation is highly related to land management practices. The findings highlight the contribution of overuse and monoculture to SOC decline under the field crops land use system. In lands where field crops once were, if they have been interplanted with permanent crops, the SOC content has improved. Overgrazing and bad management of grazing lands has led to SOC decreases. Finally, forest land use has the highest SOC content, as it is protected by forest regulation and less disturbed. ¶

Deleted: With regards

Deleted: ic

Deleted: position

Deleted: Different soil properties encountered among topographic units (slope and aspect) will affect the litter production and decomposition, which will undoubtedly have effects on SOC content.

Deleted: the

Deleted: units

Deleted: positions

639 This can be explained by **minimal erosion and even deposition of sediments from neighbouring slopes**
640 **and thus accumulation of SOC**. Erosion causes stripping of the soil in hillslope areas. As shown by Yoo
641 et al. (2006), the prevalent portion of SOC is deposited in depositional areas, with hillslopes being more
642 susceptible to sporadic mass wasting events, continuous soil erosion and production, and consequently
643 less SOC storage. In addition, the highest erodibility is related to hilly areas where soils have a tendency
644 to be shallow, coarse in texture, and low in OM, while lower erodibility is observed in flat areas with
645 organic-rich, deep, and leached soils (Lawrence, 1992).
646 From the clear difference in the variation in SOC under forest and field crop land use systems, we
647 interpret that it is the land use factor that dominates SOC distribution rather than the slope factor.
648 In general, steep slopes have a lower SOC content than flat land, as they are more vulnerable to erosion,
649 especially when associated with inappropriate management and overuse (Reza et al., 2016; Bouraima
650 et al., 2016). Cropland in sloping areas is highly vulnerable to water erosion, which leads to extensive
651 soil disturbance, while land use patterns affect vegetation cover, soil physical properties such as SOC,
652 and surface litter. Therefore, this provokes the runoff and soil erosion processes that accompany
653 **nutrients loss** (Dagnew et al., 2017; Montenegro et al., 2013). Therefore, the extent of **nutrients loss**
654 differs according to land use systems, as is the case with cereal monoculture in the study site.
655 Hence, in order to improve and maintain soil quality parameters for sustainable productivity, it is
656 crucial to reduce intensive cultivation and integrate the use of inorganic and organic fertilizers.
657 In agricultural areas, continuous intensive cultivation without appropriate soil management practices
658 has contributed to loss of SOC. Kravchenko et al. (2002) and Jiang and Thelen (2004) found that within
659 variability in topography, slope was considered to be a major crop yield limiting factor.
660 Correspondingly, after **inter-planting** permanent crops with field crops, SOC content was enhanced.
661 Herrick and Wander (1997) showed that after introducing permanent crops, slope significantly affected
662 SOC content.
663 **Our results also showed** less SOC in south-facing areas, **and confirms findings from other studies on**
664 **the interaction effects of slope and aspect on OM decomposition** (Griffiths et al., 2009).
665 According to McCune and Keon (2002), the reason for these results is that slope and aspect play a
666 significant role in solar radiation redistribution, hence the solar radiation heterogeneity on hillslopes
667 leading to differences in soil moisture and temperature. Huang et al. (2015) stated that the **SOC content**
668 in shaded aspect areas was significantly higher than in sunny aspect areas. Therefore, **increases in SOC**
669 **and OM accumulation are supported by** increased moisture and reduced temperature **while decreased**
670 **soil temperature usually results in decreased OM decomposition rates and litter decay rates**
671 (Blankinship et al., 2011).
672 For grazing lands, all the variables (slope and aspect) revealed significant effects on SOC content as also
673 shown in the study of Bird et al. (2001). SOC content is generally low, though it is higher in flat areas.
674 This is explained by overgrazing and pressure **on** the different topographic **features**, as they are all
675 easily accessible to livestock. Even on steep slopes there is pressure and overgrazing, in addition to the
676 exposure of these areas to erosion by wind and rain. This **highlights the susceptibility** of this land use
677 system to erosion and deterioration of soil quality.

Deleted: the fact that soils on flat slopes tend to be thicker as a result of deposition.

Deleted: nutrition

Deleted: nutrition

Deleted: Therefore, under different land use systems, the difference in SOC content is related to the effect of variation in land use system intensity along the toposequences. As shown by our results, higher SOC content was recorded in the forest where there is less disturbance and use, and statistically slope has no significant effect on SOC variation. In the field cropping area, the fact that soils are overused and subject to continuous intensive cultivation without appropriate soil management practices has contributed to the degradation of important soil quality indicators such as SOC.

Deleted: ation of

Deleted: in combination

Deleted: , in keeping with the tendency for the highest SOC content to be in flat areas and the lowest in steep areas

Deleted: According to the obtained results on the impact of all factors on SOC variation, with

Deleted: revealed

Deleted: our finding lends strong

Deleted: support to

Deleted: , as the difference between north- and south-facing areas is solar radiation, wind, and rainfall

Field Code Changed

Deleted: concentration

Deleted: as discussed previously,

Deleted: generated by means of

Deleted: . D

Deleted: in

Deleted: positions

Deleted: may reveal

Deleted: vulnerability

714 Why grazing land use systems are the most sensitive to all the tested variables (slope and aspect), can
715 be explained as follows: in the study area, grazing land was generally open grassland and it is evident
716 that soils are more sensitive in open grassland than under tree canopies, as SOC stocks under tree
717 canopies are in general higher than in open grassland (e.g. Seddaiu et al., 2013). Moreno et al. (2007)
718 also found that the SOC content in the topsoil beneath the tree canopies as being “around twice as high
719 as beyond the tree canopy”. This can also be related to overgrazing, as shown in a literature review of
720 the effects of overgrazing in the Mediterranean basin (Sanjari et al., 2008; Costa et al., 2012).
721 Furthermore, the semi-arid climate and inclined topography prevailing in the Mediterranean grazing
722 lands render ecosystems vulnerable to SOC losses. As shown by Ryan et al. (2008), the higher the level
723 of grazing, or the greater the residue removal, the greater the decline in mean SOC level. The reason
724 behind the decrease in carbon and nutrient cycling is mainly that SOC in grassland is accumulated in
725 roots, which leads to its loss with every removal of aboveground biomass.

726 The most likely clarification for the results obtained on decreased SOC content in steep south-facing
727 areas under the field crops land use system is that soils are affected by soil degradation initiated by
728 inappropriate land management and consequently have weak vegetation cover. This condition makes
729 these soils more sensitive to the south-facing exposition characterized by higher solar radiation and
730 evaporation, and thus decreases soil moisture, biological activity, and SOC loss. Wakene and Heluf
731 (2004) have also indicated that intensive cultivation aggravates OM oxidation and hence reduces SOC
732 content.

733
734 Therefore, some options for sustainable land management practices can be recommended, such as
735 establishment of enclosures (Mekuria and Aynekulu, 2013), which could be efficient in recovering the
736 degraded grazing land areas of the watershed. In addition to protecting trees against damage caused
737 by uncontrolled grazing animals by installing fences and trunk protection, mixing of animal species,
738 mostly sheep and goats, but also cows and horses and setting additional fodder provision could be a
739 feature of the summer season.

740 In order to maintain improved soil quality and sustainable productivity in croplands, there is a need to
741 reduce intensive cultivation, agroforestry, and practice of fallow, integrate use of inorganic and organic
742 fertilizers, and pay more attention to the most vulnerable areas (steep and south-facing areas).

743 There are strong indications that agroforestry has been successful in retaining and even improving SOC
744 and soil fertility: the results show that introducing an agroforestry system – e.g. combining an olive
745 plantation with annual field crops – has increased SOC content in the most vulnerable areas. Thus, such
746 types of sustainable land use should be the focus of land managers and land use planners.

747

748 5. Conclusions

749

Deleted: are

Deleted:

Deleted: ?

Deleted: This

Deleted: by the fact that

Deleted: case

Deleted: stated:

Deleted: projected

Deleted:

Deleted: “T

Deleted: amount

Deleted: content of

Deleted: projection was

Field Code Changed

Deleted: OM

Deleted: OM

Deleted: (

Deleted: ping

767 Land management can profoundly affect SOC stocks. In areas with exceedingly erodible soils, such as
768 those on steep slopes and south-facing zones as shown in this study, application of soil and water
769 conservation measures is crucial to sustain agricultural fields and prevent or reduce soil degradation.
770 Greater efforts to reduce SOC decline are required on steep slopes and south-facing land than in flat
771 areas and north-facing land. However, a further study of the area is recommended, especially land use
772 in combination with other topographic features such as altitude and curvature and their effects on SOC
773 content.
774 By far the best option, however, is to identify land management practices that increase C stocks whilst
775 at the same time enhancing other aspects of the environment, e.g. improved soil fertility, decreased
776 erosion, greater profitability, or improved yield of agricultural and forestry products. There are a
777 number of management practices available that could be implemented to protect and enhance existing
778 C sinks now and in the future.
779 Since such practices are consistent with, and may even be encouraged by, many current international
780 agreements and conventions, their rapid adoption should be as widely encouraged as possible.
781 Finally, this paper contributes towards filling a gap in investigation on the impacts of various land uses
782 on SOC in Tunisia. The results presented in this paper are valid for calibration of further soil spectral
783 libraries in north-western Tunisia; this was the first soil spectral library collated in Tunisia and the
784 methodology can be replicated and applied to other areas. Further studies on SOC variation depending
785 on land use and topographic features are needed to inform sustainable land management in Tunisia.

786 6. Acknowledgments

787 This research was funded by the Islamic Development Bank, Grant/Award Number: 78/TUN/P34 and
788 the Swiss Federal Commission for Scholarships for Foreign Students, Grant/Award Number:
789 2014.0968/Tunesien/OP. The authors thank Tina Hirschbuehl, Dee Cooke, and William Critchley for
790 English proofreading.

791 7. Conflict of Interest Statement

792 The authors affirm that there are no conflicts of interest regarding the publication of this paper.

793 8. References:

794 Achiba, W.B., Gabteni, N., Lakhdar, A., Laing, G.D., Verloo, M., Jedidi, N., & Gallali T.: Effects of 5-year
795 application of municipal solid waste compost on the distribution and mobility of heavy metals in a
796 Tunisian calcareous soil. Agric. Ecosyst. Environ. 130, 156 – 163, 2009.
797

Deleted: soil C

Deleted: and appropriate management

Deleted: can be used to sequester soil C

Deleted: As with all human activities, the social dimension, especially land management, needs to be considered when implementing soil C sequestration practices. Since there will be increasing competition for limited land resources in the coming century, soil C sequestration cannot be viewed in isolation from other environmental and social needs. In order to increase soil C sequestration, as part of wider plans to enhance sustainable land management, more attention should be paid to the importance of land use and the different topographic factors.

Deleted: to reduce SOC decline

Deleted: s

Deleted: factors

Deleted: factors

816 Agence de la Vulgarisation et de la Formation Agricoles (AVFA) : <http://www.avfa.agrinet.tn>.
817 Informations Régionales > Beja. Agricoles. Tunisie, Ministère de l'Agriculture des ressources
818 hydrauliques et de la pêche, 2016.
819 Álvaro-Fuentes, J., López, M., Cantero-Martínez, C., & Arrúe, J. : Tillage effects on soil organic carbon
820 fractions in Mediterranean dryland agroecosystems. *Soil Science Society of America Journal*, 72(2), 541-
821 547, 2008.
822 Amare, T., Hergarten, C., Hurni, H., Wolfgramm, B., Yitafaru, B., & Selassie, Y. G.: Prediction of soil
823 organic carbon for Ethiopian highlands using soil spectroscopy. *ISRN Soil Science*, 2013.
824 Arrouays, D., Kicin, J., Péliissier, P., & Vion, I. : Evolution des stocks de carbone des sols après
825 déforestation: analyse spatio-temporelle à l'échelle d'un paysage pédologique. *Étude et gestion des sols*,
826 1(2), 7-15, 1994.
827 Bale, C.L. Williams, B.J. & Charley. J.L.: The impact of aspect on forest structure and floristics in some
828 Eastern Australian sites. *For. Ecol. Manag.*, 110, pp. 363-377, 1998.
829 Bird S.B., Herrick J.E. & Wander M.M.: Exploiting heterogeneity of soil organic matter in rangelands:
830 benefits for carbon sequestration. R.F. Follett, J.M. Kimble, R. Lal (Eds.), *The Potential of U.S. Grazing*
831 *Lands to Sequester Carbon and Mitigate the Greenhouse Effect*, CRC Press, Boca Raton FL, USA, 2001
832
833 Blankinship, J.C., Niklaus, P.A. & Hungate, B.A.: A meta-analysis of responses of soil biota to global
834 change. *Oecologia* 165: 553- 565, 2011.
835 Bouraima, A.K., He, B.& Tian T.: Runoff, nitrogen (N) and phosphorus (P) losses from purple slope
836 cropland soil under rating fertilization in Three Gorges Region Environ. Sci. Pollut. Res. Int., 23 (5)
837 (2016), pp. 4541-4550, 2016.
838 Brahim, N., Bernoux, M., Blavet, D. & Gallali, T.: Tunisian soil organic carbon stocks. *International*
839 *Journal of Soil Science* 5: 34-40, 2010.
840 Cerdà, A., González-Pelayo, Ó., Giménez-Morera, A., Jordán, A., Pereira, P., Novara, A., Brevik, E.C.,
841 Prosdoci, M., Mahmoodabadi, M., Keesstra, S., García Orenes, F. & Ritsema, C.: The use of barley
842 straw residues to avoid high erosion and runoff rates on persimmon plantations in Eastern Spain under
843 low frequency - high magnitude simulated rainfall events. *Soil Research*. 54: 154-165.
844 DOI:10.1071/SR15092, 2016.
845 Cerri, C. : Dynamique de la matiere organique du sol après défrichement et mise em culture. Utilisation
846 du traçage isotopique naturel em 13C. *Cah. ORSTOM*, 24, 335-336, 1988.
847 Corral-Fernández, R., Parras-Alcántara L. & Lozano-García, B.: Stratification ratio of soil organic C, N
848 and C: N in Mediterranean evergreen oak woodland with conventional and organic tillage. *Agric.*
849 *Ecosyst. Environ.*, 164, pp. 252-259, 2013.
850 Costa, C., Papatheodorou, E.M., Monokrousos, N., Stamou, G.P.: Spatial variability of soil organic C,
851 inorganic N and extractable P in a Mediterranean grazed area. *Land Degradation & Development*
852 <http://dx.doi.org/10.1002/ldr.2188>, 2012.
853 Dagnew, D.C., Guzman, C.D., Akal, A.T., Tebebu, T.Y., Zegeye, A.D., Mekuria, W., Tilahun, S.A. &
854 Steenhuis, T.S.: Effects of land use on catchment runoff and soil loss in the sub-humid ethiopian
855 highlands. *Ecophysiol. Hydrobiol.*, 17 (4), pp. 274-282, 2017. Diodato, N., & Ceccarelli, M.: Multivariate
856 indicator Kriging approach using a GIS to classify soil degradation for Mediterranean agricultural
857 lands. *Ecological Indicators*, 4(3), 177-187, 2004.

Deleted: .

Field Code Changed

859 Doran, J. W.: Soil health and global sustainability: translating science into practice. *Agriculture,*
860 *Ecosystems & Environment*, 88(2), 119-127, 2002.

861 Doran, J. W., & M. R. Zeiss.: Soil health and sustainability: managing the biotic component of soil
862 quality. *Appl. Soil Ecol.* 15: 3 - 11, 2000.

863 FAO.: *Global Forest Resources Assessment 2010.* (378 pp.), 2010.

864 Ferreira, A.O., Sá, J.C.M., Harms, M.G., Miara, S., Briedis, C., Quadros, C., Santos, J.B., Canalli, L.B.S.
865 & Dias, C.T.S.: Stratification ratio as soil carbon sequestration indicator in macroaggregates of Oxisol
866 under no-tillage. *Cienc. Rural*, 42, pp. 645-652. (in Portuguese), 2012.

867 García Ruiz, J., Lana Renault, N., Nadal Romero, E., & Beguería, S. (2012). Erosion in Mediterranean
868 Ecosystems: changes and future challenges. Paper presented at the EGU General Assembly Conference
869 Abstracts.

870 Garcia-Pausas, J. , Casals, P. , Camarero, L. , Huguet, C. , Sebastià, M.-T. , Thompson, R. , Romanyà, J.:
871 Soil organic carbon storage in mountain grasslands of the Pyrenees: effects of climate and topography.
872 *Biogeochemistry* 82, 279–289, 2007.

873 Griffiths, R. P., Madritch, M. D., & Swanson, A. K.: The effects of topography on forest soil
874 characteristics in the Oregon Cascade Mountains (USA): Implications for the effects of climate change
875 on soil properties. *Forest Ecology and Management*, 257(1), 1-7, 2009.

876 Hamza, MA. & Anderson, WK.: Soil compaction in cropping systems - a review of the nature, causes
877 and possible solutions. *Soil & Tillage Research*, 82, 121–145, 2005.

878 Hassine, H.B., Aloui, T., Gallali, T., Bouzid, T., El Amri, S., & Hassen, R. : Évaluation quantitative et
879 rôles de la matière organique dans les sols cultivés en zones subhumides et semi-arides
880 méditerranéennes de la Tunisie. *Agrosolutions*, 19, 4-14, 2008.

881 Herrick, J.E. & Wander, M., Relationships between soil organic carbon and soil quality in cropped and
882 rangeland soils: the importance of distribution, composition, and soil biological activity. R. Lal, J.M.
883 Kimble, R.F. Follett, B.A. Stewart (Eds.), *Soil Processes and the Carbon Cycle*, CRC Press, Boca Raton
884 (1997), pp. 405-425, 1997.

885 Hill, J., & Schütt, B.: Mapping complex patterns of erosion and stability in dry Mediterranean
886 ecosystems. *Remote Sensing of Environment*, 74(3), 557-569, 2000.

887 Hill, J., Stellmes, M., Udelhoven, T., Röder, A., & Sommer, S.: Mediterranean desertification and land
888 degradation: mapping related land use change syndromes based on satellite observations. *Global and*
889 *Planetary Change*, 64(3-4), 146-157, 2008.

890 Huang, Y.M., Liu, D. & An, S.S.: An Effect of slope aspect on soil nitrogen and microbial properties in
891 the Chinese Loess region. *Catena*, 125 (2015), pp. 135-145, 2015.

892 Irvin, B.J.: *Spatial Information Tools for Delineating Landform Elements to Support Soil/Landscape*
893 *Analysis.* PhD Thesis, University of Wisconsin-Madison.
894 grunwald.ifas.ufl.edu/Nat_resources/soil_forming_factors/formation.htm (Accessed on
895 05/12/2007), 1996.

896 Jendoubi, D. & Khemiri H. : Le système d'Agroforesterie pour la protection des terres et l'amélioration
897 des revenus des exploitants dans les zones montagneuses de Nord Ouest Tunisien.
898 https://qcat.wocat.net/en/wocat/technologies/view/technologies_3722/, 2018.

899 Jendoubi, D., Hodel, E., Liniger, H.P. & Subhatu, A.T.: Land degradation assessment using landscape
900 unit approach and normalized difference vegetation index in Northwest of Tunisia. *Journal of*
901 *Mediterranean Ecology* vol. 17, 2019. 67-79. © Firma Effe Publisher, Reggio Emilia, Italy, 2019.

Deleted: Hartemink, A. E.: Soil fertility decline in the tropics: with case studies on plantations: Cabi., 2003. ¶

904 Jianga, P. & Thelen, K.D.: Effect of Soil and Topographic Properties on Crop Yield in a North-Central
905 Corn-Soybean Cropping System. *Agronomy Journal Abstract - SITE-SPECIFIC ANALYSIS*. Vol. 96 No.
906 1, p. 252-258. doi:10.2134/agronj2004.0252, 2004.

907 Jobbagy, E.G. & Jackson, R. B.: The vertical distribution of soil organic carbon and its relation to climate
908 and vegetation. *Ecol. Appl.* 10, 423--436, 2000.

909 Karamesouti, M., Detsis, V., Kounalaki, A., Vasiliou, P., Salvati, L., & Kosmas, C.: Land-use and land
910 degradation processes affecting soil resources: evidence from a traditional Mediterranean cropland
911 (Greece). *Catena*, 132, 45-55, 2015.

912 Kosmas, C., Detsis, V., Karamesouti, M., Kounalaki, K., Vassiliou, P., & Salvati, L.: Exploring long-term
913 impact of grazing management on land degradation in the socio-ecological system of Asteroussia
914 Mountains, Greece. *Land*, 4(3), 541-559, 2015.

915 Kosmas, C., Gerontidis, S. & Marathanou, M.: The effect of land use change on soils and vegetation
916 over various lithological formations on Lesbos (Greece). *Catena*, 40, pp. 51-68, 2000.

917 Kravchenko, A. N. & Bullock, D. G.: Spatial variability of soybean quality data as a function of field
918 topography: II. A Proposed technique for calculating the size of the area for differential soybean
919 harvest. *Crop Science* 42, 816-821, 2002.

920 Lal, R.: Soil quality and sustainability. In *Methods for Assessment of Soil Degradation*. *Advances in*
921 *Soil Science*. R. Lal, W.H. Blum, C. Valentine, and B.A. Stewart (eds.). CRC Press, Boca Raton, FL, pp.
922 17-30, 1998.

923 Lal, R.: Forest soils and carbon sequestration. *Forest Ecology and Management*, 220(1-3), 242-258., 2005

924 ▾
925 Lawrence, W.M.: The variation of soil erodibility with slope position in a cultivated Canadian prairie
926 landscape. *Earth Surface Processes and Landforms*. Vol. 17, Issue 6, Wiley and Sons, Ltd. pp. 543-556,
927 1992.

928 Lemenih, M., & Itanna, F.: Soil carbon stocks and turnovers in various vegetation types and arable lands
929 along an elevation gradient in southern Ethiopia. *Geoderma*, 123(1-2), 177-188, 2004.

930 McCune, B. & Keon, D.: Equations for potential annual direct incident radiation and heat load *J. Veg.*
931 *Sci.*, 13, pp. 603-606, 2002.

932 Mekuria, W., & Aynekulu, E.: Enclosure land management for restoration of the soils in degraded
933 communal grazing lands in northern Ethiopia. *Land Degradation & Development*, 24(6), 528-538, 2013.

934 ▾
935 Montenegro, A.A.A., Abrantes, J.R.C.B., Lima, J.L.M.P.D., Singh, V.P. & Santos, T.E.M.: Impact of
936 mulching on soil and water dynamics under intermittent simulated rainfall. *Catena*, 109 (10), pp. 139-
937 14; 2013.

938 Moreno, G., Obrador, J.J. & García, V.: Impact of evergreen oaks on soil fertility and crop production in
939 intercropped dehesas. *Agr. Ecosyst. Environ.*, 119, pp. 270-280, 2007.

940 Mutuo, PK., Shepherd, KD., Albrecht, A. & Cadisch, G.: Prediction of carbon mineralization rates from
941 different soil physical fractions using diffuse reflectance spectroscopy. *Soil Biology & Biochemistry* 38:
942 1658-1664, 2006.

943 Muñoz-Rojas, M., Jordán, A., Zavala, L., De la Rosa, D., Abd-Elmabod, S., & Anaya-Romero, M.: Impact
944 of land use and land cover changes on organic carbon stocks in Mediterranean soils (1956-2007). *Land*
945 *Degradation & Development*, 26(2), 168-179, 2015.

Deleted: .

Deleted:

948 Muñoz-Rojas, M., Jordán, A., Zavala, L.M., De la Rosa, D., Abd-Elmabod, S.K. & Anaya-Romero, M.:
949 Organic carbon stocks in Mediterranean soil types under different land uses (Southern Spain). *Solid*
950 *Earth*, 3, pp. 375-386, 10.5194/se-3-375-2012, 2012.
951 Nachtergaele, F., Biancalani, R., Bunning, S. & George, H.: Land degradation assessment: the LADA
952 approach. In 19th World Congress of Soil Science, Brisbane, Australia 1st Aug 2010.
953 Nair, P.K.R., Nair, V.D., Kumar, B.M. & Showalter, J.M.: Carbon sequestration in agroforestry systems.
954 *Adv. Agron.*, 108, pp. 237-307, 2011.
955 Neff, J.C., Reynolds, R.L., Belnap, J. & Lamothe, P.: Multi-decadal impacts of grazing on soil physical
956 and biogeochemical properties in southeast Utah. *Ecological Applications*, 15 (1), pp. 87-95, 2005.
957 Parras-Alcántara, L., Lozano-García, B., Keesstra, S., Cerdà, A. & Brevik, E.C.: Long-term effects of soil
958 management on ecosystem services and soil loss estimation in olive grove top soils. *Sci. Total Environ.*,
959 571, pp. 498-506, 2016.
960 Ping, C.L., Jastrow, J.D., Jorgenson, M.T., Michaelson, G.J. & Shur, Y.L.: Permafrost soils and carbon
961 cycling. *Soil*, 1, pp. 147-171, 2015.
962 Post, W. M., & Kwon, K. C.: Soil carbon sequestration and land-use change: processes and potential.
963 *Global change biology*, 6(3), 317-327, 2000.
964 Reza, A., Eum, J., Jung, S., Choi, Y., Owen, J.S. & Kim, B.: Export of non-point source suspended
965 sediment, nitrogen, and phosphorus from sloping highland agricultural fields in the East Asian
966 monsoon region. *Environ. Monit. Assess.*, 188 (12), p. 692, 2016.
967 Robert, M.: La séquestration du carbone dans le sol pour une meilleure gestion de terres, 2012.
968 Ryan, J., Masri, S., Ibriki, H., Singh, M., Pala, M. & Harris, H.: Implications of cereal-based crop
969 rotations, nitrogen fertilization, and stubble grazing on soil organic matter in a Mediterranean-type
970 environment. *Turkish J. Agric. For*, 32, 289-297, 2008.
971 Sanjari, G., Ghadire, H., Ciesiolka, CAA. & Yu, B.: Comparing the effects of continuous and time-
972 controlled grazing systems on soil characteristics in Southeast Queensland. *Australian Journal of Soil*
973 *Research* 46: 348-358. 10.1071/SR07220, 2008.
974 Sarraf, M. Larsen B. & Owaygen, M.: Cost of Environmental Degradation – The Case of Lebanon and
975 Tunisia. *Environmental Economics series*. Paper no. 97. World Bank publications. June 2004.
976 Scarascia-Mugnozza, G., Oswald, H., Piussi, P., & Radoglou, K.: Forests of the Mediterranean region:
977 gaps in knowledge and research needs. *Forest Ecology and Management*, 132(1), 97-109, 2000.
978 Scowcroft, P., Turner, D.R. & Vitousek, P.M.: Decomposition of *Metrosideros polymorpha* leaf litter
979 along elevational gradients in Hawaii *Glob. Chang. Biol.*, 6, pp. 73-85, 2008.
980 Seddaiu, G. Porcu, G., Ledda, L., Roggero, P.P., Agnelli, A., et al.: Soil organic matter content and
981 composition as influenced by soil management in a semi-arid Mediterranean agro-silvo-pastoral
982 system. *Agric Ecosyst Environ* 167: 1-11 doi <http://dx.doi.org/10.1016/j.agee.2013.01.002>, 2013.
983 Shepherd, K. D., & Walsh, M. G.: Development of reflectance spectral libraries for characterization of
984 soil properties. *Soil Science Society of America Journal*, 66(3), 988-998, 2002.
985 Shiferaw, A., & Hergarten, C.: Visible near infrared (VisNIR) spectroscopy for predicting soil organic
986 carbon in Ethiopia. *Journal of Ecology and the Natural Environment*, 6(3), 126-139, 2014.
987 Shukla, M., Lal, R., & Ebinger, M.: Determining soil quality indicators by factor analysis. *Soil and Tillage*
988 *Research*, 87(2), 194-204, 2006.

Deleted:

990 Soussana, J.F., Loiseau, P., Vuichard, N., Ceschia, E., Balesdent, J., Chevallier, T., Arrouays, D.: Carbon
991 cycling and sequestration opportunities in temperate grasslands. *Soil Use and Management*, 20, pp.
992 219-230, 2004.

993 Stanners, D., & Bourdeau, P.: Europe's environment: the Dobris assessment Europe's environment: the
994 Dobris assessment: Office for Official Publication of the European Communities, 1995.

995 Sullivan, M., & Verhoosel, J.C.: *Statistics: Informed decisions using data*. New York: Pearson; 2013.

996 Van-Camp, L., Bujarrabal, B., Gentile, A.-R., Jones, R.J.A., Montanarella, L., Olazabal, C. & Selvaradjou,
997 S.-K.: Reports of the Technical Working Groups Established under the Thematic Strategy for Soil
998 Protection. EUR 21319 EN/3. Office for Official Publications of the European Communities,
999 Luxembourg 872 pp. 2004.

1000 Verheye, W. & De la Rosa, D.: Mediterranean soils, in *Land Use and Land Cover, from Encyclopedia of*
1001 *Life Support Systems (EOLSS)*, Developed under the Auspices of the UNESCO, EOLSS Publishers,
1002 Oxford, UK, 2005.

1003 Viscarra Rossel, R. A., Walvoort, D. J. J., McBratney, A.B., Janik, L. J. & Skjemstad, J. O.: "Visible, Near
1004 Infrared, Mid Infrared or Combined Diffuse Reflectance Spectroscopy for Simultaneous Assessment of
1005 Various Soil Properties." *Geoderma* 131 (1-2): 59-75. doi:10.1016/j.geoderma.2005.03.007, 2006.

1006 Wakene, N. & Heluf, G.: The impact of different land use systems on soil quality of western Ethiopia
1007 Alfisols. *International Research on Food Security: Natural Resource Management and Rural Poverty*
1008 *Reduction through Research for Development and Transformation*. Deutcher Tropentage-Berlin 5-7
1009 October 2004. pp. 1-7. [http://www. Tropentage.de/2004/abstracts/full/265.pdf](http://www.Tropentage.de/2004/abstracts/full/265.pdf), 2004.

1010 Wolfgramm, B., Seiler, B., Kneubühler, M., & Liniger, H.: Spatial assessment of erosion and its impact
1011 on soil fertility in the Tajik foothills. *EARSel eProceedings*, 6(1), 12-25, 2007.

1012 [Wolfgramm, B.: Land use, soil degradation and soil conservation in the loess hills of central Tajikistan. PhD thesis, Bern University, Switzerland. 2007.](#)

1013 Wu H., Guo A. & Peng C. : Land use induced changes
1014 in organic carbon storage in soils of China. *Glob. Change Biol.* 9: 305-315, 2003.

1015 Yimer, F., Ledin, S. & Abdelkadir, A.: Soil organic carbon and total nitrogen stocks as affected by
1016 topographic aspect and vegetation in the Bale Mountains, Ethiopia. *Geoderma*, 135, pp. 335-344, 2006.

1017 Yimer, F., Ledin, S. & Abdelkadir, V.: Changes in soil organic carbon and total nitrogen contents in
1018 three adjacent land use types in the Bale Mountains, south-eastern highlands of Ethiopia. *Fores. Ecol.*
1019 *Manage.*, 242, pp. 337-342, 2007.

1020 Yoo, K., Amundson, R., Heimsath, A.M. & Dietrich, W.E.: Spatial patterns of soil organic carbon on
1021 hillslopes: integrating geomorphic processes and the biological C cycle. *Geoderma*, 130, pp. 47-65, 2006.

1022

Deleted: ¶

Deleted: ¶

Deleted: ¶

Deleted: ¶

Formatted: English (United Kingdom)

Deleted: