

Arable soil formation and erosion: a hillslope-based cosmogenic-nuclide study in the United Kingdom

Daniel L. Evans¹, John N. Quinton¹, Andrew M. Tye², Ángel Rodés³, Jessica A.C. Davies¹, Simon. M. Mudd⁴, and Timothy A. Quine⁵

¹ Lancaster Environment Centre, Lancaster University, Lancaster, Lancashire, UK

² British Geological Survey, Keyworth, Nottinghamshire, UK

10 ³ Scottish Universities Environmental Research Centre, East Kilbride, UK

⁴ School of GeoSciences, University of Edinburgh, Edinburgh, UK

⁵ College of Life and Environmental Sciences, University of Exeter, Exeter, UK

Correspondence to: D. L. Evans (d.evans3@lancaster.ac.uk)

15

20

25

30

5 Abstract

Arable soils are critical resources that support multiple ecosystem services. They are frequently threatened, however, by accelerated erosion. Subsequently, policy to ensure their long-term security is an urgent societal priority. Although long-term security relies upon a balance between the rates of soil loss and formation, there have been few investigations of the formation rates of soils supporting arable agriculture. This paper addresses this knowledge gap by presenting the first isotopically-constrained soil formation rates for an arable (Nottinghamshire, UK) and coniferous woodland hillslope (Shropshire, UK). Rates ranged from 0.026 mm year⁻¹ to 0.096 mm year⁻¹ across the two sites. These rates fall within the range of previously published rates for soils in temperate climates and on sandstone lithologies but significantly differed to those measured in the only other UK-based study. We suggest this is due to the parent material at our sites being more susceptible to weathering. Furthermore, soil formation rates were found to be greatest for aeolian-derived sandstone when compared with fluvially-derived lithology raising questions about the extent to which the petrographic composition of the parent material governs rates of soil formation. On the hillslope currently supporting arable agriculture, we utilised cosmogenically-derived rates of soil formation and erosion in a first-order lifespan model and found, in a worst-case scenario, that the backslope A horizon could be eroded in 138 years with bedrock exposure occurring in 212 years under the current management regime. These findings represent the first quantitative estimate of cultivated soil lifespans in the UK.

Copyright statement.

25

30

5 1.0 Introduction

Soil erosion is a significant threat to society (Pimentel et al., 1995; UNCCD, 2017). Whilst uncultivated ‘pristine’ soils may develop steady-state thicknesses, where erosion and production are in dynamic equilibrium (Phillips, 2010), human-induced erosion has led to soil thinning across many landscapes (Montgomery, 2007). Soil erosion, left unchecked, can ultimately lead to the removal of the soil cover and the exposure of the underlying parent material (Amundson et al., 2015). The development of soil conservation strategies has long been an active field for research and practice (Panagos et al., 2016; Govers et al., 2017). Given any long-term strategy to preserve soil resources relies upon a balance between the rates of soil loss and soil renewal (Hancock et al., 2015), the measurement of soil formation is a fundamental component in these conservation efforts.

The mechanisms associated with soil formation have been studied for over a century, with a focus on the development of soil horizons and the evolution of soil properties (Dokuchaev, 1879; Jenny, 1941; Bryan and Teakle, 1949; Tugel et al., 2005). Efforts to quantify the rates at which soils form from parent materials have included studying how soil properties change across chronosequences (Turner et al., 2018), developing chemical weathering models (Burke et al., 2007) and, in particular, employing terrestrial cosmogenic radionuclide analysis (Heimsath et al., 1997). In the latter, the concentrations of radioactive isotopes in the bedrock, which are partly dependent upon the rate at which bedrock transforms into soil, are measured and assumed to equal the rates of soil formation.

Despite the recent advancements in cosmogenic radionuclide analysis, their application in soil science has, arguably, not been fully realized. Moreover, there are three research challenges that may explain this. First, there is a dearth of soil formation rate data. Whilst there have been many attempts at calculating a global average soil formation rate from collating multiple inventories (Alexander, 1988; Montgomery, 2007; Stockmann et al., 2014; Minasny et al., 2015), these datasets often omit more than 100 countries, particularly in Africa and Europe, presenting a clear rationale for more studies to take place in these areas of the world. Second, over 80% of the soil formation rate inventory, comprising data from Montgomery (2007), Portenga and Bierman (2011) and Stockmann et al. (2014), is attributed to samples taken from outcrops and stream sediments procured from drainage basins. Moreover, only 252 ^{10}Be -derived rates from this inventory of 1850 stem from samples extracted from underneath the soil mantle. In addition, the majority of these stem from mountain regions and deserts (Heimsath et al., 1997; Wilkinson et al., 2005; Zhao et al., 2018; Struck et al., 2018). This is partly because the observation and estimation of bedrock weathering rates is most commonly carried out by the geomorphological community, principally

to identify the mechanisms behind long-term landscape evolution (Heimsath, 2006; Heimsath and Burke, 2013; Ackerer et al., 2016; Zhao et al., 2018). As a result, there has been no investment in deriving rates of soil formation for soils that support arable agriculture (Heimsath, 2014), despite these soils being identified as a societal priority (FAO, 2015). Such soils are critical to the delivery of multiple ecosystem services and, for many countries, are one of the most critical resources in ensuring the health of society and sustained economic growth. They are also often intensely managed and thus the loci for accelerated erosion (Quinton et al., 2010; Borrelli et al., 2017). However, in the absence of soil formation rate data, the magnitude of the threat erosion places on the sustainability of soils and arable production is unknown, amounting to a critical knowledge gap. Third, although the distributions of inventoried soil erosion and formation rates are often presented together to demonstrate the severity of soil erosion (Montgomery, 2007; Minasny et al., 2015), the spread of globally-compiled data is such that it cannot offer a useful forecast of the sustainability of soil at a site scale. Both distributions are platykurtic and there is substantial overlap in these rates: 0—28.8 mm year⁻¹ for soil formation (Minasny et al., 2015) and 0—52.9 mm year⁻¹ for soil erosion (Montgomery, 2007). For a greater understanding into the sustainability of soil resources at the local scale, we argue that soil scientists should undertake empirical measurements of both soil formation and erosion in parallel.

In this UK-based study, we present ¹⁰Be-derived soil formation rates for two catena sequences in an arable and coniferous woodland setting. The former are the first of their kind globally and the latter are the first of their kind in Europe. We place our results in the context of the rates previously derived in similar climatic and petrographic settings around the world. Finally, using previously measured soil erosion rates at the arable site, we calculate first-order soil productive lifespans to infer the long-term sustainability of the soil resource.

2.0 Materials and Methods

2.1 Site Description

This study measures soil formation down two catena sequences (Figure 1). The first is an arable hillslope at Rufford Forest Farm (RFF), east of Mansfield in Nottinghamshire, UK (53°7'13.43" N, 1°4'39.61" W). The second is a woodland hillslope at Comer Wood (CW), north of Quatford in Shropshire, UK (52°30'30.43" N, 2°22'45.68" W). RFF was selected as it is the site of previous tillage and water-based erosion studies (Quine and Walling, 1991; Walling and Quine, 1991; Govers et al., 1996). Electing CW as a sister site is justified based on its similarities in parent geology, macroclimate and soil physical properties with RFF as detailed below. A Trimble S6 Total Station was used to measure the relative elevation and slope of the catenas at both sites (Figure 1b).

A reconnaissance study of the parent materials and their feasibility for cosmogenic radionuclide analysis was undertaken in spring 2017. Both sites are underlain by Triassic sandstone. In RFF, the Sherwood sandstone (Chester formation; Olenekian, 247—251 Ma) is described as pinkish to red, medium to coarse grained, pebbly, cross-bedded, and friable. In CW, the New

Red sandstone (Bridgnorth formation; Cisuralian, 273—299 Ma) is described as brick-red, medium grained, cross-bedded and aeolian based. Both RFF and CW are south-facing slopes, and sit in a temperate oceanic climate (Cfb), between 96—99 m a.s.l. and 50—71 m a.s.l., respectively. The mean annual precipitation and temperature is 709 mm and 9.8°C at RFF and 668 mm and 9.9°C at CW, respectively (Met Office, 2018).

5

Both sites are positioned beyond the areal limits of the Late Devensian ice sheet, but studies conducted on similar formations of Triassic Sherwood Sandstone nearby suggest that the weathering of the parent material was partly induced by freeze-thaw processes associated with periglacial active layer development possibly during this period (Tye et al., 2012). Although proglacial glaciogenic deposits have been found in the vicinity of CW, the prevalence of similar deposits on the study
10 hillslope has not been studied. However, unpublished work conducted by the authors suggests that the upper (3—5 m) of the lithosphere at both sites was subject to high-magnitude sediment transport at least 200,000 BP or before, potentially during the Anglian glaciation (~450,000 BP). The complex land-use and vegetation change in the Sherwood Sandstone outcrop, within which RFF is based, has been extensively studied and mapped by Tye et al. (2013). Following the onset of the Holocene, the area has been dominated by a complex sequence of land-use change including broadleaf woodland (6000—
15 2000 BC), heathland (43—409 AD) and landscaped heathland for hunting (1600 AD). From at least 1855 AD, RFF has been under an arable regime and in the last twelve years, the dominant crops have been Winter Wheat and Rye. CW is understood to have been an open field until 1903—1926 and then heathland until 1954. Between 1954 and the present day, however, the site has been continuously occupied by coniferous forest (Evans, 2018).

20

25




Figure 1: Locations of the study sites in this paper (a) with elevation profiles (b) for both Comer Woodland (CW; green) and Rufford Forest Farm (RFF; blue). The position of summit (triangles), shoulder (diamonds), backslope (circles) and toeslope (squares) sampling positions are indicated on each profile. Photographs of RFF (c) and CW (d) were taken by the author at the time of sampling.

5

10

15

2.2 Saprolite Extraction and Soil Sampling

- Four positions (summit, shoulder, backslope and toeslope) along a catena transect were selected for depth to bedrock surveys and saprolite extraction. First, a dynamic cone penetrometer was used to estimate the depth of the soil-saprolite interface. At RFF, a percussion drilling rig then proceeded to extract a series of vertical undisturbed core samples of the soil and saprolite.
- 5 Cores were later halved lengthways, and by observing the changes in the consolidation and physical integrity of the extracted material (i.e. whether it remained intact when removed from the core), together with the penetration resistance data acquired in the field, the soil-saprolite interface was demarcated. Two samples of saprolite (5 cm thickness) were then subsampled for cosmogenic radionuclide analysis; one at this interface and one from 50 cm below. At CW, following the use of the dynamic cone penetrometer to locate suitable sites, a soil pit was manually dug vertically at each of the four sampling locations.
- 10 Observing the changes in the consolidation and physical integrity of the material down the profile wall, together with the penetration resistance data, the soil-saprolite interface was ascertained. A sample of saprolite (5 cm thickness) was then extracted from this interface for cosmogenic isotope analysis.

The bombardment of quartz minerals in the uppermost metres of bedrock with cosmic rays leads to the production of ^{10}Be . Assuming the intensity of these cosmic rays and the in situ weathering of bedrock (ϵ) is constant, the concentration of ^{10}Be (N) in a sample of bedrock, Eq. (1), is dependent upon the balance of two factors: the time that the bedrock has been exposed to cosmic rays with longer durations leading to greater concentrations and the weathering of this bedrock into mobile regolith (soil) with greater rates of bedrock weathering leading to smaller concentrations (Lal, 1991; Stockmann et al., 2014). We assume here that the production of ^{10}Be and the erosion of the bedrock is at an equilibrium:

$$N = \sum_{i=sp, \mu_f, \mu^-} \frac{P_i(\theta) \cdot e^{-\frac{x}{\Lambda_i}}}{\lambda + \frac{\epsilon \rho}{\Lambda_i}} \left(1 - e^{-t \left(\lambda + \frac{\epsilon \rho}{\Lambda_i}\right)}\right)$$

(1)

where: P are the annual production rates of ^{10}Be by spallation, fast muons and stopping muons (sp, μf and $\mu\text{-}$) at a surface with slope Θ ; x is the mass sample depth ($\rho \cdot z$); ρ is the density of overburden material; z is the depth of the sample; t is the age of the bedrock surface (the age when the original surface was generated) λ is the decay constant of ^{10}Be with λ equalling $\ln 2 / ^{10}\text{Be}$ half-life; and Λ are the mean attenuation of cosmic radiations (Lal, 1991). t is usually considered infinite. At RFF paper, we took two samples from the same depth profile at each catena position to test if the data support these assumptions. RFF data is compatible with landscape ages >200 ka. Production rates, decay constants and attenuation lengths were calculated using field data and the CRONUS-Earth online calculator v2.3 Matlab code for the St scheme (Balco, 2008). As N can be measured using Accelerator Mass Spectrometry (AMS), Eq. (1) can be solved for ε by simple interpolation of N.

10 A total of twelve samples of saprolite (eight from RFF and four from CW) were prepared for AMS at the Cosmogenic Isotope Analysis Facility, East Kilbride, Scotland. This comprised of mineral separation, quartz cleaning and procedures leading to the preparation of BeO sample cathodes (Kohl and Nishiizumi, 1992; Fifield, 1999; Corbett et al., 2016). The AMS measurements were carried out at the SUERC AMS laboratory (Xu et al., 2010). ^{10}Be concentrations are based on 2.79×10^{-11} $^{10}\text{Be}/^9\text{Be}$ ratio for the NIST Standard Reference Material 4325. The processed blank ratio ranged between 6 and 13%
15 of the sample $^{10}\text{Be}/^9\text{Be}$ ratios. The uncertainty of this correction is included in the stated standard uncertainties. Concentrations of ^{10}Be were subsequently determined, following Balco (2006) (see Supplementary Table 1).

Previous work (e.g. Heimsath, 1997) has assumed that the bulk density of the soil above the bedrock surface is either equal to that of the bedrock, or constant with depth. For this paper, we developed a model called 'coSOILcal' to calculate soil formation rates using empirically measured bulk density data from each catena position at both RFF and CW. The local
20 annual production rate of ^{10}Be at each study site must also account for any obstructions that reduce the cosmic ray flux to the parent material (Phillips et al., 2016). For an obstruction to cause this reduction, it is required to be several metres thick which equates, in practice, to topographic features at the scale of tens of meters or greater. The shielding factor, therefore, is a ratio of the ^{10}Be production rate at the obstructed site to that at an identical site but with a flat surface and a clear horizon (Balco, 2008). To calculate both shielding factors and subsequently normalize local ^{10}Be production rates, site elevation,
25 latitude and longitude were inputted into the CRONUS-Earth Matlab code v2.3 using Lal/Stone (St) scaling (Balco, 2008).

Soil samples were sub-sampled every 5 cm from each core at RFF and on each profile wall at CW. All samples were then oven dried overnight (105°C for 12 hours), grounded with a pestle and mortar, and sieved to discard the >2 mm fraction before being subject to particle size analysis and loss on ignition (LOI). Particle size analysis was conducted using a
30 Beckman Coulter Laser Diffraction Particle Sizing Analyser LS 13 320 (pump speed: 70 %; sonication: 10 seconds; run-length: 30 seconds). For LOI, 5 g of each sample was placed in a Carbolite furnace CWF 1300 (550°C for 12 hours).

The soils at RFF are classified as Arenosols (IUSS Working Group WRB, 2015) with weak horizonisation. An Ap loamy-sand horizon (82% sand, 16% silt, 2% clay) thickens from 30 to 75 cm and increases in LOI content from 3.65 to 3.91% from summit to toeslope, respectively. Despite being subject to arable practices for over 150 years, the presence of a 30 cm Ap horizon may be explained in part by the incorporation of mineral matter with the remaining organic material after harvest, although further isotopic work is required to verify this for RFF. This Ap horizon is underlain by a 5 cm fluvial pebble-bed, typical of the Bunter pebble-beds found in the vicinity (Ambrose et al., 2014). An undifferentiated, weakly-consolidated subsoil steadily grades into saprolitic, moderately-consolidated sandstone. The soils at CW are classified as Arenosols (IUSS Working Group WRB, 2015). Similar to RFF, there is little evidence for horizonisation down the profile at CW. A thin (<5 cm) litter-fermentation-humus layer overlays an undifferentiated, weakly-consolidated, sandy subsoil (94% sand, 5% silt, 1% clay) and grades into moderately-consolidated saprolitic sandstone. The sandy composition of these soils suggests that proglacial outwash deposits have not contributed to the soils of the study sites and that, instead, the soils are largely residual.

2.3 Lifespan analysis at Rufford Forest Farm

To provide an insight into the sustainability of the soil profiles at RFF under arable agriculture, in terms of the balance of erosion and formation, a first-order lifespan model was employed. Calculating the sustainability of a net-eroding soil in first-order terms has been attempted in the past (Elwell and Stocking, 1984; Sparovek and Schnug, 2001; Montgomery, 2007; Medeiros et al. 2016). Early models (Stocking and Pain, 1983), however, did not account for mass inputs into the soil system, such as that derived from bedrock weathering. In this study, this omission was addressed by using soil formation rates empirically measured at RFF. Furthermore, in previous models, the solum thickness used to calculate the soil lifespan is not universally consistent. Some authors constrain the lifespan by the minimum depth required for primary production (Stocking and Pain, 1983; Elwell and Stocking, 1984). Notwithstanding the fact that this soil threshold depth will, in part, be crop-dependent, soils that fall below this threshold may still be able to fulfil some of the ecosystem services, such as the sequestration of carbon. To address this here, two lifespan (L) scenarios were calculated, both of which are based on the continuation of contemporary arable agriculture. The first referred to the expected lifespan of the current A horizon (D = 30 cm across the catena). At the toeslope, an additional lifespan was calculated to account for the greater depth (75 cm) of the A horizon. Here, we did not account for any transformation of subsoil into topsoil, which could occur if erosion rates are sufficiently low, nor did we account for any allochthonous inputs to the profile such as aeolian additions and organic amendments. The second estimated the time until the underlying parent material is exposed. Here, the observed depth to the soil-saprolite interface at each catena position was employed.

Both lifespan scenarios were calculated for summit, shoulder, backslope and toeslope catena positions. Three different erosion rates (E) were applied. First, a mean annual erosion rate of $1.19 \text{ mm year}^{-1}$ was used based on ^{137}Cs -based data ($n = 103$) measured by Quine and Walling (1991) at RFF. This mean value represents all erosion processes, including water-based and tillage-based erosion. Two additional lifespans were calculated using rates from the 5th and 95th percentiles of this dataset ($0.19 \text{ mm year}^{-1}$ and 2.2 mm year^{-1} , respectively). It should be acknowledged here that the rates of soil formation represent timescales four orders of magnitude greater than those of soil erosion. However, if lifespans are to provide an insight into the sustainability of the soil profiles at RFF, the soil erosion rates must represent those from contemporary arable agriculture.

The soil formation rates, as empirically measured in this paper, were then plotted to derive the soil production function P ; Eq. (2):

$$P = W e^{\left(\frac{-h}{\gamma}\right)} \quad (2)$$

where W is the production rate at zero soil thickness (h) and γ is a parameter that determines the thickness of soil when soil formation falls off by $1/e$. The data for both the production rate (P) and the thickness of the soil (h) was used to calculate W and gamma using least squares regression. In this study, γ was calculated as being 2.26 m, which is substantially greater than that previously reported (e.g. Heimsath, 1997). It was therefore concluded that soil formation rates at RFF are relatively insensitive to changes in soil thickness. As a result, constant soil formation rates (F) for each catena position, together with two additional rates representing upper and lower standard deviations, were used to calculate soil lifespans. Furthermore, the expected increase in soil formation rates as a result of soil thinning were captured within these upper and lower uncertainties. Soil lifespans were thus calculated using Eq. (3):

$$L = \frac{D}{E - F} \quad (3)$$

where D is depth in mm, E is gross annual soil erosion rate in mm year⁻¹ and F is gross annual soil formation rate in mm year⁻¹.

3.0 Results and Discussion

5 3.1 Soil Formation Rates

Soil formation rates calculated from measured ¹⁰Be concentrations at RFF range from 0.026 mm year⁻¹ to 0.084 mm year⁻¹, with the mean soil formation rate being 0.048 ± 0.008 mm year⁻¹ (Table 1). At CW, soil formation rates range from 0.053 mm year⁻¹ to 0.096 mm year⁻¹, with the mean soil formation rate being 0.070 ± 0.010 mm year⁻¹, which is 0.022 mm year⁻¹ greater than that at RFF. These rates indicate declining soil formation rates with increasing soil thickness (Fig. 2—3). In accordance with geomorphological theory (Conacher and Dalrymple, 1977; King et al., 1983; Pennock, 2003; Schaeztl, 2013), soils are thinner on the slope convexities and the steepest gradients where surface erosion is considered most prevalent. In contrast, soil thicknesses are greater at the summit where surface erosion has been less extensive and the toeslope zone where sediment is deposited. In RFF, the fastest soil formation rates were found on the backslope where soils are thinnest. These results are consistent with many theorized mechanisms that demonstrate how parent material overlain by shallower soils is more affected by diurnal thermal stresses, contact with water and physical disturbance which can together proliferate physical and chemical weathering processes and thus the conversion of saprolite into soil. Conversely, it was found the slowest formation rates were associated with the deepest soils at the summit where the increasing thickness of the soil mantle buffers the parent material from any subaerial factors that may otherwise proliferate weathering (Carson and Kirby, 1972; Cox et al., 1980; Dietrich et al., 1995; Minasny and McBratney, 1999; Wilkinson and Humphreys, 2005). At CW, the difference in soil thickness between eroding and non-eroding zones is less pronounced. On the shoulder and backslope positions, where soils are thinnest, the soil formation rates were 0.03 mm year⁻¹ faster than summit and toeslope positions.

[Table 1]

5

10

15

20

25

30




Figure 2: Soil formation rates and the depths to saprolite for the four sampling positions along the catena transects at Rufford Forest Farm (blue; n = 4) and Comer Woodland (green; n = 4). The error bars represent one sigma uncertainties. At RFF, two ^{10}Be concentrations down the same depth profile have been used in the coSOILcal model to derive a 'best fit' soil formation rate. Depth here refers to that for the midpoint (between the top and bottom) of the sample.

10

15

5

10

15

20

25




Figure 3: Soil formation rates against sampling depth for Rufford Forest Farm (blue; n = 8) and Comer Woodland (green; n = 4). Depth here refers to that for the midpoint (between the top and bottom) of the sample.

5

- 10 Comparing data between RFF and CW demonstrates that there are other factors besides soil thickness that govern soil formation rates. For example, at the shoulder the soil thickness at CW is greater by 25 cm than that at RFF which would suggest slower formation rates. Instead soil formation rates are faster by $0.038 \text{ mm year}^{-1}$ at CW. One possible explanation is the petrographic composition of the parent material and the susceptibility of that parent material to weathering. Whilst both RFF and CW are underlain by sandstone, the bedrock at RFF is fluvially-derived whereas that at CW is aeolian-derived.
- 15 Petrological studies on fluvially-derived sandstone report a greater concentration of cementing clays in the matrix material which ultimately reduces the porosity and decreases its susceptibility to particle detachment, leading to slower soil formation rates (Wakatsuki et al., 2005; Mareschal et al., 2015).

In studies where cosmogenic methodologies have not been applied, it has been found that land use regime can promote or retard rates of bedrock weathering. Humphreys (1994) found that root channels and mesofaunal pedotubules in both the topsoil and subsoil can enhance the surface to bedrock hydrological connectivity. Similarly, Dong et al. (2018) demonstrated how an interconnected network of ecohydrologic interactions controls the supply and transport of acid to the bedrock. When a greater proportion of root mass was distributed in the uppermost horizons of the soil profile, CO₂ was predominantly emitted as gas whereas when roots were distributed in the subsoil, more CO₂ moved downwards to increase acid production and enhance chemical weathering. Other work has sought to identify the mechanisms that affect the thermal regime of soil profiles and the consequential impacts on the weathering susceptibility of the parent material (Ahnert, 1967; Minasny and McBratney, 1999). At CW, the roots are deeper than those found observed at RFF and this is likely to proliferate weathering processes. However, given the fact that the ¹⁰Be derived soil formation rates are millennial scale averages, it is unlikely that relatively recent (decadal-centennial) variances in the site's land use regime would be captured in the isotopic data (Darvill et al., 2013).

3.2 Derived soil formation rates in reference to the global inventory

Figure 4 compares soil formation rates for the study sites to an inventory of soil formation rates extracted from the published literature (n = 252; Fig. 4a; Supplementary Table 2). The median soil formation rate in this study (0.051 mm year⁻¹) is 0.028 mm year⁻¹ faster than that of the mantled inventory, a statistically significant difference (U test; P < 0.05). However, this global inventory comprises studies conducted on a range of geologies and climates, which are both influences on bedrock weathering rates.

Isolating the data from temperate climates (n = 187; Fig. 4b) presents a median soil formation rate of 0.035 mm year⁻¹, which is 0.016 mm year⁻¹ slower than that measured for RFF and CW, although there is no statistically significant difference between those data and those we have measured at the UK study sites presented in this paper (U test; P > 0.05). It is likely that the inventory's median soil formation rate for temperate climates is slower as 44% of the temperate-based data has been collected from regions that have lower mean annual precipitation than RFF and CW which can lead to less weathering activity at the parent material (Heimsath et al., 2001; Heimsath et al., 2005; Dixon et al., 2009; Heimsath et al., 2012).

Isolating the sandstone-derived data from the inventory (n = 57; Fig. 4c) presents a median soil formation rate of 0.045 mm year⁻¹ which is 0.006 mm year⁻¹ slower than that measured for RFF and CW, although there is no statistically significant difference (U test; P > 0.05). Although the sandstone-derived data were derived from the global soil-mantled database, all data stem from sites in temperate climates which reduces the influence that climate may have otherwise had in this analysis on lithology. We suggest that faster formation rates at RFF and CW may be explained by the fact that the specific varieties of sandstone at these study sites are generally more susceptible to weathering than those within the sandstone-based

inventory, of which the dominant form is the greywacke, characterised by a hard, fine-grained argillaceous matrix, with greater resistance to weathering (Cummins et al., 1962). Although there has been substantial work on the susceptibilities of major geological rock types to weathering (Stockmann et al., 2014; Wilson et al., 2017), we do not know of any study which seeks to identify whether the susceptibility of specific varieties of sandstone have an influence on soil formation rates.

- 5 The only other study to measure soil formation rates in the UK is that of Riggins et al. (2011) where rates were derived for Bodmin Moor, Cornwall (n = 5; Fig. 4d). In that study, the median soil formation rate was $0.015 \text{ mm year}^{-1}$, which is $0.036 \text{ mm year}^{-1}$ slower than that for RFF and CW and statistically significant (U test; $P < 0.05$), despite the fact that Bodmin Moor receives about 300 mm more precipitation per year than the sites in this study which should increase soil formation rates (Riggins et al., 2011). This is explained by the parent material at Bodmin Moor (coarse-grained granite) being generally less
10 prone to weathering than the varieties of sandstone evident at RFF and CW (Portenga and Bierman, 2011).

15

20

25

30

5

10

15

20

25

30




Figure 4: Soil formation rates from a globally compiled inventory (grey circles) and from this study at Rufford Forest Farm (blue triangles) and Comer Woodland (green diamonds) plotted against sampling depth. The depth here refers to that for the midpoint (between the top and bottom) of the sample. Rates in grey are from (a) the total mantled inventory (n = 252); (b) studies from temperate climates (n = 187); (c) studies on sandstone geology (n = 57); and (d) the UK, exclusively from Riggins et al. (2011) (n = 5). Error bars indicate the standard error.

10

3.3 Lifespan analysis at Rufford Forest Farm

Based on a mean annual erosion rate of $1.19 \text{ mm year}^{-1}$ under arable agriculture, the lifespans of the A horizon across the catena at RFF range between 258—272 years (Figure 5). This range expands to 138—3000 years when the 5th and 95th

percentile soil erosion rates are applied. However, further examination of the A horizon from cores extracted down the catena suggest that the toeslope is in a phase of aggradation rather than thinning. This is supported by the fact that the depth of the Ap horizon at the toeslope is 75 cm, whereas it is 30 cm on all other observed landscape positions. Moreover, comprised within the upper stratigraphy of the soil profile down the catena is the Bunter Pebble Bed which can be found at approximately 30 cm on summit, shoulder and backslope positions but 70 cm at the toeslope. The depth to which this pebble bed occurs at the toeslope suggests that either colluviation has occurred or is still occurring. In a scenario where colluviation is no longer active, the lifespan of this 75 cm A horizon is finite and ranges from 347 – 5245 years, but lifespans here could be longer or indefinite if colluviation continues. This demonstrates the difficulty of calculating lifespans using soil formation rates derived from bedrock alone and not from other system inflows of soil mass such as that from colluviation and soil carbon additions.

Soil lifespans indicating the time until the exposure of the parent material span between 407—1334 years. The range of these lifespans can be explained by the fact that unlike scenario one, where a constant A horizon thickness of 30 cm was applied across the catena, the soil thickness applied here is the depth to the soil-saprolite interface measured at each catena position (see Table 1). Applying upper and lower confidence intervals in the soil formation term and the 5th and 95th percentiles in the soil erosion term further widens the breadth of lifespans to 212—9688 years. The shortest lifespans are found on the backslope where bedrock exposure is expected to occur between 212—4500 years. In contrast, the greatest lifespans are found at the summit where soil thickness is 155 cm (713—9688 years). Although soil formation rates are greater at the toeslope, the depth to bedrock is 40 cm greater at the summit and, as a result, longer durations are required for bedrock to become exposed at this position. The soil detached and transported from the backslope is expected, in part, to continue to be a contributory source of the colluvium observed at the toeslope. Although the growth of soil profiles due to colluvium is not considered in the lifespan equation, it suggests that lifespans at the toeslope may either be longer than the calculated maximum of 8042 years or indefinite.

25




Figure 5: First-order soil lifespans calculated at four catena positions at Rufford Forest Farm for Scenario 1 (the time until the erosion of a 30 cm A horizon) and Scenario 2 (the time until bedrock exposure). The centre diagram indicates the thickness of the A horizon (dark brown), the subsoil (light brown) and the depth to the soil-saprolite interface (bricks). Red diamonds denote lifespans calculated using a mean annual soil erosion rate of $1.19 \text{ mm year}^{-1}$ from Quine and Walling (1991) and soil formation rates from this study. Black dots denote the minimum and maximum lifespans calculated using the 5th and 95th percentile of the soil erosion dataset and the one sigma uncertainties in the soil formation dataset.

5

10

15

The first-order lifespans presented here are based on a number of assumptions. Notwithstanding the fact that the land management regime may change within the cited time spans altering the protection the soils receive from wind and water, the erosion rates employed neither reflects the increase in the erodibility of subsoil horizons, characterised by a relatively weaker soil structure (Tanner et al., 2018) nor the potential role that the Bunter Pebble Bed may play in armouring the soil surface in the future. Moreover, they do not reflect the expected shift in erosivity, commensurate with more intense precipitation events (Burt et al., 2015). Acknowledging these factors, the lifespans presented here are likely to be overestimated. However, the fate of eroded soil upslope may contribute to the up-building of soil profiles in downslope concavities, extending the lifespans in the colluvial zone. In this respect the lifespans presented here, particularly those for the toeslope, are likely to be underestimated.

10 4.0 Conclusions

We have presented the first isotopically-derived rates of soil formation for soils currently supporting arable agriculture. Rates derived for two UK catena sequences using cosmogenic radionuclide analysis range from 0.026 mm year⁻¹ to 0.084 mm year⁻¹, with mean rates being 0.048 ± 0.008 mm year⁻¹ and 0.070 ± 0.010 mm year⁻¹ for Rufford Forest Farm and Comer Wood, respectively. By combining soil formation rates from Rufford Forest Farm with soil erosion rates derived from a prior isotopic study in a first-order lifespan model, we estimate that in a worst-case scenario the soil that currently comprises the A horizon on the backslope may be eroded in 138 years and bedrock exposure may occur in 212 years. Assessing gross soil erosion with measured rates of soil formation is important because soils that support arable agriculture are under threat from accelerated soil erosion. We have therefore shown that both the derivation and application of soil formation rates must become a fundamental component in future discussions of soil sustainability.

20

This work also represents the second of all isotopic studies of soil formation in the UK and therefore a significant contribution to our knowledge of pedogenesis. Soil formation rates were found to fall within the range of those previously published for soils in temperate climates and on sandstone lithologies, but were found to be significantly greater than those measured previously at Bodmin Moor. This is explained by the fact that the parent material at Bodmin Moor is a coarse-grained granite and therefore less susceptible to weathering than the sandstone materials underlying Rufford Forest Farm and Comer Wood. Such petrographic controls may also explain the greater rates of soil formation at Comer Wood where the sandstone matrix is largely devoid of the cementing agents present at Rufford Forest Farm and, therefore, more susceptible to particle detachment during physical and chemical weathering. Given that petrographic variability has not been thoroughly investigated in pedogenesis work, greater investment is warranted to better understand how the geochemical composition of the parent material governs the rates of soil formation.

30

Data availability

The authors declare no restrictions on the availability of materials or information.

Supplement link

5

Author contribution

D. L. E., J. N. Q., A. M. T. and J. A. C. D. designed research. D. L. E. and A. M. T. conducted sampling. D. L. E. and A. R. conducted laboratory work and analysed results. D. L. E. prepared the manuscript with contributions from all co-authors.

10 **Competing interests**

J. N. Q is a member of the editorial board of the journal.

Acknowledgments

The authors wish to thank Mr Annis (National Trust) for permission to carry out fieldwork on Comer Wood and Mr and Mrs
15 King (TAG Farming) for permission to carry out fieldwork on Rufford Forest Farm. We thank Vassil Karloukovski for
assistance in surveying and Andrew Binley, Paul McLachlan, Jonathan Riley, Carl Horabin and the BGS Dando Drilling Rig
Team for the acquisition of samples. We also wish to thank Allan Davidson, Ángel Rodés, Derek Fabel at the NERC
Cosmogenic Isotope Analysis Facility for preparing samples for AMS and their subsequent assistance in data analysis. We
thank Tim Quine for sharing multiple datasets from fieldwork conducted at Rufford Forest Farm. Finally, we wish to thank
20 the anonymous referees and Dan Morgan for their very constructive feedback on the manuscript. This work was partly
supported by BBSRC and NERC through a Soils Training and Research Studentships (STARS) (Grant number:
NE/M009106/1) and partly by a NERC research grant (Grant number: CIAF 9179/1017). STARS is a consortium consisting
of Bangor University, British Geological Survey, Centre for Ecology and Hydrology, Cranfield University, James Hutton
Institute, Lancaster University, Rothamsted Research and the University of Nottingham.

References

- Ackerer, J., Chabaux, F., Van der Woerd, J., Viville, D., Pelt, E., Kali, E., Lerouge, C., Ackerer, P., di Chiara Roupert, R. and Négrel, P. 'Regolith evolution on the millennial timescale from combined U-Th-Ra isotopes and in situ cosmogenic ^{10}Be analysis in a weathering profile (Strengbach catchment, France)', *Earth and Planetary Science Letters*, 453, pp. 33-43, 2016
- Ahnert, F. 'The role of the equilibrium concept in the interpretation of landforms of fluvial erosion and deposition', in Macar, P. (ed.) *L'évolution des versants*. Liege: Universite de Liege, pp. 23-41, 1967.
- Alexander, E. B. 'Rates of Soil Formation: Implications for Soil-Loss Tolerance', *Soil Science*, 145(1), pp. 37-45.
- Ambrose, K., Hough, E. and Smith, N. J. P. (2014) Lithostratigraphy of the Sherwood Sandstone Group of England, Wales and south-west Scotland. Available at: <http://nora.nerc.ac.uk/id/eprint/507530> (Accessed: 30 September 2018), 1988.
- Amundson, R., Berhe, A. A., Hopmans, J. W., Olson, C., Sztein, A. E. and Sparks, D. L. 'Soil and human security in the 21st century', *Science*, 348(6235), doi: 10.1126/science.1261071, 2015.
- Balco, G. Converting Al and Be isotope ratio measurements to nuclide concentrations in quartz. Available at: http://hess.ess.washington.edu/math/docs/common/ams_data_reduction/ (Accessed: 30 September 2018), 2006.
- Balco, G., Stone, J. O., Lifton, N. A. and Dunai, T. J. 'A complete and easily accessible means of calculating surface exposure ages or erosion rates from ^{10}Be and ^{26}Al measurements', *Quaternary Geochronology*, 3, pp. 174-195, 2008.
- Borrelli, P., Robinson, D. A., Fleischer, L. R., Lugato, E., Ballabio, C., Alewell, C., Meusburger, K., Modugno, S., Schütt, B., Ferro, V., Bagarello, V., Van Oost, K., Montanarella, L. and Panagos, P. 'An assessment of the global impact of 21st century land use change on soil erosion', *Nature Communications*, 8, pp. 1-13, 2017.
- Bryan, W. H. and Teakle, L. J. H. 'Pedogenic Inertia: a Concept in Soil Science', *Nature*, 164, p. 969, 1949.
- Burke, B. C., Heimsath, A. and White, A. F. 'Coupling chemical weathering with soil production across soil-mantled landscapes', *Earth Surface Processes and Landforms*, 32, pp. 853-873, 2007.

- Burt, T., Boardman, J., Foster, I. and Howden, N. 'More rain, less soil: long-term changes in rainfall intensity with climate change', *Earth Surface Processes and Landforms*, 41(4), pp. 563-566, 2015.
- Carson, M. A. and Kirkby, M. J. *Hillslope form and process*. Cambridge: Cambridge University Press, 1972.
- 5
- Conacher, A. J. and Dalrymple, J. B. 'The nine unit landsurface model: an approach to pedogeomorphic research', *Geoderma*, 18, pp. 3-154, 1977.
- Corbett, L. B., Bierman, P. and Rood, D. H. 'An approach for optimizing in situ cosmogenic ¹⁰Be sample preparation',
- 10 *Quaternary Geochronology*, 33, pp. 24-34, 2016.
- Cox, N. J. 'On the relationship between bedrock lowering and regolith thickness', *Earth Surface Processes*, 5, pp. 271-274, 1980.
- 15 Cummins, W. A. 'The Greywacke problem', *Geological Journal*, 3(1), pp. 51-72, 1962.
- Darvill, C. M. 'Cosmogenic nuclide analysis', in Clarke, L. (ed.) *Geomorphological Techniques*. London: British Society for Geomorphology, ch. 4, sec. 2.10, 2013.
- 20 Dietrich, W. E., Reiss, R., Hsu, M. and Montgomery, D. R. 'A process-based model for colluvial soil depth and shallow landsliding using digital elevation data', *Hydrological Processes*, 9, pp. 383-400, 1995.
- Dixon, J. I., Heimsath, A. M. and Amundson, R. 'The critical role of climate and saprolite weathering in landscape evolution', *Earth Surface Processes and Landforms*, 34, pp. 1507-1521, 2009.
- 25
- Dokuchaev, V. V. *Mapping the Russian Soils*. Russia: Imperial University of St. Petersburg, 1879.
- Dong, X., Cohen, M. J., Martin, J. B., McLaughlin, D. L., Murray, A. B., Ward, N. D., Flint, M. K. and Heffernan, J. B. 'Ecohydrologic processes and soil thickness feedbacks control limestone-weathering rates in a Karst landscape'. To be
- 30 published in *Chemical Geology* [Preprint]. Available at: <https://www.sciencedirect.com/science/article/pii/S0009254118302419> (Accessed: 30 September 2018).
- Elwell, H. A. and Stocking, M. A. 'Estimating soil life-span for conservation planning', *Tropical Agriculture*, 61(2), pp. 148-150, 1984.

Evans, D. L. Telephone conversation with Mike Annis, 8 October, 2018.

5 F.A.O. and I.T.P.S. Status of the World's Soil Resources (SWSR) Main Report. Rome, Italy: Food and Agriculture Organisation of the United Nations and Intergovernmental Technical Panel on Soils, 2015.

Fifield, L. K. (1999) 'Accelerator mass spectrometry and its application', *Reports on Progress in Physics*, 62, pp. 1223-1274.

Gosse, J. C. and Phillips, F. M.: 'Terrestrial in situ cosmogenic nuclides: theory and application', *Quaternary Science Reviews*, 20(14), pp. 1475-1560, 2001.

10

Govers, G., Merckx, R., van Wesemael, B. and Van Oost, K.: 'Soil conservation in the 21st century: why we need smart agricultural intensification', *Soil*, 3(1), pp. 45-59, 2017.

15 Govers, G., Quine, T. A., Desmet, P. J. J. and Walling, D. E.: 'The relative contribution of soil tillage and overland flow erosion to soil redistribution on agricultural land', *Earth Surface Processes and Landforms*, 21(10), pp. 929-946, 1996.

Hancock, G. R., Wells, T., Martinez, C. and Dever, C.: 'Soil erosion and tolerable soil loss: insights into erosion rates for a well-managed grassland catchments', *Geoderma*, 237, pp. 256-265, 2015.

20 Heimsath, A. M.: 'Limits of Soil Production?', *Science*, 343, pp. 617-618, 2014.

Heimsath, A. M.: 'Eroding the land: steady-state and stochastic rates and processes through a cosmogenic lens', *Geological Society of America*, 415, pp. 111-129, 2006.

25 Heimsath, A. M. and Burke, B. C.: 'The impact of local geochemical variability on quantifying hillslope soil production and chemical weathering', *Geomorphology*, 200, pp. 75-88, 2013.

Heimsath, A. M., Dietrich, W. E., Nishiizumi, K. and Finkal, R. C.: 'The Soil Production Function and Landscape Equilibrium', *Nature*, 388, pp. 358-361, 1997.

30

Heimsath, A. M., Dietrich, W. E., Nishiizumi, K. and Finkel, R. C.: 'Stochastic processes of soil production and transport: erosion rates, topographic variation and cosmogenic nuclides in the Oregon Coast Range', *Earth Surface Processes and Landforms*, 26, pp. 531-532, 2001.

- Heimsath, A. M., Furbish, D. J. and Dietrich, W. E.: 'The illusion of diffusion: field evidence for depth-dependent sediment transport', *Geology*, 33(12), pp. 949-952, 2005.
- 5 Heimsath, A. M., DiBiase, R. A. and Whipple, K. X.: 'Soil production limits and the transition to bedrock-dominated landscapes', *Nature Geoscience*, 5(3), pp. 210-214, 2012.
- Humphreys, G. S.: 'Bioturbation, biofabrics and the biomantle: an example from the Sydney Basin', in Ringrose-Voase and Humphreys, G. S (eds) *Soil Micromorphology: studies in management and genesis*. Amsterdam: Elsevier, pp. 421-436, 1994.
- 10 IUSS Working Group WRB.: *World Reference Base for Soil Resources 2014, update 2015 International soil classification system for naming soils and creating legends for soil maps*. World Soil Resources Reports No. 106. FAO: Rome, 2015.
- Jenny, H.: *Factors of Soil Formation: A System of Quantitative Pedology*. New York: McGraw-Hill, 1941.
- 15 King, G. J., Acton, D. F. and St. Arnaud, R. J.: 'Soil-landscape analysis in relation to soil redistribution and mapping at a site within the Weyburn association', *Canadian Journal of Soil Science*, 63, pp. 657-670, 1983.
- Kohl, C. P. and Nishiizumi, K.: 'Chemical isolation of quartz for measurement of in-situ produced cosmogenic nuclides', *Geochimica et Cosmochimica Acta*, 56(9), pp. 3583-3587, 1992.
- 20 Lal, D.: 'Cosmic ray labelling of erosion surfaces: in situ nuclide production rates and erosion models', *Earth and Planetary Science Letters*, 104, pp. 424-439, 1991.
- 25 Mareschal, L., Turpault, M. P. and Ranger, J.: 'Effect of granite crystal grain size on soil properties and pedogenic processes along a lithosequence', *Geoderma*, 249, pp. 12-20, 2015.
- Medeiros, G. O. R., Giarolla, A., Sampalo, G. and Marinho, M. A.: 'Diagnosis of the Accelerated Soil Erosion in São Paulo State (Brazil) by the Soil Lifetime Index Methodology', *Revista Brasileira de Ciência do Solo*, 40, pp. 1-15, 2016.
- 30 Minasny, B. and McBratney, A. B.: 'A rudimentary mechanistic model for soil production and landscape development', *Geoderma*, 90, pp. 3-21, 1999.

- Minasny, B., Finke, P., Stockmann, U., Vanwalleggem, T. and Bratney, A. B.: 'Resolving the integral connection between pedogenesis and landscape evolution', *Earth-Science Reviews*, 150, pp. 102-120, 2015.
- Montgomery, D. R.: 'Soil erosion and agricultural sustainability', *Proceedings of the National Academy of Sciences of the United States of America*, 104(33), pp. 13268-13272, 2007.
- Panagos, P., Imeson, A., Meusburger, K., Borrelli, P., Poesen, J. and Alewell, C.: 'Soil conservation in Europe: Wish or Reality?', *Land Degradation and Development*, 27(6), pp. 1547-1551, 2016.
- Pennock, D. J.: 'Terrain attributes, landform segmentation, and soil redistribution', *Soil Tillage Research*, 69, pp. 15-26, 2003.
- Phillips, J. D.: 'The convenient fiction of steady-state soil thickness', *Geoderma* 156:389-398, 2010.
- Phillips, F. M., Argento, D. C., Balco, G., Caffee, M. W., Clem, J., Dunai, T. J., Finkel, R., Goehring, B., Gosse, J. C., Hudson, A. M., Jull, A. J. T., Kelly, M. A., Kurz, M., Lal, D., Lifton, N., Marrero, S. M., Nishiizumi, K., Reedy, R. C., Schaefer, J., Stone, J. O. H., Swanson, T. and Zreda, M. G.: 'The CRONUS-Earth Project: A synthesis', *Quaternary Geochronology*, 31, pp. 119-154, 2016.
- Pimentel, D., Harvey, C., Resosudarmo, P., Sinclair, K., Kurz, D., McNair, M., Crist, S., Shpritz, L., Fitton, L., Saffouri, R. and Blair, R.: 'Environmental and Economic Costs of Soil Erosion and Conservation Benefits', *Science*, 267(5201), pp. 1117-1123, 1995.
- Portenga, E. W. and Bierman, P. R.: 'Understanding Earth's eroding surface with ^{10}Be ', *GSA Today*, 21(8), pp. 4-10, 2011.
- Quine, T. A. and Walling, D. E.: 'Rates of soil erosion on arable fields in Britain: quantitative data from caesium-137 measurements', *Soil Use and Management*, 7(4), pp. 169-176, 1991.
- Quinton, J. N., Govers, G., Van Oost, K. and Bardgett, R. D.: 'The impact of agricultural soil erosion on biogeochemical cycling', *Nature Geosciences*, 3, pp. 311-314, 2010.
- Riggins, S. G., Anderson, R. S., Anderson, S. P. and Tye, A. M.: 'Solving a conundrum of a steady-state hilltop with variable soil depths and production rates, Bodmin Moor, UK', *Geomorphology*, 128, pp. 73-84, 2011.

- Rodés, Á., Pallàs, R., Braucher, R., Moreno, X., Masana, E. and Bourlés, D. L.: 'Effect of density uncertainties in cosmogenic ^{10}Be depth-profiles: Dating a cemented Pleistocene alluvial fan (Carboneras Fault, SE Iberia)', *Quaternary Geochronology*, 6, pp. 186-194, 2011.
- 5 Schaetzl, R. J.: 'Catenas and Soils', in Shroder, J. F. (ed.) *Treatise on Geomorphology*. San Diego, California: Academic Press, pp. 145-158, 2013.
- Sparovek, G. and Schnug, E.: 'Temporal Erosion-Induced Soil Degradation and Yield Loss', *Soil Science Society of America Journal*, 65, pp. 1479-1486, 2001.
- 10 Stocking, M. A. and Pain, A.: *Soil Life and the Minimum Soil Depth for Productive Yields: Developing a New Concept*. Norwich: University of East Anglia, School of Development Studies, 1983.
- Stockmann, U., Minasny, B. and McBratney, A. B.: 'How fast does soil grow?', *Geoderma*, 216, pp. 48-61, 2014.
- 15 Struck, M., Jansen, J. D., Fujioka, T., Codilean, A. T., Fink, D., Egholm, D. L., Fülöp, R., Wilcken, K. M. and Kotevski, S.: 'Soil production and transport on postorogenic desert hillslopes quantified with ^{10}Be and ^{26}Al ', *GSA Bulletin*, 130(5), pp. 1017-1040, 2018.
- 20 Tanner, S., Katra, I., Argaman and Ben-Hur, M.: 'Erodibility of waste (Loess) soils from construction sites under water and wind erosional forces', *Science of the Total Environment*, 616, pp. 1524-1532, 2018.
- Tugel, A. J., Herrick, J. E., Brown, J. R., Mausbach, M. J., Puckett, W. and Hipple, K.: 'Soil change, soil survey and natural resources decision making: a blueprint for action', *Soil Science Society of America Journal*, 69, pp. 738-747, 2005.
- 25 Turner, B. L., Hayes, P. E. and Laliberté, E.: 'A climosequence of chronosequences in southwestern Australia', *European Journal of Soil Science*, 69(1), pp. 69-86, 2018.
- Tye, A. M., Kemp, S. J., Lark, R. M. and Milodowski, A. E.: 'The role of peri-glacial active layer development in determining soil-regolith thickness across a Triassic sandstone outcrop in the UK', *Earth Surface Processes and Landforms*, 37(9), pp. 971-983, 2012.
- 30 Tye, A. M., Robinson, D. A. and Lark, R. M.: 'Gradual and anthropogenic soil change for fertility and carbon on marginal sandy soils', *Geoderma*, 207, pp. 35-48., 2013.

Wakatsuki, T., Tanaka, Y. and Matsukura, Y.: 'Soil slips on weathering-limited slopes underlain by coarse-grained granite or fine-grained gneiss near Seoul, Republic of Korea', *Catena*, 60(2), pp. 181-203, 2005.

- 5 Walling, D. E. and Quine, T. A.: 'The use of ^{137}Cs measurements to investigate soil erosion on arable fields in the UK: potential applications and limitations', *Journal of Soil Science*, 42, pp. 147-165, 1991.

Wilkinson, M. T. and Humphreys, G. S.: 'Exploring pedogenesis via nuclide-based soil production rates and OSL-based bioturbation rates', *Australian Journal of Soil Research*, 43, pp. 767-779, 2005.

10

Wilkinson, M. T., Chappell, J., Humphreys, G. S., Fifield, K., Smith, B., Hesse, P., Heimsath, A. M. and Ehlers, T. A.: 'Soil production in heath and forest, Blue Mountains, Australia: influence of lithology and palaeoclimate', *Earth Surface Processes and Landforms*, 30(8), pp. 923-934, 2005.

- 15 Wilson, S. G., Lambert, J., Nanzyo, M. and Dahlgren, R. A.: 'Soil genesis and mineralogy across a volcanic lithosequence', *Geoderma*, 285, pp. 301-312, 2017.

Zhao, T., Liu, W., Xu, Z., Liu, T., Xu, S., Cui, L. and Shi, C.: 'Cosmogenic nuclides (^{10}Be and ^{26}Al) erosion rate constraints in the Badain Jaran Desert, northwest China: implications for surface erosion mechanisms and landform evolution',

- 20 *Geosciences Journal*, pp. 1-10, doi: 10.1007/s1230, 2018.

25

30



Copernicus Publications

The Innovative Open Access Publisher

5

10

15

Table 1: ^{10}Be concentrations and calculated maximum soil formation rates for Rufford Forest Farm (RFF) and Comer Wood (CW)

Site	Catena Position	Elevation, m	Horizon Position	Depth, cm	^{10}Be atoms, g	Uncertainty of ^{10}Be atoms, g	^{10}Be production rate at surface, $\text{g}^{-1} \text{year}^{-1}$	Soil Formation Rates, (Best Fit) mm ka^{-1}	Uncertainty, mm ka^{-1}
RFF	Summit	98.7	A	150	35266	2364	4.63	30	29 - 33
RFF	Summit	98.7	B	203	22683	1586	4.63	26	24 - 28
RFF	Shoulder	99.3	A	53	54380	2030	4.63	38	36 - 41
RFF	Shoulder	99.3	B	100	30064	1850	4.63	38	36 - 40
RFF	Backslope	97.9	A	43	45603	1833	4.63	80	77 - 83
RFF	Backslope	97.9	B	93	28876	1661	4.63	84	77 - 88
RFF	Toeslope	95.7	A	108	32738	2006	4.62	49	46 - 53
RFF	Toeslope	95.7	B	160	25237	1562	4.62	36	34 - 39
CW	Summit	70.6	A	148	24507	1696	4.49	57	52 - 59
CW	Shoulder	65.3	A	78	24811	1333	4.46	96	90 - 99
CW	Backslope	58.9	A	78	31263	2035	4.42	73	69 - 78
CW	Toeslope	50.1	A	88	41276	1522	4.39	53	51 - 54

Horizon Position ‘A’ denotes the sample was taken at the soil-saprolite interface. Horizon Position ‘B’ denotes an additional sample was taken ~50cm below the interface from the same depth profile. The depth here refers to that for the midpoint (between the top and bottom) of the sample. The shielding correction was calculated as 1.0 (to 1 d.p) for all samples and ^{10}Be production rates are corrected for elevation and location (see Supplementary Table 1). All uncertainties are one standard deviation and are based on uncertainties in the measurement of ^{10}Be concentration as outlined in Rodés et al. (2011).

