



1 Targeting the soil quality and soil health concepts when aiming for the 2 United Nations Sustainable Development Goals and the EU Green 3 Deal

4 Antonello Bonfante¹, Angelo Basile¹, Johan Bouma²

5 ¹Institute for Mediterranean Agricultural and Forest Systems - CNR-ISAFOM, Ercolano, 80056, Italy

6 ²Em. Prof Soil Science, Wageningen University, the Netherlands

7 *Correspondence to:* Antonello Bonfante (antonello.bonfante@cnr.it)

8 **Abstract.** The soil quality and soil health concepts are widely used as soils receive more attention in the worldwide policy
9 arena. So far, however, the distinction between the two concepts is unclear and operational procedures for measurement are
10 still being developed. A proposal is made to focus soil health on actual soil conditions, as determined by a limited set of
11 indicators that reflect favourable rooting conditions. In addition, soil quality can express inherent soil conditions in a given
12 soil type (genoform) reflecting the effects of past and present soil management (expressed by various phenoforms). Soils
13 contribute to ecosystem services that, in turn, contribute to the UN Sustainable Development Goals and, more recently, to the
14 EU Green Deal. Relevant soil ecosystem services are biomass production (SDG2: zero hunger), providing clean water (SDG6);
15 climate mitigation by carbon capture and reduction of greenhouse gas emissions (SDG13: climate action) and biodiversity
16 preservation (SDG15: life on land). The use of simulation models for the soil-water-atmosphere-plant system is proposed as a
17 quantitative and reproducible procedure to derive single values for soil health and soil quality for current and future climate
18 conditions. Crop production parameters from the international: “yield-gap” program are used combined with soil-specific
19 parameters expressing the effects of phenoforms. These procedures focus on the ecosystem service: biomass production Other
20 ecosystem services are determined by soil-specific management to be based on experiences obtained in similar soils elsewhere
21 or by new research. A case study, covering three Italian soil series, illustrates the application of the proposed concepts, showing
22 that soil types (soil series) acted significantly different to effects of management also in their reaction to climate change.

23 1 Introduction

24 The soil receives increasing attention in the research and policy arena focusing on its capability to perform a number of
25 functions. The concepts of soil quality and soil health are often used to express this capability, but this is only meaningful
26 when these two concepts are clearly defined and can be established with operational and reproducible methods. So far, this
27 methodology has not been developed. Moreover, methods to assess soil health and soil quality derive their significance from
28 societal relevance in a broad ecosystem context as defined by the United Nations in 2015, in terms of seventeen Sustainable
29 Development Goals (<https://www.un.org/sustainabledevelopment-goals>) and by the 2019 Green Deal of the European Union



30 (<https://ec.europa.eu/info/strategy/european-green-deal>). In the United States, Soil health is supported by the policy arena and
31 is being studied by at least three Institutes: Cornell University, The National Soil Health Institute and the US Dept. of
32 Agriculture. The new research and innovation program of the European Union for the period 2021-2027, “Horizon Europa”
33 has defined five MISSION areas, among them: “Soil Health and Food”, recognizing the importance of soils for sustainable
34 development. Soils are now clearly on the international research agenda!

35 To allow operational use of the soil health concept, a clear measurement methodology is needed. So far, Cornell University
36 has proposed a method to measure soil health, defining a set of indicators and a procedure resulting in a number between 1
37 and 100 ranging from highly unhealthy to shinningly healthy. This procedure will be discussed in this paper. The term soil
38 health is attractive not only because of its analogy with human health that facilitates communication with the public but also,
39 and particularly, because soils are biologically active as are humans. The older term soil quality that has been used for decades
40 (e.g., Bünemann et al. 2018) has a more sterile character that could also apply to, e.g., nuts and bolts. According to some (e.g.,
41 USDA, 2019), soil health and soil quality have the same meaning. This, however, is not logical because why introduce a new
42 term when it has the same meaning as the old one? The objective of this article is to propose that both terms can be distinguished
43 allowing a useful distinction between actual versus inherent conditions. The proposed concepts have been illustrated in an
44 Italian case study.

45 **1.1 The soil quality concept**

46 Soil quality has been defined as: “*the capacity of a soil to function within ecosystem and land-use boundaries to sustain*
47 *biological productivity, maintain environmental quality and promote plant and animal health*” as quoted by Bünemann et al.
48 (2018) in a comprehensive review of more than 250 scientific papers covering soil quality. The authors conclude that, in
49 contrast to the quality of water, air, and nature, there still is no universally accepted method to measure soil quality. This is a
50 serious problem, limiting application in practice and in environmental rules and regulations.

51 **1.2 The soil health concept**

52 Soil health has been defined in the US as “*the continued capacity of the soil to function as a vital living ecosystem that sustains*
53 *plants, animals and humans*”. Indicators for soil health have been defined in the USA: 19 by Cornell University (Moebius-
54 Clune et al., 2017), 31 by the National Soil Health Institute (<http://soilhealthinstitute.org>) and 11 by the US Department of
55 Agriculture (USDA, 2019). How these indicators are combined into a single soil health parameter for a given soil is presented
56 by the Cornell protocol. Only three texture classes of soils are distinguished: coarse, medium and fine. For each texture class,
57 measurements for each indicator are assembled for soils at different locations in that particular texture class and a frequency
58 curve of values is constructed. Obviously, such curves become more diagnostic as more data become available. When placed
59 on the frequency curve, any new observation of the indicator will obtain a number between 0 and 100. This procedure is
60 repeated for every indicator and in the end all numbers will be averaged producing one characteristic number for soil health
61 for that particular soil, which is quite attractive for communication purposes. The frequency curve also allows the distinction



62 of a threshold frequency value above which the particular indicator exceeds a critical environmental threshold value, sometimes
63 defined by environmental laws and regulations. In their reporting red, orange, yellow and green colours are used to indicate
64 whether or not this occurs. A red label indicates that a given threshold is exceeded and that action is needed, possibly to be
65 based on favourable management experiences obtained elsewhere in soils of the same texture class or by new research. This
66 is attractive because it can directly result in management advice. In an example presented by Moebius-Clune et al. (2017) on
67 page 73, values for twelve indicators are presented, three of which with a red label: “surface hardness”, “aggregate stability”
68 and “active carbon content”, suggesting a need for corrective measures. But what does this imply for soil health? A soil is
69 unhealthy if only one or more indicators are red? And how to interpret an average value for all twelve quite different indicators
70 with different colours?

71 Also, a question can be raised about the large number of indicators for soil health in the three US systems. Why not primarily
72 consider demands by roots as they link plants with the soil? A number of conditions do not allow root growth: e.g., presence
73 of excessive amounts of chemical pollutants, salty soils (solonchack), alkaline soils (solonetz) and very acid soils with low pH
74 values. Soils with such properties are clearly unhealthy. Otherwise, roots require: (i) temperatures that allow growth; (ii) soil
75 structure that allows easy accessibility of the entire soil volume, allowing roots to reach their genetically determined depth;
76 (iii) adequate water, air and nutrient availability during the growing season; (iv) adequate infiltration rates of water at the soil
77 surface; and (v) adequate organic matter content and the associated biological activity that is essential for many soil functions,
78 including nutrient uptake by plants. These five parameters can be measured at a given time and place and the reports by
79 Moebius Clune, (2017) and USDA (2019) contain detailed descriptions of measurement methods.

80 Parameters to be measured at a given point in time should have a semi-permanent character to be diagnostic. Temperature and
81 nutrient status are quite variable, the latter high at the moment of fertilization and increasingly lower as the crop adsorbs
82 nutrients. Of course, this is different in nature areas where inherent nutrient contents are important to allow particular types of
83 vegetation to develop. However, nutrient deficiencies in agricultural soils can be rapidly corrected by fertilization and the
84 nutrient status, though essential for root growth, is therefore less suitable as a parameter in agricultural soils. Soil structure,
85 excluding a limited period after soil tillage, is more permanent and governs infiltration rates and soil water and air regimes as
86 a function of weather conditions and groundwater dynamics. Soil structure is therefore suitable as a parameter. Aggregate
87 stability is a measure for soil resistance to deformation but the method has been criticised as being unrepresentative (e.g.,
88 Baveye, 2020). The use of penetrometers may be more effective to measure mechanical resistance affecting root penetration.
89 Biological activity is subject to an even longer time span than compaction: increasing the organic matter content of soils may
90 take several years. The organic matter content is, therefore, a suitable parameter and many measurement methods are available,
91 including rapid methods applying proximal sensors. More detailed measurements of biodiversity have been defined by
92 Moebius-Clune, (2017) and for the LUCAS soil database (Orgiazzi et al., 2018), requiring laboratory measurements.

93 In conclusion, parameters for soil health for a given soil type at a given time and place, are: (i) soil structure, expressed by
94 descriptions in soil survey reports and supported by bulk density values and measured infiltration rates, and, possibly, by
95 penetrometer values, (ii) water and air regimes, as estimated by drainage class in soil survey reports, can be expressed indirectly



96 by the widely used but static parameter: “available water” defining the water content between two pressure heads, which,
97 however, poorly represent natural dynamic soil water and air regimes. Dynamic modeling presents more realistic data as will
98 be discussed later (e.g., Bouma, 2018, Bonfante et al., 2019) and (iii) organic matter contents.
99 Nevertheless, the procedure based on the three parameters mentioned above produces three separate values. Back, therefore,
100 to the definition of soil health that mentions “functioning of soils”, whereby soil contributions to biomass production is a key
101 function, among six other defined functions (EC, 2006). The degree by which biomass production is affected by the three
102 separate parameters remains unclear. An integrated approach is therefore needed and can be obtained by simulating the soil-
103 water-atmosphere-plant system.

104 **1.3 Still a role for soil quality?**

105 The soil health concept offers one fundamental problem. A sandy soil and a clay soil can both be healthy, but they obviously
106 have quite different water and nutrient regimes and use- potentials. But differences among soils can be expressed by the soil
107 quality concept when considering inherent properties of soils as expressed in soil classification, like texture, which is most
108 stable among all soil parameters (see also Moebius-Clune, 2017). In analogy with human health, soil health for a given soil at
109 a given time expresses the actual condition expressed by the parameters discussed above, just like a doctor assesses the health
110 of a patient at a given time applying a set of tests. As discussed, different health values can be found in the same soil type as a
111 function of past management, such as compaction, soil crusting followed by runoff, erosion, etc., as illustrated in the Italian
112 case study presented below. However, the range of such soil health values is characteristically different for every soil type and
113 can, therefore, function as a measure of soil quality. Droogers and Bouma (1997) have distinguished genoforms, expressing a
114 given soil classification, but also phenoforms of that particular genoform, as a function of different forms of management with
115 strong effects on soil functioning (e.g., erosion, compaction, crust formation). Traditional soil survey interpretations are based
116 on so-called “representative profiles” for each mapping unit on the soil map, based on permanent Taxonomic soil criteria,
117 correctly ignoring in the classification context the effects of management which would lead to highly variable classifications.
118 Different phenoforms of a given genoform can, however, function quite differently and this cannot be ignored when
119 considering soil health.

120 **1.4 Simulating the soil-water-atmosphere-plant system to obtain a single soil health value**

121 Application of simulation models of the soil-water-atmosphere-plant system can integrate the values of the parameters
122 mentioned above as they function as input data for the model, producing a single, integrated value for biomass production.
123 Many operational models are available (e.g., Reynolds et al., 2018; SWAP by Kroes et al., 2017; SWAP-WOFOST by Hack-
124 ten Broeke et al., 2019; ICASA by White et al., 2013; APSIM by Holzworth et al., 2018; Ma et al., 2012 and others). These
125 models use rooting depth, weather data and when the required hydraulic conductivity and moisture retention data are not
126 available, these values can be estimated with pedotransfer functions using texture (as defined by the soil type), % organic
127 matter and bulk density as input data, the soil health parameters identified above (Bouma, 1989; Van Looy et al., 2017). So



128 rather than have sets of separate parameters for soil health, an integrated expression is obtained by the model that directly
129 addresses a key soil function, which is its contribution to the ecosystem service “biomass production”. The term “contribution”
130 needs to be emphasized as “biomass production” is not determined by soils alone but by many other factors and, certainly, by
131 management. Applying modelling, an alternative procedure to define soil health was proposed by Bonfante et al. (2019) where
132 biomass production forms the starting point. Following the agronomic Yield Gap program (van Ittersum et al., 2013) yields
133 are calculated by simulation models of the soil-water-atmosphere-plant system: Y_p = potential production determined for a
134 representative crop considering radiation and temperature regimes in a given climate region, assuming that adequate water and
135 nutrients are available and pest and diseases do not occur. This is a science- based value that applies everywhere on earth and
136 yields unique, quantitative and reproducible data. Y_w is the water-limited yield, as Y_p , but expressing the effect of the actual
137 soil water regime under local conditions, and Y_a is the actual yield. The yield gap is $Y_w - Y_a$. These parameters of the Yield-
138 gap program can be applied to define soil health and soil quality parameters to be discussed in the next section but need to be
139 modified to express the specific impact of the soil.

140 Simulation modelling offers the possibility to express soil functioning, as mentioned in the definition of soil health, by an
141 interdisciplinary modelling effort with input by agronomists, hydrologists and climatologists, each providing basic data for the
142 models. This yields one number, based on an interdisciplinary analysis, which is preferable to a series of separate numbers for
143 soil parameters only as in the US systems. The soil science discipline presents the parameters, mentioned above, to the
144 interdisciplinary research team in the context of a well defined soil type that defines moisture regimes and rooting patterns.
145 This way, the soil type functions as a “carrier of information” or a “class-pedotransfer function” (Bouma, 1989).
146 Moreover, and more importantly, modelling is the only option to explore possible future effects of climate change on soil
147 health and soil quality, as will be demonstrated below. Procedures to define single soil health and soil quality parameters will
148 be presented in the materials and methods section of the paper.

149 **1.5 Targeting soil health and soil quality towards the SDGs and the Green Deal by focusing on ecosystem services**

150 The discussion of soil health and soil quality so far focused on the soil and the way it functions, mentioning goals such as
151 “biological productivity and environmental quality” (soil quality) and “vital soils that sustain plants, animals and humans”
152 (soil health). As mentioned in the introduction, since 2015, 193 countries have made a United Nations-initiated commitment
153 to reach seventeen Sustainable Development Goals (SDGs). The European Union launched its Green Deal in 2019. The soil
154 quality and soil health concepts are no meaningful goals by themselves and can obtain societal significance when linked to the
155 SDGs and the Green Deal. But there is no direct link, if only because soil management plays a key role in achieving the SDGs
156 and the goals of the Green Deal. The challenge for soil science is to explore ways in which healthy soils can contribute to
157 improving a number of key ecosystem services, that, in turn, contribute to the SDGs (e.g., Bouma, 2014; Keesstra, 2016). This
158 is important because SDGs and goals of the Green Deal are not only determined by ecosystem services but also by e.g., socio-
159 economic and political factors that are beyond control by sciences studying crop growth. Attention for the SDGs and the Green
160 deal implies attention for not only biomass production (SDG 2: zero hunger) but also for other ecosystem services that relate



161 directly to environmental quality, such as the quality of ground and surface water (SDG6: clean water and sanitation), carbon
162 sequestration and reduction of greenhouse-gas emissions for climate mitigation (SDG 13: climate action) and biodiversity
163 preservation (SDG 15: life on land). That is why the following definitions of soil health and soil quality are proposed:

- 164 • *Soil health is the actual capacity of a particular soil to function, contributing to ecosystem services*
- 165 • *Soil quality is the inherent capacity of a particular soil to function, contributing to ecosystem services.*

166 Both general definitions focus on soil contributions to ecosystem services that, in turn, contribute at this point in time to the
167 realization of the United Nations Sustainable Development Goals and the goals of the EU- Green Deal.

168 The four ecosystem services, mentioned above, have a different character. Biomass production (SDG 2) is governed by climatic
169 conditions and soil water regimes as characterized by modelling that yields quantitative and reproducible results for Y_p and
170 Y_w . Management plays a key role in determining Y_a , and the other ecosystem services and is characteristically different for
171 different soil types. Clean water (SDG 6) can e.g., be obtained by precision fertilization, minimizing nutrient leaching to the
172 groundwater, while combatting erosion can minimize surface water pollution. But there are, in contrast to Y_p or Y_w values
173 for biomass production, no theoretical reference values for this ecosystem service, only threshold values of water quality by
174 environmental laws and regulations. This also applies to carbon sequestration and reduction of greenhouse gas emissions (SDG
175 13) and to life on land (SDG 15) for which as yet no environmental laws have been introduced. Different soils in different
176 climate zones will offer different challenges and opportunities to be met by appropriate management.

177 **2 Materials and methods**

178 **2.1 The Soil–Water–Atmosphere–Plant (SWAP) model**

179 The Soil–Water–Atmosphere–Plant (SWAP) model (Kroes et al., 2017) was applied to solve the soil water balance during
180 maize cultivation under estimated climate change and soil % SOM scenarios of A_p horizons. SWAP is an integrated physically-
181 based simulation model of water, transport in the saturated–unsaturated zone in relation to crop growth. It assumes
182 unidimensional vertical flow processes and calculates the soil water flow through the Richards equation. Soil water retention
183 $\theta(h)$ and hydraulic conductivity $k(\theta)$ relationships as proposed by van Genuchten (1980) were applied. The unit gradient was
184 set as the condition at the bottom boundary. The upper boundary conditions of SWAP in agricultural crops are generally
185 described by the potential evapotranspiration ET_p , irrigation and daily precipitation. Potential evapotranspiration was then
186 partitioned into potential evaporation and potential transpiration according to the LAI evolution, following the approach of
187 Ritchie (1972). The water uptake and actual transpiration were modeled according to Feddes et al., (1978), where the actual
188 transpiration declines from its potential value through the parameter α , varying between 0 and 1 according to the soil water
189 potential.

190



191 2.2 Soil Health and Soil Quality indicators

192 Application of the soil-water-atmosphere-plant simulation model and the yield-gap parameters results in four characteristics:

193 (i) a measure for actual soil health of a given soil type in a given climate zone at a given time by the SH index:

$$194 \quad SH = (Y_w - \text{phenoform} / Y_w - \text{ref}) \cdot 100 \quad [1]$$

195 where Y_w -phenoform expresses Y_w for a given phenoform and Y_w -ref represents the undisturbed soil phenoform. This
196 index expresses the effect of the soil on the measured yield Y_a , a value that is affected by many other factors than the soil;

197 (ii) a measure for intrinsic soil quality (SQ_p) for a given soil type in a given climate zone, reflecting a characteristic range of
198 soil health values obtained at different locations (SHL) as a function of different types of management (SHM) applied to
199 that particular soil type, resulting in different phenoforms (p).

$$200 \quad SQ_p = f(\text{SHL}, \text{SHM}) \quad [2]$$

201 An example for three Italian soils will be shown later in figure 2.

202 (iii) a measure for intrinsic soil quality for all soils occurring in a given region in the same climate zone (SQ_r):

$$203 \quad SQ_r = (Y_w / Y_p) \cdot 100 \quad [3]$$

204 allowing comparisons among different soils in the region, with an option to again express effects of different phenoforms,
205 and:

206 (iv) a measure for intrinsic soil quality allowing comparisons among all soils in the world in different climate zones (SQ_w):

$$207 \quad SQ_w = (Y_w / Y_{\text{max}}) \cdot 100 \quad [4]$$

208 Values (ii) through (iv) can also be derived for different climate scenarios up to the year 2100, as reported by the
209 Intergovernmental Panel on Climate Change (IPCC, 2014).

210 2.3 An Italian case study

211 Six prominent Italian soil series were analysed to illustrate the proposed method to define soil health and soil quality. Because
212 of space constraints results of three soils will be discussed in this paper. The modeling process and the background of the IPCC
213 scenarios have been presented elsewhere (Bonfante et al., 2019, 2020; Bonfante and Bouma, 2015) and will be summarized
214 below.

215 The maize was simulated from May (emergence) to the end of August (harvest) with a peak of leaf area index (LAI) of 5.8 m²
216 m⁻². Finally, the above ground biomass (AGB) to determine the yield values (Y_w) was estimated using the normalized water
217 productivity concept (WP; 33 g m⁻² for maize; Steduto et al., 2012).

218 The simulation runs were performed for six selected soils using a future climate scenario of a site of southern Italy (Destra
219 Sele plain) where half of the analysed soils occur. The future climate scenarios were obtained by using the high resolution
220 regional climate model (RCM) COSMO-CLM (Rockel et al., 2008), with a configuration employing a spatial resolution of
221 0.0715° (about 8 km), which was optimized over the Italian area. The validations performed showed that model data agree



222 closely with different regional high-resolution observational datasets, in terms of both average temperature and precipitation
223 (Bucchignani et al., 2015) and in terms of extreme events (Zollo et al., 2015).

224 The severe Representative Concentration Pathway (RCP) 8.5 scenario was applied, based on the IPCC modelling approach to
225 generate greenhouse gas concentrations (Meinshausen et al., 2011).

226 The results were performed on reference climate RC (1971–2005) and RCP 8.5, the latter divided into three different time
227 periods (2010–2040, 2040–2070 and 2070–2100). Daily reference evapotranspiration (ET_0) was evaluated according to the
228 Hargreaves and Samani (1985) equation.

229 Under the RCP 8.5 scenario, the temperature in Destra Sele is expected to increase approximately two degrees Celsius,
230 respectively, every 30 years to 2100, starting from the RC. The differences in temperature between RC and the period 2070–
231 2100 showed an average increase in the minimum and maximum temperatures of about 6.2°C (for both min and max over the
232 year). The projected increase in temperatures produces an increase in the expected ET_0 . In particular, during the maize growing
233 season, an average increase of ET_0 of about 18% is expected until 2100 (Bonfante et al., 2020).

234 Simulations were run considering an undisturbed soil (the reference) and three phenofoms: two expressing degradation
235 phenomena (erosion and compacted plowpan) and one considering an increase of % OM in the first soil horizon (Ap), as a
236 possible result of combatting a low % OM due to soil degradation.

237 In particular:

238 (i) The compacted plowlayer was applied at 30 cm depth (10 cm of thickness) with the following physical characteristics:
239 $\theta_s=0.30 \text{ cm}^3\text{cm}^{-3}$, $n=1.12$, $\alpha=0.004$ and $k_0=2 \text{ cm day}^{-1}$, following the notation of van Genuchten (1980). Roots were
240 restricted to the upper 30 cm of the soil.

241 (ii) Erosion was simulated for the Ap horizon, reducing the upper soil layer to 20 cm. The maximum rooting depth was
242 assumed to be 60 cm (A+B horizons) with a higher root density in the Ap horizon.

243 (iii) The effect of the increase of SOM to 4% on the first soil horizon (Ap) on hydraulic properties was realized applying
244 the procedure developed and reported in Bonfante et al. (2020) on hydraulic properties measured in the lab.

245 2.3.1 Soil characteristics

246 The Italian soils are located in a plain in an alluvial environments, two in the Campania region (P5 and P6) and P4 in the
247 Lombardy Region. The physical properties of the three selected soils are presented in Table 1. Soil texture range from sandy
248 loam to loamy sand and organic matter contents in Ap horizons are relatively low, ranging from 1.4 to 2.6%, justifying runs
249 for hypothetical contents of 4%. Based on field observations, the rooting depth of maize was estimated to be 80 cm, implying
250 that not the only Ap horizon but also subsoil horizons contribute to the water supply to maize.

251 The soil hydraulic properties applied in the simulation runs, water retention, $\theta(h)$, and hydraulic conductivity, $k(\theta)$, curves
252 were measured in the laboratory. Undisturbed soil samples (volume $\approx 750 \text{ ml}$) were collected from all of the recognized
253 horizons of the six soil profiles. Samples were slowly saturated from the bottom and the saturated hydraulic conductivity
254 measured by a falling head permeameter (Reynolds et al., 2002). Then, both couples of θ -h and k - θ data were obtained by



255 means of the evaporation method (Arya, 2002) consisting of an automatically recorded of the pressure head at three different
256 depths and the weight of the sample during a 1-dimensional transient upward flow. From these information, i) the water
257 retention data θ -h were obtained applying an iterative method (Basile et al., 2012) and ii) the unsaturated hydraulic conductivity
258 data were obtained by applying the instantaneous profile method, requiring the spatio-temporal distribution of θ and h, namely
259 $\theta(z,t)$ and $h(z,t)$, being z and t the depth and time, respectively (Basile et al., 2006). Additional points of the dry branch of the
260 water retention curve were determined using a dewpoint potentiometer (WP4-T, Decagon Devices, Washington, USA).
261 The parameters of the van Genuchten-Mualem model for water retention and hydraulic conductivity functions were obtained
262 by fitting the experimental θ -h and k- θ data points (Van Genuchten, 1980).

263 **3 Results**

264 The emphasis in this paper will be on the application of the soil health and soil quality definitions presented above. Initially,
265 three adverse effects of management were considered: surface runoff caused by relatively low infiltration rates, erosion of 20
266 cm of topsoils (while soil classification remains the same), and formation of a plowpan at 30 cm depth (see Bonfante et al.,
267 2019). Results showed, however, that under prevailing current and future climate conditions surface runoff was negligible.
268 Results will therefore only be presented for phenoforms showing effects of erosion and the plowpan and for increased %OM,
269 as mentioned above.

270 **3.1 Water-limited yields (Y_w)**

271 Water-limited yields (Y_w) for four climate periods and three phenoforms for each soil are shown in Figure 1a for soil P4,
272 Figure 1b for soil P5, and Figure 1c for soil P6. Y_w values drop for all soils and their phenoforms in the period from the RC
273 to the 2070 -2100 climate scenario, particularly for climate scenarios beyond 2040, but due to relatively high standard
274 deviations, not all differences are significant. However, each soil shows significant drops of Y_w for the erosion and plowpan
275 phenoforms, again particularly beyond 2040, when comparing values with Y_w undisturbed. Soils P4 and P5 show rather
276 identical behavior but soil P6 has significantly higher values for Y_w for the erosion and plowpan phenoforms beyond 2040.
277 An increase of % OM has minimal effect as explained by Bonfante et al. (2020) when considering hydraulic conductivity and
278 moisture retention data.

279 **3.2 Soil health values for different climate periods**

280 The SH index applies to soil health parametera measurements for a given soil at a given time, defining actual conditions with
281 reference to the particular production potential of the soil type that is present as expressed by Y_w calculated with optimal soil
282 parameters as discussed above. Y_w -phenoform conveys conditions, expressed by the three soil parameters observed at the site.
283 When Y_w -phenoform is equal to Y_w , the soil health value will be 100, but this is highly improbable. Lower values indicate
284 room for improvement but offer no information as to factors that lead to these low values (see next section). Calculated SH



285 indexes for three Italian soil series in four climate periods are reported in Table 2. In this study, four soil conditions were
286 simulated that are common in the field, considering four climate periods: a non-degraded soil characterized by optimal soil
287 parameters (producing Yw-ref), and two Yw-phenoform values: erosion of topsoil, formation of a plowpan, and an increase
288 to 4% OM. As actual conditions are discussed here, the current climate of 2010-2040 should be considered. Erosion reduces
289 SH to appr. 88, while the plowpan has much stronger effect with significantly different values of 55 (soil P4), 66 (P5), and 75
290 (P6). Increasing % OM does not deviate from the value of 100, which corresponds with data reported in Figures 1, 2, and 3.
291 To determine the health index at a given time and place in a given soil, the three soil parameters discussed above are measured
292 and the model is used to calculate a (Yw-phenoform) value that is next compared with the Yw-ref value calculated with optimal
293 soil parameter values for that particular soil. Management practices should be documented that have resulted in the Yw-
294 phenoform being considered.

295 **3.3 Soil quality (SQp) in terms of characteristic ranges of soil health values**

296 The SH index, mentioned in the previous section, characterizes soil health at a given time and location, as measured in a
297 particular soil type. A gap may become obvious between Yw-phenoform and Yw-ref but it is not clear what can be done to
298 close the gap. Soil health values for a given soil series can also be obtained at different locations in the same climate zone
299 where different forms of management have resulted in different phenoforms representing a characteristic range of values that
300 can be seen as a measure for inherent soil quality (SQp). Figure 2 shows a range of values obtained for a given soil type
301 assuming, in this case, the occurrence of only three phenoforms. This only illustrates a principle and many observations in the
302 field can and should extend the number of points for Yw-phenoform. This range offers a point of reference for each
303 observation, as discussed in the previous section, and allows conclusions as to advisable management procedures associated
304 with the different phenoforms that, together, determine the observed ranges in Figure 2.

305 Figure 2 shows a decreasing sensitivity for soil degradation moving from soil P4 to soil P6. Soil health ratios change from 56
306 (P4), 66 (P5) to 78 (P6). The effects of climate change on the index are, again, strongest for soil P4. Figure 2 shows that not
307 only the ranges of the health index are significantly different for the three soils but also their resilience to climate change. A
308 particular soil health measurement in a given soil, as described in the previous section, can now be placed into the bar shown
309 in Figure 2 indicating possible room for improvement. As every measurement is combined with an assessment of soil use and
310 management that has resulted in the particular phenoform being observed, the system allows the generation of useful
311 management information for the land user.

312 **3.4 Comparing different soils in a given region (SQr).**

313 So far, particular soil types have been considered. The analysis can be extended to all soils in a given region and climate zone
314 and this comparison of different soils can be valuable for regional land use planning. This requires the definition of Yp for the
315 area that is used for the simulations. For the Italian soils being considered Yp=18 tons ha⁻¹ and this value is maintained for all
316 climate scenarios considered, implicitly assuming that other factors affecting biomass production will not change. Table 3



317 shows significant differences among the soils providing a valuable quantitative assessment. Differences are maintained when
318 different climate periods are considered. Soil P4 scores again the lowest values, with soil P5 intermediate and soil P6 with the
319 highest values but even this soil has a low score of 50 for the last climate period when a plowpan is present.

320 **3.4 How to assess soil quality in a global context? (SQw).**

321 Questions about potential food production in future, considering the effects of climate change require a mechanism to compare
322 different soils in the world in their capacity to produce biomass. Assuming a maximum production to be achieved in the world
323 (Y_{max}) considering theoretical photosynthesis under particular climate conditions, values of Y_p and Y_w can be expressed as
324 a function of Y_{max} . Use of Y_w will produce the most realistic values in view of the limited water availability in many areas
325 of the world. Areas with relatively high values have a higher potential than areas with low values and this analysis can be
326 helpful input from soil science contributing to global food production scenarios. Based on current evaluations, a Y_{max} of 20
327 tons ha^{-1} is used here as a reference and this results in SQw values that can also be expressed for various phenoforms, showing
328 effects of different forms of degradation Table 4. As in Table 3, differences between the three soils are significant. How these
329 values are to be judged will depend on comparable values to be assembled for other areas of the world.

330 **4 Discussion**

331 The Soil Health concept, as defined in the literature and as modified in this study, is inadequate to allow a comparison of the
332 capacity of different soils to function. Two soils may be healthy in their own way, but a healthy clay soil has a significantly
333 different “capacity to function” as compared with a healthy sandy soil. Still, the soil health concept is suitable to express the
334 actual condition of a given soil by comparing Y_w -phenoform with Y_w -ref as discussed in this paper, producing a soil health
335 index SH. The advantage of this procedure is its basis in a quantitative and reproducible scientific analysis of the plant
336 production process as a function of soil moisture regimes, made possible by applying soil-water-atmosphere-plant simulation
337 models. Y_w -ref and Y_w -phenoform reflect the impact of soil conditions on Y_a , the measured yield, as water and nutrients are
338 assumed to be optimal and pests and diseases do not occur. Observing the difference between Y_a on the one hand and Y_w -
339 phenoform and Y_w -ref on the other can result in fruitful interaction between soil scientists and agronomists applying a common
340 language as an effective means of communication.

341 When applied to three Italian soils, defined by soil classification in terms of three genoforms, a range of values is obtained not
342 only for an undisturbed soil but also for soils affected by poor forms of soil management resulting in erosion and compaction
343 (two “phenoforms”), and a third phenoform following “good” management increasing % OM. All of these phenoforms still
344 maintain their genoform classification (Bouma, 1989; Rossiter and Bouma, 2018). In this study effects of only three
345 hypothetical phenoforms were explored. In future, field work is required to distinguish a number of characteristic phenoforms
346 for every genoform, as a function of current and past soil management. Existing soil maps can be used to identify sampling
347 spots (e.g., Pulleman et al., 2000; Sonneveld et al., 2002).



348 Again, the different soils show significantly different behavior and the ranges for each soil series, reflecting the effects of
349 management, are different. This range represents an inherent property of the soil series being considered and it is a de facto
350 measure for soil quality (SQp) as expressed in Figure 2 and adds an important element to soil survey interpretations that are
351 now empirical and qualitative in terms of “general suitabilities or limitations for various forms of land use”. This requires that
352 properties of phenoforms are explained in terms of management practices. In this context, Pullemnan et al. (2000) and
353 Sonneveld et al. (2002) successfully correlated present and past management with % organic matter in topsoil.

354 When considering the use of soils in a given region, the SQr, as defined above, is helpful to compare the production potential
355 of different soils in that particular region

356 Finally, analyses on the world level can be made by considering the SQw index, expressing local Yw-ref values (if so desired
357 subdivided in terms of relevant phenoform values) versus a global upper limit. This could be a valuable absolute procedure to
358 compare soils on world level which may be relevant when considering future world food supply scenarios, allowing a focus
359 on potentially favorable locations. This provides an added value to the “yield-gap” program that focuses on reducing the gap
360 (van Ittersum et al., 2013).

361 However, as stated in the introduction, soil health and soil quality are no objectives in themselves. Achieving the UN
362 Sustainable Development Goals and the goals of the EU Green Deal require that soils provide effective contributions to various
363 ecosystem services that, in turn, contribute to SDGs and the Green Deal. Soils function in an interdisciplinary context and the
364 implicit hypothesis of soil health assumes that healthy soils will make better contributions to ecosystem services than unhealthy
365 ones and soils with low quality in a regional and world context. But a healthy soil can still make a poor contribution to
366 ecosystem services when poorly managed, illustrating the overriding importance of the management factor.

367 Application of soil-water-atmosphere-plant models is focused on the ecosystem service: “biomass or primary production”.
368 However, at the same time, other services have to be provided as well as discussed earlier: water quality protection, reduction
369 of greenhouse gas emissions, carbon capture and biodiversity preservation. Here, applying appropriate management is crucial
370 and, in contrast to the calculations of biomass production, there is no underlying basic theory to identify options. That is why
371 defining a characteristic range of soil health values for any given soil types a measure for inherent soil quality (SQp) is
372 important to link the land user with experiences obtained elsewhere on similar soils in the same climate zone.

373 **5 Conclusions**

- 374 1. Focusing on actual conditions when defining soil health and on inherent conditions when defining soil quality allows a
375 meaningful distinction between the two concepts that are both needed.
- 376 2. Introduction of the terminology of the agronomic “yield gap” program, allows quantitative and reproducible expressions
377 for the soil health and soil quality concepts. The distinction of Yw-ref and Yw-phenoform allows independent estimates
378 of soil contributions to Ya, the actual yield (=ecosystem service: biomass production) that is determined by many other
379 factors disciplines than soil. Applying the “yield-gap” terminology will facilitate interaction with agronomists.



- 380 3. The soil health and soil quality concepts have societal relevance as they contribute to defining ecosystem services that,
381 in turn, contribute to the UN-SDGs and the EU Green Deal.
- 382 4. Soil types were effective “carriers of information” (class-pedotransfer functions) showing distinctly different values for
383 the soils being considered.
- 384 5. Effects of climate change on Yw were significant for the Italian soils being considered with projected reductions in
385 productivity, also for non-degraded soils including soils with higher organic matter contents that may not allow
386 economically viable forms of agriculture by the end of the 21th century if irrigation is not feasible.
- 387 6. Even healthy soils can fail in making significant contributions to ecosystem services when poor management is applied.
388 Soil use and management play a key role when interpreting soil health and soil quality indexes by providing advice as
389 to how to increase indexes. The effects of soil use and management on a given type of soil (genoform) can be expressed
390 by defining phenoforms of particular genoforms. This will require new fieldwork guided by existing soil maps.
- 391 7. Effects of climate change on Yw were significant for the Italian soils being considered with projected reductions in
392 productivity, also for non-degraded soils including soils with higher organic matter contents, that may not allow
393 economically viable forms of agriculture by the end of the 21th century if irrigation is not feasible.

394 **6 Acknowledgements.**

395 Acknowledge Mrs. N. Orefice and Dr. R. De Mascellis for soil hydraulic property measurements and Dr. Eugenia Monaco for
396 the support in the analysis of climate scenarios. Climate data from the “Regional Models and Geo-Hydrogeological Impacts
397 Division” of the Centro Euro-Mediterraneo sui Cambiamenti Climatici (CMCC), Capua (CE) – Italy, were applied in this
398 study, with support by Dr. Paola Mercogliano and dr. Edoardo Bucchignani. Finally, a special thanks to Dr. Guido Rianna for
399 climate data analysis support.

400 LANDSUPPORT

401 **Funding:** This research was funded by EC H2020 LANDSUPPORT project, grant number 774234

402 **References**

- 403 Arya, L. M.: Wind and hot-air methods, in *Physical Methods*, pp. 916–926, Soil Science Society of America, Inc., 2002.
- 404 Basile, A., Coppola, A., De Mascellis, R. and Randazzo, L.: Scaling approach to deduce field unsaturated hydraulic properties
405 and behavior from laboratory measurements on small cores, *Vadose Zo. J.*, 5(3), 1005–1016, doi:10.2136/vzj2005.0128, 2006.
- 406 Basile A., Buttafuoco G., Mele G. and Tedeschi A.: Complementary techniques to assess physical properties of a fine soil
407 irrigated with saline water, *Environ. earth Sci.*, 66, 1797–1807, doi:10.1007/s12665-011-1404-2, 2012.
- 408 Baveye, P. C.: Bypass and hyperbole in soil research: Worrying practices critically reviewed through examples, *Eur. J. Soil*
409 *Sci.*, doi:10.1111/ejss.12941, 2020.



- 410 Bonfante, A. and Bouma, J.: The role of soil series in quantitative land evaluation when expressing effects of climate change
411 and crop breeding on future land use, *Geoderma*, 259–260, 187–195, 2015.
- 412 Bonfante, A., Terribile, F. and Bouma, J.: Refining physical aspects of soil quality and soil health when exploring the effects
413 of soil degradation and climate change on biomass production: An Italian case study, *SOIL*, 5(1), 1–14, doi:10.5194/soil-5-1-
414 2019, 2019.
- 415 Bonfante, A., Basile, A. and Bouma, J.: Exploring the effect of varying soil organic matter contents on current and future
416 moisture supply capacities of six Italian soils, *Geoderma*, 361, 114079, doi:10.1016/j.geoderma.2019.114079, 2020.
- 417 Bouma, J.: *Using Soil Survey Data for Quantitative Land Evaluation*, pp. 177–213, Springer, New York, NY., 1989.
- 418 Bouma, J.: Soil science contributions towards sustainable development goals and their implementation: linking soil functions
419 with ecosystem services, *J. plant Nutr. soil Sci.*, 177(2), 111–120, 2014.
- 420 Bouma, J.: Comment on: B. Minasny & A.B. Mc Bratney. 2018. Limited effect of organic matter on soil available water
421 capacity, *Eur. J. Soil Sci.*, 69(1), 154–154, doi:10.1111/ejss.12509, 2018.
- 422 Bucchignani, E., Montesarchio, M., Zollo, A. L. and Mercogliano, P.: High-resolution climate simulations with COSMO-
423 CLM over Italy: performance evaluation and climate projections for the 21st century, *Int. J. Climatol.*, 36(2), 735–756, 2015.
- 424 Bünemann, E. K., Bongiorno, G., Bai, Z., Creamer, R. E., De Deyn, G., de Goede, R., Fleskens, L., Geissen, V., Kuyper, T.
425 W., Mäder, P., Pulleman, M., Sukkel, W., van Groenigen, J. W. and Brussaard, L.: Soil quality – A critical review, *Soil Biol.*
426 *Biochem.*, 120, 105–125, doi:10.1016/j.soilbio.2018.01.030, 2018.
- 427 Droogers, P. and Bouma, J.: Soil survey input in exploratory modeling of sustainable soil management practices, *Soil Sci. Soc.*
428 *Am. J.*, 61(6), 1704–1710, 1997.
- 429 European Commission (EC). Communication from the Commission to the Council, the European Parliament, the European
430 Economic and Social Committee and the Committee of the Regions, Thematic Strategy for Soil Protection, COM 231 Final,
431 Brussels, Belgium, 2006.
- 432 Feddes, R. A., Kowalik, P. J., Zaradny, H. and others: *Simulation of field water use and crop yield.*, Centre for Agricultural
433 Publishing and Documentation., 1978.
- 434 Van Genuchten, M. T.: A closed-form equation for predicting the hydraulic conductivity of unsaturated soils, *Soil Sci. Soc.*
435 *Am. J.*, 44(5), 892–898, 1980.
- 436 Hargreaves, G. H. and Samani, Z. A.: Reference crop evapotranspiration from temperature, *Appl. Eng. Agric.*, 1(2), 96–99,
437 1985.
- 438 Holzworth, D., Huth, N. I., Fainges, J., Brown, H., Zurcher, E., Cichota, R., Verrall, S., Herrmann, N. I., Zheng, B. and Snow,
439 V.: APSIM Next Generation: Overcoming challenges in modernising a farming systems model, *Environ. Model. Softw.*, 103,
440 43–51, doi:10.1016/j.envsoft.2018.02.002, 2018.
- 441 IPCC: *Climate Change 2014--Impacts, Adaptation and Vulnerability: Regional Aspects*, edited by T. E. B. [Field, C.B., V.R.
442 Barros, D.J. Dokken, K.J. Mach, M.D. Mastrandrea, A. N. L. M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma,
443 E.S. Kissel, and L. L. W. (eds.)]. S. MacCracken, P.R. Mastrandrea, Cambridge University Press., 2014.



- 444 van Ittersum, M. K., Cassman, K. G., Grassini, P., Wolf, J., Titttonell, P. and Hochman, Z.: Yield gap analysis with local to
445 global relevance a review, *F. Crop. Res.*, 143, 4–17, 2013.
- 446 Keesstra, S. D.: The significance of soils and soil science towards realization of the UN sustainable development goals, *Soil*,
447 (2), 111–128, doi:10.5194/soil-2-111-2016-supplement, 2016.
- 448 Kroes, J. G., Van Dam, J. C., Bartholomeus, R. P., Groenendijk, P., Heinen, M., Hendriks, R. F. A., Mulder, H. M., Supit, I.
449 and Van Walsum, P. E. V: Theory description and user manual SWAP version 4, <http://www.swap.alterra.nl>, Wageningen
450 [online] Available from: www.wur.eu/environmental-research (Accessed 24 July 2019), 2017.
- 451 Van Looy, K., Bouma, J., Herbst, M., Koestel, J., Minasny, B., Mishra, U., Montzka, C., Nemes, A., Pachepsky, Y. A.,
452 Padarian, J., Schaap, M. G., Tóth, B., Verhoef, A., Vanderborght, J., van der Ploeg, M. J., Weihermüller, L., Zacharias, S.,
453 Zhang, Y. and Vereecken, H.: Pedotransfer Functions in Earth System Science: Challenges and Perspectives, *Rev. Geophys.*,
454 55(4), 1199–1256, doi:10.1002/2017RG000581, 2017.
- 455 Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L. T., Lamarque, J. F., Matsumoto, K., Montzka, S. A.,
456 Raper, S. C. B., Riahi, K. and others: The RCP greenhouse gas concentrations and their extensions from 1765 to 2300, *Clim.*
457 *Change*, 109(1–2), 213, 2011.
- 458 Moebius-Clune, B. N., Moebius-Clune, D. J., Gugino, B. K., Idowu, O. J., Schindelbeck, R. R., Ristow, A. J. and others:
459 Comprehensive assessment of soil health: The Cornell Framework Manual, Edition 3.1, Cornell Univ., Ithaca, NY, 2016.
- 460 Orgiazzi, A., Ballabio, C., Panagos, P., Jones, A. and Fernández-Ugalde, O.: LUCAS Soil, the largest expandable soil dataset
461 for Europe: a review, *Eur. J. Soil Sci.*, 69(1), 140–153, doi:10.1111/EJSS.12499@10.1111/(ISSN)13652389.BSS-
462 ANNIVERSARY-COLLECTION, 2018.
- 463 Pulleman, M. M., Bouma, J., Van Essen, E. A. and Meijles, E. W.: Soil organic matter content as a function of different land
464 use history, *Soil Sci. Soc. Am. J.*, 64(2), 689–693, 2000.
- 465 Reynolds, M., Kropff, M., Crossa, J., Koo, J., Kruseman, G., Molero Milan, A., Rutkoski, J., Schulthess, U., Balwinder-Singh,
466 Sonder, K., Tonnang, H., Vadez, V., Reynolds, M., Kropff, M., Crossa, J., Koo, J., Kruseman, G., Molero Milan, A., Rutkoski,
467 J., Schulthess, U., Balwinder-Singh, Sonder, K., Tonnang, H. and Vadez, V.: Role of Modelling in International Crop
468 Research: Overview and Some Case Studies, *Agronomy*, 8(12), 291, doi:10.3390/agronomy8120291, 2018.
- 469 Reynolds, W. D., Elrick, D. E., Youngs, E. G., Booltink, H. W. G., Bouma, J. and Dane, J. H.: Saturated and field-saturated
470 water flow parameters. 2. Laboratory methods, [online] Available from: [http://agris.fao.org/agris-
471 search/search.do?recordID=NL2003682903](http://agris.fao.org/agris-
471 search/search.do?recordID=NL2003682903) (Accessed 28 January 2019), 2002.
- 472 Ritchie, J. T.: Model for predicting evaporation from a row crop with incomplete cover, *Water Resour. Res.*, 8(5), 1204–1213,
473 1972.
- 474 Rockel, B., Will, A. and Hense, A.: The regional climate model COSMO-CLM (CCLM), *Meteorol. Zeitschrift*, 17(4), 347–
475 348, 2008.
- 476 Rossiter, D. G. and Bouma, J.: A new look at soil phenoforms--Definition, identification, mapping, *Geoderma*, 314, 113–121,
477 2018.



478 Sonneveld, M. P. W., Bouma, J. and Veldkamp, A.: Refining soil survey information for a Dutch soil series using land use
479 history, *Soil Use Manag.*, 18(3), 157–163, doi:10.1111/j.1475-2743.2002.tb00235.x, 2002.

480 Steduto, P., Hsiao, T. C., Fereres, E. and Raes, D.: Crop yield response to water, FAO Roma., 2012.


481 White, J. W., Hunt, L. A., Boote, K. J., Jones, J. W., Koo, J., Kim, S., Porter, C. H., Wilkens, P. W. and Hoogenboom, G.:
482 Integrated description of agricultural field experiments and production: The ICASA Version 2.0 data standards, *Comput.*
483 *Electron. Agric.*, 96, 1–12, doi:10.1016/j.compag.2013.04.003, 2013.

484 Zollo, A. L., Turco, M. and Mercogliano, P.: Assessment of hybrid downscaling techniques for precipitation over the Po river
485 basin, in *Engineering Geology for Society and Territory-Volume 1*, pp. 193–197, Springer., 2015.

486


487

488



489

490 **Figure 1: The average Y_w of four soil phenoforms of three soils (a) P4, (b) P5, and (c) P6 under reference (RC) and future climate**
 491 **scenario (RCP 8.5). Y_p is the local current potential production and Y_{max} is the maximum potential production under no stressed**
 492 **field conditions (water, nutrient and pests disease).**



493

494 **Figure 2. Range of soil health indexes – $SH=(Yw\text{-phenofom}/Yw\text{-ref}) \times 100$ - for the three soils demonstrating differences among**
495 **soils and projected effects of climate change. This range characterizes the inherent soil quality SQp for these particular soil types.**

496



Tab. 1 Physical characteristics and classifications of the three Italian soils being studied (from Bonfante et al., 2020).

ID	Soil		Hor.	Thick. (cm)	Clay	Silt	Sand	S.O.M.
	Series	Classification						
P4	Sordio ⁺	Ultic Haplustalf, coarse loamy, mixed, mesic (Sandy Loam)	Ap1	0-18	17.9	32.6	49.5	1.4
			Ap2	18-30	17.7	33.2	49.1	1.4
			Bt1	30-56	21.8	31.4	46.8	0.4
			Bt2	56-83	13.4	12.1	74.5	0.2
			BC	83+	10.0	6.3	83.7	0.1
P5	Masseria Manfredi ⁺⁺	Typic Ustivitrands, sandy, mixed, thermic (Sandy Loam)	Ap1	0-10	10.5	38.5	51.0	2.6
			Ap2	10-40	5.9	43.6	50.5	2.6
			Bw	40-80	3.9	31.1	65.0	-
			BC	80-110	11.6	15.4	73.0	-
			C	110+	4.6	9.4	86.0	-
P6	Masseria Battaglia ⁺⁺	Vitrandic Haplustept, sandy, mixed (Loamy Sand)	Ap1	0-20	4.1	18.6	77.3	1.7
			Ap2	20-53	6.1	18.4	75.5	1.6
			Bw1	53-61	1.4	12.4	86.2	0.9
			Bw2	61-106	2.2	8.7	89.1	0.9
			C	106+	1.0	24.6	74.4	0.2

⁺ Soil series *The soil map of Lodi plain (1:37.500) (ERSAL, 2000)*

⁺⁺ Closed to soil series of *"The soil map of province of Naples" (1:75.000) (Di Gennaro and Terribile, 1999)*

497

498



Tab.2. Table 2. Soil health indexes - SH ((Yw-phenoform/Yw-ref) x 100), defining actual conditions, for three selected soils being studied for four climate periods as indicated. Values are reported for the non-degraded soil and for hypothetical phenoforms representing , erosion of 20 cm of topsoil without a change of soil classification (Yw-erosion) and occurrence of a plowpan at 30 cm depth (Yw-plowpan) Indexes are also included for hypothetically increased % organic matter to levels of 4% ,(Yw-4% O.M.).

Soil	Climate scenario	Yw-erosion	Yw-Plowpan	Yw- 4% O.M.
P4	RC (1971-2005)	88.4 (± 2.0)	55.4 (± 1.9)	101.1 (± 1.8)
	RCP 8.5 (2010-2040)	88.0 (± 1.9)	55.4 (± 1.9)	101.0 (± 1.7)
	RCP 8.5 (2040-2070)	85.1 (± 2.0)	51.0 (± 1.8)	101.1 (± 1.9)
	RCP 8.5 (2070-2100)	83.7 (± 2.3)	49.2 (± 2.0)	101.2 (± 2.2)
P5	RC (1971-2005)	88.9 (± 1.7)	66.1 (± 1.8)	100.7 (± 1.6)
	RCP 8.5 (2010-2040)	88.9 (± 1.6)	66.3 (± 1.7)	100.7 (± 1.5)
	RCP 8.5 (2040-2070)	87.0 (± 1.7)	62.3 (± 1.7)	100.8 (± 1.7)
	RCP 8.5 (2070-2100)	86.7 (± 2.0)	61.3 (± 2.0)	100.8 (± 1.9)
P6	RC (1971-2005)	85.5 (± 1.4)	75.4 (± 1.4)	102.4 (± 1.0)
	RCP 8.5 (2010-2040)	84.9 (± 1.4)	75.0 (± 1.4)	102.7 (± 1.0)
	RCP 8.5 (2040-2070)	82.5 (± 1.5)	72.2 (± 1.5)	103.7 (± 1.3)
	RCP 8.5 (2070-2100)	82.1 (± 1.8)	71.8 (± 1.8)	104.2 (± 1.5)

499
 500



Table 3. SQR index ((Yw/Yp) x100) for the three selected soils and the four climate periods.
 Yp is assumed to be 18 tons ha⁻¹.

Soil	Climate scenario	Soil phenoform				
		Undisturbed	4% O.M.	Erosion	Plowpan	
		(Yw/Yp) x 100				
P4	RC	(1971-2005)	76.3 (± 1.8)	77.2 (± 1.8)	67.4 (± 2.1)	42.3 (± 2.0)
	RCP 8.5	(2010-2040)	74.9 (± 1.7)	75.7 (± 1.7)	66.0 (± 2.1)	41.5 (± 2.0)
		(2040-2070)	65.2 (± 1.8)	65.9 (± 1.8)	55.4 (± 2.0)	33.2 (± 1.7)
	RCP	(2070-2100)	59.7 (± 2.1)	60.4 (± 2.1)	50.0 (± 2.3)	29.4 (± 1.9)
P5	RC	(1971-2005)	83.1 (± 1.6)	83.6 (± 1.5)	73.8 (± 1.8)	54.9 (± 1.9)
	RCP 8.5	(2010-2040)	81.4 (± 1.4)	82.0 (± 1.4)	72.4 (± 1.8)	53.9 (± 1.9)
		(2040-2070)	72.9 (± 1.6)	73.5 (± 1.6)	63.5 (± 1.7)	45.4 (± 1.7)
	RCP	(2070-2100)	67.8 (± 1.9)	68.4 (± 1.9)	58.8 (± 2.1)	41.5 (± 2.0)
P6	RC	(1971-2005)	92.0 (± 1.1)	94.2 (± 0.9)	78.7 (± 1.7)	69.4 (± 1.7)
	RCP 8.5	(2010-2040)	90.6 (± 1.0)	93.0 (± 0.8)	76.9 (± 1.6)	67.9 (± 1.6)
		(2040-2070)	83.4 (± 1.3)	86.5 (± 1.2)	68.8 (± 1.6)	60.2 (± 1.5)
	RCP	(2070-2100)	78.2 (± 1.5)	81.5 (± 1.4)	64.2 (± 2.0)	56.1 (± 1.9)

501
 502



Table 4. SQw index ((Yw/Ymax)x100) for the three selected soils and the four climate periods. Ymax is assumed to be 20 tons ha⁻¹.

Soil	Climate scenario	Soil phenoform				
		Undisturbed	4% O.M.	Erosion	Plowpan	
(Yw/Ymax) x 100						
P4	RC	(1971-2005)	68.7 (± 1.8)	69.4 (± 1.8)	60.7 (± 2.1)	38.0 (± 2.0)
	RCP 8.5	(2010-2040)	67.4 (± 1.7)	68.1 (± 1.7)	59.4 (± 2.1)	37.4 (± 2.0)
		(2040-2070)	58.6 (± 1.8)	59.3 (± 1.8)	49.9 (± 2.0)	29.9 (± 1.7)
		(2070-2100)	53.7 (± 2.1)	54.4 (± 2.1)	45.0 (± 2.3)	26.4 (± 1.9)
P5	RC	(1971-2005)	74.8 (± 1.6)	75.3 (± 1.5)	66.4 (± 1.8)	49.4 (± 1.9)
	RCP 8.5	(2010-2040)	73.3 (± 1.4)	73.8 (± 1.4)	65.1 (± 1.8)	48.5 (± 1.9)
		(2040-2070)	65.6 (± 1.6)	66.2 (± 1.6)	57.1 (± 1.7)	40.9 (± 1.7)
		(2070-2100)	61.0 (± 1.9)	61.5 (± 1.9)	52.9 (± 2.1)	37.4 (± 2.0)
P6	RC	(1971-2005)	82.8 (± 1.1)	84.8 (± 0.9)	70.9 (± 1.7)	62.5 (± 1.7)
	RCP 8.5	(2010-2040)	81.5 (± 1.0)	83.7 (± 0.8)	69.2 (± 1.6)	61.1 (± 1.6)
		(2040-2070)	75.0 (± 1.3)	77.8 (± 1.2)	61.9 (± 1.6)	54.2 (± 1.5)
		(2070-2100)	70.4 (± 1.5)	73.3 (± 1.4)	57.8 (± 2.0)	50.5 (± 1.9)