

Microbial communities and their predictive functional profiles in the arid soil of Saudi Arabia

Munawwar A. Khan¹, Shams T. Khan²

¹Department of Life and Environmental Sciences, College of Natural and Health Sciences, Zayed University, Dubai, 19282, UAE

²Department of Agricultural Microbiology, Faculty of Agricultural Science, Aligarh Muslim University, Aligarh, U.P, India

Correspondence to: Shams T. Khan (shamsalig75@gmail.com)

Abstract. Saudi Arabia has the world's fifth-largest desert and is the biggest importer of food and agricultural products. Understanding soil microbial communities is key to improving the agricultural potential of the region. Therefore, soil microbial communities of the semi-arid region of Abha known for agriculture and arid regions of Hafr Al-Batin and Muzahmiyah were studied using Illumina sequencing. Microbial community composition varied remarkably from other deserts and from one place to another. The highest diversity was found in the rhizospheric soil of Muzahmiyah followed by Abha. Firmicutes, Proteobacteria, and Actinobacteria were three main phyla detected in all the samples. Unlike other deserts, Bacteroidetes was not a major constituent and the population of Firmicutes was quite high. Soils from the agricultural region of Abha were significantly different from other samples in containing only 1% Firmicutes and three to six times higher population of Actinobacteria and Bacteroidetes, respectively. The presence of photosynthetic bacteria, ammonia oxidizers, and nitrogen fixers along with bacteria capable of surviving on simple and unlikely carbon sources like dimethylformamide was indicative of their survival strategies under harsh environmental conditions in the arid soil. Functional inference using PICRUSt analysis shows an abundance of genes involved in photosynthesis and nitrogen fixation. Microbial communities show greater similarity with hot Namib desert than with the cold Antarctic desert.

Keywords: Saudi Arabian Desert; soil microbiome; arid soil; Actinobacteria; PICRUSt

1 Introduction

Saudi Arabian desert also referred to as the Sahara Arabian desert is the fifth-largest desert of the world bordering with Yemen, the Persian Gulf, and Iraq (Holm, 1960). The desert is characterized by the presence of vast barren areas of sand referred to as empty quarters or Rub-al Khali and Wadi al-Batin. Most of the area is barren with almost no vegetation and the growing population is dependent on imported agricultural products for food (Fiaz et al., 2016). According to the government of the USA, Saudi Arabia imports 14.8 billion US dollar worth of agricultural products every year (<https://www.export.gov/apex/article2?id=Saudi-Arabia-Market-Overview>). Although the climatic conditions are not favorable, the Saudi government has launched various programs to promote agriculture. In fact, 52.7 million ha area, which is

25% of the total country's area is currently cultivable (Fiaz et al., 2016). Especially the Asir region, with Abha as its capital is well known for agriculture and receives more rainfall than the rest of the country.

The microbial communities of Arid regions are largely uncharacterized, to the best of our knowledge no report from Saudi-Arabia is available (Makhalanyane et al., 2015; Schulze-Makuch et al., 2018). Although the vast desert lacks vegetation and therefore are expected to be devoid of macromolecules and the microbial communities involved in the recycling of the nutrients. However, active microbial communities have been detected even in hyper-arid deserts of the Atacama where the rain is received only once in a decade (Schulze-Makuch et al., 2018). Such studies are crucial in improving the agricultural potential in these extreme habitats. And to design strategies for the modification of soil with microbial consortia for improving the agricultural potential of the arid soil (Fierer et al., 2012; Fierer, 2017). These microorganisms may alter soil fertility through sustaining the soil nutrient cycling, carbon sequestration, and by influencing other geochemical processes. ~~Consequently, improving soil fertility and the agricultural potential of the soil.~~ This study, therefore, was aimed at comparing the soil microbial communities of Abha from the Asir region and arid regions of Mazamiyah and Hafr Al-Batin. The knowledge of the microbial communities present in these regions may provide some insights into the role of microorganisms in various geochemical processes.

45

2 Materials and Methods

2.1 Soil Sample Collection

The desert soil samples were collected from three regions namely Abha of Asir region (semi-arid), and the arid regions of Muzahmiyah (near Riyadh) and Hafr Al-Batin. Samples were collected between 26 Jan and 18 Feb 2019. Figure 1 and table 1 shows the locations of the sampling sites and the climatic conditions at these sites. ~~Each sample was a mixture of three subsamples each collected from a distance of 1m from each other. Debris (2 cm) from the surface was removed at the time of sampling and a soil core from a depth of 5 cm was then obtained.~~ From Muzahmiyah two samples namely a rhizospheric soil and a non-rhizospheric soil sample was collected. Following collection, samples were transported to the lab at room temperature and were processed immediately after arriving at the lab. Portions of soils were stored at -20°C for DNA extractions. Biogeochemical properties and other details of sampling sites are given in table 1. The pH, of the soil, was determined by using a pH meter. While total aerobic counts in the soil were determined on 1/5th diluted nutrient agar plates using the dilution plating method.

55

2.2 DNA extraction and HiSeq analysis

Genomic DNA from the soil was prepared using the direct lysis method of Robe et al (Robe et al., 2003). For obtaining enough DNA, extraction was carried out in replicates and the DNA was pooled and concentrated. The composition and diversity of bacterial communities in soil were determined by amplifying the V3-V4 regions of bacterial 16S ribosomal RNA (rRNA) genes. A set of 341F and 806R primers and the DNA extracted from the samples as a template were used for amplification.

60

65 PCR reactions were carried out at 50 µl scale containing ~40 ng of DNA template, 25 µl Dream Taq Green PCR Master Mix (2×), 20.5 µl H₂O, 0.5 µl of 1% bovine serum albumin, and 0.2 µM of each primer. The PCR was carried out by programming the thermal regime as initial denaturation at 95 °C for 5 min; followed by 35 cycles of denaturation at 95 °C for 30 s, annealing at 56 °C for 30 s, and extension at 72 °C for 30 s; and a final extension step at 72 °C for 7 min. Amplicon sequencing was conducted on an Illumina HiSeq 2500 platform. Further processing of the reads and quality filtering was conducted as described earlier (Yuan et al., 2018).

70 **2.3 Data analysis**

Raw data obtained from the sequencing was processed using QIIME (**Quantitative Insights Into Microbial Ecology**) (Caporaso et al., 2010). Sequences were clustered into operational taxonomic units using UCLUST and an identity threshold of 97%. Sequences were assigned to their phylogenetic groups using the QIIME pipeline and the green genes database ver. 13.5 (Santamaria et al., 2012). Further processing of the sequences was carried out using Calypso (Zakrzewski et al., 2017).
75 Rarefaction curves were calculated for the number of species present in each sample. Alpha diversity was determined using both taxonomic metrics (numbers of phylotypes). ~~Principal coordinates analyses were conducted as input the pairwise distances between bacterial communities (Bray-Curtis and UniFrac distances calculated from the 16S rRNA gene amplicon data).~~ To test whether sample categories harbored significantly different microbial communities, we used an analysis of similarities (ANOSIM). To determine whether the relative abundances of individual taxa were significantly different between sample
80 categories pairwise *t*-tests with P values were calculated. The abundance of various taxa in the samples as correlation charts were also calculated using Calypso. While abundance pie charts were calculated using Krona (Ondov et al., 2013). **Predictive functional analysis of microbial communities using 16S rRNA gene sequences was carried out using PICRUSt (Phylogenetic Investigation of Communities by Reconstruction of Unobserved States).and STAMP (statistical analysis of taxonomic and functional profiles) for functional inference using PICRUSt and STAMP software v 2.1.3 was used for statistical analyses and to detect differentially abundant OTUs between two sample groups (Parks et al., 2014).**

85 The sequences have been submitted to SRA with the accession numbers from SAMN12651127-12651133.

3 Results and discussions

3.1 Properties of soil samples

The details of the collected soil samples are given in table 1. The pH of all the soil samples was alkaline and ranged from 7.9
90 to 8.4. Abha located at high elevation near Yemen is a semi-arid region. **The weather of the** city is generally mild throughout the year and is especially cooler during the “low-sun” season. The annual average temperature of Abha is only 17.5 °C and seldom rises above 35 °C. The city receives an annual rainfall of about 230 mm most of which occurs between February and April. The soil type in Abha is sand and gravel and the pH of the soil sample was slightly alkaline i.e. 7.9. The total CFU counts obtained on nutrient agar medium was $3.63 \pm 1.9 \times 10^5$ CFU/g of soil. While, Muzahmiah is an arid region with an
95 average annual temperature of 25.3 °C, and in summer it may cross 45 °C. The area receives annual precipitation of only 88

mm. The soil type in Muzahmiyah is reported to be sandy loam (Siham, 2007). Two samples were collected from Muzahmiyah the pH of both samples was alkaline (8.2). The CFU counts of the rhizospheric soil of *Haloxylon persicum* was fifty times higher ($5.5 \pm 1.9 \times 10^5$ CFU/g of soil) than the non-rhizospheric ($1.1 \pm 0.9 \times 10^4$ CFU/g of soil) soil at a distance of a few centimeters. Hafr Al-Batin is also an arid region with an average annual rainfall of only 126 mm. The soil was alkaline with a pH of 8.4 and the aerobic bacterial count on nutrient agar was $8.1 \pm 2.7 \times 10^4$ CFU/g of soil.

3.2 Microbial diversity of soil samples

Reads per sample varied between 68640 to 420570. For analysis, a total of 12,790 data points were included for each sample.

The alpha diversity in the samples was calculated using species as OTUs, rarefaction curves and Shannon diversity index show the highest diversity is associated with the samples collected from the rhizospheric soil (M15) of *Haloxylon persicum* collected from Muzahmiyah (Figure 2). While the microbial population was least diverse in the sample collected from the non-rhizospheric soil from the same region (M5). It is to be noted that the diversity of rhizospheric soil from Muzahmiyah was comparable to the diversity observed in the samples collected from Abha, a region known for agriculture. Aerobic plate counts obtained from the two samples were also comparable. The species richness of Hafr Al-Batin (HB) was also comparable to the samples collected from Abha and Muzahmiyah although the aerobic plate count was as low as found in non-rhizospheric soil from Muzahmiyah (M5). R values of 0.85 obtained in ANOISM analysis show that the studied microbial communities are significantly different from each other (Figure 3 A)

Between 69-88% of the total reads were assigned to Bacteria and the majority of the reads can be assigned to phylum Actinobacteria, Firmicutes and Proteobacteria. While remaining reads were assigned to unknown groups detailed in supplementary figure 1. Populations of Firmicutes in samples collected from Muzahmiyah M15, M5, Hafr Al-batin, and Abha were 50, 0.7, 13, and 1% of the total bacteria, respectively. While the population of proteobacteria was 25, 90, 31, and 53% in soil samples from Muzahmiyah M15, M5, Hafr Al-batin, and Abha, respectively. The population of Actinobacteria was highest in soil from Abha (31% of the total bacteria). While a comparable population of Actinobacteria in soils of Hafr Al-Batin (10%) and rhizospheric soil of Muzahmiyah (9%) was observed (Figure 3 B). Earlier reports show that the major soil bacteria found in desert soil belong to Actinobacteria, Bacteroidetes, and Proteobacteria (Fierer et al., 2012; Andrew et al., 2012). Soil sample from Abha shows almost the same pattern where the population of Proteobacteria, Actinobacteria, and Bacteroidetes was 53, 31, and 6% of the total bacteria, respectively. While samples from Muzahmiyah and Hafr Al-Batin vary in not having the significant populations of Bacteroidetes and an increased population of Firmicutes (Figure 3 B). A high population of Bacteroidetes in samples other than Abha may be due to the unavailability of complex organic matter in these soils. As members of the Bacteroidetes are known to degrade various macromolecules in the soil. The high populations of Firmicutes and Actinobacteria in desert soil may be due to their ability to produce spores under high temperature and aridity. It is to be noted that temperatures do not cross 35 °C in Abha which has the lowest population of Firmicutes which may not be the case with other deserts. A high population of actinobacteria has been found in both the cold Antarctica desert as well as hot Namib

130 desert (Aislabie et al., 2006; Armstrong et al., 2016). Interestingly, the highest population of Actinobacteria (34% of the total
bacteria) was observed in Abha which is Geographically close to the Namib desert. Acidobacteria were only found in the
rhizospheric soil of Muzahmiyah (2% of the total), while planctomycetes were found in the soils of Abha and Hafr Al-Batin.
Among other minor phyla, notably present in soil samples were the members of Phylum *Gemmatimonadetes*, like Gemm 3 in
135 some cases constituting as much as 4% (Muzahmiyah rhizospheric soil) of the total reads. These bacteria have been reported
earlier also in the desert soil and recently strains from the phylum with photosynthetic capability have been cultured (Meola
et al., 2015; Zeng et al., 2014). The ammonia-oxidizing Archaea Candidatus *Nitrososphaera gargensis* was also found in most
of the samples. Reads belonging to iii115 constituted up to 3% of the total population at least in two samples (M15 and Hafr
Al-Batin), **these sequences have been reported** from the soil in earlier studies also (Marasco et al., 2018). The Antarctica desert
soil survey shows that Actinobacteria were present prominently along with *Bacillus* spp., *Flavobacterium* spp. and
140 *Acinetobacter* spp. Deinococcus–Thermus and Gemmatimonadetes clades, which have low or no representation in other
surface soils, are relatively common in Dry Valley clone libraries. Members of 14 phyla have been found in the soil of
Antarctica desert including Actinobacteria, *Gemmatimonas*, *Proteobacteria*, *Bacteroidetes*, *Deinococcus* and *Thermus*,
Planctomycetes, *Chloroflexi*, *Verrucomicrobia*, *Acidobacteria*, *Cyanobacteria*, TM7, and OP11. The most dominant were
Acidobacteria, *Actinobacteria*, and *Bacteroidetes* (Cary et al., 2010). In the case of the hot Namib desert, 19 different phyla
145 were observed as shown in Figure 3B. The most abundant phyla were *Bacteroidetes*, *Proteobacteria*, and *Actinobacteria*
(Armstrong et al., 2016).

The relative abundance of the major genus found in the soil samples is shown as a heatmap in figure 4B. A detailed microbial
community composition generated by the Calypso program is shown in Supplementary figures 1-4. The Venn diagram (Figure
4A) shows that the core genera found in all the samples were 272, while Abha and Rhizospheric soil sample from Muzahmiyah
150 (M15) shared the maximum number of genera (300). While M15 and HB also shared 295 genera and Abha and HB shared 279
genera. Some of the most abundant genera include *Pseudomonas*, *Paenibacillus*, *Bacillus*, Candidatus *Nitrososphaera*,
Devosia, *Adhaeribacter*, and others. The bacterial genera found in desert soil are expected to withstand extreme climatic
conditions and to perform some vital functions. For example, genus like *Bacillus* and *Paenibacillus* or *Actinobacteria* like
Nocardioidea, and *Streptomyces* are spore-forming bacteria and hence can survive extreme heat and arid conditions. Many of
155 these genera have been isolated from desert already. *Ramlibacter* one of the genera found in our samples forms cyst and its
genome analysis shows adaptation to **arid** conditions (De Luca et al., 2011). *Ramlibacter* was originally isolated from the
meteorite fragments buried in the sands of a desert (De Luca et al., 2011). *Modestobacter* another dominant bacteria found in
our samples was isolated from the Atacama desert of Chile, South America (Busarakam et al., 2016). The genera like
Pseudomonas and *Adhaeribacter*, may produce extracellular polysaccharide to survive under the arid condition and to form
160 strong biofilms or may also contribute to water retention in soil promoting the formation of soil crust.

It is to be noted that some bacteria may play important roles besides tolerating the extreme conditions. Candidatus
Nitrososphaera and *Planctomyces* found in all the samples in high numbers **are known to oxidize ammonia** (Stieglmeier et al.,

2014). Notably, it was observed that the desert soil catalyzes ammonia formation (Schrauzer, 1978). Nitrite oxidizing bacteria
165 *Nitrospira* were also found in all the samples but the reads were especially higher in Rhizospheric soil from Muzahimiyah. The
phototrophic bacterium *Rhodoplanes* present in all the samples may be one of the bacteria involved in carbon fixation in the
nutrient-deficient arid soil. Members of *Devosia* are known to nodulate *Neptunia* for nitrogen fixation and therefore, may serve
as a nitrogen source in otherwise nitrogen-deficient arid soils. Members of the genus *Mesorhizobium*, *Bradyrhizobium*, and
Sinorhizobium were also found in all the samples. While the members of the genus *Rhizobium* were detected only in
170 Rhizospheric soil from Muzahimiyah. Another genus found in significant numbers in all the soil samples and its population
was especially high in Abha was *Balneimonas*. Which is the member of family *Bradyrhizobiaceae* and is known to produce
extracellular material playing an important role in the formation of soil crust (Matthews et al., 2019).

Abha soil sample was also distinct in having high populations of bacteria with the ability to survive on the simple
source of nutrients and extreme conditions or the ability to perform an important geochemical or agricultural function. The
175 population of genera *Adhaeribacter*, *Modestobacter*, *Ramlibacter*, radiation-resistant *Geodermatophilus*, *Pseudonocardia*, and
Flavobacterium was high in samples from Abha. The population of *Paracoccus* and *Phenyllobacterium* capable of growing
optimally on artificial compounds like **dimethylformamide**, chloridazon, antipyrin, and pyramidon was also significantly
higher (Eberspächer and Lingens, 2006). N₂-fixing *Azospirillum*, *Agrobacterium*, and phototrophic *Rhodobacter* were also
present in high numbers in soil from Abha. While, the rhizospheric soil of Muzahimiyah also shows similar pattern containing
180 a high population of Candidatus *Nitrososphaera*, *Ramlibacter*, *Bradyrhizobium*, Phototrophic *Rhodoplanes*. But the sample
was different in having the significantly high populations of bacteria like *Paenibacillus*, *Alicyclobacillus*, and *Sporosarcina*.
Soil samples collected from Hafr Al-Batin have a completely different community with high populations of *Pseudomonas*,
Propionibacterium, *Brevundimonas*, *Staphylococcus*, and *Burkholderia*. **The microbial community in Hafr Al-Batin soil is
completely different probably due to the completely different environmental conditions in Hafr Al-Batin. This region is well
185 known for its extreme arid condition in Saudi Arabia.** The functional inference using PICRUSt analysis show similar results
(Figure 5 and 6). Some of the most abundant genes belong to transporters, peptidases, housekeeping genes, and general
function. Genes involved in prokaryotic photosynthesis, chlorophyll metabolism constitute more than 2% of the total genes
(Figures 5 and 6). Furthermore, genes involved in the metabolism of simple substrates like methane, butanoate, and benzoate
were also predicted to have a high proportion. This indicates the survival strategy of the microbial community under nutrient-
190 deficient harsh environmental conditions. **Interestingly, it was found that the proportion of genes for prokaryotic
photosynthesis was lowest (1/4th) in samples from Abha compared to other samples. Probably comparatively higher soil
fertility and semiarid nature of soil do not require a high population of photosynthetic bacteria for maintaining and providing
carbon to other soil organisms. Similarly, the proportion of genes involved in methane and nitrogen metabolism and
peptidoglycan biosynthesis were lowest (~1/3rd) in Abha samples. Indicating that the population of Gram-positive bacteria in
195 all the samples other than the sample from Abha is high. Production of methane is a characteristic of arid soils and the presence
of these genes in high proportion in all the samples other than Abha further confirms the fact observed in previous studies.**

Bacterial

4. Conclusion

200 Understanding the composition of the desert microbial community may help us in understanding the role of different
microorganisms in extreme environments. The analysis shows that the microbial communities of Saudi desert were
characterized by the presence of high numbers of Actinobacteria, Proteobacteria, and Firmicutes. These microbial communities
besides showing Saharan desert signature phyla like *Gemmatimonas* also show biogeochemically important microorganisms
205 exemplified by primary producers like *Rhodoplanes* and Cyanobacteria, nitrogen-fixing members of the genus *Rhizobium*,
Bradyrhizobium, and ammonia oxidizers *Candidatus Nitrososphaera*. Communities were also characterized by the presence
of microbes capable of growing on simple and unlikely carbon sources such as methane, butanoate, dimethylformamide
indicating the survival strategies adopted by microbial communities under nutrient-deficient condition.

Appendix A: Supplementary data: Figure 1-4 show the diversity of microbial community as Krona pie charts generated
210 using Calypso for M15, M5, Abha and Hafr Al-Batin, respectively.

Code and data availability. The nucleotide sequence data is submitted to GenBank with the accession number
SAMN12651127-12651133.

215 **Author contributions.** MAK performed data analysis and prepared the manuscript. STK collected the samples, prepared
genomic DNA and carried out sequencing.

Competing interests. The authors declare that they have no conflict of interest.

220 **Acknowledgements.** We would like to thank Zayed University Office of Research for financial support.

Financial support. This research has been supported by the Zayed University Microbial Diversity and Eco-genomics research
cluster project (Activity code number: R16092).

225 **Review statement.** This paper was edited by and reviewed by two referees.

References

1. Holm DA (1960) Desert Geomorphology in the Arabian Peninsula. Science 132 (3437):1369-1379

2. Fiaz S, Noor MA, Aldosri FO (2018) Achieving food security in the Kingdom of Saudi Arabia through innovation: Potential role of agricultural extension. *Journal of the Saudi Society of Agricultural Sciences* 17 (4):365-375. doi:<https://doi.org/10.1016/j.jssas.2016.09.001>
3. Makhalanyane TP, Valverde A, Gunnigle E, Frossard A, Ramond J-B, Cowan DA (2015) Microbial ecology of hot desert edaphic systems. *FEMS Microbiology Reviews* 39 (2):203-221. doi:10.1093/femsre/fuu011 %J *FEMS Microbiology Reviews*
4. Schulze-Makuch D, Wagner D, Kounaves SP, Mangelsdorf K, Devine KG, de Vera J-P, Schmitt-Kopplin P, Grossart H-P, Parro V, Kaupenjohann M, Galy A, Schneider B, Airo A, Frösler J, Davila AF, Arens FL, Cáceres L, Cornejo FS, Carrizo D, Dartnell L, DiRuggiero J, Flury M, Ganzert L, Gessner MO, Grathwohl P, Guan L, Heinz J, Hess M, Keppler F, Maus D, McKay CP, Meckenstock RU, Montgomery W, Oberlin EA, Probst AJ, Sáenz JS, Sattler T, Schirmack J, Sephton MA, Schloter M, Uhl J, Valenzuela B, Vestergaard G, Wörmer L, Zamorano P (2018) Transitory microbial habitat in the hyperarid Atacama Desert. 115 (11):2670-2675. doi:10.1073/pnas.1714341115 %J *Proceedings of the National Academy of Sciences*
5. Fierer N (2017) Embracing the unknown: disentangling the complexities of the soil microbiome. *Nature Reviews Microbiology* 15:579. doi:10.1038/nrmicro.2017.87
6. Fierer N, Leff JW, Adams BJ, Nielsen UN, Bates ST, Lauber CL, Owens S, Gilbert JA, Wall DH, Caporaso JG (2012) Cross-biome metagenomic analyses of soil microbial communities and their functional attributes. 109 (52):21390-21395. doi:10.1073/pnas.1215210110 %J *Proceedings of the National Academy of Sciences*
7. Robe P, Nalin R, Capellano C, Vogel TM, Simonet P (2003) Extraction of DNA from soil. *European Journal of Soil Biology* 39 (4):183-190. doi:[https://doi.org/10.1016/S1164-5563\(03\)00033-5](https://doi.org/10.1016/S1164-5563(03)00033-5)
8. Yuan H-Y, Liu P-P, Wang N, Li X-M, Zhu Y-G, Khan ST, Alkhedhairy AA, Sun G-X (2018) The influence of soil properties and geographical distance on the bacterial community compositions of paddy soils enriched on SMFC anodes. *Journal of Soils and Sediments* 18 (2):517-525. doi:10.1007/s11368-017-1769-2
9. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Pena AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE, Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Turnbaugh PJ, Walters WA, Widmann J, Yatsunenko T, Zaneveld J, Knight R (2010) QIIME allows analysis of high-throughput community sequencing data. *Nature methods* 7 (5):335-336. doi:10.1038/nmeth.f.303
10. Santamaria M, Fosso B, Consiglio A, De Caro G, Grillo G, Licciulli F, Liuni S, Marzano M, Alonso-Aleman D, Valiente G, Pesole G (2012) Reference databases for taxonomic assignment in metagenomics. *Briefings in Bioinformatics* 13 (6):682-695. doi:10.1093/bib/bbs036 %J *Briefings in Bioinformatics*
11. Zakrzewski M, Proietti C, Ellis JJ, Hasan S, Brion M-J, Berger B, Krause L (2017) Calypso: a user-friendly web-server for mining and visualizing microbiome-environment interactions. *Bioinformatics* 33 (5):782-783. doi:10.1093/bioinformatics/btw725
12. Ondov BD, Bergman NH, Phillippy AM (2011) Interactive metagenomic visualization in a Web browser. *BMC bioinformatics* 12:385. doi:10.1186/1471-2105-12-385

13. Parks DH, Tyson GW, Hugenholtz P, Beiko RG (2014) STAMP: statistical analysis of taxonomic and functional profiles. *Bioinformatics* (Oxford, England) 30 (21):3123-3124. doi:10.1093/bioinformatics/btu494
14. A. Siham A-K (2007) Soil Analysis of Contaminated Soil from Riyadh City, Saudi Arabia and Influence of Aluminium and Cobalt Ions on the Growth of Fungi Isolated, vol 7. doi:10.3923/jbs.2007.549.553
- 265 15. Andrew DR, Fitak RR, Munguia-Vega A, Racolta A, Martinson VG, Dontsova K (2012) Abiotic Factors Shape Microbial Diversity in Sonoran Desert Soils. 78 (21):7527-7537. doi:10.1128/AEM.01459-12 %J *Applied and Environmental Microbiology*
16. Aislabie JM, Chhour K-L, Saul DJ, Miyauchi S, Ayton J, Paetzold RF, Balks MR (2006) Dominant bacteria in soils of Marble Point and Wright Valley, Victoria Land, Antarctica. *Soil Biology and Biochemistry* 38 (10):3041-3056. doi:<https://doi.org/10.1016/j.soilbio.2006.02.018>
- 270 17. Armstrong A, Valverde A, Ramond J-B, Makhalanyane TP, Jansson JK, Hopkins DW, Aspray TJ, Seely M, Trindade MI, Cowan DA (2016) Temporal dynamics of hot desert microbial communities reveal structural and functional responses to water input. *Scientific Reports* 6:34434. doi:10.1038/srep34434
<https://www.nature.com/articles/srep34434#supplementary-information>
- 275 18. Meola M, Lazzaro A, Zeyer J (2015) Bacterial Composition and Survival on Sahara Dust Particles Transported to the European Alps. 6 (1454). doi:10.3389/fmicb.2015.01454
19. Zeng Y, Feng F, Medová H, Dean J, Koblížek M (2014) Functional type 2 photosynthetic reaction centers found in the rare bacterial phylum Gemmatimonadetes. 111 (21):7795-7800. doi:10.1073/pnas.1400295111 %J *Proceedings of the National Academy of Sciences*
- 280 20. Marasco R, Rolli E, Fusi M, Michoud G, Daffonchio D (2018) Grapevine rootstocks shape underground bacterial microbiome and networking but not potential functionality. *Microbiome* 6 (1):3. doi:10.1186/s40168-017-0391-2
21. Cary SC, McDonald IR, Barrett JE, Cowan DA (2010) On the rocks: the microbiology of Antarctic Dry Valley soils. *Nature Reviews Microbiology* 8:129. doi:10.1038/nrmicro2281
22. De Luca G, Barakat M, Ortet P, Fochesato S, Jourlin-Castelli C, Ansaldi M, Py B, Fichant G, Coutinho P, Voulhoux R, 285 Bastien O, Maréchal E, Henrissat B, Quentin Y, Noirot P, Filloux A, Méjean V, S DuBow M, Barras F, Heulin T (2011) The Cyst-Dividing Bacterium *Ramlibacter tataouinensis* TTB310 Genome Reveals a Well-Stocked Toolbox for Adaptation to a Desert Environment, vol 6. doi:10.1371/journal.pone.0023784
23. Busarakam K, Bull AT, Trujillo ME, Riesco R, Sangal V, van Wezel GP, Goodfellow M (2016) *Modestobacter caceresii* sp. nov., novel actinobacteria with an insight into their adaptive mechanisms for survival in extreme hyper-arid Atacama Desert 290 soils. *Systematic and Applied Microbiology* 39 (4):243-251. doi:<https://doi.org/10.1016/j.syapm.2016.03.007>
24. Stieglmeier M, Klingl A, Alves RJE, Rittmann SKMR, Melcher M, Leisch N, Schleper C (2014) *Nitrososphaera viennensis* gen. nov., sp. nov., an aerobic and mesophilic, ammonia-oxidizing archaeon from soil and a member of the archaeal phylum Thaumarchaeota. *Int J Syst Evol Microbiol* 64 (Pt 8):2738-2752. doi:10.1099/ijms.0.063172-0

25. Schrauzer GN (1978) Desert sands catalyze ammonia formation. *Chemical & Engineering News Archive* 56 (46):7.
295 doi:10.1021/cen-v056n046.p007a
26. Matthews A, Pierce S, Hipperson H, Raymond B (2019) Rhizobacterial Community Assembly Patterns Vary Between
Crop Species. *Frontiers in microbiology* 10:581-581. doi:10.3389/fmicb.2019.00581
27. Eberspächer J, Lingens F (2006) The Genus *Phenylobacterium*. In: Dworkin M, Falkow S, Rosenberg E, Schleifer K-H,
Stackebrandt E (eds) *The Prokaryotes: Volume 5: Proteobacteria: Alpha and Beta Subclasses*. Springer New York, New York,
300 NY, pp 250-256. doi:10.1007/0-387-30745-1_13


305

310

315

320

325




330

Figure 1: Location of sampling sites (A) and the climatic conditions at these sampling sites (B).

335


340



345

350



355



360

365

Figure 2: Estimates of Alpha diversity. Samples are rarified to read depth of 68640. Panel A shows the rarefaction curves obtained determining the species richness. While panel B shows the Shannon diversity indices for the desert soil samples studied.




375

Figure 3: Analysis of similarity (ANOSIM) within and between samples (A). The abundance of different phyla in desert soil samples studied and their comparison with the hot (Namib) and cold (Antarctica) desert samples are shown in panel B. Data for Antarctic desert and Namib desert are taken from Armstrong et al., 2016 and Cary et al., 2010, respectively.

380

A



B





Figure 4: Venn diagram showing the number of genera present in different desert soil samples (A). Heat map showing 20 most dominant genera present in the desert soil samples. M5 & M15; Samples collected from Muzamiyah, HB; sample collected from Hafr Al-Batin.



Figure 5: Predicted functional gene abundance in four different samples obtained through PICRUSt analysis. M5 & M15; Samples collected from Muzamiyah, HB; sample collected from Hafr Al-Batin.



400

Figure 6. Statistical comparison (Welch's *t*-test) of the predicted function gene abundance between M5 and M15 samples obtained from Mazamiyah (A) and between ABHA and Haftral Batin (B). P-value correction; $P < 0.05$.

405

Table 1. The properties of soil samples collected for the study

Properties	Abha	Muzamiyah		Hafr Al-Batin
		M15	M5	
Coordinates	18°15'24.4"N	24°24'43.6"N		28°02'05.3"N
	42°32'45.4"E	46°15'31.4"E		45°44'03.3"E
Date of Collection	28 January 2017	6 February 2017		13 February 2017
pH	7.9	8.2		8.4
Total Aerobic Count on NA (CFU/g of soil)	$3.63 \pm 1.9 \times 10^5$	$1.1 \pm 0.9 \times 10^4$	$5.5 \pm 1.9 \times 10^5$	$8.1 \pm 2.7 \times 10^4$
*Average yearly Temp (°C)	17.5	25.3		24.8
Maximum Temperature (°C)	30.5	44		45
*Average yearly rain fall (mm)	230	88		126
*Soil type	Sand and silty	Sandy loam		Sand and Gravel
† <i>Other deserts</i>	pH	Temperature		precipitation
		(°C)		(mm/yr)
Namib desert	7.9-8.5	5 to 45		5-100
Antarctica desert	-	-15 to -30		Less than 100

*Al-Zahrani 2017; Tarawneh other literature, †Makhalanyane et al., 2015; Cary et al., 2010

- Aislabie, J. M., Chhour, K.-L., Saul, D. J., Miyauchi, S., Ayton, J., Paetzold, R. F., and Balks, M. R.: Dominant bacteria in soils of Marble Point and Wright Valley, Victoria Land, Antarctica, *Soil Biology and Biochemistry*, 38, 3041-3056, <https://doi.org/10.1016/j.soilbio.2006.02.018>, 2006.
- 415 Andrew, D. R., Fitak, R. R., Munguia-Vega, A., Racolta, A., Martinson, V. G., and Dontsova, K.: Abiotic Factors Shape Microbial Diversity in Sonoran Desert Soils, 78, 7527-7537, 10.1128/AEM.01459-12 %J *Applied and Environmental Microbiology*, 2012.
- 420 Armstrong, A., Valverde, A., Ramond, J.-B., Makhalanyane, T., Jansson, J., Hopkins, D., Aspray, T., Seely, M., Tuffin, M., and Cowan, D.: Temporal dynamics of hot desert microbial communities reveal structural and functional responses to water input, *Scientific Reports*, 6, 34434, 10.1038/srep34434, 2016.
- Busarakam, K., Bull, A. T., Trujillo, M. E., Riesco, R., Sangal, V., van Wezel, G. P., and Goodfellow, M.: *Modestobacter caceresii* sp. nov., novel actinobacteria with an insight into their adaptive mechanisms for survival in extreme hyper-arid Atacama Desert soils, *Systematic and applied microbiology*, 39, 243-251, 10.1016/j.syapm.2016.03.007, 2016.
- 425 Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., Fierer, N., Pena, A. G., Goodrich, J. K., Gordon, J. I., Huttley, G. A., Kelley, S. T., Knights, D., Koenig, J. E., Ley, R. E., Lozupone, C. A., McDonald, D., Muegge, B. D., Pirrung, M., Reeder, J., Sevinsky, J. R., Turnbaugh, P. J., Walters, W. A., Widmann, J., Yatsunenko, T., Zaneveld, J., and Knight, R.: QIIME allows analysis of high-throughput community sequencing data, *Nature methods*, 7, 335-336, 10.1038/nmeth.f.303, 2010.
- 430 Cary, S. C., McDonald, I. R., Barrett, J. E., and Cowan, D. A.: On the rocks: the microbiology of Antarctic Dry Valley soils, *Nature Reviews Microbiology*, 8, 129-138, 10.1038/nrmicro2281, 2010.
- De Luca, G., Barakat, M., Ortet, P., Fochesato, S., Jourlin-Castelli, C., Ansaldi, M., Py, B., Fichant, G., Coutinho, P. M., Voulhoux, R., Bastien, O., Marechal, E., Henrissat, B., Quentin, Y., Noirot, P., Filloux, A., Mejean, V., DuBow, M. S., Barras, F., Barbe, V., Weissenbach, J., Mihalcescu, I., Vermeglio, A., Achouak, W., and Heulin, T.: The Cyst-Dividing Bacterium *Ramlibacter tataouinensis* TTB310 Genome Reveals a Well-Stocked Toolbox for Adaptation to a Desert Environment, *PLoS ONE*, 6, e23784, 2011.
- 435 Eberspächer, J., and Lingens, F.: The Genus *Phenylobacterium*, in: *The Prokaryotes: Volume 5: Proteobacteria: Alpha and Beta Subclasses*, edited by: Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K.-H., and Stackebrandt, E., Springer New York, New York, NY, 250-256, 2006.
- 440 Fiaz, S., Noor, M. A., and Aldosari, F.: Achieving food security in the Kingdom of Saudi Arabia through innovation: Potential role of agricultural extension, *Journal of the Saudi Society of Agricultural Sciences*, 17, 10.1016/j.jssas.2016.09.001, 2016.
- Fierer, N., Leff, J. W., Adams, B. J., Nielsen, U. N., Bates, S. T., Lauber, C. L., Owens, S., Gilbert, J. A., Wall, D. H., and Caporaso, J. G.: Cross-biome metagenomic analyses of soil microbial communities and their functional attributes, 109, 21390-21395, 10.1073/pnas.1215210110 %J *Proceedings of the National Academy of Sciences*, 2012.
- 445 Fierer, N.: Embracing the unknown: disentangling the complexities of the soil microbiome, *Nature Reviews Microbiology*, 15, 579-590, 10.1038/nrmicro.2017.87, 2017.
- Holm, D. A.: Desert Geomorphology in the Arabian Peninsula, 132, 1369-1379, 10.1126/science.132.3437.1369 %J *Science*, 1960.
- Makhalanyane, T. P., Valverde, A., Gunnigle, E., Frossard, A., Ramond, J. B., and Cowan, D. A.: Microbial ecology of hot desert edaphic systems, *FEMS microbiology reviews*, 39, 203-221, 10.1093/femsre/fuu011, 2015.
- 450 Marasco, R., Rolli, E., Fusi, M., Michoud, G., and Daffonchio, D.: Grapevine rootstocks shape underground bacterial microbiome and networking but not potential functionality, *Microbiome*, 6, 3, 10.1186/s40168-017-0391-2, 2018.
- Matthews, A., Pierce, S., Hipperson, H., and Raymond, B.: Rhizobacterial Community Assembly Patterns Vary Between Crop Species, 10, 10.3389/fmicb.2019.00581, 2019.
- 455 Meola, M., Lazzaro, A., and Zeyer, J.: Bacterial Composition and Survival on Sahara Dust Particles Transported to the European Alps, *Frontiers in microbiology*, 6, 1454-1454, 10.3389/fmicb.2015.01454, 2015.
- Ondov, B. D., Bergman, N. H., and Phillippy, A. M.: Krona: Interactive Metagenomic Visualization in a Web Browser, in: *Encyclopedia of Metagenomics*, edited by: Nelson, K. E., Springer New York, New York, NY, 1-8, 2013.
- Parks, D. H., Tyson, G. W., Hugenholtz, P., and Beiko, R. G.: STAMP: statistical analysis of taxonomic and functional profiles, *Bioinformatics (Oxford, England)*, 30, 3123-3124, 10.1093/bioinformatics/btu494, 2014.
- 460 Robe, P., Nalin, R., Capellano, C., Vogel, T. M., and Simonet, P.: Extraction of DNA from soil, *European Journal of Soil Biology*, 39, 183-190, [https://doi.org/10.1016/S1164-5563\(03\)00033-5](https://doi.org/10.1016/S1164-5563(03)00033-5), 2003.

- Santamaria, M., Fosso, B., Consiglio, A., De Caro, G., Grillo, G., Licciulli, F., Liuni, S., Marzano, M., Alonso-Aleman, D., Valiente, G., and Pesole, G.: Reference databases for taxonomic assignment in metagenomics, *Briefings in Bioinformatics*, 13, 682-695, 10.1093/bib/bbs036 %J Briefings in Bioinformatics, 2012.
- 465 Schrauzer, D. G. N.: Desert sands catalyze ammonia formation, *Chemical & Engineering News Archive*, 56, 7, 10.1021/cen-v056n046.p007a, 1978.
- Schulze-Makuch, D., Wagner, D., Kounaves, S. P., Mangelsdorf, K., Devine, K. G., de Vera, J.-P., Schmitt-Kopplin, P., Grossart, H.-P., Parro, V., Kaupenjohann, M., Galy, A., Schneider, B., Airo, A., Frösler, J., Davila, A. F., Arens, F. L., Cáceres, L., Cornejo, F. S., Carrizo, D., Dartnell, L., DiRuggiero, J., Flury, M., Ganzert, L., Gessner, M. O., Grathwohl, P., Guan, L.,
470 Heinz, J., Hess, M., Keppler, F., Maus, D., McKay, C. P., Meckenstock, R. U., Montgomery, W., Oberlin, E. A., Probst, A. J., Sáenz, J. S., Sattler, T., Schirmack, J., Sephton, M. A., Schlöter, M., Uhl, J., Valenzuela, B., Vestergaard, G., Wörmer, L., and Zamorano, P.: Transitory microbial habitat in the hyperarid Atacama Desert, 115, 2670-2675, 10.1073/pnas.1714341115 %J Proceedings of the National Academy of Sciences, 2018.
- Siham, A.-K.: Soil Analysis of Contaminated Soil from Riyadh City, Saudi Arabia and Influence of Aluminium and Cobalt
475 Ions on the Growth of Fungi Isolated, *Journal of Biological Sciences*, 7, 10.3923/jbs.2007.549.553, 2007.
- Stieglmeier, M., Klingl, A., Alves, R. J., Rittmann, S. K., Melcher, M., Leisch, N., and Schleper, C.: *Nitrososphaera viennensis* gen. nov., sp. nov., an aerobic and mesophilic, ammonia-oxidizing archaeon from soil and a member of the archaeal phylum Thaumarchaeota, *International journal of systematic and evolutionary microbiology*, 64, 2738-2752, 10.1099/ijs.0.063172-0, 2014.
- 480 Yuan, H.-Y., Liu, P.-P., Wang, N., Li, X.-M., Zhu, Y.-G., Khan, S. T., Alkhedhairy, A. A., and Sun, G.-X.: The influence of soil properties and geographical distance on the bacterial community compositions of paddy soils enriched on SMFC anodes, *Journal of Soils and Sediments*, 18, 517-525, 10.1007/s11368-017-1769-2, 2018.
- Zakrzewski, M., Proietti, C., Ellis, J. J., Hasan, S., Brion, M. J., Berger, B., and Krause, L.: Calypso: a user-friendly web-server for mining and visualizing microbiome-environment interactions, *Bioinformatics (Oxford, England)*, 33, 782-783,
485 10.1093/bioinformatics/btw725, 2017.
- Zeng, Y., Feng, F., Medova, H., Dean, J., and Koblizek, M.: Functional type 2 photosynthetic reaction centers found in the rare bacterial phylum Gemmatimonadetes, *Proc Natl Acad Sci U S A*, 111, 7795-7800, 10.1073/pnas.1400295111, 2014.