

Geogenic organic carbon in terrestrial sediments and its contribution to total soil carbon

Fabian Kalks^{a*}, Gabriel Noren^b, Carsten W. Mueller^{cd}, Mirjam Helfrich^a, Janet Rethemeyer^b, Axel Don^a

^a Thünen Institute of Climate-Smart Agriculture, Bundesallee 65, 38116 Braunschweig, Germany

^b Institute of Geology and Mineralogy, University of Cologne, Institute of Geology and Mineralogy, Zùlpicher Str. 49b, 50674 Cologne, Germany

^c Technical University of Munich, Chair of Soil Science, Emil-Ramann Strasse 2, 85354 Freising-Weihenstephan, Germany

^d University of Copenhagen, Department of Geosciences and Natural Resource Management, Øster Voldgade 10, DK-1350 Copenhagen K, Denmark

*Corresponding author: axel.don@thuenen.de + 49 531 596 2641

Abstract

Geogenic organic carbon (GOC) from sedimentary rocks is an overlooked fraction in soils that has not yet been quantified, but influences the composition, age and stability of total organic carbon (OC) in soils. In this context, GOC is the OC in bedrocks deposited during sedimentation. However, the contribution of GOC to total soil OC varies depending on the type of bedrock. As yet, no far studies have investigated the contribution of GOC derived from different terrestrial sedimentary rocks to soil OC contents.

In order to fill this knowledge gap, 10-m long sediment cores from three sites recovered from Pleistocene Loess, Miocene Sand and Triassic Red Sandstone were analysed and the amount of GOC calculated based on ^{14}C measurements. The ^{14}C ages of bulk sedimentary OC revealed that OC is comprised of both biogenic and geogenic components. The biogenic component relates to OC that recently entered the sediments from plant sources. All the sediments contained considerable amounts of GOC (median amounts of 0.10 g kg^{-1} in Miocene Sand, 0.27 g kg^{-1} in Pleistocene Loess and 0.17 in Red Sandstone) compared with subsoil OC contents (between 0.53 - 15.21 g kg^{-1}). Long-term incubation experiments revealed that the GOC appeared comparatively stable against biodegradation. Its possible contribution to subsoil OC stocks (0.3-1.5 m depth) ranged from 1 to 26 % in soil developed in the Miocene Sand, from 16 to 21 % in the Loess soil and from 6 to 36 % at the Red Sandstone site. Thus, GOC with no detectable ^{14}C content influenced the ^{14}C ages of subsoil OC, and may partly explain the strong increase in ^{14}C ages observed in many subsoils. This could be particularly important in young soils on terrestrial sediments with comparatively low amounts of OC, where GOC can make a large contribution to total OC stocks.

Keywords Geogenic organic carbon; sedimentary organic carbon; ^{14}C ; terrestrial sediments; incubation experiment

1. Introduction

On average, the world's soils store more than 50 % of OC in the subsoil below 30 cm depth (Batjes, 2014). This type of carbon is considered a highly stable carbon pool due to its apparently high ^{14}C ages (Mathieu et al., 2015, Schrumpf et al., 2013). However, another explanation for this could be the contribution of geogenic organic carbon (GOC), which is defined here as OC that originates from deposition during sedimentation and rock formation, and may increasingly influence subsoil OC with depth (Graz et al., 2010, Kögel-Knabner et al., 2008, Schrumpf et al., 2013, Trumbore, 2009). GOC in most cases is devoid of ^{14}C and thus may lead to an overestimation of ancient OC sources although a number of studies showed the importance of root derived, young OC inputs to subsoils (Angst et al., 2016, Crow et al., 2009). Therefore GOC may significantly influence and affect the overall ^{14}C signal, particularly in OC-poor subsoils. Vindušková et al. (2015) investigated the contribution of GOC to soils in reclaimed mine soils, and found GOC contributions to total soil OC of between 26 and 99 %. Furthermore OC-rich sediments with contents of 2-7 g kg⁻¹ (Hemingway et al., 2018) and 28-105 g kg⁻¹ (Frouz et al., 2011) have been investigated with regard to the stability of OC in these sediments, but no conclusion reached about GOC contributions in soils. However, the impact of GOC on soils derived from sediments or sedimentary rocks with lower OC contents has not yet been investigated. Considering that approximately 65 % of the continental earth's surface is covered with sediments and sedimentary rocks (Amiotte Suchet et al., 2003), a potentially large fraction of soils could contain GOC that contributes to soil OC stocks, even though a large portion might be derived from recent sedimentation processes. There is not yet much literature about sediments containing only low amounts of OC. There are estimations that assume sandstones to be free of GOC (van der Voort et al., 2018) or, in contrast, a storage model that assumes generally high GOC amounts of 2.4 g kg⁻¹ for all sandy deposits (Copard et al., 2007). Therefore, more information about the amounts of GOC in sediments is needed.

To estimate the possible contribution of GOC to subsoil OC stocks, it is important to establish the amount of OC in sediments that comes from sedimentation (GOC) and to distinguish it from OC derived from current vegetation (biogenic OC). There are many soil- and substrate-specific factors that might influence the OC contribution from current vegetation to sedimentary OC, such as potential rooting depth or pore distribution. No method has yet been established to allow a direct quantification of GOC in different soils or sediments, apart from promising methods to quantify the graphitic part of GOC in soils (Zethof et al., 2019). The only reliable approach to distinguish between both sources is the use of ^{14}C . Since deposition of sediments mostly took place > 50,000 years BP, they do not contain ^{14}C , which has a mean half-life time of 5,730 years (Libby, 1952). In addition, the $\delta^{13}\text{C}$ values of OC in the sediments allow carbonaceous sources with $\delta^{13}\text{C}$ values around 0 ‰ to be distinguished from organic sources with $\delta^{13}\text{C}$ values < -22 ‰. Thus, the use of both carbon isotopes could reveal whether the OC is a mixture of GOC and OC from vegetation that is less than 50,000 years old.

One important question about the possible contribution of GOC in soils is whether the GOC is mineralised when it becomes part of the soil. As GOC resists degradation once it has been deposited, it can be assumed that it already exhibits a strong inherent recalcitrance. Nevertheless, this could also be due to a physical protection that prevents microbial accessibility. However, when it becomes part of the subsoil during soil development, this OC pool could be degraded by the infiltration of water, oxygen, fresh nutrients and microorganisms. Direct microbial coal degradation has already been observed in incubation experiments on mine soils (Rumpel and Kögel-Knabner, 2002, Waschies and Huttli, 1999) or shale bedrocks directly exposed to the surface (Soulet et al., 2017). There has been no study to establish whether GOC is degradable in OC-poor sediments or sedimentary rocks has not been investigated, but it could differ since the amount of available OC can also drive microbial respiration (Colman and Schimel, 2013). Therefore, these sediments might contain fewer microorganisms that can be spatially separated from the GOC, which may hamper its respiration.

To the best of the author's knowledge, there has only been one study by van der Voort et al. (2018) that has investigated the amount of GOC in soils. They estimated that GOC makes up around 80 % of soil OC in a moraine-derived soil, suggesting that GOC's contribution to soil OC is large. However, apart from this study by van der Voort et al. (2018) on a very specific sediment, there have been no direct calculations of the amount of GOC in soils.

The aim of this study was to quantify GOC in different terrestrial sediments and a sedimentary rock, and investigate its stability in incubation experiments in order to make assumptions about its possible contribution to soil OC stocks in soil profiles at the same site. The main research questions were: i) What is the relationship between sedimentary and subsoil OC contents? ii) Is OC in sediments ¹⁴C-free and how much is really geogenic? iii) Will sedimentary GOC be degraded? and iv) How much does GOC contribute to soil OC?

2. Material and methods

2.1 Site description

Three sites were selected with different sedimentary bedrocks derived from a single geologic substrate that can be found close to the surface and is homogeneous down to 10 m depth. The sites represented one sedimentary rock and two soft sediments. The sedimentary rock was a sandstone (Solling Formation, Triassic) under European beech forest (*Fagus sylvatica*) 11.5 km north-east of Göttingen (51°35.012' N; 10°3.960' O), referred to below as “Red Sandstone”. The soil is classified as a Folic Brunic Arenosol according to the World Reference Base for Soil Resources (WRB, 2006). The sediments were loessic deposits (Weichselian Glacial) that have been under agricultural land use for the past decades, 30 km north of Göttingen (51°48.101 N; 9°58.002' O), referred to as “Loess”, and terrestrial sandy deposits from the Miocene (Neogene formerly named Tertiary) in a European beech forest 13 km south-west of Göttingen (51°28.673 N; 9°45.323' O) referred to as “Miocene Sand”. The respective soils have been classified as a Haplic Luvisol and a Dystric Chromic Arenosol accordingly. The sediments at the Loess site were deposited between the last glacial and interglacial periods between 115,000 and 400,000 years BP, according to Jordan and Schwartau (1993). To the best of the author’s knowledge, these forest sites have never been under agricultural use. The associated soils are classified as a Luvisol and a Cambisol respectively. Mean annual air temperature and precipitation were 9.2 °C and 647 mm (1981-2010) according to the nearby weather station that covers all three sites.

2.2 Sampling and sample preparation

Two 10-m long sediment cores with a diameter of 15 cm were drilled at each site in April 2017. For the soft sediments (Loess and Miocene Sand), drilling was conducted as percussion drilling, and for the hard sediment (Red Sandstone) as cable core drilling with water as the flushing solution. The replicates per site were drilled approximately 10 m apart. The sampled cores were subdivided into 1-m increments. For further chemical analysis, material was taken from a depth of 85-95 cm for each 1-m increment, and the outer 5 cm removed to avoid possible contamination. Thus the increment from 1-2 m for example, was represented by a sample from a 1.85-1.95 m depth.

One sample from each depth increment (1-2 m, 2-3 m, 3-4 m, 4-5 m, 5-6 m, 6-7 m, 7-8 m, 8-9 m, 9-10 m) for chemical analysis was oven dried at 60 °C and sieved to pass 2 mm. The Red Sandstone samples were crushed with a hammer before being dried and sieved. Additionally, approximately 1-m deep soil profiles were dug and soil samples taken from the different classified layers to obtain corresponding soil parameters (Fig. 1).




Fig. 1: Respective soil profiles at the Loess (A), Miocene Sand (B) and Red Sandstone (C) sites. Soil classification was conducted using the German classification system for soil horizons (Eckelmann et al., 2006) and later transferred to the WRB (2006). Depth transitions to the starting sediment were 82 cm for Loess (Al + Bt + elCv), 72 cm for Miocene Sand (Bv - ilCv) and 68 cm for Red Sandstone (ilCv₁).

Samples were oven dried at 60°C and stored and sieved to pass 2 mm. For the determination of OC and ¹³C, samples were ground in a planetary ball mill.

2.3 Chemical analysis and calculations

Three aliquots of each sieved sample were analysed by dry combustion for total C and total N content (TruMac CN LECO, St. Joseph, MI, USA). Samples with a pH value of > 6 (measured in 0,01 mol L⁻¹ CaCl) were analysed for carbonates after the sample was ignited for 16 h in a muffle kiln at 450 °C according to Nelson and Summers (1983), to remove the organic part of total C. The OC concentration was calculated by subtracting carbonaceous C from total C, expressed as g OC kg⁻¹ dry matter. Homogenised samples were further analysed for δ¹³C values after carbonate removal in an isotope ratio mass spectrometer (Delta Plus, Thermo Fisher, Waltham, MA, USA) coupled to an elemental analyser (FLASH EA 1122 NA 1500; Wigan, United Kingdom). Resulting δ¹³C values (‰) were expressed relative to the international standard of Vienna Pee Dee Belemnite. The bulk densities for the soil samples were obtained with 250 cm³ sampling rings from each layer of the soil profile. For the sedimentary Loess and Miocene Sand samples, the bulk density of the deepest soil sample (94 and 100 cm respectively) was used. The bulk densities and densities without pore space of the intact Red Sandstone cores were determined on four subsamples (from 1.6, 3.6, 7 and 9 m depth) with a Dryflow-pycnometer (GeoPyc 1360) and a gas pycnometer (AccuPyc 1330) respectively. Bulk densities for the missing depth increments were linearly interpolated. For radiocarbon (¹⁴C) analysis, the sediment samples were first treated with 1% HCl to remove inorganic C, and then transferred into pre-combusted

quartz ampoules containing copper oxide and silver wool. The ampoules were evacuated, flame-sealed, then combusted at 900 °C, and the CO₂ evolved was purified on a vacuum ring (Rethemeyer et al., 2019). The ¹⁴C contents were measured with the MICADAS accelerator mass spectrometry (AMS) system at ETH Zürich, Switzerland. Where possible, one sedimentary sample per depth increment and site and one sample per soil layer was analysed. Due to the very low OC contents in some sediment samples, ¹⁴C contents could only be determined for three samples from the Miocene Sand (from 1.9, 4.9 and 7.9 m depth) and four from the Red Sandstone (1.9, 4.9, 7.9 and 9.9 m depth). For the Loess, ¹⁴C of bulk OC was measured in all depth intervals (1.9-9.9 m).

Total OC stocks (Mg ha⁻¹) were calculated according to Eq. 1:

$$OC\ stock = OC \cdot BD \cdot (1 - stone\ content) \cdot depth \cdot 0.1 \quad Eq. 1$$

where *OC* is the weight-based OC content, either in the fine soil <2-mm fraction of the soil profiles or in the sediments (g kg⁻¹), *BD* is the bulk density of the fine soil (g cm⁻³), stone content is the volume-based proportion of stones (cm³ cm⁻³) and depth is the thickness of the depth increment (cm). To be able to compare OC stocks and contributions from GOC, it was decided to set the borders between the topsoils- and subsoils at 0.3 m and the transition from subsoils to the sediments at 1.5 m. According to Richter and Markewitz (1995) this represents a common border for the transition from soil to sediment. The sediments were further subdivided into an upper and a lower part at a depth of 4 m.

In a second step, the amount of GOC and biogenic OC in the sediments was calculated, considering GOC as one carbon pool free of ¹⁴C. For the sediments, the proportion of biogenic OC (*f_{biogenic}*) of the total amount of OC was calculated in a two pool model (Eq. 2) used by Cerri et al. (1985):

$$f_{biogenic} (\%) = \frac{F_{biogenic\ OC} - F_{GOC}}{F_{sample} - F_{GOC}} \cdot 100 \quad Eq. 2$$

where *F* represents the ¹⁴C content in the fraction modern carbon (F¹⁴C) from a source compared with the ¹⁴C content of an oxalic standard (Stuiver and Polach, 1977, Torn et al., 2009). The sources were the GOC fraction (*F_{GOC}*), the sample (*F_{sample}*) and the biogenic OC fraction (*F_{biogenic OC}*). Since the ¹⁴C content of the GOC fraction can be set to zero, this equation could be simplified to:

$$f_{biogenic} (\%) = \frac{F_{biogenic\ OC}}{F_{sample}} \cdot 100 \quad Eq. 3$$

For the biogenic OC in the sediments, an average ¹⁴C age ranging from 1,000-4,000 years BP was assumed, based on published ¹⁴C results of dissolved OC reaching greater depths (Artinger et al., 1996, Schiff et al., 1997). The ¹⁴C contents in the sediment from 2 to 4 m depth of the loess led to ages < 3,000 years BP, and were therefore even younger than at 74 cm depth (4,413 years BP). Thus, they were treated

like the soil part for the calculation below of a GOC fraction. Respective times were converted into ^{14}C contents ($F_{\text{biogenic OC}}$) according to Torn et al. (2009):

$$F_{\text{biogenic OC}} = e^{\left(\frac{t}{-8033}\right)} \quad \text{Eq. 4}$$

where t represents the ^{14}C age (1,000 or 4,000 years BP respectively) and 8033 represents the mean life of radiocarbon in years. The proportion of GOC in the sediments (f_{GOC}) is therefore the portion left (Eq. 5):

$$f_{\text{GOC}} = 100 \% - f_{\text{biogenic}} \quad \text{Eq. 5}$$

For the depth increments without measured ^{14}C ages (Tab. S1), the calculated amounts were linearly interpolated with measured ^{14}C ages from samples above and below. This was done by assuming a depth-dependent correlation and using the adjacent values. To calculate the amount of GOC in the soil profiles, the weight-based amount of GOC in the sediments was first calculated by multiplying its fraction (f_{GOC}) by the respective OC content (in g OC per kg^{-1} dry mass). The median amount of GOC (g GOC per kg^{-1} dry mass) of these sedimentary values was then taken and its proportion of soil OC content calculated (g OC per kg^{-1} fine soil) in the soil profile. This was done for the proportion of GOC in the sediments calculated with a 1,000 and a 4,000-year-old biogenic OC fraction ($F_{\text{biogenic OC}}$ in Eq. 3) to obtain a range of GOC contributions. It was assumed that the GOC fraction resisted degradation during soil formation. Therefore, this proportion represents the largest possible amount of GOC that could contribute to soil OC stocks. On this assumption, it was also possible to define the influence of GOC in the soil profile on the resulting ^{14}C ages. Since the calculated ^{14}C ages represent a mixture of the ^{14}C content from GOC and the biogenic fraction (Eq. 5), the GOC fraction has the same influence on the soil ^{14}C age as on bulk OC (according to Eq. 3). Reducing the age by this fraction would therefore represent an “unbiased” age of soil-derived-OC.

2.4 Incubation experiment

To assess the potential stability of OC in the sediments against microbial decay, two laboratory incubation experiments were conducted at 20 °C for 50 and 63 days respectively. This was done to reveal the potential degradation of OC from the sediments. A temperature of 20 °C was chosen because only very low degradation rates were expected at lower temperatures. The first, 50-day experiment was conducted with intact Red Sandstone core samples, while the second experiment was performed after the Red Sandstone was crushed to pieces < 2 mm to simulate the process of weathering when the intact sediment or sedimentary rock becomes part of the (sub)soil. For the incubations, four subsamples of 1,340-6,890 g per sample from different depth intervals were used. The sample material was stored at room temperature until the start of the incubation experiment. Four samples were taken from each sediment and from each of four depth ranges for Miocene Sand (1.2-2.8, 3.2-4.8, 6.1-7.8 and 8.2-9.9 m), Red Sandstone (2-3, 4-5.8, 7-8 and 8.4-9.8 m) and Loess (1.4-2, 2.7-3, 4.8-6 and 9.1-10 m). Water

content was adjusted to correspond to 40 % of water-holding capacity based on the poured bulk density, determined by filling the loose material into a defined volume and measuring its weight. The material for the four repetitions was mixed in large plastic vats and water added before the respective four subsamples were transferred to incubation vessels. There were also four blank samples with no material. Based on preliminary tests and their calculated bulk density and porosity, the intact Red Sandstone samples were kept in a barrel with pure water for 14 hours to reach a water content of nearly 40 %. Samples were placed in polycarbonate vessels with a volume of 7069 cm³ and made airtight. The lids contained two tube connectors so that the samples could be flushed with ambient air. After flushing, samples were set to a starting pressure of about 1,300 mbar and kept closed until the end of the incubation. Nine gas samples were taken in evacuated glass vials (20 mL), 0, 3, 7, 13, 21, 30, 59 and 63 days after the start of incubation. Samples were analysed for CO₂ concentrations by gas chromatography (Agilent 7890A, GC, Agilent Technologies, Santa Clara, USA) to account for the amount of accumulated CO₂. Three additional gas samples were taken 0, 30 and 63 days after the incubation started and analysed with an isotope ratio mass spectrometer (Delta Plus XP, Thermo Fisher Scientific, Bremen, Germany) to account for the development of $\delta^{13}\text{C}$ of CO₂ during the respiration. The corresponding pressure was measured on each sampling date. When the over-pressure of a vessel was lost due to leakages, it was removed from the sampling because contamination with ambient air could no longer be ruled out. This happened for a third of all the samples.

The amount of respired CO₂-C (mg CO₂-C d⁻¹) was calculated using Eq 6.

$$CO_2 - C = \frac{0.1 \cdot p \cdot x_i \cdot M \cdot V}{R \cdot T \cdot t} \quad \text{Eq. 6}$$

where p is the pressure (mbar), x_i is the difference in CO₂ concentration between the samplings (ppm), M is the molar mass of C (g mol⁻¹), V is the air volume of the sample (m³), R is the molar gas constant (J kmol⁻¹ K⁻¹), T is the incubation temperature (K) and t is the elapsed time (d) between the samplings. Based on the $\delta^{13}\text{C}$ values of CO₂ the proportion of CO₂ derived from carbonates was subtracted where necessary according to Bertrand et al. (2007). This respiration rate was related to the OC content of the samples (called “OC-normalised respiration”) by being divided by the total amount of OC in g in the sample.

Since we observed an almost linear respiration behaviour was observed in both incubation experiments, a simple linear regression model was fitted to describe the mineralisation rate per time (mineralised OC (%)) = $k \cdot t$.

2.5 Statistics

Statistical analyses were conducted using the statistical environment R (R Core Team (2018)) including the function “*lm*” to fit linear models and the package *ggplot2* (Wickham, 2016) for graphical presentation. The models were tested for deviations from homoscedasticity, normality of residuals and absence of collinearity. The tests revealed heteroscedasticity of the residuals, which can be explained by an increasing standard deviation with time. It subsequently became apparent that the residuals were not normally distributed since the dependent variable, representing the proportion of OC being mineralised, only allows values between 0 and 1. Bearing this in mind, the results should be treated as an indicator of differences between the samples and a scale for the mineralisation of the OC pool. Therefore, calculations of standard errors and significance of the parameters were omitted since this would not produce reasonable results with the model used.

3. Results

3.1 Relationship between sedimentary and subsoil organic carbon

Organic carbon was detectable in all the sediments analysed using three laboratory replicates per sample and comparing them with the muffled samples, measurable. The mean relative standard deviation of the laboratory replicates was 9.5 %. In terms of the detection limit, the sample with the lowest total C content (mean of 0.04 g C kg⁻¹ soil) showed values between 0.00 and 0.01 g C kg⁻¹ soil after removal of OC at 450 °C. Thus the range of 0.00 to 0.01 g C kg⁻¹ soil was assumed to be the mean standard error from the measurement. Despite having the same material down to 10 m depth at each site, there were still some inhomogeneities that were visible and measurable. This was especially true for Loess and Red Sandstone. The amount of OC in the sediments from 1 to 10 m depth was comparatively low in Miocene Sand and Red Sandstone (0.04-0.71 g kg⁻¹ and 0.01-0.53 g kg⁻¹, respectively), while much higher OC contents of 0.21-9.71 g C kg⁻¹ were found in Loess (Fig. 2 a). The median OC content of the sediments was within a range comparable to that of the respective deepest subsoil horizon. This deepest horizon was a Cv horizon at 94 cm depth for Loess, 100 cm depth for Miocene Sand and 74 cm depth for Red Sandstone. In detail, the median OC content in the sediments, compared with the respective Cv horizons, corresponded to 27 % for Loess, 29 % for Miocene Sand and 39 % for Red Sandstone. The Loess OC contents were highly variable, highlighting the changing sedimentary conditions during the past glacial and interglacial periods (Jordan and Schwartau, 1993). At 4-5 m depth, the OC contents of Loess were even higher (9.7 g kg⁻¹) than in the subsoil (3 g kg⁻¹). This was not an outlier because the high OC content could be visually confirmed by the very dark colour of the sample. In Miocene Sand and Red Sandstone, no clear depth gradient of OC was found at 2-10 m depth (Fig. 2 a). Even though the OC content in the sediments were low, the OC stocks could be very large. A comparison of OC stocks in topsoils (0-0.3 m), subsoils (0.3-1.5 m) and the sediments down to 10 m depth revealed quite high OC stocks in the sediments. For Loess, OC in the sediment contributed up to 71 % of the total OC amount while it was 51 % for Red Sandstone and 21 % for Miocene Sand (Table 1).

The distribution of the $\delta^{13}\text{C}$ values of OC in the soil and sediment profiles showed an increase in $\delta^{13}\text{C}$ with depth in the soil down to 1 m (Fig. 2 b). In contrast, the $\delta^{13}\text{C}$ values of OC in the sediments showed no clear trend with increasing depth, but they all were within the range of C_3 plant material. A value above -25 ‰ for Red Sandstone at 4 m depth could be explained by the high values of inorganic carbon (IC) at this depth. It can be assumed that decarbonisation of this sample was not completely successful. Unexpectedly high amounts of IC were found in parts of Red Sandstone, indicating the presence of calcareous deposits in this terrestrial material (Fig. 2 c). The Loess also included some distinct calcareous layers at 5 and 10 m depth, while there were only small amounts of around 0.1 mg IC g^{-1} soil at the other depths. This could be due to the fact that the investigated loess deposits belong to the “Leine Ilme Basin”, a region with aeolian loamy loess that has been decalcified during weathering and soil genesis (Wagner, 2011), while the contents in the soil profile can be explained by liming. No IC was present in Miocene Sand.




Fig. 2: Depth distribution of different bulk properties of the soil profiles and deep drilling cores. Presented parameters include the log scale organic carbon (OC) and inorganic carbon (IC) content (a and c) and the $\delta^{13}\text{C}$ values of the organic carbon (b) related to the amount of fine soil or dry mass respectively. Filled and unfilled symbols represent the two different cores. For the Loess, only one core could be analysed.

Table 1: OC stocks and proportions for the three sites down to 10 m depth. Proportions of biogenic and geogenic OC were calculated based on ^{14}C results and assumptions described in the material and methods section. Represented ranges are calculated based on the assumption of a 1,000 or 4,000-year-old biogenic OC fraction reaching the sediments.

Substrate	Layer	Depth (m)	TOC		OC stocks (Mg ha ⁻¹)				Proportion of OC (%) ^b			
			(Mg·ha ⁻¹)	(%) ^a	geogenic		biogenic		geogenic		biogenic	
					4,000 yrs.	1,000 yrs.	1,000 yrs.	4,000 yrs.	4,000 yrs.	1,000 yrs.	1,000 yrs.	4,000 yrs.
Loess	Topsoil	0.0 - 0.3	40.6	11	1.1 - 1.2		40 - 39		3 - 3		97 - 97	
	Subsoil	0.3 - 1.5	66.0	17	5.1 - 5.3		61 - 61		8 - 8		92 - 92	
	Upper Sediment	1.5 - 4.0	218.6	57	10.6 - 11.0		208 - 208		5 - 5		95 - 95	
	Lower Sediment	4.0 - 10.0	55.1	15	42.9 - 46.7		8 - 12		78 - 85		15 - 22	
Red Sandstone	Topsoil	0.0 - 0.3	22.2	30	0.6 - 0.7		22 - 22		3 - 3		97 - 97	
	Subsoil	0.3 - 1.5	13.7	19	1.6 - 1.7		12 - 12		12 - 12		88 - 88	
	Upper Sediment	1.5 - 4.0	18.7	25	13.1 - 14.9		4 - 6		70 - 80		20 - 30	
	Lower Sediment	4.0 - 10.0	19.3	26	14.1 - 15.7		4 - 5		73 - 82		18 - 27	
Miocene Sand	Topsoil	0.0 - 0.3	39.1	31	0.3 - 0.3		39 - 39		1 - 1		99 - 99	
	Subsoil	0.3 - 1.5	60.8	48	1.3 - 1.6		59 - 60		2 - 3		97 - 98	
	Upper Sediment	1.5 - 4.0	10.6	8	3.7 - 5.8		5 - 7		34 - 55		45 - 66	
	Lower Sediment	4.0 - 10.0	16.3	13	8.3 - 10.8		6 - 8		51 - 66		34 - 49	

309 ^a % of total OC stock from 0-10 m

310 ^b % of OC stock in the respective depth increment

311 3.2 Ages of organic carbon in soils and sediments and contributions from geogenic organic carbon

312 The ages of OC in the Loess soil profiles revealed a modern carbon signature (0 years BP) at 0.3 m
313 depth, with a sharp increase up to $4,413 \pm 51$ years BP at 0.7 m depth (Fig. 3 a). For the Red Sandstone
314 soil profile there was only an increase in the ages from a modern signature at 0.04 m to 532 ± 41 years
315 BP at 0.3 m depth. The Miocene Sand soil profile at 0.4 m depth showed an increase from $1,277 \pm 41$
316 to $1,771 \pm 44$ years BP at 0.6 m depth. Thus, OC of the subsoil (around 0.6 m depth) in Loess was more
317 than twice as old as in Miocene Sand. In contrast, Loess had a modern signature at 0.3 m, while the soil
318 developed in Red Sandstone showed an average age of 532 ± 41 years BP at 0.3 m depth.

319 The ages of OC in the sediments ranged from 2,200-30,730 years BP, with respective mean ages of
320 $9,077 \pm 3,234$ years BP for Miocene Sand, $13,674 \pm 9,632$ years BP for Loess, and $14,463 \pm 1,992$ years
321 BP for Red Sandstone. For all sediments, 11 out of 16 samples had a ¹⁴C content that led to an ¹⁴C age
322 older than 11,600 years BP, which we assumed to be the time after the latest glacial period when soil
323 development started (Litt et al., 2007). Therefore, the sediments contained a mixture of geogenic (¹⁴C-
324 free) and biogenic (with ¹⁴C) OC. Despite being the youngest sediment, Loess partly revealed the highest
325 apparent ¹⁴C ages up to $30,730 \pm 631$ years BP (Fig. 3 a). The ages of OC in the sediment of Red
326 Sandstone and Miocene Sand ranged from $12,940 \pm 132$ to $17,390 \pm 206$ years BP and from $6,750 \pm 86$
327 to $12,770 \pm 151$ years BP respectively, revealing no depth trend with age in the deeper sediment. The
328 calculated GOC fraction in the sediments was highest for Red Sandstone, ranging from 67 to 87 %, with
329 a mean of 77 % (Fig. 3 b). For the three Miocene Sand samples, the GOC fraction ranged from 29 to 77

% with a mean of 53 %. Loess showed a sharp increase at a depth of five metres where GOC contribution rising to between 71 to 98 % while it was only 5 to 19 % at the 2 to 4 m depth.

The calculated weight-based content of GOC in the sediment revealed a comparatively uniform distribution for all the sediments with depth, except the extremely high contents of Loess at 5 m depth (Fig. 3 c). The investigated sediment depths revealed quite a narrow range of GOC contents with $0.10 \pm 0.03 \text{ g kg}^{-1}$ for Miocene Sand, $0.17 \pm 0.12 \text{ g kg}^{-1}$ for Red Sandstone and $0.27 \pm 0.08 \text{ g kg}^{-1}$ for Loess.




Fig. 3: Depth distribution of apparent ^{14}C ages (in years BP) (a), GOC contribution to OC contents in the sediments (b) and resulting weight based amounts of GOC (c) in the sediments. The range in the contribution from GOC is due to the assumption of an average biogenic OC age of 4,000 (empty shapes) and 1,000 years (filled shapes).

3.3 Biodegradability of sedimentary derived organic carbon

The incubation experiment revealed a potential, but low biodegradability of OC for all samples, but without a clear depth gradient from 1 to 10 m (Fig. 4). To compare the effect of crushing on the respiration of Red Sandstone samples, the mineralisation rates from the first incubation experiment were compared with those of the second incubation experiment (Table 2). While Red Sandstone showed very low mineralisation when the samples were incubated as intact cores ($0.3\text{--}1.0 \text{ mg CO}_2\text{-C g}^{-1} \text{ OC y}^{-1}$), the mineralisation rate constants were up to five times higher when the samples were crushed ($1.0\text{--}2.1 \text{ mg CO}_2\text{-C g}^{-1} \text{ OC y}^{-1}$). In a comparison of Loess and Miocene Sand, both revealed large differences between samples with the lowest (1.2 and $1.8 \text{ mg CO}_2\text{-C g}^{-1} \text{ OC y}^{-1}$) and highest (34.2 and $12 \text{ mg CO}_2\text{-C g}^{-1} \text{ OC y}^{-1}$) respiration rate constants. Interestingly, there was no depth gradient for the different substrates, but

samples from 7 and 9 m depth of Miocene Sand tended to have up to 6.8 times lower respiration rate constants than samples from the 2 and 4 m depth. There was also a very wide variation between samples from the same depth (see Fig. A1) as revealed by the high standard deviations. Assuming a constant mineralisation rate over time, the results of the incubation experiment would result in mean residence times of between 29 and 135 years for the Loess, 83 and 556 years for the Miocene Sand and between 476 and 1,000 years for the crushed Red Sandstone. Since the CH₄ levels of the samples remained at a low level (Fig. A3) there were no indications of oxygen-limited conditions during the incubation.

Table 2: Comparison of mineralisation rate constants between the first (intact Red Sandstone samples) and second incubation experiments (crushed Red Sandstone samples) calculated using linear models.

Substrate	Depth (m)	Mineralisation rate constant (mg CO ₂ -C g ⁻¹ OC y ⁻¹) ^a
Miocene Sand	2	12.0
	4	10.5
	7	1.8
	9	2.9
Loess	1.7	7.4
	2.9	34.2
	5.4	10.2
	9.6	1.3
Uncrushed Red Sandstone	2.5	0.3
	5	1.0
	7.5	0.4
	9	0.4
Crushed Red Sandstone	2.5	1.3
	5	2.1
	7.5	2.0
	9	1.0

^a represents the slope of the fitted linear model for respiration during the incubation experiment




Fig. 4: Potential degradability of sedimentary OC from three sites. Results represent cumulative respiration from Loess and Miocene Sand samples and both Red Sandstone incubation experiments with uncrushed and crushed samples, with respective standard deviations ($n = 4$). Dashed lines represent a fitted linear model to the respiration data.

3.4 Possible contribution from geogenic organic carbon to soil organic carbon

As bedrock weathers, it becomes part of the soil and GOC also becomes soil OC. The potential amount of GOC in the subsoil was dependent on the sedimentary bedrock (Fig. 5). In the subsoil of Miocene Sand the proportion of GOC amounted to 2-3 %, at the Loess site it was 8 %, and at the Red Sandstone site it was 12 % (Table 1). For the defined topsoils (0-30 cm depth), contributions of GOC to soil OC were smaller with 0.7-0.9 % for Miocene Sand, 2.8-2.9 % for Loess and 2.8-3.0 % for Red Sandstone. The possible contribution of GOC to soil OC was also calculated under the assumption that the biogenic fraction had an unrealistically high average age of 10,000 years (Fig. A5). This resulted in the GOC fraction falling to 0 for Miocene Sand. For Loess the geogenic fraction fell by ~ 25 %, and for Red Sandstone by ~ 16 %.

The presence of ¹⁴C-free GOC in soil OC reduced the mean bulk soil OC ¹⁴C ages, depending on its proportion in soil OC content. Topsoils that developed in Loess and Red Sandstone had a modern ¹⁴C content of 1.029 and 1.035 F¹⁴C, similar to the atmospheric ¹⁴C content in 1950. Due to the large proportion of biogenic OC, no influence of a geogenic fraction was detected in the topsoil of these sites. No ¹⁴C data were available for the topsoil of Miocene Sand. For all subsoils, the influence of GOC on

bulk soil OC ^{14}C contents depended greatly on the depth and corresponding OC contents. For Loess, the possible influence on ^{14}C ages in the subsoils was quite high, with an average of 10 % reduction in mean apparent ^{14}C ages in the subsoil. Thus it would reduce the measured age of 4,413 years BP in 74 cm depth by 532-555 years BP. Geogenic OC potentially reduced the mean apparent radiocarbon age of 1,277 years BP at 0.39 m depth in Miocene Sand by about 7-9 years, and the radiocarbon age of 1,771 years BP at 0.61 m depth by 20-24 years. These reductions are below the respective standard deviations of the measurement. Nevertheless, at 1 m depth a given possible proportion of 13.1-16.3 % would reduce an interpolated ^{14}C age of 3,053 years BP by 399-497 years. For Red Sandstone the influence of GOC on ^{14}C ages would be highest in the subsoil. At 74 cm depth it would increase an age of 1,453 years BP by 451-490 years BP. Due to the low amounts of soil OC at 90 cm depth at the Red Sandstone site, the weight-based median amount of GOC in the sediments was four times higher than the biogenic amount of soil OC.




Fig. 5: Largest possible contribution of GOC to OC (red area) in relation to bulk OC content, taking into account the median GOC contents of the sediments for the respective horizontal weight based OC contents. The contribution of GOC is the mean amount based on the assumption of an average biogenic OC age of between 4,000 and 1,000 years

4. Discussion

4.1 Geogenic OC in the sediments

Generally, our calculations on the GOC fraction in the sediments are based on the assumption that biogenic OC in the sediments is not older than 4,000 yrs BP on average. And we also excluded the influence of a biogenic OC fraction that derives from soils that developed before the latest glacial period. Thus, there is uncertainty of a biogenic OC fraction in the sediments since it is unknown when biogenic OC entered the sediments. We assumed a mean age of 1000 to 4000 years based on DO^{14}C data that was leached from the soil. Nevertheless, even with an assumed age of 10,000 years for the biogenic OC fraction, the highest possible contribution was 15 % for the Loess (94 cm depth) and 22 % for the Red Sandstone (74 cm depth) (Fig. A5). A mean age of 10,000 years would be an unrealistic assumption since sediments are open systems and may receive OC input throughout the pedogenic period if vegetation is present and not only at the start of pedogenesis.

4.1.1 Site dependent contents of GOC

Regarding the calculated contribution of GOC to OC in the sediments, the assumed range of biogenic ^{14}C ages from 1,000-4,000 years BP was within the typical range for ages of dissolved OC leaching from soils (Artinger et al., 1996, Jia et al., 2019). Nevertheless, the range of 1,000-4,000 years BP did not greatly influence the range of calculated sedimentary contribution from GOC, especially for Loess (~ 6 % difference) and Red Sandstone (~ 9 % difference). The calculated GOC contribution for Miocene Sand was comparatively low (29-77 %) compared with the contribution for Red Sandstone and Loess, especially at 5 m depth. Compared with the Red Sandstone, this could be due to deep biogenic carbon inputs, such as roots and root exudates from the trees (Angst et al., 2016, John et al., 2016, Kirfel et al., 2017, Tückmantel et al., 2017), since the loosely bedded Miocene Sand allows for deep infiltration compared with the Red Sandstone site with its shallow bedrock as a root-restricting layer (Schneider and Don, 2019). Although it was not possible to define the exact rooting depth for Miocene Sand, this depth probably does not exceed 4 m, according to Schenk and Jackson (2005). Therefore, the increase in sedimentary contribution from 5 to 8 m depth of Miocene Sand could be due to the decreasing influence of roots and root exudates at 8 m depth.

In Loess, the low ^{14}C ages at 2-4 m depth (2,200-2,770 years BP) in contrast to a ^{14}C age of 4,413 years BP at 0.74 m depth were surprising. This might indicate past anthropogenic activities or erosion-driven material movement that might have led to a mixing of the upper part of the profile. Furthermore, the modern ^{14}C signature at 21 cm depth could be due to the plough layer at the Loess site mixing the upper 30 cm. Nevertheless, at depths below 4 m the high ^{14}C ages of OC in the sediments indicated a large proportion of GOC. This could be due to different sedimentation periods, soil forming and also soil burial processes (Chaopricha and Marin-Spiotta, 2014) that took place during the Pleistocene. These processes can lead to the presence of buried layers in Loess with varying amounts of rather recalcitrant OC, as shown by Marin-Spiotta et al. (2014). For the investigated Loess site, different sedimentation and soil forming processes can be expected due to the presence of completely different material in the cores in terms of colour and measured OC contents. For example, the very dark Loess at 4 m depth with

its high OC content support the assumption of sedimentation circumstances that favoured the accumulation and preservation of OC. This is in accordance with Jordan and Schwartau (1993) who investigated the same site and assigned the different layers to specific Pleistocene sedimentation periods.

In summary, the contribution of GOC to sedimentary OC was substrate dependent. A loosely bedded sediment like Miocene Sand with extremely low concentrations of OC could be more prone to infiltration of biogenic OC and dilution of GOC. This resulted in contributions of biogenic OC to the sediments of about 50 %. In contrast, the Loess site with comparatively low infiltration rates or the Red Sandstone site with reduced possibilities for deep rooting seemed to contain relative constant contributions from GOC of around 80 %.

The GOC contribution within the sediments was not found to increase with soil depth. This is in contrast to the results of Frouz et al. (2011) which showed that different sediment types from a Miocene clay sediment had higher weight-based carbon contents at 150 m compared with 30 m depth. However, a comparison with the present study is difficult since Frouz et al. (2011) did not distinguish between the geogenic and biogenic OC fractions, and OC contents were much higher (28-112 g kg⁻¹ dry mass compared to 0.008-10 g kg⁻¹ dry mass in our results). However, it also underlines the importance of different sedimentation processes for the amount and depth distribution of OC in sediments and sedimentary rocks. *4.1.2 Geogenic OC in Miocene Sand and Red Sandstone compared with other studies*

It is hard to compare the weight-based amounts of GOC in terrestrial sediments since most studies in this field rarely determine the amounts of OC in terrestrial sediments, or presume that sandy sediments, for example, do not contain large amounts of OC (Artinger et al., 1996). Quite high amounts of OC have been found in the skeleton part of different soils on sandstones with 0.61-1.97 t ha⁻¹ (Corti et al., 2002). Nevertheless, they mentioned the possible strong influence of organic substances from the soil solution without quantifying it, and did not directly investigate OC in the sediments. Additionally, Copard et al. (2007) assumed an OC amount of 2.4 g kg⁻¹ from an unknown source for all sandy sediments in a global storage modelling approach for the first metre of sediments. This would fit with the amounts of around 1 and 5 mg OC g⁻¹ found by Krummholz et al. (1997) for Dakota sandstone layers at > 180 m depth, but is much greater than the median GOC amount in Red Sandstone (0.2 g kg⁻¹) and Miocene Sand (0.1 g kg⁻¹) in the present study.

4.1.3 Geogenic OC in Loess compared with other studies

Loess deposits are relatively well investigated because they provide a record of paleoenvironmental conditions (Hatté et al., 1998, Head et al., 1989, Murton et al., 2015, Wang et al., 1996). The median amount of 0.27 g kg⁻¹ from the present study was low compared with studies by Hatté et al. (1998), Wang et al. (1996) and Strauss et al. (2012). Hatté et al. (1998) investigated 20-m deep loess deposits in the Rhine valley and found OC contents between 1.0-8.6 g kg⁻¹, Wang et al. (1996) investigated 12-m deep loess deposits in China and found OC contents of 31.2 ± 30.5 g kg⁻¹ and Strauss et al. (2012)

found OC contents of $15 \pm 14 \text{ g kg}^{-1}$ in Yedoma loess deposits in Siberia. This shows that the deposits from the site in the present study stored comparatively low GOC contents, although most of the above-mentioned studies do not distinguish between a biogenic and geogenic OC pool. Nevertheless, as also shown in the present study, Loess had a high OC content compared with other sediments. This was in line with highest OC contents in subsoils at the Loess site and may indicate the importance and contribution of bedrock OC to subsoil OC.

4.2 Is sedimentary derived organic carbon biodegradable?

The incubation experiment revealed a mineralisation of OC within the sediments with values between 0.1 and 3.4 % of total OC being mineralised after one year, assuming a constant mineralisation rate. The incubation temperature of 20 °C is above the typical mean temperatures in the subsoils underlining incubation derived mineralisation rates to be potential rates. For subsoils with comparable climatic conditions, Wordell-Dietrich et al. (2019) found temperatures at 150 cm depth ranging from 4 to 14.4°C over a two-year period. Assuming a Q_{10} value of around two for the assumable difference (Hamdi et al., 2013), the respiration rate at typical subsoil temperatures would be roughly half that. A direct mineralisation of OC from sediments is in agreement with several studies investigating the direct mineralisation from outcrops (Copard et al., 2007, Horan et al., 2017, Petsch et al., 2000, Soulet et al., 2017) Seifert et al., 2001). The difference to the present study is, that those studies observed this mineralisation when the sediments were directly exposed to the surface or/and were part of a very rapidly eroding area with GOC from the sediments getting inputs of recent vegetation. However, Frouz et al. (2011) conducted an incubation experiment with sedimentary samples from OC-rich Miocene clay sediments and found quite high respiration rate, with values between 0.4-1.2 % OC loss per year during a 91 day incubation experiment.. Kieft and Rosacker (1991) also found high respiration rates of sedimentary samples, with values between 0.1-1 % OC loss per year. Those results are in fairly good agreement with the respiration rate constants observed in the present study. Compared with subsoil incubation experiments, the mineralisation we found for sediments was quite low. For example, in a subsoil incubation experiments at 20 °C, Wordell-Dietrich et al. (2017) found that between 5-9.5 mg $\text{CO}_2\text{-C g}^{-1}$ OC of OC are mineralised after incubation for 63 days, Wang et al. (2013) report values between 5-15 mg $\text{CO}_2\text{-C g}^{-1}$ OC after 28 days, and Soucemarianadin et al. (2018) report values between 10- and 12.5 mg $\text{CO}_2\text{-C g}^{-1}$ OC after 70 days. The difference between the respiration rates observed in the present incubation experiment and the results from sedimentary and subsoil incubation experiments could be due to different microbial communities, OC quality and the physical disconnection between OC and potential decomposers.

Our study indicated that there was a considerable portion of biogenic OC in the sediments. The observed mineralisation could therefore be mainly due to the consumption of this biogenic OC part. Furthermore, the low mineralisation rate of Red Sandstone during the incubation as intact cores (Fig. A1) promoted the stability of GOC when it is part of the sediments.

The low mineralisation rates of the OC in the sediments might be caused by a lack of fresh substrates and/or microorganisms that could enhance the degradation of OC (Fontaine et al., 2007). Seifert et al. (2011) have shown that microorganisms are able to degrade sedimentary OC after the addition of glucose in black slate outcrops. Nevertheless, mineralisation of OC could only be observed with the addition of water, which indicates the widely recognised fact of the presence of an active microbial community in the sediments (Bomberg et al., 2017, Joergensen and Wichern, 2018, Magnabosco et al., 2018).

The incubation experiment was unable to answer the question of whether GOC is mineralised when it becomes part of the (sub-)soil. Assuming that a large part of the biogenic OC in the sediments was mineralised during the incubation experiment, geogenic OC could still be preserved during soil formation especially in the subsoils. This is in accordance with the indirect approach taken by Graz et al. (2010) to determine the mineralisation of sedimentary OC when it becomes part of the subsoil. They stated that 30 % of GOC resists degradation when it becomes part of the soil due to the results of a quantitative palynofacies analysis of bedrock and soil samples. Hemingway et al. (2018) found that sedimentary OC directly exposed to the surface in a rapidly eroding tropical mountain area exhibits considerable mineralisation down to 1 m below the surface, also leading to around 30 % of GOC remaining in the soil. Based on ^{14}C measurements they found out that on average 67 ± 11 % of the OC fraction in the sediments could be lost during soil formation, but no distinction was made between biogenic and geogenic OC fractions. This indicates that a microbial mineralisation of bedrock OC takes place but may be partly restricted to biogenic OC.

Regarding the depth distribution of GOC in the sediments, the amount of GOC (in g kg^{-1}) did not increase with depth, but there were clear differences. This reflects the sedimentation history with different initial amounts of OC and degradation during sedimentation, which is particularly evident in the high amounts of GOC at 5 m depth in Loess. However, the contents of GOC, especially in Red Sandstone and Miocene Sand, were within the same range for the whole depth. This might indicate that degradation of GOC is not depth-dependent within the sediments. If there were a stronger degradation of GOC with decreasing depth, a decreasing amount of GOC could be expected due to the input of water, microorganisms and fresh nutrients from above. Furthermore, there was a relatively constant contribution of biogenic OC within the sediments, meaning that if biogenic OC enters the sediments with possibly degrading microorganisms, this biogenic OC might also be largely mineralised. A study by Heitkötter et al. (2018) demonstrate effectively that degradation of OC in subsoils is primarily limited to small hotspot areas. Also in sediments, Krummholz et al. (1997) show that microbial communities can particularly be found in spatially discrete areas. Nevertheless, Heitkötter et al. (2018) also show that microorganisms outside the hotspots can be activated when substrate is supplied. Thus, the bioavailability of GOC might be site-dependent since root channels as microbial hotspots, for example, are less abundant and stable in sandy soils (Schneider and Don, 2019). With regard to the sites

investigated here, the solid Red Sandstone might only obtain water in preferential flow paths (Swanson, 2006) leading to a comparatively stable OC pool in the sediment. Meanwhile for weathered, poorly structured sandy soils, prevalent matrix flow conditions can be assumed (Flury et al., 1994). Thus, the bulk GOC might be supplied with fresh substrates and water from above more often than for the well-structured Loess soil with more frequent and stable preferential flow paths (Schneider and Don, 2019). This might lead to a lower accessibility and therefore slower turnover rates of OC (Dungait et al., 2012). The conclusion can be drawn that sediment derived OC can and will be degraded when it becomes part of the subsoil but probably to a comparatively limited extent and low rates with dominant mineralisation of biogenic OC.

4.3 How much does GOC contributes to soil organic carbon?

The contribution of GOC to soil OC stocks in this study was driven by the amount of OC in the soil, the amount of GOC in the respective sediment and also the turnover of GOC when it becomes part of the soil. Our results revealed that despite differing between sediments, GOC content varied in a quite narrow range between 0.1 and 0.3 g kg⁻¹. The contribution of GOC to topsoil OC was negligible. Assuming no degradation of GOC, the greatest possible contributions to total subsoil OC were found for Red Sandstone (~30 %) and the lowest for Miocene Sand (0.6 %). This was due to the range of OC contents in the subsoils (0.53 g kg⁻¹-15.21 g kg⁻¹). When soil OC contents were low, the possible contributions from GOC were high and vice versa. For the investigated soils, the OC contents of 3 g kg⁻¹ soil allowed for possible GOC contributions of between 5- and 10 %. For OC contents around 1 g kg⁻¹ soil, a GOC contribution of between 10- and 20 % seemed possible. Thus, greater contributions were made by GOC rich sediments such as Loess, while smaller contributions from sandy sediments. In comparison, van der Voort et al. (2018) estimated the contribution of GOC in a soil derived from glacial deposits (flysch) at between 80-100 cm depth to be around 40 %. For a soil developed from a poorly consolidated sedimentary rock (calcareous and shaly moraine), they calculate the contribution of GOC to range from 20 % at 145 cm depth to 80 % at 310 cm depth. An attempt has also been made to fractionate subsoils to extract the most stable OC that may be derived from GOC. Paul et al. (2001) investigated a soil developed on loess over till with 30 % of subsoil OC as a non-hydrolysable fraction showing a ¹⁴C age of 13,000 years BP. They also concluded that this high ¹⁴C age can partly be explained by a GOC fraction. These results indicate that deposits from the past glacial periods, such as flysch or till in particular, have much greater potential for OC contributions from GOC, possibly due to the higher amounts of GOC in their sediments. Since only terrestrial sediments were investigated in the present study, it should be noted that marine sediments or shales also contain much higher amounts of OC, up to 250 g kg⁻¹ (Hemingway et al., 2018, Petsch et al., 2000). Thus, the amount of GOC they contain and their possible contribution to subsoil OC stocks might therefore be much higher.

Nevertheless, the ¹⁴C ages of OC in the subsoil can be high in soils derived from igneous parent materials without GOC (Rumpel et al., 2002), although even crystalline bedrocks contain microbial communities

(Purkamo et al., 2020). Furthermore, on a global scale, the ^{14}C ages of soil OC are primarily driven by climatic conditions, clay content and age of the soil (Mathieu et al., 2015). However, for terrestrial sediments with comparatively low amounts of GOC that started their soil development after the latest glacial period, a scale of possible contributions could be obtained when the amount of OC is known. Thus, on a global scale, the high ^{14}C age of subsoils is not driven just by the GOC fraction but the presence of GOC may greatly influence subsoil ^{14}C .

5. Conclusions

This approach of estimating the GOC contribution to soil OC showed that common and abundant terrestrial sediments with low amounts of sedimentary OC can make a considerable contribution to subsoil OC stocks. One fraction of OC in the sediments is of geogenic origin and could therefore influence measured ^{14}C ages in soil, particularly in subsoils. Subsoils are known for their high ^{14}C ages and slow turnover rates and slow reaction to changing environmental condition. These subsoil OC properties may partly be derived from the GOC in the subsoil. The sediments at the investigated sites contained OC in a range of 0.1-0.3 g kg⁻¹, allowing for contributions from GOC of between 10-30 % in subsoils. Incubation of sediments indicated that this geogenic contribution presents a quite stable OC pool, especially for subsoils. Thus, even sediments with comparatively low amounts of OC were also able to demonstrate the large contribution of GOC.

Data availability

The data will be made available on request

Author contribution

AD conceived of and designed the study, FK performed the sampling and analysis, and wrote the first draft. All the authors contributed to generating and reviewing the subsequent versions of the manuscript.

Competing interest

The authors declare that they have no conflict of interest

Acknowledgements

This study was funded by the Deutsche Forschungsgemeinschaft (DFG) (DO1734/4-2) within the framework of the research unit SUBSOM (FOR1806)-“The Forgotten Part of Carbon Cycling: Organic Matter Storage and Turnover in Subsoils”. We would like to thank Frank Hegewald and the student assistants for their support in the field and in the laboratory, Jens Dyckmanns and Lars Szvec from the Centre for Stable Isotope Research and Analysis at the University of Göttingen for $^{13}\text{CO}_2$ measurements, Reiner Dohrmann from the Federal Institute for Geosciences and Natural Resources for density

610 measurements of sedimentary rocks and Roland Prietz for his excellent soil classification assistance.
611 We would also like to thank the AK laboratory team for their support.
612

References

- Amiotte Suchet, P.; Probst, J. L. and Ludwig, W.: Worldwide distribution of continental rock lithology: Implications for the atmospheric/soil CO₂ uptake by continental weathering and alkalinity river transport to the oceans, *Global Biogeochemical Cycles*, 17, 2003
- Angst, G.; John, S.; Mueller, C. W.; Kogel-Knabner, I. and Rethemeyer, J.: Tracing the sources and spatial distribution of organic carbon in subsoils using a multi-biomarker approach, *Scientific Reports*, 6, 29478, 2016
- Artinger, R.; Buckau, G.; Kim, J.; Geyer, S. and Wolf, M.: Influence of sedimentary organic matter on dissolved fulvic acids in groundwater. Significance for groundwater dating with ¹⁴C in dissolved organic matter, 1996
- Batjes, N. H.: Total carbon and nitrogen in the soils of the world, *Eur J Soil Sci*, 65, 10-21, 2014
- Bertrand, I.; Delfosse, O. and Mary, B.: Carbon and nitrogen mineralization in acidic, limed and calcareous agricultural soils: Apparent and actual effects, *Soil Biology and Biochemistry*, 39, 276-288, 2007
- Bomberg, M.; Raulio, M.; Jylha, S.; Mueller, C. W.; Hoschen, C.; Rajala, P.; Purkamo, L.; Kietavainen, R.; Ahonen, L. and Itavaara, M.: CO₂ and carbonate as substrate for the activation of the microbial community in 180 m deep bedrock fracture fluid of Outokumpu Deep Drill Hole, Finland, *AIMS Microbiol*, 3, 846-871, 2017
- Cerri, C.; Feller, C.; Balesdent, J.; Victoria, R. and Plenecassagne, A.: PARTICLE-SIZE FRACTIONATION AND STABLE CARBON ISOTOPE DISTRIBUTION APPLIED TO THE STUDY OF SOIL ORGANIC-MATTER DYNAMICS, *Comptes Rendus De L Academie Des Sciences Serie Ii*, 300, 423-+, 1985
- Chaopricha, N. T. and Marin-Spiotta, E.: Soil burial contributes to deep soil organic carbon storage, *Soil Biol Biochem*, 69, 251-264, 2014
- Colman, B. P. and Schimel, J. P.: Drivers of microbial respiration and net N mineralization at the continental scale, *Soil Biology Biochemistry*, 60, 65-76, 2013
- Copard, Y.; Amiotte-Suchet, P. and Di-Giovanni, C.: Storage and release of fossil organic carbon related to weathering of sedimentary rocks, *Earth Planet Sc Lett*, 258, 345-357, 2007
- Corti, G.; Ugolini, F. C.; Agnelli, A.; Certini, G.; Cuniglio, R.; Berna, F. and Sanjurjo, M. J. F.: The soil skeleton, a forgotten pool of carbon and nitrogen in soil, *Eur J Soil Sci*, 53, 283-298, 2002
- Crow, S. E.; Lajtha, K.; Filley, T. R.; Swanston, C. W.; Bowden, R. D. and Caldwell, B. A.: Sources of plant-derived carbon and stability of organic matter in soil: implications for global change, *Global Change Biology*, 15, 2003-2019, 2009
- Eckelmann, W.; Sponagel, H.; Grottenthaler, W.; Hartmann, K.-J.; Hartwich, R.; Janetzko, P.; Joisten, H.; Kühn, D.; Sabel, K.-J. and Traidl, R.: *Bodenkundliche Kartieranleitung*. KA5, 2006
- Fontaine, S.; Barot, S.; Barre, P.; Bdioui, N.; Mary, B. and Rumpel, C.: Stability of organic carbon in deep soil layers controlled by fresh carbon supply, *Nature*, 450, 277-280, 2007
- Frouz, J.; Cajthaml, T.; Kribek, B.; Schaeffer, P.; Bartuska, M.; Galertova, R.; Rojik, P. and Kristufek, V.: Deep, subsurface microflora after excavation respiration and biomass and its potential role in degradation of fossil organic matter, *Folia Microbiologica*, 56, 389-396, 2011
- Graz, Y.; Di-Giovanni, C.; Copard, Y.; Laggoun-Defarge, F.; Boussafir, M.; Lallier-Verges, E.; Baillif, P.; Perdereau, L. and Simonneau, A.: Quantitative palynofacies analysis as a new tool to study transfers of fossil organic matter in recent terrestrial environments, *Int J Coal Geol*, 84, 49-62, 2010
- Hatté, C.; Fontugne, M.; Rousseau, D.-D.; Antoine, P.; Zöller, L.; Laborde, N. T. r. and Bentaleb, I.: $\delta^{13}\text{C}$ variations of loess organic matter as a record of the vegetation response to climatic changes during the Weichselian, *Geology*, 26, 583-586, 1998
- Head, M.; Zhou, W. and Zhou, M.: Evaluation of ¹⁴C ages of organic fractions of paleosols from loess-paleosol sequences near Xian, China, *Radiocarbon*, 31, 680-695, 1989
- Hemingway, J. D.; Hilton, R. G.; Hovius, N.; Eglinton, T. I.; Haghipour, N.; Wacker, L.; Chen, M.-C. and Galy, V. V.: Microbial oxidation of lithospheric organic carbon in rapidly eroding tropical mountain soils, *Science*, 360, 209-212, 2018

665 Horan, K.; Hilton, R. G.; Selby, D.; Ottley, C. J.; Gröcke, D. R.; Hicks, M. and Burton, K. W.:
 666 Mountain glaciation drives rapid oxidation of rock-bound organic carbon, *Science advances*, 3,
 667 e1701107, 2017
 668 Jia, J.; Feng, X.; Pannatier, E. G.; Wacker, L.; McIntyre, C.; van der Voort, T.; Montlucon, D. and
 669 Eglinton, T.: ¹⁴C characteristics of dissolved lignin along a forest soil profile, *Soil Biology*
 670 *Biochemistry*, 2019
 671 Joergensen, R. G. and Wichern, F.: Alive and kicking: Why dormant soil microorganisms matter, *Soil*
 672 *Biology and Biochemistry*, 116, 419-430, 2018
 673 John, S.; Angst, G.; Kirfel, K.; Preusser, S.; Mueller, C. W.; Leuschner, C.; Kandeler, E. and
 674 Rethemeyer, J.: Which are important soil parameters influencing the spatial heterogeneity of ¹⁴C in
 675 soil organic matter, *Biogeosciences Discuss*, 123, 2016
 676 Jordan, H. and Schwartau, W.: Das Lößprofil von Ahlshausen und weitere tiefe Quartäraufschlüsse
 677 entlang der Bundesbahn-Neubaustrecke bei Northeim, Südniedersachsen, *Quaternary Science Journal*,
 678 43, 12, 1993
 679 Kieft, T. L. and Rosacker, L. L.: Application of respiration-and adenylate-based soil microbiological
 680 assays to deep subsurface terrestrial sediments, *Soil biology Biochemistry*, 23, 563-568, 1991
 681 Kirfel, K.; Leuschner, C.; Hertel, D. and Schuldt, B.: Influence of Root Diameter and Soil Depth on
 682 the Xylem Anatomy of Fineto Medium-Sized Roots of Mature Beech Trees in the Top- and Subsoil,
 683 *Front Plant Sci*, 8, 2017
 684 Kögel-Knabner, I.; Guggenberger, G.; Kleber, M.; Kandeler, E.; Kalbitz, K.; Scheu, S.; Eusterhues, K.
 685 and Leinweber, P.: Organo-mineral associations in temperate soils: Integrating biology, mineralogy,
 686 and organic matter chemistry, *Journal of Plant Nutrition and Soil Science*, 171, 61-82, 2008
 687 Libby, W. F.: Radiocarbon Dating, *The Society for American Archaeology*, 132, 1952
 688 Litt, T.; Behre, K.-E.; Meyer, K.-D.; Stephan, H.-J. and Wansa, S.: Stratigraphische Begriffe für das
 689 Quartär des norddeutschen Vereisungsgebietes, *Quaternary Science Journal*, 56, 7-65, 2007
 690 Magnabosco, C.; Lin, L.-H.; Dong, H.; Bomberg, M.; Ghiorse, W.; Stan-Lotter, H.; Pedersen, K.;
 691 Kieft, T.; Van Heerden, E. and Onstott, T. C.: The biomass and biodiversity of the continental
 692 subsurface, *Nature Geoscience*, 11, 707-717, 2018
 693 Marin-Spiotta, E.; Chaopricha, N. T.; Plante, A. F.; Diefendorf, A. F.; Mueller, C. W.; Grandy, A. S.
 694 and Mason, J. A.: Long-term stabilization of deep soil carbon by fire and burial during early Holocene
 695 climate change, *Nature Geoscience*, 7, 428-432, 2014
 696 Mathieu, J. A.; Hatté, C.; Balesdent, J. and Parent, É.: Deep soil carbon dynamics are driven more by
 697 soil type than by climate: a worldwide meta-analysis of radiocarbon profiles, *Global change biology*,
 698 21, 4278-4292, 2015
 699 Murton, J. B.; Goslar, T.; Edwards, M. E.; Bateman, M. D.; Danilov, P. P.; Savvinov, G. N.; Gubin, S.
 700 V.; Ghaleb, B.; Haile, J. and Kanevskiy, M.: Palaeoenvironmental Interpretation of Yedoma Silt (Ice
 701 Complex) Deposition as Cold-Climate Loess, Duvanny Yar, Northeast Siberia, *Permafrost Periglacial*
 702 *Processes*, 26, 208-288, 2015
 703 Nelson, D.W., Sommers, L.: *Methods of soil analysis: Part 2 chemical and microbiological properties*,
 704 *Wiley Online Library*, 9, 539-579, 1983
 705 Paul, E.; Collins, H. and Leavitt, S.: Dynamics of resistant soil carbon of Midwestern agricultural soils
 706 measured by naturally occurring ¹⁴C abundance, *Geoderma*, 104, 239-256, 2001
 707 Petsch, S.; Berner, R. and Eglinton, T.: A field study of the chemical weathering of ancient
 708 sedimentary organic matter, *Org Geochem*, 31, 475-487, 2000
 709 R Core Team: *R: A language and environment for statistical computing*, 2018
 710 Rethemeyer, J.; Gierga, M.; Heinze, S.; Stolz, A.; Wotte, A.; Wischhöfer, P.; Berg, S.; Melchert, J.
 711 and Dewald, A.: Current sample preparation and analytical capabilities of the radiocarbon laboratory
 712 at CologneAMS, *Radiocarbon*, 61, 1449-1460, 2019
 713 Rumpel, C. and Kögel-Knabner, I.: The role of lignite in the carbon cycle of lignite-containing mine
 714 soils: evidence from carbon mineralisation and humic acid extractions, *Org Geochem*, 33, 393-399,
 715 2002
 716 Rumpel, C.; Kögel-Knabner, I. and Bruhn, F.: Vertical distribution, age, and chemical composition of
 717 organic carbon in two forest soils of different pedogenesis, *Org Geochem*, 33, 1131-1142, 2002
 718 Schiff, S.; Aravena, R.; Trumbore, S. E.; Hinton, M.; Elgood, R. and Dillon, P.: Export of DOC from
 719 forested catchments on the Precambrian Shield of Central Ontario: clues from ¹³C and ¹⁴C,
 720 *Biogeochemistry*, 36, 43-65, 1997

721 Schneider, F. and Don, A.: Root-restricting layers in German agricultural soils. Part I: extent and
 722 cause, *Plant Soil*, 442, 433-451, 2019
 723 Schrumpf, M.; Kaiser, K.; Guggenberger, G.; Persson, T.; Kögel-Knabner, I. and Schulze, E.-D.:
 724 Storage and stability of organic carbon in soils as related to depth, occlusion within aggregates, and
 725 attachment to minerals, *Biogeosciences*, 10, 1675-1691, 2013
 726 Soucemarianadin, L. N.; Cecillon, L.; Guenet, B.; Chenu, C.; Baudin, F.; Nicolas, M.; Girardin, C. and
 727 Barre, P.: Environmental factors controlling soil organic carbon stability in French forest soils, *Plant*
 728 *Soil*, 426, 267-286, 2018
 729 Soulet, G.; Hilton, R. G.; Garnett, M. H.; Dellinger, M.; Croissant, T.; Ogrič, M. and Klotz, S.: in situ
 730 measurement of flux and isotopic composition of CO₂ released during oxidative weathering of
 731 sedimentary rocks, *Biogeosciences*, 2017
 732 Strauss, J.; Schirrmeister, L.; Wetterich, S.; Borchers, A. and Davydov, S.: Grain-size properties and
 733 organic-carbon stock of Yedoma Ice Complex permafrost from the Kolyma lowland, northeastern
 734 Siberia, *Global biogeochemical cycles*, 26, 2012
 735 Stuiver, M. and Polach, H. A.: Reporting of C-14 data-Discussion, *Radiocarbon*, 19, 355-363, 1977
 736 Swanson, S. K.; Bahr, J. M.; Bradbury, K. R. and Anderson, K. M. J. S. G.: Evidence for preferential
 737 flow through sandstone aquifers in Southern Wisconsin, 184, 331-342, 2006
 738 Torn, M.; Swanston, C.; Castanha, C. and Trumbore, S., Wiley, Hoboken, 2009
 739 Trumbore, S.: Radiocarbon and soil carbon dynamics, *Annual Review of Earth Planetary Sciences*, 37,
 740 47-66, 2009
 741 Tückmantel, T.; Leuschner, C.; Preusser, S.; Kandeler, E.; Angst, G.; Mueller, C. W. and Meier, I. C.:
 742 Root exudation patterns in a beech forest: dependence on soil depth, root morphology, and
 743 environment, *Soil Biology Biochemistry*, 107, 188-197, 2017
 744 van der Voort, T.; Mannu, U.; Hagedorn, F.; McIntyre, C.; Walthert, L.; Schleppi, P.; Haghypour, N.
 745 and Eglinton, T.: Dynamics of deep soil carbon - insights from ¹⁴C time series across a climatic
 746 gradient, *Biogeosciences*, 2018
 747 Vindušková, O.; Sebağ, D.; Cailleau, G.; Brus, J. and Frouz, J.: Methodological comparison for
 748 quantitative analysis of fossil and recently derived carbon in mine soils with high content of aliphatic
 749 kerogen, *Org Geochem*, 89, 14-22, 2015
 750 Wang, X.; Cammeraat, L. H.; Wang, Z.; Zhou, J.; Govers, G. and Kalbitz, K.: Stability of organic
 751 matter in soils of the Belgian Loess Belt upon erosion and deposition, *Eur J Soil Sci*, 64, 219-228,
 752 2013
 753 Wang, Y.; Amundson, R. and Trumbore, S.: Radiocarbon dating of soil organic matter, *Quaternary*
 754 *Research*, 45, 282-288, 1996
 755 Waschkes, C. and Huttel, R. F.: Microbial degradation of geogenic organic C and N in mine spoils,
 756 *Plant Soil*, 213, 221-230, 1999
 757 Wordell-Dietrich, P.; Don, A. and Helfrich, M.: Controlling factors for the stability of subsoil carbon
 758 in a Dystric Cambisol, *Geoderma*, 304, 40-48, 2017
 759 Wordell-Dietrich, P.; Don, A.; Wotte, A.; Rethemeyer, J.; Bachmann, J.; Helfrich, M.; Kirfel, K. and
 760 Leuschner, C.: Vertical partitioning of CO₂ production in a Dystric Cambisol, *Biogeosciences*
 761 *Discussion*, 1-27, 2019
 762 WRB, I. W. G., Food and Agriculture Organization of the United Nations, Rome, 2006
 763 Zethof, J. H. T.; Leue, M.; Vogel, C.; Stoner, S. W. and Kalbitz, K.: Identifying and quantifying
 764 geogenic organic carbon in soils – the case of graphite, *SOIL*, 5, 383-398, 2019