

What do we know about how the terrestrial multicellular soil fauna reacts to microplastic?

Frederick Büks¹, Nicolette Loes van Schaik², Martin Kaupenjohann¹

¹Chair of Soil Science, Dept. of Ecology, Technische Universität Berlin, 10587 Berlin, Germany

²Chair of Ecohydrology & Landscape Evaluation, Dept. of Ecology, Technische Universität Berlin, 10587 Berlin, Germany

Correspondence to: Frederick Büks (frederick.bueks@tu-berlin.de)

Final responses to all referees plus marked-up version

Dear Referee #1

First I would like to express my sincere thanks to you for thoroughly reviewing our manuscript and for your very helpful and precise suggestions. In the following I will answer your points. Our corrections are marked-up with **yellow numbers** within the corrected manuscript at the end of this document.

Best regards,
Frederick Büks

Abstract

[1] Lines 20-21: "Most of the studies applied MP concentrations similar to amounts in slightly to very heavily polluted soils." This sentence makes the reader expect that generally, the concentrations in the experimental environments are mostly the same as expected in the environment, but is this really the case? I would suggest showing the percentage of experiments with high microplastic exposure that is not representative of most soils.

Thanks a lot for this point. We now write: "About 58 % of the studies thereby use inappropriate concentrations or units, but 42 % applied MP concentrations similar to amounts in slightly to very heavily polluted soils."

Introduction

[2] Line 53: Instead of "microbial decay", I'd suggest "processing by soil organisms", since this includes any process relevant for the generation of smaller plastic particles.

Done.

[3] Line 61: I'd suggest changing the sentence to "intensive use of plastic mulching and sewage sludge", for the former, Huang et al. (2020) show an increase in microplastic by approx. 1 order of magnitude between fields with 5 and 24 continuous years of plastic mulching.

Done.

[4] Line 95: Suggest changing "feed on" to "inadvertently ingest", otherwise it sounds like the organisms are actually able to metabolize the microplastics.

Done and reference added.

Search pattern

[5] The cut-off dates (time period that was considered) of the search should be mentioned somewhere.

Information added to this chapter (see the answer to referee #3)

[6] Figure 1: This figure shows the phylogenetic tree of edaphic fauna, rather than "edaphic tree of faunal life".

Thank you. And done.

Data collection

[7] Line 113-122: I've been having some difficulties understanding the search methodology and table 8 (table 8 should be moved at the appropriate place to become table 1).

We moved the table to line 124 and mentioned that it contains the number of found studies. All table numbers were adjusted within the text.

It would be great if the authors could re-word this, specifying:

What does it mean that some combinations would have caused too much search effort?

It means e.g. that searching for a taxon only in combination with "PET" gives results for PET bottles for cultivation and experiments and also the "use" as pets, if the search is not case sensitive. We now tried to clarify this in our text.

“Organism-plastic” is not a type-shape combination.
Oh, yes, that's right. Corrected.

What exactly does the number of studies in table 8 mean? The number of articles or single experiments (sometimes more than one taxon or plastic type is used in one article)?

The number counts for how often type-shape combinations were used in all reviewed experimental setups independently of organism.

Some articles are included that studied the uptake of macroplastics by organisms, mainly termites and ant species. It is reasonable to include these studies, but it should be mentioned more prominently, in the abstract and aims of the review, that macroplastics are included.

Where macroplastics were used in the reviewed studies, the size was explicitly mentioned in the article text, so we do not see a necessity for elaborating the text. We did add a mention of macroplastics to the abstract.

Maybe also in the synthesis, a sentence about the proportions of experiments using macro-, micro-, and nanoplastic would be a helpful piece of information.

Now mentioned in “4.2 Limitations of previous studies”

[8] Tables 1-7: What does N/A mean in the tables? In some cases I assume “not analysed” (e.g., passive transport), but in other cases it should mean “not mentioned” (e.g., aging, coating, etc.) or “not observed” (e.g., measured adverse effects). I think this needs to be specified. Usually, N/A refers to “not applicable”, but this doesn't fit in the tables.

In this work it means “(data) not available”. We marked it at the tables.

Synthesis

Lines 549-550: Could you cite the studies that imitated weathering in the described way?

We did so. Tsunoda et al. (2010) artificially aged their plastic by soaking in hot water at 90°C for 21 days, and then it was sanded/scratched with medium-grade paper prior to the test. Gebhard and Forster (2018) incubated particles in seawater for 4 weeks to stimulate the formation of biofilms.

[9] Lines 555-557: This is true, but it should be acknowledged that these additives are mainly present in commercial plastics, and therefore, mentioning of additives is not expected for “clean” microbeads specifically synthesized for the experiments. Nevertheless, the disadvantages of using these microbeads has been clearly discussed earlier in this section.

Done.

Conclusions

[10] Line 620-621: I am a little concerned about describing the results as “alarming”. Is it really? The following sentences actually refute this rather strong statement.

Replaced with “considerable”.

[11] Lines 624-629: I would suggest changing the sentence to: “To elucidate [...], the most exact reproduction of plastic concentrations and properties [. . .]”. However, the difficulty here is that very scarce data of limited quality is available on concentrations of microplastic in soils, so a range of concentrations need to be used for future experiments in order to match the “real world” concentrations in soil, while expecting a decrease in uncertainty in analytic results in the future. Especially in the lower size ranges (<100µm) quantification is currently challenging. Therefore, little is known about size distributions occurring in soils. It might be worth mentioning this dilemma in a sentence.

Done.

[12] Technical corrections:

All done.

Dear PD Dr. Werner Kratz (referee #2)

Thank you very much for your review. In the following I will try to answer your comments at my best. Our corrections are marked-up with green numbers within the corrected manuscript at the end of this document.

*Best regards,
Frederick Büks*

[1] Line 53: Is that only "microbial" decay?

We agree, we will change this to "processing by soil organisms" as it is actually micro- as well as macroorganisms.

[2] Figure 1: The taxonomic group "further Panarthropods" is placed centrally, the other groups are not.
Done.

[3] Table 7: The last three experiments within this table were conducted by feeding the mice with a MP suspension. You might write "(food)" behind the concentration data as in the other tables.
Thanks a lot. Done.

[4] Line 507: "Preferably" instead of "preferrably".
Done.

[5] Table 8: Could you explain the meaning of the numbers within the table. Are these the numbers of experiments with the named type-shape combinations?
Yes. Please see the answer to referee #1 (Table 8 is now Table 1).

[6] Lines 549-507: Is that proved that carboxylation of microspheres decreases hydrophobicity in an appreciable extent?
We ask the manufacturers of Polysciences Europe GmbH, a leading producer of PS microspheres, and they said no. We added this important information to the review.

Dear Referee #3

Thank you very much for your critical review of our manuscript. It has helped us to see some points which still need clarification. In the following, we want to explain how we propose to adjust our article based on the reviewer's comments and also explain why in some cases we do not agree with the reviewer's proposed changes. Our corrections are marked-up with **purple numbers** within the corrected manuscript at the end of this document.

(1) First and foremost, please have the manuscript edited by a professional (!) native (!)biologist (!). The English of your text is largely understandable, but rough. Apart from annoying typos, I found sentences the meaning of which I only understood when trying to translate them to German (my native language). So, your text will heavily benefit from thorough native editing.

Rereading our article we did indeed see that some typos had escaped our notice. We are slightly surprised by the request of the reviewer to have the manuscript edited by a "professional (!) native (!) biologist (!)". We **rephrased some stiff sentences and corrected grammatical errors**. If a proofreading is indeed wished, we will have a scientific translator (English native speaker) correct the article.

(2) Then, the text lacks conciseness, it is overly long. For example, I suggest to omit all biological/ecological details you provide when introducing a taxon. This is per se interesting, but not to the point here (except when the reader needs background to understand microplastic effects). Then, figure 1 does not contribute to the understanding of your presentation, omit it. And I do not think it necessary to present taxa for which there is no information available, especially if the taxa are of minor or no importance in soil (e.g. line 183ff, 205, 220, 227, 450ff) or if the literature is not on edaphic species (435ff). As a reviewer, you are of course required to address blind spots of research (thus pointing out important taxa that are missing in literature), but you need to better balance completeness with a concise presentation.

Your suggestion to omit the ecological presentation of some key taxa is understandable. If we would expect all readers to be well acquainted with the soil fauna, we would definitely go along with this. However, SOIL is a multi-disciplinary journal connecting a broad spectrum of soil scientists. Therefore, we think it is helpful to provide a short overview of information on the soil fauna, such as ecological functionalities (marker function, transport, degradation, habitat and food selection), which might influence how they cope with microplastics. We have critically gone through the article and here we summarize which parts we will shorten.

- **[1] Proposal:** We shortened the introduction of the springtail section, as it is indeed oversized.

For the same reason we illustrated the phylogenetic tree of soil life.

- **[2] Proposal:** We would agree with moving it to the supplements in order to save space, in case this is wished.

We also do not fully agree with your suggestion to delete taxonomic groups that have not yet been subject of studies on microplastics. The reason is, that the aim of this work is not only to review effects on studied taxa, but also to show gaps of knowledge especially apart from the common model organisms. In fact, their importance for the current ecological research should be shortly mentioned.

- **[2] Proposal:** Unstudied taxa are still presented, but their importance for future research is now additionally mentioned in section 4.3 to better "balance completeness".
- **[3] Proposal:** We shortened the chapter about Onychophora.

Potamopyrgus antipodarum in fact is a benthic snail.

- **[4] Proposal:** We use this benthic species to show more clearly how inconsistent the few results for benthic and terrestrial snails are.

(3) I miss a convincing argumentation why you focus on multicellular animals (but then, you provide many details about bacteria, fungi, algae, plant roots in 72ff...omit this). A good line of reasoning could be that you follow up on the Rillig and Bonkowski (2018) paper.

The aim of this review is to depict the influence of microplastic contamination in soils to the soil fauna. But, to present a holistic view on the food web, we refer to microorganisms, plant roots and biofilms within the introduction section. Being large fields of knowledge on their own, these organisms are not part of the focus in this review, however they are food sources for meso- and macroorganisms and, thus, worthy of mention. Given that we only use 22 lines to describe these other parts of the phylogenetic tree of soil life, we think this is merited and wish to leave this part in the review.

Unfortunately, we do not understand how Rillig and Bonkowski (2018), a paper on soil protozoa, matches your point. We have read this paper and do mention it elsewhere in the review.

(4) Please provide details of your literature search (123ff). When did you search? Which time span did you cover? Which search strings? Please consider the literature on meta-analyses how to properly specify these technical aspects.

The search was applied between June 2019 and January 2020, repeated in the first week of January 2020 and covers publications until January 2020. The search strings result from combinations of taxon, plastic type and particle shape shown in Table 1 (formerly Table 8).

- [5] Proposal: Information added to section 2.

(5) Thank you very much for the positive note.

(6) line 636f: Please reconsider including your supervisor as a co-author. What "supervision" means is nowhere clearly defined, however, co-authorship is only justified for significant contributions to the manuscript. Honorary authorship violates the principle of scientific honesty.

We understand this point completely and agree that it is not good practice to include scientists who have not contributed significantly to a paper. We also acknowledge that supervision is a very broad term and would like to specify the contribution of Martin Kaupenjohann to the paper. [6] Martin Kaupenjohann was involved in the development of the idea and concept for this paper. During the literature reading and writing phase he has supported the work with frequent discussions of the contents of the article. And finally he has critically revised the manuscript.

Best regards,

Dr. Frederick Büks
Dr. Loes van Schaik
Prof. Dr. Martin Kaupenjohann

Dear Dr. Maha Deeb (referee #4)

Thank you very much for the repeated check and your friendly report.

Best regards,

Dr. Frederick Büks

What do we know about how the terrestrial multicellular soil fauna reacts to microplastic?

Frederick Büks¹, Nicolette Loes van Schaik², Martin Kaupenjohann¹

¹Chair of Soil Science, Dept. of Ecology, Technische Universität Berlin, 10587 Berlin, Germany

²Chair of Ecohydrology & Landscape Evaluation, Dept. of Ecology, Technische Universität Berlin, 10587 Berlin, Germany

Correspondence to: Frederick Büks (frederick.bueks@tu-berlin.de)

Abstract. The ubiquitous accumulation of microplastic (MP) particles across all global ecosystems comes along with the uptake into soil food webs. In this review, we analyzed studies on passive translocation, active ingestion, bioaccumulation and adverse effects within the phylogenetic tree of multicellular soil faunal life. The representativity of these studies for natural soil ecosystems was assessed using data on the type of plastic, shape, composition, concentration and time of exposure.

Available studies cover a wide range of soil organisms, with emphasis on earthworms, nematodes, springtails, beetles and lugworms, each focused on well-known model organisms. [1] About 58 % of the studies thereby used inappropriate concentrations or units, but 42 % applied MP concentrations similar to amounts in slightly to very heavily polluted soils. In many cases, however, polystyrene microspheres have been used, a combination of plastic type and shape, that is easily available, but does not represent the main plastic input into soil ecosystems. In turn, MP fibers are strongly underrepresented compared to their high abundance within contaminated soils. [7] A few studies also examined the comminution of macroplastic by the soil fauna. Further properties of plastic such as aging, coating and additives were insufficiently documented. Despite these limitations, there is a recurring pattern of active intake followed by a population shift within the gut microbiome and adverse effects on motility, growth, metabolism, reproduction and mortality in various combinations, especially at high concentrations and small particle sizes.

For the improvement of future studies, we identified problems of past experiments and recommend that coming studies take into account the type, shape, grade of aging, specific concentrations of MP fractions and long-term incubation, in natural and contaminated soils.

34 1 Introduction

35 Imagine a compact plastic cube of nearly 2 km side length and a weight of 7300000000 tons,
36 with major percentages by weight of 36 % polyethylene (PE), 21 % polypropylene (PP), 12 %
37 polyvinyl chloride (PVC) and 10 % of each polyethylene terephthalate (PET), polyurethane
38 (PU) and polystyrene (PS). That is the accumulated global non-fiber production of the six
39 main plastic types until 2015. It accounts to 87 % of the all-time plastic production, which
40 evolved exponentially, since the early 1950s, from some megatons (Mt) to 8300 Mt in 2015,
41 with only 260 Mt annual output in 2009 increased to 380 Mt in 2015 (Thompson et al., 2009;
42 Geyer et al., 2017). Of this ever produced plastic, 6300 Mt became waste until 2015, of which
43 only 21 % were recycled or incinerated, whereas 5000 Mt ended up in landfills and nature
44 (Geyer et al., 2017). As a corollary of production, use and disposal, a certain part of plastic
45 waste is constantly released into the environment through various pathways, but our
46 knowledge about rates of mass flow into global ecosystems is very limited. Based on waste
47 generation in coastal countries, Jambeck et al. (2015) calculated the global plastic input to
48 marine ecosystems to be roughly 4.8 to 12.7 Mt in 2010. Such data on soils are lacking, but
49 Nizzetto et al. (2016) estimated that the load of microplastic (MP) to agricultural sites in
50 Europe is in the same order of magnitude as that in marine environments.

51 By littering, plastic mulching, the application of sewage sludge, digestates and composts as
52 well as windblown dispersal (Bertling et al., 2018; Weithmann et al., 2018; Zhang et al., 2019;
53 Wang et al., 2019a), plastic from our technosphere arrives in soil ecosystems in various forms
54 as large and small fragments, fibers and particles. Exposed to UV radiation, mechanical
55 stress and ~~211~~ processing by soil organisms, plastic items become weathered and prone to a
56 successive comminution towards the size range of MP with increased surface, charge and
57 biofilm cover (Kale et al., 2015; Andrady, 2017). However, the resistance of plastic to
58 metabolization causes a constant accumulation in soils as long as the release rate from
59 human processes is above the very slow rate of degradation.

60 Due to a lack of monitoring programs, data on MP concentrations in terrestrial soils are rare,
61 and those using w/w concentrations represent only a small part compared to item
62 concentrations. In soils with only slightly contaminated conditions, amounts seem to average
63 about 1 mg kg⁻¹ soil dry weight (and approx. 200 items kg⁻¹ dry soil) (Rezaei et al., 2019). On
64 sites with industrial activity or ~~13~~ intensive use of plastic mulching and sewage sludge in
65 agriculture, concentrations can be increased by 2 to 4 orders of magnitude (Fuller and
66 Gautam, 2016; Zhang and Liu, 2018; Huang et al., 2020). Semisubhydic soils such as
67 beaches, mudflats, mangroves or lagoons, that are additionally contaminated from the aquatic
68 side, contain MP of the order of 10 to 100 items kg⁻¹ dry soil and single extreme samplings
69 contained several thousand items (Nor and Obbard, 2014; Naji et al., 2017; Garcés-Ordóñez
70 et al., 2019; Li et al., 2018a). More informative data using mg kg⁻¹ are only available for
71 beaches and coastal deconstruction yards in municipal neighborhoods and amount to 0.5 and

72 70 mg kg⁻¹ dry soil, 0.00005 and 0.007 % w/w, respectively (Reddy et al., 2006; Claessens et
73 al., 2011). All these concentration data represent a wide range of particle sizes between 0 and
74 5000 µm with different materials, shapes and degrees of aging.

75 Plastic particles can possibly enter and accumulate in the food web by either direct uptake
76 from soil or by consumption of other soil biota contaminated by adhesion or ingestion (Huerta
77 Lwanga et al., 2017a). There is evidence, that MP is incorporated even by plants and
78 unicellular organisms at the base of the food web. **Bacteria**, for example, that are reasonably
79 assumed to avoid MP uptake due to their minor size and the prevalent lack of phagocytosis,
80 were shown to take up inorganic nanoparticles of a few nanometers (Kumar et al., 2011).
81 Although the physiochemical properties of weathered nanoparticulate plastics might differ
82 from these, also their uptake seems [12] likely.

83 A similar argument can be made for **fungi** and soil **algae**, but studies on incorporation are
84 lacking, whereas the transfer into a freshwater food web by adhesion of nanoplastic on algae
85 has been shown by Chae et al. (2018). The uptake of MP into **plant roots** is also inhibited
86 (Rillig et al., 2019), but occurred for nanoplastics that permeate into the plant tissue (Li et al.,
87 2019). Also the integration into root tissue after adsorption to the rhizodermis [12] has yet to be
88 studied.

89 In contrast, **protozoa** feature phagocytosis for the active ingestion of particles. Diverse soil,
90 freshwater and marine ciliates ingest PS/latex beads of 0.1 to 14.4 µm in laboratory
91 experiments, with preferences to their natural prey size (Fenchel, 1980; Jonsson, 1986; Lavin
92 et al., 1990). Soil amoebas act similarly, but additionally select according to food quality
93 (Weisman and Korn, 1967; Vogel et al., 1980; Bowers and Olszewski, 1983; Avery et al.,
94 1995; Elloway et al., 2006).

95 Finally, many soil microbiota live protected within biofilms. Plastic particles were shown to be
96 a potential surface for the formation of those biofilms (Lobelle and Cunliffe, 2011), which are a
97 food source for grazing primary consumers. [4] Inadvertent ingestion might also transfer
98 occluded or abraded MP to higher trophic levels.

99 But what about the larger organisms that feed on all these, free plastic particles,
100 contaminated microorganisms, biofilms and one another? Recent work discussed the effects
101 of MP on soil biota (Chae and An, 2018) or called for intensified research on certain
102 taxonomic groups (Rillig and Bonkowski, 2018). Thus, we were motivated to give on our part
103 a review with focus on the most-produced plastics and their passive translocation, ingestion,
104 bioaccumulation and adverse effects on the multicellular soil fauna. The types, sizes and
105 shapes of plastic used in former laboratory studies were compared with the available
106 knowledge on plastic in the environment, and recommendations are given for future research.
107 [12] This analysis aims to support the assessment of the influence of MP on the ecosystem
108 services provided by diverse soil organisms.

109 2 Search pattern

110 Within the tree of life, edaphic branches were identified comprising taxa that permanently
 111 inhabit the soil, are both-sided part of the soil food web and/or the burrowing macro- and
 112 megafauna or have active subterranean larval stages. The resulting tree of soil life based on
 113 the NCBI taxonomy database (Fig. 1) was drawn using the software [phyloT](#) and shows the
 114 leading taxonomic rank, which is mainly the family, but in exceptions – e.g. if one species
 115 represents the only soil-born between many aquatic – a lower rank.




Figure 1: Tree of edaphic fauna. Taxonomic ranks, that were examined in this qualitative study, are placed at the outer rim of the diagram. The length of the connecting line between two taxa represents the grade of phylogenetic relationship.

117 A pattern of search terms was established (71 Table 1), consisting of „taxon“ (Linné’s
 118 binominal nomenclature, common name, plural-sensitive search), „plastic type“ (plastic,
 119 microplastic, nanoplastic, PE or polyethylene, PP or polypropylene, PVC or polyvinyl chloride,
 120 PS or polystyrene, PU or polyurethane, PET or polyethylene terephthalate and latex) and
 121 „common shapes“ (fragments, particles, fibers, microfibrs, beads, microbeads,
 122 microspheres). 71 Some type- shape combinations caused problems, as they led to a very
 123 large amount of unuseful, off-topic papers – e.g. using any taxon combined with PET, papers
 124 with the use of PET bottles in experimental set-ups were selected or also studies on pets.
 125 Those combinations of search terms were excluded from this pattern. Further plastic types
 126 and shapes occurring within the found studies were also included in the review. Data on
 127 microspheres and microbeads were pooled, as both names describe one and the same.

128

Table 1: 51 Types and shapes of microplastic particles in edaphon studies within this review. (X) symbolizes combinations excluded from the search pattern. 71 The number counts for how often type-shape combinations were used in all reviewed experimental setups independently of organism. Empty fields stand for zero results. Microbeads and microspheres are often mixed up terms and, thus, counted together.

Organism: Linné’s systematic names OR common name	fragments	particles	fibrs	microfibrs	beads	microbeads	microspheres	other, diverse, N/A
plastic	X							
microplastic								
nanoplastic								
PE OR polyethylene	X	4	10	1	1	1	4	7
PP OR polypropylene	X	1						
PVC OR polyvinyl chloride	X	4	6	1				
PS OR polystyrene	X	6	3				24	4
PU OR polyurethane	X							
PET OR polyethylene terephthalate	X	3	2				X	
latex	X			X			6	
other		6	3	1				1
N/A		1	1		2			3

129

130

131 The search was applied 5151 between June 2019 and January 2020 within the Web of Science
 132 Core Collection Database, repeated in the first week of January 2020 and covers publications
 133 until January 2020. The search strings result from combinations of taxon, plastic type and
 134 particle shape shown in Table 1. Based on the search pattern, data on passive transport,

135 ingestion, bioaccumulation and adverse effects were collected for each edaphic group.
136 Studies that only use uncommon, local, outdated, weird or nicknames are excluded by the
137 search pattern. Studies testing injection to tissues, lymph or blood were excluded, as they do
138 not represent natural ways to incorporate MPs. Data on inhalation by the megafauna in fact
139 represent a natural way of uptake, but were also excluded as they are exclusively related to
140 above-ground organisms, that only occur on the outer edge of the food-web. Also running
141 debates on phylogenetic classifications are not part of this work and the [12] taxonomists will
142 be able to adjust the branches accordingly to [12] their purpose.

143 The data of related taxonomic groups were pooled and evaluated for their environmental
144 representativity based on exposure time, plastic concentrations and properties used. From
145 this synthesis recommendations for a structured experimental design were derived for
146 application in future studies.

147 3 Data collection

148 149 3.1 Insects

150 Within the Panarthropoda, the insects comprise the highest taxonomic diversity. And,
151 regarding MPs, they represent an unevenly studied taxonomic group.

152 Within the Insecta, the **Coleoptera** (beetles) are an extraordinarily diverse and abundant
153 taxon. Studies on plastic uptake into adult individuals mainly focus on the subfamily of
154 Scarabaeinae (dung beetles). Comprehensive experiments with latex microbeads showed,
155 that many species only ingest [12] smaller particles with maximum diameters of about 10 to
156 83 µm and retain them within the gut – with a slightly positive dependency on body size.
157 Larger particles were rejected by a filtering mechanism within the mouth region and not
158 ground with the mandibles (Holter, 2000; Holter et al., 2002; Holter and Scholtz, 2005).
159 Beside those on Nematods, these data comprise by far the most detailed information about
160 [12] size-dependent uptake of MP particles compared to other edaphic taxa. This gives a good
161 foundation for future studies on adverse concentrations. In addition, several studies with
162 plastic as predominant food source could show chewing, ingestion and intestinal degradation
163 of different PS and PE foams in feeding experiments with *Tenebrio sp.* larvae (mealworms).
164 These experiments also pointed out an alteration of the gut microbiome, but no adverse
165 effects on reproduction and survival, with only one case of a non-significant tendency to
166 higher mortality after 1 month of exposure (Yang et al., 2015; Brandon et al., 2018; Yang et
167 al., 2018; Peng et al., 2019).

168 The **Isoptera** (termites), recently categorized as part of the order Blattodea, are the oldest
169 social insects with a tribal history of about 130 million years (Korb, 2008). Especially in arid
170 ecosystems with a lack of earthworms they play an important role in homogenization of soils,
171 but also in sorting of soil mineral particles for building mounds as well as decomposition and
172 distribution of organic matter (De Bruyn and Conacher, 1990). Tsunoda et al. (2010) and Lenz
173 et al. (2012) could show, that different termite species are picky feeders and erode PE, but
174 avoid other plastic cable sheathings. This suggests the excretion of ground MP particles by
175 termites, but metabolic impacts are unknown. In contrast to termites, data on **other Blattodea**
176 (e.g. cockroaches) were not found.

177 The suborder **Apocrita** comprises some flying insects, that inhabit burrows within the soil,
178 such as ground-dwelling wasps within the **Vespidae** superfamily, mining bees within the
179 **Apoidea** superfamily and the **Spheciformes**. They mostly do not prey and feed on
180 subterrestrial organisms, but may move MP particles into the ground, as implied by a report of
181 Allasino et al. (2019) on [12] solitary bees, which built nests fully made of plastic fragments.
182 The Apocrita also contain the **Formicidae** (ants). Some ant species are considered an
183 important factor for seed dispersal, a behavior, that could also be shown for artificial plastic
184 seeds with ~2 mm diameter (Hughes and Westoby, 1992; Angotti et al., 2018). Robins and

185 Robins (2011) found that this also includes differently shaped cultural objects: *Rhytidoponera*
186 *metallica*, a representative of ground-nesting, omnivore ants, is capable not only of a
187 remarkable bioturbation, but also of an active, apparently random burying of anthropogenic
188 plastic artifacts >1 mm. Seeds are used as a food source, thus, the ingestion of plastic bites is
189 conceivable, but not documented. The uptake of latex microspheres $\geq 0.88 \mu\text{m}$ with liquids by
190 larvae of *Solenopsis invicta* seems to be prevented by filtration within the mouth and the
191 particles are released as larger aggregates, whereas other species ingest by far larger
192 particles up to $150 \mu\text{m}$ (Glancey et al., 1981). However, also here data on adverse effects are
193 missing.

194 Further insects with edaphic adult stages, e.g. **Dermaptera** (earwigs), **Heteroptera** (true
195 bugs) and **Zygentoma** (silverfish, fishmoth, firebrat) or soil- or litter-dwelling larvae such as
196 **Embiopoda** (webspinners, footspinners), **Thysanoptera** (thrips), **Psocoptera** (booklice,
197 barklice, barkflies), **Neuroptera** (lacewings), **Raphidioptera** (snakeflies) or **Zoraptera** (angel
198 insects) were not yet researched with focus on soil MP.

199 Regarding insects, mainly studies on translocation and uptake of MP were carried out. In
200 contrast, work on bioaccumulation is completely lacking and adverse effects are sparsely
201 tested using *Tenebrio sp.* larvae. Such studies could provide information whether or not the
202 input of MP in soil ecosystems is one of many factors causing the global decline of the
203 entomofauna (Oliveira et al., 2019; Sánchez-Bayo and Wyckhuys, 2019).

Table 2: Microplastic studies on Coleoptera, Blattodea (Blattod.), Apoidea (A.) and Formicidae (mb=microbeads, fr=fragments, ms=microspheres, b=beads, [8] N/A=information not available). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

	organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
Coleoptera	<i>Aphodius erraticus</i>	Petri dish	latex	N/A	N/A	N/A	mb	5	N/A	45 min	N/A	no	N/A	N/A	Holter (2000)
	<i>Aphodius rufipes</i>							2..39				≤14 µm			
	<i>Aphodius ater</i>							2..39				≤14 µm			
	<i>Aphodius fimetarius</i>							2..39				≤18 µm			
	<i>Aphodius contaminatus</i>							2..39				≤18 µm			
	<i>Aphodius fossor</i>							2..39				≤18 µm			
	diverse dung beetles	vial	latex	N/A	N/A	N/A	mb	2..83	N/A	45 min	N/A	≤10..≤60 µm	N/A	N/A	Holter et al. (2002)
	diverse dung beetles	N/A	latex	N/A	N/A	N/A	mb	2..83	N/A	45 min	N/A	≤4..≤95 µm	N/A	N/A	Holter and Scholtz (2005)
	<i>Tenebrio molitor</i> larvae	container	PS	N/A	N/A	no	foam	N/A	100% w/w (food)	31 d	N/A	yes	biodegrad.	N/A	Yang et al. (2015)
Blattod.	<i>Tenebrio molitor</i> larvae	container	LD-PE PS	N/A	N/A	no flame retardant	foam	8..27 cm ³	50..100% w/w (food)	32 d	N/A	yes	biodegrad.	microbiome	Brandon et al. (2018)
	<i>Tenebrio molitor</i> larvae	container	PS	N/A	N/A	N/A	foam	N/A	4..100% w/w (food)	32 d	N/A	yes	biodegrad.	no	Yang et al. (2018)
	<i>Tenebrio molitor</i> larvae	N/A	PS	N/A	N/A	no	foam	N/A	86..100% w/w (food)	31 d	N/A	yes	biodegrad.	microbiome	Peng et al. (2019)
	<i>Tenebrio obscurus</i> larvae														
A.	<i>Coptotermes formosanus</i>	mesocosm	LD-PE others	yes/no	N/A	N/A	cable sheets	4 cm, Ø 0.8 cm	N/A	42 d	N/A	yes no	N/A	N/A	Tsunoda et al. (2010)
	diverse termites	in situ	MD-PE PA	no	N/A	anti-oxidant stabilizer	cable sheets	30 cm, Ø 1.4 cm	N/A	6 yr.	N/A	yes no	N/A	N/A	Lenz et al. (2012)
Formicidae	<i>Megachile</i> sp.	in situ	N/A	N/A	N/A	N/A	fr	N/A	N/A	N/A	yes	N/A	N/A	N/A	Allasino et al. (2019)
	<i>Solenopsis invicta</i>	Petri dish	latex	N/A	N/A	fluorescence	ms	0.9..4.5	2.5% w/w (food)	direct	N/A	filtration	N/A	N/A	Glancey et al. (1982)
	<i>Rhytidoponera metallica</i>	in situ	N/A	N/A	N/A	N/A	b	N/A	50 items per nest	3 d	yes	N/A	N/A	N/A	Hughes and Westoby (1992)
	<i>Aphaenogaster longiceps</i>														
	<i>Pheidole</i> sp.														
Formicidae	<i>Rhytidoponera metallica</i>	mesocosm	N/A	N/A	N/A	N/A	diverse	<75.5 cm	N/A	26 mos.	yes	N/A	N/A	N/A	Robins and Robins (2011)
	diverse ants	in situ	N/A	N/A	attractant	N/A	b	1.8 cm	N/A	1 d	yes	N/A	N/A	N/A	Angotti et al. (2018)

205 3.2 Other panarthropods

206 Apart from the insects, **Acari** (mites) comprise many abundant soil-living taxa, that feed on
207 litter, fungi and fauna as predators and parasites and are bioindicators, as they are sensitive
208 to changes in the soil physiochemical environment (Gulvik, 2007). Experiments indicated, that
209 mites passively transport MP due to pushing and dragging after attachment to their cuticle, as
210 shown with 80 to 250 μm sized PVC particles in a **Petri** dish experiment without soil (Zhu et
211 al., 2018a). The population within manure pats slightly declines when exposed to mm-sized
212 unweathered PE and PS particles at concentrations of 5 % v/v and declines strongly at
213 ≥ 60 % v/v (Stamatiadis and Dindal, 1990). This could probably be an effect of moisture
214 deficiency due to a reduced water holding capacity in an unnaturally enriched substrate, but
215 not necessarily through plastic intake. In contrast, no data was found on their arachnoid,
216 preying relatives, the order of **Pseudoscorpiones** (false scorpions).

217 Just as many other highly abundant and diverse representatives of the soil mesofauna, the
218 **Oniscideae** (woodlice) contribute to the decomposition of litter by chewing and passage
219 through their digestive system (Warburg, 1987) and react strongly to environmental pollution,
220 thus, they are potentially used as bioindicators (van Gestel et al., 2018). They practice a strict
221 selection of natural food sources (Hassall and Rushton, 1984). This is also demonstrated for
222 starch and cellulose based plastic films (4 cm^2), which were consumed and digested in
223 experiments with the model organism *Porcellio scaber*, in contrast to PHB
224 (polyhydroxybutyrate) films, that reduces the feeding rate (Wood and Zimmer, 2014). Smaller
225 PE particles ($137 \pm 51 \mu\text{m}$ and $183 \pm 93 \mu\text{m}$) embedded into food pellets (0.4 % w/w) were taken
226 up easily by *Porcellio scaber*, and the smaller fraction caused a slight and non-significant
227 reduction of body mass after 14 days of exposure, but not of feeding, defecation or energy
228 reserves (Kokalj et al., 2018).

229 Other panarthropodean groups are even less studied in terms of MP. We did not find literature
230 on the subphylum of Myriapoda containing the classes of **Diplopoda** (millipedes), **Chilopoda**
231 (centipedes), **Paupoda** and **Symphyla** (pseudocentipedes or symphilids), which are
232 important litter-feeders and predators within various soil ecosystems.

233 The situation is nearly similar with the phylum of **Tardigrada** (water-bears or tardigrades), that
234 has many ecologically relevant and well studied species feeding on microorganisms and
235 detritus particles. Sparse field research in semisubhydric environments showed no uptake of
236 MP **fibers** by tardigrada (Gusmão et al., 2016), but comprehensive data on terrestrial soils are
237 lacking. ¹³ Similarly, the related phylum of **Onychophora** (velvet worms), primordial
238 invertebrates that are mainly native in litter and soils with high water holding capacity under
239 pleistocene-like forest vegetation within tropical and moderate regions (Monge-Nájera, 1994).

240 ~~Another branch within the panarthropoda, the phylum of **Onychophora** (velvet worms),~~
241 ~~comprises primordial invertebrates that are mainly native in litter and soils with high water~~

242 holding capacity under pleistocene-like forest vegetation within tropical and moderate regions
243 (Monge-Nájera, 1994). As predators, they most likely take up plastic debris appearing within
244 or on their prey, but no studies on MP are available, most likely due to their remote habitats,
245 low abundance and little scientific focus.

246 The phylum of **Collembola** (springtails) [11], together with the **Diplura** and **Protura** (Westheide
247 and Rieger, 1996; Pass et al., 2011), an abundant, diverse and ubiquitous soil-borne phylum
248 with a broad spectrum of food sources (Hopkin, 1997), also represent an intensively studied
249 group within the Arthropoda. Together with the **Diplura** (which mainly live in tropic and
250 subtropic regions in litter and humid topsoil and feed on fungal hyphae, POM and prey)
251 (Westheide and Rieger, 1996) and the **Protura** (Pass et al., 2011), the **Collembola** build an
252 intensively studied morphological group, that [12] exhibits similar ecological functions, such as
253 distribution and decomposition of organic matter as well as the control of fungal abundance
254 (Hopkin, 1997). Springtails provide up to 27 % of the soil biomass and up to 33 % of the total
255 soil respiration (with higher shares in colder ecosystems) (Petersen, 1994) with up to 100000
256 individuals per square meter (Hopkin, 1997). Thus, their well-being plays an important role for
257 ecosystem functioning.

258 In a Petri dish experiment without soil, Maaß et al. (2017) showed the passive transport of
259 urea-formaldehyde particles <400 µm and undefined PET fragments by two Collembola
260 species (*Folsomia candida* and *Proisotoma minuta*) due to attachment, but found no
261 ingestion. Within a soil matrix, trials of Kim and An (2019) indicated hindrance of collembolan
262 migration by larger PS particles (44±39, 282±131 and 676±479 µm) at concentrations of
263 1000 mg kg⁻¹ corresponding to highly contaminated soils. In addition, they found suppressed
264 mobility due to the attachment of even smaller PS microbeads (0.47 to 0.53 µm) at
265 concentrations of 8 mg kg⁻¹ dry soil, which is equivalent to values found in nature. Small
266 particles <50 µm were moved, while larger particles were most likely [12] cast off. When *F.*
267 *candida* encounters two of its predators, the mites *Damaeus exspinosus* and *Hypoaspis*
268 *aculeifer*, the dispersal of 80 to 250 µm PVC particles is enhanced as shown by Zhu et al.
269 (2018a) in a Petri dish experiment. Without proving the ingestion or the minimal effective MP
270 concentration, Zhu et al. (2018b) published an alteration of the gut microbiome and adverse
271 effects on growth and reproduction of *F. candida* by 80 to 250 µm PVC particles mixed in soil
272 at concentrations of 1000 mg kg⁻¹ dry soil. These data were not considered robust (van Gestel
273 and Selonen, 2018), but match with a later study that found inhibited reproduction at
274 ≥1000 mg kg⁻¹ and avoidance behavior as well as microbiome alteration at ≥5000 mg kg⁻¹ (Ju
275 et al., 2019). Such concentrations can occur in highly contaminated soils (Fuller and Gautam,
276 2016). However, documentations on the active uptake, gnawing and grinding of MP by
277 springtails proposed by Rillig (2012) is still lacking and also studies on Diplura and Protura
278 were not found.

Table 3: Microplastic studies on Acari, Oniscidea (Onisc.), Tardigrada (T.) and Collembola (fr=fragments, p=particles, mf=microfibers, mb=microbeads, ms=microspheres, ^s=semisubhydic, [8] N/A=information not available). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

	organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
Acari	diverse mites	microcosm	PE PS	no	N/A	N/A	fr	<4800 >2000	0..90% v/v (manure)	16 d	N/A	N/A	N/A	≥5% v/v: abundance ↓	Stamatiadis and Dindal (1990)
	<i>Hypoaspis aculeifer</i> <i>Damaeus exspinosus</i>	Petri dish	PVC	N/A	no	N/A	p	80..250	5000 items per dish	N/A	yes	N/A	N/A	N/A	Zhu et al. (2018a)
Onisc	<i>Porcellio scaber</i>	mesocosm	PHB	no	N/A	N/A	fr	4 cm ²	1 item per cosm	14 d	N/A	yes	N/A	feeding ↓	Wood and Zimmer (2014)
	<i>Porcellio scaber</i>	Petri dish	PE	N/A	N/A	N/A	fr	183±93 137±51	0.4% w/w (food)	14 d	N/A	yes	N/A	no	Kokalj et al. (2018)
T.	diverse tardigrades ^s	in situ	N/A	N/A	N/A	N/A	mf	N/A	N/A	N/A	N/A	no	N/A	N/A	Gusmão et al. (2016)
Collembola	<i>Folsomia candida</i> <i>Proisotoma minuta</i>	cup	UF, PET	N/A	no	N/A	p,fr	<400	2.5..5 mg per cup	N/A	yes	N/A	N/A	N/A	Maaß et al. (2017)
	<i>Folsomia candida</i>	Petri dish	PVC	N/A	no	N/A	p	80..250	5000 items per dish	N/A	yes	N/A	N/A	N/A	Zhu et al. (2018a)
	<i>Folsomia candida</i>	microcosm	PVC	N/A	no	N/A	p	80..250	1000	56 d	N/A	N/A	N/A	microbiome, growth ↓, reproduction ↓	Zhu et al. (2018b)
	<i>Folsomia candida</i>	microcosm	PE	N/A	no	N/A	mb	<500	0..10000 0..10000 0..5000	7 d 28 d 28 d	N/A	N/A	N/A	≥5000: avoidance ≥1000: reproduction ↓ ≥5000: microbiome	Ju et al. (2019)
	<i>Lobelia sokamensis</i>	soil sample	PS	N/A	carboxyl	fluorescence	mb	0.5	4..8		yes				
			PE	no	N/A	fluorescence	ms	27..32	1000		yes				
			PE	no	N/A	fluorescence	ms	250..300	1000		N/A				
			PS	no	N/A	no	fr	44±39	1000	≤3 min	yes				
			PS	no	N/A	no	fr	282±131	1000		N/A				
			PS	no	N/A	no	fr	676±479	1000		N/A				
												N/A	N/A	avoidance, motivity ↓	Kim and An (2019)

280 3.3 Annelida

281 Land-based Annelida comprise another large group of invertebrates. The Lumbricidae
282 (earthworms) are a well-studied family (Darwin, 1881; Lavelle et al., 2006), represented in
283 high abundance and diversity in many ecosystems all around the world (Phillips et al., 2019).
284 Earthworms are often used as indicators for soil health (Fründ et al., 2011; Pulleman et al.,
285 2012), as they are ecosystem engineers which through their burrowing activity influence
286 various soil physical, chemical and biological processes (Jouquet et al., 2006; Lavelle et al.,
287 2006).

288 By far the most of the studies on the influence of MP on earthworms are performed with PE
289 and the species *Lumbricus terrestris* or *Eisenia fetida*, but there are also single studies with
290 *Aporrectodea rosea* (Boots et al., 2019) and *Eisenia andrei* (Rodriguez-Seijo et al., 2017) and
291 with the less common species *Metaphire californica* (Wang et al., 2019b). We found one field
292 study of earthworms and MPs (Huerta Lwanga et al., 2017a) among many laboratory
293 experiments with MPs mixed into soil volumes (concentrations ranging up to 20000 mg kg⁻¹
294 dry soil) or applied with litter on top of the soil surface (≤60% w/w). The particles sizes were
295 usually <1 mm in diameter, but some were even up to 2x2 cm², and the duration of
296 experiments was generally 14 to 28 days, few lasted up to 60 days.

297 The uptake of MPs of a broad size range by earthworms was shown in studies based on
298 particles in earthworm casts of *Lumbricus terrestris* (Huerta Lwanga et al., 2016; Cao et al.,
299 2017; Hodson et al., 2017; Rillig et al., 2017; Prendergast-Miller et al., 2019; Yu et al., 2019;
300 Huerta Lwanga et al., 2017a), *Eisenia fetida* (Rodríguez-Seijo et al., 2018; Chen et al., 2020;
301 Wang et al., 2019c), *Eisenia andrei* (Rodriguez-Seijo et al., 2017) and *Metaphire californica*
302 (Wang et al., 2019b). Zhang et al. (2018) showed that relatively large PE particles of
303 1.5 x1.5 cm² are not ingested by *Lumbricus terrestris*, but partial ingestion of such large
304 particles of biodegradable MPs does take place after initial weathering in soil or in compost
305 has occurred. In some laboratory experiments, MPs were found in the gut of dissected
306 earthworms (Huerta Lwanga et al., 2016; Hodson et al., 2017; Rodriguez-Seijo et al., 2017),
307 but the concentration of MPs in the gut was not significantly different between treatments nor
308 significantly different from the bulk soil concentration, so there was no evidence of
309 accumulation of MPs in the earthworm bodies (Hodson et al., 2017). Chen et al. (2020)
310 assume an accumulation of MP takes place in *Eisenia fetida*, based on an observed increase
311 of MP concentrations in the casts in the course of 4 weeks. Huerta Lwanga et al. (2017a)
312 supposed an accumulation of MPs in the food chain as the concentration of MPs in chicken
313 gizzards is strongly increased compared to that in the earthworm casts in the same
314 experiments. However, mainly the amount of large particles, i.e. macroplastics, in the gizzards
315 was very large, thus it seems likely that the chicken directly fed on plastics and an
316 accumulation through the food chain cannot be proven with the current knowledge and should
317 be further investigated.

318 Several studies did not find significant negative effects of MPs on earthworms' avoidance
319 behaviour (Judy et al., 2019), nor on growth (Hodson et al., 2017; Rodriguez-Seijo et al.,
320 2017; Judy et al., 2019; Wang et al., 2019c), mortality (Hodson et al. (2017); Rillig et al.
321 (2017); Rodriguez-Seijo et al. (2017); Judy et al. (2019); Prendergast-Miller et al. (2019) or
322 reproduction (Huerta Lwanga et al., 2016; Rodriguez-Seijo et al., 2017). However, other
323 studies do show adverse effects of the uptake of MP in different degrees and on different
324 aspects of earthworms' fitness: A reduced growth was shown by Cao et al. (2017) for *Eisenia*
325 *Fetida* and the mortality increased at an exposure of concentrations ≥ 10000 mg kg⁻¹ dry soil.
326 At lower concentrations no significant effects were found. The growth of *Aporrectodea rosea*
327 was also inhibited when exposed to biodegradable polylactic acid, conventional high-density
328 polyethylene (at 1000 mg kg⁻¹ dry soil), and MP clothing fibers (at 10 mg kg⁻¹ dry soil) (Boots
329 et al., 2019). Huerta Lwanga et al. (2016) showed a decrease in growth and increased
330 mortality at concentrations $\geq 28\%$ w/w in litter and after 60 days, though after just 14 days no
331 mortality occurred in these experiments.

332 In some studies, additional effects such as histopathological changes or stress biomarkers
333 were measured. For *Eisenia fetida* Chen et al. (2020) observed skin damage at
334 1500 mg MP kg⁻¹ in soil, measured an increase in catalase activity and malondialdehyde
335 content at 1000 mg kg⁻¹ and at ≥ 1000 mg kg⁻¹ acetylcholine esterase was significantly
336 stimulated. Wang et al. (2019c) tested *Eisenia fetida* and found that MPs only increased the
337 catalase and peroxidase levels as well as the level of lipid peroxidation and decreased the
338 activity of superoxide dismutase and glutathione S-transferase at an exposure of
339 200000 mg kg⁻¹ dry soil for 14 days. No discernible influence was found at 100000 mg kg⁻¹.
340 However, Rodríguez-Seijo et al. (2018) also found for *Eisenia fetida* a significant positive
341 correlation of MP concentration with different biomarker responses: catalase, glutathione S-
342 transferase, lactate dehydrogenase and thiobarbituric acid reactive substances. In addition,
343 Rodriguez-Seijo et al. (2017) observed histological damage of the gut and occurrence of
344 inflammatory processes as well as an increase of stress response indicators associated with
345 MP exposure of *Eisenia andrei*. For *Lumbricus terrestris* Prendergast-Miller et al. (2019)
346 showed an increase in metallothionein expression at an exposure with ≥ 1000 mg kg⁻¹ dry soil
347 and a decrease in heat shock protein 70 at a concentration of ≥ 10000 mg kg⁻¹.

348 Due to the large differences in experimental conditions – e.g. size of the MPs, addition of MPs
349 to soil or to litter, duration of experiments, earthworm species – the current knowledge is not
350 sufficient to detect whether there is a threshold in MP size and concentration at which the MP
351 become harmful for earthworms and how this threshold differs for different earthworms
352 species and MP shapes. The results of Huerta Lwanga et al. (2016), who found no effects of
353 MPs on earthworms at 14 days, but significant influence on growth and mortality after
354 60 days, indicate the importance of longer measurements. This is consistent with Pelosi et al.

355 (2015), who concluded that the influence of pesticides on earthworm communities should be
356 tested in long term field experiments.

357 Earthworms activity also increased the transport of MP in soil columns to deeper soil layers
358 (Rillig et al., 2017; Yu et al., 2019; Huerta Lwanga et al., 2017b). The smaller the MP the
359 stronger the transport. Particles are transported both actively – ingested and later cast out –
360 and passively after attachment to the earthworm's body or by water flow through the biopores.
361 As Huerta Lwanga et al. (2018) showed that the bacteria in the gut of *Lumbricus terrestris* can
362 decompose MPs, it seems likely that particles taken up at the surface are egested as smaller
363 particles in deeper layers.

364 Microplastics might well serve as a vector for contaminant transport to soil organisms. Though
365 adsorption on plastics was seen to be lower than on the soil matrix, the desorption of Zn was
366 seen to be higher in synthetic earthworm guts. However, there was no measurable negative
367 effect of Zn or the PE on *Lumbricus terrestris* (Hodson et al., 2017). Wang et al. (2019b)
368 studied the influence of MP on arsenic uptake and negative effects on *Metaphire californica*
369 and concluded that MPs decreased the uptake of arsenic and that MPs reduced the influence
370 of arsenic on the gut bacterial communities. Rodríguez-Seijo et al. (2019) showed altered
371 enzyme activities and enhanced avoidance behavior in face of LD-PE pellets spiked with the
372 insecticide chlorpyrifos. Yang et al. (2019a) studied the influence of MPs on the transport of
373 glyphosate, however they mainly showed that the glyphosate transport was increased by
374 earthworm activity, the role of MPs in this transport could not be determined with this study.
375 These studies show that MP might have very different influences on the uptake and the
376 adverse effects of different pollutants on earthworms and further investigation is needed in
377 order to understand the influence of MPs on pollutant transport.

378 In contrast to the recently well-researched Lumbricidae, a near relative, the family of
379 **Megascolecidae** (giant earthworms), is not yet mentioned in literature. Another branch within
380 the Annelida, the small **Enchytraeidae** (potworms), were shown to suffer adverse effects on
381 body weight and microbiome with PS microspheres (0.05 to 0.1 μm) at concentrations of
382 $\geq 10\%$ w/w within their food source, but an unexpected increase of reproduction at 0.5 % w/w
383 (Zhu et al., 2018b). The reproduction was reduced at abnormal concentrations of
384 90 g kg^{-1} dry soil of polyamid particles (13 to 150 μm), but not with PVC (Lahive et al., 2019).

385 The edaphon of semisubhydric soils is often treated as a marginal group between the area of
386 interest of soil and aquatic scientists. As a highly diverse soil biocenosis outside the focus of
387 this paper, the benthos along seashores and fresh waters is also affected by MPs and should
388 therefore be shortly mentioned by reviewing the lugworm **Arenicola marina**, a well examined
389 deposit-feeder of the tidal flats. In situ, MP accumulates within its tissue and feces (Van
390 Cauwenberghe et al., 2015). In laboratory experiments, PS particles $\geq 500\ \mu\text{m}$ were avoided
391 as food-source and passively translocated within the sediment at concentrations of $\sim 2\ \text{g}\ \text{kg}^{-1}$

392 (Gebhardt and Forster, 2018), but were measured within the feces at $\sim 74 \text{ g kg}^{-1}$ causing
393 effects on feeding activity and body weight with no influence on the survival rate (Besseling et
394 al., 2012). PS microspheres $\leq 30 \mu\text{m}$ remained within the animal without any adverse effects
395 regardless of particle size (Van Cauwenberghe et al., 2015). Other studies found adverse
396 effects on respiration, energy reserves, feeding, egestion and casting after uptake of PVC
397 particles $\leq 478 \mu\text{m}$ at different sediment concentrations of $> 2 \text{ g kg}^{-1}$, but neither on biomass
398 and survival nor due to HD-PE (Wright et al., 2013; Green et al., 2016). There is further a
399 difficulty in distinguishing between the adverse effects of MPs and substances adsorbed on or
400 leached from MPs (Besseling et al., 2012). When adding PCB-spiked PE to mud flat sediment
401 with concentrations up to **121 5000 mg kg⁻¹** dry mass, there was no significant change of
402 survival rate or body weight. The decreased feeding activity and heap mass could be
403 attributed to increasing plastic concentrations, but not to enhanced PCB bioaccumulation via
404 PE uptake (Besseling et al., 2017). However, all these studies found adverse effects at MP
405 concentrations orders of magnitude above natural values.

Table 4: Microplastic studies on Lumbricidae (p=particles, ms=microspheres, b=beads, f=fibers, ms=microfibers, [8] N/A=information not available). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
<i>Lumbricus terrestris</i>	mesocosm	PE	washed (C ₈ H ₁₂ , C ₈ H ₁₈)	N/A	N/A	p	<150	0..60% w/w (litter)	14 d / 60 d	yes	yes	N/A	at 60 d, ≥28% w/w: survival ↑, growth ↓	Huerta Lwanga et al. (2016)
<i>Eisenia fetida</i>	glass beaker	PS	N/A	N/A	N/A	ms	50..80	0..20000	30 d	N/A	yes	N/A	≥5000: survival ↓ ≥10000: weight ↓	Cao et al. (2017)
<i>Lumbricus terrestris</i>	bag	PE	N/A	N/A	N/A	p	0.92±1.09 mm ²	3500	28 d	N/A	yes	no	no	Hodson et al. (2017)
<i>Lumbricus terrestris</i>	home yard	diverse	yes	N/A	N/A	N/A	N/A	0.87±1.9 items g ⁻¹	N/A	N/A	yes	conc. in chickens > in earthworms	N/A	Huerta Lwanga et al. (2017a)
<i>Lumbricus terrestris</i>	mesocosm	PE	washed (C ₈ H ₁₂ , C ₈ H ₁₈)	N/A	N/A	p	<150	0..60% w/w (litter)	14 d	yes	yes	N/A	N/A	Huerta Lwanga et al. (2017b)
<i>Lumbricus terrestris</i>	mesocosm	PE	N/A	no	no	b	710..2800	750 µg on 2.5 kg soil	21 d	yes	yes	N/A	no	Rillig et al. (2017)
<i>Eisenia andrei</i>	mesocosm	LD-PE	N/A	N/A	N/A	pellets	250..1000	0..1000	28 d	N/A	yes	N/A	≥62.5: intestinal damage	Rodriguez-Seijo et al (2017)
<i>Lumbricus terrestris</i> (gut bacteria)	mesocosm glass bottle	PE	washed (C ₈ H ₁₂ , C ₈ H ₁₈)	N/A	N/A	p	150	7% w/w (litter) 10000	60 d (earthworms) 21 d (bacteria)	N/A	yes	N/A	N/A	Huerta Lwanga et al. (2018)
<i>Eisenia fetida</i>	mesocosm	LD-PE	washed (EtOH)	N/A	N/A	pellets	250..1000	0..1000	28 d	N/A	yes	N/A	≥125: altered enzyme activity	Rodriguez-Seijo et al (2018)
<i>Aporrectodea rosea</i>	mesocosm	PLA, PE N/A	N/A	N/A	N/A	p f	N/A	1000 10	30 d	N/A	yes	N/A	growth ↓	Boots et al. (2019)
<i>Eisenia fetida</i>	mesocosm	HD-PE, PET, PVC	no	N/A	no	f	<2000	soil extract	48 h / 56 d	N/A	N/A	N/A	no	Judy et al. (2019)
<i>Lumbricus terrestris</i>	bag	PE	N/A	N/A	N/A	mf	Ø40.7±3.8 x 361.6±387.0	0..10000	35 d	N/A	yes	N/A	≥1000: metallothionein expression ↑ ≥10000: heat shock protein 70 ↓	Prendergast-Miller et al. (2019)
<i>Eisenia fetida</i>	mesocosm	LD-PE	washed (EtOH)	N/A	chlorpyrifos (CPF)	pellets	5000 250..1000	40 items on 0.5 kg soil 180..200 items on 0.5 kg soil	14 d	N/A	N/A	N/A	with CPF: altered enzyme activity, avoidance of MPs	Rodriguez-Seijo et al (2019)
<i>Metaphire californica</i>	mesocosm	PVC	N/A	N/A	sodium arsenate	p	N/A	2000	28 d	yes	yes	N/A	microbiome	Wang et al. (2019b)
<i>Eisenia fetida</i>	glass beaker	PE PS	washed (MeOH)	N/A	PAHs, PCBs, Nile Red (NR)	p	<300 <250	0..200000 0..100	14 d 28 d	N/A	yes	N/A	≥200000: altered enzyme activity	Wang et al. (2019c)
<i>Lumbricus terrestris</i>	mesocosm	PE	washed (C ₈ H ₁₂ , C ₈ H ₁₈)	N/A	glyphosate	p	<150	0..7% w/w (litter)	14 d	N/A	N/A	N/A	N/A	Yang et al. (2019a)
<i>Lumbricus terrestris</i>	mesocosm	PE	N/A	N/A	N/A	N/A	<1000	7% w/w (litter)	14 d	yes	yes	N/A	N/A	Yu et al. (2019)
<i>Lumbricus terrestris</i>	Petri dish mesocosm	PE and div. biodegradables	unweathered, field or compost	N/A	N/A	p	1.5x1.5 cm ² 2x2 cm ²	4 items per dish 10 items per dish	14 d 50 d	yes yes	no yes	N/A	N/A	Zhang et al. (2018)
<i>Eisenia fetida</i>	bag	PE	washed (EtOH)	N/A	N/A	p	<400	0..1500	28 d	N/A	yes	yes	skin damage, ≥250 mg/kg: oxidative stress ≥1000 mg/kg: neurotoxicity ↑	Chen et al. (2020)

Table 5: Microplastic studies on Enchytraeidae and *Arenicola marina* (mb=microbeads, p=particles, ms=microspheres, sed.=sediment, ^s=semisubhydryc, [8] N/A=information not available). Concentrations refer to mg kg⁻¹ dry soil in terrestrial soils and mg kg⁻¹ dry sediment in semisubhydryc soils, if not specially marked.

	organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
Enchytr.	<i>Enchytraeus crypticus</i>	Petri dish	PS	N/A	N/A	N/A	mb	0.05..0.1	0..10% w/w (food)	7	N/A	yes	N/A	at 0.5% w/w: reproduction ↑ ≥10% w/w: microbiome, weight ↓	Zhu et al. (2018c)
	<i>Enchytraeus crypticus</i>	microcosm	PA PVC	N/A	N/A	fluorescence N/A	p	13..150 106..150	20000..120000 90000	20 h / 21 d	N/A	yes	N/A	≥90000: reproduction ↓ no	Lahive et al. (2018)
	<i>Arenicola marina</i> ^s	in situ liquid culture	N/A PS	N/A no	N/A N/A	N/A N/A	N/A ms	N/A 10..90	N/A 10000..50000 items kg ⁻¹	N/A 14 d	N/A N/A	yes yes	1.2±2.8 items g ⁻¹ 10 µm: 9600±1800 items kg ⁻¹ 30 µm: 800±700 items kg ⁻¹	N/A no	Cauwenberghe et al. (2015)
Arenicola	<i>Arenicola marina</i> ^s	mesocosm	PS PA	yes	biofilm	N/A	p	500..1000	~2000 ~1000	106..240 d	yes	no	N/A	N/A	Gebhardt and Forster (2018)
	<i>Arenicola marina</i> ^s	mesocosm	PS	N/A	N/A	N/A	p	400..1300	0..74000	28 d	N/A	≥400 µm	no	≥74000: feeding ↓, weight ↓	Besseling et al., (2012)
	<i>Arenicola marina</i> ^s	mesocosm	PVC HD-PE	N/A	N/A	N/A	p	9..478 3..316	0..20000 mg kg ⁻¹ wet sed.	31 d	N/A	N/A	N/A	>2000: respiration ↓, casting ↓ no	Green et al. (2016)
	<i>Arenicola marina</i> ^s	mesocosm	PE	N/A	PCBs	fluorescence	mb	10..180	0..5000	28 d	N/A	yes	no	feeding activity ↓, heap mass ↓	Besseling et al. (2017)
	<i>Arenicola marina</i> ^s	mesocosm	PVC	N/A	N/A	not leaching	p	~130	0..50000	28 d	N/A	N/A	N/A	≥10000: energy reserves ↓ ≥50000: feeding ↓, egestion ↓, casting ↓	Wright et al. (2013)

3.4 Further invertebrates

As part of the microfauna, the phylum **Nematoda** (nematodes or roundworms) is an ecologically important branch containing >25000 species (Zhang, 2013) in freshwater, marine, endobiotic and soil habitats. Due to their diverse trophic interactions nematodes hold a central position in both bottom-up and top-down controlled food webs (Yeates, 2001; Ferris, 2010) and thus most likely the uptake and transfer of MP.

Active feeding of adults and larvae of different species on 0.5 to 6 μm PS/latex microspheres (the size of their bacterial prey) was proven by Nika et al. (2016) and Fueser et al. (2019). However, most MP experiments on Nematodes are based on the bacterial-feeding model organism *Caenorhabditis elegans*. Kiyama et al. (2012) showed the favored uptake of PS microspheres with sizes of 0.5 to 3 μm by adult and 0.5 μm by larval *C. elegans*. The ingestion of MP decreased in the presence of bacteria as the natural food source.

When larval stages and adults ingested PS between 0.05 and 5 μm within an aqueous suspension or on agar plates, adverse effects such as oxidative stress, neurodegeneration, intestinal and DNA damage or dysfunction in motility, growth, life span, defecation, reproduction or energy metabolism appeared from a wide spectrum of concentrations from $\geq 1 \mu\text{g l}^{-1}$ up to $\geq 86.3 \text{ mg l}^{-1}$ (Zhao et al., 2017; Dong et al., 2018; Kim et al., 2019; Lei et al., 2018a; Lei et al., 2018b; Qu et al., 2019a). These effects are **not seen** below $1 \mu\text{g l}^{-1}$ (Qu et al., 2019b), and are enhanced due to amino modifications on micropsphere surfaces (Qu et al., 2019c). The incubation on agar plates with PE, PP and PVC particles $< 70 \mu\text{m}$ caused similar influences on survival, fertility, brood size and intestinal function (Lei et al., 2018b). Leachates from soils amended with 5 mg kg^{-1} dry soil of HD-PE and PVC decreased reproduction in laboratory cultures, but there was no effect shown on survival and after application of PET (Judy et al., 2019). Furthermore, silica nanoparticles (0.05 μm) are not only taken up orally but also via the vulva and spermathecae and migrate into gonad cells (Scharf et al., 2013). This process was confirmed for PS nanoparticles with the potential of a transfer to the progeny (Zhao et al., 2017).

The clear adverse effects of these studies are limited in their representativity by a narrow restriction to liquid cultures and a single model organism. Broader studies like on prominent soil-born nematodes such as *Acrobeloides buetschlii* (Frey, 1971) are still lacking. When assuming in first proximity mg l^{-1} solution = mg kg^{-1} dry soil, the applied concentrations between 0.001 and 86.8 mg l^{-1} match lower levels of soil contamination.

Feeding studies on the phylum **Rotifera** with MPs are fully based on PS microbeads and model organisms of the planktonic genus *Brachionus*. However, this data can carefully be transferred to soil environments as also soil rotifers are aquatic organisms living in water-filled pores and waterfilms. Different *Brachionus sp.* ingest microbeads $< 10 \mu\text{m}$ with strong preference for particles the size of their natural food source, namely bacteria and algae with

445 2 to 5 μm in diameter (Vadstein et al., 1993; Heerkloß and Hlawka, 1995; Baer et al., 2008;
446 Jeong et al., 2016). The uptake appears to be selective as microbeads are fewer incorporated
447 than bacteria and algae (Vadstein et al., 1993). The egestion of particles $\leq 0.5 \mu\text{m}$ is hindered
448 compared to $6 \mu\text{m}$ (Jeong et al., 2016). In suspension, microbeads $\leq 0.5 \mu\text{m}$ cause adverse
449 effects on fertility and life span at $\geq 0.1 \text{ mg l}^{-1}$ as well as oxidative stress and less growth at
450 $\geq 10 \text{ mg l}^{-1}$ (Jeong et al., 2016; Sun et al., 2019).

451 Terrestrial mollusks comprise snails and slugs within the class of **Gastropoda**. These grazers
452 feed on bacterial biofilms, fungi and plant tissue (Parkyn and Newell, 2013). Studies on
453 terrestrial species are sparse, but data on the benthic *Littorina sp.* imply passive transport and
454 non-selective MP uptake by feeding on surfaces with contaminated feces and mucus trails of
455 other snails (Gutow et al., 2019). With focus on benthic snails, Imhof and Laforsch (2016)
456 found no significant influence on growth parameters and fertility of juveniles and adult
457 *Potamopyrgus antipodarum* even when a food source with 70 % w/w of 5 to $600 \mu\text{m}$ sized
458 fragments was given (a mixture of PA, PC, PET, PS, PVC). In contrast, adverse effects were
459 found in recent work on the terrestrial snail *Achatina fulica*, that showed uptake and complete
460 gastrointestinal passage within 48 h with partial degradation of PET fibers (appr.
461 $1258 \times 76 \mu\text{m}$), but reduced excretion and food intake as well as increased oxidative stress at
462 concentrations of $\geq 0.01 \text{ g kg}^{-1}$, $\geq 0.14 \text{ g kg}^{-1}$ and $\geq 0.71 \text{ g kg}^{-1}$ dry soil, respectively (Song et al.,
463 2019).

Table 6: Microplastic studies on nematods (ms=microspheres, fr=fragments, np=nanoparticles, mb=microbeads, ms=microspheres, ox.=oxidative, [8] N/A=information not available). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

Microplastic studies on nematods (ms=microspheres, fr=fragments, np=nanoparticles, mb=microbeads, ms=microspheres, ox.=oxidative). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

	organism	experimental environment	plastic type	aging	coating	additives	shape	size span [μm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
	<i>Caenorhabditis elegans</i>	agar plate	PS	N/A	carboxyl sulfate amino	fluorescence	ms	0.1..6.6	N/A	0.5..2 h	N/A	yes	0.5..3 μm	N/A	Kiyama et al. (2012)
	<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	carboxyl	fluorescence	ms	0.1	0.001..10 mg l ⁻¹	4.5 d	N/A	Yes	N/A	≥0.01 mg l ⁻¹ : motility ↓, growth ↓, defecation ↓, within gonads	Zhao et al. (2017)
	<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	ζ=-10mV	fluorescence	ms	0.1	0.00001..0.001 mg l ⁻¹	N/A	N/A	Yes	N/A	≥0.001 mg l ⁻¹ : motility ↓, ox. stress ↑	Dong et al. (2018)
	<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	N/A	preservatives, fluorescence	ms	0.05..0.2	0.001..86.8 mg l ⁻¹ 17.3..86.8 mg l ⁻¹	24 h	N/A	Yes	N/A	≥17.3 mg l ⁻¹ : motility ↓, reproduction ↓ ≥86.3 mg l ⁻¹ : ox. stress ↑ ≥17.3 mg l ⁻¹ : metabolic dysf.	Kim et al. (2019)
	<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	ζ=-10mV	fluorescence	ms	0.1	0.001..1 mg l ⁻¹	N/A	N/A	Yes	N/A	≥1 mg l ⁻¹ : neurodegeneration ≥0.01 mg l ⁻¹ : motility ↓	Qu et al. (2019a)
	<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	N/A	N/A	ms	0.1..5	1 mg l ⁻¹	3 d	N/A	Yes	N/A	motility ↓, survival ↓, growth ↓, ox. stress ↑, neurotoxicity	Lei et al. (2018a)
Nematoda	<i>Caenorhabditis elegans</i>	agar plate	PE, PP, PVC, PS	no	N/A	N/A	fr, ms	0.1..200	0.5..10.0 mg m ⁻²	2 d	N/A	Yes	N/A	≥0.5 mg m ⁻² : survival ↓ at 5 mg m ⁻² : growth ↓, reproduction ↓, ox. stress ↑, intestinal damage	Lei et al. (2018b)
	<i>Caenorhabditis elegans</i>	agar plate	PS	N/A	N/A	fluorescence	ms	0.1..5	2500 mg l ⁻¹	7 d	N/A	Yes	N/A	mainly 1μm: intestinal damage	Scharf et al. (2013)
	<i>Caenorhabditis elegans</i>	liquid culture	HD-PE, PET, PVC	no	N/A	no	fr	<2000	soil extract	72 h	N/A	N/A	N/A	within tissue and gonads	Judy et al. (2019)
	<i>Caenorhabditis elegans</i>	agar plates	latex	N/A	N/A	fluorescence	mb	0.5	N/A	30 min	N/A	yes	N/A	reproduction ↓	Nika et al. (2016)
	<i>Caenorhabditis elegans</i> <i>Panagrolaimus thienemanni</i> <i>Plectus acuminatus</i> <i>Poikilolaimus regenfussi</i> <i>Acrobeloides nanus</i> <i>Pristionchus pacificus</i> <i>Aphelenchoides parietinus</i>	liquid culture	PS	N/A	N/A	fluorescence	ms	0.5..6	3·10 ⁹ ..10 ¹⁰ items l ⁻¹ (~0.2..1200 mg l ⁻¹)	4..73 h	N/A	≤3μm ≤0.5μm ≤1μm ≤1μm ≤1μm ≤6μm no	N/A	N/A	Fueser et al. (2019)
	<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	N/A	N/A	ms	0.1	0.0001..0.001 mg l ⁻¹	N/A	N/A	N/A	N/A	no	Qu et al. (2019b)
	<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	no amino	N/A	ms	0.1	0.001..1 mg l ⁻¹	N/A	N/A	yes	N/A	≥0.01 mg l ⁻¹ : reproduction ↓, DNA damage ≥0.001 mg l ⁻¹ : reproduction ↓, DNA damage	Qu et al. (2019c)

Table 7: Microplastic studies on Rotifera and Gastropoda (ms=microspheres, mb=microbeads, fr=fragments, f=fibers, ox.=oxidative, pref.=preferential, ^p=planctic, ^b=benthic, [8] N/A=information not available). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

	organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
Rotifera	<i>Brachionus plicatilis</i> ^p	liquid culture	PS	N/A	carboxyl	fluorescence	ms	1.6..20	5·10 ⁹ µm ³ l ⁻¹ (~5.25 mg l ⁻¹)	35 min	N/A	≤10 µm	pref. 4.5 µm	N/A	Bear et al. (2008)
	<i>Brachionus plicatilis</i> ^p	liquid culture	latex	N/A	N/A	fluorescence	mb	0.3..3.1	3·10 ⁷ ..7·10 ⁸ items l ⁻¹ (~0.0004..11 mg l ⁻¹)	20 min	N/A	yes	pref. ≥2 µm	N/A	Vadstein et al. (1993)
	<i>Brachionus koreanus</i> ^p	liquid culture	PS	no	N/A	fluorescence	mb	0.05..6	0...20 mg l ⁻¹	1 d	N/A	yes	egestion rate 0.05 µm < 0.5 µm < 6 µm	≤0.5 µm, ≥0.1 mg l ⁻¹ : reproduction ↓, survival ↓ ≤0.5 µm, 10 mg l ⁻¹ : oxidative stress ↑	Jeong et al. (2016)
	<i>Brachionus plicatilis</i> ^p	liquid culture	PS	N/A	N/A	N/A	mb	0.07..7	0..20 mg l ⁻¹	N/A	N/A	yes	N/A	≤0.07 µm, ≥10 mg l ⁻¹ : reproduction ↓, growth ↓ ≤0.07 µm and ≥0.1 mg l ⁻¹ : survival ↓	Sun et al. (2019)
	<i>Brachionus quadridentatus</i> ^p <i>Brachionus plicatilis</i> ^p	liquid culture	PS	N/A	N/A	N/A	ms	2..10	N/A	8..10 d	N/A	pref. 3..5 µm pref. 2 µm	N/A	N/A	Heerkloß and Hlawa (1993)
Gastropoda	<i>Littorina littorea</i> ^b	microcosm	PMMA	N/A	N/A	fluorescence	fr	10..100	increasing	16 h	N/A	yes	N/A	N/A	Gutow et al. (2019)
	<i>Potamopyrgus antipodarum</i> ^b	aquarium	PET, PS, PVC, PA, PC	N/A	N/A	no	fr	5..600	0..70% w/w (food)	≤141 d	N/A	yes	N/A	no	Imhof and Laforsch (2016)
	<i>Achatina fulica</i>	mesocosm	PET	N/A	N/A	no / stained	f	approx. 1258x76 µm	10..710	28 d	N/A	yes	excretion after 48 hours	≥140: food intake ↓ ≥10: excretion ↓ ≥710: ox. stress ↑, gastrointestinal damage	Song et al. (2019)

466 3.5 Vertebrates

467 Different taxa of the class of Amphibia have a predator function within the edaphic food web
468 (e.g. preying on invertebrates) (Hebrard et al., 1992). While no data on the reaction to soil
469 MPs are available neither for the legless **Gymnophiona** nor for adults of the order **Anura**,
470 sparse data on tadpoles of aquatic frogs suggest uptake followed by regular excretion of PS
471 microspheres as shown with *Xenopus tropicalis* (Hu et al., 2016). Further, there exist no data
472 on the families **Serpentes** (snakes) and **Anguillidae** within the class of Reptilia, residing at the
473 outer rim of the food web.

474 Within the broad field of Mammalia, studies on MP ingestion are sparse and focus on **mice** as
475 a rodent model organism. Feeding of mice with PS microspheres of 1 to 14 μm in
476 concentrations of 1.49×10^6 to 4.55×10^7 particles at a volume of 10 ml kg^{-1} body weight for
477 4 weeks showed no adverse effects (Stock et al., 2019). In contrast, longer exposition
478 (6 weeks) with lower concentrations of particles with the same shape and size range changed
479 the mouse microbiome and caused metabolic and intestinal dysfunction (Lu et al., 2018; Jin et
480 al., 2019), which comes along with bioaccumulation within organs (Yang et al., 2019b). These
481 studies are regularly conducted with passive feeding and exclude active foraging on
482 perceptible plastic particles. However, the uptake via prey or feeding on contaminated roots
483 and litter is highly probable. Further Rodentia – **Cricetidae** (hamsters, lemmings, voles),
484 **Bathyergidae** (blesmols, mole-rats), **Octodontidae** as well as **Spermophilus** (ground
485 squirrels) and **Marmota** (marmots) within the family of **Sciuridae** – were not yet studied, just
486 as other mammalian (sub)orders like **Chrysochloridae** (golden moles), **Cingulata**
487 (armadillos), **Macroscelidea** (elephant shrews), **Notoryctemorphia** and **Peramelemorphia**.

Table 8: [3] Microplastic studies on Anura (An.) and Rodentia (ms=microspheres, ^a=aquatic, [8] N/A=information not available).

	organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
An.	<i>Xenopus tropicalis</i> ^a	Petri dish	PS	N/A	N/A	fluorescence	ms	1..10	100..10 ⁸ items l ⁻¹ (55·10 ⁻⁹ ..55 mg l ⁻¹)	48 h	N/A	yes	egestion within days	N/A	Hu et al. (2016)
					carboxyl			1	4.55·10 ⁷ items per mouse (0.025 mg per mouse)						
	transgenic mice	in vivo	PS	N/A	sulfate	fluorescence	ms	4	4.55·10 ⁷ items per mouse (1.6 mg per mouse)	28 d	N/A	yes	N/A	no	Stock et al. (2019)
					sulfate			10	1.49·10 ⁶ items per mouse (0.8 mg per mouse)						
Rodentia	mice	in vivo	PS	N/A	N/A	fluorescence	ms	5	0.1..1 mg l ⁻¹ (food)	42 d	N/A	yes	N/A	≥0.1 mg l ⁻¹ : microbiome, metabolic dysfunction	Jin et al. (2019)
	mice	in vivo	PS	N/A	N/A	N/A	ms	0.5..50	0.1..1 mg l ⁻¹ (food)	35 d	N/A	N/A	N/A	≥0.1 mg l ⁻¹ : microbiome, metabolic dysfunction	Lu et al. (2018)
	<i>Mus musculus</i>	in vivo	PS	N/A	N/A	fluorescence	ms	5..20	200 mg l ⁻¹ (food)	28 d	N/A	yes	8x, 8±5 and 0.71±0.14 mg kg ⁻¹ body weight	≥1 mg l ⁻¹ : body weight ↓	Yang et al. (2019b)

489 4 Synthesis

490 4.1 Summarized observations

491 Our systematic search comprised recent research on the interaction of soil organisms with
492 MP, but also studies with focus on feeding experiments, that are published much earlier than
493 the awareness on plastic in the environment appeared. The numerous studies found with
494 focus on the ingestion of MPs consistently showed the active uptake by diverse soil
495 organisms with few exceptions spread over the whole branch of invertebrates. In addition,
496 also studies on adverse effects caused by the intake of MP contaminated food (e.g. of food
497 pellets by dung beetles) imply the ingestion into the test organism. Distinct size preferences
498 are [12] observed in dung beetles, nematodes, rotifers and ants showing that mainly particles
499 are ingested, that are small enough to enter the gastrointestinal tract. In contrast, active
500 comminution by gnawing on larger particles was tested only for a few taxa and confirmed for
501 woodlice, termites and mealworms, and in the case of earthworms only after initial
502 weathering.

503 After the ingestion, MP is [12] translocated actively until excretion or death of the transporting
504 organism, which was only directly shown in experiments with earthworms. The passive
505 transport by attachment, dragging and pushing was [12] investigated in a few experiments with
506 earthworms, mites and springtails that partly worked without soil substrate and consistently
507 showed positive results.

508 After exposition to MP, a pattern of adverse effects can be seen: Across various taxa, altered
509 microbiomes, reduced motility, body mass, fertility and life span as well as increased oxidative
510 stress and metabolic malfunctioning occur in different combinations mainly due to μm -sized
511 MP in and above the whole known natural range of [12] concentrations. For some taxa such as
512 Nematodes, Gastropoda and Rotifera these effects appear at natural and increased MP
513 concentrations ($<100 \text{ mg kg}^{-1}$ dry soil), for Collembola and Lumbricidae at concentrations like
514 in highly contaminated sites ($\geq 1000 \text{ mg kg}^{-1}$ dry soil) and for Enchytraeidae, *Arenicola marina*
515 and in further experiments with earthworms at implausibly high values. The data show a
516 tendency, that the effects occur at lower concentrations, when the added particles are smaller.
517 Small sized particles also provide the highest surface/volume ratio and thus the highest
518 reactive surface per weight.

519 Most studies work with defined increasing MP concentrations and particle sizes in soil
520 substrates and food sources, which can be used to determine relationships between
521 environmental concentrations and adverse effects. However, the lack of information about
522 intake rates, grades of accumulation and effective prey-predator transfer leads to a gap within
523 the chain of explanation for toxic effects on the soil organisms. In some experiments, the
524 intestinal passage of MP and sizes [4] preferably retained within the gut were shown, but there
525 are no experiments that could demonstrate quantitative bioaccumulation. In contrast,

526 quantification of the retained and egested MP particle size fractions might be biased due to
527 gnawing and intestinal comminution as shown for woodlice, termites, mealworms, snails and
528 earthworms.

529 In order to improve our understanding of processes underlying adverse effects of MP on soil
530 organisms, data on ingestion rates, dwell times, biodegradation and egestion rates are
531 important bricks e.g. to reveal bioaccumulation dynamics. However, there are only a few data
532 on biodegradation (mealworms, snails, earthworms), egestion (rotifers, frogs, snails,
533 earthworms) and remaining concentrations in the body (lugworm, mice, earthworms).

534 4.2 Limitations of previous studies

535 The available studies worked with items within the full size span of micro- and nanoplastics
536 ($\leq 5000 \mu\text{m}$). [17] Approximately 72 % of the experiments used microplastic (0.1 to $5000 \mu\text{m}$),
537 only 6 % nanoplastic ($< 0.1 \mu\text{m}$), 10 % included macroplastic ($> 5000 \mu\text{m}$) and 12 % used
538 microplastic of undefined size. When $\text{MP} \geq 50 \mu\text{m}$ was applied, mainly particles and fragments
539 made of PE and PVC were used, whereas PS/latex microspheres were mainly applied for
540 sizes $\leq 10 \mu\text{m}$ ([17] Table 1). The latter are readily available, highly standardized and are mostly
541 used with fluorescent dyes and either without additional functional groups, carboxylated or,
542 more rarely, with amino or sulfate groups. However, there are indications that the spectrum of
543 particle type and shape used in experiments does not correspond to the properties of
544 particles in soils. In different natural as well as agriculturally and industrially contaminated
545 terrestrial and semisubhydric sites, fibers and fragments of PE and PP, mostly $\leq 100 \mu\text{m}$, were
546 much more abundant than PVC, PET and PS items (Claessens et al., 2011; Vianello et al.,
547 2013; Nor and Obbard, 2014; Naji et al., 2017; Zhang and Liu, 2018; Li et al., 2018a). This is
548 probably caused by high loads of MP fibers in discharged waste water and sewage sludge,
549 which is used in agricultural sites worldwide (Mahon et al., 2016; Li et al., 2018b). It is likely
550 that shape plays an important role for the ingestion of MP items. Unfortunately, we did not find
551 studies that have carried out a complete classification of sampling sites according to plastic
552 origin, size and type, that could help to evaluate differences between former experimental and
553 natural plastic composition to achieve the most realistic experimental conditions. Little
554 knowledge about the size distribution of MP in soils furthermore complicates the
555 determination of realistic concentrations for the addition of a certain particle size spectrum. All
556 reviewed studies either arbitrarily set their applied concentrations or had to base them on
557 measurements of total specific MP masses, regardless of how much of this mass is in the
558 tested size range. This may lead to a false estimation of total adverse MP concentrations.

559 In contrast to particle type and shape, the documentation of chemical properties of MP
560 samples in most of these studies is fragmentary. Some experiments explicitly mentioned that
561 the added plastic was unweathered, whereas most studies lack information about the degree
562 of aging implying that unweathered items were used. Only a few experiments involved aging
563 of MP, but without comparison to results of natural weathering (Tsunoda et al., 2010;
564 Gebhardt and Forster, 2018). That is in conflict with natural conditions, as plastic that [12]
565 remains within the soil after littering, sewage sludge application or plastic mulching shows
566 signs of weathering, e.g. modified carbonyl indices (Andrady, 2017), while unweathered soil
567 MP might be rare. In addition, Zhang et al. (2018) showed that earthworms actively
568 comminute only weathered bioplastics. [6] In experiments using PS microspheres,
569 carboxylation is often used to imitate a reduced hydrophobicity due to weathering. However,
570 according to manufacturer information microplastics only have little influence on
571 hydrophobicity.

572 Weathering of MP surfaces within soils comes along with biofilm growth and adsorption of
573 organic molecules, which could potentially affect the attractiveness or toxicity for grazers and
574 other organisms. Such coatings were applied only in a few cases (Besseling et al., 2017;
575 Angotti et al., 2018; Gebhardt and Forster, 2018), but were not documented in most studies.
576 Similarly, the type and concentration of additives such as flame retardants, anti-oxidants or
577 stabilizers often remained undocumented, with exception of fluorescent dyes, that are well
578 mentioned. The release of additives can have a harmful effect on the test organism, as shown
579 for aquatic environments (e Silva et al., 2016). Some studies on the ingestion of MP by the
580 soil mesofauna indicate that the diameter of the gastrointestinal tract is a useful upper size
581 limit for added particles, as far as the organism is unable to crush them (Heerkloß and Hlawa,
582 1995; Holter, 2000; Holter et al., 2002; Holter and Scholtz, 2005; Baer et al., 2008; Fueser et
583 al., 2019). However, using only ingestible particle sizes in their natural concentrations neglect
584 the adverse effects of plastic leachates, which can also get into the soil solution and onto the
585 mineral phase from larger particles and affect soil life.

586 The conditions of incubation differ considerably in terms of habitats and duration of exposure.
587 In most studies, the exposure ranges from a few minutes to a few days in experiments with
588 micro- and small mesofauna and hours to several weeks in experiments with large meso- and
589 macrofauna and is mainly based on excretion or reproductive cycles. Long-term studies,
590 which are indeed difficult to carry out in mesocosms, practically do not exist. However, certain
591 adverse effects might only establish themselves after long term trials, as was shown for the
592 influence of pesticides (Pelosi et al., 2015).

593 Some experiments were carried out in soil-free test environments such as liquid cultures or
594 Petri dishes with nutrient solutions or a specific food source (nematods, rotifers, mice).
595 Therefore, motivity is less restricted and feeding behavior can be altered compared to
596 cultivation within soil environments. For example, the ingestion of MP by nematodes
597 decreases in the presence of an alternative and more natural food source like bacteria, which
598 can significantly reduce the bioaccumulation and thus the effective toxicity (Kiyama et al.,
599 2012). This can lead to less consumption of MP in soil environments and an overestimation of
600 the toxicity in liquid culture experiments. Also, all laboratory feeding experiments were carried
601 out by use of only one species. The complexity of the food web in soils is thereby excluded
602 and the potential accumulation from prey to predators still unexplored.

603 4.3 Pinpoints for future research

604 Most studies reviewed in this work have a pioneering role in MP research and, thus, are
605 subject to some experimental limitations caused by an early state of knowledge. The adverse
606 effects recently found are alarming, but must be considered under the restrictions named
607 above. We propose the following points as part of a *modus operandi* for future MP research.

608 In past studies, particular adverse effects of MP were measured only for certain sizes,
609 shapes, coatings, leachates or adsorbed substances (*Tables 2 to 8*). Experimental
610 concentrations were assumed randomly or derived from cumulative concentrations of one or
611 more MP types measured in natural soils (approx. 1 to some 1000 mg kg⁻¹ dry soil),
612 regardless of size. For those specific experiments coming, the spectrum of concentrations
613 used should be adapted to the quantities of the size spectrum, that occurs within the soil. For
614 future studies on mixed contaminations, we recommend to evaluate the overall adverse
615 effects of PE, PP, PVC, PET, PU and PS to certain test organisms by use of typical MP-
616 specific spans of concentration, size and shape distribution in natural soils or food samples.
617 This previously requires well-structured data of appropriate MP type, shape and size for
618 different soils in differently contaminated areas.

619 Experiments on adverse effects should be applied within soil matrices to allow the interplay of
620 plastic, natural organic and mineral matter. The MP should be weathered, as plastic in soils
621 underlie broad environmental aging. Pre-weathering of MP should therefor not only be
622 performed in climate chambers (e.g. following DIN EN ISO 4892-2/3), but also include
623 subsequent leaching and equilibration of additives or coatings within the soil matrix before the
624 main experiment. Furthermore, the experimental design may consider coatings with biofilms
625 or attractants and even particle color to regulate the preference of the test organisms.

626 Most detailed information about ingestion are available for dung beetles, nematods and
627 earthworms, data on adverse effects on nematods, earthworms, lugworms and collembola.
628 Future experiments should focus on **a larger variety of ecologically relevant taxa like**
629 **Coleoptera, Formicidae, Acari, Oniscidea, Collembola, Lumbricidae, Enchytraeidae,**
630 **Nematoda and Gastropoda.** The studies are recommended to conduct with emphasis on
631 uptake, accumulation and key adverse effects like on survival rate, motility, growth and fertility
632 as well as on the stability of the intestinal microbiome. Further studies with more than one test
633 organism are important to foster our understanding of MP within certain food chains. Also
634 long-term experiments might reveal adverse effects, which evolve slowly within populations.
635 This may enable the **assessment** of the distribution and effects of MP within the food web and
636 the resulting long-term impact on soil ecosystems.

637 5 Conclusion

638 Our review of 77 studies on the impact of microplastic on the soil fauna shows a [10]
639 considerable diversity and distribution of adverse effects within the soil tree of life. However,
640 these effects have to be considered carefully, as many experiments [12] did not use plastic
641 matching properties within natural soils and found adverse effects only at concentrations like
642 in highly contaminated soils or above. To elucidate effective concentrations [11] and properties
643 for short and long-term effects on soil faunal health, the most exact reproduction of plastic
644 properties within the soil matrix and natural living conditions of the test organisms is
645 necessary [11] together with a better knowledge on common concentrations and size
646 distributions of soil microplastic. For future experiments we therefore recommend to choose
647 compositions of type, shape, size, [11] concentration, grade of weathering, leachability and
648 coating with biofilms and other organic matter as expected in the habitat to be examined.
649 Furthermore, coming studies should include long-term exposure and food chain experiments
650 to get a better look at the effect of even smaller MP concentrations and their enrichment
651 within the food web. This may give us a better way of assessing the impact of global
652 microplastic contamination on e.g. soil biodiversity, soil carbon cycles and soil quality.

653 **Author contribution**

654 Frederick Büks developed the review concept, collected data and prepared the manuscript
655 except for earthworms. Nicolette Loes van Schaik did all the work on earthworms. Martin
656 Kaupenjohann supervised the study **by participating in structural discussions on the idea**
657 **and concept of the paper as well as the final corrections.**

658 The authors declare that they have no conflict of interest.

659

660

661 **Acknowledgement**

662 Many thanks to Ivica Letunic, who kindly gave us access to phyloT and made our work in face
663 of New Year much easier. I will not forget to invite you for a coffee.

664

665

666 **Competing interests**

667 The authors declare that they have no conflict of interest.

668 **References**

- 669 Allasino M. L., Marrero H. J., Dorado J. and Torretta J. P.: Scientific note: first global report of
670 a bee nest built only with plastic, *Apidologie*, 50, 230-233, doi:10.1007/s13592-019-00635-6,
671 2019.
- 672 Andrady A. L.: The plastic in microplastics: a review, *Marine Pollution Bulletin*, 119, 12-22,
673 doi:10.1016/j.marpolbul.2017.01.082, 2017.
- 674 Angotti M. A., Rabello A., Santiago G. and Ribas C.: Seed removal by ants in Brazilian
675 savanna: optimizing fieldwork, *Sociobiology*, 65, 155-161,
676 doi:10.13102/sociobiology.v65i2.1938, 2018.
- 677 Avery S. V., Harwood J. L. and Lloyd D.: Quantification and Characterization of Phagocytosis
678 in the Soil Amoeba *Acanthamoeba castellanii* by Flow Cytometry, *Appl. Environ. Microbiol.*,
679 61, 1124-1132, doi:N/A, 1995.
- 680 Baer A., Langdon C., Mills S., Schulz C. and Hamre K.: Particle size preference, gut filling and
681 evacuation rates of the rotifer *Brachionus* "Cayman" using polystyrene latex beads,
682 *Aquaculture*, 282, 75-82, doi:10.1016/j.aquaculture.2008.06.020, 2008.
- 683 Bertling J., Bertling R. and Hamann L.: Kunststoffe in Der Umwelt: Mikro- und Makroplastik.
684 Ursachen, Mengen, Umweltschicksale, Wirkungen, Lösungsansätze, Empfehlungen,
685 doi:10.24406/UMSICHT-N-497117, 2018.
- 686 Besseling E., Foekema E. M., van den Heuvel-Greve M. J. and Koelmans A. A.: The effect of
687 microplastic on the uptake of chemicals by the lugworm *Arenicola marina* (L.) under
688 environmentally relevant exposure conditions, *Environmental Science & Technology*, 51,
689 8795-8804, doi:10.1021/acs.est.7b02286, 2017.
- 690 Besseling E., Wegner A., Foekema E. M., Van Den Heuvel-Greve M. J. and Koelmans A. A.:
691 Effects of microplastic on fitness and PCB bioaccumulation by the lugworm *Arenicola marina*
692 (L.), *Environmental Science & Technology*, 47, 593-600, doi:10.1021/es302763x, 2012.
- 693 Boots B., Russell C. W. and Green D. S.: Effects of Microplastics in Soil Ecosystems: Above
694 and Below Ground, *Environmental Science & Technology*, 53, 11496-11506,
695 doi:10.1021/acs.est.9b03304, 2019.
- 696 Bowers B. and Olszewski T. E.: *Acanthamoeba* discriminates internally between digestible
697 and indigestible particles, *The Journal of Cell Biology*, 97, 317-322, doi:10.1083/jcb.97.2.317,
698 1983.
- 699 Brandon A. M., Gao S.-H., Tian R., Ning D., Yang S.-S., Zhou J., Wu W.-M. and Criddle C. S.:
700 Biodegradation of polyethylene and plastic mixtures in mealworms (larvae of *Tenebrio molitor*)

701 and effects on the gut microbiome, *Environmental Science & Technology*, 52, 6526-6533,
 702 doi:10.1021/acs.est.8b02301, 2018.

703 Cao D., Wang X., Luo X., Liu G. and Zheng H.: Effects of polystyrene microplastics on the
 704 fitness of earthworms in an agricultural soil, In: *IOP Conference Series: Earth and*
 705 *Environmental Science*, doi:10.1088/1755-1315/61/1/012148, 2017.

706 Chae Y. and An Y.-J.: Current research trends on plastic pollution and ecological impacts on
 707 the soil ecosystem: A review, *Environmental Pollution*, 240, 387-395,
 708 doi:10.1016/j.envpol.2018.05.008, 2018.

709 Chae Y., Kim D., Kim S. W. and An Y.-J.: Trophic transfer and individual impact of nano-sized
 710 polystyrene in a four-species freshwater food chain, *Scientific Reports*, 8, 284,
 711 doi:10.1038/s41598-017-18849-y, 2018.

712 Chen Y., Liu X., Leng Y. and Wang J.: Defense responses in earthworms (*Eisenia fetida*)
 713 exposed to low-density polyethylene microplastics in soils, *Ecotoxicology and Environmental*
 714 *Safety*, 187, 109788, doi:10.1016/j.ecoenv.2019.109788, 2020.

715 Claessens M., De Meester S., Van Landuyt L., De Clerck K. and Janssen C. R.: Occurrence
 716 and distribution of microplastics in marine sediments along the Belgian coast, *Marine*
 717 *Pollution Bulletin*, 62, 2199-2204, doi:10.1016/j.marpolbul.2011.06.030, 2011.

718 Darwin C.: The formation of vegetable mould through the action of worms, with observations
 719 on their habits, John Murray, London, 1881.

720 De Bruyn L. and Conacher A. J.: The role of termites and ants in soil modification-a review,
 721 *Soil Research*, 28, 55-93, doi:10.1071/SR9900055, 1990.

722 Dong S., Qu M., Rui Q. and Wang D.: Combinational effect of titanium dioxide nanoparticles
 723 and nanopolystyrene particles at environmentally relevant concentrations on nematode
 724 *Caenorhabditis elegans*, *Ecotoxicology and Environmental Safety*, 161, 444-450, doi:10.1016/
 725 j.ecoenv.2018.06.021, 2018.

726 Elloway E. A., Bird R. A., Hewitt C. J., Kelly S. L. and Smith S. N.: Characterization of
 727 *Acanthamoeba*-microsphere association by multiparameter flow cytometry and confocal
 728 microscopy, *Cytometry Part A*, 69, 266-272, doi:10.1002/cyto.a.20210, 2006.

729 Fenchel T.: Suspension feeding in ciliated protozoa: functional response and particle size
 730 selection, *Microbial Ecology*, 6, 1-11, doi:10.1007/BF02020370, 1980.

731 Ferris H.: Contribution of nematodes to the structure and function of the soil food web, *Journal*
 732 *of Nematology*, 42, 63, doi:N/A, 2010.

733 Frey F.: The suitability of *Acrobeloides buetschlii* for nematological experiments,
 734 *Nematologica*, 17, 474-477, doi:N/A, 1971.

735 Fründ H.-C., Graefe U. and Tischer S.: Earthworms as bioindicators of soil quality, In: Biology
736 of earthworms, Springer, doi:10.1007/978-3-642-14636-7_16, 2011.

737 Fueser H., Mueller M.-T., Weiss L., Höss S. and Traunspurger W.: Ingestion of microplastics
738 by nematodes depends on feeding strategy and buccal cavity size, Environmental Pollution,
739 255, 113227, doi:10.1016/j.envpol.2019.113227, 2019.

740 Fuller S. and Gautam A.: A procedure for measuring microplastics using pressurized fluid
741 extraction, Environmental Science & Technology, 50, 5774-5780,
742 doi:10.1021/acs.est.6b00816, 2016.

743 Garcés-Ordóñez O., Castillo-Olaya V. A., Granados-Briceño A. F., Garcéa L. M. B. and Díaz
744 L. F. E.: Marine litter and microplastic pollution on mangrove soils of the Ciénaga Grande de
745 Santa Marta, Colombian Caribbean, Marine Pollution Bulletin, 145, 455-462,
746 doi:10.1016/j.marpolbul.2019.06.058, 2019.

747 Gebhardt C. and Forster S.: Size-selective feeding of *Arenicola marina* promotes long-term
748 burial of microplastic particles in marine sediments, Environmental Pollution, 242, 1777-1786,
749 doi:10.1016/j.envpol.2018.07.090, 2018.

750 van Gestel C. A., Loureiro S. and others: Terrestrial isopods as model organisms in soil
751 ecotoxicology: a review, ZooKeys, 127, doi:10.3897/zookeys.801.21970, 2018.

752 van Gestel C. A. and Selonen S.: Ecotoxicological effects of microplastics in soil: Comments
753 on the paper by Zhu et al.(2018) 'Exposure of soil collembolans to microplastics perturbs their
754 gut microbiota and alters their isotopic composition', Soil Biology & Biochemistry 116,
755 302-310, Soil Biology and Biochemistry, 124, 116-117, doi:10.1016/j.soilbio.2018.05.032,
756 2018.

757 Geyer R., Jambeck J. R. and Law K. L.: Production, use, and fate of all plastics ever made,
758 Science Advances, 3, e1700782, doi:10.1126/sciadv.1700782, 2017.

759 Glancey B. M., Vander Meer R., Glover A., Lofgren C. and Vinson S.: Filtration of
760 microparticles from liquids ingested by the red imported fire ant *Solenopsis invicta* Buren,
761 Insectes Sociaux, 28, 395-401, doi:10.1007/BF02224196, 1981.

762 Green D. S., Boots B., Sigwart J., Jiang S. and Rocha C.: Effects of conventional and
763 biodegradable microplastics on a marine ecosystem engineer (*Arenicola marina*) and
764 sediment nutrient cycling, Environmental Pollution, 208, 426-434,
765 doi:10.1016/j.envpol.2015.10.010, 2016.

766 Gulvik M.: Mites (Acari) as indicators of soil biodiversity and land use monitoring: a review,
767 Polish Journal of Ecology, 55, 415-440, doi:N/A, 2007.

768 Gusmão F., Di Domenico M., Amaral A. C. Z., Martínez A., Gonzalez B. C., Worsaae K., do
769 Sul J. A. I. and da Cunha Lana P.: In situ ingestion of microfibres by meiofauna from sandy
770 beaches, *Environmental Pollution*, 216, 584-590, doi:10.1016/j.envpol.2016.06.015, 2016.

771 Gutow L., Bartl K., Saborowski R. and Beermann J.: Gastropod pedal mucus retains
772 microplastics and promotes the uptake of particles by marine periwinkles, *Environmental*
773 *Pollution*, 246, 688-696, doi:10.1016/j.envpol.2018.12.097, 2019.

774 Hassall M. and Rushton S.: Feeding behaviour of terrestrial isopods in relation to plant
775 defences and microbial activity, In: *Symposia of the Zoological Society of London*, doi:N/A,
776 1984.

777 Hebrard J. J., Maloiy G. M. and Alliangana D. M.: Notes on the habitat and diet of *Afrocaecilia*
778 *taimana* (Amphibia: Gymnophiona), *Journal of Herpetology*, 513-515, doi:10.2307/1565136,
779 1992.

780 Heerkloß R. and Hlawa S.: Feeding biology of two brachionid rotifers: *Brachionus*
781 *quadridentatus* and *Brachionus plicatilis*, *Hydrobiologia*, 313, 219-221,
782 doi:10.1007/BF00025954, 1995.

783 Hodson M. E., Duffus-Hodson C. A., Clark A., Prendergast-Miller M. T. and Thorpe K. L.:
784 Plastic bag derived-microplastics as a vector for metal exposure in terrestrial invertebrates,
785 *Environmental Science & Technology*, 51, 4714-4721, doi:10.1021/acs.est.7b00635, 2017.

786 Holter P.: Particle feeding in *Aphodius* dung beetles (Scarabaeidae): old hypotheses and new
787 experimental evidence, *Functional Ecology*, 631-637, doi:N/A, 2000.

788 Holter P. and Scholtz C.: Are ball-rolling (Scarabaeini, Gymnopleurini, Sisyphini) and
789 tunnelling scarabaeine dung beetles equally choosy about the size of ingested dung
790 particles?, *Ecological Entomology*, 30, 700-705, doi:10.1111/j.0307-6946.2005.00746.x, 2005.

791 Holter P., Scholtz C. and Wardhaugh K.: Dung feeding in adult scarabaeines (tunnellers and
792 endocoprids): even large dung beetles eat small particles, *Ecological Entomology*, 27, 169-
793 176, doi:10.1046/j.1365-2311.2002.00399.x, 2002.

794 Hopkin S. P.: *Biology of the springtails (Insecta: Collembola)*, OUP Oxford, 1997.

795 Hu L., Su L., Xue Y., Mu J., Zhu J., Xu J. and Shi H.: Uptake, accumulation and elimination of
796 polystyrene microspheres in tadpoles of *Xenopus tropicalis*, *Chemosphere*, 164, 611-617,
797 doi:https://doi.org/10.1016/j.chemosphere.2016.09.002, 2016.

798 Huang, Y., Liu, Q., Jia, W., Yan, C. and Wang, J.: Agricultural plastic mulching as a source of
799 microplastics in the terrestrial environment. *Environmental Pollution*, 260, 114096,
800 doi: 10.1016/j.envpol.2020.114096, 2020.

801 Huerta Lwanga E., Gertsen H., Gooren H., Peters P., Salánki T., van der Ploeg M., Besseling
802 E., Koelmans A. A. and Geissen V.: Microplastics in the terrestrial ecosystem: implications for
803 *Lumbricus terrestris* (Oligochaeta, Lumbricidae), *Environmental Science & Technology*, 50,
804 2685-2691, doi:10.1021/acs.est.5b05478, 2016.

805 Huerta Lwanga E., Gertsen H., Gooren H., Peters P., Salánki T., van der Ploeg M., Besseling
806 E., Koelmans A. A. and Geissen V.: Incorporation of microplastics from litter into burrows of
807 *Lumbricus terrestris*, *Environmental Pollution*, 220, 523-531,
808 doi:10.1016/j.envpol.2016.09.096, 2017b.

809 Huerta Lwanga E., Thapa B., Yang X., Gertsen H., Salánki T., Geissen V. and Garbeva P.:
810 Decay of low-density polyethylene by bacteria extracted from earthworm's guts: A potential for
811 soil restoration, *Science of the Total Environment*, 624, 753-757,
812 doi:10.1016/j.scitotenv.2017.12.144, 2018.

813 Huerta Lwanga E., Vega J. M., Quej V. K., de los Angeles Chi J., del Cid L. S., Chi C., Segura
814 G. E., Gertsen H., Salánki T., van der Ploeg M. and others: Field evidence for transfer of
815 plastic debris along a terrestrial food chain, *Scientific Reports*, 7, 14071, doi:10.1038/s41598-
816 017-14588-2, 2017a.

817 Hughes L. and Westoby M.: Effect of diaspore characteristics on removal of seeds adapted
818 for dispersal by ants, *Ecology*, 73, 1300-1312, doi:10.2307/1940677, 1992.

819 Imhof H. K. and Laforsch C.: Hazardous or not - Are adult and juvenile individuals of
820 *Potamopyrgus antipodarum* affected by non-buoyant microplastic particles?, *Environmental*
821 *Pollution*, 218, 383-391, doi:10.1016/j.envpol.2016.07.017, 2016.

822 Jambeck J. R., Geyer R., Wilcox C., Siegler T. R., Perryman M., Andrady A., Narayan R. and
823 Law K. L.: Plastic waste inputs from land into the ocean, *Science*, 347, 768-771, doi:10.1126/
824 science.1260352, 2015.

825 Jeong C.-B., Won E.-J., Kang H.-M., Lee M.-C., Hwang D.-S., Hwang U.-K., Zhou B., Souissi
826 S., Lee S.-J. and Lee J.-S.: Microplastic size-dependent toxicity, oxidative stress induction,
827 and p-JNK and p-p38 activation in the monogonont rotifer (*Brachionus koreanus*),
828 *Environmental Science & Technology*, 50, 8849-8857, doi:10.1021/acs.est.6b01441., 2016.

829 Jin Y., Lu L., Tu W., Luo T. and Fu Z.: Impacts of polystyrene microplastic on the gut barrier,
830 microbiota and metabolism of mice, *Science of the Total Environment*, 649, 308-317,
831 doi:10.1016/j.scitotenv.2018.08.353, 2019.

832 Jonsson P. R.: Particle size selection, feeding rates and growth dynamics of marine
833 planktonic oligotrichous ciliates (Ciliophora: Oligotrichina), *Mar Ecol Prog Ser*, 33, 265-277,
834 doi:N/A, 1986.

835 Jouquet P., Dauber J., Lagerlöf J., Lavelle P. and Lepage M.: Soil invertebrates as ecosystem
836 engineers: intended and accidental effects on soil and feedback loops, *Applied Soil Ecology*,
837 32, 153-164, doi:10.1016/j.apsoil.2005.07.004, 2006.

838 Ju H., Zhu D. and Qiao M.: Effects of polyethylene microplastics on the gut microbial
839 community, reproduction and avoidance behaviors of the soil springtail, *Folsomia candida*,
840 *Environmental Pollution*, 247, 890-897, doi:10.1016/j.envpol.2019.01.097, 2019.

841 Judy J. D., Williams M., Gregg A., Oliver D., Kumar A., Kookana R. and Kirby J. K.:
842 Microplastics in municipal mixed-waste organic outputs induce minimal short to long-term
843 toxicity in key terrestrial biota, *Environmental Pollution*, 252, 522-531,
844 doi:10.1016/j.envpol.2019.05.027, 2019.

845 Kale S. K., Deshmukh A. G., Dudhare M. S. and Patil V. B.: Microbial degradation of plastic: a
846 review, *Journal of Biochemical Technology*, 6, 952-961, doi:N/A, 2015.

847 Kim H. M., Lee D.-K., Long N. P., Kwon S. W. and Park J. H.: Uptake of nanopolystyrene
848 particles induces distinct metabolic profiles and toxic effects in *Caenorhabditis elegans*,
849 *Environmental Pollution*, 246, 578-586, doi:10.1016/j.envpol.2018.12.043, 2019.

850 Kim S. W. and An Y.-J.: Soil microplastics inhibit the movement of springtail species,
851 *Environment International*, 126, 699-706, doi:10.1016/j.envint.2019.02.067, 2019.

852 Kiyama Y., Miyahara K. and Ohshima Y.: Active uptake of artificial particles in the nematode
853 *Caenorhabditis elegans*, *Journal of Experimental Biology*, 215, 1178-1183,
854 doi:10.1242/jeb.067199, 2012.

855 Kokalj A. J., Horvat P., Skalar T. and Kržan A.: Plastic bag and facial cleanser derived
856 microplastic do not affect feeding behaviour and energy reserves of terrestrial isopods,
857 *Science of the Total Environment*, 615, 761-766, doi:10.1016/j.scitotenv.2017.10.020, 2018.

858 Korb J.: The ecology of social evolution in termites, In: *Ecology of social evolution*, Springer,
859 doi:10.1007/978-3-540-75957-7_7, 2008.

860 Kumar A., Pandey A. K., Singh S. S., Shanker R. and Dhawan A.: Cellular uptake and
861 mutagenic potential of metal oxide nanoparticles in bacterial cells, *Chemosphere*, 83, 1124-
862 1132, doi:10.1016/j.chemosphere.2011.01.025, 2011.

863 Lahive E., Walton A., Horton A. A., Spurgeon D. J. and Svendsen C.: Microplastic particles
864 reduce reproduction in the terrestrial worm *Enchytraeus crypticus* in a soil exposure,
865 *Environmental Pollution*, 255, 113174, doi:10.1016/j.envpol.2019.113174, 2019.

866 Lavelle P., Decaëns T., Aubert M., Barot S., Blouin M., Bureau F., Margerie P., Mora P. and
867 Rossi J.-P.: Soil invertebrates and ecosystem services, *European Journal of Soil Biology*, 42,
868 S3-S15, doi:10.1016/j.ejsobi.2006.10.002, 2006.

869 Lavin D. P., Fredrickson A. and Srien F.: Flow cytometric measurement of rates of particle
 870 uptake from dilute suspensions by a ciliated protozoan, *Cytometry*, 11, 875-882, doi:10.1002/
 871 cyto.990110804, 1990.

872 Lei L., Liu M., Song Y., Lu S., Hu J., Cao C., Xie B., Shi H. and He D.: Polystyrene (nano)
 873 microplastics cause size-dependent neurotoxicity, oxidative damage and other adverse
 874 effects in *Caenorhabditis elegans*, *Environmental Science: Nano*, 5, 2009-2020, doi:10.1039/
 875 C9EN00473D, 2018a.

876 Lei L., Wu S., Lu S., Liu M., Song Y., Fu Z., Shi H., Raley-Susman K. M. and He D.:
 877 Microplastic particles cause intestinal damage and other adverse effects in zebrafish *Danio*
 878 *rerio* and nematode *Caenorhabditis elegans*, *Science of the Total Environment*, 619, 1-8,
 879 doi:10.1016/j.scitotenv.2017.11.103, 2018b.

880 Lenz M., Creffield J. W., Evans T. A., Kard B., Vongkaluang C., Sornnuwat Y., Lee C.-Y.,
 881 Yoshimura T. and Tsunoda K.: Resistance of polyamide and polyethylene cable sheathings to
 882 termites in Australia, Thailand, USA, Malaysia and Japan: a comparison of four field
 883 assessment methods, *International Biodeterioration & Biodegradation*, 66, 53-62,
 884 doi:10.1016/j.ibiod.2011.11.001, 2012.

885 Li J., Zhang H., Zhang K., Yang R., Li R. and Li Y.: Characterization, source, and retention of
 886 microplastic in sandy beaches and mangrove wetlands of the Qinzhou Bay, China, *Marine*
 887 *Pollution Bulletin*, 136, 401-406, doi:10.1016/j.marpolbul.2018.09.025, 2018a.

888 Li L., Zhou Q., Yin N., Tu C. and Luo Y.: Uptake and accumulation of microplastics in an
 889 edible plant, *Chinese Science Bulletin*, 64, 928-934, doi:10.1360/N972018-00845, 2019.

890 Li X., Chen L., Mei Q., Dong B., Dai X., Ding G. and Zeng E. Y.: Microplastics in sewage
 891 sludge from the wastewater treatment plants in China, *Water Research*, 142, 75-85,
 892 doi:10.1016/j.watres.2018.05.034, 2018b.

893 Lobelle D. and Cunliffe M.: Early microbial biofilm formation on marine plastic debris, *Marine*
 894 *Pollution Bulletin*, 62, 197-200, doi:10.1016/j.marpolbul.2010.10.013, 2011.

895 Lu L., Wan Z., Luo T., Fu Z. and Jin Y.: Polystyrene microplastics induce gut microbiota
 896 dysbiosis and hepatic lipid metabolism disorder in mice, *Science of the Total Environment*,
 897 631, 449-458, doi:10.1016/j.scitotenv.2018.03.051, 2018.

898 Maaß S., Daphi D., Lehmann A. and Rillig M. C.: Transport of microplastics by two
 899 collembolan species, *Environmental Pollution*, 225, 456-459,
 900 doi:10.1016/j.envpol.2017.03.009, 2017.

901 Mahon A. M., O'Connell B., Healy M. G., O'Connor I., Officer R., Nash R. and Morrison L.:
 902 Microplastics in sewage sludge: effects of treatment, *Environmental Science & Technology*,
 903 51, 810-818, doi:10.1021/acs.est.6b04048, 2016.

904 Monge-Nájera J.: Ecological biogeography in the Phylum Onychophora, *Biogeographica*, 70,
905 111-123, doi:N/A, 1994.

906 Naji A., Esmaili Z., Mason S. A. and Vethaak A. D.: The occurrence of microplastic
907 contamination in littoral sediments of the Persian Gulf, Iran, *Environmental Science and*
908 *Pollution Research*, 24, 20459-20468, doi:10.1007/s11356-017-9587-z, 2017.

909 Nika L., Gibson T., Konkus R. and Karp X.: Fluorescent beads are a versatile tool for staging
910 *Caenorhabditis elegans* in different life histories, *G3: Genes, Genomes, Genetics*, 6, 1923-
911 1933, doi:10.1534/g3.116.030163, 2016.

912 Nizzetto L., Futter M. and Langaas S.: Are agricultural soils dumps for microplastics of urban
913 origin?, *Environmental Science & Technology*, doi:10.1021/acs.est.6b04140, 2016.

914 Nor N. H. M. and Obbard J. P.: Microplastics in Singapore's coastal mangrove ecosystems,
915 *Marine Pollution Bulletin*, 79, 278-283, doi:10.1016/j.marpolbul.2013.11.025, 2014.

916 Oliveira M., Ameixa O. M. and Soares A. M.: Are ecosystem services provided by insects"
917 bugged" by micro (nano) plastics?, *TrAC Trends in Analytical Chemistry*,
918 doi:10.1016/j.trac.2019.02.018, 2019.

919 Parkyn J. and Newell D. A.: Australian land snails: a review of ecological research and
920 conservation approaches, *Molluscan Research*, 33, 116-129,
921 doi:10.1080/13235818.2013.782793, 2013.

922 Pass G., Szucsich N. U. and others: 100 years of research on the Protura: many secrets still
923 retained, *Soil Organisms*, 83, 309-334, doi:N/A, 2011.

924 Pelosi C., Bertrand M., Thénard J. and Mougin C.: Earthworms in a 15 years agricultural trial,
925 *Applied Soil Ecology*, 88, 1-8, doi:10.1016/j.apsoil.2014.12.004, 2015.

926 Peng B.-Y., Su Y., Chen Z., Chen J., Zhou X., Benbow M. E., Criddle C., Wu W.-M. and Zhang
927 Y.: Biodegradation of Polystyrene by Dark (*Tenebrio obscurus*) and Yellow (*Tenebrio molitor*)
928 Mealworms (Coleoptera: Tenebrionidae), *Environmental Science & Technology*,
929 doi:10.1021/acs.est.8b06963, 2019.

930 Petersen H.: A review of collembolan ecology in ecosystem context, *Acta Zoologica Fennica*,
931 195, 111-118, doi:N/A, 1994.

932 Phillips H. R., Guerra C. A., Bartz M. L., Briones M. J., Brown G., Ferlian O., Gongalsk K.,
933 Krebs J., Orgiazzi A., Schwarz B. and others: Global distribution of earthworm diversity,
934 *bioRxiv*, 587394, doi:10.1126/science.aax4851, 2019.

935 Prendergast-Miller M. T., Katsiamides A., Abbass M., Sturzenbaum S. R., Thorpe K. L. and
936 Hodson M. E.: Polyester-derived microfibre impacts on the soil-dwelling earthworm *Lumbricus*
937 *terrestris*, *Environmental Pollution*, 251, 453-459, doi:10.1016/j.envpol.2019.05.037, 2019.

938 Pulleman M., Creamer R., Hamer U., Helder J., Pelosi C., Peres G. and Rutgers M.: Soil
 939 biodiversity, biological indicators and soil ecosystem services—an overview of European
 940 approaches, *Current Opinion in Environmental Sustainability*, 4, 529-538,
 941 doi:10.1016/j.cosust.2012.10.009, 2012.

942 Qu M., Kong Y., Yuan Y. and Wang D.: Neuronal damage induced by nanopolystyrene
 943 particles in nematode *Caenorhabditis elegans*, *Environmental Science: Nano*,
 944 doi:10.1039/C9EN00473D, 2019a.

945 Qu M., Nida A., Kong Y., Du H., Xiao G. and Wang D.: Nanopolystyrene at predicted
 946 environmental concentration enhances microcystin-LR toxicity by inducing intestinal damage
 947 in *Caenorhabditis elegans*, *Ecotoxicology and Environmental Safety*, 183, 109568,
 948 doi:10.1016/j.ecoenv.2019.109568, 2019b.

949 Qu M., Qiu Y., Kong Y. and Wang D.: Amino modification enhances reproductive toxicity of
 950 nanopolystyrene on gonad development and reproductive capacity in nematode
 951 *Caenorhabditis elegans*, *Environmental Pollution*, 254, 112978,
 952 doi:10.1016/j.envpol.2019.112978, 2019c.

953 Reddy M. S., Basha S., Adimurthy S. and Ramachandraiah G.: Description of the small
 954 plastics fragments in marine sediments along the Alang-Sosiya ship-breaking yard, India,
 955 *Estuarine, Coastal and Shelf Science*, 68, 656-660, doi:10.1016/j.ecss.2006.03.018, 2006.

956 Rezaei M., Riksen M. J., Sirjani E., Sameni A. and Geissen V.: Wind erosion as a driver for
 957 transport of light density microplastics, *Science of The Total Environment*, 669, 273-281,
 958 doi:10.1016/j.scitotenv.2019.02.382, 2019.

959 Rillig M. C.: Microplastic in terrestrial ecosystems and the soil?, *Environmental Science &*
 960 *Technology*, 6453–6454, doi:10.1021/es302011r, 2012.

961 Rillig M. C. and Bonkowski M.: Microplastic and soil protists: a call for research,
 962 *Environmental Pollution*, 241, 1128-1131, doi:10.1016/j.envpol.2018.04.147, 2018.

963 Rillig M. C., Lehmann A., de Souza Machado A. A. and Yang G.: Microplastic effects on
 964 plants, *New Phytologist*, doi:10.1111/nph.15794, 2019.

965 Rillig M. C., Ziersch L. and Hempel S.: Microplastic transport in soil by earthworms, *Scientific*
 966 *Reports*, 7, 1362, doi:10.1038/s41598-017-01594-7, 2017.

967 Robins R. and Robins A.: The antics of ants: ants as agents of bioturbation in a midden
 968 deposit in south-east Queensland, *Environmental Archaeology*, 16, 151-161,
 969 doi:10.1179/174963111X13110803261010, 2011.

970 Rodríguez-Seijo A., da Costa J. P., Rocha-Santos T., Duarte A. C. and Pereira R.: Oxidative
 971 stress, energy metabolism and molecular responses of earthworms (*Eisenia fetida*) exposed

972 to low-density polyethylene microplastics, *Environmental Science and Pollution Research*, 25,
 973 33599-33610, doi:10.1007/s11356-018-3317-z, 2018.

974 Rodriguez-Seijo A., Lourenço J., Rocha-Santos T., Da Costa J., Duarte A., Vala H. and
 975 Pereira R.: Histopathological and molecular effects of microplastics in *Eisenia andrei* Bouché,
 976 *Environmental Pollution*, 220, 495-503, doi:10.1016/j.envpol.2016.09.092, 2017.

977 Rodríguez-Seijo A., Santos B., da Silva E. F., Cachada A. and Pereira R.: Low-density
 978 polyethylene microplastics as a source and carriers of agrochemicals to soil and earthworms,
 979 *Environmental Chemistry*, 16, 8-17, doi:10.1071/EN18162, 2019.

980 Sánchez-Bayo F. and Wyckhuys K. A.: Worldwide decline of the entomofauna: A review of its
 981 drivers, *Biological Conservation*, 232, 8-27, doi:10.1016/j.biocon.2019.01.020, 2019.

982 Scharf A., Piechulek A. and von Mikecz A.: Effect of nanoparticles on the biochemical and
 983 behavioral aging phenotype of the nematode *Caenorhabditis elegans*, *Acs Nano*, 7, 10695-
 984 10703, doi:10.1021/nn403443r, 2013.

985 e Silva P. P. G., Nobre C. R., Resaffe P., Pereira C. D. S. and Gusmão F.: Leachate from
 986 microplastics impairs larval development in brown mussels, *Water Research*, 106, 364-370,
 987 doi:10.1016/j.watres.2016.10.016, 2016.

988 Song Y., Cao C., Qiu R., Hu J., Liu M., Lu S., Shi H., Raley-Susman K. M. and He D.: Uptake
 989 and adverse effects of polyethylene terephthalate microplastics fibers on terrestrial snails
 990 (*Achatina fulica*) after soil exposure, *Environmental Pollution*, 250, 447-455,
 991 doi:10.1016/j.envpol.2019.04.066, 2019.

992 Stamatiadis S. and Dindal D.: Coprophilous mite communities as affected by concentration of
 993 plastic and glass particles, *Experimental & Applied Acarology*, 8, 1-12,
 994 doi:10.1007/BF01193377, 1990.

995 Stock V., Böhmert L., Lisicki E., Block R., Cara-Carmona J., Pack L. K., Selb R., Lichtenstein
 996 D., Voss L., Henderson C. J. and others: Uptake and effects of orally ingested polystyrene
 997 microplastic particles in vitro and in vivo, *Archives of Toxicology*, 1-17, doi:10.1007/s00204-
 998 019-02478-7, 2019.

999 Sun Y., Xu W., Gu Q., Chen Y., Zhou Q., Zhang L., Gu L., Huang Y., Lyu K. and Yang Z.:
 1000 Small-Sized Microplastics Negatively Affect Rotifers: Changes in the Key Life-History Traits
 1001 and Rotifer--*Phaeocystis* Population Dynamics, *Environmental Science & Technology*, 53,
 1002 9241-9251, doi:10.1021/acs.est.9b02893, 2019.

1003 Thompson R. C., Swan S. H., Moore C. J. and Vom Saal F. S.: Our plastic age,
 1004 doi:10.1098/rstb.2009.0054, 2009.

1005 Tsunoda K., Rosenblat G. and Dohi K.: Laboratory evaluation of the resistance of plastics to
 1006 the subterranean termite *Coptotermes formosanus* (Blattodea: Rhinotermitidae), *International*
 1007 *Biodeterioration & Biodegradation*, 64, 232-237, doi:10.1016/j.ibiod.2009.12.008, 2010.

1008 Vadstein O., Øie G. and Olsen Y.: Particle size dependent feeding by the rotifer *Brachionus*
 1009 *plicatilis*, *Hydrobiologia*, 255, 261-267, doi:10.1007/BF00025847, 1993.

1010 Van Cauwenberghe L., Claessens M., Vandegehuchte M. B. and Janssen C. R.: Microplastics
 1011 are taken up by mussels (*Mytilus edulis*) and lugworms (*Arenicola marina*) living in natural
 1012 habitats, *Environmental Pollution*, 199, 10-17, doi:10.1016/j.envpol.2015.01.008, 2015.

1013 Vianello A., Boldrin A., Guerriero P., Moschino V., Rella R., Sturaro A. and Da Ros L.:
 1014 Microplastic particles in sediments of Lagoon of Venice, Italy: First observations on
 1015 occurrence, spatial patterns and identification, *Estuarine, Coastal and Shelf Science*, 130, 54-
 1016 61, doi:10.1016/j.ecss.2013.03.022, 2013.

1017 Vogel G., Thilo L., Schwarz H. and Steinhart R.: Mechanism of phagocytosis in *Dictyostelium*
 1018 *discoideum*: phagocytosis is mediated by different recognition sites as disclosed by mutants
 1019 with altered phagocytotic properties, *The Journal of Cell Biology*, 86, 456-465,
 1020 doi:10.1083/jcb.86.2.456, 1980.

1021 Wang H.-T., Ding J., Xiong C., Zhu D., Li G., Jia X.-Y., Zhu Y.-G. and Xue X.-M.: Exposure to
 1022 microplastics lowers arsenic accumulation and alters gut bacterial communities of earthworm
 1023 *Metaphire californica*, *Environmental Pollution*, 251, 110-116,
 1024 doi:10.1016/j.envpol.2019.04.054, 2019b.

1025 Wang J., Coffin S., Sun C., Schlenk D. and Gan J.: Negligible effects of microplastics on
 1026 animal fitness and HOC bioaccumulation in earthworm *Eisenia fetida* in soil, *Environmental*
 1027 *Pollution*, 249, 776-784, doi:10.1016/j.envpol.2019.03.102, 2019c.

1028 Wang J., Liu X., Li Y., Powell T., Wang X., Wang G. and Zhang P.: Microplastics as
 1029 contaminants in the soil environment: A mini-review, *Science of The Total Environment*,
 1030 doi:10.1016/j.scitotenv.2019.07.209, 2019a.

1031 Warburg M. R.: Isopods and their terrestrial environment, In: *Advances in ecological research*,
 1032 Elsevier, doi:10.1016/S0065-2504(08)60246-9, 1987.

1033 Weisman R. A. and Korn E. D.: Phagocytosis of latex beads by *Acanthamoeba*. I. Biochemical
 1034 properties, *Biochemistry*, 6, 485-497, doi:10.1021/bi00854a017, 1967.

1035 Weithmann N., Möller J. N., Löder M. G., Piehl S., Laforsch C. and Freitag R.: Organic
 1036 fertilizer as a vehicle for the entry of microplastic into the environment, *Science Advances*, 4,
 1037 eaap8060, doi:10.1126/sciadv.aap8060, 2018.

1038 Westheide W. and Rieger R.: Spezielle Zoologie. 1. Einzeller und Wirbellose Tiere, Gustav
1039 Fischer, 1996.

1040 Wood C. T. and Zimmer M.: Can terrestrial isopods (Isopoda: Oniscidea) make use of
1041 biodegradable plastics?, *Applied Soil Ecology*, 77, 72-79, doi:10.1016/j.apsoil.2014.01.009,
1042 2014.

1043 Wright S. L., Rowe D., Thompson R. C. and Galloway T. S.: Microplastic ingestion decreases
1044 energy reserves in marine worms, *Current Biology*, 23, R1031-R1033,
1045 doi:10.1016/j.cub.2013.10.068, 2013.

1046 Yang S.-S., Brandon A. M., Flanagan J. C. A., Yang J., Ning D., Cai S.-Y., Fan H.-Q., Wang
1047 Z.-Y., Ren J., Benbow E. and others: Biodegradation of polystyrene wastes in yellow
1048 mealworms (larvae of *Tenebrio molitor* Linnaeus): Factors affecting biodegradation rates and
1049 the ability of polystyrene-fed larvae to complete their life cycle, *Chemosphere*, 191, 979-989,
1050 doi:10.1016/j.chemosphere.2017.10.117, 2018.

1051 Yang X., Lwanga E. H., Bemani A., Gertsen H., Salanki T., Guo X., Fu H., Xue S., Ritsema C.
1052 and Geissen V.: Biogenic transport of glyphosate in the presence of LDPE microplastics: A
1053 mesocosm experiment, *Environmental Pollution*, 245, 829-835,
1054 doi:10.1016/j.envpol.2018.11.044, 2019a.

1055 Yang Y., Yang J., Wu W.-M., Zhao J., Song Y., Gao L., Yang R. and Jiang L.: Biodegradation
1056 and mineralization of polystyrene by plastic-eating mealworms: Part 1. Chemical and physical
1057 characterization and isotopic tests, *Environmental Science & Technology*, 49, 12080-12086,
1058 doi:10.1021/acs.est.5b02661, 2015.

1059 Yang Y.-F., Chen C.-Y., Lu T.-H. and Liao C.-M.: Toxicity-based toxicokinetic/toxicodynamic
1060 assessment for bioaccumulation of polystyrene microplastics in mice, *Journal of Hazardous*
1061 *Materials*, 366, 703-713, doi:10.1016/j.jhazmat.2018.12.048, 2019b.

1062 Yeates G. W.: Nematodes in ecological webs, e LS, doi:10.1002/9780470015902.a0021913,
1063 2001.

1064 Yu M., Van Der Ploeg M., Lwanga E. H., Yang X., Zhang S., Ma X., Ritsema C. J. and
1065 Geissen V.: Leaching of microplastics by preferential flow in earthworm (*Lumbricus terrestris*)
1066 burrows, *Environmental Chemistry*, 16, 31-40, doi:10.1071/EN18161, 2019.

1067 Zhang G. and Liu Y.: The distribution of microplastics in soil aggregate fractions in
1068 southwestern China, *Science of the Total Environment*, 642, 12-20,
1069 doi:10.1016/j.scitotenv.2018.06.004, 2018.

1070 Zhang L., Sintim H. Y., Bary A. I., Hayes D. G., Wadsworth L. C., Anunciado M. B. and Flury
1071 M.: Interaction of *Lumbricus terrestris* with macroscopic polyethylene and biodegradable

1073 plastic mulch, Science of The Total Environment, 635, 1600-1608,
 1074 doi:10.1016/j.scitotenv.2018.04.054, 2018.

1075 Zhang Y., Gao T., Kang S. and Sillanpää M.: Importance of atmospheric transport for
 1076 microplastics deposited in remote areas, Environmental Pollution, 254, 112953,
 1077 doi:10.1016/j.envpol.2019.07.121, 2019.

1078 Zhang Z.-Q.: Animal biodiversity: an update of classification and diversity in 2013, Zootaxa,
 1079 3703, 5-11, doi:10.11646/zootaxa.3703.1.3, 2013.

1080 Zhao L., Qu M., Wong G. and Wang D.: Transgenerational toxicity of nanopolystyrene
 1081 particles in the range of $\mu\text{g L}^{-1}$ in the nematode *Caenorhabditis elegans*, Environmental
 1082 Science: Nano, 4, 2356-2366, doi:10.1039/C7EN00707H, 2017.

1083 Zhu B.-K., Fang Y.-M., Zhu D., Christie P., Ke X. and Zhu Y.-G.: Exposure to nanoplastics
 1084 disturbs the gut microbiome in the soil oligochaete *Enchytraeus crypticus*, Environmental
 1085 Pollution, 239, 408-415, doi:10.1016/j.envpol.2018.04.017, 2018b.

1086 Zhu D., Bi Q.-F., Xiang Q., Chen Q.-L., Christie P., Ke X., Wu L.-H. and Zhu Y.-G.: Trophic
 1087 predator-prey relationships promote transport of microplastics compared with the single
 1088 *Hypoaspis aculeifer* and *Folsomia candida*, Environmental Pollution, 235, 150-154,
 1089 doi:10.1016/j.envpol.2017.12.058, 2018a.

What do we know about how the terrestrial multicellular soil fauna reacts to microplastic?

Frederick Büks¹, Nicolette Loes van Schaik², Martin Kaupenjohann¹

¹Chair of Soil Science, Dept. of Ecology, Technische Universität Berlin, 10587 Berlin, Germany

²Chair of Ecohydrology & Landscape Evaluation, Dept. of Ecology, Technische Universität Berlin, 10587 Berlin, Germany

Correspondence to: Frederick Büks (frederick.bueks@tu-berlin.de)

Final responses to all referees plus marked-up version

Dear Referee #1

First I would like to express my sincere thanks to you for thoroughly reviewing our manuscript and for your very helpful and precise suggestions. In the following I will answer your points. Our corrections are marked-up with **yellow numbers** within the corrected manuscript at the end of this document.

Best regards,
Frederick Büks

Abstract

[1] Lines 20-21: "Most of the studies applied MP concentrations similar to amounts in slightly to very heavily polluted soils." This sentence makes the reader expect that generally, the concentrations in the experimental environments are mostly the same as expected in the environment, but is this really the case? I would suggest showing the percentage of experiments with high microplastic exposure that is not representative of most soils.

Thanks a lot for this point. We now write: "About 58 % of the studies thereby use inappropriate concentrations or units, but 42 % applied MP concentrations similar to amounts in slightly to very heavily polluted soils."

Introduction

[2] Line 53: Instead of "microbial decay", I'd suggest "processing by soil organisms", since this includes any process relevant for the generation of smaller plastic particles.

Done.

[3] Line 61: I'd suggest changing the sentence to "intensive use of plastic mulching and sewage sludge", for the former, Huang et al. (2020) show an increase in microplastic by approx. 1 order of magnitude between fields with 5 and 24 continuous years of plastic mulching.

Done.

[4] Line 95: Suggest changing "feed on" to "inadvertently ingest", otherwise it sounds like the organisms are actually able to metabolize the microplastics.

Done and reference added.

Search pattern

[5] The cut-off dates (time period that was considered) of the search should be mentioned somewhere.

Information added to this chapter (see the answer to referee #3)

[6] Figure 1: This figure shows the phylogenetic tree of edaphic fauna, rather than "edaphic tree of faunal life".

Thank you. And done.

Data collection

[7] Line 113-122: I've been having some difficulties understanding the search methodology and table 8 (table 8 should be moved at the appropriate place to become table 1).

We moved the table to line 124 and mentioned that it contains the number of found studies. All table numbers were adjusted within the text.

It would be great if the authors could re-word this, specifying:

What does it mean that some combinations would have caused too much search effort?

It means e.g. that searching for a taxon only in combination with "PET" gives results for PET bottles for cultivation and experiments and also the "use" as pets, if the search is not case sensitive. We now tried to clarify this in our text.

“Organism-plastic” is not a type-shape combination.
Oh, yes, that's right. Corrected.

What exactly does the number of studies in table 8 mean? The number of articles or single experiments (sometimes more than one taxon or plastic type is used in one article)?

The number counts for how often type-shape combinations were used in all reviewed experimental setups independently of organism.

Some articles are included that studied the uptake of macroplastics by organisms, mainly termites and ant species. It is reasonable to include these studies, but it should be mentioned more prominently, in the abstract and aims of the review, that macroplastics are included.

Where macroplastics were used in the reviewed studies, the size was explicitly mentioned in the article text, so we do not see a necessity for elaborating the text. We did add a mention of macroplastics to the abstract.

Maybe also in the synthesis, a sentence about the proportions of experiments using macro-, micro-, and nanoplastic would be a helpful piece of information.

Now mentioned in “4.2 Limitations of previous studies”

[8] Tables 1-7: What does N/A mean in the tables? In some cases I assume “not analysed” (e.g., passive transport), but in other cases it should mean “not mentioned” (e.g., aging, coating, etc.) or “not observed” (e.g., measured adverse effects). I think this needs to be specified. Usually, N/A refers to “not applicable”, but this doesn't fit in the tables.

In this work it means “(data) not available”. We marked it at the tables.

Synthesis

Lines 549-550: Could you cite the studies that imitated weathering in the described way?

We did so. Tsunoda et al. (2010) artificially aged their plastic by soaking in hot water at 90°C for 21 days, and then it was sanded/scratched with medium-grade paper prior to the test. Gebhard and Forster (2018) incubated particles in seawater for 4 weeks to stimulate the formation of biofilms.

[9] Lines 555-557: This is true, but it should be acknowledged that these additives are mainly present in commercial plastics, and therefore, mentioning of additives is not expected for “clean” microbeads specifically synthesized for the experiments. Nevertheless, the disadvantages of using these microbeads has been clearly discussed earlier in this section.

Done.

Conclusions

[10] Line 620-621: I am a little concerned about describing the results as “alarming”. Is it really? The following sentences actually refute this rather strong statement.

Replaced with “considerable”.

[11] Lines 624-629: I would suggest changing the sentence to: “To elucidate [...], the most exact reproduction of plastic concentrations and properties [. . .]”. However, the difficulty here is that very scarce data of limited quality is available on concentrations of microplastic in soils, so a range of concentrations need to be used for future experiments in order to match the “real world” concentrations in soil, while expecting a decrease in uncertainty in analytic results in the future. Especially in the lower size ranges (<100µm) quantification is currently challenging. Therefore, little is known about size distributions occurring in soils. It might be worth mentioning this dilemma in a sentence.

Done.

[12] Technical corrections:

All done.

Dear PD Dr. Werner Kratz (referee #2)

Thank you very much for your review. In the following I will try to answer your comments at my best. Our corrections are marked-up with green numbers within the corrected manuscript at the end of this document.

*Best regards,
Frederick Büks*

[1] Line 53: Is that only "microbial" decay?

We agree, we will change this to "processing by soil organisms" as it is actually micro- as well as macroorganisms.

[2] Figure 1: The taxonomic group "further Panarthropods" is placed centrally, the other groups are not.
Done.

[3] Table 7: The last three experiments within this table were conducted by feeding the mice with a MP suspension. You might write "(food)" behind the concentration data as in the other tables.
Thanks a lot. Done.

[4] Line 507: "Preferably" instead of "preferrably".
Done.

[5] Table 8: Could you explain the meaning of the numbers within the table. Are these the numbers of experiments with the named type-shape combinations?
Yes. Please see the answer to referee #1 (Table 8 is now Table 1).

[6] Lines 549-507: Is that proved that carboxylation of microspheres decreases hydrophobicity in an appreciable extent?
We ask the manufacturers of Polysciences Europe GmbH, a leading producer of PS microspheres, and they said no. We added this important information to the review.

Dear Referee #3

Thank you very much for your critical review of our manuscript. It has helped us to see some points which still need clarification. In the following, we want to explain how we propose to adjust our article based on the reviewer's comments and also explain why in some cases we do not agree with the reviewer's proposed changes. Our corrections are marked-up with **purple numbers** within the corrected manuscript at the end of this document.

(1) First and foremost, please have the manuscript edited by a professional (!) native (!)biologist (!). The English of your text is largely understandable, but rough. Apart from annoying typos, I found sentences the meaning of which I only understood when trying to translate them to German (my native language). So, your text will heavily benefit from thorough native editing.

Rereading our article we did indeed see that some typos had escaped our notice. We are slightly surprised by the request of the reviewer to have the manuscript edited by a "*professional (!) native (!) biologist (!)*". We **rephrased some stiff sentences and corrected grammatical errors**. If a proofreading is indeed wished, we will have a scientific translator (English native speaker) correct the article.

(2) Then, the text lacks conciseness, it is overly long. For example, I suggest to omit all biological/ecological details you provide when introducing a taxon. This is per se interesting, but not to the point here (except when the reader needs background to understand microplastic effects). Then, figure 1 does not contribute to the understanding of your presentation, omit it. And I do not think it necessary to present taxa for which there is no information available, especially if the taxa are of minor or no importance in soil (e.g. line 183ff, 205, 220, 227, 450ff) or if the literature is not on edaphic species (435ff). As a reviewer, you are of course required to address blind spots of research (thus pointing out important taxa that are missing in literature), but you need to better balance completeness with a concise presentation.

Your suggestion to omit the ecological presentation of some key taxa is understandable. If we would expect all readers to be well acquainted with the soil fauna, we would definitely go along with this. However, SOIL is a multi-disciplinary journal connecting a broad spectrum of soil scientists. Therefore, we think it is helpful to provide a short overview of information on the soil fauna, such as ecological functionalities (marker function, transport, degradation, habitat and food selection), which might influence how they cope with microplastics. We have critically gone through the article and here we summarize which parts we will shorten.

- **[1] Proposal:** We shortened the introduction of the springtail section, as it is indeed oversized.

For the same reason we illustrated the phylogenetic tree of soil life.

- **[2] Proposal:** We would agree with moving it to the supplements in order to save space, in case this is wished.

We also do not fully agree with your suggestion to delete taxonomic groups that have not yet been subject of studies on microplastics. The reason is, that the aim of this work is not only to review effects on studied taxa, but also to show gaps of knowledge especially apart from the common model organisms. In fact, their importance for the current ecological research should be shortly mentioned.

- **[2] Proposal:** Unstudied taxa are still presented, but their importance for future research is now additionally mentioned in section 4.3 to better "balance completeness".
- **[3] Proposal:** We shortened the chapter about Onychophora.

Potamopyrgus antipodarum in fact is a benthic snail.

- **[4] Proposal:** We use this benthic species to show more clearly how inconsistent the few results for benthic and terrestrial snails are.

(3) I miss a convincing argumentation why you focus on multicellular animals (but then, you provide many details about bacteria, fungi, algae, plant roots in 72ff...omit this). A good line of reasoning could be that you follow up on the Rillig and Bonkowski (2018) paper.

The aim of this review is to depict the influence of microplastic contamination in soils to the soil fauna. But, to present a holistic view on the food web, we refer to microorganisms, plant roots and biofilms within the introduction section. Being large fields of knowledge on their own, these organisms are not part of the focus in this review, however they are food sources for meso- and macroorganisms and, thus, worthy of mention. Given that we only use 22 lines to describe these other parts of the phylogenetic tree of soil life, we think this is merited and wish to leave this part in the review.

Unfortunately, we do not understand how Rillig and Bonkowski (2018), a paper on soil protozoa, matches your point. We have read this paper and do mention it elsewhere in the review.

(4) Please provide details of your literature search (123ff). When did you search? Which time span did you cover? Which search strings? Please consider the literature on meta-analyses how to properly specify these technical aspects.

The search was applied between June 2019 and January 2020, repeated in the first week of January 2020 and covers publications until January 2020. The search strings result from combinations of taxon, plastic type and particle shape shown in Table 1 (formerly Table 8).

- [5] Proposal: Information added to section 2.

(5) Thank you very much for the positive note.

(6) line 636f: Please reconsider including your supervisor as a co-author. What "supervision" means is nowhere clearly defined, however, co-authorship is only justified for significant contributions to the manuscript. Honorary authorship violates the principle of scientific honesty.

We understand this point completely and agree that it is not good practice to include scientists who have not contributed significantly to a paper. We also acknowledge that supervision is a very broad term and would like to specify the contribution of Martin Kaupenjohann to the paper. [6] Martin Kaupenjohann was involved in the development of the idea and concept for this paper. During the literature reading and writing phase he has supported the work with frequent discussions of the contents of the article. And finally he has critically revised the manuscript.

Best regards,

Dr. Frederick Büks
Dr. Loes van Schaik
Prof. Dr. Martin Kaupenjohann

Dear Dr. Maha Deeb (referee #4)

Thank you very much for the repeated check and your friendly report.

Best regards,

Dr. Frederick Büks

What do we know about how the terrestrial multicellular soil fauna reacts to microplastic?

Frederick Büks¹, Nicolette Loes van Schaik², Martin Kaupenjohann¹

¹Chair of Soil Science, Dept. of Ecology, Technische Universität Berlin, 10587 Berlin, Germany

²Chair of Ecohydrology & Landscape Evaluation, Dept. of Ecology, Technische Universität Berlin, 10587 Berlin, Germany

Correspondence to: Frederick Büks (frederick.bueks@tu-berlin.de)

Abstract. The ubiquitous accumulation of microplastic (MP) particles across all global ecosystems comes along with the uptake into soil food webs. In this review, we analyzed studies on passive translocation, active ingestion, bioaccumulation and adverse effects within the phylogenetic tree of multicellular soil faunal life. The representativity of these studies for natural soil ecosystems was assessed using data on the type of plastic, shape, composition, concentration and time of exposure.

Available studies cover a wide range of soil organisms, with emphasis on earthworms, nematodes, springtails, beetles and lugworms, each focused on well-known model organisms. [1] About 58 % of the studies thereby used inappropriate concentrations or units, but 42 % applied MP concentrations similar to amounts in slightly to very heavily polluted soils. In many cases, however, polystyrene microspheres have been used, a combination of plastic type and shape, that is easily available, but does not represent the main plastic input into soil ecosystems. In turn, MP fibers are strongly underrepresented compared to their high abundance within contaminated soils. [7] A few studies also examined the comminution of macroplastic by the soil fauna. Further properties of plastic such as aging, coating and additives were insufficiently documented. Despite these limitations, there is a recurring pattern of active intake followed by a population shift within the gut microbiome and adverse effects on motility, growth, metabolism, reproduction and mortality in various combinations, especially at high concentrations and small particle sizes.

For the improvement of future studies, we identified problems of past experiments and recommend that coming studies take into account the type, shape, grade of aging, specific concentrations of MP fractions and long-term incubation, in natural and contaminated soils.

34 1 Introduction

35 Imagine a compact plastic cube of nearly 2 km side length and a weight of 7300000000 tons,
36 with major percentages by weight of 36 % polyethylene (PE), 21 % polypropylene (PP), 12 %
37 polyvinyl chloride (PVC) and 10 % of each polyethylene terephthalate (PET), polyurethane
38 (PU) and polystyrene (PS). That is the accumulated global non-fiber production of the six
39 main plastic types until 2015. It accounts to 87 % of the all-time plastic production, which
40 evolved exponentially, since the early 1950s, from some megatons (Mt) to 8300 Mt in 2015,
41 with only 260 Mt annual output in 2009 increased to 380 Mt in 2015 (Thompson et al., 2009;
42 Geyer et al., 2017). Of this ever produced plastic, 6300 Mt became waste until 2015, of which
43 only 21 % were recycled or incinerated, whereas 5000 Mt ended up in landfills and nature
44 (Geyer et al., 2017). As a corollary of production, use and disposal, a certain part of plastic
45 waste is constantly released into the environment through various pathways, but our
46 knowledge about rates of mass flow into global ecosystems is very limited. Based on waste
47 generation in coastal countries, Jambeck et al. (2015) calculated the global plastic input to
48 marine ecosystems to be roughly 4.8 to 12.7 Mt in 2010. Such data on soils are lacking, but
49 Nizzetto et al. (2016) estimated that the load of microplastic (MP) to agricultural sites in
50 Europe is in the same order of magnitude as that in marine environments.

51 By littering, plastic mulching, the application of sewage sludge, digestates and composts as
52 well as windblown dispersal (Bertling et al., 2018; Weithmann et al., 2018; Zhang et al., 2019;
53 Wang et al., 2019a), plastic from our technosphere arrives in soil ecosystems in various forms
54 as large and small fragments, fibers and particles. Exposed to UV radiation, mechanical
55 stress and ~~211~~ processing by soil organisms, plastic items become weathered and prone to a
56 successive comminution towards the size range of MP with increased surface, charge and
57 biofilm cover (Kale et al., 2015; Andrady, 2017). However, the resistance of plastic to
58 metabolization causes a constant accumulation in soils as long as the release rate from
59 human processes is above the very slow rate of degradation.

60 Due to a lack of monitoring programs, data on MP concentrations in terrestrial soils are rare,
61 and those using w/w concentrations represent only a small part compared to item
62 concentrations. In soils with only slightly contaminated conditions, amounts seem to average
63 about 1 mg kg⁻¹ soil dry weight (and approx. 200 items kg⁻¹ dry soil) (Rezaei et al., 2019). On
64 sites with industrial activity or ~~13~~ intensive use of plastic mulching and sewage sludge in
65 agriculture, concentrations can be increased by 2 to 4 orders of magnitude (Fuller and
66 Gautam, 2016; Zhang and Liu, 2018; Huang et al., 2020). Semisubhydic soils such as
67 beaches, mudflats, mangroves or lagoons, that are additionally contaminated from the aquatic
68 side, contain MP of the order of 10 to 100 items kg⁻¹ dry soil and single extreme samplings
69 contained several thousand items (Nor and Obbard, 2014; Naji et al., 2017; Garcés-Ordóñez
70 et al., 2019; Li et al., 2018a). More informative data using mg kg⁻¹ are only available for
71 beaches and coastal deconstruction yards in municipal neighborhoods and amount to 0.5 and

72 70 mg kg⁻¹ dry soil, 0.00005 and 0.007 % w/w, respectively (Reddy et al., 2006; Claessens et
73 al., 2011). All these concentration data represent a wide range of particle sizes between 0 and
74 5000 µm with different materials, shapes and degrees of aging.

75 Plastic particles can possibly enter and accumulate in the food web by either direct uptake
76 from soil or by consumption of other soil biota contaminated by adhesion or ingestion (Huerta
77 Lwanga et al., 2017a). There is evidence, that MP is incorporated even by plants and
78 unicellular organisms at the base of the food web. **Bacteria**, for example, that are reasonably
79 assumed to avoid MP uptake due to their minor size and the prevalent lack of phagocytosis,
80 were shown to take up inorganic nanoparticles of a few nanometers (Kumar et al., 2011).
81 Although the physiochemical properties of weathered nanoparticulate plastics might differ
82 from these, also their uptake seems [12] likely.

83 A similar argument can be made for **fungi** and soil **algae**, but studies on incorporation are
84 lacking, whereas the transfer into a freshwater food web by adhesion of nanoplastic on algae
85 has been shown by Chae et al. (2018). The uptake of MP into **plant roots** is also inhibited
86 (Rillig et al., 2019), but occurred for nanoplastics that permeate into the plant tissue (Li et al.,
87 2019). Also the integration into root tissue after adsorption to the rhizodermis [12] has yet to be
88 studied.

89 In contrast, **protozoa** feature phagocytosis for the active ingestion of particles. Diverse soil,
90 freshwater and marine ciliates ingest PS/latex beads of 0.1 to 14.4 µm in laboratory
91 experiments, with preferences to their natural prey size (Fenchel, 1980; Jonsson, 1986; Lavin
92 et al., 1990). Soil amoebas act similarly, but additionally select according to food quality
93 (Weisman and Korn, 1967; Vogel et al., 1980; Bowers and Olszewski, 1983; Avery et al.,
94 1995; Elloway et al., 2006).

95 Finally, many soil microbiota live protected within biofilms. Plastic particles were shown to be
96 a potential surface for the formation of those biofilms (Lobelle and Cunliffe, 2011), which are a
97 food source for grazing primary consumers. [4] Inadvertent ingestion might also transfer
98 occluded or abraded MP to higher trophic levels.

99 But what about the larger organisms that feed on all these, free plastic particles,
100 contaminated microorganisms, biofilms and one another? Recent work discussed the effects
101 of MP on soil biota (Chae and An, 2018) or called for intensified research on certain
102 taxonomic groups (Rillig and Bonkowski, 2018). Thus, we were motivated to give on our part
103 a review with focus on the most-produced plastics and their passive translocation, ingestion,
104 bioaccumulation and adverse effects on the multicellular soil fauna. The types, sizes and
105 shapes of plastic used in former laboratory studies were compared with the available
106 knowledge on plastic in the environment, and recommendations are given for future research.
107 [12] This analysis aims to support the assessment of the influence of MP on the ecosystem
108 services provided by diverse soil organisms.

109 2 Search pattern

110 Within the tree of life, edaphic branches were identified comprising taxa that permanently
 111 inhabit the soil, are both-sided part of the soil food web and/or the burrowing macro- and
 112 megafauna or have active subterranean larval stages. The resulting tree of soil life based on
 113 the NCBI taxonomy database (Fig. 1) was drawn using the software [phyloT](#) and shows the
 114 leading taxonomic rank, which is mainly the family, but in exceptions – e.g. if one species
 115 represents the only soil-born between many aquatic – a lower rank.




Figure 1: Tree of edaphic fauna. Taxonomic ranks, that were examined in this qualitative study, are placed at the outer rim of the diagram. The length of the connecting line between two taxa represents the grade of phylogenetic relationship.

117 A pattern of search terms was established (71 Table 1), consisting of „taxon“ (Linné’s
 118 binominal nomenclature, common name, plural-sensitive search), „plastic type“ (plastic,
 119 microplastic, nanoplastic, PE or polyethylene, PP or polypropylene, PVC or polyvinyl chloride,
 120 PS or polystyrene, PU or polyurethane, PET or polyethylene terephthalate and latex) and
 121 „common shapes“ (fragments, particles, fibers, microfibrs, beads, microbeads,
 122 microspheres). 71 Some type- shape combinations caused problems, as they led to a very
 123 large amount of unuseful, off-topic papers – e.g. using any taxon combined with PET, papers
 124 with the use of PET bottles in experimental set-ups were selected or also studies on pets.
 125 Those combinations of search terms were excluded from this pattern. Further plastic types
 126 and shapes occurring within the found studies were also included in the review. Data on
 127 microspheres and microbeads were pooled, as both names describe one and the same.

128

Table 1: 51 Types and shapes of microplastic particles in edaphon studies within this review. (X) symbolizes combinations excluded from the search pattern. 71 The number counts for how often type-shape combinations were used in all reviewed experimental setups independently of organism. Empty fields stand for zero results. Microbeads and microspheres are often mixed up terms and, thus, counted together.

Organism: Linné’s systematic names OR common name	fragments	particles	fibrs	microfibrs	beads	microbeads	microspheres	other, diverse, N/A
plastic	X							
microplastic								
nanoplastic								
PE OR polyethylene	X	4	10	1	1	1	4	7
PP OR polypropylene	X	1						
PVC OR polyvinyl chloride	X	4	6	1				
PS OR polystyrene	X	6	3				24	4
PU OR polyurethane	X							
PET OR polyethylene terephthalate	X	3	2				X	
latex	X			X			6	
other		6	3	1				1
N/A		1	1		2			3

129

130

131 The search was applied 5151 between June 2019 and January 2020 within the Web of Science
 132 Core Collection Database, repeated in the first week of January 2020 and covers publications
 133 until January 2020. The search strings result from combinations of taxon, plastic type and
 134 particle shape shown in Table 1. Based on the search pattern, data on passive transport,

135 ingestion, bioaccumulation and adverse effects were collected for each edaphic group.
136 Studies that only use uncommon, local, outdated, weird or nicknames are excluded by the
137 search pattern. Studies testing injection to tissues, lymph or blood were excluded, as they do
138 not represent natural ways to incorporate MPs. Data on inhalation by the megafauna in fact
139 represent a natural way of uptake, but were also excluded as they are exclusively related to
140 above-ground organisms, that only occur on the outer edge of the food-web. Also running
141 debates on phylogenetic classifications are not part of this work and the [12] taxonomists will
142 be able to adjust the branches accordingly to [12] their purpose.

143 The data of related taxonomic groups were pooled and evaluated for their environmental
144 representativity based on exposure time, plastic concentrations and properties used. From
145 this synthesis recommendations for a structured experimental design were derived for
146 application in future studies.

147 3 Data collection

148 149 3.1 Insects

150 Within the Panarthropoda, the insects comprise the highest taxonomic diversity. And,
151 regarding MPs, they represent an unevenly studied taxonomic group.

152 Within the Insecta, the **Coleoptera** (beetles) are an extraordinarily diverse and abundant
153 taxon. Studies on plastic uptake into adult individuals mainly focus on the subfamily of
154 Scarabaeinae (dung beetles). Comprehensive experiments with latex microbeads showed,
155 that many species only ingest [12] smaller particles with maximum diameters of about 10 to
156 83 µm and retain them within the gut – with a slightly positive dependency on body size.
157 Larger particles were rejected by a filtering mechanism within the mouth region and not
158 ground with the mandibles (Holter, 2000; Holter et al., 2002; Holter and Scholtz, 2005).
159 Beside those on Nematods, these data comprise by far the most detailed information about
160 [12] size-dependent uptake of MP particles compared to other edaphic taxa. This gives a good
161 foundation for future studies on adverse concentrations. In addition, several studies with
162 plastic as predominant food source could show chewing, ingestion and intestinal degradation
163 of different PS and PE foams in feeding experiments with *Tenebrio sp.* larvae (mealworms).
164 These experiments also pointed out an alteration of the gut microbiome, but no adverse
165 effects on reproduction and survival, with only one case of a non-significant tendency to
166 higher mortality after 1 month of exposure (Yang et al., 2015; Brandon et al., 2018; Yang et
167 al., 2018; Peng et al., 2019).

168 The **Isoptera** (termites), recently categorized as part of the order Blattodea, are the oldest
169 social insects with a tribal history of about 130 million years (Korb, 2008). Especially in arid
170 ecosystems with a lack of earthworms they play an important role in homogenization of soils,
171 but also in sorting of soil mineral particles for building mounds as well as decomposition and
172 distribution of organic matter (De Bruyn and Conacher, 1990). Tsunoda et al. (2010) and Lenz
173 et al. (2012) could show, that different termite species are picky feeders and erode PE, but
174 avoid other plastic cable sheathings. This suggests the excretion of ground MP particles by
175 termites, but metabolic impacts are unknown. In contrast to termites, data on **other Blattodea**
176 (e.g. cockroaches) were not found.

177 The suborder **Apocrita** comprises some flying insects, that inhabit burrows within the soil,
178 such as ground-dwelling wasps within the **Vespidae** superfamily, mining bees within the
179 **Apoidea** superfamily and the **Spheciformes**. They mostly do not prey and feed on
180 subterrestrial organisms, but may move MP particles into the ground, as implied by a report of
181 Allasino et al. (2019) on [12] solitary bees, which built nests fully made of plastic fragments.
182 The Apocrita also contain the **Formicidae** (ants). Some ant species are considered an
183 important factor for seed dispersal, a behavior, that could also be shown for artificial plastic
184 seeds with ~2 mm diameter (Hughes and Westoby, 1992; Angotti et al., 2018). Robins and

185 Robins (2011) found that this also includes differently shaped cultural objects: *Rhytidoponera*
186 *metallica*, a representative of ground-nesting, omnivore ants, is capable not only of a
187 remarkable bioturbation, but also of an active, apparently random burying of anthropogenic
188 plastic artifacts >1 mm. Seeds are used as a food source, thus, the ingestion of plastic bites is
189 conceivable, but not documented. The uptake of latex microspheres $\geq 0.88 \mu\text{m}$ with liquids by
190 larvae of *Solenopsis invicta* seems to be prevented by filtration within the mouth and the
191 particles are released as larger aggregates, whereas other species ingest by far larger
192 particles up to $150 \mu\text{m}$ (Glancey et al., 1981). However, also here data on adverse effects are
193 missing.

194 Further insects with edaphic adult stages, e.g. **Dermaptera** (earwigs), **Heteroptera** (true
195 bugs) and **Zygentoma** (silverfish, fishmoth, firebrat) or soil- or litter-dwelling larvae such as
196 **Embiopoda** (webspinners, footspinners), **Thysanoptera** (thrips), **Psocoptera** (booklice,
197 barklice, barkflies), **Neuroptera** (lacewings), **Raphidioptera** (snakeflies) or **Zoraptera** (angel
198 insects) were not yet researched with focus on soil MP.

199 Regarding insects, mainly studies on translocation and uptake of MP were carried out. In
200 contrast, work on bioaccumulation is completely lacking and adverse effects are sparsely
201 tested using *Tenebrio sp.* larvae. Such studies could provide information whether or not the
202 input of MP in soil ecosystems is one of many factors causing the global decline of the
203 entomofauna (Oliveira et al., 2019; Sánchez-Bayo and Wyckhuys, 2019).

Table 2: Microplastic studies on Coleoptera, Blattodea (Blattod.), Apoidea (A.) and Formicidae (mb=microbeads, fr=fragments, ms=microspheres, b=beads, [8] N/A=information not available). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

	organism	experimental environment	plastic type	aging	coating	additives	shape	size span [μm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
Coleoptera	<i>Aphodius erraticus</i>	Petri dish	latex	N/A	N/A	N/A	mb	5	N/A	45 min	N/A	no	N/A	N/A	Holter (2000)
	<i>Aphodius rufipes</i>							2..39				≤14 μm			
	<i>Aphodius ater</i>							2..39				≤14 μm			
	<i>Aphodius fimetarius</i>							2..39				≤18 μm			
	<i>Aphodius contaminatus</i>							2..39				≤18 μm			
	<i>Aphodius fossor</i>							2..39				≤18 μm			
	diverse dung beetles	vial	latex	N/A	N/A	N/A	mb	2..83	N/A	45 min	N/A	≤10..≤60 μm	N/A	N/A	Holter et al. (2002)
	diverse dung beetles	N/A	latex	N/A	N/A	N/A	mb	2..83	N/A	45 min	N/A	≤4..≤95 μm	N/A	N/A	Holter and Scholtz (2005)
	<i>Tenebrio molitor</i> larvae	container	PS	N/A	N/A	no	foam	N/A	100% w/w (food)	31 d	N/A	yes	biodegrad.	N/A	Yang et al. (2015)
Blattod.	<i>Tenebrio molitor</i> larvae	container	LD-PE PS	N/A	N/A	no flame retardant	foam	8..27 cm ³	50..100% w/w (food)	32 d	N/A	yes	biodegrad.	microbiome	Brandon et al. (2018)
	<i>Tenebrio molitor</i> larvae	container	PS	N/A	N/A	N/A	foam	N/A	4..100% w/w (food)	32 d	N/A	yes	biodegrad.	no	Yang et al. (2018)
	<i>Tenebrio molitor</i> larvae	N/A	PS	N/A	N/A	no	foam	N/A	86..100% w/w (food)	31 d	N/A	yes	biodegrad.	microbiome	Peng et al. (2019)
	<i>Tenebrio obscurus</i> larvae														
A.	<i>Coptotermes formosanus</i>	mesocosm	LD-PE others	yes/no	N/A	N/A	cable sheets	4 cm, Ø 0.8 cm	N/A	42 d	N/A	yes no	N/A	N/A	Tsunoda et al. (2010)
	diverse termites	in situ	MD-PE PA	no	N/A	anti-oxidant stabilizer	cable sheets	30 cm, Ø 1.4 cm	N/A	6 yr.	N/A	yes no	N/A	N/A	Lenz et al. (2012)
Formicidae	<i>Megachile</i> sp.	in situ	N/A	N/A	N/A	N/A	fr	N/A	N/A	N/A	yes	N/A	N/A	N/A	Allasino et al. (2019)
	<i>Solenopsis invicta</i>	Petri dish	latex	N/A	N/A	fluorescence	ms	0.9..4.5	2.5% w/w (food)	direct	N/A	filtration	N/A	N/A	Glancey et al. (1982)
	<i>Rhytidoponera metallica</i>	in situ	N/A	N/A	N/A	N/A	b	N/A	50 items per nest	3 d	yes	N/A	N/A	N/A	Hughes and Westoby (1992)
	<i>Aphaenogaster longiceps</i>														
	<i>Pheidole</i> sp.														
Formicidae	<i>Rhytidoponera metallica</i>	mesocosm	N/A	N/A	N/A	N/A	diverse	<75.5 cm	N/A	26 mos.	yes	N/A	N/A	N/A	Robins and Robins (2011)
	diverse ants	in situ	N/A	N/A	attractant	N/A	b	1.8 cm	N/A	1 d	yes	N/A	N/A	N/A	Angotti et al. (2018)

205 3.2 Other panarthropods

206 Apart from the insects, **Acari** (mites) comprise many abundant soil-living taxa, that feed on
207 litter, fungi and fauna as predators and parasites and are bioindicators, as they are sensitive
208 to changes in the soil physiochemical environment (Gulvik, 2007). Experiments indicated, that
209 mites passively transport MP due to pushing and dragging after attachment to their cuticle, as
210 shown with 80 to 250 µm sized PVC particles in a **Petri** dish experiment without soil (Zhu et
211 al., 2018a). The population within manure pats slightly declines when exposed to mm-sized
212 unweathered PE and PS particles at concentrations of 5 % v/v and declines strongly at
213 ≥60 % v/v (Stamatiadis and Dindal, 1990). This could probably be an effect of moisture
214 deficiency due to a reduced water holding capacity in an unnaturally enriched substrate, but
215 not necessarily through plastic intake. In contrast, no data was found on their arachnoid,
216 preying relatives, the order of **Pseudoscorpiones** (false scorpions).

217 Just as many other highly abundant and diverse representatives of the soil mesofauna, the
218 **Oniscideae** (woodlice) contribute to the decomposition of litter by chewing and passage
219 through their digestive system (Warburg, 1987) and react strongly to environmental pollution,
220 thus, they are potentially used as bioindicators (van Gestel et al., 2018). They practice a strict
221 selection of natural food sources (Hassall and Rushton, 1984). This is also demonstrated for
222 starch and cellulose based plastic films (4 cm²), which were consumed and digested in
223 experiments with the model organism *Porcellio scaber*, in contrast to PHB
224 (polyhydroxybutyrate) films, that reduces the feeding rate (Wood and Zimmer, 2014). Smaller
225 PE particles (137±51 µm and 183±93 µm) embedded into food pellets (0.4 % w/w) were taken
226 up easily by *Porcellio scaber*, and the smaller fraction caused a slight and non-significant
227 reduction of body mass after 14 days of exposure, but not of feeding, defecation or energy
228 reserves (Kokalj et al., 2018).

229 Other panarthropodean groups are even less studied in terms of MP. We did not find literature
230 on the subphylum of Myriapoda containing the classes of **Diplopoda** (millipedes), **Chilopoda**
231 (centipedes), **Paupoda** and **Symphyla** (pseudocentipedes or symphilids), which are
232 important litter-feeders and predators within various soil ecosystems.

233 The situation is nearly similar with the phylum of **Tardigrada** (water-bears or tardigrades), that
234 has many ecologically relevant and well studied species feeding on microorganisms and
235 detritus particles. Sparse field research in semisubhydric environments showed no uptake of
236 MP **fibers** by tardigrada (Gusmão et al., 2016), but comprehensive data on terrestrial soils are
237 lacking. ¹³ Similarly, the related phylum of **Onychophora** (velvet worms), primordial
238 invertebrates that are mainly native in litter and soils with high water holding capacity under
239 pleistocene-like forest vegetation within tropical and moderate regions (Monge-Nájera, 1994).

240 ~~Another branch within the panarthropoda, the phylum of **Onychophora** (velvet worms),~~
241 ~~comprises primordial invertebrates that are mainly native in litter and soils with high water~~

242 holding capacity under pleistocene-like forest vegetation within tropical and moderate regions
243 (Monge-Nájera, 1994). As predators, they most likely take up plastic debris appearing within
244 or on their prey, but no studies on MP are available, most likely due to their remote habitats,
245 low abundance and little scientific focus.

246 The phylum of **Collembola** (springtails) [11], together with the **Diplura** and **Protura** (Westheide
247 and Rieger, 1996; Pass et al., 2011), an abundant, diverse and ubiquitous soil-borne phylum
248 with a broad spectrum of food sources (Hopkin, 1997), also represent an intensively studied
249 group within the Arthropoda. Together with the **Diplura** (which mainly live in tropic and
250 subtropic regions in litter and humid topsoil and feed on fungal hyphae, POM and prey)
251 (Westheide and Rieger, 1996) and the **Protura** (Pass et al., 2011), the **Collembola** build an
252 intensively studied morphological group, that [12] exhibits similar ecological functions, such as
253 distribution and decomposition of organic matter as well as the control of fungal abundance
254 (Hopkin, 1997). Springtails provide up to 27 % of the soil biomass and up to 33 % of the total
255 soil respiration (with higher shares in colder ecosystems) (Petersen, 1994) with up to 100000
256 individuals per square meter (Hopkin, 1997). Thus, their well-being plays an important role for
257 ecosystem functioning.

258 In a Petri dish experiment without soil, Maaß et al. (2017) showed the passive transport of
259 urea-formaldehyde particles <400 µm and undefined PET fragments by two Collembola
260 species (*Folsomia candida* and *Proisotoma minuta*) due to attachment, but found no
261 ingestion. Within a soil matrix, trials of Kim and An (2019) indicated hindrance of collembolan
262 migration by larger PS particles (44±39, 282±131 and 676±479 µm) at concentrations of
263 1000 mg kg⁻¹ corresponding to highly contaminated soils. In addition, they found suppressed
264 mobility due to the attachment of even smaller PS microbeads (0.47 to 0.53 µm) at
265 concentrations of 8 mg kg⁻¹ dry soil, which is equivalent to values found in nature. Small
266 particles <50 µm were moved, while larger particles were most likely [12] cast off. When *F.*
267 *candida* encounters two of its predators, the mites *Damaeus exspinosus* and *Hypoaspis*
268 *aculeifer*, the dispersal of 80 to 250 µm PVC particles is enhanced as shown by Zhu et al.
269 (2018a) in a Petri dish experiment. Without proving the ingestion or the minimal effective MP
270 concentration, Zhu et al. (2018b) published an alteration of the gut microbiome and adverse
271 effects on growth and reproduction of *F. candida* by 80 to 250 µm PVC particles mixed in soil
272 at concentrations of 1000 mg kg⁻¹ dry soil. These data were not considered robust (van Gestel
273 and Selonen, 2018), but match with a later study that found inhibited reproduction at
274 ≥1000 mg kg⁻¹ and avoidance behavior as well as microbiome alteration at ≥5000 mg kg⁻¹ (Ju
275 et al., 2019). Such concentrations can occur in highly contaminated soils (Fuller and Gautam,
276 2016). However, documentations on the active uptake, gnawing and grinding of MP by
277 springtails proposed by Rillig (2012) is still lacking and also studies on Diplura and Protura
278 were not found.

Table 3: Microplastic studies on Acari, Oniscidea (Onisc.), Tardigrada (T.) and Collembola (fr=fragments, p=particles, mf=microfibers, mb=microbeads, ms=microspheres, ^s=semisubhydic, [8] N/A=information not available). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

	organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
Acari	diverse mites	microcosm	PE PS	no	N/A	N/A	fr	<4800 >2000	0..90% v/v (manure)	16 d	N/A	N/A	N/A	≥5% v/v: abundance ↓	Stamatiadis and Dindal (1990)
	<i>Hypoaspis aculeifer</i> <i>Damaeus exspinosus</i>	Petri dish	PVC	N/A	no	N/A	p	80..250	5000 items per dish	N/A	yes	N/A	N/A	N/A	Zhu et al. (2018a)
Onisc	<i>Porcellio scaber</i>	mesocosm	PHB	no	N/A	N/A	fr	4 cm ²	1 item per cosm	14 d	N/A	yes	N/A	feeding ↓	Wood and Zimmer (2014)
	<i>Porcellio scaber</i>	Petri dish	PE	N/A	N/A	N/A	fr	183±93 137±51	0.4% w/w (food)	14 d	N/A	yes	N/A	no	Kokalj et al. (2018)
T.	diverse tardigrades ^s	in situ	N/A	N/A	N/A	N/A	mf	N/A	N/A	N/A	N/A	no	N/A	N/A	Gusmão et al. (2016)
Collembola	<i>Folsomia candida</i> <i>Proisotoma minuta</i>	cup	UF, PET	N/A	no	N/A	p,fr	<400	2.5..5 mg per cup	N/A	yes	N/A	N/A	N/A	Maaß et al. (2017)
	<i>Folsomia candida</i>	Petri dish	PVC	N/A	no	N/A	p	80..250	5000 items per dish	N/A	yes	N/A	N/A	N/A	Zhu et al. (2018a)
	<i>Folsomia candida</i>	microcosm	PVC	N/A	no	N/A	p	80..250	1000	56 d	N/A	N/A	N/A	microbiome, growth ↓, reproduction ↓	Zhu et al. (2018b)
	<i>Folsomia candida</i>	microcosm	PE	N/A	no	N/A	mb	<500	0..10000 0..10000 0..5000	7 d 28 d 28 d	N/A	N/A	N/A	≥5000: avoidance ≥1000: reproduction ↓ ≥5000: microbiome	Ju et al. (2019)
	<i>Lobelia sokamensis</i>	soil sample	PS	N/A	carboxyl	fluorescence	mb	0.5	4..8		yes				
			PE	no	N/A	fluorescence	ms	27..32	1000		yes				
			PE	no	N/A	fluorescence	ms	250..300	1000		N/A				
			PS	no	N/A	no	fr	44±39	1000	≤3 min	yes				
			PS	no	N/A	no	fr	282±131	1000		N/A				
			PS	no	N/A	no	fr	676±479	1000		N/A				
												N/A	N/A	avoidance, motivity ↓	Kim and An (2019)

280 3.3 Annelida

281 Land-based Annelida comprise another large group of invertebrates. The Lumbricidae
282 (earthworms) are a well-studied family (Darwin, 1881; Lavelle et al., 2006), represented in
283 high abundance and diversity in many ecosystems all around the world (Phillips et al., 2019).
284 Earthworms are often used as indicators for soil health (Fründ et al., 2011; Pulleman et al.,
285 2012), as they are ecosystem engineers which through their burrowing activity influence
286 various soil physical, chemical and biological processes (Jouquet et al., 2006; Lavelle et al.,
287 2006).

288 By far the most of the studies on the influence of MP on earthworms are performed with PE
289 and the species *Lumbricus terrestris* or *Eisenia fetida*, but there are also single studies with
290 *Aporrectodea rosea* (Boots et al., 2019) and *Eisenia andrei* (Rodriguez-Seijo et al., 2017) and
291 with the less common species *Metaphire californica* (Wang et al., 2019b). We found one field
292 study of earthworms and MPs (Huerta Lwanga et al., 2017a) among many laboratory
293 experiments with MPs mixed into soil volumes (concentrations ranging up to 20000 mg kg⁻¹
294 dry soil) or applied with litter on top of the soil surface (≤60% w/w). The particles sizes were
295 usually <1 mm in diameter, but some were even up to 2x2 cm², and the duration of
296 experiments was generally 14 to 28 days, few lasted up to 60 days.

297 The uptake of MPs of a broad size range by earthworms was shown in studies based on
298 particles in earthworm casts of *Lumbricus terrestris* (Huerta Lwanga et al., 2016; Cao et al.,
299 2017; Hodson et al., 2017; Rillig et al., 2017; Prendergast-Miller et al., 2019; Yu et al., 2019;
300 Huerta Lwanga et al., 2017a), *Eisenia fetida* (Rodríguez-Seijo et al., 2018; Chen et al., 2020;
301 Wang et al., 2019c), *Eisenia andrei* (Rodriguez-Seijo et al., 2017) and *Metaphire californica*
302 (Wang et al., 2019b). Zhang et al. (2018) showed that relatively large PE particles of
303 1.5 x1.5 cm² are not ingested by *Lumbricus terrestris*, but partial ingestion of such large
304 particles of biodegradable MPs does take place after initial weathering in soil or in compost
305 has occurred. In some laboratory experiments, MPs were found in the gut of dissected
306 earthworms (Huerta Lwanga et al., 2016; Hodson et al., 2017; Rodriguez-Seijo et al., 2017),
307 but the concentration of MPs in the gut was not significantly different between treatments nor
308 significantly different from the bulk soil concentration, so there was no evidence of
309 accumulation of MPs in the earthworm bodies (Hodson et al., 2017). Chen et al. (2020)
310 assume an accumulation of MP takes place in *Eisenia fetida*, based on an observed increase
311 of MP concentrations in the casts in the course of 4 weeks. Huerta Lwanga et al. (2017a)
312 supposed an accumulation of MPs in the food chain as the concentration of MPs in chicken
313 gizzards is strongly increased compared to that in the earthworm casts in the same
314 experiments. However, mainly the amount of large particles, i.e. macroplastics, in the gizzards
315 was very large, thus it seems likely that the chicken directly fed on plastics and an
316 accumulation through the food chain cannot be proven with the current knowledge and should
317 be further investigated.

318 Several studies did not find significant negative effects of MPs on earthworms' avoidance
319 behaviour (Judy et al., 2019), nor on growth (Hodson et al., 2017; Rodriguez-Seijo et al.,
320 2017; Judy et al., 2019; Wang et al., 2019c), mortality (Hodson et al. (2017); Rillig et al.
321 (2017); Rodriguez-Seijo et al. (2017); Judy et al. (2019); Prendergast-Miller et al. (2019) or
322 reproduction (Huerta Lwanga et al., 2016; Rodriguez-Seijo et al., 2017). However, other
323 studies do show adverse effects of the uptake of MP in different degrees and on different
324 aspects of earthworms' fitness: A reduced growth was shown by Cao et al. (2017) for *Eisenia*
325 *Fetida* and the mortality increased at an exposure of concentrations ≥ 10000 mg kg⁻¹ dry soil.
326 At lower concentrations no significant effects were found. The growth of *Aporrectodea rosea*
327 was also inhibited when exposed to biodegradable polylactic acid, conventional high-density
328 polyethylene (at 1000 mg kg⁻¹ dry soil), and MP clothing fibers (at 10 mg kg⁻¹ dry soil) (Boots
329 et al., 2019). Huerta Lwanga et al. (2016) showed a decrease in growth and increased
330 mortality at concentrations $\geq 28\%$ w/w in litter and after 60 days, though after just 14 days no
331 mortality occurred in these experiments.

332 In some studies, additional effects such as histopathological changes or stress biomarkers
333 were measured. For *Eisenia fetida* Chen et al. (2020) observed skin damage at
334 1500 mg MP kg⁻¹ in soil, measured an increase in catalase activity and malondialdehyde
335 content at 1000 mg kg⁻¹ and at ≥ 1000 mg kg⁻¹ acetylcholine esterase was significantly
336 stimulated. Wang et al. (2019c) tested *Eisenia fetida* and found that MPs only increased the
337 catalase and peroxidase levels as well as the level of lipid peroxidation and decreased the
338 activity of superoxide dismutase and glutathione S-transferase at an exposure of
339 200000 mg kg⁻¹ dry soil for 14 days. No discernible influence was found at 100000 mg kg⁻¹.
340 However, Rodríguez-Seijo et al. (2018) also found for *Eisenia fetida* a significant positive
341 correlation of MP concentration with different biomarker responses: catalase, glutathione S-
342 transferase, lactate dehydrogenase and thiobarbituric acid reactive substances. In addition,
343 Rodriguez-Seijo et al. (2017) observed histological damage of the gut and occurrence of
344 inflammatory processes as well as an increase of stress response indicators associated with
345 MP exposure of *Eisenia andrei*. For *Lumbricus terrestris* Prendergast-Miller et al. (2019)
346 showed an increase in metallothionein expression at an exposure with ≥ 1000 mg kg⁻¹ dry soil
347 and a decrease in heat shock protein 70 at a concentration of ≥ 10000 mg kg⁻¹.

348 Due to the large differences in experimental conditions – e.g. size of the MPs, addition of MPs
349 to soil or to litter, duration of experiments, earthworm species – the current knowledge is not
350 sufficient to detect whether there is a threshold in MP size and concentration at which the MP
351 become harmful for earthworms and how this threshold differs for different earthworms
352 species and MP shapes. The results of Huerta Lwanga et al. (2016), who found no effects of
353 MPs on earthworms at 14 days, but significant influence on growth and mortality after
354 60 days, indicate the importance of longer measurements. This is consistent with Pelosi et al.

355 (2015), who concluded that the influence of pesticides on earthworm communities should be
356 tested in long term field experiments.

357 Earthworms activity also increased the transport of MP in soil columns to deeper soil layers
358 (Rillig et al., 2017; Yu et al., 2019; Huerta Lwanga et al., 2017b). The smaller the MP the
359 stronger the transport. Particles are transported both actively – ingested and later cast out –
360 and passively after attachment to the earthworm's body or by water flow through the biopores.
361 As Huerta Lwanga et al. (2018) showed that the bacteria in the gut of *Lumbricus terrestris* can
362 decompose MPs, it seems likely that particles taken up at the surface are egested as smaller
363 particles in deeper layers.

364 Microplastics might well serve as a vector for contaminant transport to soil organisms. Though
365 adsorption on plastics was seen to be lower than on the soil matrix, the desorption of Zn was
366 seen to be higher in synthetic earthworm guts. However, there was no measurable negative
367 effect of Zn or the PE on *Lumbricus terrestris* (Hodson et al., 2017). Wang et al. (2019b)
368 studied the influence of MP on arsenic uptake and negative effects on *Metaphire californica*
369 and concluded that MPs decreased the uptake of arsenic and that MPs reduced the influence
370 of arsenic on the gut bacterial communities. Rodríguez-Seijo et al. (2019) showed altered
371 enzyme activities and enhanced avoidance behavior in face of LD-PE pellets spiked with the
372 insecticide chlorpyrifos. Yang et al. (2019a) studied the influence of MPs on the transport of
373 glyphosate, however they mainly showed that the glyphosate transport was increased by
374 earthworm activity, the role of MPs in this transport could not be determined with this study.
375 These studies show that MP might have very different influences on the uptake and the
376 adverse effects of different pollutants on earthworms and further investigation is needed in
377 order to understand the influence of MPs on pollutant transport.

378 In contrast to the recently well-researched Lumbricidae, a near relative, the family of
379 **Megascolecidae** (giant earthworms), is not yet mentioned in literature. Another branch within
380 the Annelida, the small **Enchytraeidae** (potworms), were shown to suffer adverse effects on
381 body weight and microbiome with PS microspheres (0.05 to 0.1 μm) at concentrations of
382 $\geq 10\%$ w/w within their food source, but an unexpected increase of reproduction at 0.5 % w/w
383 (Zhu et al., 2018b). The reproduction was reduced at abnormal concentrations of
384 90 g kg^{-1} dry soil of polyamid particles (13 to 150 μm), but not with PVC (Lahive et al., 2019).

385 The edaphon of semisubhydric soils is often treated as a marginal group between the area of
386 interest of soil and aquatic scientists. As a highly diverse soil biocenosis outside the focus of
387 this paper, the benthos along seashores and fresh waters is also affected by MPs and should
388 therefore be shortly mentioned by reviewing the lugworm **Arenicola marina**, a well examined
389 deposit-feeder of the tidal flats. In situ, MP accumulates within its tissue and feces (Van
390 Cauwenberghe et al., 2015). In laboratory experiments, PS particles $\geq 500\ \mu\text{m}$ were avoided
391 as food-source and passively translocated within the sediment at concentrations of $\sim 2\ \text{g}\ \text{kg}^{-1}$

392 (Gebhardt and Forster, 2018), but were measured within the feces at $\sim 74 \text{ g kg}^{-1}$ causing
393 effects on feeding activity and body weight with no influence on the survival rate (Besseling et
394 al., 2012). PS microspheres $\leq 30 \mu\text{m}$ remained within the animal without any adverse effects
395 regardless of particle size (Van Cauwenberghe et al., 2015). Other studies found adverse
396 effects on respiration, energy reserves, feeding, egestion and casting after uptake of PVC
397 particles $\leq 478 \mu\text{m}$ at different sediment concentrations of $> 2 \text{ g kg}^{-1}$, but neither on biomass
398 and survival nor due to HD-PE (Wright et al., 2013; Green et al., 2016). There is further a
399 difficulty in distinguishing between the adverse effects of MPs and substances adsorbed on or
400 leached from MPs (Besseling et al., 2012). When adding PCB-spiked PE to mud flat sediment
401 with concentrations up to **121 5000 mg kg⁻¹** dry mass, there was no significant change of
402 survival rate or body weight. The decreased feeding activity and heap mass could be
403 attributed to increasing plastic concentrations, but not to enhanced PCB bioaccumulation via
404 PE uptake (Besseling et al., 2017). However, all these studies found adverse effects at MP
405 concentrations orders of magnitude above natural values.

Table 4: Microplastic studies on Lumbricidae (p=particles, ms=microspheres, b=beads, f=fibers, ms=microfibers, [8] N/A=information not available). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
<i>Lumbricus terrestris</i>	mesocosm	PE	washed (C ₅ H ₁₂ , C ₈ H ₁₈)	N/A	N/A	p	<150	0..60% w/w (litter)	14 d / 60 d	yes	yes	N/A	at 60 d, ≥28% w/w: survival ↑, growth ↓	Huerta Lwanga et al. (2016)
<i>Eisenia fetida</i>	glass beaker	PS	N/A	N/A	N/A	ms	50..80	0..20000	30 d	N/A	yes	N/A	≥5000: survival ↓ ≥10000: weight ↓	Cao et al. (2017)
<i>Lumbricus terrestris</i>	bag	PE	N/A	N/A	N/A	p	0.92±1.09 mm ²	3500	28 d	N/A	yes	no	no	Hodson et al. (2017)
<i>Lumbricus terrestris</i>	home yard	diverse	yes	N/A	N/A	N/A	N/A	0.87±1.9 items g ⁻¹	N/A	N/A	yes	conc. in chickens > in earthworms	N/A	Huerta Lwanga et al. (2017a)
<i>Lumbricus terrestris</i>	mesocosm	PE	washed (C ₅ H ₁₂ , C ₈ H ₁₈)	N/A	N/A	p	<150	0..60% w/w (litter)	14 d	yes	yes	N/A	N/A	Huerta Lwanga et al. (2017b)
<i>Lumbricus terrestris</i>	mesocosm	PE	N/A	no	no	b	710..2800	750 µg on 2.5 kg soil	21 d	yes	yes	N/A	no	Rillig et al. (2017)
<i>Eisenia andrei</i>	mesocosm	LD-PE	N/A	N/A	N/A	pellets	250..1000	0..1000	28 d	N/A	yes	N/A	≥62.5: intestinal damage	Rodriguez-Seijo et al (2017)
<i>Lumbricus terrestris</i> (gut bacteria)	mesocosm glass bottle	PE	washed (C ₅ H ₁₂ , C ₈ H ₁₈)	N/A	N/A	p	150	7% w/w (litter) 10000	60 d (earthworms) 21 d (bacteria)	N/A	yes	N/A	N/A	Huerta Lwanga et al. (2018)
<i>Eisenia fetida</i>	mesocosm	LD-PE	washed (EtOH)	N/A	N/A	pellets	250..1000	0..1000	28 d	N/A	yes	N/A	≥125: altered enzyme activity	Rodriguez-Seijo et al (2018)
<i>Aporrectodea rosea</i>	mesocosm	PLA, PE N/A	N/A	N/A	N/A	p f	N/A	1000 10	30 d	N/A	yes	N/A	growth ↓	Boots et al. (2019)
<i>Eisenia fetida</i>	mesocosm	HD-PE, PET, PVC	no	N/A	no	f	<2000	soil extract	48 h / 56 d	N/A	N/A	N/A	no	Judy et al. (2019)
<i>Lumbricus terrestris</i>	bag	PE	N/A	N/A	N/A	mf	Ø40.7±3.8 x 361.6±387.0	0..10000	35 d	N/A	yes	N/A	≥1000: metallothionein expression ↑ ≥10000: heat shock protein 70 ↓	Prendergast-Miller et al. (2019)
<i>Eisenia fetida</i>	mesocosm	LD-PE	washed (EtOH)	N/A	chlorpyrifos (CPF)	pellets	5000 250..1000	40 items on 0.5 kg soil 180..200 items on 0.5 kg soil	14 d	N/A	N/A	N/A	with CPF: altered enzyme activity, avoidance of MPs	Rodriguez-Seijo et al (2019)
<i>Metaphire californica</i>	mesocosm	PVC	N/A	N/A	sodium arsenate	p	N/A	2000	28 d	yes	yes	N/A	microbiome	Wang et al. (2019b)
<i>Eisenia fetida</i>	glass beaker	PE PS	washed (MeOH)	N/A	PAHs, PCBs, Nile Red (NR)	p	<300 <250	0..200000 0..100	14 d 28 d	N/A	yes	N/A	≥200000: altered enzyme activity	Wang et al. (2019c)
<i>Lumbricus terrestris</i>	mesocosm	PE	washed (C ₅ H ₁₂ , C ₈ H ₁₈)	N/A	glyphosate	p	<150	0..7% w/w (litter)	14 d	N/A	N/A	N/A	N/A	Yang et al. (2019a)
<i>Lumbricus terrestris</i>	mesocosm	PE	N/A	N/A	N/A	N/A	<1000	7% w/w (litter)	14 d	yes	yes	N/A	N/A	Yu et al. (2019)
<i>Lumbricus terrestris</i>	Petri dish mesocosm	PE and div. biodegradables	unweathered, field or compost	N/A	N/A	p	1.5x1.5 cm ² 2x2 cm ²	4 items per dish 10 items per dish	14 d 50 d	yes yes	no yes	N/A	N/A	Zhang et al. (2018)
<i>Eisenia fetida</i>	bag	PE	washed (EtOH)	N/A	N/A	p	<400	0..1500	28 d	N/A	yes	yes	skin damage, ≥250 mg/kg: oxidative stress ≥1000 mg/kg: neurotoxicity ↑	Chen et al. (2020)

Table 5: Microplastic studies on Enchytraeidae and *Arenicola marina* (mb=microbeads, p=particles, ms=microspheres, sed.=sediment, ^s=semisubhydryc, [8] N/A=information not available). Concentrations refer to mg kg⁻¹ dry soil in terrestrial soils and mg kg⁻¹ dry sediment in semisubhydryc soils, if not specially marked.

	organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
Enchytr.	<i>Enchytraeus crypticus</i>	Petri dish	PS	N/A	N/A	N/A	mb	0.05..0.1	0..10% w/w (food)	7	N/A	yes	N/A	at 0.5% w/w: reproduction ↑ ≥10% w/w: microbiome, weight ↓	Zhu et al. (2018c)
	<i>Enchytraeus crypticus</i>	microcosm	PA PVC	N/A	N/A	fluorescence N/A	p	13..150 106..150	20000..120000 90000	20 h / 21 d	N/A	yes	N/A	≥90000: reproduction ↓ no	Lahive et al. (2018)
Arenicola	<i>Arenicola marina</i> ^s	in situ	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	yes	1.2±2.8 items g ⁻¹	N/A	Cauwenberghe et al. (2015)
		liquid culture	PS	no	N/A	N/A	ms	10..90	10000..50000 items kg ⁻¹	14 d	N/A	yes	10 µm: 9600±1800 items kg ⁻¹ 30 µm: 800±700 items kg ⁻¹	no	
	<i>Arenicola marina</i> ^s	mesocosm	PS PA	yes	biofilm	N/A	p	500..1000	~2000 ~1000	106..240 d	yes	no	N/A	N/A	Gebhardt and Forster (2018)
	<i>Arenicola marina</i> ^s	mesocosm	PS	N/A	N/A	N/A	p	400..1300	0..74000	28 d	N/A	≥400 µm	no	≥74000: feeding ↓, weight ↓	Besseling et al., (2012)
	<i>Arenicola marina</i> ^s	mesocosm	PVC HD-PE	N/A	N/A	N/A	p	9..478 3..316	0..20000 mg kg ⁻¹ wet sed.	31 d	N/A	N/A	N/A	>2000: respiration ↓, casting ↓ no	Green et al. (2016)
	<i>Arenicola marina</i> ^s	mesocosm	PE	N/A	PCBs	fluorescence	mb	10..180	0..5000	28 d	N/A	yes	no	feeding activity ↓, heap mass ↓	Besseling et al. (2017)
	<i>Arenicola marina</i> ^s	mesocosm	PVC	N/A	N/A	not leaching	p	~130	0..50000	28 d	N/A	N/A	N/A	≥10000: energy reserves ↓ ≥50000: feeding ↓, egestion ↓, casting ↓	Wright et al. (2013)

3.4 Further invertebrates

As part of the microfauna, the phylum **Nematoda** (nematodes or roundworms) is an ecologically important branch containing >25000 species (Zhang, 2013) in freshwater, marine, endobiotic and soil habitats. Due to their diverse trophic interactions nematodes hold a central position in both bottom-up and top-down controlled food webs (Yeates, 2001; Ferris, 2010) and thus most likely the uptake and transfer of MP.

Active feeding of adults and larvae of different species on 0.5 to 6 μm PS/latex microspheres (the size of their bacterial prey) was proven by Nika et al. (2016) and Fueser et al. (2019). However, most MP experiments on Nematodes are based on the bacterial-feeding model organism *Caenorhabditis elegans*. Kiyama et al. (2012) showed the favored uptake of PS microspheres with sizes of 0.5 to 3 μm by adult and 0.5 μm by larval *C. elegans*. The ingestion of MP decreased in the presence of bacteria as the natural food source.

When larval stages and adults ingested PS between 0.05 and 5 μm within an aqueous suspension or on agar plates, adverse effects such as oxidative stress, neurodegeneration, intestinal and DNA damage or dysfunction in motility, growth, life span, defecation, reproduction or energy metabolism appeared from a wide spectrum of concentrations from $\geq 1 \mu\text{g l}^{-1}$ up to $\geq 86.3 \text{ mg l}^{-1}$ (Zhao et al., 2017; Dong et al., 2018; Kim et al., 2019; Lei et al., 2018a; Lei et al., 2018b; Qu et al., 2019a). These effects are **not seen** below $1 \mu\text{g l}^{-1}$ (Qu et al., 2019b), and are enhanced due to amino modifications on micropsphere surfaces (Qu et al., 2019c). The incubation on agar plates with PE, PP and PVC particles $< 70 \mu\text{m}$ caused similar influences on survival, fertility, brood size and intestinal function (Lei et al., 2018b). Leachates from soils amended with 5 mg kg^{-1} dry soil of HD-PE and PVC decreased reproduction in laboratory cultures, but there was no effect shown on survival and after application of PET (Judy et al., 2019). Furthermore, silica nanoparticles (0.05 μm) are not only taken up orally but also via the vulva and spermathecae and migrate into gonad cells (Scharf et al., 2013). This process was confirmed for PS nanoparticles with the potential of a transfer to the progeny (Zhao et al., 2017).

The clear adverse effects of these studies are limited in their representativity by a narrow restriction to liquid cultures and a single model organism. Broader studies like on prominent soil-born nematodes such as *Acrobeloides buetschlii* (Frey, 1971) are still lacking. When assuming in first proximity mg l^{-1} solution = mg kg^{-1} dry soil, the applied concentrations between 0.001 and 86.8 mg l^{-1} match lower levels of soil contamination.

Feeding studies on the phylum **Rotifera** with MPs are fully based on PS microbeads and model organisms of the planktonic genus *Brachionus*. However, this data can carefully be transferred to soil environments as also soil rotifers are aquatic organisms living in water-filled pores and waterfilms. Different *Brachionus* sp. ingest microbeads $< 10 \mu\text{m}$ with strong preference for particles the size of their natural food source, namely bacteria and algae with

445 2 to 5 μm in diameter (Vadstein et al., 1993; Heerkloß and Hlawka, 1995; Baer et al., 2008;
446 Jeong et al., 2016). The uptake appears to be selective as microbeads are fewer incorporated
447 than bacteria and algae (Vadstein et al., 1993). The egestion of particles $\leq 0.5 \mu\text{m}$ is hindered
448 compared to $6 \mu\text{m}$ (Jeong et al., 2016). In suspension, microbeads $\leq 0.5 \mu\text{m}$ cause adverse
449 effects on fertility and life span at $\geq 0.1 \text{ mg l}^{-1}$ as well as oxidative stress and less growth at
450 $\geq 10 \text{ mg l}^{-1}$ (Jeong et al., 2016; Sun et al., 2019).

451 Terrestrial mollusks comprise snails and slugs within the class of **Gastropoda**. These grazers
452 feed on bacterial biofilms, fungi and plant tissue (Parkyn and Newell, 2013). Studies on
453 terrestrial species are sparse, but data on the benthic *Littorina sp.* imply passive transport and
454 non-selective MP uptake by feeding on surfaces with contaminated feces and mucus trails of
455 other snails (Gutow et al., 2019). With focus on benthic snails, Imhof and Laforsch (2016)
456 found no significant influence on growth parameters and fertility of juveniles and adult
457 *Potamopyrgus antipodarum* even when a food source with 70 % w/w of 5 to $600 \mu\text{m}$ sized
458 fragments was given (a mixture of PA, PC, PET, PS, PVC). In contrast, adverse effects were
459 found in recent work on the terrestrial snail *Achatina fulica*, that showed uptake and complete
460 gastrointestinal passage within 48 h with partial degradation of PET fibers (appr.
461 $1258 \times 76 \mu\text{m}$), but reduced excretion and food intake as well as increased oxidative stress at
462 concentrations of $\geq 0.01 \text{ g kg}^{-1}$, $\geq 0.14 \text{ g kg}^{-1}$ and $\geq 0.71 \text{ g kg}^{-1}$ dry soil, respectively (Song et al.,
463 2019).

Table 6: Microplastic studies on nematods (ms=microspheres, fr=fragments, np=nanoparticles, mb=microbeads, ms=microspheres, ox.=oxidative, [8] N/A=information not available). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

Microplastic studies on nematods (ms=microspheres, fr=fragments, np=nanoparticles, mb=microbeads, ms=microspheres, ox.=oxidative). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

	organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
	<i>Caenorhabditis elegans</i>	agar plate	PS	N/A	carboxyl sulfate amino	fluorescence	ms	0.1..6.6	N/A	0.5..2 h	N/A	yes	0.5..3 µm	N/A	Kiyama et al. (2012)
	<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	carboxyl	fluorescence	ms	0.1	0.001..10 mg l ⁻¹	4.5 d	N/A	Yes	N/A	≥0.01 mg l ⁻¹ : motility ↓, growth ↓, defecation ↓, within gonads	Zhao et al. (2017)
	<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	ζ=-10mV	fluorescence	ms	0.1	0.00001..0.001 mg l ⁻¹	N/A	N/A	Yes	N/A	≥0.001 mg l ⁻¹ : motility ↓, ox. stress ↑	Dong et al. (2018)
	<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	N/A	preservatives, fluorescence	ms	0.05..0.2	0.001..86.8 mg l ⁻¹ 17.3..86.8 mg l ⁻¹	24 h	N/A	Yes	N/A	≥17.3 mg l ⁻¹ : motility ↓, reproduction ↓ ≥86.3 mg l ⁻¹ : ox. stress ↑ ≥17.3 mg l ⁻¹ : metabolic dysf.	Kim et al. (2019)
	<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	ζ=-10mV	fluorescence	ms	0.1	0.001..1 mg l ⁻¹	N/A	N/A	Yes	N/A	≥1 mg l ⁻¹ : neurodegeneration ≥0.01 mg l ⁻¹ : motility ↓	Qu et al. (2019a)
	<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	N/A	N/A	ms	0.1..5	1 mg l ⁻¹	3 d	N/A	Yes	N/A	motility ↓, survival ↓, growth ↓, ox. stress ↑, neurotoxicity	Lei et al. (2018a)
Nematoda	<i>Caenorhabditis elegans</i>	agar plate	PE, PP, PVC, PS	no	N/A	N/A	fr, ms	0.1..200	0.5..10.0 mg m ⁻²	2 d	N/A	Yes	N/A	≥0.5 mg m ⁻² : survival ↓ at 5 mg m ⁻² : growth ↓, reproduction ↓, ox. stress ↑, intestinal damage	Lei et al. (2018b)
	<i>Caenorhabditis elegans</i>	agar plate	PS	N/A	N/A	fluorescence	ms	0.1..5	2500 mg l ⁻¹	7 d	N/A	Yes	N/A	mainly 1µm: intestinal damage	Scharf et al. (2013)
	<i>Caenorhabditis elegans</i>	liquid culture	HD-PE, PET, PVC	no	N/A	no	fr	<2000	soil extract	72 h	N/A	N/A	N/A	within tissue and gonads	Judy et al. (2019)
	<i>Caenorhabditis elegans</i>	agar plates	latex	N/A	N/A	fluorescence	mb	0.5	N/A	30 min	N/A	yes	N/A	reproduction ↓	Nika et al. (2016)
	<i>Caenorhabditis elegans</i> <i>Panagrolaimus thienemanni</i> <i>Plectus acuminatus</i> <i>Poikilolaimus regenfussi</i> <i>Acrobeloides nanus</i> <i>Pristionchus pacificus</i> <i>Aphelenchoides parietinus</i>	liquid culture	PS	N/A	N/A	fluorescence	ms	0.5..6	3·10 ⁹ ..10 ¹⁰ items l ⁻¹ (~0.2..1200 mg l ⁻¹)	4..73 h	N/A	≤3µm ≤0.5µm ≤1µm ≤1µm ≤1µm ≤6µm no	N/A	N/A	Fueser et al. (2019)
	<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	N/A	N/A	ms	0.1	0.0001..0.001 mg l ⁻¹	N/A	N/A	N/A	N/A	no	Qu et al. (2019b)
	<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	no amino	N/A	ms	0.1	0.001..1 mg l ⁻¹	N/A	N/A	yes	N/A	≥0.01 mg l ⁻¹ : reproduction ↓, DNA damage ≥0.001 mg l ⁻¹ : reproduction ↓, DNA damage	Qu et al. (2019c)

Table 7: Microplastic studies on Rotifera and Gastropoda (ms=microspheres, mb=microbeads, fr=fragments, f=fibers, ox.=oxidative, pref.=preferential, ^p=planctic, ^b=benthic, [8] N/A=information not available). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

	organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
Rotifera	<i>Brachionus plicatilis</i> ^p	liquid culture	PS	N/A	carboxyl	fluorescence	ms	1.6..20	5·10 ⁹ µm ³ l ⁻¹ (~5.25 mg l ⁻¹)	35 min	N/A	≤10 µm	pref. 4.5 µm	N/A	Bear et al. (2008)
	<i>Brachionus plicatilis</i> ^p	liquid culture	latex	N/A	N/A	fluorescence	mb	0.3..3.1	3·10 ⁷ ..7·10 ⁸ items l ⁻¹ (~0.0004..11 mg l ⁻¹)	20 min	N/A	yes	pref. ≥2 µm	N/A	Vadstein et al. (1993)
	<i>Brachionus koreanus</i> ^p	liquid culture	PS	no	N/A	fluorescence	mb	0.05..6	0...20 mg l ⁻¹	1 d	N/A	yes	egestion rate 0.05 µm < 0.5 µm < 6 µm	≤0.5 µm, ≥0.1 mg l ⁻¹ : reproduction ↓, survival ↓ ≤0.5 µm, 10 mg l ⁻¹ : oxidative stress ↑	Jeong et al. (2016)
	<i>Brachionus plicatilis</i> ^p	liquid culture	PS	N/A	N/A	N/A	mb	0.07..7	0..20 mg l ⁻¹	N/A	N/A	yes	N/A	≤0.07 µm, ≥10 mg l ⁻¹ : reproduction ↓, growth ↓ ≤0.07 µm and ≥0.1 mg l ⁻¹ : survival ↓	Sun et al. (2019)
	<i>Brachionus quadridentatus</i> ^p <i>Brachionus plicatilis</i> ^p	liquid culture	PS	N/A	N/A	N/A	ms	2..10	N/A	8..10 d	N/A	pref. 3..5 µm pref. 2 µm	N/A	N/A	Heerkloß and Hlawa (1993)
Gastropoda	<i>Littorina littorea</i> ^b	microcosm	PMMA	N/A	N/A	fluorescence	fr	10..100	increasing	16 h	N/A	yes	N/A	N/A	Gutow et al. (2019)
	<i>Potamopyrgus antipodarum</i> ^b	aquarium	PET, PS, PVC, PA, PC	N/A	N/A	no	fr	5..600	0..70% w/w (food)	≤141 d	N/A	yes	N/A	no	Imhof and Laforsch (2016)
	<i>Achatina fulica</i>	mesocosm	PET	N/A	N/A	no / stained	f	approx. 1258x76 µm	10..710	28 d	N/A	yes	excretion after 48 hours	≥140: food intake ↓ ≥10: excretion ↓ ≥710: ox. stress ↑, gastrointestinal damage	Song et al. (2019)

466 3.5 Vertebrates

467 Different taxa of the class of Amphibia have a predator function within the edaphic food web
468 (e.g. preying on invertebrates) (Hebrard et al., 1992). While no data on the reaction to soil
469 MPs are available neither for the legless **Gymnophiona** nor for adults of the order **Anura**,
470 sparse data on tadpoles of aquatic frogs suggest uptake followed by regular excretion of PS
471 microspheres as shown with *Xenopus tropicalis* (Hu et al., 2016). Further, there exist no data
472 on the families **Serpentes** (snakes) and **Anguillidae** within the class of Reptilia, residing at the
473 outer rim of the food web.

474 Within the broad field of Mammalia, studies on MP ingestion are sparse and focus on **mice** as
475 a rodent model organism. Feeding of mice with PS microspheres of 1 to 14 μm in
476 concentrations of 1.49×10^6 to 4.55×10^7 particles at a volume of 10 ml kg^{-1} body weight for
477 4 weeks showed no adverse effects (Stock et al., 2019). In contrast, longer exposition
478 (6 weeks) with lower concentrations of particles with the same shape and size range changed
479 the mouse microbiome and caused metabolic and intestinal dysfunction (Lu et al., 2018; Jin et
480 al., 2019), which comes along with bioaccumulation within organs (Yang et al., 2019b). These
481 studies are regularly conducted with passive feeding and exclude active foraging on
482 perceptible plastic particles. However, the uptake via prey or feeding on contaminated roots
483 and litter is highly probable. Further Rodentia – **Cricetidae** (hamsters, lemmings, voles),
484 **Bathyergidae** (blesmols, mole-rats), **Octodontidae** as well as **Spermophilus** (ground
485 squirrels) and **Marmota** (marmots) within the family of **Sciuridae** – were not yet studied, just
486 as other mammalian (sub)orders like **Chrysochloridae** (golden moles), **Cingulata**
487 (armadillos), **Macroscelidea** (elephant shrews), **Notoryctemorphia** and **Peramelemorphia**.

Table 8: [3] Microplastic studies on Anura (An.) and Rodentia (ms=microspheres, ^a=aquatic, [8] N/A=information not available).

	organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
An.	<i>Xenopus tropicalis</i> ^a	Petri dish	PS	N/A	N/A	fluorescence	ms	1..10	100..10 ⁸ items l ⁻¹ (55·10 ⁻⁹ ..55 mg l ⁻¹)	48 h	N/A	yes	egestion within days	N/A	Hu et al. (2016)
					carboxyl			1	4.55·10 ⁷ items per mouse (0.025 mg per mouse)						
	transgenic mice	in vivo	PS	N/A	sulfate	fluorescence	ms	4	4.55·10 ⁷ items per mouse (1.6 mg per mouse)	28 d	N/A	yes	N/A	no	Stock et al. (2019)
					sulfate			10	1.49·10 ⁶ items per mouse (0.8 mg per mouse)						
Rodentia	mice	in vivo	PS	N/A	N/A	fluorescence	ms	5	0.1..1 mg l ⁻¹ (food)	42 d	N/A	yes	N/A	≥0.1 mg l ⁻¹ : microbiome, metabolic dysfunction	Jin et al. (2019)
	mice	in vivo	PS	N/A	N/A	N/A	ms	0.5..50	0.1..1 mg l ⁻¹ (food)	35 d	N/A	N/A	N/A	≥0.1 mg l ⁻¹ : microbiome, metabolic dysfunction	Lu et al. (2018)
	<i>Mus musculus</i>	in vivo	PS	N/A	N/A	fluorescence	ms	5..20	200 mg l ⁻¹ (food)	28 d	N/A	yes	8x, 8±5 and 0.71±0.14 mg kg ⁻¹ body weight	≥1 mg l ⁻¹ : body weight ↓	Yang et al. (2019b)

489 4 Synthesis

490 4.1 Summarized observations

491 Our systematic search comprised recent research on the interaction of soil organisms with
492 MP, but also studies with focus on feeding experiments, that are published much earlier than
493 the awareness on plastic in the environment appeared. The numerous studies found with
494 focus on the ingestion of MPs consistently showed the active uptake by diverse soil
495 organisms with few exceptions spread over the whole branch of invertebrates. In addition,
496 also studies on adverse effects caused by the intake of MP contaminated food (e.g. of food
497 pellets by dung beetles) imply the ingestion into the test organism. Distinct size preferences
498 are [12] observed in dung beetles, nematodes, rotifers and ants showing that mainly particles
499 are ingested, that are small enough to enter the gastrointestinal tract. In contrast, active
500 comminution by gnawing on larger particles was tested only for a few taxa and confirmed for
501 woodlice, termites and mealworms, and in the case of earthworms only after initial
502 weathering.

503 After the ingestion, MP is [12] translocated actively until excretion or death of the transporting
504 organism, which was only directly shown in experiments with earthworms. The passive
505 transport by attachment, dragging and pushing was [12] investigated in a few experiments with
506 earthworms, mites and springtails that partly worked without soil substrate and consistently
507 showed positive results.

508 After exposition to MP, a pattern of adverse effects can be seen: Across various taxa, altered
509 microbiomes, reduced motility, body mass, fertility and life span as well as increased oxidative
510 stress and metabolic malfunctioning occur in different combinations mainly due to μm -sized
511 MP in and above the whole known natural range of [12] concentrations. For some taxa such as
512 Nematodes, Gastropoda and Rotifera these effects appear at natural and increased MP
513 concentrations ($<100 \text{ mg kg}^{-1}$ dry soil), for Collembola and Lumbricidae at concentrations like
514 in highly contaminated sites ($\geq 1000 \text{ mg kg}^{-1}$ dry soil) and for Enchytraeidae, *Arenicola marina*
515 and in further experiments with earthworms at implausibly high values. The data show a
516 tendency, that the effects occur at lower concentrations, when the added particles are smaller.
517 Small sized particles also provide the highest surface/volume ratio and thus the highest
518 reactive surface per weight.

519 Most studies work with defined increasing MP concentrations and particle sizes in soil
520 substrates and food sources, which can be used to determine relationships between
521 environmental concentrations and adverse effects. However, the lack of information about
522 intake rates, grades of accumulation and effective prey-predator transfer leads to a gap within
523 the chain of explanation for toxic effects on the soil organisms. In some experiments, the
524 intestinal passage of MP and sizes [4] preferably retained within the gut were shown, but there
525 are no experiments that could demonstrate quantitative bioaccumulation. In contrast,

526 quantification of the retained and egested MP particle size fractions might be biased due to
527 gnawing and intestinal comminution as shown for woodlice, termites, mealworms, snails and
528 earthworms.

529 In order to improve our understanding of processes underlying adverse effects of MP on soil
530 organisms, data on ingestion rates, dwell times, biodegradation and egestion rates are
531 important bricks e.g. to reveal bioaccumulation dynamics. However, there are only a few data
532 on biodegradation (mealworms, snails, earthworms), egestion (rotifers, frogs, snails,
533 earthworms) and remaining concentrations in the body (lugworm, mice, earthworms).

534 4.2 Limitations of previous studies

535 The available studies worked with items within the full size span of micro- and nanoplastics
536 ($\leq 5000 \mu\text{m}$). [71] Approximately 72 % of the experiments used microplastic (0.1 to $5000 \mu\text{m}$),
537 only 6 % nanoplastic ($< 0.1 \mu\text{m}$), 10 % included macroplastic ($> 5000 \mu\text{m}$) and 12 % used
538 microplastic of undefined size. When $\text{MP} \geq 50 \mu\text{m}$ was applied, mainly particles and fragments
539 made of PE and PVC were used, whereas PS/latex microspheres were mainly applied for
540 sizes $\leq 10 \mu\text{m}$ ([71] Table 1). The latter are readily available, highly standardized and are mostly
541 used with fluorescent dyes and either without additional functional groups, carboxylated or,
542 more rarely, with amino or sulfate groups. However, there are indications that the spectrum of
543 particle type and shape used in experiments does not correspond to the properties of
544 particles in soils. In different natural as well as agriculturally and industrially contaminated
545 terrestrial and semisubhydric sites, fibers and fragments of PE and PP, mostly $\leq 100 \mu\text{m}$, were
546 much more abundant than PVC, PET and PS items (Claessens et al., 2011; Vianello et al.,
547 2013; Nor and Obbard, 2014; Naji et al., 2017; Zhang and Liu, 2018; Li et al., 2018a). This is
548 probably caused by high loads of MP fibers in discharged waste water and sewage sludge,
549 which is used in agricultural sites worldwide (Mahon et al., 2016; Li et al., 2018b). It is likely
550 that shape plays an important role for the ingestion of MP items. Unfortunately, we did not find
551 studies that have carried out a complete classification of sampling sites according to plastic
552 origin, size and type, that could help to evaluate differences between former experimental and
553 natural plastic composition to achieve the most realistic experimental conditions. Little
554 knowledge about the size distribution of MP in soils furthermore complicates the
555 determination of realistic concentrations for the addition of a certain particle size spectrum. All
556 reviewed studies either arbitrarily set their applied concentrations or had to base them on
557 measurements of total specific MP masses, regardless of how much of this mass is in the
558 tested size range. This may lead to a false estimation of total adverse MP concentrations.

559 In contrast to particle type and shape, the documentation of chemical properties of MP
560 samples in most of these studies is fragmentary. Some experiments explicitly mentioned that
561 the added plastic was unweathered, whereas most studies lack information about the degree
562 of aging implying that unweathered items were used. Only a few experiments involved aging
563 of MP, but without comparison to results of natural weathering (Tsunoda et al., 2010;
564 Gebhardt and Forster, 2018). That is in conflict with natural conditions, as plastic that [12]
565 remains within the soil after littering, sewage sludge application or plastic mulching shows
566 signs of weathering, e.g. modified carbonyl indices (Andrady, 2017), while unweathered soil
567 MP might be rare. In addition, Zhang et al. (2018) showed that earthworms actively
568 comminute only weathered bioplastics. [6] In experiments using PS microspheres,
569 carboxylation is often used to imitate a reduced hydrophobicity due to weathering. However,
570 according to manufacturer information microplastics only have little influence on
571 hydrophobicity.

572 Weathering of MP surfaces within soils comes along with biofilm growth and adsorption of
573 organic molecules, which could potentially affect the attractiveness or toxicity for grazers and
574 other organisms. Such coatings were applied only in a few cases (Besseling et al., 2017;
575 Angotti et al., 2018; Gebhardt and Forster, 2018), but were not documented in most studies.
576 Similarly, the type and concentration of additives such as flame retardants, anti-oxidants or
577 stabilizers often remained undocumented, with exception of fluorescent dyes, that are well
578 mentioned. The release of additives can have a harmful effect on the test organism, as shown
579 for aquatic environments (e Silva et al., 2016). Some studies on the ingestion of MP by the
580 soil mesofauna indicate that the diameter of the gastrointestinal tract is a useful upper size
581 limit for added particles, as far as the organism is unable to crush them (Heerkloß and Hlawa,
582 1995; Holter, 2000; Holter et al., 2002; Holter and Scholtz, 2005; Baer et al., 2008; Fueser et
583 al., 2019). However, using only ingestible particle sizes in their natural concentrations neglect
584 the adverse effects of plastic leachates, which can also get into the soil solution and onto the
585 mineral phase from larger particles and affect soil life.

586 The conditions of incubation differ considerably in terms of habitats and duration of exposure.
587 In most studies, the exposure ranges from a few minutes to a few days in experiments with
588 micro- and small mesofauna and hours to several weeks in experiments with large meso- and
589 macrofauna and is mainly based on excretion or reproductive cycles. Long-term studies,
590 which are indeed difficult to carry out in mesocosms, practically do not exist. However, certain
591 adverse effects might only establish themselves after long term trials, as was shown for the
592 influence of pesticides (Pelosi et al., 2015).

593 Some experiments were carried out in soil-free test environments such as liquid cultures or
594 Petri dishes with nutrient solutions or a specific food source (nematods, rotifers, mice).
595 Therefore, motivity is less restricted and feeding behavior can be altered compared to
596 cultivation within soil environments. For example, the ingestion of MP by nematodes
597 decreases in the presence of an alternative and more natural food source like bacteria, which
598 can significantly reduce the bioaccumulation and thus the effective toxicity (Kiyama et al.,
599 2012). This can lead to less consumption of MP in soil environments and an overestimation of
600 the toxicity in liquid culture experiments. Also, all laboratory feeding experiments were carried
601 out by use of only one species. The complexity of the food web in soils is thereby excluded
602 and the potential accumulation from prey to predators still unexplored.

603 4.3 Pinpoints for future research

604 Most studies reviewed in this work have a pioneering role in MP research and, thus, are
605 subject to some experimental limitations caused by an early state of knowledge. The adverse
606 effects recently found are alarming, but must be considered under the restrictions named
607 above. We propose the following points as part of a *modus operandi* for future MP research.

608 In past studies, particular adverse effects of MP were measured only for certain sizes,
609 shapes, coatings, leachates or adsorbed substances (*Tables 2 to 8*). Experimental
610 concentrations were assumed randomly or derived from cumulative concentrations of one or
611 more MP types measured in natural soils (approx. 1 to some 1000 mg kg⁻¹ dry soil),
612 regardless of size. For those specific experiments coming, the spectrum of concentrations
613 used should be adapted to the quantities of the size spectrum, that occurs within the soil. For
614 future studies on mixed contaminations, we recommend to evaluate the overall adverse
615 effects of PE, PP, PVC, PET, PU and PS to certain test organisms by use of typical MP-
616 specific spans of concentration, size and shape distribution in natural soils or food samples.
617 This previously requires well-structured data of appropriate MP type, shape and size for
618 different soils in differently contaminated areas.

619 Experiments on adverse effects should be applied within soil matrices to allow the interplay of
620 plastic, natural organic and mineral matter. The MP should be weathered, as plastic in soils
621 underlie broad environmental aging. Pre-weathering of MP should therefor not only be
622 performed in climate chambers (e.g. following DIN EN ISO 4892-2/3), but also include
623 subsequent leaching and equilibration of additives or coatings within the soil matrix before the
624 main experiment. Furthermore, the experimental design may consider coatings with biofilms
625 or attractants and even particle color to regulate the preference of the test organisms.

626 Most detailed information about ingestion are available for dung beetles, nematods and
627 earthworms, data on adverse effects on nematods, earthworms, lugworms and collembola.
628 Future experiments should focus on **a larger variety of ecologically relevant taxa like**
629 **Coleoptera, Formicidae, Acari, Oniscidea, Collembola, Lumbricidae, Enchytraeidae,**
630 **Nematoda and Gastropoda.** The studies are recommended to conduct with emphasis on
631 uptake, accumulation and key adverse effects like on survival rate, motility, growth and fertility
632 as well as on the stability of the intestinal microbiome. Further studies with more than one test
633 organism are important to foster our understanding of MP within certain food chains. Also
634 long-term experiments might reveal adverse effects, which evolve slowly within populations.
635 This may enable the **assessment** of the distribution and effects of MP within the food web and
636 the resulting long-term impact on soil ecosystems.

637 5 Conclusion

638 Our review of 77 studies on the impact of microplastic on the soil fauna shows a [10]
639 considerable diversity and distribution of adverse effects within the soil tree of life. However,
640 these effects have to be considered carefully, as many experiments [12] did not use plastic
641 matching properties within natural soils and found adverse effects only at concentrations like
642 in highly contaminated soils or above. To elucidate effective concentrations [11] and properties
643 for short and long-term effects on soil faunal health, the most exact reproduction of plastic
644 properties within the soil matrix and natural living conditions of the test organisms is
645 necessary [11] together with a better knowledge on common concentrations and size
646 distributions of soil microplastic. For future experiments we therefore recommend to choose
647 compositions of type, shape, size, [11] concentration, grade of weathering, leachability and
648 coating with biofilms and other organic matter as expected in the habitat to be examined.
649 Furthermore, coming studies should include long-term exposure and food chain experiments
650 to get a better look at the effect of even smaller MP concentrations and their enrichment
651 within the food web. This may give us a better way of assessing the impact of global
652 microplastic contamination on e.g. soil biodiversity, soil carbon cycles and soil quality.

653 **Author contribution**

654 Frederick Büks developed the review concept, collected data and prepared the manuscript
655 except for earthworms. Nicolette Loes van Schaik did all the work on earthworms. Martin
656 Kaupenjohann supervised the study **by participating in structural discussions on the idea**
657 **and concept of the paper as well as the final corrections.**

658 The authors declare that they have no conflict of interest.

659

660

661 **Acknowledgement**

662 Many thanks to Ivica Letunic, who kindly gave us access to phyloT and made our work in face
663 of New Year much easier. I will not forget to invite you for a coffee.

664

665

666 **Competing interests**

667 The authors declare that they have no conflict of interest.

668 **References**

- 669 Allasino M. L., Marrero H. J., Dorado J. and Torretta J. P.: Scientific note: first global report of
670 a bee nest built only with plastic, *Apidologie*, 50, 230-233, doi:10.1007/s13592-019-00635-6,
671 2019.
- 672 Andrady A. L.: The plastic in microplastics: a review, *Marine Pollution Bulletin*, 119, 12-22,
673 doi:10.1016/j.marpolbul.2017.01.082, 2017.
- 674 Angotti M. A., Rabello A., Santiago G. and Ribas C.: Seed removal by ants in Brazilian
675 savanna: optimizing fieldwork, *Sociobiology*, 65, 155-161,
676 doi:10.13102/sociobiology.v65i2.1938, 2018.
- 677 Avery S. V., Harwood J. L. and Lloyd D.: Quantification and Characterization of Phagocytosis
678 in the Soil Amoeba *Acanthamoeba castellanii* by Flow Cytometry, *Appl. Environ. Microbiol.*,
679 61, 1124-1132, doi:N/A, 1995.
- 680 Baer A., Langdon C., Mills S., Schulz C. and Hamre K.: Particle size preference, gut filling and
681 evacuation rates of the rotifer *Brachionus* "Cayman" using polystyrene latex beads,
682 *Aquaculture*, 282, 75-82, doi:10.1016/j.aquaculture.2008.06.020, 2008.
- 683 Bertling J., Bertling R. and Hamann L.: Kunststoffe in Der Umwelt: Mikro- und Makroplastik.
684 Ursachen, Mengen, Umweltschicksale, Wirkungen, Lösungsansätze, Empfehlungen,
685 doi:10.24406/UMSICHT-N-497117, 2018.
- 686 Besseling E., Foekema E. M., van den Heuvel-Greve M. J. and Koelmans A. A.: The effect of
687 microplastic on the uptake of chemicals by the lugworm *Arenicola marina* (L.) under
688 environmentally relevant exposure conditions, *Environmental Science & Technology*, 51,
689 8795-8804, doi:10.1021/acs.est.7b02286, 2017.
- 690 Besseling E., Wegner A., Foekema E. M., Van Den Heuvel-Greve M. J. and Koelmans A. A.:
691 Effects of microplastic on fitness and PCB bioaccumulation by the lugworm *Arenicola marina*
692 (L.), *Environmental Science & Technology*, 47, 593-600, doi:10.1021/es302763x, 2012.
- 693 Boots B., Russell C. W. and Green D. S.: Effects of Microplastics in Soil Ecosystems: Above
694 and Below Ground, *Environmental Science & Technology*, 53, 11496-11506,
695 doi:10.1021/acs.est.9b03304, 2019.
- 696 Bowers B. and Olszewski T. E.: *Acanthamoeba* discriminates internally between digestible
697 and indigestible particles, *The Journal of Cell Biology*, 97, 317-322, doi:10.1083/jcb.97.2.317,
698 1983.
- 699 Brandon A. M., Gao S.-H., Tian R., Ning D., Yang S.-S., Zhou J., Wu W.-M. and Criddle C. S.:
700 Biodegradation of polyethylene and plastic mixtures in mealworms (larvae of *Tenebrio molitor*)

701 and effects on the gut microbiome, *Environmental Science & Technology*, 52, 6526-6533,
702 doi:10.1021/acs.est.8b02301, 2018.

703 Cao D., Wang X., Luo X., Liu G. and Zheng H.: Effects of polystyrene microplastics on the
704 fitness of earthworms in an agricultural soil, In: *IOP Conference Series: Earth and*
705 *Environmental Science*, doi:10.1088/1755-1315/61/1/012148, 2017.

706 Chae Y. and An Y.-J.: Current research trends on plastic pollution and ecological impacts on
707 the soil ecosystem: A review, *Environmental Pollution*, 240, 387-395,
708 doi:10.1016/j.envpol.2018.05.008, 2018.

709 Chae Y., Kim D., Kim S. W. and An Y.-J.: Trophic transfer and individual impact of nano-sized
710 polystyrene in a four-species freshwater food chain, *Scientific Reports*, 8, 284,
711 doi:10.1038/s41598-017-18849-y, 2018.

712 Chen Y., Liu X., Leng Y. and Wang J.: Defense responses in earthworms (*Eisenia fetida*)
713 exposed to low-density polyethylene microplastics in soils, *Ecotoxicology and Environmental*
714 *Safety*, 187, 109788, doi:10.1016/j.ecoenv.2019.109788, 2020.

715 Claessens M., De Meester S., Van Landuyt L., De Clerck K. and Janssen C. R.: Occurrence
716 and distribution of microplastics in marine sediments along the Belgian coast, *Marine*
717 *Pollution Bulletin*, 62, 2199-2204, doi:10.1016/j.marpolbul.2011.06.030, 2011.

718 Darwin C.: The formation of vegetable mould through the action of worms, with observations
719 on their habits, John Murray, London, 1881.

720 De Bruyn L. and Conacher A. J.: The role of termites and ants in soil modification-a review,
721 *Soil Research*, 28, 55-93, doi:10.1071/SR9900055, 1990.

722 Dong S., Qu M., Rui Q. and Wang D.: Combinational effect of titanium dioxide nanoparticles
723 and nanopolystyrene particles at environmentally relevant concentrations on nematode
724 *Caenorhabditis elegans*, *Ecotoxicology and Environmental Safety*, 161, 444-450, doi:10.1016/
725 j.ecoenv.2018.06.021, 2018.

726 Elloway E. A., Bird R. A., Hewitt C. J., Kelly S. L. and Smith S. N.: Characterization of
727 *Acanthamoeba*-microsphere association by multiparameter flow cytometry and confocal
728 microscopy, *Cytometry Part A*, 69, 266-272, doi:10.1002/cyto.a.20210, 2006.

729 Fenchel T.: Suspension feeding in ciliated protozoa: functional response and particle size
730 selection, *Microbial Ecology*, 6, 1-11, doi:10.1007/BF02020370, 1980.

731 Ferris H.: Contribution of nematodes to the structure and function of the soil food web, *Journal*
732 *of Nematology*, 42, 63, doi:N/A, 2010.

733 Frey F.: The suitability of *Acrobeloides buetschlii* for nematological experiments,
734 *Nematologica*, 17, 474-477, doi:N/A, 1971.

735 Fründ H.-C., Graefe U. and Tischer S.: Earthworms as bioindicators of soil quality, In: Biology
736 of earthworms, Springer, doi:10.1007/978-3-642-14636-7_16, 2011.

737 Fueser H., Mueller M.-T., Weiss L., Höss S. and Traunspurger W.: Ingestion of microplastics
738 by nematodes depends on feeding strategy and buccal cavity size, Environmental Pollution,
739 255, 113227, doi:10.1016/j.envpol.2019.113227, 2019.

740 Fuller S. and Gautam A.: A procedure for measuring microplastics using pressurized fluid
741 extraction, Environmental Science & Technology, 50, 5774-5780,
742 doi:10.1021/acs.est.6b00816, 2016.

743 Garcés-Ordóñez O., Castillo-Olaya V. A., Granados-Briceño A. F., Garcéa L. M. B. and Díaz
744 L. F. E.: Marine litter and microplastic pollution on mangrove soils of the Ciénaga Grande de
745 Santa Marta, Colombian Caribbean, Marine Pollution Bulletin, 145, 455-462,
746 doi:10.1016/j.marpolbul.2019.06.058, 2019.

747 Gebhardt C. and Forster S.: Size-selective feeding of *Arenicola marina* promotes long-term
748 burial of microplastic particles in marine sediments, Environmental Pollution, 242, 1777-1786,
749 doi:10.1016/j.envpol.2018.07.090, 2018.

750 van Gestel C. A., Loureiro S. and others: Terrestrial isopods as model organisms in soil
751 ecotoxicology: a review, ZooKeys, 127, doi:10.3897/zookeys.801.21970, 2018.

752 van Gestel C. A. and Selonon S.: Ecotoxicological effects of microplastics in soil: Comments
753 on the paper by Zhu et al.(2018) 'Exposure of soil collembolans to microplastics perturbs their
754 gut microbiota and alters their isotopic composition', Soil Biology & Biochemistry 116,
755 302-310, Soil Biology and Biochemistry, 124, 116-117, doi:10.1016/j.soilbio.2018.05.032,
756 2018.

757 Geyer R., Jambeck J. R. and Law K. L.: Production, use, and fate of all plastics ever made,
758 Science Advances, 3, e1700782, doi:10.1126/sciadv.1700782, 2017.

759 Glancey B. M., Vander Meer R., Glover A., Lofgren C. and Vinson S.: Filtration of
760 microparticles from liquids ingested by the red imported fire ant *Solenopsis invicta* Buren,
761 Insectes Sociaux, 28, 395-401, doi:10.1007/BF02224196, 1981.

762 Green D. S., Boots B., Sigwart J., Jiang S. and Rocha C.: Effects of conventional and
763 biodegradable microplastics on a marine ecosystem engineer (*Arenicola marina*) and
764 sediment nutrient cycling, Environmental Pollution, 208, 426-434,
765 doi:10.1016/j.envpol.2015.10.010, 2016.

766 Gulvik M.: Mites (Acari) as indicators of soil biodiversity and land use monitoring: a review,
767 Polish Journal of Ecology, 55, 415-440, doi:N/A, 2007.

768 Gusmão F., Di Domenico M., Amaral A. C. Z., Martínez A., Gonzalez B. C., Worsaae K., do
769 Sul J. A. I. and da Cunha Lana P.: In situ ingestion of microfibres by meiofauna from sandy
770 beaches, *Environmental Pollution*, 216, 584-590, doi:10.1016/j.envpol.2016.06.015, 2016.

771 Gutow L., Bartl K., Saborowski R. and Beermann J.: Gastropod pedal mucus retains
772 microplastics and promotes the uptake of particles by marine periwinkles, *Environmental*
773 *Pollution*, 246, 688-696, doi:10.1016/j.envpol.2018.12.097, 2019.

774 Hassall M. and Rushton S.: Feeding behaviour of terrestrial isopods in relation to plant
775 defences and microbial activity, In: *Symposia of the Zoological Society of London*, doi:N/A,
776 1984.

777 Hebrard J. J., Maloiy G. M. and Alliangana D. M.: Notes on the habitat and diet of *Afrocaecilia*
778 *taimana* (Amphibia: Gymnophiona), *Journal of Herpetology*, 513-515, doi:10.2307/1565136,
779 1992.

780 Heerkloß R. and Hlawa S.: Feeding biology of two brachionid rotifers: *Brachionus*
781 *quadridentatus* and *Brachionus plicatilis*, *Hydrobiologia*, 313, 219-221,
782 doi:10.1007/BF00025954, 1995.

783 Hodson M. E., Duffus-Hodson C. A., Clark A., Prendergast-Miller M. T. and Thorpe K. L.:
784 Plastic bag derived-microplastics as a vector for metal exposure in terrestrial invertebrates,
785 *Environmental Science & Technology*, 51, 4714-4721, doi:10.1021/acs.est.7b00635, 2017.

786 Holter P.: Particle feeding in *Aphodius* dung beetles (Scarabaeidae): old hypotheses and new
787 experimental evidence, *Functional Ecology*, 631-637, doi:N/A, 2000.

788 Holter P. and Scholtz C.: Are ball-rolling (Scarabaeini, Gymnopleurini, Sisyphini) and
789 tunnelling scarabaeine dung beetles equally choosy about the size of ingested dung
790 particles?, *Ecological Entomology*, 30, 700-705, doi:10.1111/j.0307-6946.2005.00746.x, 2005.

791 Holter P., Scholtz C. and Wardhaugh K.: Dung feeding in adult scarabaeines (tunnellers and
792 endocoprids): even large dung beetles eat small particles, *Ecological Entomology*, 27, 169-
793 176, doi:10.1046/j.1365-2311.2002.00399.x, 2002.

794 Hopkin S. P.: *Biology of the springtails (Insecta: Collembola)*, OUP Oxford, 1997.

795 Hu L., Su L., Xue Y., Mu J., Zhu J., Xu J. and Shi H.: Uptake, accumulation and elimination of
796 polystyrene microspheres in tadpoles of *Xenopus tropicalis*, *Chemosphere*, 164, 611-617,
797 doi:https://doi.org/10.1016/j.chemosphere.2016.09.002, 2016.

798 Huang, Y., Liu, Q., Jia, W., Yan, C. and Wang, J.: Agricultural plastic mulching as a source of
799 microplastics in the terrestrial environment. *Environmental Pollution*, 260, 114096,
800 doi: 10.1016/j.envpol.2020.114096, 2020.

801 Huerta Lwanga E., Gertsen H., Gooren H., Peters P., Salánki T., van der Ploeg M., Besseling
802 E., Koelmans A. A. and Geissen V.: Microplastics in the terrestrial ecosystem: implications for
803 *Lumbricus terrestris* (Oligochaeta, Lumbricidae), *Environmental Science & Technology*, 50,
804 2685-2691, doi:10.1021/acs.est.5b05478, 2016.

805 Huerta Lwanga E., Gertsen H., Gooren H., Peters P., Salánki T., van der Ploeg M., Besseling
806 E., Koelmans A. A. and Geissen V.: Incorporation of microplastics from litter into burrows of
807 *Lumbricus terrestris*, *Environmental Pollution*, 220, 523-531,
808 doi:10.1016/j.envpol.2016.09.096, 2017b.

809 Huerta Lwanga E., Thapa B., Yang X., Gertsen H., Salánki T., Geissen V. and Garbeva P.:
810 Decay of low-density polyethylene by bacteria extracted from earthworm's guts: A potential for
811 soil restoration, *Science of the Total Environment*, 624, 753-757,
812 doi:10.1016/j.scitotenv.2017.12.144, 2018.

813 Huerta Lwanga E., Vega J. M., Quej V. K., de los Angeles Chi J., del Cid L. S., Chi C., Segura
814 G. E., Gertsen H., Salánki T., van der Ploeg M. and others: Field evidence for transfer of
815 plastic debris along a terrestrial food chain, *Scientific Reports*, 7, 14071, doi:10.1038/s41598-
816 017-14588-2, 2017a.

817 Hughes L. and Westoby M.: Effect of diaspore characteristics on removal of seeds adapted
818 for dispersal by ants, *Ecology*, 73, 1300-1312, doi:10.2307/1940677, 1992.

819 Imhof H. K. and Laforsch C.: Hazardous or not - Are adult and juvenile individuals of
820 *Potamopyrgus antipodarum* affected by non-buoyant microplastic particles?, *Environmental*
821 *Pollution*, 218, 383-391, doi:10.1016/j.envpol.2016.07.017, 2016.

822 Jambeck J. R., Geyer R., Wilcox C., Siegler T. R., Perryman M., Andrady A., Narayan R. and
823 Law K. L.: Plastic waste inputs from land into the ocean, *Science*, 347, 768-771, doi:10.1126/
824 science.1260352, 2015.

825 Jeong C.-B., Won E.-J., Kang H.-M., Lee M.-C., Hwang D.-S., Hwang U.-K., Zhou B., Souissi
826 S., Lee S.-J. and Lee J.-S.: Microplastic size-dependent toxicity, oxidative stress induction,
827 and p-JNK and p-p38 activation in the monogonont rotifer (*Brachionus koreanus*),
828 *Environmental Science & Technology*, 50, 8849-8857, doi:10.1021/acs.est.6b01441., 2016.

829 Jin Y., Lu L., Tu W., Luo T. and Fu Z.: Impacts of polystyrene microplastic on the gut barrier,
830 microbiota and metabolism of mice, *Science of the Total Environment*, 649, 308-317,
831 doi:10.1016/j.scitotenv.2018.08.353, 2019.

832 Jonsson P. R.: Particle size selection, feeding rates and growth dynamics of marine
833 planktonic oligotrichous ciliates (Ciliophora: Oligotrichina), *Mar Ecol Prog Ser*, 33, 265-277,
834 doi:N/A, 1986.

835 Jouquet P., Dauber J., Lagerlöf J., Lavelle P. and Lepage M.: Soil invertebrates as ecosystem
836 engineers: intended and accidental effects on soil and feedback loops, *Applied Soil Ecology*,
837 32, 153-164, doi:10.1016/j.apsoil.2005.07.004, 2006.

838 Ju H., Zhu D. and Qiao M.: Effects of polyethylene microplastics on the gut microbial
839 community, reproduction and avoidance behaviors of the soil springtail, *Folsomia candida*,
840 *Environmental Pollution*, 247, 890-897, doi:10.1016/j.envpol.2019.01.097, 2019.

841 Judy J. D., Williams M., Gregg A., Oliver D., Kumar A., Kookana R. and Kirby J. K.:
842 Microplastics in municipal mixed-waste organic outputs induce minimal short to long-term
843 toxicity in key terrestrial biota, *Environmental Pollution*, 252, 522-531,
844 doi:10.1016/j.envpol.2019.05.027, 2019.

845 Kale S. K., Deshmukh A. G., Dudhare M. S. and Patil V. B.: Microbial degradation of plastic: a
846 review, *Journal of Biochemical Technology*, 6, 952-961, doi:N/A, 2015.

847 Kim H. M., Lee D.-K., Long N. P., Kwon S. W. and Park J. H.: Uptake of nanopolystyrene
848 particles induces distinct metabolic profiles and toxic effects in *Caenorhabditis elegans*,
849 *Environmental Pollution*, 246, 578-586, doi:10.1016/j.envpol.2018.12.043, 2019.

850 Kim S. W. and An Y.-J.: Soil microplastics inhibit the movement of springtail species,
851 *Environment International*, 126, 699-706, doi:10.1016/j.envint.2019.02.067, 2019.

852 Kiyama Y., Miyahara K. and Ohshima Y.: Active uptake of artificial particles in the nematode
853 *Caenorhabditis elegans*, *Journal of Experimental Biology*, 215, 1178-1183,
854 doi:10.1242/jeb.067199, 2012.

855 Kokalj A. J., Horvat P., Skalar T. and Kržan A.: Plastic bag and facial cleanser derived
856 microplastic do not affect feeding behaviour and energy reserves of terrestrial isopods,
857 *Science of the Total Environment*, 615, 761-766, doi:10.1016/j.scitotenv.2017.10.020, 2018.

858 Korb J.: The ecology of social evolution in termites, In: *Ecology of social evolution*, Springer,
859 doi:10.1007/978-3-540-75957-7_7, 2008.

860 Kumar A., Pandey A. K., Singh S. S., Shanker R. and Dhawan A.: Cellular uptake and
861 mutagenic potential of metal oxide nanoparticles in bacterial cells, *Chemosphere*, 83, 1124-
862 1132, doi:10.1016/j.chemosphere.2011.01.025, 2011.

863 Lahive E., Walton A., Horton A. A., Spurgeon D. J. and Svendsen C.: Microplastic particles
864 reduce reproduction in the terrestrial worm *Enchytraeus crypticus* in a soil exposure,
865 *Environmental Pollution*, 255, 113174, doi:10.1016/j.envpol.2019.113174, 2019.

866 Lavelle P., Decaëns T., Aubert M., Barot S., Blouin M., Bureau F., Margerie P., Mora P. and
867 Rossi J.-P.: Soil invertebrates and ecosystem services, *European Journal of Soil Biology*, 42,
868 S3-S15, doi:10.1016/j.ejsobi.2006.10.002, 2006.

869 Lavin D. P., Fredrickson A. and Srien F.: Flow cytometric measurement of rates of particle
870 uptake from dilute suspensions by a ciliated protozoan, *Cytometry*, 11, 875-882, doi:10.1002/
871 cyto.990110804, 1990.

872 Lei L., Liu M., Song Y., Lu S., Hu J., Cao C., Xie B., Shi H. and He D.: Polystyrene (nano)
873 microplastics cause size-dependent neurotoxicity, oxidative damage and other adverse
874 effects in *Caenorhabditis elegans*, *Environmental Science: Nano*, 5, 2009-2020, doi:10.1039/
875 C9EN00473D, 2018a.

876 Lei L., Wu S., Lu S., Liu M., Song Y., Fu Z., Shi H., Raley-Susman K. M. and He D.:
877 Microplastic particles cause intestinal damage and other adverse effects in zebrafish *Danio*
878 *rerio* and nematode *Caenorhabditis elegans*, *Science of the Total Environment*, 619, 1-8,
879 doi:10.1016/j.scitotenv.2017.11.103, 2018b.

880 Lenz M., Creffield J. W., Evans T. A., Kard B., Vongkaluang C., Sornnuwat Y., Lee C.-Y.,
881 Yoshimura T. and Tsunoda K.: Resistance of polyamide and polyethylene cable sheathings to
882 termites in Australia, Thailand, USA, Malaysia and Japan: a comparison of four field
883 assessment methods, *International Biodeterioration & Biodegradation*, 66, 53-62,
884 doi:10.1016/j.ibiod.2011.11.001, 2012.

885 Li J., Zhang H., Zhang K., Yang R., Li R. and Li Y.: Characterization, source, and retention of
886 microplastic in sandy beaches and mangrove wetlands of the Qinzhou Bay, China, *Marine*
887 *Pollution Bulletin*, 136, 401-406, doi:10.1016/j.marpolbul.2018.09.025, 2018a.

888 Li L., Zhou Q., Yin N., Tu C. and Luo Y.: Uptake and accumulation of microplastics in an
889 edible plant, *Chinese Science Bulletin*, 64, 928-934, doi:10.1360/N972018-00845, 2019.

890 Li X., Chen L., Mei Q., Dong B., Dai X., Ding G. and Zeng E. Y.: Microplastics in sewage
891 sludge from the wastewater treatment plants in China, *Water Research*, 142, 75-85,
892 doi:10.1016/j.watres.2018.05.034, 2018b.

893 Lobelle D. and Cunliffe M.: Early microbial biofilm formation on marine plastic debris, *Marine*
894 *Pollution Bulletin*, 62, 197-200, doi:10.1016/j.marpolbul.2010.10.013, 2011.

895 Lu L., Wan Z., Luo T., Fu Z. and Jin Y.: Polystyrene microplastics induce gut microbiota
896 dysbiosis and hepatic lipid metabolism disorder in mice, *Science of the Total Environment*,
897 631, 449-458, doi:10.1016/j.scitotenv.2018.03.051, 2018.

898 Maaß S., Daphi D., Lehmann A. and Rillig M. C.: Transport of microplastics by two
899 collembolan species, *Environmental Pollution*, 225, 456-459,
900 doi:10.1016/j.envpol.2017.03.009, 2017.

901 Mahon A. M., O'Connell B., Healy M. G., O'Connor I., Officer R., Nash R. and Morrison L.:
902 Microplastics in sewage sludge: effects of treatment, *Environmental Science & Technology*,
903 51, 810-818, doi:10.1021/acs.est.6b04048, 2016.

904 Monge-Nájera J.: Ecological biogeography in the Phylum Onychophora, *Biogeographica*, 70,
905 111-123, doi:N/A, 1994.

906 Naji A., Esmaili Z., Mason S. A. and Vethaak A. D.: The occurrence of microplastic
907 contamination in littoral sediments of the Persian Gulf, Iran, *Environmental Science and*
908 *Pollution Research*, 24, 20459-20468, doi:10.1007/s11356-017-9587-z, 2017.

909 Nika L., Gibson T., Konkus R. and Karp X.: Fluorescent beads are a versatile tool for staging
910 *Caenorhabditis elegans* in different life histories, *G3: Genes, Genomes, Genetics*, 6, 1923-
911 1933, doi:10.1534/g3.116.030163, 2016.

912 Nizzetto L., Futter M. and Langaas S.: Are agricultural soils dumps for microplastics of urban
913 origin?, *Environmental Science & Technology*, doi:10.1021/acs.est.6b04140, 2016.

914 Nor N. H. M. and Obbard J. P.: Microplastics in Singapore's coastal mangrove ecosystems,
915 *Marine Pollution Bulletin*, 79, 278-283, doi:10.1016/j.marpolbul.2013.11.025, 2014.

916 Oliveira M., Ameixa O. M. and Soares A. M.: Are ecosystem services provided by insects"
917 bugged" by micro (nano) plastics?, *TrAC Trends in Analytical Chemistry*,
918 doi:10.1016/j.trac.2019.02.018, 2019.

919 Parkyn J. and Newell D. A.: Australian land snails: a review of ecological research and
920 conservation approaches, *Molluscan Research*, 33, 116-129,
921 doi:10.1080/13235818.2013.782793, 2013.

922 Pass G., Szucsich N. U. and others: 100 years of research on the Protura: many secrets still
923 retained, *Soil Organisms*, 83, 309-334, doi:N/A, 2011.

924 Pelosi C., Bertrand M., Thénard J. and Mougin C.: Earthworms in a 15 years agricultural trial,
925 *Applied Soil Ecology*, 88, 1-8, doi:10.1016/j.apsoil.2014.12.004, 2015.

926 Peng B.-Y., Su Y., Chen Z., Chen J., Zhou X., Benbow M. E., Criddle C., Wu W.-M. and Zhang
927 Y.: Biodegradation of Polystyrene by Dark (*Tenebrio obscurus*) and Yellow (*Tenebrio molitor*)
928 Mealworms (Coleoptera: Tenebrionidae), *Environmental Science & Technology*,
929 doi:10.1021/acs.est.8b06963, 2019.

930 Petersen H.: A review of collembolan ecology in ecosystem context, *Acta Zoologica Fennica*,
931 195, 111-118, doi:N/A, 1994.

932 Phillips H. R., Guerra C. A., Bartz M. L., Briones M. J., Brown G., Ferlian O., Gongalsk K.,
933 Krebs J., Orgiazzi A., Schwarz B. and others: Global distribution of earthworm diversity,
934 *bioRxiv*, 587394, doi:10.1126/science.aax4851, 2019.

935 Prendergast-Miller M. T., Katsiamides A., Abbass M., Sturzenbaum S. R., Thorpe K. L. and
936 Hodson M. E.: Polyester-derived microfibre impacts on the soil-dwelling earthworm *Lumbricus*
937 *terrestris*, *Environmental Pollution*, 251, 453-459, doi:10.1016/j.envpol.2019.05.037, 2019.

938 Pulleman M., Creamer R., Hamer U., Helder J., Pelosi C., Peres G. and Rutgers M.: Soil
 939 biodiversity, biological indicators and soil ecosystem services—an overview of European
 940 approaches, *Current Opinion in Environmental Sustainability*, 4, 529-538,
 941 doi:10.1016/j.cosust.2012.10.009, 2012.

942 Qu M., Kong Y., Yuan Y. and Wang D.: Neuronal damage induced by nanopolystyrene
 943 particles in nematode *Caenorhabditis elegans*, *Environmental Science: Nano*,
 944 doi:10.1039/C9EN00473D, 2019a.

945 Qu M., Nida A., Kong Y., Du H., Xiao G. and Wang D.: Nanopolystyrene at predicted
 946 environmental concentration enhances microcystin-LR toxicity by inducing intestinal damage
 947 in *Caenorhabditis elegans*, *Ecotoxicology and Environmental Safety*, 183, 109568,
 948 doi:10.1016/j.ecoenv.2019.109568, 2019b.

949 Qu M., Qiu Y., Kong Y. and Wang D.: Amino modification enhances reproductive toxicity of
 950 nanopolystyrene on gonad development and reproductive capacity in nematode
 951 *Caenorhabditis elegans*, *Environmental Pollution*, 254, 112978,
 952 doi:10.1016/j.envpol.2019.112978, 2019c.

953 Reddy M. S., Basha S., Adimurthy S. and Ramachandraiah G.: Description of the small
 954 plastics fragments in marine sediments along the Alang-Sosiya ship-breaking yard, India,
 955 *Estuarine, Coastal and Shelf Science*, 68, 656-660, doi:10.1016/j.ecss.2006.03.018, 2006.

956 Rezaei M., Riksen M. J., Sirjani E., Sameni A. and Geissen V.: Wind erosion as a driver for
 957 transport of light density microplastics, *Science of The Total Environment*, 669, 273-281,
 958 doi:10.1016/j.scitotenv.2019.02.382, 2019.

959 Rillig M. C.: Microplastic in terrestrial ecosystems and the soil?, *Environmental Science &*
 960 *Technology*, 6453–6454, doi:10.1021/es302011r, 2012.

961 Rillig M. C. and Bonkowski M.: Microplastic and soil protists: a call for research,
 962 *Environmental Pollution*, 241, 1128-1131, doi:10.1016/j.envpol.2018.04.147, 2018.

963 Rillig M. C., Lehmann A., de Souza Machado A. A. and Yang G.: Microplastic effects on
 964 plants, *New Phytologist*, doi:10.1111/nph.15794, 2019.

965 Rillig M. C., Ziersch L. and Hempel S.: Microplastic transport in soil by earthworms, *Scientific*
 966 *Reports*, 7, 1362, doi:10.1038/s41598-017-01594-7, 2017.

967 Robins R. and Robins A.: The antics of ants: ants as agents of bioturbation in a midden
 968 deposit in south-east Queensland, *Environmental Archaeology*, 16, 151-161,
 969 doi:10.1179/174963111X13110803261010, 2011.

970 Rodríguez-Seijo A., da Costa J. P., Rocha-Santos T., Duarte A. C. and Pereira R.: Oxidative
 971 stress, energy metabolism and molecular responses of earthworms (*Eisenia fetida*) exposed

972 to low-density polyethylene microplastics, *Environmental Science and Pollution Research*, 25,
 973 33599-33610, doi:10.1007/s11356-018-3317-z, 2018.

974 Rodriguez-Seijo A., Lourenço J., Rocha-Santos T., Da Costa J., Duarte A., Vala H. and
 975 Pereira R.: Histopathological and molecular effects of microplastics in *Eisenia andrei* Bouché,
 976 *Environmental Pollution*, 220, 495-503, doi:10.1016/j.envpol.2016.09.092, 2017.

977 Rodríguez-Seijo A., Santos B., da Silva E. F., Cachada A. and Pereira R.: Low-density
 978 polyethylene microplastics as a source and carriers of agrochemicals to soil and earthworms,
 979 *Environmental Chemistry*, 16, 8-17, doi:10.1071/EN18162, 2019.

980 Sánchez-Bayo F. and Wyckhuys K. A.: Worldwide decline of the entomofauna: A review of its
 981 drivers, *Biological Conservation*, 232, 8-27, doi:10.1016/j.biocon.2019.01.020, 2019.

982 Scharf A., Piechulek A. and von Mikecz A.: Effect of nanoparticles on the biochemical and
 983 behavioral aging phenotype of the nematode *Caenorhabditis elegans*, *Acs Nano*, 7, 10695-
 984 10703, doi:10.1021/nn403443r, 2013.

985 e Silva P. P. G., Nobre C. R., Resaffe P., Pereira C. D. S. and Gusmão F.: Leachate from
 986 microplastics impairs larval development in brown mussels, *Water Research*, 106, 364-370,
 987 doi:10.1016/j.watres.2016.10.016, 2016.

988 Song Y., Cao C., Qiu R., Hu J., Liu M., Lu S., Shi H., Raley-Susman K. M. and He D.: Uptake
 989 and adverse effects of polyethylene terephthalate microplastics fibers on terrestrial snails
 990 (*Achatina fulica*) after soil exposure, *Environmental Pollution*, 250, 447-455,
 991 doi:10.1016/j.envpol.2019.04.066, 2019.

992 Stamatiadis S. and Dindal D.: Coprophilous mite communities as affected by concentration of
 993 plastic and glass particles, *Experimental & Applied Acarology*, 8, 1-12,
 994 doi:10.1007/BF01193377, 1990.

995 Stock V., Böhmert L., Lisicki E., Block R., Cara-Carmona J., Pack L. K., Selb R., Lichtenstein
 996 D., Voss L., Henderson C. J. and others: Uptake and effects of orally ingested polystyrene
 997 microplastic particles in vitro and in vivo, *Archives of Toxicology*, 1-17, doi:10.1007/s00204-
 998 019-02478-7, 2019.

999 Sun Y., Xu W., Gu Q., Chen Y., Zhou Q., Zhang L., Gu L., Huang Y., Lyu K. and Yang Z.:
 1000 Small-Sized Microplastics Negatively Affect Rotifers: Changes in the Key Life-History Traits
 1001 and Rotifer--*Phaeocystis* Population Dynamics, *Environmental Science & Technology*, 53,
 1002 9241-9251, doi:10.1021/acs.est.9b02893, 2019.

1003 Thompson R. C., Swan S. H., Moore C. J. and Vom Saal F. S.: Our plastic age,
 1004 doi:10.1098/rstb.2009.0054, 2009.

1005 Tsunoda K., Rosenblat G. and Dohi K.: Laboratory evaluation of the resistance of plastics to
 1006 the subterranean termite *Coptotermes formosanus* (Blattodea: Rhinotermitidae), *International*
 1007 *Biodeterioration & Biodegradation*, 64, 232-237, doi:10.1016/j.ibiod.2009.12.008, 2010.

1008 Vadstein O., Øie G. and Olsen Y.: Particle size dependent feeding by the rotifer *Brachionus*
 1009 *plicatilis*, *Hydrobiologia*, 255, 261-267, doi:10.1007/BF00025847, 1993.

1010 Van Cauwenberghe L., Claessens M., Vandegehuchte M. B. and Janssen C. R.: Microplastics
 1011 are taken up by mussels (*Mytilus edulis*) and lugworms (*Arenicola marina*) living in natural
 1012 habitats, *Environmental Pollution*, 199, 10-17, doi:10.1016/j.envpol.2015.01.008, 2015.

1013 Vianello A., Boldrin A., Guerriero P., Moschino V., Rella R., Sturaro A. and Da Ros L.:
 1014 Microplastic particles in sediments of Lagoon of Venice, Italy: First observations on
 1015 occurrence, spatial patterns and identification, *Estuarine, Coastal and Shelf Science*, 130, 54-
 1016 61, doi:10.1016/j.ecss.2013.03.022, 2013.

1017 Vogel G., Thilo L., Schwarz H. and Steinhart R.: Mechanism of phagocytosis in *Dictyostelium*
 1018 *discoideum*: phagocytosis is mediated by different recognition sites as disclosed by mutants
 1019 with altered phagocytotic properties, *The Journal of Cell Biology*, 86, 456-465,
 1020 doi:10.1083/jcb.86.2.456, 1980.

1021 Wang H.-T., Ding J., Xiong C., Zhu D., Li G., Jia X.-Y., Zhu Y.-G. and Xue X.-M.: Exposure to
 1022 microplastics lowers arsenic accumulation and alters gut bacterial communities of earthworm
 1023 *Metaphire californica*, *Environmental Pollution*, 251, 110-116,
 1024 doi:10.1016/j.envpol.2019.04.054, 2019b.

1025 Wang J., Coffin S., Sun C., Schlenk D. and Gan J.: Negligible effects of microplastics on
 1026 animal fitness and HOC bioaccumulation in earthworm *Eisenia fetida* in soil, *Environmental*
 1027 *Pollution*, 249, 776-784, doi:10.1016/j.envpol.2019.03.102, 2019c.

1028 Wang J., Liu X., Li Y., Powell T., Wang X., Wang G. and Zhang P.: Microplastics as
 1029 contaminants in the soil environment: A mini-review, *Science of The Total Environment*,
 1030 doi:10.1016/j.scitotenv.2019.07.209, 2019a.

1031 Warburg M. R.: Isopods and their terrestrial environment, In: *Advances in ecological research*,
 1032 Elsevier, doi:10.1016/S0065-2504(08)60246-9, 1987.

1033 Weisman R. A. and Korn E. D.: Phagocytosis of latex beads by *Acanthamoeba*. I. Biochemical
 1034 properties, *Biochemistry*, 6, 485-497, doi:10.1021/bi00854a017, 1967.

1035 Weithmann N., Möller J. N., Löder M. G., Piehl S., Laforsch C. and Freitag R.: Organic
 1036 fertilizer as a vehicle for the entry of microplastic into the environment, *Science Advances*, 4,
 1037 eaap8060, doi:10.1126/sciadv.aap8060, 2018.

1038 Westheide W. and Rieger R.: Spezielle Zoologie. 1. Einzeller und Wirbellose Tiere, Gustav
1039 Fischer, 1996.

1040 Wood C. T. and Zimmer M.: Can terrestrial isopods (Isopoda: Oniscidea) make use of
1041 biodegradable plastics?, *Applied Soil Ecology*, 77, 72-79, doi:10.1016/j.apsoil.2014.01.009,
1042 2014.

1043 Wright S. L., Rowe D., Thompson R. C. and Galloway T. S.: Microplastic ingestion decreases
1044 energy reserves in marine worms, *Current Biology*, 23, R1031-R1033,
1045 doi:10.1016/j.cub.2013.10.068, 2013.

1046 Yang S.-S., Brandon A. M., Flanagan J. C. A., Yang J., Ning D., Cai S.-Y., Fan H.-Q., Wang
1047 Z.-Y., Ren J., Benbow E. and others: Biodegradation of polystyrene wastes in yellow
1048 mealworms (larvae of *Tenebrio molitor* Linnaeus): Factors affecting biodegradation rates and
1049 the ability of polystyrene-fed larvae to complete their life cycle, *Chemosphere*, 191, 979-989,
1050 doi:10.1016/j.chemosphere.2017.10.117, 2018.

1051 Yang X., Lwanga E. H., Bemani A., Gertsen H., Salanki T., Guo X., Fu H., Xue S., Ritsema C.
1052 and Geissen V.: Biogenic transport of glyphosate in the presence of LDPE microplastics: A
1053 mesocosm experiment, *Environmental Pollution*, 245, 829-835,
1054 doi:10.1016/j.envpol.2018.11.044, 2019a.

1055 Yang Y., Yang J., Wu W.-M., Zhao J., Song Y., Gao L., Yang R. and Jiang L.: Biodegradation
1056 and mineralization of polystyrene by plastic-eating mealworms: Part 1. Chemical and physical
1057 characterization and isotopic tests, *Environmental Science & Technology*, 49, 12080-12086,
1058 doi:10.1021/acs.est.5b02661, 2015.

1059 Yang Y.-F., Chen C.-Y., Lu T.-H. and Liao C.-M.: Toxicity-based toxicokinetic/toxicodynamic
1060 assessment for bioaccumulation of polystyrene microplastics in mice, *Journal of Hazardous*
1061 *Materials*, 366, 703-713, doi:10.1016/j.jhazmat.2018.12.048, 2019b.

1062 Yeates G. W.: Nematodes in ecological webs, *e LS*, doi:10.1002/9780470015902.a0021913,
1063 2001.

1064 Yu M., Van Der Ploeg M., Lwanga E. H., Yang X., Zhang S., Ma X., Ritsema C. J. and
1065 Geissen V.: Leaching of microplastics by preferential flow in earthworm (*Lumbricus terrestris*)
1066 burrows, *Environmental Chemistry*, 16, 31-40, doi:10.1071/EN18161, 2019.

1067 Zhang G. and Liu Y.: The distribution of microplastics in soil aggregate fractions in
1068 southwestern China, *Science of the Total Environment*, 642, 12-20,
1069 doi:10.1016/j.scitotenv.2018.06.004, 2018.

1070 Zhang L., Sintim H. Y., Bary A. I., Hayes D. G., Wadsworth L. C., Anunciado M. B. and Flury
1071 M.: Interaction of *Lumbricus terrestris* with macroscopic polyethylene and biodegradable

1073 plastic mulch, Science of The Total Environment, 635, 1600-1608,
 1074 doi:10.1016/j.scitotenv.2018.04.054, 2018.

1075 Zhang Y., Gao T., Kang S. and Sillanpää M.: Importance of atmospheric transport for
 1076 microplastics deposited in remote areas, Environmental Pollution, 254, 112953,
 1077 doi:10.1016/j.envpol.2019.07.121, 2019.

1078 Zhang Z.-Q.: Animal biodiversity: an update of classification and diversity in 2013, Zootaxa,
 1079 3703, 5-11, doi:10.11646/zootaxa.3703.1.3, 2013.

1080 Zhao L., Qu M., Wong G. and Wang D.: Transgenerational toxicity of nanopolystyrene
 1081 particles in the range of $\mu\text{g L}^{-1}$ in the nematode *Caenorhabditis elegans*, Environmental
 1082 Science: Nano, 4, 2356-2366, doi:10.1039/C7EN00707H, 2017.

1083 Zhu B.-K., Fang Y.-M., Zhu D., Christie P., Ke X. and Zhu Y.-G.: Exposure to nanoplastics
 1084 disturbs the gut microbiome in the soil oligochaete *Enchytraeus crypticus*, Environmental
 1085 Pollution, 239, 408-415, doi:10.1016/j.envpol.2018.04.017, 2018b.

1086 Zhu D., Bi Q.-F., Xiang Q., Chen Q.-L., Christie P., Ke X., Wu L.-H. and Zhu Y.-G.: Trophic
 1087 predator-prey relationships promote transport of microplastics compared with the single
 1088 *Hypoaspis aculeifer* and *Folsomia candida*, Environmental Pollution, 235, 150-154,
 1089 doi:10.1016/j.envpol.2017.12.058, 2018a.