



1 What do we know about how the terrestrial 2 multicellular soil fauna reacts to microplastic? 3

4 Frederick Büks¹, Nicolette Loes van Schaik², Martin Kaupenjohann¹
5

6 ¹Chair of Soil Science, Dept. of Ecology, Technische Universität Berlin, 10587 Berlin, Germany

7 ²Chair of Ecohydrology & Landscape Evaluation, Dept. of Ecology, Technische Universität Berlin, 10587 Berlin, Germany
8

9 Correspondence to: Frederick Büks (frederick.bueks@tu-berlin.de)
10
11

12 **Abstract.** The ubiquitous accumulation of microplastic (MP) particles across all global
13 ecosystems comes along with the uptake into soil food webs. In this review, we analyzed
14 studies on passive translocation, active ingestion, bioaccumulation and adverse effects within
15 the phylogenetic tree of multicellular soil faunal life. The representativity of these studies for
16 natural soil ecosystems was assessed using data on the type of plastic, shape, composition,
17 concentration and time of exposure.

18 Available studies cover a wide range of soil organisms, with emphasis on earthworms,
19 nematodes, springtails, beetles and lugworms, each focused on well known model organisms.
20 Most of the studies applied MP concentrations similar to amounts in slightly to very heavily
21 polluted soils. In many cases, however, polystyrene microspheres have been used, a
22 combination of plastic type and shape, that is easily available, but does not represent the
23 main plastic input into soil ecosystems. In turn, MP fibres are strongly underrepresented
24 compared to their high abundance within contaminated soils. Further properties of plastic
25 such as aging, coating and additives were insufficiently documented. Despite these
26 limitations, there is a recurring pattern of active intake followed by a population shift within the
27 gut microbiome and adverse effects on motility, growth, metabolism, reproduction and
28 mortality in various combinations, especially at high concentrations and small particle sizes.

29 For the improvement of future studies, we identified problems of past experiments and give
30 recommendations that take into account the type, shape, grade of aging, specific
31 concentrations of MP fractions and long-term incubation in natural and contaminated soils.



32 **1 Introduction**

33 Imagine a compact plastic cube of nearly 2 km side length and a weight of 7300000000 tons,
34 with major percentages by weight of 36 % polyethylene (PE), 21 % polypropylene (PP), 12 %
35 polyvinyl chloride (PVC) and 10 % of each polyethylene terephthalate (PET), polyurethane
36 (PU) and polystyrene (PS). That is the cumulated global non-fibre production of the six main
37 plastic types until 2015. It accounts to 87 % of the all-time plastic production, which evolved,
38 since the early 1950s, exponentially from some megatons (Mt) to 8300 Mt in 2015, with only
39 260 Mt annual output in 2009 increased to 380 Mt in 2015 (Thompson et al., 2009; Geyer et
40 al., 2017). Of this ever produced plastic, 6300 Mt became waste until 2015, of which only
41 21 % were recycled or incinerated, whereas 5000 Mt ended up in landfills and nature (Geyer
42 et al., 2017). As a corollary of production, use and disposal, a certain part of plastic waste is
43 constantly released into the environment on various paths, but our knowledge about mass
44 flow rates into global ecosystems is very limited. Based on waste generation in coastal
45 countries, Jambeck et al. (2015) calculated the global plastic input to marine ecosystems to
46 be roughly 4.8 to 12.7 Mt in 2010. Such data on soils are lacking, but Nizzetto et al. (2016)
47 estimated that the load of microplastic (MP) to agricultural sites in Europe is in the same order
48 of magnitude as to marine environments.

49 By littering, plastic mulching, the application of sewage sludge, digestates and composts as
50 well as windblown dispersal (Bertling et al., 2018; Weithmann et al., 2018; Zhang et al., 2019;
51 Wang et al., 2019a), plastic from our technosphere arrives in soil ecosystems in various forms
52 as large and small fragments, fibers and particles. Exposed to UV radiation, mechanical
53 stress and microbial decay, plastic items become weathered and prone to a successive
54 comminution towards the size range of MP with increased surface, charge and biofilm cover
55 (Kale et al., 2015; Andrady, 2017). However, the resistance of plastic to metabolization
56 causes a constant accumulation in soils as long as the release rate from human processes is
57 above the very slow rate of degradation.

58 Due to a lack of monitoring programs, data on MP concentrations in terrestrial soils are rare,
59 and those using w/w concentrations are even sparser. Under less contaminated conditions,
60 amounts seem to average about 1 mg kg⁻¹ soil dry weight (and approx. 200 items kg⁻¹ dry soil)
61 (Rezaei et al., 2019). On sites with industrial activity or use of plastic mulching and sewage
62 sludge in agriculture, concentrations can be increased by 2 to 4 orders of magnitude (Fuller
63 and Gautam, 2016; Zhang and Liu, 2018). Semisubhydric soils such as beaches, mudflats,
64 mangroves or lagoons, that are additionally contaminated from the aquatic side, contain MP
65 of the order of 10 to 100 items kg⁻¹ dry soil and single extreme samplings contained several
66 thousand items (Nor and Obbard, 2014; Naji et al., 2017; Garcés-Ordóñez et al., 2019; Li et
67 al., 2018a). More informative data using mg kg⁻¹ are only available for beaches and coastal
68 deconstruction yards in municipal neighbourhood and amount to 0.5 and 70 mg kg⁻¹ dry soil,
69 0.00005 and 0.007 % w/w, respectively (Reddy et al., 2006; Claessens et al., 2011). All these



70 concentration data represent a wide range of particle sizes between 0 and 5000 μm with
71 different materials, shapes and grades of aging.

72 Plastic particles can possibly enter and accumulate within the food web by either direct
73 uptake from soil or consumption of other soil biota contaminated by adhesion or ingestion
74 (Huerta Lwanga et al., 2017a). There is evidence, that MP is incorporated even by plants and
75 unicellular organisms at the base of the food web. **Bacteria**, for example, that are reasonably
76 assumed to avoid MP uptake due to their minor size and the prevalent lack of phagocytosis,
77 were shown to take up inorganic nanoparticles of a few nanometers (Kumar et al., 2011).
78 Although the physiochemical properties of weathered nanoparticulate plastics might differ from
79 these, also their uptake seems reasonable.

80 A similar argument can be made for **fungi** and soil **algae**, but studies on incorporation are
81 lacking, whereas the transfer into a freshwater food web by adhesion of nanoplastic on algae
82 has been shown by Chae et al. (2018). The uptake of MP into **plant roots** is also inhibited
83 (Rillig et al., 2019), but occurred for nanoplastics that permeate into the plant tissue (Li et al.,
84 2019). Also the ingrowth into root tissue after adsorption to the rhizodermis should be tested.

85 In contrast, **protozoa** feature phagocytosis for the active ingestion of particles. Diverse soil,
86 freshwater and marine ciliates ingest PS/latex beads of 0.1 to 14.4 μm in laboratory
87 experiments, with preferences to their natural prey size (Fenchel, 1980; Jonsson, 1986; Lavin
88 et al., 1990). Soil amoebas act similarly, but additionally select according to food quality
89 (Weisman and Korn, 1967; Vogel et al., 1980; Bowers and Olszewski, 1983; Avery et al.,
90 1995; Elloway et al., 2006).

91 At last, many soil microbiota live protected within biofilms. Plastic particles were shown to be
92 surface for the formation of those biofilms (Lobelle and Cunliffe, 2011), which are a food
93 sources of grazing primary consumers. Feeding on them might also transfer occluded or
94 abraded MP to higher trophic levels.

95 But what about the larger organisms that feed on all these, free plastic particles,
96 contaminated microorganisms, biofilms and one another? Recent work discussed the effects
97 of MP on soil biota (Chae and An, 2018) or called for intensified research on certain
98 taxonomic groups (Rillig and Bonkowski, 2018). Thus, we were motivated to give on our part
99 a short review with focus on the most-produced plastics and their passive translocation,
100 ingestion, bioaccumulation and adverse effects on the multicellular soil fauna. The types,
101 sizes and shapes of plastic used in former laboratory studies were compared with our
102 knowledge on plastic in the environment, and recommendations are given for future research.
103 This analysis is aimed to help for assessing the influence of MP on the ecosystem services of
104 diverse soil organisms.



105 2 Search pattern

106 Within the tree of life, edaphic branches were identified comprising taxa that permanently
107 inhabit the soil, are both-sided part of the soil food web and/or the burrowing macro- and
108 megafauna or have active subterranean larval stages. The resulting tree of soil life based on
109 the NCBI taxonomy database (Fig. 1) was charted by use of the software [phyloT](#) and shows
110 the leading taxonomic rank, which is mainly the family, but in exceptions – e.g. if one species
111 represents the only soil-born between many aquatic – a lower rank.




Figure 1: Edaphic tree of faunal life. Taxonomic ranks, that were examined in this qualitative study, are placed at the outer rim of the diagram. The length of the connecting line between two taxa is representative for the grade of phylogenetic relationship.



113 A pattern of search terms was established (see *Table 8*), consisting of „taxon“ (Linné’s
114 binominal nomenclature, common name, plural-sensitive search), „plastic type“ (plastic,
115 microplastic, nanoplastic, PE or polyethylene, PP or polypropylene, PVC or polyvinyl chloride,
116 PS or polystyrene, PU or polyurethane, PET or polyethylene terephthalate and latex) and
117 „common shapes“ (fragments, particles, fibres, microfibrils, beads, microbeads,
118 microspheres). Type-shape combinations, that would had cause to much search effort (e.g.
119 organism–plastic) or did not appear within a foregoing search (e.g. PET–microbeads or latex–
120 microfibrils), were excluded from this pattern. Further plastic types and shapes occurring
121 within the found studies were also included to the review. Data on microspheres and
122 microbeads were pooled, as both names describe one and the same.

123 The search appeared within the Web of Science Core Collection Database. Based on the
124 search pattern, data on passive transport, ingestion, bioaccumulation and adverse effects
125 were collected for each edaphic group. Studies that only use uncommon, local, outdated,
126 weird or nicknames are excluded by the pattern. Studies testing injection to tissues, lymph or
127 blood were excluded, as they do not represent natural ways to incorporate MPs. Data on
128 inhalation by the megafauna in fact represent a natural way of uptake, but were also excluded
129 as they are exclusively related to above-ground organisms, that only occur on the outer edge
130 of the food-web. Also running debates on phylogenetic classifications are not part of this work
131 and the taxonomist will be able to adjust the branches accordingly to his purpose.

132 The data of related taxonomic groups were pooled and evaluated for their environmental
133 representativity based on exposure time, plastic concentrations and properties used. From
134 these data recommendations for a structured experimental design in future studies were
135 derived.



136 **3 Data collection**

137

138 **3.1 Insects**

139 Within the Panarthropoda, the insects comprise the highest taxonomic diversity. And,
140 regarding MPs, they represent an unevenly studied taxonomic group.

141 Within the Insecta, the **Coleoptera** (beetles) build an extraordinarily diverse and abundant
142 taxon. Studies on plastic uptake into adult individuals mainly focus on the subfamily of
143 Scarabaeinae (dung beetles). Comprehensive experiments with latex microbeads showed,
144 that many species only ingest fine particles with maximum diameters of about 10 to 83 μm
145 and retain them within the gut – with a slightly positive dependency on body size. Larger
146 particles were rejected by a filtering mechanism within the mouth region and not ground with
147 the mandibles (Holter, 2000; Holter et al., 2002; Holter and Scholtz, 2005). Beside those on
148 Nematods, these data comprise by far the most detailed information about size-dependend
149 uptake of MP particles compared to other edaphic taxa. This gives a good foundation for
150 future studies on adverse concentrations. In addition, several studies with plastic as
151 predominant food source could show chewing, ingestion and intestinal degradation of
152 different PS and PE foams in feeding experiments with *Tenebrio sp.* larvae (mealworms).
153 These experiments also pointed out an alteration of the gut microbiome, but no adverse
154 effects on reproduction and survival, with only in one case of non-significant tendency to
155 higher mortality after 1 month of exposure (Yang et al., 2015; Brandon et al., 2018; Yang et
156 al., 2018; Peng et al., 2019).

157 The **Isoptera** (termites), recently categorized as part of the order Blattodea, are the oldest
158 social insects having a tribal history of about 130 million years (Korb, 2008). Especially in arid
159 ecosystems with lack of earthworms they play an important role in homogenization of soils,
160 but also in sorting of soil mineral particles for building mounds as well as decomposition and
161 distribution of organic matter (De Bruyn and Conacher, 1990). Tsunoda et al. (2010) and Lenz
162 et al. (2012) could show, that different termite species are picky feeders and erode PE, but
163 avoid other plastic cable sheathings. This suggests the excretion of ground MP particles by
164 termites, but metabolic impacts are unknown. In contrast to termites, data on **other Blattodea**
165 (e.g. cockroaches) were not found.

166 The suborder **Apocrita** comprises some flying insects, that inhabit burrows within the soil,
167 such as ground-dwelling wasps within the **Vespidae** superfamily, mining bees within the
168 **Apoidea** superfamily and the **Spheciformes**. They mostly do not prey and feed on
169 subterrestrial organisms, but may move MP particles into the ground, as implied by a report of
170 Allasino et al. (2019) on solitary bees, which built nests fully made of plastic fragments. The
171 Apocrita also contain the **Formicidae** (ants). Some ant species are considered an important
172 factor for seed dispersal, a behavior, that could also be shown for artificial plastic seeds with
173 ~2 mm diameter (Hughes and Westoby, 1992; Angotti et al., 2018). Robins and Robins (2011)



174 found that this also includes differently shaped cultural objects: *Rhytidoponera metallica*, a
175 representative of ground-nesting, omnivore ants, is capable not only of a remarkable
176 bioturbation but also of an active, apparently random burying of anthropogenic plastic
177 artefacts >1 mm. Seeds are used as a food source, thus, the ingestion of plastic bites is
178 conceivable, but not documented. The uptake of latex microspheres $\geq 0.88 \mu\text{m}$ with liquids by
179 larvae of *Solenopsis invicta* seems to be prevented by filtration within the mouth and the
180 particles are released as larger aggregates, whereas other species ingest by far larger
181 particles up to $150 \mu\text{m}$ (Glancey et al., 1981). However, also here data on adverse effects are
182 missing.

183 Further insects with edaphic adult stages, e.g. **Dermaptera** (earwigs), **Heteroptera** (true
184 bugs) and **Zygentoma** (silverfish, fishmoth, firebrat) or soil- or litter-dwelling larvae such as
185 **Embioptera** (webspinners, footspinners), **Thysanoptera** (thrips), **Psocoptera** (booklice,
186 barklice, barkflies), **Neuroptera** (lacewings), **Raphidioptera** (snakeflies) or **Zoraptera** (angel
187 insects) are not yet researched with focus on soil MP.

188 Regarding insects, mainly studies on translocation and uptake of MP were carried out. In
189 contrast, work on bioaccumulation is completely lacking and adverse effects are sparsely
190 tested using *Tenebrio sp.* larvae. Such studies could provide information whether or not the
191 input of MP in soil ecosystems is one of many factors causing the global decline of the
192 entomofauna (Oliveira et al., 2019; Sánchez-Bayo and Wyckhuys, 2019).



Table 1: Microplastic studies on Coleoptera, Blattodea (Blattoid.), Apoidea (A.) and Formicidae (mb=microbeads, fr=fragments, ms=microspheres, b=beads). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
<i>Aphodius erraticus</i>							5				no			
<i>Aphodius rufipes</i>							2..39				≤14 µm			
<i>Aphodius ater</i>	petri dish	latex	N/A	N/A	N/A	mb	2..39	N/A	45 min	N/A	≤14 µm	N/A	N/A	Holter (2000)
<i>Aphodius fimetarius</i>							2..39				≤18 µm			
<i>Aphodius contaminatus</i>							2..39				≤18 µm			
<i>Aphodius fossor</i>							2..39				≤18 µm			
diverse dung beetles	vial	latex	N/A	N/A	N/A	mb	2..83	N/A	45 min	N/A	≤10..≤60 µm	N/A	N/A	Holter et al. (2002)
diverse dung beetles	N/A	latex	N/A	N/A	N/A	mb	2..83	N/A	45 min	N/A	≤4..≤95 µm	N/A	N/A	Holter and Scholiz (2005)
<i>Tenebrio molitor</i> larvae	container	PS	N/A	N/A	no	foam	N/A	100% w/w (food)	31 d	N/A	yes	biodegrad.	N/A	Yang et al. (2015)
<i>Tenebrio molitor</i> larvae	container	LD-PE PS	N/A	N/A	no flame retardant	foam	8..27 cm ³	50..100% w/w (food)	32 d	N/A	yes	biodegrad.	microbiome	Brandon et al. (2018)
<i>Tenebrio molitor</i> larvae	container	PS	N/A	N/A	N/A	foam	N/A	4..100% w/w (food)	32 d	N/A	yes	biodegrad.	no	Yang et al. (2018)
<i>Tenebrio molitor</i> larvae	N/A	PS	N/A	N/A	no	foam	N/A	86..100% w/w (food)	31 d	N/A	yes	biodegrad.	microbiome	Peng et al. (2019)
<i>Tenebrio obscurus</i> larvae														
<i>Coptotermes formosanus</i>	mesocosm	LD-PE others	yes/no	N/A	N/A	cable sheets	4 cm, Ø 0.8 cm	N/A	42 d	N/A	yes	N/A	N/A	Tsumoda et al. (2010)
diverse termites	in situ	MD-PE PA	no	N/A	anti-oxidant stabilizer	cable sheets	30 cm, Ø 1.4 cm	N/A	6 yr.	N/A	yes	N/A	N/A	Lenz et al. (2012)
<i>Megachile</i> sp.	in situ	N/A	N/A	N/A	N/A	fr	N/A	N/A	N/A	yes	N/A	N/A	N/A	Allasino et al. (2019)
<i>Solenopsis invicta</i>	petri dish	latex	N/A	N/A	fluorescence	ms	0.9..4.5	2.5% w/w (food)	direct	N/A	filtration	N/A	N/A	Glancey et al. (1982)
<i>Rhytidoponera metallica</i>	in situ	N/A	N/A	N/A	N/A	b	N/A	50 items per nest	3 d	yes	N/A	N/A	N/A	Hughes and Westoby (1992)
<i>Alphaenogaster longiceps</i>														
<i>Pheidole</i> sp.														
<i>Rhytidoponera metallica</i>	mesocosm	N/A	N/A	N/A	N/A	diverse	<75.5 cm	N/A	26 mos.	yes	N/A	N/A	N/A	Robins and Robins (2011)
diverse ants	in situ	N/A	N/A	attractant	N/A	b	1.8 cm	N/A	1 d	yes	N/A	N/A	N/A	Angotti et al. (2018)



194 3.2 Other panarthropods

195 Apart from the insects, **Acari** (mites) comprise many abundant soil-living taxa, that feed on
196 litter, fungi and fauna as predators and parasites and are bioindicators, as they are sensitive
197 to changes in the soil physiochemical environment (Gulvik, 2007). Experiments indicated, that
198 mites passively transport MP due to pushing and dragging after attachment to their cuticle, as
199 shown with 80 to 250 μm sized PVC particles in a petri dish experiment without soil (Zhu et
200 al., 2018a). The population within manure pats slightly declines when exposed to mm-sized
201 unweathered PE and PS particles at concentrations of 5 % v/v and declines strongly at
202 ≥ 60 % v/v (Stamatiadis and Dindal, 1990). This could probably be an effect of moisture
203 deficiency due to a reduced water holding capacity in an unnaturally enriched substrate, but
204 not necessarily through plastic intake. In contrast, no data was found on their arachnoid,
205 preying relatives, the order of **Pseudoscorpiones** (false scorpions).

206 Just as many other highly abundant and diverse representatives of the soil mesofauna, the
207 **Oniscideae** (woodlice) contribute to the decomposition of litter by chewing and passage
208 through their digestive system (Warburg, 1987) and react strongly to environmental pollution,
209 as such they are potentially used as bioindicators (van Gestel et al., 2018). They practice a
210 strict selection of natural food sources (Hassall and Rushton, 1984). This is also
211 demonstrated for starch and cellulose based plastic films (4 cm^2), which were consumed and
212 digested in experiments with the model organism *Porcellio scaber*, in contrast to PHB
213 (polyhydroxybutyrate) films, that reduces the feeding rate (Wood and Zimmer, 2014). Smaller
214 PE particles (137 \pm 51 μm and 183 \pm 93 μm) embedded into food pellets (0.4 % w/w) were taken
215 up easily by *Porcellio scaber*, and the smaller fraction caused a slight and non-significant
216 reduction of body mass after 14 days of exposure, but not of feeding, defecation or energy
217 reserves (Kokalj et al., 2018).

218 Other panarthropodean groups are even less studied in terms of MP. We did not find literature
219 on the subphylum of Myriapoda containing the classes of **Diplopoda** (millipedes), **Chilopoda**
220 (centipedes), **Paupoda** and **Symphyla** (pseudocentipedes or symphilids), important litter-
221 feeders and predators within various soil ecosystems.

222 The situation is nearly similar with the phylum of **Tardigrada** (water-bears or tardigrades), that
223 has many ecologically relevant and well studied species feeding on microorganisms and
224 detritus particles. Sparse field research in semisubhydric environments showed no uptake of
225 MP fibres by tardigrada (Gusmão et al., 2016), but comprehensive data on terrestrial soils are
226 lacking.

227 Another branch within the panarthropoda, the phylum of **Onychophora** (velvet worms),
228 comprises primordial invertebrates that are mainly native in litter and soils with high water
229 holding capacity under pleistocene-like forest vegetation within tropical and moderate regions
230 (Monge-Nájera, 1994). As predators, they most likely take up plastic debris appearing within



231 or on their prey, but no studies on MP are available, most likely due to their remote habitats,
232 low abundance and little scientific focus.

233 The **Collembola** (springtails), an abundant, diverse and ubiquitous soil-borne phylum with a
234 broad spectrum of food sources (Hopkin, 1997), also represent an intensively studied group
235 within the Arthropoda. Together with the **Diplura** (which mainly live in tropic and subtropic
236 regions in litter and humid topsoil and feed on fungal hyphae, POM and prey) (Westheide and
237 Rieger, 1996) and the **Protura** (Pass et al., 2011), the Collembola build a morphological
238 group, that exhibit similar ecological functions, such as distribution and decomposition of
239 organic matter as well as the control of fungal abundance (Hopkin, 1997). Springtails provide
240 up to 27 % of the soil biomass and up to 33 % of the total soil respiration (with higher shares
241 in colder ecosystems) (Petersen, 1994) with up to 100000 individuals per square meter
242 (Hopkin, 1997). Thus, their well-being plays an important role for ecosystem functioning.

243 In a petri dish experiment without soil, Maaß et al. (2017) showed the passive transport of
244 urea-formaldehyde particles <400 µm and undefined PET fragments by two Collembola
245 species (*Folsomia candida* and *Proisotoma minuta*) due to attachment, but found no
246 ingestion. Within a soil matrix, trials of Kim and An (2019) indicated hindrance of collembolan
247 migration by larger PS particles (44±39, 282±131 and 676±479 µm) at concentrations of
248 1000 mg kg⁻¹ corresponding to highly contaminated soils. In addition, they found suppressed
249 mobility due to the attachment of even smaller PS microbeads (0.47 to 0.53 µm) at
250 concentrations of 8 mg kg⁻¹ dry soil, which is equivalent to values found in nature. Small
251 particles <50 µm were moved, while larger particles were most likely peeled off. When *F.*
252 *candida* encounters two of its predators, the mites *Damaeus exspinosus* and *Hypoaspis*
253 *aculeifer*, the dispersal of 80 to 250 µm PVC particles is enhanced as shown by Zhu et al.
254 (2018a) in a Petri dish experiment. Without proving the ingestion or the minimal effective MP
255 concentration, Zhu et al. (2018b) published an alteration of the gut microbiome and adverse
256 effects on growth and reproduction of *F. candida* by 80 to 250 µm PVC particles mixed in soil
257 at concentrations of 1000 mg kg⁻¹ dry soil. These data were not considered robust (van Gestel
258 and Selonen, 2018), but fit into a later study that found inhibited reproduction at
259 ≥1000 mg kg⁻¹ and avoidance behavior as well as altered microbiome at ≥5000 mg kg⁻¹ (Ju et
260 al., 2019). Such concentrations can occur in highly contaminated soils (Fuller and Gautam,
261 2016). However, documentations on the active uptake, gnawing and grinding of MP by
262 collembolans proposed by Rillig (2012) is still lacking and also studies on Diplura and Protura.



Table 2: Microplastic studies on Acari, Oniscidea (Onisc.), Tardigrada (T.) and Collembola (fr=fragments, p=particles, mf=microfibres, mb=microbeads, ms=microspheres, s=semisubhydric). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
Acari	microcosm	PE	no	N/A	N/A	fr	<4800 >2000	0..90& v/v (manure)	16 d	N/A	N/A	N/A	≥5% v/v: abundance ↓	Stamatiadis and Dindal (1990)
	petri dish	PS	N/A	no	N/A	p	80..250	5000 items per dish	N/A	yes	N/A	N/A	N/A	Zhu et al. (2018a)
Onisc.	mesocosm	PVC	no	N/A	N/A	fr	4 cm ²	1 item per cosm	14 d	N/A	yes	N/A	feeding ↓	Wood and Zimmer (2014)
	petri dish	PHB	no	N/A	N/A	fr	183±93 137±51	0.4% w/w (food)	14 d	N/A	N/A	N/A	no	Kokaji et al. (2018)
Collembola	in situ	PE	N/A	N/A	N/A	mf	N/A	N/A	N/A	N/A	no	N/A	N/A	Gusmão et al. (2016)
	cup	N/A	N/A	no	N/A	p,fr	<400	2.5..5 mg per cup	N/A	yes	N/A	N/A	N/A	Maaß et al. (2017)
	petri dish	UF, PET	N/A	no	N/A	p	80..250	5000 items per dish	N/A	yes	N/A	N/A	N/A	Zhu et al. (2018a)
	microcosm	PVC	N/A	no	N/A	p	80..250	1000	56 d	N/A	N/A	N/A	microbiome, growth ↑, reproduction ↓	Zhu et al. (2018b)
	microcosm	PVC	N/A	no	N/A	mb	<500	0..10000 0..10000 0..5000	7 d 28 d 28 d	N/A	N/A	N/A	≥5000: avoidance ≥1000: reproduction ↑ ≥5000: microbiome	Ju et al. (2019)
Lobelia sokamensis	soil sample	PS	N/A	carboxyl	fluorescence	mb	0.5	4..8		yes				
		PE	no	N/A	fluorescence	ms	27..32	1000		yes				
		PE	no	N/A	fluorescence	ms	250..300	1000			N/A			
		PS	no	N/A	no	fr	44±39	1000		≤3 min	yes	N/A	avoidance, motivity ↓	Kim and An (2019)
		PS	no	N/A	no	fr	282±131	1000			N/A			
PS	no	N/A	no	no	fr	676±479	1000			N/A				



264 3.3 Annelida

265 Another large group of invertebrates beside the branch of panarthropoda comprises land-
266 based Annelida. Within the Annelida, the **Lumbricidae** (earthworms) comprise a well-studied
267 family (Darwin, 1881; Lavelle et al., 2006), represented in high abundance and diversity in
268 many ecosystems all around the world (Phillips et al., 2019). Earthworms are often used as
269 indicators for soil health (Fründ et al., 2011; Pulleman et al., 2012), as they are ecosystem
270 engineers which through their burrowing activity influence various soil physical, chemical and
271 biological processes (Jouquet et al., 2006; Lavelle et al., 2006).

272 By far the most of the studies on the influence of MP on earthworms are performed with PE
273 and the species *Lumbricus terrestris* or *Eisenia fetida*, but there are also single studies with
274 *Aporrectodea rosea* (Boots et al., 2019) and *Eisenia andrei* (Rodriguez-Seijo et al., 2017) and
275 with the less common species *Metaphire californica* (Wang et al., 2019b). We found one field
276 study of earthworms and MPs (Huerta Lwanga et al., 2017a) among many laboratory
277 experiments with MPs mixed into soil volumes (concentrations ranging up to 20000 mg kg⁻¹
278 dry soil) or applied with litter on top of the soil surface ($\leq 60\%$ w/w). The particles sizes were
279 usually <1 mm in diameter, but some were even up to 2×2 cm², and the duration of
280 experiments was generally 14 to 28 days, few lasted up to 60 days.

281 The uptake of MPs of a broad size range by earthworms was shown in studies based on
282 particles in earthworm casts of *Lumbricus terrestris* (Huerta Lwanga et al., 2016; Cao et al.,
283 2017; Hodson et al., 2017; Rillig et al., 2017; Prendergast-Miller et al., 2019; Yu et al., 2019;
284 Huerta Lwanga et al., 2017a), *Eisenia fetida* (Rodríguez-Seijo et al., 2018; Chen et al., 2020;
285 Wang et al., 2019c), *Eisenia andrei* (Rodriguez-Seijo et al., 2017) and *Metaphire californica*
286 (Wang et al., 2019b). Zhang et al. (2018) showed that relatively large PE particles of
287 1.5×1.5 cm² are not ingested by *Lumbricus terrestris*, but partial ingestion of such large
288 particles of biodegradable MPs does take place after initial weathering in soil or in compost
289 has occurred. In some laboratory experiments, MPs were found in the gut of dissected
290 earthworms (Huerta Lwanga et al., 2016; Hodson et al., 2017; Rodriguez-Seijo et al., 2017),
291 but the concentration of MPs in the gut was not significantly different between treatments nor
292 significantly different from the bulk soil concentration, so there was no evidence of
293 accumulation of MPs in the earthworm bodies (Hodson et al., 2017). Chen et al. (2020)
294 assume an accumulation of MP takes place in *Eisenia fetida*, based on an observed increase
295 of MP concentrations in the casts in the course of 4 weeks. Huerta Lwanga et al. (2017a)
296 supposed an accumulation of MPs in the food chain as the concentration of MPs in chicken
297 gizzards is strongly increased compared to that in the earthworm casts in the same
298 experiments. However, mainly the amount of large particles, i.e. macroplastics, in the gizzards
299 was very large, thus it seems likely that the chicken directly fed on plastics and an
300 accumulation through the food chain cannot be proven with the current knowledge and should
301 be further investigated.



302 Several studies did not find significant negative effects of MPs on earthworms' avoidance
303 behaviour (Judy et al., 2019), nor on growth (Hodson et al., 2017; Rodriguez-Seijo et al.,
304 2017; Judy et al., 2019; Wang et al., 2019c), mortality (Hodson et al. (2017); Rillig et al.
305 (2017); Rodriguez-Seijo et al. (2017); Judy et al. (2019); Prendergast-Miller et al. (2019) or
306 reproduction (Huerta Lwanga et al., 2016; Rodriguez-Seijo et al., 2017). However, other
307 studies do show adverse effects of the uptake of MP in different degrees and on different
308 aspects of earthworms' fitness: A reduced growth was shown by Cao et al. (2017) for *Eisenia*
309 *Fetida* and the mortality increased at an exposure of concentrations ≥ 10000 mg kg⁻¹ dry soil.
310 At lower concentrations no significant effects were found. The growth of *Aporrectodea rosea*
311 was also inhibited when exposed to biodegradable polylactic acid, conventional high-density
312 polyethylene (at 1000 mg kg⁻¹ dry soil), and MP clothing fibers (at 10 mg kg⁻¹ dry soil) (Boots
313 et al., 2019). Huerta Lwanga et al. (2016) showed a decrease in growth and increased
314 mortality at concentrations $\geq 28\%$ w/w in litter and after 60 days, though after just 14 days no
315 mortality occurred in these experiments.

316 In some studies, additional effects such as histopathological changes or stress biomarkers
317 were measured. For *Eisenia fetida* Chen et al. (2020) observed skin damage at
318 1500 mg MP kg⁻¹ in soil, measured an increase in catalase activity and malondialdehyde
319 content at 1000 mg kg⁻¹ and at ≥ 1000 mg kg⁻¹ acetylcholine esterase was significantly
320 stimulated. Wang et al. (2019c) tested *Eisenia fetida* and found that MPs only increased the
321 catalase and peroxidase levels as well as the level of lipid peroxidation and decreased the
322 activity of superoxide dismutase and glutathione S-transferase at an exposure of
323 200000 mg kg⁻¹ dry soil for 14 days. No discernible influence was found at 100000 mg kg⁻¹.
324 However, Rodríguez-Seijo et al. (2018) also found for *Eisenia fetida* a significant positive
325 correlation of MP concentration with different biomarker responses: catalase, glutathione S-
326 transferase, lactate dehydrogenase and thiobarbituric acid reactive substances. In addition,
327 Rodríguez-Seijo et al. (2017) observed histological damage of the gut and occurrence of
328 inflammatory processes as well as an increase of stress response indicators associated with
329 MP exposure of *Eisenia andrei*. For *Lumbricus terrestris* Prendergast-Miller et al. (2019)
330 showed an increase in metallothionein expression at an exposure with ≥ 1000 mg kg⁻¹ dry soil
331 and a decrease in heat shock protein 70 at a concentration of ≥ 10000 mg kg⁻¹.

332 Due to the large differences in experimental conditions – e.g. size of the MPs, addition of MPs
333 to soil or to litter, duration of experiments, earthworm species – the current knowledge is not
334 sufficient to detect whether there is a threshold in MP size and concentration at which the MP
335 become harmful for earthworms and how this threshold differs for different earthworms
336 species and MP shapes. The results of Huerta Lwanga et al. (2016), who found no effects of
337 MPs on earthworms at 14 days, but significant influence on growth and mortality after
338 60 days, indicate the importance of longer measurements. This is consistent with Pelosi et al.



339 (2015), who concluded that the influence of pesticides on earthworm communities should be
340 tested in long term field experiments.

341 Earthworms activity also increased the transport of MP in soil columns to deeper soil layers
342 (Rillig et al., 2017; Yu et al., 2019; Huerta Lwanga et al., 2017b). The smaller the MP the
343 stronger the transport. Particles are transported both actively – ingested and later cast out –
344 and passively after attachment to the earthworm's body or by water flow through the biopores.
345 As Huerta Lwanga et al. (2018) showed that the bacteria in the gut of *Lumbricus terrestris* can
346 decompose MPs, it seems likely that particles taken up at the surface are egested as smaller
347 particles in deeper layers.

348 Microplastics might well serve as a vector for contaminant transport to soil organisms. Though
349 adsorption on plastics was seen to be lower than on the soil matrix, the desorption of Zn was
350 seen to be higher in synthetic earthworm guts. However, there was no measurable negative
351 effect of Zn or the PE on *Lumbricus terrestris* (Hodson et al., 2017). Wang et al. (2019b)
352 studied the influence of MP on arsenic uptake and negative effects on *Metaphire californica*
353 and concluded that MPs decreased the uptake of arsenic and that MPs reduced the influence
354 of arsenic on the gut bacterial communities. Rodríguez-Seijo et al. (2019) showed altered
355 enzyme activities and enhanced avoidance behavior in face of LD-PE pellets spiked with the
356 insecticide chlorpyrifos. Yang et al. (2019a) studied the influence of MPs on the transport of
357 glyphosate, however they mainly showed that the glyphosate transport was increased by
358 earthworm activity, the role of MPs in this transport could not be determined with this study.
359 These studies show that MP might have very different influences on the uptake and the
360 adverse effects of different pollutants on earthworms and further investigation is needed in
361 order to understand the influence of MPs on pollutant transport.

362 In contrast to the recently well-researched Lumbricidae, a near relative, the family of
363 **Megascolecidae** (giant earthworms), is not yet mentioned in literature. Another branch within
364 the Annelida, the small **Enchytraeidae** (potworms), were shown to suffer adverse effects on
365 body weight and microbiome with PS microspheres (0.05 to 0.1 μm) at concentrations of
366 $\geq 10\%$ w/w within their food source, but an unexpected increase of reproduction at 0.5 % w/w
367 (Zhu et al., 2018b). The reproduction was reduced at abnormal concentrations of
368 90 g kg^{-1} dry soil of polyamid particles (13 to 150 μm), but not with PVC (Lahive et al., 2019).

369 The edaphon of semisubhydric soils often became a marginal group between the area of
370 interest of soil and aquatic scientists. As a highly diverse soil biocoenosis outside the focus of
371 this paper, the benthos along seashores and fresh waters is also affected by MPs and should
372 therefore be shortly mentioned by reviewing the lugworm **Arenicola marina**, a well examined
373 deposit-feeder of the tidal flats. In situ, MP accumulates within its tissue and feces (Van
374 Cauwenberghe et al., 2015). In laboratory experiments, PS particles $\geq 500\ \mu\text{m}$ were avoided
375 as food-source and passively translocated within the sediment at concentrations of $\sim 2\text{ g kg}^{-1}$



376 (Gebhardt and Forster, 2018), but were measured within the feces at $\sim 74 \text{ g kg}^{-1}$ causing
377 effects on feeding activity and body weight with no influence on the survival rate (Besseling et
378 al., 2012). PS microspheres $\leq 30 \mu\text{m}$ remained within the animal without any adverse effects
379 regardless of particle size (Van Cauwenberghe et al., 2015). Other studies found adverse
380 effects on respiration, energy reserves, feeding, egestion and casting after uptake of PVC
381 particles $\leq 478 \mu\text{m}$ at different sediment concentrations of $> 2 \text{ g kg}^{-1}$, but neither due to HD-PE
382 nor on biomass and survival (Wright et al., 2013; Green et al., 2016). There is a difficulty in
383 distinguishing between the adverse effects of MPs and substances adsorbed on or leached
384 from MPs (Besseling et al., 2012). When adding PCB-spiked PE to mud flat sediment with
385 concentrations up to 5 g kg^{-1} dry mass, there was no significant change of survival rate or
386 body weight. The decreased feeding activity and heap mass could be attributed to increasing
387 plastic concentrations, but not to enhanced PCB bioaccumulation via PE uptake (Besseling et
388 al., 2017). However, all these studies found adverse effects at MP concentrations orders of
389 magnitude above natural values.



Table 3: Microplastic studies on Lumbricidae (p=particles, ms=microspheres, b=beads, f=fibres, mf=microfibres). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

organism	experimental environment	plastic type	aging (C ₁₂ H ₂₂ O ₄ , C ₁₂ H ₁₆)	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
<i>Lumbricus terrestris</i>	mesocosm	PE	washed (C ₁₂ H ₂₂ O ₄ , C ₁₂ H ₁₆)	N/A	N/A	p	<150	0..60% w/w (litter)	14 d / 60 d	yes	yes	N/A	at 60 d. ≥28% w/w survival ↓, growth ↑	Huerta Lwanga et al. (2016)
<i>Eisenia fetida</i>	glass beaker	PS	N/A	N/A	N/A	ms	50..80	0..20000	30 d	N/A	yes	N/A	≥5000: survival ↓ ≥100000: weight ↓	Cao et al. (2017)
<i>Lumbricus terrestris</i>	bag	PE	N/A	N/A	N/A	p	0.92±1.09 mm ²	3500	28 d	N/A	yes	no	no	Hodson et al. (2017)
<i>Lumbricus terrestris</i>	home yard	diverse	yes	N/A	N/A	N/A	N/A	0.87±1.9 items g ⁻¹	N/A	N/A	yes	conc. in chickens > in earthworms	N/A	Huerta Lwanga et al. (2017a)
<i>Lumbricus terrestris</i>	mesocosm	PE	washed (C ₁₂ H ₂₂ O ₄ , C ₁₂ H ₁₆)	N/A	N/A	p	<150	0..60% w/w (litter)	14 d	yes	yes	N/A	N/A	Huerta Lwanga et al. (2017b)
<i>Lumbricus terrestris</i>	mesocosm	PE	N/A	no	no	b	71.0..2800	750 µg on 2.5 kg soil	21 d	yes	yes	N/A	no	Rillig et al. (2017)
<i>Eisenia andrei</i>	mesocosm	LD-PE	N/A	N/A	N/A	pellets	250..1000	0..1000	28 d	N/A	yes	N/A	≥62.5: intestinal damage	Rodríguez-Sejor et al (2017)
<i>Lumbricus terrestris</i> (gut bacteria)	mesocosm	PE	washed (C ₁₂ H ₂₂ O ₄ , C ₁₂ H ₁₆)	N/A	N/A	p	150	7% w/w (litter)	60 d (earthworms)	N/A	yes	N/A	N/A	Huerta Lwanga et al. (2018)
<i>Eisenia fetida</i>	mesocosm	LD-PE	washed (EtOH)	N/A	N/A	pellets	250..1000	0..1000	28 d	N/A	yes	N/A	≥125: altered enzyme activity	Rodríguez-Sejor et al (2018)
<i>Aporectodea rosea</i>	mesocosm	PLA, PE	N/A	N/A	N/A	p	N/A	1000	30 d	N/A	yes	N/A	growth ↑	Boots et al. (2019)
<i>Eisenia fetida</i>	mesocosm	HD-PE, PET, PVC	no	N/A	no	f	<2000	soil extract	48 h / 56 d	N/A	N/A	N/A	no	Judy et al. (2019)
<i>Lumbricus terrestris</i>	bag	PE	N/A	N/A	N/A	mf	∅40-7±3.8 x 361.6±387.0	0..10000	35 d	N/A	yes	N/A	≥1000: metallothionein expression ↑ ≥10000: heat shock protein 70 ↓	Prendergast-Miller et al. (2019)
<i>Eisenia fetida</i>	mesocosm	LD-PE	washed (EtOH)	N/A	chlopyrifos (CPF)	pellets	5000	40 items on 0.5 kg soil	14 d	N/A	N/A	N/A	with CPF: altered enzyme activity, avoidance of MPs	Rodríguez-Sejor et al (2019)
<i>Metaphire californica</i>	mesocosm	PVC	N/A	N/A	sodium arsenate	p	N/A	2000	28 d	yes	yes	N/A	microbiome	Wang et al. (2019b)
<i>Eisenia fetida</i>	glass beaker	PE	washed (MeOH)	N/A	PAHs, PCBs, NiR Rad (Nf)	p	<300	0..200000	14 d	N/A	yes	N/A	≥200000: altered enzyme activity	Wang et al. (2019c)
<i>Lumbricus terrestris</i>	mesocosm	PE	washed (C ₁₂ H ₂₂ O ₄ , C ₁₂ H ₁₆)	N/A	glyphosate	p	<150	0..7% w/w (litter)	14 d	N/A	N/A	N/A	N/A	Yang et al. (2019a)
<i>Lumbricus terrestris</i>	mesocosm	PE	N/A	N/A	N/A	N/A	<1000	7% w/w (litter)	14 d	yes	yes	N/A	N/A	Yu et al. (2019)
<i>Lumbricus terrestris</i>	petri dish	PE and div. biodegradable	unweathered, field or compost	N/A	N/A	p	1.5x4.5 cm ²	4 items per dish	14 d	yes	no	N/A	N/A	Zhang et al. (2018)
<i>Eisenia fetida</i>	mesocosm	PE	washed (EtOH)	N/A	N/A	p	2x2 cm ²	10 items per dish	50 d	yes	yes	N/A	N/A	Zhang et al. (2018)
<i>Eisenia fetida</i>	bag	PE	washed (EtOH)	N/A	N/A	p	<400	0..1500	28 d	N/A	yes	yes	skin damage, ≥250 mg/kg: oxidative stress ≥1000 mg/kg: neurotoxicity ↑	Chen et al. (2020)



Table 4: Microplastic studies on Enchytraeidae and Arenicola marina (mb=microbeads, p=particles, ms=microspheres, sed.=sediment, s=semisubhydryc). Concentrations refer to mg kg⁻¹ dry soil in terrestrial and dry sediment in semisubhydryc soils, if not specially marked.

organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
Enchytraeus crypticus	petri dish	PS	N/A	N/A	N/A	mb	0.05..0.1	0..10% w/w (food)	7	N/A	yes	N/A	at 0.5% w/w: reproduction ↑ ≥10% w/w: microbiome, weight ↓	Zhu et al. (2018c)
	microcosm	PVC	N/A	N/A	fluorescence	p	13..150 106..150	20000..120000 90000	20 h / 21 d	N/A	yes	N/A	≥90000: reproduction ↓ no	Lahve et al. (2018)
Arenicola marina ^s	in situ	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	yes	1.2±2.8 items g ⁻¹	N/A	Cauwenbergh et al. (2015)
	liquid culture	PS	no	N/A	N/A	ms	10..90	10000..50000 items kg ⁻¹	14 d	N/A	yes	10 µm: 9600±1800 items kg ⁻¹ 30 µm: 800±700 items kg ⁻¹	no	
Arenicola marina ^s	mesocosm	PS	yes	biofilm	N/A	p	500..1000	~2000	106..240 d	yes	no	N/A	N/A	Gebhardt and Forster (2018)
Arenicola marina ^s	mesocosm	PS	N/A	N/A	N/A	p	400..1300	0..74000	28 d	N/A	≥400 µm	no	≥74000: feeding ↑, weight ↓ >2000: respiration ↑, casting ↓	Besseling et al., (2012)
Arenicola marina ^s	mesocosm	HD-PE	N/A	N/A	N/A	p	3..316	0..20000 mg kg ⁻¹ wet sed.	31 d	N/A	N/A	N/A	no	Green et al. (2016)
Arenicola marina ^s	mesocosm	PE	N/A	PCBs	fluorescence	mb	10..180	0..5000	28 d	N/A	yes	no	feeding activity ↑, heap mass ↓	Besseling et al. (2017)
Arenicola marina ^s	mesocosm	PVC	N/A	N/A	not leaching	p	~130	0..50000	28 d	N/A	N/A	N/A	≥10000: energy reserves ↓ ≥50000: feeding ↑, egestion ↑, casting ↓	Wright et al. (2013)



392 3.4 Further invertebrates

393 As part of the microfauna, the phylum **Nematoda** (nematodes or roundworms) is an
394 ecologically important branch containing >25000 species (Zhang, 2013) in freshwater, marine,
395 endobiotic and soil habitats. Due to their diverse trophic interactions nematodes hold a
396 central position in both bottom-up and top-down controlled food webs (Yeates, 2001; Ferris,
397 2010) and thus most likely the uptake and transfer of MP.

398 Active feeding of adults and larvae of different species on 0.5 to 6 μm PS/latex microspheres
399 (the size of their bacterial prey) was proven by Nika et al. (2016) and Fueser et al. (2019).
400 However, most MP experiments on Nematodes are based on the bacterial-feeding model
401 organism *Caenorhabditis elegans*. Kiyama et al. (2012) showed the favored uptake of PS
402 microspheres with sizes of 0.5 to 3 μm by adult and 0.5 μm by larval *C. elegans*. The
403 ingestion of MP decreased in the presence of bacteria as the natural food source.

404 When larval stages and adults ingested PS between 0.05 and 5 μm within an aqueous
405 suspension or on agar plates, adverse effects such as oxidative stress, neurodegeneration,
406 intestinal and DNA damage or dysfunction in motility, growth, life span, defecation,
407 reproduction or energy metabolism appeared from a wide spectrum of concentrations from
408 $\geq 1 \mu\text{g l}^{-1}$ up to $\geq 86.3 \text{ mg l}^{-1}$ (Zhao et al., 2017; Dong et al., 2018; Kim et al., 2019; Lei et al.,
409 2018a; Lei et al., 2018b; Qu et al., 2019a). These effects are missed below $1 \mu\text{g l}^{-1}$ (Qu et al.,
410 2019b), and are enhanced due to amino modifications on micropshere surfaces (Qu et al.,
411 2019c). The incubation on agar plates with PE, PP and PVC particles <70 μm caused similar
412 influences on survival, fertility, brood size and intestinal function (Lei et al., 2018b) Leachates
413 from soils amended with 5 mg kg^{-1} dry soil of HD-PE and PVC decreased reproduction in
414 laboratory cultures, but there was no effect shown on survival and after application of PET
415 (Judy et al., 2019). Furthermore, silica nanoparticles (0.05 μm) are not only taken up orally
416 but also via the vulva and spermathecae and migrate into gonad cells (Scharf et al., 2013),
417 This process was confirmed for PS nanoparticles with the potential of a transfer to the
418 progeny (Zhao et al., 2017).

419 The clear adverse effects of these studies are limited in their representativity by a narrow
420 restriction to liquid cultures and a single model organism lacking broader studies on
421 prominent soil-born nematodes such as *Acrobeloides buetschlii* (Frey, 1971). When assuming
422 in first proximity mg l^{-1} solution = mg kg^{-1} dry soil, the applied concentrations between 0.001
423 and 86.8 mg l^{-1} match lower levels of soil contamination.

424 Feeding studies on the phylum **Rotifera** with MPs are fully based on PS microbeads and
425 model organisms of the planktonic genus *Brachionus*. However, this data can carefully be
426 transferred to soil environments as also soil rotifers are aquatic organisms living in water-filled
427 pores and waterfilms. Different *Brachionus sp.* ingest microbeads <10 μm with strong
428 preference for particles the size of their natural food source, namely bacteria and algae with



429 2 to 5 μm in diameter (Vadstein et al., 1993; Heerkloß and Hlawa, 1995; Baer et al., 2008;
430 Jeong et al., 2016). The uptake appears to be selective as the microbeads are less
431 incorporated than bacteria and algae (Vadstein et al., 1993). The egestion of particles
432 $\leq 0.5 \mu\text{m}$ is hindered compared to 6 μm (Jeong et al., 2016). In suspension, microbeads
433 $\leq 0.5 \mu\text{m}$ cause adverse effects on fertility and life span at $\geq 0.1 \text{ mg l}^{-1}$ as well as oxidative
434 stress and less growth at $\geq 10 \text{ mg l}^{-1}$ (Jeong et al., 2016; Sun et al., 2019).

435 Terrestrial molluscs comprise snails and slugs within the class of **Gastropoda**. These grazers
436 feed on bacterial biofilms, fungi and plant tissue (Parkyn and Newell, 2013). Studies on
437 terrestrial species are sparse, but data on the benthic *Littorina sp.* imply passive transport and
438 non-selective MP uptake by feeding on surfaces with contaminated feces and mucus trails of
439 other snails (Gutow et al., 2019). However, Imhof and Laforsch (2016) found no significant
440 influence on growth parameters and fertility of juveniles and adult *Potampoyrgus antipodarum*
441 even when a food source with 70 % w/w of 5 to 600 μm sized fragments was given (a mixture
442 of PA, PC, PET, PS, PVC). In contrast, adverse effects were found in recent work on the
443 terrestrial snail *Achatina fulica*, that showed uptake and complete gastrointestinal passage
444 within 48 h with partial degradation of PET fibres (appr. 1258x76 μm), but reduced excretion
445 and food intake as well as increased oxidative stress at concentrations of $\geq 0.01 \text{ g kg}^{-1}$,
446 $\geq 0.14 \text{ g kg}^{-1}$ and $\geq 0.71 \text{ g kg}^{-1}$ dry soil, respectively (Song et al., 2019).



Table 5: Microplastic studies on nematods (ms=microspheres, fr=fragments, np=nanoparticles, mb=microbeads, ms=microspheres, ox.=oxidative). Concentrations refer to mg kg⁻¹ dry soil, if not specially marked.

organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
<i>Caenorhabditis elegans</i>	agar plate	PS	N/A	sulfate amino	fluorescence	ms	0.1..6.6	N/A	0.5..2 h	N/A	yes	0.5..3 µm	N/A	Kiyama et al. (2012)
<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	carboxyl	fluorescence	ms	0.1	0.001..10 mg l ⁻¹	4.5 d	N/A	Yes	N/A	≥0.01 mg l ⁻¹ ; motivity ↑, growth ↑, defecation ↓, within gonads	Zhao et al. (2017)
<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	ζ=10mv	fluorescence	ms	0.1	0.00001..0.001 mg l ⁻¹	N/A	N/A	Yes	N/A	≥0.001 mg l ⁻¹ ; motivity ↑, ox. stress ↑	Dong et al. (2018)
<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	N/A	preservatives, fluorescence	ms	0.05..0.2	0.001..86.8 mg l ⁻¹	24 h	N/A	Yes	N/A	≥86.3 mg l ⁻¹ ; ox. stress ↑	Kim et al. (2019)
<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	ζ=10mv	fluorescence	ms	0.1	0.001..1 mg l ⁻¹	N/A	N/A	Yes	N/A	≥17.3 mg l ⁻¹ ; metabolic dyst.	Qu et al. (2019a)
<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	N/A	N/A	ms	0.1..5	1 mg l ⁻¹	3 d	N/A	Yes	N/A	motivity ↑, survival ↓, growth ↑, ox. stress ↑, neurotoxicity	Lei et al. (2018a)
<i>Caenorhabditis elegans</i>	agar plate	PE, PP, PVC, PS	no	N/A	N/A	fr, ms	0.1..200	0.5..10.0 mg m ⁻²	2 d	N/A	Yes	N/A	≥0.5 mg m ⁻² survival ↓ at 5 mg m ⁻² ; growth ↓, fertility ↓, ox. stress ↑, intestinal damage	Lei et al. (2018b)
<i>Caenorhabditis elegans</i>	agar plate	PS	N/A	N/A	fluorescence	ms	0.1..5	2500 mg l ⁻¹	7 d	N/A	Yes	N/A	mainly 2µm; intestinal damage	Scharf et al. (2013)
<i>Caenorhabditis elegans</i>	liquid culture	HD-PE, PET, P/C	no	N/A	no	fr	<2000	soil extract	72 h	N/A	N/A	N/A	fertility ↓	Judy et al. (2019)
<i>Caenorhabditis elegans</i>	agar plates	latex	N/A	N/A	fluorescence	mb	0.5	N/A	30 min	N/A	yes	N/A	N/A	Nika et al. (2016)
<i>Panagrolaimus thienemanni</i>											≤3µm			
<i>Plectus acuminatus</i>											≤0.5µm			
<i>Poikilolaimus regenfussi</i>											≤1µm			
<i>Acrobeloides nanus</i>											≤1µm			
<i>Pristionchus pacificus</i>											no			
<i>Aphelenchoides parietinus</i>											no			
<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	N/A	fluorescence	ms	0.5..6	3·10 ⁸ ..10 ¹⁰ items l ⁻¹ (~0.2..1200 mg l ⁻¹)	4..73 h	N/A	N/A	N/A	N/A	Fueser et al. (2019)
<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	N/A	N/A	ms	0.1	0.0001..0.001 mg l ⁻¹	N/A	N/A	N/A	N/A	no	Qu et al. (2019b)
<i>Caenorhabditis elegans</i>	liquid culture	PS	N/A	amino	N/A	ms	0.1	0.001..1 mg l ⁻¹	N/A	N/A	yes	N/A	≥0.01 mg l ⁻¹ ; fertility ↓, DNA damage	Qu et al. (2019c)



449 3.5 Vertebrates

450 Different taxa of the class of Amphibia have a predator function within the edaphic food web
451 (e.g. preying on invertebrates) (Hebrard et al., 1992). While no data on the reaction to soil
452 MPs are available neither for the legless **Gymnophiona** nor for adults of the order **Anura**,
453 sparse data on tadpoles of aquatic frogs suggest uptake followed by regular excretion of PS
454 microspheres as shown with *Xenopus tropicalis* (Hu et al., 2016). Further, there exist no data
455 on the families **Serpentes** (snakes) and **Anguillidae** within the class of Reptilia, residing at the
456 outer rim of the food web.

457 Within the broad field of Mammalia, studies on MP ingestion are sparse and focus on **mice** as
458 a rodent model organism. Feeding of mice with PS microspheres of 1 to 14 μm in
459 concentrations of 1.49×10^6 to 4.55×10^7 particles at a volume of 10 ml kg^{-1} body weight for
460 4 weeks showed no adverse effects (Stock et al., 2019). In contrast, longer exposition
461 (6 weeks) with lower concentrations of particles with the same shape and size range changed
462 the mouse microbiome and caused metabolic and intestinal dysfunction (Lu et al., 2018; Jin et
463 al., 2019), which comes along with bioaccumulation within organs (Yang et al., 2019b). These
464 studies are regularly conducted with passive feeding and exclude active foraging on
465 perceptible plastic particles. However, the uptake via prey or feeding on contaminated roots
466 and litter is highly probable. Further Rodentia – **Cricetidae** (hamsters, lemmings, voles),
467 **Bathyergidae** (blesmols, mole-rats), **Octodontidae** as well as **Spermophilus** (ground
468 squirrels) and **Marmota** (marmots) within the family of **Sciuridae** – were not yet studied, just
469 as other mammalian (sub)orders like **Chrysochloridae** (golden moles), **Cingulata**
470 (armadillos), **Macroscelidea** (elephant shrews), **Notoryctemorphia** and **Peramelemorphia**.



Table 7: Microplastic studies on Anura (An.) and Rodentia (ms=microspheres, a=aquatic).

organism	experimental environment	plastic type	aging	coating	additives	shape	size span [µm]	concentrations	exposure time	passive transport	active uptake	bioaccum. dynamics	measured adverse effects	reference
<i>Xenopus tropicalis</i> ^a	petri dish	PS	N/A	N/A	fluorescence	ms	1..10	100..10 ⁹ items l ⁻¹ (55..10 ⁹ ..55 mg l ⁻¹)	48 h	N/A	yes	egestion within days	N/A	Hu et al. (2016)
Rodentia	in vivo	PS	N/A	carboxyl sulfate sulfate	fluorescence	ms	1	4.55·10 ⁷ items per mouse (0.025 mg per mouse)	28 d	N/A	yes	N/A	no	Stock et al. (2019)
							4	4.55·10 ⁷ items per mouse (1.6 mg per mouse)	42 d	N/A	yes	N/A	≥0.1 mg l ⁻¹ : microbiome, metabolic dysfunction	Jin et al. (2019)
							10	1.49·10 ⁸ items per mouse (0.8 mg per mouse)	35 d	N/A	N/A	N/A	≥0.1 mg l ⁻¹ : microbiome, metabolic dysfunction	Lu et al. (2018)
<i>Mus musculus</i>	in vivo	PS	N/A	N/A	fluorescence	ms	5..20	200 mg l ⁻¹	28 d	N/A	yes	8x, 8x.5 and 0.71x0.14 mg kg ⁻¹ body weight	N/A	Yang et al. (2019b)



472 **4 Synthesis**

473 **4.1 Summarized observations**

474 Our systematic search comprised recent research on the interaction of soil organisms with
475 MP, but also studies with focus on feeding experiments, that are published much earlier than
476 the awareness on plastic in the environment appeared. The numerous studies found with
477 focus on the ingestion of MPs consistently showed the active uptake by diverse soil
478 organisms with few exceptions spread over the whole branch of invertebrates. In addition,
479 also studies on adverse effects caused by the intake of MP contaminated food (e.g. of food
480 pellets by dung beetles) imply the ingestion into the test organism. Distinct size preferences
481 are measured for dung beetles, nematodes, rotifers and ants showing that mainly particles
482 are ingested, that are small enough to enter the gastrointestinal tract. In contrast, active
483 comminution by gnawing on larger particles was tested only for a few taxa and confirmed for
484 woodlice, termites and mealworms, and in the case of earthworms only after initial
485 weathering.

486 After the ingestion, MP is understandably translocated actively until excretion or death of the
487 transporting organism, which was only directly shown in experiments with earthworms. The
488 passive transport by attachment, dragging and pushing was checked in a few experiments
489 with earthworms, mites and springtails that partly worked without soil substrate and
490 consistently showed positive results.

491 After exposition to MP, a pattern of adverse effects can be seen: Across various taxa, altered
492 microbiomes, reduced motility, body mass, fertility and life span as well as increased oxidative
493 stress and metabolic malfunctioning occur in different combinations mainly due to μm -sized
494 MP in and above the whole known natural range of concentration. For some taxa such as
495 Nematodes, Gastropoda and Rotifera these effects appear at natural and increased MP
496 concentrations ($<100 \text{ mg kg}^{-1}$ dry soil), for Collembola and Lumbricidae at concentrations like
497 in highly contaminated sites ($\geq 1000 \text{ mg kg}^{-1}$ dry soil) and for Enchytraeidae, *Arenicola marina*
498 and in further experiments with earthworms at unplausible high values. The data show a
499 tendency, that the effects occur at lower concentrations, when the added particles are smaller.
500 Small sized particles also provide the highest surface/volume ratio and thus the highest
501 reactive surface per weight.

502 Most studies work with defined increasing MP concentrations and particle sizes in soil
503 substrates and food sources, which can be used to determine relationships between
504 environmental concentrations and adverse effects. However, the lack of information about
505 intake rates, grades of accumulation and effective prey-predator transfer leads to a gap within
506 the chain of explanation for toxic effects on the soil organisms. In some experiments, the
507 intestinal passage of MP and sizes preferably retained within the gut were shown, but there
508 are no experiments that could demonstrate quantitative bioaccumulation. In contrast,



509 quantifications of the retained and egested MP particle size fractions might be biased due to
510 gnawing and intestinal comminution as shown for woodlice, termites, mealworms, snails and
511 earthworms.

512 In order to improve our understanding of processes underlying adverse effects of MP on soil
513 organisms, data on ingestion rates, dwell times, biodegradation and egestion rates are
514 important bricks e.g. to reveal bioaccumulation dynamics. However, there are only a few data
515 on biodegradation (mealworms, snails, earthworms), egestion (rotifers, frogs, snails,
516 earthworms) and remaining concentrations in the body (lugworm, mice, earthworms).



517 **4.2 Limitations of previous studies**

518 The available studies worked with items within the full size span of micro- and nanoplastics
519 ($\leq 5000 \mu\text{m}$). When MP $\geq 50 \mu\text{m}$ was applied, mainly particles and fragments made of PE and
520 PVC were used, whereas PS/latex microspheres were mainly applied for sizes $\leq 10 \mu\text{m}$
521 (*Table 8*). The latter are readily available, highly standardized and are mostly used with
522 fluorescent dyes and either hydrophobic, carboxylated or, more rarely, with amino or sulfate
523 groups. However, there are indications that the spectrum of particle type and shape used in
524 experiments does not correspond to the properties of particles in soils. In different natural as
525 well as agriculturally and industrially contaminated terrestrial and semi-subhydric sites, fibers
526 and fragments of PE and PP, mostly $\leq 100 \mu\text{m}$, were much more abundant than PVC, PET
527 and PS items (Claessens et al., 2011; Vianello et al., 2013; Nor and Obbard, 2014; Najji et al.,
528 2017; Zhang and Liu, 2018; Li et al., 2018a). This is probably caused by high loads of MP
529 fibers in discharged waste water and sewage sludge, which is used in agricultural sites
530 worldwide (Mahon et al., 2016; Li et al., 2018b). It is likely that shape plays an important role
531 for the ingestion of MP items. Unfortunately, we did not find studies that have carried out a
532 complete classification of sampling sites according to plastic origin, size and type, that could
533 help to evaluate differences between former experimental and natural plastic composition to
534 achieve the most realistic experimental conditions. Little knowledge about the size distribution
535 of MP in soils furthermore complicates the determination of realistic concentrations for the
536 addition of a certain particle size spectrum. All reviewed studies either arbitrarily set their
537 applied concentrations or had to base them on measurements of total specific MP masses,
538 regardless of how much of this mass is in the tested size range. This may lead to a
539 malestimation of total adverse MP concentrations.

540 In contrast to particle type and shape, the documentation of chemical properties of MP
541 samples in most of these studies is fragmentary. Some experiments explicitly mentioned that
542 the added plastic was unweathered, whereas most studies lack of information about the
543 degree of aging implying that unweathered items were used. Only a few experiments involved
544 aging of MP, but without comparison to results of natural weathering (Tsunoda et al., 2010;
545 Gebhardt and Forster, 2018). That is in conflict with natural conditions, as plastic that
546 remains within the soil after littering, sewage sludge application or plastic mulching shows
547 signs of weathering, e.g. modified carbonyl indices (Andrady, 2017), while unweathered soil
548 MP might be rare. In addition, Zhang et al. (2018) showed that earthworms actively
549 comminute only weathered bioplastics. In experiments using PS microspheres, the reduced
550 hydrophobicity due to weathering is therefor imitated by means of surface carboxylation.

551 Weathering of MP surfaces within soils comes along with biofilm growth and adsorption of
552 organic molecules, which could potentially affect the attractiveness or toxicity for grazers and
553 other organisms. Such coatings were applied only in a few cases (Besseling et al., 2017;
554 Angotti et al., 2018; Gebhardt and Forster, 2018), but were not documented in most studies.



555 Similarly, the type and concentration of additives such as flame retardants, anti-oxidants or
556 stabilizers often remained undocumented, with exception of fluorescent dyes, that are well
557 mentioned. The release of additives can have a harmful effect on the test organism, as shown
558 for aquatic environments (e Silva et al., 2016). Some studies on the ingestion of MP by the
559 soil mesofauna indicate that the diameter of the gastrointestinal tract is a useful upper size
560 limit for added particles, as far as the organism is unable to crush them (Heerkloß and Hlawa,
561 1995; Holter, 2000; Holter et al., 2002; Holter and Scholtz, 2005; Baer et al., 2008; Fueser et
562 al., 2019). However, using only ingestible particle sizes in their natural concentrations neglect
563 the adverse effects of plastic leachates, which can also get into the soil solution and onto the
564 mineral phase from larger particles and affect soil life.

565 The conditions of incubation differ considerably in terms of habitats and duration of exposure.
566 In most studies, the exposure ranges from a few minutes to a few days in experiments with
567 micro- and small mesofauna and hours to several weeks in experiments with large meso- and
568 macrofauna and is mainly based on excretion or reproductiv cycles. Long-term studies, which
569 are indeed difficult to carry out in mesocosms, practically do not exist. However, certain
570 adverse effects might only establish themselves after long term trials, as was shown for the
571 influence of pesiticides (Pelosi et al., 2015).

572 Some experiments were carried out in soil-free test environments such as liquid cultures or
573 petri dishes with nutrient solutions or a specific food source (nematods, rotifers, mice). By
574 this, motivity is less restricted and feeding behavior can be altered compared to cultivation
575 within soil environments. For example, the ingestion of MP by nematods decreases in the
576 presence of an alternative and more natural food source like bacteria, which can significantly
577 reduce the bioaccumulation and thus the effective toxicity (Kiyama et al., 2012). This can lead
578 to less consumption of MP in soil environments and an overestimation of the toxicity in liquid
579 culture experiments. Also, all laboratory feeding experiments were carried out by use of only
580 one species. The complexity of the food web in soils is thereby excluded and the potential
581 accumulation from prey to predators still unexplored.

582

583

584

585



Table 8: Types and shapes of microplastic particles in edaphon studies. (X) symbolizes combinations outside the search pattern, the number counts for studies, empty fields stand for zero results. Microbeads and microspheres are often mixed up terms and, thus, counted together.

Linné's systematic names OR common name		fragments	particles	fibres	microfibres	beads	microspheres	other, diverse, N/A
plastic	X							
microplastic								
nanoplastic								
PE OR polyethylene	X	4	10	1	1	1	4	7
PP OR polypropylene	X	1						
PVC OR polyvinyl chloride	X	4	6	1				
PS OR polystyrene	X	6	3				24	4
PU OR polyurethane	X							
PET OR polyethylene terephthalate	X	3		2			X	
latex	X				X		6	
other		6	3		1			1
N/A		1		1		2		3

586



587 **4.3 Pinpoints for future research**

588 Most studies reviewed in this work have a pioneering role in MP research and, thus, are
589 subject to some experimental limitations caused by an early state of knowledge. The adverse
590 effects recently found are alarming, but must be considered under the restrictions named
591 above. We propose the following points as part of a *modus operandi* for future MP research.

592 In past studies, particular adverse effects of MP were measured only for certain sizes,
593 shapes, coatings, leachates or adsorbed substances (*Tables 1 to 7*). Experimental
594 concentrations were assumed randomly or derived from cumulative concentrations of one or
595 more MP types measured in natural soils (approx. 1 to some 1000 mg kg⁻¹ dry soil),
596 regardless of size. For those specific experiments coming, the spectrum of concentrations
597 used should be adapted to the quantities of the size spectrum occurring within the soil. For
598 future studies on mixed contaminations, we recommend to evaluate the overall adverse
599 effects of PE, PP, PVC, PET, PU and PS to certain test organisms by use of typical MP-
600 specific spans of concentration, size and shape distribution in natural soils or food samples.
601 This previously requires well-structured data of appropriate MP type, shape and size for
602 different soils in differently contaminated areas.

603 Experiments on adverse effects should be applied within soil matrices to allow the interplay of
604 plastic, natural organic and mineral matter. The MP should be weathered, as plastic in soils
605 underlie broad environmental aging. Pre-weathering of MP should therefor not only be
606 performed in climate chambers (e.g. following DIN EN ISO 4892-2/3), but also include
607 subsequent leaching and equilibration of additives or coatings within the soil matrix before the
608 main experiment. Furthermore, the experimental design may consider coatings with biofilms
609 or attractants and even particle color to regulate the preference of the test organisms.

610 Most detailed information about ingestion are available for dung beetles, nematods and
611 earthworms, data on adverse effects on nematods, earthworms, lugworms and collembola.
612 Future experiments should focus on more ecologically relevant taxa with emphasis on uptake,
613 accumulation and key adverse effects like on survival rate, motility, growth and fertility as well
614 as on the stability of the intestinal microbiome. Further studies with more than one test
615 organism are important to foster our understanding of MP within certain food chains. Also
616 long-term experiments might reveal adverse effects, which evolve slowly within populations.
617 This may enable the assesement of the distribution and effects of MP within the food web
618 and the resulting long-term impact on soil ecosystems.



619 **5 Conclusion**

620 Our review of 77 available studies on the impact of microplastic on the soil fauna shows an
621 alarming diversity and distribution of adverse effects within the soil tree of life. However, these
622 effects have to be considered carefully, as many experiments did not work with plastic
623 matching properties within natural soils and found adverse effects only at concentrations like
624 in highly contaminated soils or above. To elucidate effective concentrations for short and long-
625 term effects on soil faunal health, the most exact reproduction of plastic properties within the
626 soil matrix and natural living conditions of the test organisms is necessary. For future
627 experiments we therefore recommend to choose compositions of type, shape, size, grade of
628 weathering, leachability and coating with biofilms and other organic matter as expected in the
629 habitat to be examined. Furthermore, coming studies should include long-term exposure and
630 food chain experiments to get a better look at the effect of even smaller MP concentrations
631 and their enrichment within the food web. This may give us a better way of assessing the
632 impact of global microplastic contamination on e.g. soil biodiversity, soil carbon cycles and
633 soil quality.



634 **Author contribution**

635 Frederick Büks developed the review concept, collected data and prepared the manuscript
636 except for earthworms. Nicolette Loes van Schaik did all the work on earthworms. Martin
637 Kaupenjohann supervised the whole study.

638 The authors declare that they have no conflict of interest.

639

640

641 **Acknowledgement**

642 Many thanks to Ivica Letunic, who kindly gave us access to phyloT and made our work in face
643 of New Year much easier. I will not forget to invite you for a coffee.

644

645

646 **Competing interests**

647 The authors declare that they have no conflict of interest.



648 **References**

- 649 Allasino M. L., Marrero H. J., Dorado J. and Torretta J. P.: Scientific note: first global report of
650 a bee nest built only with plastic, *Apidologie*, 50, 230-233, doi:10.1007/s13592-019-00635-6,
651 2019.
- 652 Andrad A. L.: The plastic in microplastics: a review, *Marine Pollution Bulletin*, 119, 12-22,
653 doi:10.1016/j.marpolbul.2017.01.082, 2017.
- 654 Angotti M. A., Rabello A., Santiago G. and Ribas C.: Seed removal by ants in Brazilian
655 savanna: optimizing fieldwork, *Sociobiology*, 65, 155-161,
656 doi:10.13102/sociobiology.v65i2.1938, 2018.
- 657 Avery S. V., Harwood J. L. and Lloyd D.: Quantification and Characterization of Phagocytosis
658 in the Soil Amoeba *Acanthamoeba castellanii* by Flow Cytometry, *Appl. Environ. Microbiol.*,
659 61, 1124-1132, doi:N/A, 1995.
- 660 Baer A., Langdon C., Mills S., Schulz C. and Hamre K.: Particle size preference, gut filling and
661 evacuation rates of the rotifer *Brachionus* "Cayman" using polystyrene latex beads,
662 *Aquaculture*, 282, 75-82, doi:10.1016/j.aquaculture.2008.06.020, 2008.
- 663 Bertling J., Bertling R. and Hamann L.: Kunststoffe in Der Umwelt: Mikro- und Makroplastik.
664 Ursachen, Mengen, Umweltschicksale, Wirkungen, Lösungsansätze, Empfehlungen,
665 doi:10.24406/UMSICHT-N-497117, 2018.
- 666 Besseling E., Foekema E. M., van den Heuvel-Greve M. J. and Koelmans A. A.: The effect of
667 microplastic on the uptake of chemicals by the lugworm *Arenicola marina* (L.) under
668 environmentally relevant exposure conditions, *Environmental Science & Technology*, 51,
669 8795-8804, doi:10.1021/acs.est.7b02286, 2017.
- 670 Besseling E., Wegner A., Foekema E. M., Van Den Heuvel-Greve M. J. and Koelmans A. A.:
671 Effects of microplastic on fitness and PCB bioaccumulation by the lugworm *Arenicola marina*
672 (L.), *Environmental Science & Technology*, 47, 593-600, doi:10.1021/es302763x, 2012.
- 673 Boots B., Russell C. W. and Green D. S.: Effects of Microplastics in Soil Ecosystems: Above
674 and Below Ground, *Environmental Science & Technology*, 53, 11496-11506,
675 doi:10.1021/acs.est.9b03304, 2019.
- 676 Bowers B. and Olszewski T. E.: *Acanthamoeba* discriminates internally between digestible
677 and indigestible particles, *The Journal of Cell Biology*, 97, 317-322, doi:10.1083/jcb.97.2.317,
678 1983.
- 679 Brandon A. M., Gao S.-H., Tian R., Ning D., Yang S.-S., Zhou J., Wu W.-M. and Criddle C. S.:
680 Biodegradation of polyethylene and plastic mixtures in mealworms (larvae of *Tenebrio molitor*)



- 681 and effects on the gut microbiome, *Environmental Science & Technology*, 52, 6526-6533,
682 doi:10.1021/acs.est.8b02301, 2018.
- 683 Cao D., Wang X., Luo X., Liu G. and Zheng H.: Effects of polystyrene microplastics on the
684 fitness of earthworms in an agricultural soil, In: IOP Conference Series: Earth and
685 Environmental Science, doi::10.1088/1755-1315/61/1/012148, 2017.
- 686 Chae Y. and An Y.-J.: Current research trends on plastic pollution and ecological impacts on
687 the soil ecosystem: A review, *Environmental Pollution*, 240, 387-395,
688 doi:10.1016/j.envpol.2018.05.008, 2018.
- 689 Chae Y., Kim D., Kim S. W. and An Y.-J.: Trophic transfer and individual impact of nano-sized
690 polystyrene in a four-species freshwater food chain, *Scientific Reports*, 8, 284,
691 doi:10.1038/s41598-017-18849-y, 2018.
- 692 Chen Y., Liu X., Leng Y. and Wang J.: Defense responses in earthworms (*Eisenia fetida*)
693 exposed to low-density polyethylene microplastics in soils, *Ecotoxicology and Environmental*
694 *Safety*, 187, 109788, doi:10.1016/j.ecoenv.2019.109788, 2020.
- 695 Claessens M., De Meester S., Van Landuyt L., De Clerck K. and Janssen C. R.: Occurrence
696 and distribution of microplastics in marine sediments along the Belgian coast, *Marine*
697 *Pollution Bulletin*, 62, 2199-2204, doi:10.1016/j.marpolbul.2011.06.030, 2011.
- 698 Darwin C.: The formation of vegetable mould through the action of worms, with observations
699 on their habits, John Murray, London, 1881.
- 700 De Bruyn L. and Conacher A. J.: The role of termites and ants in soil modification-a review,
701 *Soil Research*, 28, 55-93, doi:10.1071/SR9900055, 1990.
- 702 Dong S., Qu M., Rui Q. and Wang D.: Combinational effect of titanium dioxide nanoparticles
703 and nanopolystyrene particles at environmentally relevant concentrations on nematode
704 *Caenorhabditis elegans*, *Ecotoxicology and Environmental Safety*, 161, 444-450,
705 doi:10.1016/j.ecoenv.2018.06.021, 2018.
- 706 Elloway E. A., Bird R. A., Hewitt C. J., Kelly S. L. and Smith S. N.: Characterization of
707 *Acanthamoeba*--microsphere association by multiparameter flow cytometry and confocal
708 microscopy, *Cytometry Part A*, 69, 266-272, doi:10.1002/cyto.a.20210, 2006.
- 709 Fenchel T.: Suspension feeding in ciliated protozoa: functional response and particle size
710 selection, *Microbial Ecology*, 6, 1-11, doi:10.1007/BF02020370, 1980.
- 711 Ferris H.: Contribution of nematodes to the structure and function of the soil food web, *Journal*
712 *of Nematology*, 42, 63, doi:N/A, 2010.
- 713 Frey F.: The suitability of *Acroboloides buetschlii* for ncmatological experiments,
714 *Nematologica*, 17, 474-477, doi:N/A, 1971.



- 715 Fründ H.-C., Graefe U. and Tischer S.: Earthworms as bioindicators of soil quality, In: Biology
716 of earthworms, Springer, doi:10.1007/978-3-642-14636-7_16, 2011.
- 717 Fueser H., Mueller M.-T., Weiss L., Höss S. and Traunspurger W.: Ingestion of microplastics
718 by nematodes depends on feeding strategy and buccal cavity size, Environmental Pollution,
719 255, 113227, doi:10.1016/j.envpol.2019.113227, 2019.
- 720 Fuller S. and Gautam A.: A procedure for measuring microplastics using pressurized fluid
721 extraction, Environmental Science & Technology, 50, 5774-5780,
722 doi:10.1021/acs.est.6b00816, 2016.
- 723 Garcés-Ordóñez O., Castillo-Olaya V. A., Granados-Briceño A. F., Garcéa L. M. B. and Díaz
724 L. F. E.: Marine litter and microplastic pollution on mangrove soils of the Ciénaga Grande de
725 Santa Marta, Colombian Caribbean, Marine Pollution Bulletin, 145, 455-462,
726 doi:10.1016/j.marpolbul.2019.06.058, 2019.
- 727 Gebhardt C. and Forster S.: Size-selective feeding of *Arenicola marina* promotes long-term
728 burial of microplastic particles in marine sediments, Environmental Pollution, 242, 1777-1786,
729 doi:10.1016/j.envpol.2018.07.090, 2018.
- 730 van Gestel C. A., Loureiro S. and others: Terrestrial isopods as model organisms in soil
731 ecotoxicology: a review, ZooKeys, 127, doi:10.3897/zookeys.801.21970, 2018.
- 732 van Gestel C. A. and Selonen S.: Ecotoxicological effects of microplastics in soil: Comments
733 on the paper by Zhu et al.(2018) 'Exposure of soil collembolans to microplastics perturbs their
734 gut microbiota and alters their isotopic composition', Soil Biology & Biochemistry 116,
735 302-310, Soil Biology and Biochemistry, 124, 116-117, doi:10.1016/j.soilbio.2018.05.032,
736 2018.
- 737 Geyer R., Jambeck J. R. and Law K. L.: Production, use, and fate of all plastics ever made,
738 Science Advances, 3, e1700782, doi:10.1126/sciadv.1700782, 2017.
- 739 Glancey B. M., Vander Meer R., Glover A., Lofgren C. and Vinson S.: Filtration of
740 microparticles from liquids ingested by the red imported fire ant *Solenopsis invicta* Buren,
741 Insectes Sociaux, 28, 395-401, doi:10.1007/BF02224196, 1981.
- 742 Green D. S., Boots B., Sigwart J., Jiang S. and Rocha C.: Effects of conventional and
743 biodegradable microplastics on a marine ecosystem engineer (*Arenicola marina*) and
744 sediment nutrient cycling, Environmental Pollution, 208, 426-434,
745 doi:10.1016/j.envpol.2015.10.010, 2016.
- 746 Gulvik M.: Mites (Acari) as indicators of soil biodiversity and land use monitoring: a review,
747 Polish Journal of Ecology, 55, 415-440, doi:N/A, 2007.



- 748 Gusmão F., Di Domenico M., Amaral A. C. Z., Martínez A., Gonzalez B. C., Worsaae K., do
749 Sul J. A. I. and da Cunha Lana P.: In situ ingestion of microfibres by meiofauna from sandy
750 beaches, *Environmental Pollution*, 216, 584-590, doi:10.1016/j.envpol.2016.06.015, 2016.
- 751 Gutow L., Bartl K., Saborowski R. and Beermann J.: Gastropod pedal mucus retains
752 microplastics and promotes the uptake of particles by marine periwinkles, *Environmental*
753 *Pollution*, 246, 688-696, doi:10.1016/j.envpol.2018.12.097, 2019.
- 754 Hassall M. and Rushton S.: Feeding behaviour of terrestrial isopods in relation to plant
755 defences and microbial activity, In: *Symposia of the Zoological Society of London*, doi:N/A,
756 1984.
- 757 Hebrard J. J., Maloiy G. M. and Alliangana D. M.: Notes on the habitat and diet of *Afrocaecilia*
758 *taimana* (Amphibia: Gymnophiona), *Journal of Herpetology*, 513-515, doi:10.2307/1565136,
759 1992.
- 760 Heerkloß R. and Hlawka S.: Feeding biology of two brachionid rotifers: *Brachionus*
761 *quadridentatus* and *Brachionus plicatilis*, *Hydrobiologia*, 313, 219-221,
762 doi:10.1007/BF00025954, 1995.
- 763 Hodson M. E., Duffus-Hodson C. A., Clark A., Prendergast-Miller M. T. and Thorpe K. L.:
764 Plastic bag derived-microplastics as a vector for metal exposure in terrestrial invertebrates,
765 *Environmental Science & Technology*, 51, 4714-4721, doi:10.1021/acs.est.7b00635, 2017.
- 766 Holter P.: Particle feeding in *Aphodius* dung beetles (Scarabaeidae): old hypotheses and new
767 experimental evidence, *Functional Ecology*, 631-637, doi:N/A, 2000.
- 768 Holter P. and Scholtz C.: Are ball-rolling (Scarabaeini, Gymnopleurini, Sisyphini) and
769 tunnelling scarabaeine dung beetles equally choosy about the size of ingested dung
770 particles?, *Ecological Entomology*, 30, 700-705, doi:10.1111/j.0307-6946.2005.00746.x, 2005.
- 771 Holter P., Scholtz C. and Wardhaugh K.: Dung feeding in adult scarabaeines (tunnellers and
772 endocoprids): even large dung beetles eat small particles, *Ecological Entomology*, 27, 169-
773 176, doi:10.1046/j.1365-2311.2002.00399.x, 2002.
- 774 Hopkin S. P.: *Biology of the springtails (Insecta: Collembola)*, OUP Oxford, 1997.
- 775 Hu L., Su L., Xue Y., Mu J., Zhu J., Xu J. and Shi H.: Uptake, accumulation and elimination of
776 polystyrene microspheres in tadpoles of *Xenopus tropicalis*, *Chemosphere*, 164, 611-617,
777 doi:<https://doi.org/10.1016/j.chemosphere.2016.09.002>, 2016.
- 778 Huerta Lwanga E., Gertsen H., Gooren H., Peters P., Salánki T., van der Ploeg M., Besseling
779 E., Koelmans A. A. and Geissen V.: Microplastics in the terrestrial ecosystem: implications for
780 *Lumbricus terrestris* (Oligochaeta, Lumbricidae), *Environmental Science & Technology*, 50,
781 2685-2691, doi:10.1021/acs.est.5b05478, 2016.



- 782 Huerta Lwanga E., Gertsen H., Gooren H., Peters P., Salánki T., van der Ploeg M., Besseling
783 E., Koelmans A. A. and Geissen V.: Incorporation of microplastics from litter into burrows of
784 *Lumbricus terrestris*, *Environmental Pollution*, 220, 523-531,
785 doi:10.1016/j.envpol.2016.09.096, 2017b.
- 786 Huerta Lwanga E., Thapa B., Yang X., Gertsen H., Salánki T., Geissen V. and Garbeva P.:
787 Decay of low-density polyethylene by bacteria extracted from earthworm's guts: A potential for
788 soil restoration, *Science of the Total Environment*, 624, 753-757,
789 doi:10.1016/j.scitotenv.2017.12.144, 2018.
- 790 Huerta Lwanga E., Vega J. M., Quej V. K., de los Angeles Chi J., del Cid L. S., Chi C., Segura
791 G. E., Gertsen H., Salánki T., van der Ploeg M. and others: Field evidence for transfer of
792 plastic debris along a terrestrial food chain, *Scientific Reports*, 7, 14071, doi:10.1038/s41598-
793 017-14588-2, 2017a.
- 794 Hughes L. and Westoby M.: Effect of diaspore characteristics on removal of seeds adapted
795 for dispersal by ants, *Ecology*, 73, 1300-1312, doi:10.2307/1940677, 1992.
- 796 Imhof H. K. and Laforsch C.: Hazardous or not - Are adult and juvenile individuals of
797 *Potamopyrgus antipodarum* affected by non-buoyant microplastic particles?, *Environmental*
798 *Pollution*, 218, 383-391, doi:10.1016/j.envpol.2016.07.017, 2016.
- 799 Jambeck J. R., Geyer R., Wilcox C., Siegler T. R., Perryman M., Andrady A., Narayan R. and
800 Law K. L.: Plastic waste inputs from land into the ocean, *Science*, 347, 768-771,
801 doi:10.1126/science.1260352, 2015.
- 802 Jeong C.-B., Won E.-J., Kang H.-M., Lee M.-C., Hwang D.-S., Hwang U.-K., Zhou B., Souissi
803 S., Lee S.-J. and Lee J.-S.: Microplastic size-dependent toxicity, oxidative stress induction,
804 and p-JNK and p-p38 activation in the monogonont rotifer (*Brachionus koreanus*),
805 *Environmental Science & Technology*, 50, 8849-8857, doi:10.1021/acs.est.6b01441., 2016.
- 806 Jin Y., Lu L., Tu W., Luo T. and Fu Z.: Impacts of polystyrene microplastic on the gut barrier,
807 microbiota and metabolism of mice, *Science of the Total Environment*, 649, 308-317,
808 doi:10.1016/j.scitotenv.2018.08.353, 2019.
- 809 Jonsson P. R.: Particle size selection, feeding rates and growth dynamics of marine
810 planktonic oligotrichous ciliates (Ciliophora: Oligotrichina), *Mar Ecol Prog Ser*, 33, 265-277,
811 doi:N/A, 1986.
- 812 Jouquet P., Dauber J., Lagerlöf J., Lavelle P. and Lepage M.: Soil invertebrates as ecosystem
813 engineers: intended and accidental effects on soil and feedback loops, *Applied Soil Ecology*,
814 32, 153-164, doi:10.1016/j.apsoil.2005.07.004, 2006.



- 815 Ju H., Zhu D. and Qiao M.: Effects of polyethylene microplastics on the gut microbial
816 community, reproduction and avoidance behaviors of the soil springtail, *Folsomia candida*,
817 *Environmental Pollution*, 247, 890-897, doi:10.1016/j.envpol.2019.01.097, 2019.
- 818 Judy J. D., Williams M., Gregg A., Oliver D., Kumar A., Kookana R. and Kirby J. K.:
819 Microplastics in municipal mixed-waste organic outputs induce minimal short to long-term
820 toxicity in key terrestrial biota, *Environmental Pollution*, 252, 522-531,
821 doi:10.1016/j.envpol.2019.05.027, 2019.
- 822 Kale S. K., Deshmukh A. G., Dudhare M. S. and Patil V. B.: Microbial degradation of plastic: a
823 review, *Journal of Biochemical Technology*, 6, 952-961, doi:N/A, 2015.
- 824 Kim H. M., Lee D.-K., Long N. P., Kwon S. W. and Park J. H.: Uptake of nanopolystyrene
825 particles induces distinct metabolic profiles and toxic effects in *Caenorhabditis elegans*,
826 *Environmental Pollution*, 246, 578-586, doi:10.1016/j.envpol.2018.12.043, 2019.
- 827 Kim S. W. and An Y.-J.: Soil microplastics inhibit the movement of springtail species,
828 *Environment International*, 126, 699-706, doi:10.1016/j.envint.2019.02.067, 2019.
- 829 Kiyama Y., Miyahara K. and Ohshima Y.: Active uptake of artificial particles in the nematode
830 *Caenorhabditis elegans*, *Journal of Experimental Biology*, 215, 1178-1183,
831 doi:10.1242/jeb.067199, 2012.
- 832 Kokalj A. J., Horvat P., Skalar T. and Kržan A.: Plastic bag and facial cleanser derived
833 microplastic do not affect feeding behaviour and energy reserves of terrestrial isopods,
834 *Science of the Total Environment*, 615, 761-766, doi:10.1016/j.scitotenv.2017.10.020, 2018.
- 835 Korb J.: The ecology of social evolution in termites, In: *Ecology of social evolution*, Springer,
836 doi:10.1007/978-3-540-75957-7_7, 2008.
- 837 Kumar A., Pandey A. K., Singh S. S., Shanker R. and Dhawan A.: Cellular uptake and
838 mutagenic potential of metal oxide nanoparticles in bacterial cells, *Chemosphere*, 83, 1124-
839 1132, doi:10.1016/j.chemosphere.2011.01.025, 2011.
- 840 Lahive E., Walton A., Horton A. A., Spurgeon D. J. and Svendsen C.: Microplastic particles
841 reduce reproduction in the terrestrial worm *Enchytraeus crypticus* in a soil exposure,
842 *Environmental Pollution*, 255, 113174, doi:10.1016/j.envpol.2019.113174, 2019.
- 843 Lavelle P., Decaëns T., Aubert M., Barot S., Blouin M., Bureau F., Margerie P., Mora P. and
844 Rossi J.-P.: Soil invertebrates and ecosystem services, *European Journal of Soil Biology*, 42,
845 S3-S15, doi:10.1016/j.ejsobi.2006.10.002, 2006.
- 846 Lavin D. P., Fredrickson A. and Srien F.: Flow cytometric measurement of rates of particle
847 uptake from dilute suspensions by a ciliated protozoan, *Cytometry*, 11, 875-882,
848 doi:10.1002/cyto.990110804, 1990.



- 849 Lei L., Liu M., Song Y., Lu S., Hu J., Cao C., Xie B., Shi H. and He D.: Polystyrene (nano)
850 microplastics cause size-dependent neurotoxicity, oxidative damage and other adverse
851 effects in *Caenorhabditis elegans*, *Environmental Science: Nano*, 5, 2009-2020,
852 doi:10.1039/C9EN00473D, 2018a.
- 853 Lei L., Wu S., Lu S., Liu M., Song Y., Fu Z., Shi H., Raley-Susman K. M. and He D.:
854 Microplastic particles cause intestinal damage and other adverse effects in zebrafish *Danio*
855 *rerio* and nematode *Caenorhabditis elegans*, *Science of the Total Environment*, 619, 1-8,
856 doi:10.1016/j.scitotenv.2017.11.103, 2018b.
- 857 Lenz M., Creffield J. W., Evans T. A., Kard B., Vongkaluang C., Sornnuwat Y., Lee C.-Y.,
858 Yoshimura T. and Tsunoda K.: Resistance of polyamide and polyethylene cable sheathings to
859 termites in Australia, Thailand, USA, Malaysia and Japan: a comparison of four field
860 assessment methods, *International Biodeterioration & Biodegradation*, 66, 53-62,
861 doi:10.1016/j.ibiod.2011.11.001, 2012.
- 862 Li J., Zhang H., Zhang K., Yang R., Li R. and Li Y.: Characterization, source, and retention of
863 microplastic in sandy beaches and mangrove wetlands of the Qinzhou Bay, China, *Marine*
864 *Pollution Bulletin*, 136, 401-406, doi:10.1016/j.marpolbul.2018.09.025, 2018a.
- 865 Li L., Zhou Q., Yin N., Tu C. and Luo Y.: Uptake and accumulation of microplastics in an
866 edible plant, *Chinese Science Bulletin*, 64, 928-934, doi:10.1360/N972018-00845, 2019.
- 867 Li X., Chen L., Mei Q., Dong B., Dai X., Ding G. and Zeng E. Y.: Microplastics in sewage
868 sludge from the wastewater treatment plants in China, *Water Research*, 142, 75-85,
869 doi:10.1016/j.watres.2018.05.034, 2018b.
- 870 Lobelle D. and Cunliffe M.: Early microbial biofilm formation on marine plastic debris, *Marine*
871 *Pollution Bulletin*, 62, 197-200, doi:10.1016/j.marpolbul.2010.10.013, 2011.
- 872 Lu L., Wan Z., Luo T., Fu Z. and Jin Y.: Polystyrene microplastics induce gut microbiota
873 dysbiosis and hepatic lipid metabolism disorder in mice, *Science of the Total Environment*,
874 631, 449-458, doi:10.1016/j.scitotenv.2018.03.051, 2018.
- 875 Maaß S., Daphi D., Lehmann A. and Rillig M. C.: Transport of microplastics by two
876 collembolan species, *Environmental Pollution*, 225, 456-459,
877 doi:10.1016/j.envpol.2017.03.009, 2017.
- 878 Mahon A. M., O'Connell B., Healy M. G., O'Connor I., Officer R., Nash R. and Morrison L.:
879 Microplastics in sewage sludge: effects of treatment, *Environmental Science & Technology*,
880 51, 810-818, doi:10.1021/acs.est.6b04048, 2016.
- 881 Monge-Nájera J.: Ecological biogeography in the Phylum Onychophora, *Biogeographica*, 70,
882 111-123, doi:N/A, 1994.



- 883 Naji A., Esmaili Z., Mason S. A. and Vethaak A. D.: The occurrence of microplastic
884 contamination in littoral sediments of the Persian Gulf, Iran, *Environmental Science and*
885 *Pollution Research*, 24, 20459-20468, doi:10.1007/s11356-017-9587-z, 2017.
- 886 Nika L., Gibson T., Konkus R. and Karp X.: Fluorescent beads are a versatile tool for staging
887 *Caenorhabditis elegans* in different life histories, *G3: Genes, Genomes, Genetics*, 6, 1923-
888 1933, doi:10.1534/g3.116.030163, 2016.
- 889 Nizzetto L., Futter M. and Langaas S.: Are agricultural soils dumps for microplastics of urban
890 origin?, *Environmental Science & Technology*, doi:10.1021/acs.est.6b04140, 2016.
- 891 Nor N. H. M. and Obbard J. P.: Microplastics in Singapore's coastal mangrove ecosystems,
892 *Marine Pollution Bulletin*, 79, 278-283, doi:10.1016/j.marpolbul.2013.11.025, 2014.
- 893 Oliveira M., Ameixa O. M. and Soares A. M.: Are ecosystem services provided by insects"
894 bugged" by micro (nano) plastics?, *TrAC Trends in Analytical Chemistry*,
895 doi:10.1016/j.trac.2019.02.018, 2019.
- 896 Parkyn J. and Newell D. A.: Australian land snails: a review of ecological research and
897 conservation approaches, *Molluscan Research*, 33, 116-129,
898 doi:10.1080/13235818.2013.782793, 2013.
- 899 Pass G., Szucsich N. U. and others: 100 years of research on the Protura: many secrets still
900 retained, *Soil Organisms*, 83, 309-334, doi:N/A, 2011.
- 901 Pelosi C., Bertrand M., Thénard J. and Mougin C.: Earthworms in a 15 years agricultural trial,
902 *Applied Soil Ecology*, 88, 1-8, doi:10.1016/j.apsoil.2014.12.004, 2015.
- 903 Peng B.-Y., Su Y., Chen Z., Chen J., Zhou X., Benbow M. E., Criddle C., Wu W.-M. and Zhang
904 Y.: Biodegradation of Polystyrene by Dark (*Tenebrio obscurus*) and Yellow (*Tenebrio molitor*)
905 Mealworms (Coleoptera: Tenebrionidae), *Environmental Science & Technology*,
906 doi:10.1021/acs.est.8b06963, 2019.
- 907 Petersen H.: A review of collembolan ecology in ecosystem context, *Acta Zoologica Fennica*,
908 195, 111-118, doi:N/A, 1994.
- 909 Phillips H. R., Guerra C. A., Bartz M. L., Briones M. J., Brown G., Ferlian O., Gongalsk K.,
910 Krebs J., Orgiazzi A., Schwarz B. and others: Global distribution of earthworm diversity,
911 bioRxiv, 587394, doi:10.1126/science.aax4851, 2019.
- 912 Prendergast-Miller M. T., Katsiamides A., Abbass M., Sturzenbaum S. R., Thorpe K. L. and
913 Hodson M. E.: Polyester-derived microfibre impacts on the soil-dwelling earthworm *Lumbricus*
914 *terrestris*, *Environmental Pollution*, 251, 453-459, doi:10.1016/j.envpol.2019.05.037, 2019.
- 915 Pulleman M., Creamer R., Hamer U., Helder J., Pelosi C., Peres G. and Rutgers M.: Soil
916 biodiversity, biological indicators and soil ecosystem services—an overview of European



- 917 approaches, *Current Opinion in Environmental Sustainability*, 4, 529-538,
918 doi:10.1016/j.cosust.2012.10.009, 2012.
- 919 Qu M., Kong Y., Yuan Y. and Wang D.: Neuronal damage induced by nanopolystyrene
920 particles in nematode *Caenorhabditis elegans*, *Environmental Science: Nano*,
921 doi:10.1039/C9EN00473D, 2019a.
- 922 Qu M., Nida A., Kong Y., Du H., Xiao G. and Wang D.: Nanopolystyrene at predicted
923 environmental concentration enhances microcystin-LR toxicity by inducing intestinal damage
924 in *Caenorhabditis elegans*, *Ecotoxicology and Environmental Safety*, 183, 109568,
925 doi:10.1016/j.ecoenv.2019.109568, 2019b.
- 926 Qu M., Qiu Y., Kong Y. and Wang D.: Amino modification enhances reproductive toxicity of
927 nanopolystyrene on gonad development and reproductive capacity in nematode
928 *Caenorhabditis elegans*, *Environmental Pollution*, 254, 112978,
929 doi:10.1016/j.envpol.2019.112978, 2019c.
- 930 Reddy M. S., Basha S., Adimurthy S. and Ramachandraiah G.: Description of the small
931 plastics fragments in marine sediments along the Alang-Sosiya ship-breaking yard, India,
932 *Estuarine, Coastal and Shelf Science*, 68, 656-660, doi:10.1016/j.ecss.2006.03.018, 2006.
- 933 Rezaei M., Riksen M. J., Sirjani E., Sameni A. and Geissen V.: Wind erosion as a driver for
934 transport of light density microplastics, *Science of The Total Environment*, 669, 273-281,
935 doi:10.1016/j.scitotenv.2019.02.382, 2019.
- 936 Rillig M. C.: Microplastic in terrestrial ecosystems and the soil?, *Environmental Science &*
937 *Technology*, 6453-6454, doi:10.1021/es302011r, 2012.
- 938 Rillig M. C. and Bonkowski M.: Microplastic and soil protists: a call for research,
939 *Environmental Pollution*, 241, 1128-1131, doi:10.1016/j.envpol.2018.04.147, 2018.
- 940 Rillig M. C., Lehmann A., de Souza Machado A. A. and Yang G.: Microplastic effects on
941 plants, *New Phytologist*, doi:10.1111/nph.15794, 2019.
- 942 Rillig M. C., Ziersch L. and Hempel S.: Microplastic transport in soil by earthworms, *Scientific*
943 *Reports*, 7, 1362, doi:10.1038/s41598-017-01594-7, 2017.
- 944 Robins R. and Robins A.: The antics of ants: ants as agents of bioturbation in a midden
945 deposit in south-east Queensland, *Environmental Archaeology*, 16, 151-161,
946 doi:10.1179/174963111X13110803261010, 2011.
- 947 Rodríguez-Seijo A., da Costa J. P., Rocha-Santos T., Duarte A. C. and Pereira R.: Oxidative
948 stress, energy metabolism and molecular responses of earthworms (*Eisenia fetida*) exposed
949 to low-density polyethylene microplastics, *Environmental Science and Pollution Research*, 25,
950 33599-33610, doi:10.1007/s11356-018-3317-z, 2018.



- 951 Rodríguez-Seijo A., Lourenço J., Rocha-Santos T., Da Costa J., Duarte A., Vala H. and
952 Pereira R.: Histopathological and molecular effects of microplastics in *Eisenia andrei* Bouché,
953 *Environmental Pollution*, 220, 495-503, doi:10.1016/j.envpol.2016.09.092, 2017.
- 954 Rodríguez-Seijo A., Santos B., da Silva E. F., Cachada A. and Pereira R.: Low-density
955 polyethylene microplastics as a source and carriers of agrochemicals to soil and earthworms,
956 *Environmental Chemistry*, 16, 8-17, doi:10.1071/EN18162, 2019.
- 957 Sánchez-Bayo F. and Wyckhuys K. A.: Worldwide decline of the entomofauna: A review of its
958 drivers, *Biological Conservation*, 232, 8-27, doi:10.1016/j.biocon.2019.01.020, 2019.
- 959 Scharf A., Piechulek A. and von Mikecz A.: Effect of nanoparticles on the biochemical and
960 behavioral aging phenotype of the nematode *Caenorhabditis elegans*, *Acs Nano*, 7, 10695-
961 10703, doi:10.1021/nn403443r, 2013.
- 962 e Silva P. P. G., Nobre C. R., Resaffe P., Pereira C. D. S. and Gusmão F.: Leachate from
963 microplastics impairs larval development in brown mussels, *Water Research*, 106, 364-370,
964 doi:10.1016/j.watres.2016.10.016, 2016.
- 965 Song Y., Cao C., Qiu R., Hu J., Liu M., Lu S., Shi H., Raley-Susman K. M. and He D.: Uptake
966 and adverse effects of polyethylene terephthalate microplastics fibers on terrestrial snails
967 (*Achatina fulica*) after soil exposure, *Environmental Pollution*, 250, 447-455,
968 doi:10.1016/j.envpol.2019.04.066, 2019.
- 969 Stamatiadis S. and Dindal D.: Coprophilous mite communities as affected by concentration of
970 plastic and glass particles, *Experimental & Applied Acarology*, 8, 1-12,
971 doi:10.1007/BF01193377, 1990.
- 972 Stock V., Böhmert L., Lisicki E., Block R., Cara-Carmona J., Pack L. K., Selb R., Lichtenstein
973 D., Voss L., Henderson C. J. and others: Uptake and effects of orally ingested polystyrene
974 microplastic particles in vitro and in vivo, *Archives of Toxicology*, 1-17, doi:10.1007/s00204-
975 019-02478-7, 2019.
- 976 Sun Y., Xu W., Gu Q., Chen Y., Zhou Q., Zhang L., Gu L., Huang Y., Lyu K. and Yang Z.:
977 Small-Sized Microplastics Negatively Affect Rotifers: Changes in the Key Life-History Traits
978 and Rotifer--Phaeocystis Population Dynamics, *Environmental Science & Technology*, 53,
979 9241-9251, doi:10.1021/acs.est.9b02893, 2019.
- 980 Thompson R. C., Swan S. H., Moore C. J. and Vom Saal F. S.: Our plastic age,
981 doi:10.1098/rstb.2009.0054, 2009.
- 982 Tsunoda K., Rosenblat G. and Dohi K.: Laboratory evaluation of the resistance of plastics to
983 the subterranean termite *Coptotermes formosanus* (Blattodea: Rhinotermitidae), *International*
984 *Biodeterioration & Biodegradation*, 64, 232-237, doi:10.1016/j.ibiod.2009.12.008, 2010.



- 985 Vadstein O., Øie G. and Olsen Y.: Particle size dependent feeding by the rotifer *Brachionus*
986 *plicatilis*, *Hydrobiologia*, 255, 261-267, doi:10.1007/BF00025847, 1993.
- 987 Van Cauwenberghe L., Claessens M., Vandegehuchte M. B. and Janssen C. R.: Microplastics
988 are taken up by mussels (*Mytilus edulis*) and lugworms (*Arenicola marina*) living in natural
989 habitats, *Environmental Pollution*, 199, 10-17, doi:10.1016/j.envpol.2015.01.008, 2015.
- 990 Vianello A., Boldrin A., Guerriero P., Moschino V., Rella R., Sturaro A. and Da Ros L.:
991 Microplastic particles in sediments of Lagoon of Venice, Italy: First observations on
992 occurrence, spatial patterns and identification, *Estuarine, Coastal and Shelf Science*, 130, 54-
993 61, doi:10.1016/j.ecss.2013.03.022, 2013.
- 994 Vogel G., Thilo L., Schwarz H. and Steinhart R.: Mechanism of phagocytosis in *Dictyostelium*
995 *discoideum*: phagocytosis is mediated by different recognition sites as disclosed by mutants
996 with altered phagocytotic properties, *The Journal of Cell Biology*, 86, 456-465,
997 doi:10.1083/jcb.86.2.456, 1980.
- 998 Wang H.-T., Ding J., Xiong C., Zhu D., Li G., Jia X.-Y., Zhu Y.-G. and Xue X.-M.: Exposure to
999 microplastics lowers arsenic accumulation and alters gut bacterial communities of earthworm
1000 *Metaphire californica*, *Environmental Pollution*, 251, 110-116,
1001 doi:10.1016/j.envpol.2019.04.054, 2019b.
- 1002 Wang J., Coffin S., Sun C., Schlenk D. and Gan J.: Negligible effects of microplastics on
1003 animal fitness and HOC bioaccumulation in earthworm *Eisenia fetida* in soil, *Environmental*
1004 *Pollution*, 249, 776-784, doi:10.1016/j.envpol.2019.03.102, 2019c.
- 1005 Wang J., Liu X., Li Y., Powell T., Wang X., Wang G. and Zhang P.: Microplastics as
1006 contaminants in the soil environment: A mini-review, *Science of The Total Environment*,
1007 doi:10.1016/j.scitotenv.2019.07.209, 2019a.
- 1008 Warburg M. R.: Isopods and their terrestrial environment, In: *Advances in ecological research*,
1009 Elsevier, doi:10.1016/S0065-2504(08)60246-9, 1987.
- 1010 Weisman R. A. and Korn E. D.: Phagocytosis of latex beads by *Acanthamoeba*. I. Biochemical
1011 properties, *Biochemistry*, 6, 485-497, doi:10.1021/bi00854a017, 1967.
- 1012 Weithmann N., Möller J. N., Löder M. G., Piehl S., Laforsch C. and Freitag R.: Organic
1013 fertilizer as a vehicle for the entry of microplastic into the environment, *Science Advances*, 4,
1014 eaap8060, doi:10.1126/sciadv.aap8060, 2018.
- 1015 Westheide W. and Rieger R.: *Spezielle Zoologie. 1. Einzeller und Wirbellose Tiere*, Gustav
1016 Fischer, 1996.



- 1017 Wood C. T. and Zimmer M.: Can terrestrial isopods (Isopoda: Oniscidea) make use of
1018 biodegradable plastics?, *Applied Soil Ecology*, 77, 72-79, doi:10.1016/j.apsoil.2014.01.009,
1019 2014.
- 1020 Wright S. L., Rowe D., Thompson R. C. and Galloway T. S.: Microplastic ingestion decreases
1021 energy reserves in marine worms, *Current Biology*, 23, R1031-R1033,
1022 doi:10.1016/j.cub.2013.10.068, 2013.
- 1023 Yang S.-S., Brandon A. M., Flanagan J. C. A., Yang J., Ning D., Cai S.-Y., Fan H.-Q., Wang
1024 Z.-Y., Ren J., Benbow E. and others: Biodegradation of polystyrene wastes in yellow
1025 mealworms (larvae of *Tenebrio molitor* Linnaeus): Factors affecting biodegradation rates and
1026 the ability of polystyrene-fed larvae to complete their life cycle, *Chemosphere*, 191, 979-989,
1027 doi:10.1016/j.chemosphere.2017.10.117, 2018.
- 1028 Yang X., Lwanga E. H., Bemani A., Gertsen H., Salanki T., Guo X., Fu H., Xue S., Ritsema C.
1029 and Geissen V.: Biogenic transport of glyphosate in the presence of LDPE microplastics: A
1030 mesocosm experiment, *Environmental Pollution*, 245, 829-835,
1031 doi:10.1016/j.envpol.2018.11.044, 2019a.
- 1032 Yang Y., Yang J., Wu W.-M., Zhao J., Song Y., Gao L., Yang R. and Jiang L.: Biodegradation
1033 and mineralization of polystyrene by plastic-eating mealworms: Part 1. Chemical and physical
1034 characterization and isotopic tests, *Environmental Science & Technology*, 49, 12080-12086,
1035 doi:10.1021/acs.est.5b02661, 2015.
- 1036 Yang Y.-F., Chen C.-Y., Lu T.-H. and Liao C.-M.: Toxicity-based toxicokinetic/toxicodynamic
1037 assessment for bioaccumulation of polystyrene microplastics in mice, *Journal of Hazardous
1038 Materials*, 366, 703-713, doi:10.1016/j.jhazmat.2018.12.048, 2019b.
- 1039 Yeates G. W.: Nematodes in ecological webs, *e LS*, doi:10.1002/9780470015902.a0021913,
1040 2001.
- 1041 Yu M., Van Der Ploeg M., Lwanga E. H., Yang X., Zhang S., Ma X., Ritsema C. J. and
1042 Geissen V.: Leaching of microplastics by preferential flow in earthworm (*Lumbricus terrestris*)
1043 burrows, *Environmental Chemistry*, 16, 31-40, doi:10.1071/EN18161, 2019.
- 1044 Zhang G. and Liu Y.: The distribution of microplastics in soil aggregate fractions in
1045 southwestern China, *Science of the Total Environment*, 642, 12-20,
1046 doi:10.1016/j.scitotenv.2018.06.004, 2018.
- 1047 Zhang L., Sintim H. Y., Bary A. I., Hayes D. G., Wadsworth L. C., Anunciado M. B. and Flury
1048 M.: Interaction of *Lumbricus terrestris* with macroscopic polyethylene and biodegradable
1049 plastic mulch, *Science of The Total Environment*, 635, 1600-1608,
1050 doi:10.1016/j.scitotenv.2018.04.054, 2018.



- 1051 Zhang Y., Gao T., Kang S. and Sillanpää M.: Importance of atmospheric transport for
1052 microplastics deposited in remote areas, *Environmental Pollution*, 254, 112953,
1053 doi:10.1016/j.envpol.2019.07.121, 2019.
- 1054 Zhang Z.-Q.: Animal biodiversity: an update of classification and diversity in 2013, *Zootaxa*,
1055 3703, 5-11, doi:10.11646/zootaxa.3703.1.3, 2013.
- 1056 Zhao L., Qu M., Wong G. and Wang D.: Transgenerational toxicity of nanopolystyrene
1057 particles in the range of $\mu\text{g L}^{-1}$ in the nematode *Caenorhabditis elegans*, *Environmental*
1058 *Science: Nano*, 4, 2356-2366, doi:10.1039/C7EN00707H, 2017.
- 1059 Zhu B.-K., Fang Y.-M., Zhu D., Christie P., Ke X. and Zhu Y.-G.: Exposure to nanoplastics
1060 disturbs the gut microbiome in the soil oligochaete *Enchytraeus crypticus*, *Environmental*
1061 *Pollution*, 239, 408-415, doi:10.1016/j.envpol.2018.04.017, 2018b.
- 1062 Zhu D., Bi Q.-F., Xiang Q., Chen Q.-L., Christie P., Ke X., Wu L.-H. and Zhu Y.-G.: Trophic
1063 predator-prey relationships promote transport of microplastics compared with the single
1064 *Hypoaspis aculeifer* and *Folsomia candida*, *Environmental Pollution*, 235, 150-154,
1065 doi:10.1016/j.envpol.2017.12.058, 2018a.