



Short Communication: Quantifying and Correcting for Pre-Assay CO₂ Loss in Short-Term Carbon Mineralization Assays

Matthew A. Belanger^{1,2}, Carmella Vizza^{1,2}, G. Philip Robertson^{1,3}, and Sarah S. Roley²

¹W.K. Kellogg Biological Station, Michigan State University, Hickory Corners, MI 49060, USA

5 ²School of the Environment, Washington State University, Richland, WA 99354, USA

³Department of Plant, Soil, and Microbial Sciences, Michigan State University, East Lansing, MI 48824, USA

Correspondence to: Carmella Vizza (carmella.vizza@wsu.edu)

10

Abstract. The active fraction of soil organic carbon (SOC) is an important component of soil health and often is quickly assessed as the amount of CO₂ released by re-wetting dried soils in short-term (24–72 h) assays. However, soils can lose carbon (C) as they dry and if soil samples vary in moisture content at sampling, differential C loss during the pre-assay dry-down period may complicate interpretations of C availability. We examined pre-assay CO₂ loss and its influence on apparent
15 C availability in the same soil at initial moisture contents of 30, 50, and 70% water-filled pore space (WFPS). We found that 50 and 70% WFPS treatments lost more C during drying than those in the 30% WFPS treatment, which led to a 26–32% underestimate of C availability in wetter soil. We developed a soil-specific correction factor to account for these initial soil moisture effects. Future C mineralization studies may benefit from similar corrections.

1 Introduction

20

The pulse of CO₂ following the re-wetting of dried soils (Robertson et al., 1999; Franzluebbers et al., 2000) has been widely used to indicate soil C availability because of its association with soil microbial biomass C and the active fraction of SOC. This method is derived from the “Birch Effect,” whereby re-wetted dry soils release a pulse of CO₂ resulting from increased microbial activity (Birch, 1958). Drought stress drives microbial communities to dormancy or death (Borken and Matzner, 2009), and following the reintroduction of moisture, microbes burst or release solutes to avoid bursting (Schimel et al., 2007),
25 which stimulates C mineralization (Kim et al., 2012).

Although the short-term pulse of CO₂ following the re-wetting of dry soils is a widely used method for assessing soil C availability (e.g., Culman et al., 2013; Ladoni et al., 2016; Morrow et al., 2016; Sprunger and Robertson, 2018), we are unaware of efforts to quantify the potential bias introduced by assaying soils of different moisture contents at the time of sampling.
30 Soils that differ in moisture will dry down at different rates, potentially losing different amounts of available C prior to the start of the assay. If sufficiently large, differential pre-assay losses could complicate comparisons of C availability across field treatments or landscape catenas.



Here we investigate the influence of different initial soil moisture levels on pre-assay CO₂ release during drying for an Alfisol soil in the upper Midwest, USA. We test the hypothesis that moister soil will have higher pre-assay CO₂ loss because a longer
35 dry-down period results in more time for such losses to occur.

2 Materials and methods

2.1 Site description

We collected soil from the Ap horizon (0–20 cm) of an arable grass field at the W.K. Kellogg Biological Station (KBS) in Hickory Corners, MI (42°41'02" N, –85°37'34" W). KBS soils are mixed, mesic Typic Hapludalfs of co-mingled Kalamazoo
40 and Oshtemo series (Crum and Collins, 1995) developed on glacial outwash with intermixed loess (Luehmann et al., 2016). Soil collected in September 2019 for this experiment was from the Kalamazoo series, which describe well-drained fine-loams (43% sand, 38% silt, 19% clay) with ~2% total C (Grandy and Robertson, 2006) and a pH of 7.2 (Robertson et al., 1993). Average annual rainfall at KBS is 1005 mm, average annual snowfall is 1300 mm, and mean annual temperature is 10.1°C (Robertson and Hamilton, 2015). The site was in various corn-soybean-wheat rotations for the past 40 years and before that,
45 corn-soybean-small grain rotations for at least 60 years.

2.2 Experimental design

To examine the influence of initial soil moisture on the pre-assay loss of CO₂ during dry-down, we pre-wet recently collected soil to three different initial water-filled pore space (WFPS) levels: 30, 50, and 70%. Then we measured gravimetric soil moisture (GSM) and CO₂ loss while soil was air-drying, after which we re-wet them and measured the 24-hr CO₂ pulse by
50 standard methods (Robertson et al., 1999; Franzluebbbers et al., 2000).

2.3 Laboratory analyses

After collection, soil was sieved through a 4-mm mesh and mixed. We measured gravimetric soil moisture (GSM) and calculated the target volumetric water content (VWC, g H₂O cm⁻³ soil) for each treatment following Eq. 1 (Elliott et al., 1999):
$$\text{VWC} = \text{WFPS}/100 * (1 - \text{SBD}/2.65) \quad (1)$$

55 where soil bulk density (SBD) was 1.5 g soil cm⁻³, a previously assessed value from KBS soils (Robertson, 2016). Then we divided VWC by SBD to obtain a target GSM and thereby determined the amount of water to add to the field-moist soil (11% WFPS; GSM = 0.032 g H₂O g⁻¹ dry soil). We then weighed 40 g of soil into each of 75 specimen cups. Each cup was randomly assigned to an initial WFPS treatment (30, 50, or 70%), for a total of 25 replicates per treatment. We added sufficient deionized water to each cup to achieve the target initial WFPS and stirred to evenly distribute water. After soil was wet and stirred in the
60 specimen cups, the contents of each cup were transferred to a labeled paper bag. The soil was spread evenly across the bottom of the bag, and the top portion of the bag was cut off to increase air flow. Afterwards, the soil was immediately weighed and set on a laboratory bench to air-dry.



Immediately after wetting, as well as 1, 3, and 8 days later, we assessed GSM and CO₂ loss rates for five replicates per initial
65 WFPS treatment. GSM, which was determined after drying the soil at 105°C for 24 hrs, stabilized at 1.5% in the air-dried soil
(Fig. 1a), but did not reach zero even when soil was completely air-dry. Because soil in all initial WFPS treatments were air-
dry by day 3, with CO₂ loss rates close to zero, we terminated GSM and CO₂ measurements after day 8.

CO₂ loss rates at each sampling interval were measured by placing 10 g of soil into a 235 mL mason jar equipped with a gas-
70 sampling septum. Then we sampled 5 mL of headspace from each jar at 4 intervals (0, 0.5, 1, and 2 hr), injected it into an
evacuated 3 mL exetainer (Labco Limited, Lampeter, Wales, United Kingdom), and replaced the jar headspace with laboratory
air. CO₂ samples were analyzed within 24 hrs using a LI-820 CO₂ Gas Analyzer (LI-COR Biosciences, Lincoln, NE, USA).

On day 15 we re-wet the remaining five replicates of air-dried soil from each initial WFPS treatment to 50% WFPS
75 (Franzuebbers et al., 2000). We then assessed subsequent 24-hr CO₂ pulses by sampling headspaces at 0, 2, 4, 8, and 24 hrs.

2.4 Statistical analyses and correction factor

CO₂ pulses were calculated as the positive slope of the linear regression of CO₂ concentrations through time after accounting
for headspace dilution, and then converted to a standardized rate using the ideal gas law. In 17 of 75 cases, we omitted one of
the four data points within a jar, which were clear visual outliers. In two cases, we rejected jars with leaks. CO₂ loss rates
80 during the dry-down period were analyzed with a two-way analysis of covariance (ANCOVA), where initial WFPS treatment
and days elapsed since wetting were factors and GSM at the time of sampling was a covariate. Additionally, a one-way analysis
of variance (ANOVA) was used to determine whether initial WFPS treatment had an effect on the 24-hr CO₂ pulses upon re-
wetting the air-dried soil.

85 We also calculated a correction factor to account for pre-assay CO₂ loss prior to the 24-hr CO₂ pulse assay. To calculate the
total amount of CO₂ loss during dry-down for each initial WFPS treatment, we calculated a best-fit exponential decay curve
($Y = \alpha^{\beta X} + \theta$), where Y = daily CO₂-C loss and X = length of dry-down period, until soil was air-dry (i.e., immediately after
wetting through day 3). Total C loss was equivalent to the area under the curve.

90 Because we used sacrificial sampling, we could not calculate standard deviation or standard error in the usual way. Instead,
we used a bootstrapping approach in which we computed predicted values for CO₂ losses (\hat{Y}_i) and residuals ($e_i = Y_i - \hat{Y}_i$). All
zeroes for CO₂ losses were set to 1 for the sake of fitting the regression because an exponential decay curve can approach but
never attain 0 and because 1 was lower than any value we observed. Then we created a bootstrap sampling of residuals specific
to each dry-down interval (0, 1, or 3 days), sampled randomly from each interval with replacement, and added randomly
95 sampled residuals to predicted values ($Y_i^* = \hat{Y}_i + e_i^*$) for each dry-down interval (after Hesterberg, 2015). Residuals were




bootstrapped 10,000 times to derive multiple estimates of coefficients for the exponential decay curve (α , β , and θ). We also integrated under the curve 10,000 times to get an error estimate (i.e., coefficient of variation) associated with the total amount of pre-assay CO₂ loss during dry-down.

100 Then we divided the total CO₂ loss by three days to obtain the daily rate used to calculate a correction factor following Eq. 2:
CF = (daily CO₂ loss during dry-down / 24-hr CO₂ pulse after re-wetting) + 1 (2)

The correction factor for each treatment was then multiplied by each replicate's 24-hr CO₂ pulse following re-wetting. Finally, we verified that the correction factors worked by conducting a one-way ANOVA to determine whether initial WFPS treatment still had an effect on the corrected pulses. For all analyses, we confirmed that assumptions of normality and homogeneity of
105 variance were not violated.

3 Results

Soil in the 50 and 70% WFPS treatments took longer to dry than did soil in the 30% WFPS treatment (Fig 1a). A day after wetting, soil from the 30% WFPS treatment was completely air-dry, but soil had lost only 79% and 68% of its initial moisture in the 50 and 70% WFPS treatments, respectively. All soil was air-dry by three days after wetting. Pre-assay CO₂ losses



110 **Fig 1. (a) Gravimetric soil moisture (GSM) during air-drying and (b) daily CO₂ losses from each initial water-filled pore space (WFPS) treatment during the dry-down period. Error bars represent standard errors of the mean.**

mirrored soil moisture loss, reaching zero for all WFPS treatments by day 3 (Fig 1b). Both GSM at the time of sampling and day had effects on pre-assay CO₂ loss rates ($P < 0.0001$), but initial WFPS treatment did not ($P = 0.28$) probably because GSM captures more variation in soil moisture as the soil dries than WFPS treatment. However, there was an interaction between
115 treatment and day ($P = 0.0005$). Soil of even the lowest initial WFPS treatment lost C as CO₂ over three days of drying (26 µg



CO₂-C g⁻¹ soil for 30% WFPS), but losses were disproportionately higher from wetter soil (62 and 71 μg CO₂-C g⁻¹ soil for 50 and 70% WFPS, respectively).

120 Initial soil moisture (i.e., WFPS treatment) had a significant effect on 24-hr CO₂ pulses after re-wetting air-dried soil ($P = 0.007$; Fig 2). While final CO₂ pulses were lower for the 50 and 70% WFPS treatments relative to 30% WFPS (Fig. 2), the 50




Fig 2. 24-hr CO₂ pulses after the re-wetting of air-dried soil for each initial water-filled pore space (WFPS) treatment. Error bars represent standard error of the mean.

and 70% WFPS treatments also tended to have greater pre-assay CO₂ losses during three days of dry-down, which represented 125 77 and 95% of their 24-hr CO₂ pulses, respectively. After accounting for these losses with correction factors, the 24-hr CO₂ pulses were similar across initial WFPS treatments ($P = 0.28$; Fig. 3).




Fig 3. Daily CO₂ production rates for each initial water-filled pore space (WFPS) treatment. Lined bars represent the average daily rate of pre-assay CO₂ loss during a 3-day dry-down period and solid bars represent the 24-hr CO₂ pulses after re-wetting the air-dried soil. Together both bars represent the 24-hr CO₂ pulse corrected for pre-assay losses of CO₂ during dry-down. Error bars represent standard deviation as described in the “Statistical analyses” section.

4 Discussion

Initial soil moisture levels played a significant role in our ability to accurately characterize soil C availability (Fig 2) via the conventional 24-hr CO₂ pulse assay (Robertson et al., 1999; Franzluebbers et al., 2000). Wetter soil lost more C during dry-down, presumably because soil microbes remained active for a longer period of time. These losses decreased the short-term CO₂ pulses and therefore the final estimates of soil C availability.

Without knowledge of these losses, one might erroneously conclude that soil from the 30% WFPS treatment had about 35% higher soil C availability than the others (Fig 2), but this trend is instead due to higher pre-assay CO₂ losses during the dry-down period for wetter soil (Fig 3). It is striking that even short drying intervals (i.e., 1 versus 3 days) can affect soil C availability as deduced from the 24-hr CO₂ pulse after re-wetting air-dried soil. However, we were able to account for the pre-assay CO₂ losses for our soil with a correction factor that made C availability approximately equivalent across all initial WFPS treatments.

These trends suggest that efforts to characterize C availability via short-term CO₂ pulses following the re-wetting of dry soil should exercise caution if comparisons involve soils with a range of initial soil moistures. This includes soils compared across seasons; across drought, precipitation, or irrigation gradients; across landscape catenas; across crop, grazing, or forest



management practices; and as well in cross-site comparisons and meta-analyses that include soils collected at different initial soil moistures.

150

A correction factor that accounts for pre-assay CO₂ losses may help to normalize such comparisons. In our soil, pre-assay CO₂ losses led to a C mineralization bias as high as 32%, for which we could confidently correct by applying a correction factor based on measured rates of pre-assay CO₂ loss (Eq. 2). Other soils with moisture contents sufficient to oxidize available C during dry-down will require different correction factors. A soil-specific correction factor can be calculated by measuring CO₂ loss during dry-down on a subset of samples, as we described above (Eq. 2).

155

An alternate solution is to minimize the dry-down period such that little available C is lost prior to the assay. Strategies to minimize pre-assay CO₂ loss might include exposing soils to temperatures high enough to speed evaporation, but low enough to avoid sterilization (Jager, 1968) or otherwise artificially disrupt the microbial community (Evans and Wallenstein, 2012).

160

This could be performed in a closed vented chamber such as a soil incubator. Alternatively, faster and more even drying might be achieved with a steady flow of air (i.e., a fan or vented system) over exposed soil samples. Even with faster drying, however, a correction factor may be needed.

Overall, our results demonstrate that using the 24-hr CO₂ pulse following the re-wetting of a dried soil to evaluate soil C availability can be misleading for soils with different moisture contents at time of sampling. For such soils a correction factor based on pre-assay CO₂ losses can be applied with confidence.

165

Data availability

Data will be made publicly available upon publication at Dryad; a pre-publication version is available at <https://datadryad.org/stash/share/o4a-ESVy8wxpnk-kbrZicM9eMuMrX2F5oJpg039DWGQ>.

170 Author contributions

CV and GPR designed this study, MAB and CV performed the laboratory assays, MAB analyzed the CO₂ samples, CV conducted the statistical analyses, MAB and CV wrote the paper with contributions from SSR and GPR.

Competing interests

The authors declare that they have no conflict of interest.



175 Acknowledgements

We acknowledge the Roley lab group at WSU and the Robertson lab group at KBS for their helpful comments, especially Sven Bohm, whose questions inspired us to undertake this study. We also thank Stuart Jones for his advice about bootstrapping. Support for this research was provided by the National Science Foundation (NSF) Division of Environmental Biology DEB 1754212, by the NSF Long-term Ecological Research Program (DEB 1832042) at the W. K. Kellogg Biological Station (KBS),
180 by the Great Lakes Bioenergy Research Center, U.S. Department of Energy Office of Science, Office of Biological and Environmental Research (Award DE-SC0018409), and by Michigan State University AgBioResearch.

References

- Birch, H. F.: The effect of soil drying on humus decomposition and nitrogen availability, *Plant Soil*, 10(1), 9–31, doi:10.1007/BF01343734, 1958.
- 185 Borken, W. and Matzner, E.: Reappraisal of drying and wetting effects on C and N mineralization and fluxes in soils, *Global Change Biology*, 15(4), 808–824, doi:10.1111/j.1365-2486.2008.01681.x, 2009.
- Crum, J. R. and Collins, H. P.: Soils of the Kellogg Biological Station (KBS), Kellogg Biological Station Long-Term Research
190 Publication, Zenodo, <https://doi.org/10.5281/zenodo.2581504>, 1995.
- Culman, S. W., Snapp, S. S., Green, J. M. and Gentry, L. E.: Short- and long-term labile soil carbon and nitrogen dynamics reflect management and predict corn agronomic performance, *Agronomy Journal*, 105(2), 493–502, doi:10.2134/agronj2012.0382, 2013.
- 195 Elliott, E. T., Heil, J. W., Kelly, E. F. and Monger, H. C.: Soil structure and other physical properties, in: *Standard Soil Methods for Long-Term Ecological Research*, edited by: Robertson, G. P., Coleman, D., Bledsoe, C., and Sollins, P., Oxford University Press, Inc., New York, NY, USA, 77, 1999.
- 200 Evans, S. E. and Wallenstein, M. D.: Soil microbial community response to drying and rewetting stress: does historical precipitation regime matter?, *Biogeochemistry*, 109(1), 101–116, doi:10.1007/s10533-011-9638-3, 2012.
- Franzluebbers, A. J., Haney, R. L., Honeycutt, C. W., Schomberg, H. H. and Hons, F. M.: Flush of carbon dioxide following rewetting of dried soil relates to active organic pools, *Soil Science Society of America Journal*, 64(2), 613–623,
205 doi:10.2136/sssaj2000.642613x, 2000.
- Grandy, A. S. and Robertson, G. P.: Aggregation and organic matter protection following tillage of a previously uncultivated soil, *Soil Science Society of America Journal*, 70(4), 1398–1406, doi:10.2136/sssaj2005.0313, 2006.
- 210 Hesterberg, T. C.: What teachers should know about the bootstrap: resampling in the undergraduate statistics curriculum, *The American Statistician*, 69(4), 371–386, doi:10.1080/00031305.2015.1089789, 2015.
- Jager, G.: The influence of drying and freezing of soil on its organic matter decomposition, *Stikstof*, 12, 75–88, 1968.
- 215 Kim, D. G., Vargas, R., Bond-Lamberty, B. and Turetsky, M. R.: Effects of soil rewetting and thawing on soil gas fluxes: a review of current literature and suggestions for future research., *Biogeosciences*, 9(7), 2459–2483, doi: 10.5194/bg-9-2459-2012, 2012.



- 220 Ladoni, M., Basir, A., Robertson, P. G. and Kravchenko, A. N.: Scaling-up: cover crops differentially influence soil carbon in agricultural fields with diverse topography, *Agriculture, Ecosystems & Environment*, 225, 93–103, doi:10.1016/j.agee.2016.03.021, 2016.
- 225 Luehmann, M. D., Peter, B. G., Connallon, C. B., Schaetzl, R. J., Smidt, S. J., Liu, W., Kincare, K. A., Walkowiak, T. A., Thorlund, E. and Holler, M. S.: Loamy, two-storied soils on the outwash plains of southwestern lower Michigan: pedoturbation of loess with the underlying sand, *Annals of the American Association of Geographers*, 106(3), 551–572, doi:10.1080/00045608.2015.1115388, 2016.
- 230 Morrow, J. G., Huggins, D. R., Carpenter-Boggs, L. A. and Reganold, J. P.: Evaluating measures to assess soil health in long-term agroecosystem trials, *Soil Science Society of America Journal*, 80(2), 450–462, doi:10.2136/sssaj2015.08.0308, 2016.
- Robertson, G. P.: Soil carbon and nitrogen deep core surveys at the Kellogg Biological Station, Hickory Corners, MI (2001 to 2013) ver. 23, Environmental Data Initiative, <https://doi.org/10.6073/pasta/8c9f759d32699902d88c7b5d607fb2f7>, 2016.
- 235 Robertson, G. P. and Hamilton, S. K.: The ecology of agriculture landscapes: long-term research on the path to sustainability, in: *Long-Term Ecological Research at the Kellogg Biological Station LTER Site*, edited by: Robertson, G. P. and Hamilton, S. K., Oxford University Press, Inc., New York, NY, USA, 1–32, 2015.
- Robertson, G. P., Crum, J. R. and Ellis, B. G.: The spatial variability of soil resources following long-term disturbance, *Oecologia*, 96(4), 451–456, doi:10.1007/BF00320501, 1993.
- 240 Robertson, G. P., Wedin, D. A., Groffman, P. M., Blair, J. M. and Holland, E. A.: Soil carbon and nitrogen availability: nitrogen mineralization, nitrification, and soil respiration, in: *Standard Soil Methods for Long-Term Ecological Research*, edited by: Robertson, G. P., Coleman, D., Bledsoe, C., and Sollins, P., Oxford University Press, Inc., New York, NY, USA, 258–271, 1999.
- 245 Schimel, J., Balsler, T. C. and Wallenstein, M.: Microbial stress-response physiology and its implications for ecosystem function, *Ecology*, 88(6), 1386–1394, doi:10.1890/06-0219, 2007.
- Sprunger, C. D. and Robertson, G. P.: Early accumulation of active fraction soil carbon in newly established cellulosic biofuel systems, *Geoderma*, 318, 42–51, doi:10.1016/j.geoderma.2017.11.040, 2018.