# Point-by-point changes made to "Obtaining more benefits from crop residues as soil amendments by application as chemically heterogeneous mixtures" by Marijke Struijk et al.

L. 13 (RC1 ) We removed "(i.e. mixture 6= sum of the parts)" so that it now reads: "Mixing high C:N ratio with low C:N ratio amendments may result in greater carbon use efficiency and non-additive benefits in soil properties (i.e. mixture ≠ sum of the parts)."

L. 47 (RC1) We removed "(mixture > sum of the parts)" so that it now reads: "Synergistic non-additive mixing effects are frequently observed, i.e. decomposition of the mixture is greater than would be predicted from the rate of decomposition of individual litter types <del>(mixture > sum of the parts)</del>, especially when the litters are chemically heterogeneous (Pérez Harguindeguy et al., 2008; Wardle et al., 1997)"

**L. 64. (RC1)** We specified the "other soil properties" so that is now reads: *"These experiments suggest that non-additivity in decomposition rates and changes to soil C and N dynamics could go hand-in-hand"* 

**L. 92 (RC2)** We incorporate the suggestion to include mention of microbial succession during decomposition by adding the following sentence to the paragraph starting at line 88:

"... Other authors have also suggested the possibility of manipulating the functionality of the soil microbial community with soil amendments, such as Li et al. (2019) who report that eutrophic microbes are stimulated by organic carbon amendments and oligotrophic microbes are stimulated by chemical fertilisers. Studies have also demonstrated that changes in tree litter diversity affect both fungal and bacterial diversity (Otsing et al., 2018; Santonja et al., 2018). **Research on decomposition in forest systems indicates a succession in the community composition of microbial decomposers as the decomposition of residues progresses (Bastian et al., 2009; Purahong et al., 2016), and this succession is different in decomposition of litters of different qualities (Aneja et al., 2006).**"

Additional references were included:

Aneja (2006): doi.org/10.1007%2Fs00248-006-9006-3 Bastian (2009): doi.org/10.1016/j.soilbio.2008.10.024 Puraong et al. (2016): doi.org/10.1111/mec.13739

**L. 118 (RC1 and RC2)** We revised the wording of hypothesis 1 in response to comments from both reviewers. In RC1 there seemed to be some confusion over non-additive effects vs. differences between individual treatments. To clarify we reworded the hypothesis to read:

"In particular, we hypothesised faster decomposition of residue mixtures to result in a higher soil respiration rate in the short term, as well as the release of greater levels of soil available nutrients (N, P, K, Mg) and SOM compared to **what would be expected by combining the effects of** individual residues (hypothesis 1)"

Additionally, details on pH were requested in RC2, so we further revised the hypothesis to read: *"In particular, we hypothesised faster decomposition of residue mixtures to result in a higher soil respiration rate in the short term, as well as the release of greater levels of soil available nutrients (N, P, K, Mg) and SOM, compared to individual residues,* **which**  *leads to greater ammonification of resudie N (Xu et al., 2006), and, in turn, leads to a greater increase in pH* (hypothesis 1)."

The additional reference Xu et al. (2006) has been included in the reference list. This meant we needed to adjust the wording of the pH result in **line 240-45** as follows:

**"In agreement with** our hypothesis, there was a non-additive increase in pH from the mixtures relative to individual amendments (hypothesis 1), although this was not significant (Table 5) and per-treatment results (discussed in next section) show that the pH decreased in all treatments relative to the control (F = 2.238; p = 0.095; one-way ANOVA; Supplement S2)."

**L. 140 (RC1)** To avoid confusion over maturity of compost, we removed the word 'fresh' from this sentence so that it now reads:

Application rates of the different amendments were 20 t  $ha^{-1}$  fresh compost (equivalent to 7 t  $ha^{-1}$  dry matter), 13.3 t  $ha^{-1}$  woodchips (equivalent to 8.7 t  $ha^{-1}$  dry matter) and 10±0.8 t  $ha^{-1}$  straw (equivalent to 9.2±0.8 t  $ha^{-1}$  dry matter).

L. 141 (RC1) The use of ±0.8 has been clarified:

"... and  $10\pm0.8$  t ha<sup>-1</sup> straw (equivalent to  $9.2\pm0.8$  t ha<sup>-1</sup> dry matter;  $\pm$  indicates inclusive range of the straw application rate)."

**L. 144 (RC1)** We removed the word "roughly" because it results from the range of application rates addressed previously and is therefore redundant.

**L. 157 (RC1)** Table 2 has been moved to supplementary information. All table numbers have been adjusted accordingly both in the tables and in the text.

**L. 170 (RC2)** Additional analytes Mn and Fe were added to the residue characterisation table (originally Table 3, but now renumbered to Table 2) in response to RC2. The recovery rates of these analytes were added to the Methods section.

## L.259. See L.118 above

L. 248 (RC1) For clarity we rephrased the paragraph as follows:

"Both compost-residue mixtures resulted in a non-additive increase in lettuce yield, available and potentially mineralisable N, available Mg, SOM, and soil respiration, but not in available K (hypothesis 1) some of which were statistically significant, **as further specified below** (Table 5). Most notably, we observed greater available N and SOM levels in soils to which a mixture of residues was applied, compared to the available N and SOM levels in treatments receiving only individual residue amendments. The strawcompost mixture resulted in a significant (T = 4.022, p = 0.014) non-additive increase in SOM of 13.10%, and while the woodchip-compost mixture did not result in statistically significant non-additivity (T = 0.954, p = 0.205), it did result in a positive non-additive increase in crease in SOM of 6.73%."

**L. 317 (RC2)** As requested, we expanded on our description of the data presented in Table 7, to also include the significant correlation between SOM and available nutrients:

"A number of noteworthy correlations may help explain the data and are summarised in Table 7. There were some significant correlations between the amount of nutrients applied and the amount of available K and Mg in the soils at the end of the experiment, which indicates a positive effect of the residue amendments. The amount of C applied via the residue amendments was not correlated with the amount of SOM. Yield was positively correlated with the sum of available and potentially mineralisable N, available P and Mg, SOM and aggregate stability. **SOM was also positively correlated with available N, P and Mg, and with soil respiration.**"

**L. 350 (RC1)** We clarified why Solvita burst method may not be representative of in-situ respiration by adding the following in line 350, and we corrected 4 mm as the sieve size which should have been 2 mm:

"Moreover, our soil respiration measurements were taken by the Solvita burst method, on soil samples removed from the field and sieved to **2** mm **removing parts of residues and other organic matter greater than 2 mm**, which may not have been a good representation of the respiration from a soil mixed with crop residues at various stages of decomposition."

**L. 355 (RC2)** In response to RC2 we made mention of the fact that the residues introduce microbes that may not already be present in the soil by revising in line 355 as follows:

"As pointed out by Lecerf et al. (2011), niche complementarity effects, in which different groups of decomposing organisms **(already present in the soil, or newly introduced via the residues)** develop a synergistic association in residue breakdown, tend to advance with time, leading to a generally higher number of long-term litter-mixing studies finding non-additive effects."

L. 397 (RC1) In response to RC1 we revised as follows:

"Even in the straw-compost treatment, the SOM level was very close to that of the control treatment, suggesting **net** mineralisation of native SOM **as a result of the residue amendment** was negligible"

L. 482. An acknowledgement was added by one of the authors.

## Supplement

Table S1 was moved from the main text to the supplement in response to RC1, and the rest of the supplement renumbered accordingly.

Table S2 was originally Table 5 and was moved to the supplement in response to RC1. Calculations in S4 were revised and completed.

## Obtaining more benefits from crop residues as soil amendments by application as chemically heterogeneous mixtures

Marijke Struijk<sup>1,2</sup>, Andrew P. Whitmore<sup>2</sup>, Simon R. Mortimer<sup>3</sup> and Tom Sizmur<sup>1</sup>

<sup>1</sup> Department of Geography and Environmental Science, University of Reading, Reading, UK.
 <sup>2</sup> Department of Sustainable Agriculture Sciences, Rothamsted Research, Harpenden, UK.
 <sup>3</sup> School of Agriculture, Policy and Development, University of Reading, Reading, UK.

Correspondence to: Marijke Struijk (m.struijk@pgr.reading.ac.uk, permanent address: mstruijk@gmx.com)

Abstract. Crop residues are valuable soil amendments in terms of the carbon and other nutrients they contain, but incorporation of residues does not always translate into increases in nutrient availability, soil organic matter (SOM), soil structure, and overall soil fertility. Studies have demonstrated accelerated decomposition rates of chemically heterogeneous litter mixtures, compared to the decomposition of individual litters, in forest and grassland systems. Mixing high C:N ratio

- with low C:N ratio amendments may result in greater carbon use efficiency and non-additive benefits in soil properties. We hypothesised that non-additive benefits would accrue from mixtures of low-quality (straw or woodchips) and highquality (vegetable-waste compost) residues applied before lettuce planting in a full-factorial field experiment. Properties
- 15 indicative of soil structure and nutrient cycling were used to assess benefits from residue mixtures, including soil respiration, aggregate stability, bulk density, SOM, available and potentially mineralisable N, available P, K and Mg, and crop yield. Soil organic matter and mineral nitrogen levels were significantly and non-additively greater in the straw-compost mixture compared to individual residues, which mitigated the N immobilisation occurring with straw-only applications. Addition of compost significantly increased soil available N, K and Mg levels. Together, these observations suggest that greater nutrient availability improved the ability of decomposer organisms to degrade straw in the straw-compost mixture.
- We demonstrate that mixtures of crop residues can influence soil properties non-additively. Thus, greater benefits may be achieved by removing, mixing, and re-applying crop residues, than by simply returning them to the soils *in situ*.

#### 1. Introduction

- Intensive agricultural systems, with a monoculture of crops and relying on external inputs of fertilisers and 25 pesticides/herbicides, are criticised for their negative environmental impacts. These include the degradation of soil – particularly degradation of soil organic matter (SOM), biodiversity loss, and over-application of N and P (Malézieux *et al.*, 2009; Tilman *et al.*, 2002). Implementation of multispecies cropping systems (e.g. Malézieux *et al.*, 2009) and increasing functional diversity via trait-based approaches (Garnier and Navas, 2012) are some methods that have been proposed to increase biodiversity and functional complementarity of the variety of species present in arable cropping systems. These
- 30 approaches can lead to more sustainable nutrient cycling, reduced soil erosion, stabilised crop production, and improvements to a system's innate capacity to resist pests, diseases and other environmental disturbances (Gurr *et al.*, 2003). However, some farming systems prevent the cultivation of more than one crop in a field at any one time, and so applying mixtures of crop residues may provide an alternative route to obtaining the benefits of multispecies cropping within monocultural arable cropping systems.
- 35 Crop residues comprise the majority of plant materials harvested worldwide (Medina *et al.*, 2015; Smil, 1999) and are readily available on arable farms. Containing carbon and other nutrients, they present a valuable resource as soil amendments with the potential to increase SOM and nutrient levels, which feed the soil food web (Kumar and Goh, 1999) and may increase soil aggregation and improve soil structure (Cosentino *et al.*, 2006; Martin *et al.*, 1955). Unfortunately, while these changes in soil properties are likely to lead to increased crop yield, decomposition of residue soil amendments

Deleted: (i.e. mixture ≠ sum of the parts)

does not always translate into such benefits and is instead followed by loss from the system, with lower soil N retention and C levels than expected (Catt *et al.*, 1998; Powlson *et al.*, 2011; Thomsen and Christensen, 2006).

Rather than applying a single crop residue, mixtures of crop residues could form a better soil amendment. Complementarity in mixtures of different residues has been previously shown in research on the decomposition rates of mixtures of moss and

- 45 leaf litters in forest ecosystems and grass clippings in grassland ecosystems (Gartner and Cardon, 2004; Hättenschwiler *et al.*, 2005). Synergistic non-additive mixing effects are frequently observed, i.e. decomposition of the mixture is greater than would be predicted from the rate of decomposition of individual litter types, especially when the litters are chemically heterogeneous (Pérez Harguindeguy *et al.*, 2008; Wardle *et al.*, 1997).
- Suggested mechanisms for non-additive decomposition rates of mixtures include physical, chemical and biological processes (Gartner and Cardon, 2004). Frequently cited is the mechanism that N-rich residues are thought to accelerate the decomposition of N-poor residues (Seastedt, 1984) by inter-specific transfer of nutrients in the residue mixture (Berglund *et al.*, 2013; Briones and Ineson, 1996). Additionally, more heterogeneous and improved micro-environmental conditions increase habitat and resource options for decomposer organisms (Hättenschwiler *et al.*, 2005), also known as the improved micro-environmental condition theory (Makkonen *et al.*, 2013).
- 55 However, whether synergistic decomposition rates in mixtures are related to benefits in terms of soil nutrient and carbon management is unclear because studies on the C and N dynamics in decomposing residue mixtures are limited (Redin *et al.*, 2014). It has been shown that increased plant species richness can promote soil C and N stocks via higher plant productivity (Cong *et al.*, 2014) and to increased diversity and functionality of soil microbes (Lange *et al.*, 2015) as well as the whole soil food web (Eisenhauer *et al.*, 2013). Quemada and Cabrera (1995) found non-additivity in the C and N dynamics when
- 60 mixtures of leaves and stems were decomposed compared to individual residues, with the C:N ratio of the residues playing an important role in N mineralisation. Nilsson *et al.* (2008) report synergistic effects on soil available N as well as on plant productivity when mixing *Populus tremula* litter (C:N = 40, known to decompose quickly) with *Empetrum hermaphroditum* (C:N = 77, known to decompose slowly). These experiments suggest that non-additivity in decomposition rates and changes to soil C and N dynamics could go hand-in-hand.
- 65 Increasingly more evidence is emerging that SOM accumulation is primarily derived from the production of microbial residues (Ludwig *et al.*, 2015; Simpson *et al.*, 2007), and this microbially-derived SOM seems to be produced at the early stages of plant-residue decomposition (Cotrufo *et al.*, 2015). Microbial carbon use efficiency (CUE) describes a functional trait of microbes that refers to the fraction of carbon assimilated from organic matter additions to the soil system compared to C losses to the atmosphere via microbial respiration (Allison *et al.*, 2010). Different microbial species exhibit an inherent
- 70 CUE window, so that they can operate at different CUE levels to fulfil their maintenance and growth C requirements depending on environmental factors (Schimel *et al.*, 2007). Organic substrates can feed into different microbial metabolic pathways (e.g. anabolism vs. catabolism) or microbial communities that exhibit different overall inherent CUE levels (e.g. fungi vs. bacteria, or copiotrophs vs. oligotrophs) (Jones *et al.*, 2018). Therefore, an increase in the amount of SOM from microbial activity is not linearly related to CO<sub>2</sub> production, or to the quantity of C applied to the soil, but depends also on the 75 CUE of the decomposer community.

Fertilisation practices typical of intensively managed arable soils stimulate copiotrophic microorganisms (Fierer *et al.*, 2012) with boom-bust population dynamics. These microbial communities tend to exhibit a lower inherent CUE window than slower growing oligotrophic communities (Ho *et al.*, 2017; Roller and Schmidt, 2015). In intensively managed arable soils, the decomposition of soil-applied crop residues can lead to a large portion of residue-derived C being respired as CO<sub>2</sub> rather

80 than turned into SOM (Bailey *et al.*, 2002; Six *et al.*, 2006). Decomposition of high-C:N residues requires microbes with a relatively high CUE, but due to N-limitation they operate towards the lower end of their CUE window (Kallenbach *et al.*, 2019). Low-C:N residues, providing relatively more N, may increase the CUE of individual microbes, but can also shift the composition of the soil microbial community to one that exhibits an inherently lower CUE (Kallenbach *et al.*, 2019). As

Deleted: other soil properties

**Deleted:** (mixture > sum of the parts)

suggested by Kallenbach *et al.* (2019), a mixture of crop residues of different C:N ratios could therefore achieve a more diverse microbial community comprising organisms fulfilling niches of both high and low inherent CUE windows, and may enable all species to operate at their maximum CUE. Other authors have also suggested the possibility of manipulating the functionality of the soil microbial community with soil amendments, such as Li *et al.* (2019) who report that microbes in a eutrophic system are stimulated by organic carbon amendments and oligotrophic microbes are stimulated by chemical fertilisers. Studies have also demonstrated that changes in tree litter diversity affect both fungal and bacterial diversity (Otsing *et al.*, 2018; Santonja *et al.*, 2018). Research on decomposition in forest systems indicates a succession in the community composition of microbial decomposers as the decomposition of residues progresses (Bastian *et al.*, 2009; Purahong *et al.*, 2016), and this succession is different in decomposition of litters of different qualities (Aneja *et al.*, 2006).

- 95 Low-quality plant materials with high C:N ratios constitute the majority of crop residues produced by arable farming practices worldwide, typically involving cultivation of corn, wheat and rice (Medina *et al.*, 2015). The potential of crop residue soil amendments to deliver benefits to crops would be better exploited if the decomposition processes were manipulated for C to persist in the soil biomass, necromass or other forms of (semi-)stabilised SOM, such as in soil aggregates. Generally soil amendments consisting of one large amount of a single crop residue do not always deliver
- 100 benefits. We suggest that the non-additive decomposition rates observed in forest litter mixtures reinforced by recent insights into the link between CUE and the difference in C:N ratio of soil organic co-amendments, can inform strategies to obtain more benefits from crop residues as soil amendments. Mixing these crop residues to create chemically diverse crop-residue mixtures with a CUE-optimised C:N ratio to generate a greater diversity of functionally complementary microbial niches and to enable each member of the microbial community to function at a maximised CUE, could be a relatively simple method to
- 105 obtain more benefits from this precious, but ubiquitous, resource. If this approach can attain higher CUE levels for high-C:N residues, a considerable increase in net SOM could be realised in arable cropping systems, along with other beneficial changes in soil properties (e.g. nutrient retention) leading to greater soil fertility, and meanwhile increasing biodiversity in otherwise monocultural arable cropping systems.

The aim of this study was to investigate the potential of chemically heterogeneous mixtures of crop residue amendments to 110 improve soil properties for crop production. A field experiment was set up on an intensive organic arable cropping farm. Amendments of mixtures and individual crop residues were applied: vegetable waste compost was used as low-C:N (highquality) residue, and wheat straw and woodchips were used as high-C:N (low-quality) residues. Properties indicative of soil structure and nutrient cycling were used to assess benefits from residue mixtures compared to individual residues, including lettuce crop yield, soil respiration, soil aggregate stability and bulk density, SOM, available and potentially mineralisable N,

- 115 and available P, K and Mg. We predicted higher decomposition rates when mixtures of crop residues were applied compared to individual residue amendments, leading to non-additive effects in soil properties that could be beneficial for crop production. In particular, we hypothesised faster decomposition of residue mixtures to result in a higher soil respiration rate in the short term, as well as the release of greater levels of soil available nutrients (N, P, K, Mg) and SOM compared to <u>what</u> would be expected by combining the effects of individual residues, which leads to greater ammonification of residue N (Xu
- 120 <u>et al., 2006</u>), and, in turn, leads to a greater increase in pH (hypothesis 1). An increase in SOM will likely change soil physical properties, which we expected to observe as an increase in soil aggregate stability and a decrease in soil bulk density (hypothesis 2). These changes in soil physicochemical properties were subsequently expected to lead to a higher crop yield (hypothesis 3).

#### 2. Methodology

#### 125 2.1. Study site and experimental design

A field experiment was set up in an intensively managed horticultural area of lowland fen on an organic farm near Ely in Cambridgeshire, United Kingdom (52° 21' N; 0° 17' E). During the experiment, between 11 June 2018 and 26 July 2018, the field site was used for growing gem lettuce crops (*Lactuca sativa* L. var. longifolia, commercial variety 'Xamena'), following a year of celery crop in 2017, conversion to organic in 2016 (grass ley), winter wheat in 2015, and beetroot in

- 130 2014. The typical crop rotation followed by the farm is celery, followed by beetroot, celery or onion, followed by lettuce, followed by a break crop of perennial ryegrass and white clover or a cereal. The experimental plots were located on clay loam, on a roddon, a dried raised bed formed by the deposition of silt and clay from a watercourse which pushed peat to the sides. The mineral part of the soils typically do not perform as well as the surrounding organic soils because they require more fertiliser, so we expected they would respond more quickly to residue amendments.
- 135 Four replicates of six treatments, within a full-factorial randomised complete block design of the factors *compost* and *residue* (Table 1) were applied to 2 m × 6 m experimental plots within a 6 m × 48 m field site consisting of 3 × 8 = 24 plots situated between the tire tracks of farm machinery. All samples were taken from the inner 2 m × 2 m of each plot to incorporate a 4-metre long buffer zone between plots along the same strip.
- The residue amendment treatments were prepared on 17 May 2018. Application rates of the different amendments were 20 t
  ha<sup>-1</sup> compost (equivalent to 7 t ha<sup>-1</sup> dry matter), 13.3 t ha<sup>-1</sup> woodchips (equivalent to 8.7 t ha<sup>-1</sup> dry matter) and 10±0.8 t ha<sup>-1</sup> straw (equivalent to 9.2±0.8 t ha<sup>-1</sup> dry matter; ± indicates inclusive range of the straw application rate). These are within the range of application rates that are common in intensive arable cropping systems in Europe (Recous *et al.*, 1995; S. Gardner, 2018, pers. comm.), and were chosen to obtain similar amounts of dry matter for each residue. These rates were consistently applied in both individual amendment treatments and mixtures, so residue-compost treatments contained twice as much dry
- 145 matter compared to individual amendments. Applications were spread out evenly over the plots by hand on 12 June 2018 (Figure 1c), followed by power-harrowing to incorporate the residues in the soil profile. Gem lettuce plugs were sown the following day.

#### 2.2. Soil and residue characterisation

- Baseline soil samples were collected on 11 June 2018 (before organic amendments were applied). For each plot, soil samples were collected as the combination of five 30 mm diameter soil cores taken to 20 cm depth. These 24 composite samples were air-dried, disaggregated with the aid of a mortar and pestle, sieved to 2 mm and analysed for soil moisture (at 105 °C overnight), SOM by loss on ignition (LOI) (at 500 °C overnight), pH (after 2 hrs shaking 2.5 ±0.005 g soil with 25 ml Ultrapure water [> 18.2 Ω/cm]), and soil texture by laser granulometry (Malvern Mastersizer 3). A portion of each soil sample was ball milled and analysed for total C and N (Flash 2000, Thermo Fisher Scientific, Cambridge, U.K., calibrated
- 155 with aspartic acid, 104% N and 100% C recovery rates of in-house reference soil material traceable to GBW 07412). There were no significant treatment differences for any of these baseline soil variables, tested with a one-way analysis of variance (ANOVA) of treatments or a two-way ANOVA of the factors *residue* and *compost* (Supplement S2).

All amendments were provided by the farm and sourced and prepared on-site. The compost amendment was composed of the following vegetable residues from the farm: spinach, celery, several lettuce varieties, carrots, leeks, spring onions, onions and shallots, cabbage, bell peppers, beetroots, and mushrooms (Figure 1a-b). Due to the high water content of these residues, the farm co-composts with straw to provide sufficient dry matter content in the compost mixture. The straw amendment used in the treatments containing straw was winter wheat straw available on-site, and the woodchip amendment was from poplar trees commonly grown as a wind break in the local area. Dried and milled residues were analysed for total C and N (Flash 2000 as aforementioned, 109% recovery rate of both C and N of in-house reference rapeseed material, traceable to certified Deleted: fresh

Deleted: roughly

Deleted: Table

reference material GBW 07412). The total concentrations of P, K and Mg were determined by ICP-OES (inductively
coupled plasma optical emission spectroscopy, Perkin Elmer Optima 7300 Dual View, recovery rates of 99% for P, 94% for
K<sub>4</sub> 102% for Mg, 92% for Mn, and 114% for Fe of in-house hay reference material traceable to certified reference NCSDC
73349) analysis of 0.5 g residues samples digested in 8 ml of nitric acid (trace metal grade) using MARS 6 microwave digestion system (Table 2).

The amounts of C, N and other nutrients applied in each treatment were calculated based on the chemical characterisation of the residues and their application rates (Table  $\frac{3}{2}$ ).

#### 2.3. Assessment of yield

180

185

Lettuce crops were planted on 14 June 2018 and harvested from the inner 2 m  $\times$  2 m of each plot on 20 and 21 July 2018, i.e. 38 days after residue application and 36 days after planting. Each lettuce head was harvested whole and weighed to calculate the total biomass produced per treatment. Meanwhile lettuce crops were qualitatively assessed, which included screening for chlorosis, caterpillar damage, tip burn, and rotting. In some cases dried out mushrooms were found on the outer leaves, which was also noted.

#### 2.4. Assessment of soil biogeochemical properties

All soil samples were taken from the inner 2 m  $\times$  2 m of each plot on 26 July 2018, i.e. 44 days after residue application. From each plot a 10 cm deep bulk density core of 9.8 cm diameter was collected. A series of six 30 mm diameter soil cores to 20 cm depth were collected, combined and homogenised in a zip-lock bag, and used for a suite of analyses. A sub-sample

- of the fresh soil was sieved to 2 mm for analysis of available N (i.e. sum of NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup>) by 1 M KCl extraction before and after a 4-week incubation at 70% of the water-holding capacity (WHC). Extracts were filtered through a Whatman no. 2 filter and analysed colorimetrically for NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> on a Skalar San<sup>++</sup> continuous flow analyser. *Available N was taken as the sum of the* NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> *measured in the first extract. Potentially* mineralisable N was calculated as the difference in
- 190 NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> measured before and after the 4-week incubation period. A sub-sample of the fresh soil was sent to NRM laboratories (Bracknell, UK), where it was air-dried and sieved to 2 mm for measurement of available P by extraction with 0.5 M NaHCO<sub>3</sub>, available K and Mg by extraction with 1 M NH<sub>4</sub>NO<sub>3</sub>, soil particle size distribution by laser granulometry, SOM based on LOI at 430 °C, and the Solvita CO<sub>2</sub> burst test measuring the concentration of CO<sub>2</sub> produced by soils moistened to 50% of their WHC.
- 195 Earthworm and mesofauna sampling was performed, but only a few juvenile earthworms were found, which made identification difficult. The endogeic species *A. chlorotica* was identified in at least three of the 24 plots. The abundance of mesofauna (Collembola and mites) extracted from the soils using Tüllgen funnels was null. Some Collembola were observed while harvesting the lettuce crop, so their absence from the samples is probably due to the removal of plants that provided some shelter from the hot and dry weather conditions.
- 200 Wet aggregate stability was assessed as per Nimmo and Perkins (2002) using soil samples that were collected into tubs (to prevent soil compression) from the top 10 cm of each plot, and subsequently air-dried. A 4 g subsample from each plot was slowly pre-wetted on moistened filter paper. The wet sieving procedure involved a wet-sieving apparatus composed of vertically moving 250 µm sieves to hold the soil samples sitting inside a can. The cans were filled up with water such that the soil was submerged, causing the unstable soil aggregates to break apart and pass through the sieve into the can. First, the
- 205 soils were wet-sieved for 3 minutes in deionised water to collect unstable soil particles and subsequently in a solution of 2 g/L (NaPO<sub>3</sub>)<sub>6</sub> to disperse the water-stable aggregates. The stable fraction of soil (i.e. wet aggregate stability) was then calculated as the weight of soil caught by the dispersing solution divided by the sum of the weights of soil caught by both water and dispersant. Any particles larger than 250 µm did not pass the sieve and were not included in the calculation.

Deleted:	and
Deleted:	recovery rate of P, K and Mg, respectively,

Deleted: 3

### 2.5. Data analyses

We observed a gradient in the soil %C and a similar gradient in the %N content of the baseline soil samples that was not 215 well captured by our original blocking design, so the data were retrospectively blocked accordingly (Supplement S1). This was necessary because the calculation of non-additive effects, described below, relies on paired samples within blocks rather than treatment averages across blocks.

Statistical analyses were performed in R 3.5.1 (R Foundation for Statistical Computing) using RStudio 1.1.456 (RStudio, Inc.). To determine effects of treatments and/or factors on individual soil parameters, a two-way ANOVA, including

- 220 interactions, with the factors compost (compost or no compost) and residue (straw, woodchips or no residue) was performed. If a factor had a significant effect (p < 0.05), a post-hoc Tukey HSD test was run to determine which treatments were significantly different from each other. Taking into account that four replicates per treatment is a limited number of data points, assumptions of the ANOVA test were assessed both visually and via the relevant statistical tests: homoscedasticity was evaluated with a Q-Q plot of the ANOVA residuals plotted against the fitted data of the ANOVA, as well as a Levene
- 225 test of the data set. Normal distribution of the residuals was evaluated with a residuals-versus-fitted plot and a Shapiro-Wilk test of the residuals of the ANOVA. Pearson correlations were performed to investigate relationships between different variables

Properties indicative of soil structure and nutrient cycling were used to assess non-additive effects from residue mixtures compared to individual residues, including lettuce crop yield, soil respiration, soil aggregate stability and bulk density, SOM, available and potentially mineralisable N, and available P, K and Mg. The % effect of each measurement of the treatment

effects was first determined by adjusting to the measured effect of the control treatment:

$$\% effect = \frac{\text{treatment-control}}{\text{control}} \ 100\% \tag{1}$$

Next, the % non-additive effects of the residue mixtures were calculated as the difference between the % effect of the mixture and the % effect of the sum of the parts:

235  $\% \textit{ non-additive effect}_{mixture} = \ \% \textit{ effect}_{mixture} - (\% \textit{ effect}_{compost} + \% \textit{ effect}_{residue})$ where residue refers to straw or woodchips. A one-sided T-test of the % non-additive effects was performed with an alternate hypothesis (H<sub>1</sub>) of  $\mu > 0$  for yield, available N, potentially mineralisable N, available P, K, Mg, soil respiration, SOM, aggregate stability, and an alternate hypothesis of  $\mu < 0$  for bulk density and pH. Normality was tested with a Shapiro-Wilk test.

#### 240 3. Results

230

#### 3.1 Non-additive effects

Non-additive effects measured 44 days after application of the treatments were mostly synergistic (i.e. mixture > sum of the parts), although the majority of effects were not statistically significant (Figure 2). The magnitude and direction of deviation from additivity were usually similar for both the woodchip-compost and straw-compost mixtures, although non-additive

245 effects from the woodchip-compost mixture were sometimes less pronounced than those from the straw-compost mixture. Both compost-residue mixtures resulted in a non-additive increase in lettuce yield, available and potentially mineralisable N, available Mg, SOM, and soil respiration, but not in available K (hypothesis 1), some of which were statistically significant, as further specified below (Table 4). Most notably, we observed greater available N and SOM levels in soils to which a mixture of residues was applied, compared to the available N and SOM levels in treatments receiving only individual residue

amendments. The straw-compost mixture resulted in a significant (T = 4.022, p = 0.014) non-additive increase in SOM of 250 13.10%, and while the woodchip-compost mixture did not result in statistically significant non-additivity (T = 0.954, p = 0.205), it did result in a positive non-additive increase in SOM of 6.73%.

Deleted: 5

(2)

Likewise, amendment with the straw-compost mixture led to significantly (T = 3.789, p = 0.016) greater available N levels that were 55.06% higher on average than would have been expected from the available N levels in treatments receiving

- individual amendments of straw or compost only. The positive non-additive effect on available N observed in soils that received the woodchip-compost mixture was, however, smaller (7.16% increase on average) and not statistically significant (T = 0.235, p = 0.415). A non-significant non-additive increase in available P was only observed after application of the straw-compost mixture, but not after application of the woodchip-compost mixture (hypothesis 1). In agreement with, our
- 260 hypothesis, there was a non-additive increase in pH from the mixtures relative to individual amendments (hypothesis 1), although this was not significant (Table 4) and per-treatment results (discussed in next section) show that the pH decreased in all treatments relative to the control (F = 2.238; p = 0.095; one-way ANOVA; Supplement S<sup>2</sup>). We also observed non-additive effects from both compost-residue mixtures on the soil structure, i.e. a decrease in bulk density and an increase in aggregate stability (hypothesis 2), and a non-additive increase of about 10% was found for crop yield from both crop-residue mixtures (hypothesis 3). Although the effects on soil structure and yield were mostly non-significant, the decrease in bulk
- 265 mixtures (hypothesis 3). Although the effects on soil structure and yield were mostly non-significant, the decrease in bulk density after amendment with the straw-compost mixture was borderline significant (F = -2.232, p = 0.056) (Table 4). The following sections contain the per-treatment results of the soil physical and biochemical properties measured in this experiment. It should be noted that application rates of the mixtures were about twice as high as individual amendments to enable calculation of non-additivity, so measurements from residue-mixture treatments cannot be directly compared to
- 270 individual-residue treatments.

#### 3.2. Per-treatment results

295

Yield assessed by total biomass of gem lettuce produced per plot seemed to be somewhat reduced by the straw-only treatment but was not significantly affected by any of the treatments or factors (Figure 3a; see Supplement S<sup>5</sup> for statistical outputs).

275 Lettuce plants in the straw-only treatments suffered noticeably less damage, particularly from caterpillars, tip burn, and rot (Table <u>S2 in Supplement S3</u>). There was a significant interaction between *residue* and *compost* in terms of the qualities of lettuce plants harvested (F = 3.568, p = 0.050; two-way ANOVA), with the biggest difference between straw-only and straw-compost (p = 0.067; post-hoc Tukey HSD). Mushrooms were observed on the outer leaves of some lettuce heads in plots receiving woodchips, or in two cases in plots neighbouring treatments including woodchips, so fungi may have been introduced and/or promoted by woodchips.

Levels of SOM and N (available and potentially mineralisable) were negatively affected by the straw-only treatment, while treatments of woodchip-only and compost-only had little effect on SOM and N levels compared to the control (Figures 3b and 4). Residue mixtures increased SOM and N in most cases, with the exception of the effect of the straw-compost treatment on SOM. Nonetheless, there was a non-additive effect in SOM and N in the straw-compost treatment, as this non-

additivity was in fact a negation of the negative effect on SOM and N of straw applied as an individual residue.
Treatment effects on SOM or N levels were not significantly different between treatments (SOM: F = 0.981, p = 0.456; N: F = 1.81, p = 0.163; one-way ANOVA), but the factor *compost* tended to increase soil N (F = 3.88; p = 0.065; two-way ANOVA). Soil respiration in the different treatments was rather similar in all treatments and none of the treatments caused soil respiration to deviate significantly from the control or from each other (F = 1.358, p = 0.286; one-way ANOVA;
Supplement \$20.

The addition of compost, either as an individual residue or in a mixture, significantly affected soil available K (F = 7.761; p = 0.012) and Mg (F = 4.953; p = 0.039) (Figure 5a). Akin to soil N and SOM, the lowest levels of nutrients were found in soils amended with the straw-only treatment. The increases in nutrient availability were not consistent with the crop residue amendments and ranged from -242% to 57% of the nutrient added as part of the amendments (Supplement S4). If there was

an increase in nutrients, the contribution of the amendments was relatively small in most cases and exhibited very large error

Deleted: Contrary	
Deleted: to	
Deleted: 5	
Deleted: 2	

Deleted: 5

Deleted: 4

Deleted: 6

Oeleted: 2

- 305 margins. The most notable observations from these data is the consistent immobilisation of nutrients brought about by the straw-only treatment, while amendments including woodchips or compost had a tendency to modestly increase soil available nutrients. None of the nutrient increases exceeded 100% of the nutrients added, indicating that residue amendments did not result in net mobilisation of nutrients already present in the soil.
- We observed no significant effects on the aggregate stability of the differently amended soils, but the soil bulk density 310 tended to be lowered by the *residue* factor, i.e. when a low-quality residue was part of the treatment (F = 3.28; p = 0.062; two-way ANOVA) (Figure 5b).

#### 3.3. Correlations

A number of noteworthy correlations may help explain the data and are summarised in Table 5. There were some significant correlations between the amount of nutrients applied and the amount of available K and Mg in the soils at the end of the

315 experiment, which indicates a positive effect of the residue amendments. The amount of C applied via the residue amendments was not correlated with the levels of SOM. Yield was positively and significantly correlated with the sum of available and potentially mineralisable N, available P and Mg, SOM and aggregate stability. <u>SOM was also positively correlated with available N, P and Mg, and with soil respiration.</u>

#### 4. Discussion

### 320 4.1. Non-additive effects

The objective of this study was to find out if greater benefits could be obtained from crop-residue soil amendments in an arable soil by applying them as chemically heterogeneous mixtures of low-C:N vegetable waste compost with high-C:N straw or woodchips, compared to individual residue amendments. Relative benefits of the mixtures were assessed by calculating the non-additivity of a range of effects, including yield and a selection of soil properties that are likely to be

- 325 beneficial for crop production. We found some degree of non-additivity in the direction (synergy or antagonism) we predicted in most parameters (except available P in the woodchip-compost mixture and available K in both mixtures), and significant non-additive increases in available N and SOM after application of the straw-compost mixture, indicating that even after a short amount of time (44 days) beneficial effects from a mixture of residues can be greater than the sum of its parts.
- 330 Examining per-treatment effects can help further explain the non-additivity results. The per-treatment difference in terms of SOM and available N between the woodchip-compost treatment and the straw-compost treatment was relatively small. Yet, only the straw-compost mixture exhibited significant non-additivity. Comparison of the per-treatment effects on SOM and available N reveals that the significant non-additive effects observed after application of the straw-compost mixture are in fact a negation of the negative (compared to control) effect of the straw-only treatment. As suggested earlier, this indicates
- 335 that decomposition of single crop residue amendments does not always translate into agronomic benefits, and applying mixtures of crop residues could be a route to improve those benefits.

#### 4.2. Decomposition

Although we suggested that non-additive effects might be related to differences in decomposition rates in the mixtures compared to the individual residues, we have no evidence of this in terms of soil respiration measurements. At the time of

340 sampling, high microbial activity may have increased N immobilisation and therefore decreased soil mineral N availability. However, respiration rates were equally low in the straw-only (N immobilisation) and the straw-compost treatments (N mineralisation), and both were lower than the control (Supplement S3). Likewise, Redin *et al.* (2014), who studied residue mixtures of stems and leaves of 25 different arable crop species, found mostly additive effects for decomposition rates of Deleted: 7

mixtures, but unlike the results presented here they found no synergistic effects on N mineralisation. Both here and in the study by Redin *et al.* (2014), decomposition was measured in terms of C mineralisation (measured as CO<sub>2</sub> release), which does not account for the possibility of a higher CUE when chemically diverse residue mixtures are applied, and also does not distinguish between mineralisation of residues or organic matter already present in the soil. Moreover, our soil respiration measurements were taken by the Solvita burst method, on soil samples removed from the field and sieved to 2 mm removing parts of residues and other organic matter greater than 2mm, which may not have been a good representation of the

respiration produced in-situ by a soil mixed with crop residues at various stages of decomposition.

Another reason for the absence of different soil respiration rates may be the relatively short duration of this experiment, covering the short growing period of gem lettuce. As pointed out by Lecerf *et al.* (2011), niche complementarity effects, in

355 which different groups of decomposing organisms (already present in the soil, or newly introduced via the residues) develop a synergistic association in residue breakdown, tend to advance with time, leading to a generally higher number of long-term litter-mixing studies finding non-additive effects. Indeed, Ball *et al.* (2014) only observed a non-additive effect on mass loss in a five-component mixture after 193 days. Therefore an experiment of longer duration may be able to capture more and greater treatment effects and non-additive effects.

#### 360 4.3. Yield

350

Although yield, assessed by total biomass of gem lettuce produced per plot, was not significantly affected by any of the treatments or factors, there were some notable differences between treatments. Yield appeared to be somewhat depressed by the straw-only treatment, which is not surprising considering the lower concentration of soil available N, SOM, soil nutrients and aggregate stability in this treatment, compared to the control. Crops tend to require most nitrogen during the vegetative

- 365 growth stage and when this is not available, yield will be affected (Chen *et al.*, 2014). The lettuce plants were planted as plugs just after application of the treatments, so when they were introduced to the experimental plots they were already in their vegetative stage. Significant positive correlations of yield with the sum of available and potentially mineralisable N, available P and Mg, SOM, and aggregate stability suggest that these are the main benefits provided by the crop residue amendments from an agronomic perspective.
- Overall lettuce quality was least affected in the straw-only treatments, despite the location of these treatments being towards the low soil-C end of the field site (Supplement S1). Available N levels were positively correlated with overall quality impairment (i.e. % lettuce heads affected by some form of quality impairment) (p = 0.011), and in particular with yellow tips (p = 0.017) and tip burn (p = 0.041), which may indicate the crop was suffering from N deficiency (Table 5). Indeed, the N levels were relatively low compared to those recommended for lettuce crops (RB209, 2019), and N deficiency leads to
- 375 reduced plant size, which would lead to decreased biomass production, as well as chlorosis and outside leaves senescing prematurely and dropping off (Brady and Weil, 2002), all of which were observed.

#### 4.4. Nutrient dynamics and transfer

The straw-only treatment led to a notable immobilisation of N, which was unlike the other treatments. Although this could be only a temporary effect (e.g. as in Silgram and Chambers, 2002), it may be unfavourable for lettuce crop productivity and

- 380 should be taken into account when timing crop residue applications. The notable N immobilisation in the straw-only treatment suggests that straw decomposed differently as an individual residue than in a mixture with compost, which could be explained by the C:N ratio of the treatments. Chen *et al.* (2014) evaluated soil N processing during crop residue decomposition and suggested that residues with a C:N ratio below ~25 result in net mineralisation (increase in soil available N) and those with a C:N ratio above ~30 result in net immobilisation (decrease in soil available N). Therefore, in the present
- 385 study the woodchip-only (C:N = 64) and straw-only (C:N = 41) treatments are both expected to result in net N immobilisation. The reason why N immobilisation is only observed in the straw treatment could be due to a lower

Deleted: 4

decomposition rate of the woodchips and therefore lower microbial N-mining requirement at the time of sampling. Straw is likely more decomposable due to a comparatively lower C:N ratio, a higher water-holding capacity (being more friable and having a greater surface area to hold on to moisture) (Hättenschwiler *et al.*, 2005; Iqbal *et al.*, 2015) and possibly a soil microbial community that is more adapted to decomposing straw because wheat is sometimes grown in these soils.

A slight increase in soil N (available and potentially mineralisable N) observed in the straw-compost treatment and to a lesser extent in the woodchip-compost treatment, compared to the control, could be due to N derived from the compost, the
 residue, or primed native SOM. Priming of native SOM caused by the amendment seems unlikely in the woodchip-compost treatment, because SOM levels were higher compared to the control treatment. Even in the straw-compost treatment, the

SOM level was very close to that of the control treatment, suggesting <u>net</u> mineralisation of native SOM <u>as a result of the</u> <u>residue amendment</u> was negligible. Compost was the most significant factor related to higher soil N levels, which can be attributed to its low C:N ratio, allowing for easy decomposition with minimal immobilisation of native soil mineral N. In the 400 residue mixtures, it is likely that compost provided nutrients for decomposer microbes to be able to decompose the high-C:N

residues (i.e. inter-specific nutrient transfer). Therefore, the non-additive effects on soil N in the straw-compost treatment can probably be attributed to interspecific net transfer of N from high-N to low-N residues resulting in (1) the retention of compost-derived N by straw or woodchips in the

- mixture, preventing it from being leached, and (2) a higher nutrient availability in treatments including compost, enabling
  decomposer organisms to break down and release N contained in the amendment mixture more readily. The transfer of N can occur by a combination of uptake and release by microbes on the high-N residue as they produce enzymes for decomposition, and diffusion along a gradient of high N to low N (Schimel and Hättenschwiler, 2007). The woodchips likely had a higher lignin content than straw. Ligninolytic enzyme production can be inhibited by elevated N concentrations (Carreiro *et al.*, 2000; Knorr *et al.*, 2005), resulting in a relatively greater inhibition of decomposition of the woodchips.
- 410 The transfer of N in litter mixtures appears to go hand in hand with a C transfer. In a microcosm experiment by Berglund *et al.* (2013) on pine and maize litters inoculated with both forest and arable soils, mixing residues mostly increased C loss from the lower quality litter, while C released from the higher quality litter was equivalent to decomposing as an individual litter. Therefore, the non-additively higher SOM in the straw-compost treatment is likely to be the result of enhanced C release from the straw due to the addition of compost. This phenomenon could be explained by a bidirectional transfer of C
- 415 and N between high- and low-quality residues e.g. via transport of amino acids by fungal mycelia (Tlalka *et al.*, 2007) where increased N availability near the low-quality residue enhances its decomposition and subsequent C release, while increased C in the presence of the high-quality residue has little effect on its decomposition (Berglund *et al.*, 2013).

#### 4.5. Soil physical structure

Increased SOM positively affects aggregate stability because soil microbes feeding on organic substrates enhance soil 420 aggregate formation and stability by biofilm formation and the production of extracellular polymeric substances that increase cohesion between soil particles (Martens, 2000; Totsche *et al.*, 2018). Aggregate stability, in turn, is involved in the protection of mineral-associated SOM (Angst *et al.*, 2017). Therefore, with an increase in SOM, an increase in aggregate stability would be expected, and we did indeed observe a positive correlation between these variables (p = 0.028). We also observed a positive correlation between aggregate stability and soil available N (p = 0.005). This is contrary to the 425 observation that high-quality residues and/or addition of N fertilisers result in higher aggregate turnover (formation and

breakdown) compared to a greater aggregate stability when low-quality residues are applied (Chivenge *et al.*, 2011). Because we observed positive effects on both soil N and SOM from crop residue mixtures, an increased non-additive effect on the soil physical structure from application of the right residue mixtures can therefore be anticipated over time. However, in many arable cropping systems tillage may undermine the emergence of this benefit by destroying soil aggregates and

430 exposing the SOM contained within (Nath and Lal, 2017). Furthermore, bulk density was lowered by the addition of the

low-quality residues (straw and woodchips; p = 0.062), especially when combined with compost. This could be partially due to increases in the aggregate stability in most of these treatments, although some residues (with a lower density than soil) may have also been included in the bulk density ring when sampling.

#### 4.6. Potential of residue mixing to obtain more benefits from low-quality residues

- 435 Our study provides some evidence that chemically heterogeneous crop residue mixtures can provide agronomically beneficial non-additive effects. We found prevention of N immobilisation to be the most prominent effect in the short term. Positive non-additivity in SOM levels and other soil nutrients may develop over time, but a longer term experiment is necessary to investigate this.
- Other authors have also found beneficial effects on soil N levels from mixed residue amendments. For instance, Kaewpradit 440 *et al.* (2009) mixed groundnut residues (high N) and rice straw (low N), which slowed down N loss by mineralisation during the phase between two different crops, i.e. a beneficial temporary N immobilisation. McDaniel *et al.* (2016) found that nonadditive effects of soil C and N dynamics after application of residue mixtures depend on the diversity in cropping history, with non-additive effects primarily observed in monoculture soils rather than diverse crop rotations. The authors attribute this to the low respiration rates from monoculture soils after application of low-quality residues, while soil response to high-
- quality residues is similar in both monoculture and crop rotation soils (McDaniel *et al.*, 2016). These studies indicate that potential benefits from residue-mixing are dependent on the arable cropping system.
  Manipulation of the number of component residues, the mixing ratio, and the quantity applied can be used to optimise timing and amount of nutrient release for a better synchrony with crop demand (Myers *et al.*, 1997). For instance Kuo and Sainju
- (1998) demonstrated that the timing of N mineralisation can be manipulated by the proportion of leguminous cover crop residues in the mixture, while Mao and Zeng (2012) found that both the number of residue components and their mixing ratio affected non-additivity. Furthermore, the quantity of residues applied can impact on microbial CUE: while microbial CUE is often unaffected at low substrate additions, applications of high amounts of the same material can lead to diminishing CUE levels (Jones *et al.*, 2018), e.g. as shown by Roberts *et al.*, 2007 with glucose and glucosamine additions to various foraging soil types in a microcosm experiment.
- 455 The interplay of environmental factors and amendment properties affect microbial CUE and the mechanisms involved in non-additivity of decomposing residue mixtures on soil properties (Kuebbing and Bradford, 2019), which need to be accounted for to be able to create a methodology for optimised benefits from crop residues as soil amendments in arable cropping systems. Therefore, future research on residue mixtures should incorporate not only substrate quality, but also application rate (quantity), diversity (number of residue species) and mixing ratio and how these interact with different 460 arable soil types.

#### 5. Conclusions

This experiment tested agronomic benefits obtained from multi-component and chemically heterogeneous residue mixtures compared to the individual residues. Significant positive non-additive effects on available N and SOM were measured after application of a straw-compost mixture, so we can partially accept our first hypothesis predicting greater levels of soil available nutrients and SOM in mixtures compared to individual residues. However, due to variation in the total %C contents across the experimental field site, we have some reservations about this result. Nevertheless, this study provides some evidence for the potential of crop residue mixtures to provide greater agronomic benefits than single high-C residue amendments of straw or woodchips, at least in terms of preventing N immobilisation during crop growth.

#### Data availability

470 Data have been uploaded on Mendeley Data, doi:10.17632/jcrvmb8hwy.1

#### Supplement link

[See relevant document in our submission]

#### Author contributions

MS and TS designed the experiment and performed field work. MS carried out laboratory work. MS analysed the data with support from TS and AW. MS prepared the manuscript with critical review by all authors. TS secured funding and established contact with the farm where the experiment was carried out. TS supervised the project and AW and SM cosupervised the project.

#### **Competing interests**

The authors declare that they have no conflict of interest.

#### 480 Acknowledgements

This work was funded by a University of Reading Faculty of Science/SAGES Studentship awarded to Marijke Struijk, and research expenses provided by the Waitrose Agronomy Group. <u>AW acknowledges support from the BBSRC-funded Soil to Nutrition programme (BBS/E/C/00010330)</u>. We thank G's and their employees for supply of materials and access to the field site, Xin Shu and Adetunji Adekanmbi for help in the field, Omar El-Huni and Alfonso Rodriguez Vila for help with lab work, and Anne S. Dudley, Karen J. Gutteridge, Ilse Kamerling, Fengjuan Xiao and Chris Speed for technical assistance in

the lab.

#### References

485

- Al-Maliki, S. and Scullion, J.: Interactions between earthworms and residues of differing quality affecting aggregate stability and microbial dynamics, Appl. Soil Ecol., 64, 56–62, doi:10.1016/J.APSOIL.2012.10.008, 2013.
- 490 Allison, S. D., Wallenstein, M. D. and Bradford, M. A.: Soil-carbon response to warming dependent on microbial physiology, Nat. Geosci., 3(5), 336–340, doi:10.1038/ngeo846, 2010.

Aneja, M. K., Sharma, S., Fleischmann, F., Stich, S., Heller, W., Bahnweg, G., Munch, J. C. and Schloter, M.: Microbial colonization of beech and spruce litter - Influence of decomposition site and plant litter species on the diversity of microbial community, Microb. Ecol., 52(1), 127–135, doi:10.1007/s00248-006-9006-3, 2006.

- 495 Angst, G., Mueller, K. E., Kögel-Knabner, I., Freeman, K. H. and Mueller, C. W.: Aggregation controls the stability of lignin and lipids in clay-sized particulate and mineral associated organic matter, Biogeochemistry, 132(3), 307–324, doi:10.1007/s10533-017-0304-2, 2017.
  - Bailey, V. ., Smith, J. . and Bolton, H.: Fungal-to-bacterial ratios in soils investigated for enhanced C sequestration, Soil Biol. Biochem., 34(7), 997–1007, doi:10.1016/S0038-0717(02)00033-0, 2002.
- 500 Ball, B. A., Carrillo, Y. and Molina, M.: The influence of litter composition across the litter-soil interface on mass loss, nitrogen dynamics and the decomposer community, Soil Biol. Biochem., 69, 71–82, doi:10.1016/J.SOILBIO.2013.10.048, 2014.

Bastian, F., Bouziri, L., Nicolardot, B. and Ranjard, L.: Impact of wheat straw decomposition on successional patterns of soil microbial community structure, Soil Biol. Biochem., 41, 262–275, doi:10.1016/j.soilbio.2008.10.024, 2009.

505 Berglund, S. L., Ågren, G. I. and Ekblad, A.: Carbon and nitrogen transfer in leaf litter mixtures, Soil Biol. Biochem., 57, 341–348, doi:10.1016/J.SOILBIO.2012.09.015, 2013. Brady, N. C. and Weil, R. R.: The Nature and Properties of Soils, 13th ed., Prentice Hall, Upper Saddle River, New Jersey., 2002.

- Briones, M. J. I. and Ineson, P.: Decomposition of eucalyptus leaves in litter mixtures, Soil Biol. Biochem., 28(10–11), 1381–1388, doi:10.1016/S0038-0717(96)00158-7, 1996.
  - Carreiro, M. M., Sinsabaugh, R. L., Repert, D. A. and Parkhurst, D. F.: Microbial enzyme shifts explain litter decay responses to simulated nitrogen deposition, Ecology, 81(9), 2359–2365, doi:10.1890/0012-9658(2000)081[2359:MESELD]2.0.CO;2, 2000.
- Catt, J. A., Howse, K. R., Christian, D. G., Lane, P. W., Harris, G. L. and Goss, M. J.: Strategies to decrease nitrate leaching
   in the Brimstone Farm Experiment, Oxfordshire, UK, 1988–93: the effect of straw incorporation, J. Agric. Sci., 131(3), 309–320, doi:10.1017/S0021859698005905, 1998.
  - Chen, B., Liu, E., Tian, Q., Yan, C. and Zhang, Y.: Soil nitrogen dynamics and crop residues. A review, Agron. Sustain. Dev., 34(2), 429–442, doi:10.1007/s13593-014-0207-8, 2014.
- Chivenge, P., Vanlauwe, B., Gentile, R. and Six, J.: Organic resource quality influences short-term aggregate dynamics and soil organic carbon and nitrogen accumulation, Soil Biol. Biochem., 43(3), 657–666, doi:10.1016/J.SOILBIO.2010.12.002, 2011.
  - Cong, W.-F., van Ruijven, J., Mommer, L., De Deyn, G. B., Berendse, F. and Hoffland, E.: Plant species richness promotes soil carbon and nitrogen stocks in grasslands without legumes, edited by S. Lavorel, J. Ecol., 102(5), 1163–1170, doi:10.1111/1365-2745.12280, 2014.
- 525 Cosentino, D., Chenu, C. and Le Bissonnais, Y.: Aggregate stability and microbial community dynamics under dryingwetting cycles in a silt loam soil, Soil Biol. Biochem., 38(8), 2053–2062, doi:10.1016/J.SOILBIO.2005.12.022, 2006.
  - Cotrufo, M. F., Soong, J. L., Horton, A. J., Campbell, E. E., Haddix, M. L. L., Wall, D. H. and Parton, W. J.: Formation of soil organic matter via biochemical and physical pathways of litter mass loss, Nat. Geosci., 8 [online] Available from: http://www.nature.com/articles/ngeo2520 (Accessed 17 October 2019), 2015.
- 530 Eisenhauer, N., Dobies, T., Cesarz, S., Hobbie, S. E., Meyer, R. J., Worm, K. and Reich, P. B.: Plant diversity effects on soil food webs are stronger than those of elevated CO2 and N deposition in a long-term grassland experiment., Proc. Natl. Acad. Sci. U. S. A., 110(17), 6889–94, doi:10.1073/pnas.1217382110, 2013.
- Fierer, N., Lauber, C. L., Ramirez, K. S., Zaneveld, J., Bradford, M. A. and Knight, R.: Comparative metagenomic, phylogenetic and physiological analyses of soil microbial communities across nitrogen gradients, ISME J., 6(5), 1007– 1017, doi:10.1038/ismej.2011.159, 2012.
  - Garnier, E. and Navas, M.-L.: A trait-based approach to comparative functional plant ecology: concepts, methods and applications for agroecology. A review, Agron. Sustain. Dev., 32(2), 365–399, doi:10.1007/s13593-011-0036-y, 2012.
  - Gartner, T. B. and Cardon, Z. G.: Decomposition dynamics in mixed-species leaf litter, Oikos, 104(2), 230–246, doi:10.1111/j.0030-1299.2004.12738.x, 2004.
- 540 Gurr, G. M., Wratten, S. D. and Luna, J. M.: Multi-function agricultural biodiversity: pest management and other benefits, Basic Appl. Ecol., 4(2), 107–116, doi:10.1078/1439-1791-00122, 2003.
  - Hättenschwiler, S., Tiunov, A. and Scheu, S.: Biodiversity and litter decomposition in terrestrial ecosystems, Annu. Rev. Ecol. Evol., 36, 191–218, doi:10.1146/annurev.ecolsys.36.112904.151932, 2005.
- Ho, A., Lonardo, D. P. Di and Bodelier, P. L. E.: Revisiting life strategy concepts in environmental microbial ecology,
   FEMS Microbiol. Ecol., 93(3), fix006, doi:10.1093/femsec/fix006, 2017.
  - Iqbal, A., Aslam, S., Alavoine, G., Benoit, P., Garnier, P. and Recous, S.: Rain regime and soil type affect the C and N dynamics in soil columns that are covered with mixed-species mulches, Plant Soil, 393(1–2), 319–334, doi:10.1007/s11104-015-2501-x, 2015.
- Jones, D. L., Hill, P. W., Smith, A. R., Farrell, M., Ge, T., Banning, N. C. and Murphy, D. V.: Role of substrate supply on microbial carbon use efficiency and its role in interpreting soil microbial community-level physiological profiles (CLPP), Soil Biol. Biochem., 123, 1–6, doi:10.1016/j.soilbio.2018.04.014, 2018.
  - Kaewpradit, W., Toomsan, B., Cadisch, G., Vityakon, P., Limpinuntana, V., Saenjan, P., Jogloy, S. and Patanothai, A.: Mixing groundnut residues and rice straw to improve rice yield and N use efficiency, F. Crop. Res., 110(2), 130–138, doi:10.1016/J.FCR.2008.07.011, 2009.
- 555 Kallenbach, C. M., Wallenstein, M. D., Schipanksi, M. E. and Grandy, A. S.: Managing agroecosystems for soil microbial carbon use efficiency: ecological unknowns, potential outcomes, and a path forward, Front. Microbiol., 10(1146), doi:10.3389/fmicb.2019.01146, 2019.
  - Knorr, M., Frey, S. D. and Curtis, P. S.: Nitrogen additions and litter decomposition: A meta-analysis, Ecology, 86(12), 3252–3257, doi:10.1890/05-0150, 2005.

560 Kuebbing, S. E. and Bradford, M. A.: The potential for mass ratio and trait divergence effects to explain idiosyncratic impacts of non-native invasive plants on carbon mineralization of decomposing leaf litter, edited by E. Morriën, Funct. Ecol., 33(6), 1365-2435.13316, doi:10.1111/1365-2435.13316, 2019.

Kumar, K. and Goh, K. M.: Crop residues and management practices: effects on soil quality, soil nitrogen dynamics, crop yield, and nitrogen recovery, Adv. Agron., 68, 197–319, doi:10.1016/S0065-2113(08)60846-9, 1999.

565 Kuo, S. and Sainju, U. M.: Nitrogen mineralization and availability of mixed leguminous and non-leguminous cover crop residues in soil, Biol. Fertil. Soils, 26(4), 346–353, doi:10.1007/s003740050387, 1998.

Lange, M., Eisenhauer, N., Sierra, C. A., Bessler, H., Engels, C., Griffiths, R. I., Mellado-Vázquez, P. G., Malik, A. A., Roy, J., Scheu, S., Steinbeiss, S., Thomson, B. C., Trumbore, S. E. and Gleixner, G.: Plant diversity increases soil microbial activity and soil carbon storage, Nat. Commun., 6(1), 6707, doi:10.1038/ncomms7707, 2015.

- 570 Lecerf, A., Marie, G., Kominoski, J. S., LeRoy, C. J., Bernadet, C. and Swan, C. M.: Incubation time, functional litter diversity, and habitat characteristics predict litter-mixing effects on decomposition, Ecology, 92(1), 160–169, doi:10.1890/10-0315.1, 2011.
- Li, X.-M., Chen, Q.-L., He, C., Shi, Q., Chen, S.-C., Reid, B. J., Zhu, Y.-G. and Sun, G.-X.: Organic carbon amendments affect the chemodiversity of soil dissolved organic matter and its associations with soil microbial communities, Environ. Sci. Technol., 53, 50–59, doi:10.1021/acs.est.8b04673, 2019.
  - Ludwig, M., Achtenhagen, J., Miltner, A., Eckhardt, K.-U., Leinweber, P., Emmerling, C. and Thiele-Bruhn, S.: Microbial contribution to SOM quantity and quality in density fractions of temperate arable soils, Soil Biol. Biochem., 81, 311– 322, doi:10.1016/J.SOILBIO.2014.12.002, 2015.
- Makkonen, M., Berg, M. P., van Logtestijn, R. S. P., van Hal, J. R. and Aerts, R.: Do physical plant litter traits explain non-additivity in litter mixtures? A test of the improved microenvironmental conditions theory, Oikos, 122(7), 987–997, doi:10.1111/j.1600-0706.2012.20750.x, 2013.
  - Malézieux, E., Crozat, Y., Dupraz, C., Laurans, M., Makowski, D., Ozier-Lafontaine, H., Rapidel, B., Tourdonnet, S. and Valantin-Morison, M.: Mixing plant species in cropping systems: concepts, tools and models. A review, Agron. Sustain. Dev., 29(1), 43–62, doi:10.1051/agro:2007057, 2009.
- 585 Mao, R. and Zeng, D.-H.: Non-additive effects vary with the number of component residues and their mixing proportions during residue mixture decomposition: A microcosm study, Geoderma, 170, 112–117, doi:10.1016/J.GEODERMA.2011.11.008, 2012.

Martens, D. A.: Plant residue biochemistry regulates soil carbon cycling and carbon sequestration, Soil Biol. Biochem., 32(3), 361–369, doi:10.1016/S0038-0717(99)00162-5, 2000.

590 Martin, J. P., Martin, W. P., Page, J. B., Raney, W. A. and de Ment, J. D.: Soil aggregation, Adv. Agron., 7, 1–37, doi:10.1016/S0065-2113(08)60333-8, 1955.

McDaniel, M. D., Grandy, A. S., Tiemann, L. K. and Weintraub, M. N.: Eleven years of crop diversification alters decomposition dynamics of litter mixtures incubated with soil, Ecosphere, 7(8), e01426, doi:10.1002/ecs2.1426, 2016.

- Medina, J., Monreal, C., Barea, J. M., Arriagada, C., Borie, F. and Cornejo, P.: Crop residue stabilization and application to agricultural and degraded soils: A review, Waste Manag., 42, 41–54, doi:10.1016/j.wasman.2015.04.002, 2015.
- Myers, R. J. K., Noordwijk, M. van and Vityakon, P.: Synchrony of nutrient relase and plant demand: plant litter quality, soil environment and farmer management options., in Driven by Nature: Plant Litter Quality and Decomposition, edited by G. Cadish and K. E. Giller, pp. 215–229., 1997.
- Nath, A. J. and Lal, R.: Effects of tillage practices and land use management on soil aggregates and soil organic carbon in the north appalachian region, USA, Pedosphere, 27(1), 172–176, doi:10.1016/S1002-0160(17)60301-1, 2017.
- Nilsson, M.-C., Wardle, D. A. and DeLuca, T. H.: Belowground and aboveground consequences of interactions between live plant species mixtures and dead organic substrate mixtures, Oikos, 117(3), 439–449, doi:10.1111/j.2007.0030-1299.16265.x, 2008.
- Nimmo, J. R. and Perkins, K. S.: Aggregate stability and size distribution, in Methods of soil analysis, Part 4: Physical methods., edited by J. H. Dane and G. C. Topp, Soil Science Society of America, Madison, WI., 2002.
  - Otsing, E., Barantal, S., Anslan, S., Koricheva, J. and Tedersoo, L.: Litter species richness and composition effects on fungal richness and community structure in decomposing foliar and root litter, Soil Biol. Biochem., 125, 328–339, doi:10.1016/J.SOILBIO.2018.08.006, 2018.
- Pérez Harguindeguy, N., Blundo, C. M., Gurvich, D. E., Díaz, S. and Cuevas, E.: More than the sum of its parts? Assessing
   litter heterogeneity effects on the decomposition of litter mixtures through leaf chemistry, Plant Soil, 303, 151–159,
   doi:10.1007/s11104-007-9495-y, 2008.

Powlson, D. S., Glendining, M. J., Coleman, K. and Whitmore, A. P.: Implications for soil properties of removing cereal straw: Results from long-term studies, Agron. J., 103(1), 279–287, doi:10.2134/agronj2010.0146s, 2011.

- Purahong, W., Wubet, T., Lentendu, G., Schloter, M., Pecyna, M. J., Kapturska, D., Hofrichter, M., Krüger, D. and Buscot,

   615
   F.: Life in leaf litter: novel insights into community dynamics of bacteria and fungi during litter decomposition, Mol.

   Ecol., 25(16), 4059–4074, doi:10.1111/mec.13739, 2016.
  - Quemada, M. and Cabrera, M. L.: Carbon and nitrogen mineralized from leaves and stems of four cover crops, Soil Sci. Soc. Am. J., 59(2), 471, doi:10.2136/sssaj1995.03615995005900020029x, 1995.

RB209: Nutrient Management Guide (RB209). Section 6: Vegetable and bulbs., 2019.

- 620 Recous, S., Robin, D., Darwis, D. and Mary, B.: Soil inorganic N availability: Effect on maize residue decomposition, Soil Biol. Biochem., 27(12), 1529–1538, doi:10.1016/0038-0717(95)00096-W, 1995.
  - Redin, M., Recous, S., Aita, C., Dietrich, G., Skolaude, A. C., Ludke, W. H., Schmatz, R. and Giacomini, S. J.: How the chemical composition and heterogeneity of crop residue mixtures decomposing at the soil surface affects C and N mineralization, Soil Biol. Biochem., 78, 65–75, doi:10.1016/J.SOILBIO.2014.07.014, 2014.
- 625 Roberts, P., Bol, R. and Jones, D. L.: Free amino sugar reactions in soil in relation to soil carbon and nitrogen cycling, Soil Biol. Biochem., 39(12), 3081–3092, doi:10.1016/J.SOILBIO.2007.07.001, 2007.
  - Roller, B. R. and Schmidt, T. M.: The physiology and ecological implications of efficient growth, ISME J., 9(7), 1481–1487, doi:10.1038/ismej.2014.235, 2015.
- Santonja, M., Foucault, Q., Rancon, A., Gauquelin, T., Fernandez, C., Baldy, V. and Mirleau, P.: Contrasting responses of
   bacterial and fungal communities to plant litter diversity in a Mediterranean oak forest, Soil Biol. Biochem., 125, 27– 36, doi:10.1016/J.SOILBIO.2018.06.020, 2018.
  - Schimel, J. P. and Hättenschwiler, S.: Nitrogen transfer between decomposing leaves of different N status, Soil Biol. Biochem., 39(7), 1428–1436, doi:10.1016/J.SOILBIO.2006.12.037, 2007.
- Schimel, J., Balser, T. C. and Wallenstein, M.: Microbial stress-response physiology and its implications for ecosystem function, Ecology, 88(6), 1386–1394, doi:10.1890/06-0219, 2007.
  - Seastedt, T. R.: The role of microarthropods in decomposition and mineralization processes, Annu. Rev. Entomol., 29, 25– 46 [online] Available from: http://www.annualreviews.org/doi/pdf/10.1146/annurev.en.29.010184.000325 (Accessed 20 February 2018), 1984.
- Silgram, M. and Chambers, B. J.: Effects of long-term straw management and fertilizer nitrogen additions on soil nitrogen
   supply and crop yields at two sites in eastern England, J. Agric. Sci., 139(2), 115–127, doi:10.1017/S0021859602002435, 2002.
  - Simpson, A. J., Simpson, M. J., Smith, E. and Kelleher, B. P.: Microbially derived inputs to soil organic matter: are current estimates too low?, Environ. Sci. Technol., 41(23), 8070–8076, doi:10.1021/es071217x, 2007.
- Six, J., Frey, S. D., Thiet, R. K. and Batten, K. M.: Bacterial and fungal contributions to carbon sequestration in agroecosystems, Soil Sci. Soc. Am. J., 70(2), 555, doi:10.2136/sssaj2004.0347, 2006.
- Smil, V.: Crop residues : Agriculture's largest harvest, Bioscience, 49(4), 299-308, 1999.
  - Thomsen, I. K. and Christensen, B. T.: Yields of wheat and soil carbon and nitrogen contents following long-term incorporation of barley straw and ryegrass catch crops, Soil Use Manag., 20(4), 432–438, doi:10.1111/j.1475-2743.2004.tb00393.x, 2006.
- 650 Tilman, D., Cassman, K. G., Matson, P. A., Naylor, R. and Polasky, S.: Agricultural sustainability and intensive production practices, Nature, 418, 671–677, doi:10.1038/nature01014, 2002.
  - Tlalka, M., Bebber, D. P., Darrah, P. R., Watkinson, S. C. and Fricker, M. D.: Emergence of self-organised oscillatory domains in fungal mycelia, Fungal Genet. Biol., 44(11), 1085–1095, doi:10.1016/J.FGB.2007.02.013, 2007.
- Totsche, K. U., Amelung, W., Gerzabek, M. H., Guggenberger, G., Klumpp, E., Knief, C., Lehndorff, E., Mikutta, R., Peth,
   S., Prechtel, A., Ray, N. and Kögel-Knabner, I.: Microaggregates in soils, J. Plant Nutr. Soil Sci., 181(1), 104–136,
   doi:10.1002/jpln.201600451, 2018.

Wardle, D. A., Bonner, K. I. and Nicholson, K. S.: Biodiversity and plant litter: experimental evidence which does not support the view that enhanced species richness improves ecosystem function, Oikos, 79(2), 247–258, 1997.

Xu, J. M., Tang, C. and Chen, Z. L.: The role of plant residues in pH change of acid soils differing in initial pH, Soil Biol.
 Biochem., 38(4), 709–719, doi:10.1016/j.soilbio.2005.06.022, 2006.

Figures



Figure 1: Photographs of the preparation of the mixed compost (a), the final compost product (b), the treatments applied on the experimental plots (c), and the lettuce at time of harvest (d).



• straw-compost mixture • woodchip-compost mixture

Figure 2: Non-additive effects of crop-residue mixtures on soil properties. The % non-additive effect is the difference in % effect between the mixture and the sum of the parts. Positive % non-additive effects mean that the effect of the mixture is greater than the sum of the parts, and vice versa. Yield is total lettuce biomass produced per plot, Av. N is available N, Min. N is potentially mineralisable N, soil P, K, and Mg are soil available nutrients, SOM measured as LOI, soil respiration assessed by CO<sub>2</sub> burst. Error bars represent SEM (n = 4). Significant difference from zero (where 0 = no significant non-additivity) is indicated by \* (one-tailed T-test, p < 0.05).



Figure 3: (a) Gem lettuce yield as total biomass produced per 2 m  $\times$  2 m plot sampled. (b) Soil organic matter by percent loss on ignition (% LOI) after each soil amendment treatment. Lower and upper hinges correspond to the 25th and 75th percentiles; black dots represent individual data points, occasionally overlapping (n = 4).





Figure 4: Soil available and potentially mineralisable N after each soil amendment treatment. Error bars represent SEM of available and potentially mineralisable N separately (n = 4).



Figure 5: (a) Soil available nutrients after each soil amendment treatment. (b) Soil physical properties after each treatment. Lower and upper hinges correspond to the 25th and 75th percentiles; black dots represent individual data points, occasionally overlapping (n = 4).

Tables

#### 670

## Table 1: Treatment structure composed of the factors *residue* and *compost*.

$\begin{array}{c} compost \rightarrow \\ residue \downarrow \end{array}$	compost	no compost
straw	straw-compost	straw
woodchips	woodchip-comp	woodchip
none	compost	control

Table 2: Residue characterisation (SEM indicated in parentheses, n = 3).

Nutrient	compost	straw	woodchip
C (g/kg)	322.3 (0.433)	459.0 (1.012)	485.3 (1.121)
N (g/kg)	25.3 (0.167)	11.2 (0.083)	7.6 (0.105)
C:N	12.7 (0.084)	40.9 (0.368)	63.6 (0.760)
P (g/kg)	5.5 (0.076)	1.0 (0.025)	0.9 (0.024)
K (g/kg)	20.6 (0.31)	13.1 (0.22)	5.1 (0.10)
Mg (g/kg)	4.3 (0.014)	0.7 (0.015)	1.3 (0.040)
Mn (g/kg)	258 (1.68)	<u>41 (1.15)</u>	<u>41 (1.67)</u>
Fe (g/kg)	15.0 (0.051)	0.5 (0.015)	1.0 (0.060

Table 2: Amount of C, N and other nutrients applied in each treatment (g/plot).

	straw	woodchip	compost	straw-compost	woodchip-compost
С	4645	5047	2707	8197	7754
N	114	79	213	347	292
C:N ratio	41	64	13	24	27
Р	11	9	46	59	55
K	133	53	173	330	226
Mg	7	14	37	45	50

675

#### Deleted: 4

Deleted: 3

Formatted Table

Deleted: ¶ Table 2: Baseline soil data for each treatment (SEM indicated in parentheses, n = 4). .... [1

... [1]

Table 👍: Statistical o	utputs of one-tailed T-	tests of non-additive	e effects. Significand	e of deviation f	rom additivity
(0) is indicated as $p <$	<b>0.05</b> and p < 0.1.				

I

Deleted: 5

	straw-comp	oost mixtu	re	woodchip-coi	npost mix	ture
	Mean	Т	р	Mean	Т	р
	% non-additivity			% non-additivity		
Yield	9.66	1.004	0.195	9.54	0.771	0.24
Available N	55.06	3.789	0.016	7.16	0.235	0.41
Mineralisable N	39.67	1.265	0.147	8.93	0.990	0.19
Р	3.01	0.226	0.417	-8.60	-0.788	0.75
K	-0.79	-0.082	0.530	-0.86	-0.171	0.56
Mg	9.95	1.475	0.118	2.73	0.335	0.38
SOM	13.10	4.022	0.014	6.73	0.954	0.20
pН	3.04	2.006	0.931	2.41	1.118	0.82
Respiration	5.12	0.300	0.392	16.41	1.023	0.19
Bulk density	-7.80	-2.232	0.056	-3.73	-0.919	0.21
Aggregate stability	11.41	1.555	0.109	8.57	1.291	0.14

**Deleted: 1** Table 6: Qualitative assessment of lettuce plants as the % of lettuce heads per plot affected by each condition. "Overall" quality impairment is the % of lettuce head per plot affected by one or more conditions. Mean values per treatment (n = 4; SEM in parentheses). .... [2] ... [2]

	Yield	Av N	Av+Min N	Р	K	Mg	SOM	Resp
App_C	-0.10	0.17	0.17	-0.07	0.40	0.22	0.00	-0.01
App_N	0.07	0.26	0.30	0.08	0.54	0.32	0.06	-0.09
App_P	0.00	0.20	0.23	0.00	0.49	0.22	-0.01	-0.17
App_K	0.17	0.30	0.36	0.17	0.56	0.39	0.12	-0.05
App_Mg	0.19	0.33	0.38	0.19	0.56	0.45	0.16	0.02
Yield	-	0.29	0.45	0.75	0.19	0.78	0.74	0.36
Av N	0.29	-	0.91	0.42	0.27	0.55	0.58	0.36
Av + Min N	0.45	0.91	-	0.49	0.35	0.61	0.65	0.30
Р	0.75	0.42	0.49	-	0.02	0.83	0.86	0.51
K	0.19	0.27	0.35	0.02	-	0.35	0.02	-0.26
Mg	0.78	0.55	0.61	0.83	0.35	-	0.80	0.47
SOM	0.74	0.58	0.65	0.86	0.02	0.80	-	0.62
Agg stab	0.45	0.55	0.48	0.36	0.00	0.58	0.58	0.41
Overall qual.	0.20	0.51	0.46	0.10	0.34	0.23	0.21	0.14

## S1. Supporting information for blocking structure

5

Based on the strong and consistent gradient we observed in %C content of the soils (Figure S1a) and a similar gradient for the %N content of the soils (Figure S1b), we applied a retrospective blocking structure to enable a more accurate assessment of non-additive effects. The plots with the highest %C content for each treatment were grouped into one block, the plots with the second highest %C content for each treatment were grouped into another block, etc. (Figure S1c).

(a) 2.18 2.42 2.59 2.73 3.29 3.51 3.54 3.42 2.21 2.28 2.45 2.65 2.89 3.22 3.29 3.36 2.10 2.24 2.29 2.73 2.90 2.86 3.13 3.16 0.31 0.26 0.31 (b) 0.21 0.23 0.24 0.30 0.31 0.22 0.25 0.29 0.22 0.23 0.27 0.29 0.30 0.22 0.22 0.22 0.25 0.27 0.26 0.28 0.28 (c) 3 4 4 2 2 3 3 1

Figure S1: The gradient in (a) %C and (b) %N observed in the plots, and (c) the retrospective blocking structure we applied, where each box represents a plot, numbers = blocks; and colours = treatments (grey = control, yellow = straw, beige = woodchip, purple = compost, light brown = straw-compost, chestnut brown = woodchip.compost).

1

## S2. Baseline soil data

Table S1: Baseline soil data for each treatment (SEM indicated in parentheses, n = 4).

	Soil LOI	Soil C:N	<u>Soil pH</u>	Clay content
	<u>(%)</u>			<u>(%)</u>
<u>compost</u>	7.94 (0.45)	<u>10.77 (0.25)</u>	<u>8.30 (0.03)</u>	23.3 (0.75)
straw	<u>6.84 (0.03)</u>	<u>10.45 (0.09)</u>	<u>8.30 (0.05)</u>	23.8 (1.18)
straw-compost	7.78 (0.51)	<u>10.76 (0.20)</u>	8.32 (0.04)	26.0 (1.08)
woodchip	8.03 (0.51)	10.64 (0.39)	8.27 (0.03)	24.3 (0.75)
woodchip-compost	8.29 (0.47)	<u>10.95 (0.22)</u>	8.32 (0.03)	26.5 (1.26)
<u>control</u>	8.14 (0.32)	<u>10.79 (0.13)</u>	8.21 (0.02)	24.5 (1.50)

10 S3. Additional per-treatment results



Figure S2: Soil respiration measured by the Figure S3: Soil pH after different treatments. Solvita  $CO_2$ -burst method. Lower and upper Lower and upper hinges correspond to the 25th and 75th and 75th percentiles; black dots represent individual data points, occasionally overlapping (n = 4). (n = 4).



Figure S4: Earthworm abundance per plot after different treatments. Lower and upper hinges correspond to the 25th and 75th percentiles; black dots represent individual data points, occasionally overlapping (n = 4).

Table S2: Qualitative assessment of lettuce plants as the % of lettuce heads per plot affected by each condition. "Overall" quality impairment is the % of lettuce head per plot affected by one or more conditions. Mean values per treatment (n = 4; SEM in parentheses).

Treatment	Chlo	rosis	<u>Tip burn</u>	Rot	Overall
	<u>(All)</u>	(Tips only)			
<u>control</u>	<u>49.1 (16.1)</u>	<u>47.3 (16.9)</u>	<u>15.5 (4.9)</u>	<u>1.7 (1.1)</u>	<u>77.8 (12.3)</u>
straw	<u>31.5 (11.7)</u>	<u>21.7 (7.49)</u>	<u>1.9 (1.3)</u>	<u>0.0 (0.0)</u>	<u>43.1 (15.6)</u>
woodchip	<u>39.3 (9.3)</u>	<u>33.4 (8.5)</u>	12.2 (4.5)	<u>4.3 (2.0)</u>	<u>80.4 (11.6)</u>
<u>compost</u>	40.4 (7.7)	<u>34.3 (6.6)</u>	<u>14.5 (9.0)</u>	<u>0.6 (0.6)</u>	<u>69.4 (10.7)</u>
straw-compost	58.3 (14.9)	56.1 (15.2)	16.9 (8.6)	0.7 (0.7)	93.0 (7.0)
woodchip-compost	<u>61.7 (14.0)</u>	<u>54.1 (16.7)</u>	<u>18.0 (8.7)</u>	<u>0.0 (0.0)</u>	<u>82.2 (11.0)</u>

S4. Calculation of amount of nutrients added to the soil via residue mixtures Deleted: Page Break----Deleted: 3 15 First the mass of nutrients applied per plot was calculated, using the application rate of each residue (kg residue/plot) and the amount of each nutrient in the residues (mg nutrient/kg residue): Nutrients applied (mg plot<sup>-1</sup>) = Reisude applied to each plot (kg plot<sup>-1</sup>) × nutrient content of reisdues (mg kg<sup>-1</sup>) Then, using the plot volume to 20 cm depth 0.2 m  $\times$  6 m  $\times$  2 m = 2.4 m<sup>3</sup>/plot) and the bulk density (g/m<sup>3</sup>), assuming the bulk density is representative of the top 20 cm and assuming nutrients from the residues applied 20 remained in the top 20 cm of the soil (the sampling depth), we calculated the amount of nutrients added per g of soil via the residues (mg nutrients/g soil) as: Nutrient application rate (mg  $g^{-1}$ ) =  $\frac{Nutrients applied (mg plot^{-1})}{Area of the plot (m^3)}$ /Bulk density (g  $m^{-3}$ ) Then the difference between the amount of nutrients measured in each plot and the average amount of nutrients measured in the control plots was calculated as: 25 Nutrient increase (mg  $g^{-1}$ ) = Nutrients in treatment plot (mg  $g^{-1}$ ) – nutreints in control plot (mg  $g^{-1}$ ) Then we determined the increase in soil available nutrients (relative to control) as a proportion of the amount of nutrients added to the soil via residue amendments:  $\label{eq:linear} \textit{Increase in available soil nutrients (\%)} = \frac{\textit{Nutrient increase (mg \ g^{-1})}}{\textit{Nutrient application rate (mg \ g^{-1})}} \times 100\%$ Table S2: Increase in soil available nutrients (relative to control treatment) as a proportion (%) of the quantity of Deleted: 1 nutrients added to the soil (assuming nutrients added via residues remained in the top 20 cm of the soil that was sampled). Numbers in bold are significantly different (p < 0.05) from 0 (SEM indicated in parentheses).

	straw	woodchip	compost	straw-compost	woodchip-compost
Р	-95 (4)	15(65)	12 (15)	-2 (10)	1 (7)
Κ	10 (37)	57 (68)	53 (12)	31 (3)	49 (13)
Mg	-242 (42)	38 (74)	25 (30)	15 (25)	35 (10)
Ν	-19 (5)	-3 (5)	-2 (4)	2 (3)	1 (3)

## 35 S5. Statistical outputs

# Deleted: 4

## Table S4: Statistical outputs of baseline soil properties. Significance indicated as p < 0.05 and p < 0.1.

Variable	two-way ANO	VA (per factor)	Lev	/ene	Shapi	ro-Wilk
	(residue; compost	; residue*compost)			of re	siduals
	F	р	F	р	W	р
SOM (LOI)	2.433; 0.914; 0.938	0.116; 0.352; 0.410	2.092	0.114	0.966	0.578
Soil moisture	0.843; 2.425; 0.315	0.447; 0.137; 0.733	2.911	0.043	0.965	0.536
pН	1.142; 3.241; 0.345	0.341; 0.089; 0.713	0.881	0.513	0.932	0.108
C:N	0.427; 1.094; 0.328	0.659; 0.310; 0.725	0.809	0.558	0.948	0.244
Variable	one-way ANOV	A (per treatment)	Lev	/ene	Shapiro-Wilk	
					of re	siduals
	F	р	F	р	W	р
SOM (LOI)	1.206	0.350	1.727	0.175	0.966	0.574
Soil moisture	1.067	0.420	1.598	0.208	0.947	0.228
pН	1.382	0.278	0.735	0.628	0.950	0.275

Table S5: Statistical outputs of per-treatment results. Significance indicated as p < 0.05 and p < 0.1.

Variable	two-way ANOVA (per factor)		Levene		Shapiro-Wilk	
	(residues; compost; residues*compost)				of residuals	
	F	р	F	р	W	р
Available N	0.509; 2.566; 1.930	0.609; 0.127; 0.174	1.871	0.150	0.950	0.273
Mineralisable N	0.504; 2.936; 0.797	0.612; 0.104; 0.466	1.508	0.237	0.981	0.909
Mineralisable:Available	0.372; 0.597; 0.204	0.695; 0.450; 0.818	0.656	0.661	0.973	0.759
Available+Mineralisable	0.680; 3.877; 1.895	0.519; <b>0.065</b> ; 0.179	1.313	0.303	0.958	0.391
Total biomass	1.625; 1.306; 0.303	0.225; 0.268; 0.742	0.883	0.513	0.971	0.697
CO <sub>2</sub> Burst	2.289; 0.033; 1.091	0.130; 0.859; 0.357	0.323	0.893	0.906	0.029
Earthworm abundance	0.136; 1.221; 1.945	0.874; 0.284; 0.172	0.449	0.809	0.956	0.361
P (mg/g soil)	1.547; 1.214; 0.440	0.240; 0.285; 0.651	1.300	0.308	0.967	0.586
K (mg/g soil)	0.291; 7.761; 0.009	0.751; 0.012; 0.991	2.369	0.081	0.987	0.918
Mg (mg/g soil)	2.067; 4.953; 0.450	0.156; <b>0.039</b> ; 0.645	2.573	0.063	0.960	0.437
SOM (LOI)	1.219; 0.574; 0.945	0.319; 0.458; 0.407	1.434	0.260	0.954	0.331
pН	1.459; 1.459; 3.405	0.259; 0.243; <b>0.056</b>	1.600	0.211	0.902	0.024
Bulk density	3.283; 1.269; 0.994	0.062; 0.276; 0.391	1.214	0.345	0.966	0.589
Aggregate stability	0.836; 0.022; 0.646	0.449; 0.883; 0.536	0.685	0.641	0.955	0.342
Quality impairment	0.653; 2.294; 3.568	0.532; 0.147; <b>0.050</b>	0.466	0.796	0.946	0.233