

Assessing soil redistribution of forest and cropland sites in wet tropical Africa using $^{239+240}\text{Pu}$ fallout radionuclides

5 Florian Wilken^{1,2}, Peter Fiener², Michael Ketterer³, Katrin Meusburger⁴, Daniel Iragi Muhindo⁵, Kristof van Oost⁶, Sebastian Doetterl¹

¹Department of Environmental Systems Science, Eidgenössische Technische Hochschule Zürich, Zürich, Switzerland

²Institute for Geography, Universität Augsburg, Augsburg, Germany

³Chemistry and Biochemistry, Northern Arizona University, Flagstaff, USA

⁴Swiss Federal Institute for Forest, Snow and Landscape Research, Birmensdorf, Switzerland

10 ⁵Faculty of Agronomy, Université Catholique de Bukavu, Bukavu, DR Congo

⁶Earth and Life Institute, Université Catholique de Louvain, Louvain-la-Neuve, Belgium

Correspondence to: Florian Wilken (florian.wilken@usys.ethz.ch)

15 **Abstract**

Due to the rapidly growing population in tropical Africa, a substantial rise in food demand is predicted in upcoming decades, which will result in higher pressure on soil resources. However, there is limited knowledge on soil redistribution dynamics following land conversion into arable land in tropical Africa that is partly caused by infrastructure limitations for long-term landscape scale monitoring. In this study, fallout radionuclides $^{239+240}\text{Pu}$ are used to assess soil redistribution along topographic gradients at two cropland sites and at three nearby pristine forest sites located in the DR Congo, Uganda and Rwanda. In the study area, a $^{239+240}\text{Pu}$ baseline inventory is found that is higher than typically expected for tropical regions (mean forest inventory 41 Bq m⁻²). Pristine forests show no indication for soil redistribution based on $^{239+240}\text{Pu}$ along topographical gradients. In contrast, soil erosion and sedimentation on cropland reached up to 37 cm (81 Mg ha⁻¹ yr⁻¹) and 40 cm (87 Mg ha⁻¹ yr⁻¹) within the last 55 years, respectively. Cropland sites show high intra-slope variability with locations showing severe soil erosion located in direct proximity to sedimentation sites. This study shows the applicability of a valuable method to assess tropical soil redistribution and provides insight into soil degradation rates and patterns in one of the most socio-economically and ecologically vulnerable regions of the world.

20
25

1. Introduction

30 Soil erosion is considered to be the major threat to global soil resources and substantially contributes to crop yield reduction (Amundson et al., 2015; Montanarella et al., 2016; Govers et al., 2017), which challenges food security in regions facing population growth beyond sustainable limits in the 21st century. In particular, the White Nile-Congo rift (NiCo) region faces a strong impact of soil redistribution (Lewis and Nyamulinda, 1996; FAO and ITPS, 2015; Montanarella et al., 2016) and corresponding sediment delivery (Vanmaercke et al., 2014) due to steep terrain, high rainfall erosivity with a strong intra-
35 annual seasonality (Fick and Hijmans, 2017) that causes sparse vegetation cover of soil at the end of the dry seasons but also throughout the cultivation period (Lewis and Nyamulinda, 1996; Fick and Hijmans, 2017) due to non-mechanised farming. The region is also predicted to undergo substantial climate change, which might further increase soil erosion (Borrelli et al., 2020). The loss of soil resources and crop yield decline in the NiCo region happens in parallel to a rapid population growth (population of Rwanda, Uganda and Democratic Republic of Congo 2020: 150 millions – predicted 2100: 430 millions; WPR,
40 2020), which drives rising food demands that are expected to triple for the entire Sub-Saharan Africa between 2010 and 2050 (van Ittersum et al., 2016). The associated pressure on land resources leads to various problems that will have dramatic ecological and social impacts (food insecurity, political unrest, migration) in the NiCo region (Chamberlin et al., 2014; FAO and ITPS, 2015). Under current practices, an increasing demand in food is typically compensated through deforestation to assess new soil resources, which are often located in areas with steep slopes (Govers et al., 2017). This causes a loss of highly
45 valuable forest ecosystem services (e.g. carbon storage, biodiversity, imbalance of the hydrological cycle) and the onset of potential soil erosion at previously undisturbed sites (Nyssen et al., 2004) and often unsustainable use of soil resources (Wynants et al., 2019). In steep cropland sites of the NiCo region, it is frequently observed that the entire deep tropical saprolite body is removed and the bedrock is exposed at the surface, which means a quasi-permanent loss of cropland and the option to reforest areas for decadal to centennial timescales (Evans et al., 2020). A pressing need persists to predict future dynamics and
50 timescales of cropland degradation in order to understand the pace of a rising food shortage and to develop adapted agricultural management strategies at suitable locations. Smart intensification of existing cropland soils due to adapted agricultural practices in suitable locations and the conservation/restoration of soils prone to erosion (e.g. reforestation or grassland use) has been discussed as a key management strategy to combat degradation (Govers et al., 2017). To develop a smart intensification plan detailed information on soil degradation dynamics of individual regions under specific conditions is
55 essential (e.g. land use, topography, soil type, rainfall characteristics). Soil erosion plot experiments were carried out in tropical Africa (Lewis and Nyamulinda, 1996; Xiong et al., 2019) to understand the rates of soil loss. However, plot experiments are limited to soil erosion processes, while soil redistribution dynamics (interaction of erosion and sedimentation) remain unexplained, although they are crucial to understand soil degradation rates on a landscape-scale. However, catchment monitoring that provides insight into spatially distributed soil redistribution dynamics necessitates a sufficiently long time
60 series (years to decades) to integrate a statistically representative variety of erosive rainfall events impacting different land cover conditions (Fiener et al., 2019). Particularly in regions of limited infrastructure, long-term catchment monitoring projects

are challenging and typically rare. This problem can be overcome by the analysis of fallout radionuclides from nuclear weapon tests (i.e. ^{137}Cs , $^{239+240}\text{Pu}$) as soil redistribution tracers (Meusburger et al., 2016; Alewell et al., 2017; Evrard et al., 2020), which have the major advantage to provide insight into spatial patterns of both soil erosion and sedimentation processes integrated over the period since 1963-1964 (Test Ban Treaty that caused a rapid decrease of atmospheric bomb tests; Wallbrink and Murray, 1993; Evrard et al., 2020). The most widely used fallout radioisotope in soil redistribution studies is ^{137}Cs (e.g. Porto and Walling, 2012; Chartin et al., 2013; Evrard et al., 2020), which has a rather short half-life of about 30 yrs. Hence, decay has already led to a pronounced reduction (73% in 2020) of the ^{137}Cs activity until today (Alewell et al., 2017). In tropical and equatorial regions, this is a critical limitation of using ^{137}Cs for soil redistribution analysis due to much lower fallout compared to the mid latitudes of the northern hemisphere (Hardy et al., 1973; Evrard et al., 2020). Furthermore, extreme soil erosion rates in the tropics (Lewis and Nyamulinda, 1996; Angima et al., 2003; Nyesheja et al., 2019; Xiong et al., 2019) are additionally driving a depletion of the ^{137}Cs inventories. To overcome these analytical difficulties, the fallout radionuclides ^{239}Pu and ^{240}Pu have been discussed and tested as an alternative radioisotopic tracer to ^{137}Cs for soil redistribution studies over the past decade. The major advantage of both isotopes is their long half-life ($^{239}\text{Pu} = 24,110$ yrs, $^{240}\text{Pu} = 6,561$ yrs). Furthermore, ^{239}Pu and ^{240}Pu show a very limited plant uptake (Akleyev et al., 2000) and preferentially form associations with soil iron oxides (Ryan et al., 1998; Lal et al., 2013), which potentially makes the isotopes very suitable tracers for tropical environments dominated by Ferralsols. Hence, the $^{239+240}\text{Pu}$ activity in tropical soils might be high enough to successfully carry out soil redistribution studies and provide important insights on soil redistribution dynamics in tropical Africa. Few fallout radionuclide based soil redistribution studies have been carried out in the tropics (Evrard et al., 2020) and, to the best of our best knowledge, non was conducted in the wet tropics of Africa (Af, Am climate; Kottek et al., 2006). The aims of the study are (i) to test the suitability of fallout radionuclides ^{239}Pu and ^{240}Pu as tracers of soil redistribution in the wet tropics of Africa, and (ii) to analyse soil redistribution dynamics after conversion from forest into cropland for selected sites within the East African NiCo region.

2. Methods

2.1 Study sites and sampling design

The NiCo region is located in the headwater catchments of the White Nile (Lake Edward) and the Congo River (Lake Kivu) that are part of the East African Rift Valley system (Fig. 1). The region faces rapid population growth creating substantial pressure on land resources and initiating conversion of forest into cropland. Soil degradation by water erosion is a recognised problem in the region (Lewis and Nyamulinda, 1996; Montanarella et al., 2016) indicated via frequent soil erosion events resulting in ephemeral rills and gullies as well as permanent deep gully systems. The region is characterised by steep terrain (Fig. 1; Tab. 1) and tropical climate with a mean annual air temperature between 16.7 and 19.3°C and an annual rainfall ranging between 1300 and 1900 mm (time period 1970-2000; Fick and Hijmans, 2017). The seasonal rainfall distribution is subdivided into two cycles of wet and dry seasons (Fick and Hijmans, 2017). The rainfall erosivity in the region is high due to frequently

95 occurring storm events with large rainfall amounts linked to high rainfall intensities during the rain seasons (on average 20
erosive rainfall events per rain season; events exceeding 10 mm h⁻¹ of rainfall per 30 min interval)(Doetterl et al., 2021b). Soils
in the region are deeply weathered Ferralsols (> 6 m; WRB, 2006; Doetterl et al., 2021a) developed from three geochemically
distinct parent materials (DR Congo: mafic magmatic rocks; Uganda: felsic magmatic rocks; Rwanda: sedimentary rock of
mixed geochemical composition). Soils across the study area are typically classified as clay loam while at the Ugandan forest
and cropland study sites a lower clay and higher sand content is found (Doetterl et al., 2021a).

100 The forest sites in the study area are primary tropical mountain forests (for detailed information see Doetterl et al., 2021a).
Farming is documented since the 1950s for the site in DR Congo, while conversion into cropland at the Ugandan site took
place during the 1970s (personal communication with local villagers). The cropland sites represent the typical smallholder
farming found in the region, which is based on small non-terraced fields with non-mechanised tillage practices. Due to the
small fields (mean field size = 450 m²) and an individual and dynamic field management, a high patchiness of soil cover
105 conditions is present and can alternate between bare soil and fully grown vegetation cover in direct proximity to each other
(Fig. A1).

In 2018, a soil sampling campaign was carried out in three pristine forests and two cropland sites in the NiCo region in order
to collect soil samples for a soil redistribution assessment based on the fallout radionuclides ²³⁹Pu and ²⁴⁰Pu. As part of this
110 campaign, a total of 347 samples were taken. Soil sampling was carried out using a manual closed tube soil corer (VSI soil
core sampler, Vienna-Scientific, Austria) with a diameter of 6.8 cm and a length of 120 cm.

Sampling sites in forests were located within the Kahuzi Biega (DR Congo), Kibale (Uganda) and Nyungwe Forest (Rwanda)
National Parks (Fig. 1). There, the sampling scheme is aligned to a toposequence approach that covers three different landscape
positions situated along a catena: plateau, slope and foot-slope. Sampling was carried out at 19 locations within each catena
and covers different soil layers (L-horizon, O-horizon and mineral soil; see Tab. 2). Discrete sampling at plateau locations
115 took place in order to quantify spatial variation in reference fallout inventories. To average out the typical variability of fallout
radionuclide inventories, composite samples were taken at the slope and foot-slope positions. In addition to mineral soil layers,
two organic layers at two different levels of decomposition (L and O horizons) were collected over an area of 20 cm x 20 cm
at each sampling site. About 40% of the total forest samples were taken at plateau positions (high proportion due to non-
composite sampling), while 30% of the samples were taken at slope and foot-slope locations, respectively. At foot-slope
120 locations soils were additionally sampled from 60 cm to 120 cm soil depth to cover colluvial sites with ²³⁹⁺²⁴⁰Pu activity in
deeper soil layers.

Sampling sites in cropland were placed along two catenae within DR Congo and in Uganda, covering 51 individual locations
at each study site (Tab. 2). The majority (~50%) of sampling sites are distributed along slope positions (12-13° steepness in
125 both cropland sites) while 25% of sites are located at foot-slope and plateau sites. To understand the depth distribution of
²³⁹⁺²⁴⁰Pu and the variability of radionuclide inventories under stable geomorphic conditions, three depth increments of 20 cm
thickness were taken to a total soil depth of 60 cm at plateau sites. In DR Congo, a cropland plateau site (converted to grassland

approximately in 2005) located about 8 km apart from the study site was sampled, while in Uganda, flat plateau sites under arable use in direct proximity to the study site were sampled. At slope locations, a single soil increment down to 60 cm was taken. At the foot-slope locations, to cover potential sedimentation, an additional increment from 60 to 100 cm was sampled to assure full cover of the radionuclide inventory.

2.3 ²³⁹⁺²⁴⁰Pu measurements

An assessment of fallout radionuclides ²³⁹⁺²⁴⁰Pu inventories was used to estimate effective soil redistribution since the 1960s along the investigated geomorphic transects. Plutonium isotopes measurements were conducted following Calitri et al. (2019) and Ketterer et al. (2004). The chemical preparation consisted of the following sequence:

1. Soil material was milled and subsequently dry-ashed for at least eight hours at 600°C to remove organic matter. An aliquot of the dry-ashed material of up to 50 grams was weighed into a 250 mL polypropylene bottle. Samples were tested for excessive reaction of carbonates by addition of 5 mL of 8 M HNO₃.
2. Samples were spiked using 7 picograms (~ 1 mBq) of a ²⁴²Pu tracer (NIST 4334g), in the form of a solution in 4 M aqueous HNO₃.
3. 125 mL of 8 M aqueous HNO₃ were added, with caution being exercised to add acid slowly when carbonates were present.
4. The sample vessels were capped and heated at 80°C overnight (~ 16 hours) with occasional mixing.
5. Following heating, the sample leach solutions were recovered by filtration with 0.45 micron cellulose nitrate membranes.
6. The plutonium was converted to the +4 oxidation state via addition of 0.5 g FeSO₄*7H₂O dissolved in 2 mL water, followed by 2 grams of NaNO₂ dissolved in 5 mL water. Thereafter, the solutions were heated uncapped in a 90° C convection oven for 1.5 hours to release evolved NO₂ (g) and allow for conversion of the Pu into Pu(IV).
7. 300 milligrams of Pu-selective resin TEVA (EiChrom, Lisle, IL, USA) was added to the sample solution; the mixtures were agitated over a 4-hour timeframe to allow for the resin to uptake the Pu(IV).
8. The TEVA resin was collected on a 20 mL polyethylene column equipped with a glass wool plug; the pass-through solution was drained and discarded. The columns were rinsed with the following sequence: i) 50 mL of 2 M aqueous HNO₃; ii) 20 mL of 9 M aqueous HCl; and iii) 10 mL of 2 M aqueous HNO₃. The rinse sequence removes matrix elements, uranium and thorium.
9. Plutonium was eluted from the columns using the following sequence: i) 0.5 mL water; ii) 0.5 mL of 0.05 M aqueous ammonium oxalate; and iii) 0.5 mL water, all of which were collected together for analysis directly after elution.

In the preparations, quality control samples were included to assess the results; these consisted of blanks (35 g powdered sandstone devoid of detectable Pu). Blanks consisting of 35 g sandstone spiked with small quantities (50-100 milligrams) of IAEA 384 (Fangataufu Sediment) were also prepared. Eleven measurements of IAEA 384 resulted in an average of 102 Bq kg⁻¹ ²³⁹⁺²⁴⁰Pu, and a standard deviation of 10 Bq kg⁻¹; these results compare well to the reference value of 107 Bq kg⁻¹ ²³⁹⁺²⁴⁰Pu. The blanks were used to determine a detection limit of 0.01 Bq kg⁻¹ ²³⁹⁺²⁴⁰Pu.

160 Sample Pu fractions were measured with a Thermo X Series II quadrupole ICP-MS (Bremen, Germany) equipped with an
 APEX HF high-efficiency sample introduction system (ESI Scientific, Omaha, NE, USA). The APEX is equipped with a self-
 aspirating concentric fluorinated ethylene-propylene nebulizer operating at an uptake rate of ~ 0.15 mL per minute. The
 instrument is located at Northern Arizona University; the laboratory is licensed with the State of Arizona for handling ^{242}Pu
 spike solutions. The intensities of ^{235}U , ^{239}Pu , ^{240}Pu and ^{242}Pu were recorded using a peak-jump algorithm (10 ms dwell time,
 165 1000 sweeps/integration, three integrations per sample). The ^{235}U isotope was measured as a proxy for ^{238}U , the latter whose
 intensity occasionally exceeded the linear range of the ICPMS's pulse-counting detector; this was done in order to assess the
 potential interference of $^{238}\text{U}^1\text{H}^+$ on ^{239}Pu . As many samples exhibited relatively high levels of $^{238}\text{U}^1\text{H}^+$, generating a potential
 "false-positive" detection of ^{239}Pu , it was determined to be advantageous to measure Pu using the ^{240}Pu isotope, which is
 unaffected by uranium hydride species. The measured atom ratio $^{240}\text{Pu}/^{242}\text{Pu}$ was converted into a mass of ^{240}Pu detected, using
 170 the known mass of ^{242}Pu ; the ^{240}Pu activity was calculated, and converted into the corresponding $^{239+240}\text{Pu}$ activity, based on
 the known $^{240}\text{Pu}/^{239}\text{Pu}$ atom and activity ratios in stratospheric fallout Pu (Kelley et al., 1999). The method resulted in a
 detection limit of $0.01 \text{ Bq kg}^{-1} \text{ }^{239+240}\text{Pu}$, which equates to a soil inventory of ca. $5\text{-}8 \text{ Bq m}^{-2} \text{ }^{239+240}\text{Pu}$ for a bulk density of ~ 0.8
 to $\sim 1.3 \text{ Mg m}^{-3}$.

2.4 Cropland soil redistribution calculation

175 To derive topographic change and corresponding soil redistribution using $^{239+240}\text{Pu}$ inventories, a mass balance model (Zhang
 et al., 2019) was applied integrating soil erosion and sedimentation over the period 1964 to 2018. Soil erosion and
 sedimentation processes were implemented (R-Core-Team, 2019) individually to account for differences of both processes.
 Due to topsoil loss by soil erosion, former soil from deeper layers with negligible low $^{239+240}\text{Pu}$ activity gets increasingly
 incorporated into the plough layer. This exponential decay of the $^{239+240}\text{Pu}$ inventory is mainly controlled by the soil erosion
 180 magnitude and frequency and plough depth. These processes are addressed by the mass balance model that simulates the
 $^{239+240}\text{Pu}$ inventory reduction on a year by year basis over the simulation period:

$$A_{yr} = \int_{1964}^{yr} A_{yr-1963} \left(1 - \frac{R_{yr}}{d} \right)$$

where yr is the simulation year, A_{yr} is the $^{239+240}\text{Pu}$ inventory at the specific simulation year in Bq m^{-2} , $A_{yr-1963}$ is the annually
 updated $^{239+240}\text{Pu}$ inventory in Bq m^{-2} , d is the average plough depth in m, R_{yr} is soil erosion of simulation year in m. Based
 185 on these results, an individual logarithmic function between A and R for each study site was fitted that can be used to derive
 the amount of soil loss in cm (55 yrs.)⁻¹.

Sedimentation is represented as a linear increase of the inventory:

$$R = -d \left(1 - \frac{A}{A_{ref}} \right)$$

where R is the soil redistribution rate in m (55 yrs)⁻¹, A_{ref} is the ²³⁹⁺²⁴⁰Pu reference inventory and A is the ²³⁹⁺²⁴⁰Pu activity difference (reference vs. local activity) in m x m (55 yrs)⁻¹ and Bq kg⁻¹ x Bq kg⁻¹ (55 yrs)⁻¹ over the simulation period (more details on the A_{ref} implementation are given in section 2.5).

Finally, the ²³⁹⁺²⁴⁰Pu activity (Bq m⁻²) was converted from topographic change (in m) into soil redistribution rates Y in Mg ha⁻¹ (55 yrs)⁻¹ as follows (Walling et al., 2011):

$$Y = 10 d B \left(1 - \frac{R}{d} \right)$$

where B is the bulk density in kg m⁻³. B is measured for each mineral and O-horizon sample, while B for the L layer was taken from the literature, i.e. 80 kg m⁻³ (Wilcke et al., 2002).

2.5 Cropland scenario assessment using a mass balance model

The mass balance soil mixing model was used to assess different scenario assumptions and their sensitivity. First, different ²³⁹⁺²⁴⁰Pu reference inventories were determined in two ways: (i) the mean ²³⁹⁺²⁴⁰Pu inventory of all forest sites of a specific region (Ref_{for}; i.e. mean inventory of Kahuzi Biega forest for the DR Congo cropland sites and Kibale forest for the Ugandan cropland sites) and (ii) the mean ²³⁹⁺²⁴⁰Pu inventory of the cropland plateau sites of the specific region (Ref_{plt}). These land use specific reference scenarios are supposed to address potential differences between Ref_{for} and Ref_{plt} to understand uncertainties associated with the reference determination. Second, the sensitivity of the ploughing and corresponding mixing depth is assessed using a 20±5 cm ploughing depth deviation as the exact plough depth over the integration period cannot be accurately derived. Third, to address potential interannual variability of water erosion, a scenario with five extreme years producing the same total soil erosion as a 55 years continuous soil erosion rate was compared against the results of the first scenario.

3. Results

Forest sites ²³⁹⁺²⁴⁰Pu activities and inventories

The mean inventory (i.e. sum of L, O and mineral horizons ²³⁹⁺²⁴⁰Pu activities) of all forest sampling sites is 41.3 Bq m⁻², whereas the measured ²³⁹⁺²⁴⁰Pu activity in the Kahuzi Biega forest is somewhat smaller compared to the Kibale and Nyungwe forests (Fig. 2; Kahuzi Biega: 32.7 Bq m⁻², Kibale: 42.9 Bq m⁻²; Nyungwe: 48.4 Bq m⁻²). The majority of L horizon samples fall below the ²³⁹⁺²⁴⁰Pu detection limit (0.01 Bq kg⁻¹; 40 out of 55 samples). O horizon samples show distinctively higher ²³⁹⁺²⁴⁰Pu activities (Fig. 3). However, the contribution of the O horizons to the ²³⁹⁺²⁴⁰Pu inventory of the soil profile is small (mean: 1.2%), because of low bulk density (approx. 0.2 Mg m⁻³) and O horizon thickness (mean 5 cm). Few (13%, 7 of 55 samples) of ²³⁹⁺²⁴⁰Pu activities of the 0 to 60 cm depth mineral soil layers fall below the detection limit, while the ²³⁹⁺²⁴⁰Pu activities of samples within the detectable range are at least three times higher than the detection limit (Fig. 2). In contrast, almost no activity is detected in the deeper soil layer from 60 to 120 cm.

Comparing the $^{239+240}\text{Pu}$ activities at different topographic positions does not result in a consistent $^{239+240}\text{Pu}$ activity to topography relation (Fig. 3). While the foot-slopes in Rwanda show the highest $^{239+240}\text{Pu}$ activities, the opposite is the case for foot-slopes in Uganda and DR Congo. At the plateau sites in Uganda and Rwanda, a lower $^{239+240}\text{Pu}$ activity compared to the slope sites is found.

A large $^{239+240}\text{Pu}$ activity standard deviation of the forest sites was found for all topographic positions (Fig. 3). The mean $^{239+240}\text{Pu}$ inventories found at the slope and foot-slope sites fall within the range of one standard deviation \pm mean $^{239+240}\text{Pu}$ inventory of the corresponding plateau sites. The only exception is the foot-slope in the Ugandan forest that falls in range by two standard deviations of the plateau mean.

Cropland sites $^{239+240}\text{Pu}$ activities and inventories

At both cropland study sites, a distinctively lower $^{239+240}\text{Pu}$ activity relative to the forest sites is found. The lowest activity of $^{239+240}\text{Pu}$ is found at slope positions in DR Congo where 50% (n = 16) of sampling sites fall below the detection limit. The measurable slope samples show a mean and standard deviation of $0.019 \pm < 0.01 \text{ Bq kg}^{-1}$. A pronounced increase of the $^{239+240}\text{Pu}$ activities can be observed at foot-slope positions with activity also detectable in the sampled 60-100 cm deeper soil layer (Fig. 3). Hence, the $^{239+240}\text{Pu}$ activity at the DR Congo cropland site follows a topography related spatial pattern from low activities at slope to elevated activities at foot-slope positions.

In comparison to the cropland study site in DR Congo, the activities at the Ugandan cropland site are much higher (mean $^{239+240}\text{Pu}$ activity at slope sites DR Congo: 0.012 Bq kg^{-1} , Uganda: 0.046 Bq kg^{-1}) and do rarely (3 of 44 samples) fall below the detection limit. Variability of $^{239+240}\text{Pu}$ activities at Ugandan site is extremely high and shows for slope positions a coefficient of variation of 76%. In contrast to DR Congo cropland, lower $^{239+240}\text{Pu}$ activities are found at Ugandan foot-slope sites compared to slope positions (Fig. 3). Furthermore, the Ugandan foot-slope positions showed almost no $^{239+240}\text{Pu}$ activity in the deeper soil layer of 60-100 cm soil depth (Fig. 3).

DR Congo plateau sites (Ref_{plt}), assumed to represent the preserved full inventory of the global fallout, show a substantially lower $^{239+240}\text{Pu}$ inventory compared to the about 30 km apart located forest sites (DR Congo Ref_{plt} : 8.0 Bq m^{-2} vs. mean forest Ref_{for} : 32.7 Bq m^{-2}). Similarly, Uganda cropland plateau sites $^{239+240}\text{Pu}$ inventory (24.4 Bq m^{-2}) are lower compared to nearby ($\sim 10 \text{ km}$) forest sites, but with a lower relative difference compared to their DR Congo counterparts (relative difference Ref_{for} vs. Ref_{plt} : Uganda 43%, DR Congo 75%). The $^{239+240}\text{Pu}$ activity below the 0 - 20 cm topsoil layer at the plateau sites in DR Congo show a sharp reduction of the $^{239+240}\text{Pu}$ activity in deeper soil layers (Fig. 3), while at the Ugandan cropland sites, soil layers down to 40 cm show significant $^{239+240}\text{Pu}$ activity.

Cropland soil erosion and sedimentation

An important piece of information that is provided by the erosion module of the mass balance model is the minimum quantity of soil loss that is required to cause a reduction of the $^{239+240}\text{Pu}$ inventory that falls below the detection limit after the model integration period of 55 yrs. The difference between the Ref_{for} and the Ref_{plt} $^{239+240}\text{Pu}$ baseline reference leads to substantial

differences in modelled erosion. The model indicates that at the DR Congo cropland sites, soil loss of at least 37 cm (55 yrs.)⁻¹ is necessary before the ²³⁹⁺²⁴⁰Pu activity falls below detection limit using Ref_{for}, while using Ref_{plt} at least 10 cm (55 yrs.)⁻¹ of soil loss must have taken place before the detection limit is reached. At the Ugandan cropland sites, a ²³⁹⁺²⁴⁰Pu inventory reduction to reduce activity below detection limit is found for 43 cm soil loss (55 yrs.)⁻¹ when applying Ref_{for} and 32 cm (55 yrs.)⁻¹ when applying Ref_{plt}, respectively (Fig. 4).

Also, a pronounced sensitivity of the mass balance model on the tillage depth parameter is found. A deviation from an assumed 20 cm plough depth of ±5 cm causes a change of the required soil loss until the detection limit is reached of about ±24%. Testing the low frequency but high magnitude soil erosion scenario (only 5 extreme erosion years within 55 years simulation period), showed that detection limit is reached already at 81% of soil loss compared to the continuous 55 years erosion rate. Hence, the sensitivity of the ±5 cm plough depth exceeds the impact of the erosion year frequency, even for this extreme scenario assumption in the NiCo ecosystem (approx. 20 erosive rainfall events per rain season).

The number of sampling sites that are considered to be subject to sedimentation is strongly controlled by the assumption on the applied reference (Ref_{for} and Ref_{plt}). When using Ref_{for}, 4 slope sites (DR Congo: 0, Uganda: 4) show sedimentation greater than 5 cm (55 yrs.)⁻¹. This number increased to 24 sites following Ref_{plt} (DR Congo: 9, Uganda: 15). For both reference scenarios, sedimentation at slope positions is much lower in DR Congo as compared to Ugandan cropland (Fig. 5). The foot-slope sites in DR Congo show distinctively higher ²³⁹⁺²⁴⁰Pu inventories compared to the slope sites (Fig. 5). However, the mean inventory of foot-slope positions is still lower than the Ref_{for} ²³⁹⁺²⁴⁰Pu inventory (28 vs. 32 Bq m⁻²), which would be interpreted as an indicator for weak soil erosion considering Ref_{for} for soil redistribution calculation. In contrast, if Ref_{plt} is used in the calculation, the same foot-slope positions would be interpreted as sites that received substantial sedimentation exceeding 40 cm (55 yrs.)⁻¹ (Fig. 5). The foot-slope sites in DR Congo show a pronounced ²³⁹⁺²⁴⁰Pu activity in many samples of deeper soil layers (60-100 cm), while no such deeper soil layer ²³⁹⁺²⁴⁰Pu activity is found at foot-slope sites of the Ugandan study site (Fig. 3). Simulated sedimentation that exceeds the site specific sampling depth minus the plough depth (i.e. 40 cm at the slope locations assuming a 20 cm plough depth) is rarely found for both reference assumptions (DR Congo Ref_{plt}: 1, Ref_{for}: 0; Uganda Ref_{plt}: 6, Ref_{for}: 2) and therefore suggests a limited impact of enrichment processes by selective transport of fine soil particles.

4. Discussion

Applicability of ²³⁹⁺²⁴⁰Pu as soil erosion tracer in tropical Africa

In this study, a ²³⁹⁺²⁴⁰Pu based soil redistribution analysis at three forest and two cropland sites in the NiCo region was carried out. It is shown that for the majority of samples the topsoil ²³⁹⁺²⁴⁰Pu activity is high enough to be successfully measured and provide insight on soil redistribution in tropical Africa over the past decades. To our knowledge, this is the first ²³⁹⁺²⁴⁰Pu based soil redistribution study in tropical Africa.

The $^{239+240}\text{Pu}$ inventories found in this study are much higher than expected based on the global fallout estimates reported by Kelley et al. (1999) and Hardy et al. (1973) (4.8 Bq m^{-2} and 11.1 Bq m^{-2} for 10°N and 10°S). It is well established that the initial deposition of $^{239+240}\text{Pu}$ fallout was strongly dependent upon latitude; the study of Kelley et al. (1999) compared ^{237}Np , ^{239}Pu and ^{240}Pu inventories from soil cores collected from undisturbed, worldwide locations. One data point (Muguga, Kenya) included in the results of Kelley et al. (1999) yields a $^{239+240}\text{Pu}$ inventory of 19.2 Bq m^{-2} , which is significantly lower than the inventories for soils in the mid-latitude regions of the Northern Hemisphere, e.g., Munich, Germany exhibits a $^{239+240}\text{Pu}$ inventory of 104 Bq m^{-2} . Although the results of Kelley et al. (1999) are drawn from a limited set of samples, they nevertheless make the case for a strong dependency of inventory vs. latitude, as well as annual precipitation, as factors that control the depositional inventory. For the latitudinal classification of Hardy et al. (1973) only two measurements between 0° and 10°S were located in Africa (1.2°S Muguga, Kenya & 9.0°S Luanda, Angola; Hardy et al., 1973; Kelley et al., 1999). Both stations receive a substantially lower annual precipitation (960 and 430 mm for Kenya and Angola, respectively) than the NiCo region ($>1400 \text{ mm yr}^{-1}$; Fick and Hijmans, 2017) and show contrasting $^{239+240}\text{Pu}$ inventories of 19.2 Bq m^{-2} in Kenya and 3.4 Bq m^{-2} in Angola. Hence, it is not surprising to find higher baseline $^{239+240}\text{Pu}$ inventories within the NiCo region than in Kenya or Angola. The three pristine forests show mean $^{239+240}\text{Pu}$ inventories between 33 and 48 Bq m^{-2} (DR Congo: $32.7 \pm 7.7 \text{ Bq m}^{-2}$; Uganda: $42.91 \pm 15.5 \text{ Bq m}^{-2}$; Rwanda: $48.4 \pm 18.2 \text{ Bq m}^{-2}$), which is sufficiently high for conducting soil redistribution studies. However, half of the slope sites (14 of 28) at the cropland site in DR Congo fall below the detection limit (0.01 Bq kg^{-1} ; $\sim 5 \text{ Bq m}^{-2}$). This is partly caused by the sampling design of this study, which is based on large and deep single soil increments that cover the soil depth from 0 to 60 cm and 60 to 100 cm at the slope and foot-slope positions. A straightforward way to increase the $^{239+240}\text{Pu}$ activity in the sample is the reduction of the sample increment depth for a corresponding increase of the topsoil proportion that has a higher $^{239+240}\text{Pu}$ activity (see 20 cm increments taken at cropland plateau sites Fig. 3). However, a reduction of the sampling increments necessarily requires an additional analysis of deeper soil layers in highly degraded soil systems, particularly in regions with complex soil redistribution patterns to include the full $^{239+240}\text{Pu}$ inventory. For future $^{239+240}\text{Pu}$ based soil redistribution investigations in tropical cropland areas, a two-layered sampling scheme is suggested that individually samples the plough layer and the deeper soil layer (e.g. $0\text{-}20 \text{ cm}$ and $20\text{-}60 \text{ cm}$ at a slope and an additional $60\text{-}100 \text{ cm}$ sample at foot slope sites). The two-layered sampling scheme accounts for the large number of samples falling below the detection limit and allows for a better understanding of the $^{239+240}\text{Pu}$ depth distribution and thereby depletion status of the sediment source area. In tropical forests, the sampling scheme can be focused on mineral soil as the activity in the litter and organic layer does not substantially contribute to the $^{239+240}\text{Pu}$ inventory. Sedimentation at the three forest study sites did not exceed 60 cm at the foot slope, suggesting that a sampling depth of $0\text{-}60 \text{ cm}$ is sufficient to include the full inventory. Nevertheless, additional deeper soil sampling, with reduced increment depth (e.g. 20 cm) at foot slope locations, is suggested to account for different environmental conditions and potentially higher foot slope sedimentation. The determination of appropriate reference sites is critical. At plateau sites, three depth increments ($0\text{-}20 \text{ cm}$; $20\text{-}40 \text{ cm}$; $40\text{-}60 \text{ cm}$) were sampled to understand the depth distribution of sites without substantial water erosion. At the plateau sites, no $^{239+240}\text{Pu}$ activity was found in the $40\text{-}60 \text{ cm}$ soil layer. Hence, a single sample that integrates soil from 0 to 40 cm depth seems to be sufficient. However,

the number of reference samples should be high enough to represent a robust mean with regards to $^{239+240}\text{Pu}$ variability typically associated with the corresponding land use.

320

$^{239+240}\text{Pu}$ reference inventory

In this study, two different reference scenarios are taken into account to address potential differences of $^{239+240}\text{Pu}$ inventories between stable forest and cropland plateau positions: (i) Ref_{for} mean of specific forest sites (ii) and Ref_{plt} mean of cropland plateau positions of the specific sites. A high variability of $^{239+240}\text{Pu}$ activities and corresponding inventories in three pristine tropical forests within similar topographic positions that exceeds differences in Pu inventories along slope positions is found (Fig. 3). The variation of forest $^{239+240}\text{Pu}$ inventories due to bioturbation and fallout infiltration patterns (e.g. caused by throughfall or stem flow patterns) exceeds a potential soil redistribution impact, which is illustrated by the standard deviation of the plateau sites that covers the variability of the slope and foot-slope composite samples (Fig. 3). Additional evidence that soil redistribution processes in the studied forest systems are low, is that no major differences between chemical and physical soil properties are found along geomorphic gradients (Reichenbach et al., 2021). This finding is in line with global erosion plot studies from tropical forest plots (mean erosion $0.2 \text{ Mg ha}^{-1} \text{ yr}^{-1}$, 39 plots with 116 plot years; Xiong et al., 2019). Observations on sediment delivery monitoring in the NiCo region show that the amount of sediment delivery from pristine forests is typically less than $1 \text{ Mg ha}^{-1} \text{ yr}^{-1}$ (personal communication with Simon Baumgartner, UCL runoff monitoring FORSEDCO project). Furthermore, Drake et al. (2019) exemplarily showed in the NiCo region that particular matter export within pristine tropical forest catchments are dominated by organic matter export with little to no mineral sediment being transported. In contrast, partly deforested catchments with agricultural use show substantial carbon delivery by organo-mineral complexes that indicates detachment and transport of the mineral soil layers, which is again in line with the soil erosion results of this study (Drake et al., 2019). Hence, the forest sites are assumed to represent almost the entire $^{239+240}\text{Pu}$ inventory of the baseline inventory without substantial loss due to soil erosion. The basic assumption behind the reference sites is that the full inventory is preserved as no soil redistribution has taken place and the $^{239+240}\text{Pu}$ inventories of both Ref_{for} and Ref_{plt} are supposed to be similar. However, the mean cropland plateau $^{239+240}\text{Pu}$ inventory in Uganda is about half ($24.4 \pm 7.6 \text{ Bq m}^{-2}$) and in DR Congo only a quarter ($8.0 \pm 1.0 \text{ Bq m}^{-2}$) of the mean inventories found in the nearby (<30 km) pristine forest sites (Fig. 2), which cannot be explained by local rainfall heterogeneities and corresponding spatial patterns of the baseline inventory. Deeper soil layers below 60 cm depth at cropland foot-slope positions show $^{239+240}\text{Pu}$ activity, which is a clear proxy for substantial sedimentation (Fig. 3). However, the $^{239+240}\text{Pu}$ inventory of these locations are not exceeding the forest reference inventory Ref_{for} and would therefore be interpreted as weak soil erosion applying the mass balance model. This is unexpected and can point at a variety of different processes at play that were not investigated by this study. The measured $^{239+240}\text{Pu}$ activity at the foot-slope positions may underestimate the $^{239+240}\text{Pu}$ inventory due to three potential processes: (i) Sedimentation exceeds the sampling depths that does not include the full inventory, (ii) deposition of sediments that are already depleted in $^{239+240}\text{Pu}$ as the inventory at the eroded source area has been subject to pronounced soil and corresponding $^{239+240}\text{Pu}$ loss and (iii) sedimentation sites may temporarily turn into erosion sites. However, an indication for these processes would be an increasing

330
335
340
345
350

²³⁹⁺²⁴⁰Pu activity with soil depth, which is not reflected in the data (Fig. 3). Another potential explanation is that ²³⁹⁺²⁴⁰Pu inventories are reduced due to plant uptake and subsequent plant harvest. However, a substantial plant uptake by crops, like that observed for the fallout radionuclide ¹³⁷Cs (White and Broadley, 2000; Zhu and Smolders, 2000), is unlikely as no elevated ²³⁹⁺²⁴⁰Pu activity was found in harvested crops of other studies (Akleyev et al., 2000). Another potential pathway of soil and ²³⁹⁺²⁴⁰Pu leaving the cropland plateau sites is harvest erosion associated to commonly cultivated root crops (i.e. cassava, sweet potato, groundnuts). In temperate regions, harvest erosion rates up to 12 Mg ha⁻¹ yr⁻¹ have been reported for different crop types (potato: 2.5 - 6 Mg ha⁻¹ yr⁻¹; Auerswald and Schmidt, 1986; Belotserkovsky and Larinovo, 1988; Ruyschaert et al., 2007) (sugarbeet: 5 - 8 Mg ha⁻¹ yr⁻¹; Auerswald et al., 2006) (chicory: 8.1 - 11.8 Mg ha⁻¹ yr⁻¹; Poesen et al., 2001). With cassava and sweet potato being the main food crops within the NiCo region (cassava has a higher proportion on the less fertile soils in the DR Congo, while more sweet potato is cultivated in Uganda), this is a likely source of reduction of ²³⁹⁺²⁴⁰Pu inventories. To illustrate the potential effect of harvest erosion, a simple example shows that 5.5 Mg ha⁻¹ yr⁻¹ of sediment delivery would roughly cause a 20% reduction of the baseline reference over 55 years (assuming a 20 cm plough depth and 1.35 Mg m⁻² bulk density). Harvest erosion can be assumed as a process that has a limited spatial distribution as long as the land use and crop yields are not causing pronounced spatial patterns. Therefore, in systems where harvest erosion is a relevant driver of ²³⁹⁺²⁴⁰Pu export, Ref_{plt} would be the valid reference for soil redistribution estimations. However, an accurate estimation of the contribution of ²³⁹⁺²⁴⁰Pu loss due to harvest erosion since the 1960s is impossible as limited information is available on soil harvesting loss of cassava and potato by hand cultivation. Therefore, both Ref_{for} and Ref_{plt} are taken into account within this study to cover the range from a fully preserved to a depleted ²³⁹⁺²⁴⁰Pu reference inventory in the study sites.

370

Soil redistribution in cropland of the NiCo region

Both cropland sites show indications of soil redistribution. Particularly the cropland study site in DR Congo shows evidence for (i) soil loss due to a high number of slope samples with activities falling below the detection limit (50%) and (ii) sedimentation as evidenced by a clear ²³⁹⁺²⁴⁰Pu fingerprint in the deeper soil layers of the foot-slope samples. Compared to the Ugandan cropland, the DR Congo cropland shows a much stronger difference between ²³⁹⁺²⁴⁰Pu inventories of slope positions and Ref_{for} (Fig. 2). We relate this discrepancy of the different duration since DR Congo and Uganda tropical forest has been converted into cropland. Forest into cropland conversion at the Ugandan study site took place during the 1970's. Hence, the area was under arable use for about 40 years compared to 55 years (since the test ban treaty) at the DR Congo study site. Therefore, the Ugandan cropland was exposed to soil erosion for a roughly 27% shorter time compared to DR Congo cropland. However, the relative ²³⁹⁺²⁴⁰Pu inventory reduction at slope sites in Uganda is about 29% compared to Ref_{for}, while in DR Congo the relative reduction is 83%. The much stronger relative ²³⁹⁺²⁴⁰Pu reduction in DR Congo cannot be just explained by the shorter cropland cultivation period of the Ugandan cropland site. In direct comparison between the two sites, no major difference regarding slope steepness (12°-13° in both study sites) and rainfall erosivity (Fenta et al., 2017) was observed. The different land use during the main time of atmospheric nuclear weapon tests (DR Congo=cropland; Uganda=forest) might have caused higher spatial variability of the initial ²³⁹⁺²⁴⁰Pu inventory associated with complex infiltration (e.g. stem flow) and

385

troughfall patterns in forests (Hofhansl et al., 2012). However, these small-scale variation is likely to be homogenised after 40 years of lateral soil movement by plough-based arable use. The main cultivated crops differ between the Ugandan and DR Congo study sites substantially. In Uganda sweet potato and maize are the major crops, while DR Congo cropland is dominated by cassava. This difference may have an impact on water and harvest driven soil erosion processes.

390 The determined mean soil redistribution rates at cropland slopes found in this study (negative values indicate net erosion and positive values net sedimentation; DR Congo: $-51.4 \text{ Mg ha}^{-1} \text{ yr}^{-1} \text{ Ref}_{\text{for}}$, $-1.4 \text{ Mg ha}^{-1} \text{ yr}^{-1} \text{ Ref}_{\text{plt}}$; Uganda (referred to 40 yrs. of arable land use and corresponding soil redistribution): $-18.4 \text{ Mg ha}^{-1} \text{ yr}^{-1} \text{ Ref}_{\text{for}}$, $27.8 \text{ Mg ha}^{-1} \text{ yr}^{-1} \text{ Ref}_{\text{plt}}$) show contrasting results with respect to the assumed reference (Fig. 5). In comparison to values observed (Boardman and Poesen, 2006) and modelled globally (Borrelli et al., 2017), the high erosion Ref_{for} simulations are in good agreement with plot monitoring results
395 observed in the region (mean soil loss of $68.2 \text{ Mg ha}^{-1} \text{ yr}^{-1}$; Lewis and Nyamulinda, 1996). Over very short distances, high soil redistribution heterogeneities ranging from sedimentation to heavy soil loss are found, which might be an effect of smallholder farming structures. These farming structures are potentially mitigating soil loss rates in the NiCo region due to decreasing hydrologic connectivity (Nunes et al., 2018; Baartman et al., 2020) along slopes due to a high degree of “patchiness” and a large number of field boundaries (mean field size 450 m^2 ; Fig. 5). Any conversion of this smallholder farming structure into
400 large scale farming structures, as known from regions with mechanised agriculture, will have devastating effects on soil degradation rates in the region.

5. Conclusions

This study demonstrates the feasibility of analysing fallout radionuclides ^{239}Pu and ^{240}Pu as a tool to assess soil degradation processes in tropical Africa. Interpreting $^{239+240}\text{Pu}$ activity and inventories in soils and organic layers, we assessed soil
405 redistribution rates along three pristine forests catenae and in two cropland catchments in the White Nile-Congo rift region. $^{239+240}\text{Pu}$ inventories in forest did not follow a topography related distribution, which points at minor soil erosion. In contrast, cropland sites show signs for substantial soil erosion and sedimentation that exceeds 40 cm over a period of 55 years. However, the selection of an appropriate reference is critical due to a potential $^{239+240}\text{Pu}$ inventory reduction associated with cropland use other than water erosion. Very high intra-slope variability of the $^{239+240}\text{Pu}$ inventories in cropland was found (coefficient of
410 variation up to 67%) with sites of pronounced sedimentation in close distance to highly eroded sites, potentially a result of soil cover dynamics due to smallholder farming structures with small fields and individual management. Keeping smallholder farming structures active is essential to mitigate soil degradation in the region, also under current agricultural intensification efforts. Particularly in regions with limited infrastructure and challenging monitoring conditions, $^{239+240}\text{Pu}$ based soil redistribution analysis can shed light on the pace of soil degradation, which remains a major challenge for future food security
415 in tropical Africa.

Data availability

The data is available within the project TropSOC database publication:

420 Doetterl, S., Asifiwe, R. K., Baert, G., Bamba, F., Bauters, M., Bukombe, B., Cadisch, G., Cizungu, L., Cooper, M., Hoyt, A., Kabaske, C., Kalbitz, K., Kidinda, K. L., Maier, A., Mainka, M., Mayrock, J., Muhindo, D., Mujinya, B., Mukotanyi, S. M., Nabahungu, L., Reichenbach, M., Rewald, B., Six, J., Stegmann, A., Summerauer, L., Unseld, R., van Oost, K., Verheyen, K., Vogel, C., Wilken, F., and Fiener, P.: Organic matter cycling along geochemical, geomorphic and disturbance gradients in forests and cropland of the African Tropics - Project TropSOC DATABASE_v1.0., Earth System Science Data Discussions, <https://doi.org/10.5194/essd-2021-73>, 2021a.

425 Doetterl, S., Bukombe, B., Cooper, M., Kidinda, L., Muhindo, D., Reichenbach, M., Stegmann, A., Summerauer, L., Wilken, F., and Fiener, P.: P. TropSOC Database. Version 1.0, GFZ Data Services, <https://doi.org/10.5880/figeo.2021.009>, 2021b.

Author contribution

This paper represents a result of collegial teamwork. FW, SD, PF and KvO developed the study design. ²³⁹⁺²⁴⁰Pu activity analysis was carried out by MK. DM carried out the field campaign. Data processing and illustration was carried out by FW.
430 FW, SD, PF, KVO and KM contributed to data analysis and interpretation. FW drafted the manuscript, while all authors reviewed and approved the final version.

Competing interest

The authors declare that they have no conflict of interest.

Acknowledgements

435 We gratefully acknowledge the outstanding commitment for project TropSOC and fieldwork support of Landry Cizugu, Benjamin Bukombe, Laurent Kidinda, Mario Reichenbach and for labwork Anna Stegmann, Julia Mayrock, Moritz Mainka and Robin Unseld.

References

- 440 Akleyev, A. V., Kostyuchenko, V. A., Peremyslova, L. M., Baturin, V. A., and Popova, I. Y.: Radioecological impacts of the Techa River contamination, *Health Phys.*, 79, 36-47, 10.1097/00004032-200007000-00008, 2000.
- Alewell, C., Pitois, A., Meusburger, K., Ketterer, M., and Mabit, L.: Pu239+240 from "contaminant" to soil erosion tracer: Where do we stand?, *Earth-Sci. Rev.*, 172, 107-123, 10.1016/j.earscirev.2017.07.009, 2017.
- 445 Amundson, R., Berhe, A. A., Hopmans, J. W., Olson, C., Sztein, A. E., and Sparks, D. L.: Soil and human security in the 21st century, *Science*, 348, 12610711-12610716, 2015.
- Angima, S. D., Stott, D. E., O'Neill, M. K., Ong, C. K., and Weesies, G. A.: Soil erosion prediction using RUSLE for central Kenyan highland conditions, *Agric., Ecosyst. Environ.*, 97, 295-308, [https://doi.org/10.1016/S0167-8809\(03\)00011-2](https://doi.org/10.1016/S0167-8809(03)00011-2), 2003.
- Auerswald, K., and Schmidt, F.: Atlas der Erosionsgefährdung in Bayern, GLA-Fachberichte, 1, 1986.
- 450 Auerswald, K., Gerl, G., and Kainz, M.: Influence of cropping system on harvest erosion under potato, *Soil Till. Res.*, 89, 22-34, 2006.
- Baartman, J. E. M., Nunes, J. P., Masselink, R., Darboux, F., Bieters, C., Degré, A., Cantreul, V., Cerdan, O., Grangeon, T., Fiener, P., Wilken, F., Schindewolf, M., and Wainwright, J.: What do models tell us about water and sediment connectivity?, *Geomorphology*, 367, 1-17, <https://doi.org/10.1016/j.geomorph.2020.107300>, 2020.
- 455 Belotserkovsky, Y., and Larinovo, A.: Removal of soil by harvest of potatoes and root crops (in Russian), *Vestnik Moskovskogo Universiteta Seriya 5: Geografia*, 4, 49-54, 1988.
- Boardman, J., and Poesen, J.: Soil Erosion in Europe: Major Processes, Causes and Consequences, in: *Soil Erosion in Europe*, John Wiley & Sons, Ltd, 477-487, 2006.
- Borrelli, P., Robinson, D. A., Fleischer, L. R., Lugato, E., Ballabio, C., Alewell, C., Meusburger, K., Modugno, S., Schütt, B., Ferro, V., Bagarello, V., Oost, K. V., Montanarella, L., and Panagos, P.: An assessment of the global impact of 21st century land use change on soil erosion, *Nature Communications*, 8, 1-13, 10.1038/s41467-017-02142-7, 2017.
- 460 Borrelli, P., Robinson, D. A., Panagos, P., Lugato, E., Yang, J. E., Alewell, C., Wuepper, D., Montanarella, L., and Ballabio, C.: Land use and climate change impacts on global soil erosion by water (2015-2070), *Proceedings of the National Academy of Sciences*, 117, 21994-22001, 10.1073/pnas.2001403117, 2020.
- 465 Calitri, F., Sommer, M., Norton, K., Temme, A., Brandova, D., Portes, R., Christl, M., Ketterer, M. E., and Egli, M.: Tracing the temporal evolution of soil redistribution rates in an agricultural landscape using Pu239+240 and Be-10, *Earth Surf. Process. Landf.*, 44, 1783-1798, 10.1002/esp.4612, 2019.
- Chamberlin, J., Jayne, T. S., and Headey, D.: Scarcity amidst abundance? Reassessing the potential for cropland expansion in Africa, *Food Policy*, 48, 51-65, 10.1016/j.foodpol.2014.05.002, 2014.
- 470 Chartin, C., Evrard, O., Salvador-Blanes, S., Hirschberger, F., Van Oost, K., Lefevre, I., Daroussin, J., and Macaire, J. J.: Quantifying and modelling the impact of land consolidation and field borders on soil redistribution in agricultural landscapes (1954-2009), *Catena*, 110, 184-195, 10.1016/j.catena.2013.06.006, 2013.
- Doetterl, S., Asifiwe, R. K., Baert, G., Bamba, F., Bauters, M., Bukombe, B., Cadisch, G., Cizungu, L., Cooper, M., Hoyt, A., Kabaske, C., Kalbitz, K., Kidinda, K. L., Maier, A., Mainka, M., Mayrock, J., Muhindo, D., Mujinya, B., Mukotanyi, S., M., Nabahungu, L., Reichenbach, M., Rewald, B., Six, J., Stegmann, A., Summerauer, L., Unseld, R., van Oost, K., Verheyen, K., Vogel, C., Wilken, F., and Fiener, P.: Organic matter cycling along geochemical, geomorphic and disturbance gradients in forests and cropland of the African Tropics - Project TropSOC DATABASE_v1.0., *Earth System Science Data Discussions*, <https://doi.org/10.5194/essd-2021-73>, 2021a.
- 475 Doetterl, S., Bukombe, B., Cooper, M., Kidinda, L., Muhindo, D., Reichenbach, M., Stegmann, A., Summerauer, L., Wilken, F., and Fiener, P.: P. TropSOC Database. Version 1.0, GFZ Data Services, <https://doi.org/10.5880/figeo.2021.009>, 2021b.
- 480 Drake, T. W., Van Oost, K., Barthel, M., Bauters, M., Hoyt, A. M., Podgorski, D. C., Six, J., Boeckx, P., Trumbore, S. E., Ntaboba, L. C., and Spencer, R. G. M.: Mobilization of aged and biolabile soil carbon by tropical deforestation, *Nat. Geosci.*, 12, 541-547, 10.1038/s41561-019-0384-9, 2019.
- 485 Evans, D. L., Quinton, J. N., Davies, J. A. C., Zhao, J., and Govers, G.: Soil lifespans and how they can be extended by land use and management change, *Environmental Research Letters*, 15, 1-10, 10.1088/1748-9326/aba2fd, 2020.

- Evrard, O., Chaboche, P.-A., Ramon, R., Foucher, A., and Lacey, J. P.: A global review of sediment source fingerprinting research incorporating fallout radiocesium (^{137}Cs), *Geomorphology*, 362, 1-22, <https://doi.org/10.1016/j.geomorph.2020.107103>, 2020.
- FAO, and ITPS: Status of the World's Soil Resources – Main Report, FAO, Rome, Italy, 2015.
- 490 Fenta, A. A., Yasuda, H., Shimizu, K., Haregeweyn, N., Kawai, T., Sultan, D., Ebabu, K., and Belay, A. S.: Spatial distribution and temporal trends of rainfall and erosivity in the Eastern Africa region, *Hydrol. Processes*, 31, 4555-4567, 10.1002/hyp.11378, 2017.
- Fick, S. E., and Hijmans, R. J.: WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas, *Int. J. Climatol.*, 37, 4302-4315, <https://doi.org/10.1002/joc.5086>, 2017.
- 495 Fiener, P., Wilken, F., and Auerswald, K.: Filling the gap between plot and landscape scale – eight years of soil erosion monitoring in 14 adjacent watersheds under soil conservation at Scheyern, Southern Germany, *Adv. Geosci.*, 48, 31-48, 10.5194/adgeo-48-31-2019, 2019.
- Govers, G., Merckx, R., van Wesemae, B., and Van Oost, K.: Soil conservation in the 21st century: why we need smart agricultural intensification, *Soil*, 3, 45-59, 10.5194/soil-3-45-2017, 2017.
- 500 Hardy, E. P., Krey, P. W., and Volchok, H. L.: Global inventory and distribution of fallout plutonium, *Nature*, 241, 444-445, 10.1038/241444a0, 1973.
- Hofhansl, F., Wanek, W., Drage, S., Huber, W., Weissenhofer, A., and Richter, A.: Controls of hydrochemical fluxes via stemflow in tropical lowland rainforests: Effects of meteorology and vegetation characteristics, *J. Hydrol.*, 452, 247-258, 10.1016/j.jhydrol.2012.05.057, 2012.
- 505 Kelley, J. M., Bond, L. A., and Beasley, T. M.: Global distribution of Pu isotopes and Np-237, *Science of the Total Environment*, 238, 483-500, 10.1016/s0048-9697(99)00160-6, 1999.
- Ketterer, M. E., Hafer, K. M., Link, C. L., Kolwaite, D., Wilson, J., and Mietelski, J. W.: Resolving global versus local/regional Pu sources in the environment using sector ICP-MS, *J. Anal. At. Spectrom.*, 19, 241-245, 10.1039/b302903d, 2004.
- Kotteck, M., Grieser, J., Beck, C., Rudolf, B., and Rubel, F.: World map of the Koppen-Geiger climate classification updated, *Meteorol. Z.*, 15, 259-263, 10.1127/0941-2948/2006/0130, 2006.
- 510 Lal, R., Tims, S. G., Fifield, L. K., Wasson, R. J., and Howe, D.: Applicability of Pu-239 as a tracer for soil erosion in the wet-dry tropics of northern Australia, *Nuclear Instruments & Methods in Physics Research Section B-Beam Interactions with Materials and Atoms*, 294, 577-583, 10.1016/j.nimb.2012.07.041, 2013.
- Lewis, L. A., and Nyamulinda, V.: The critical role of human activities in land degradation in Rwanda, *Land Degrad. Dev.*, 7, 47-55, 1996.
- 515 Meusburger, K., Mabit, L., Ketterer, M., Park, J. H., Sandor, T., Porto, P., and Alewell, C.: A multi-radionuclide approach to evaluate the suitability of Pu $^{239+240}$ as soil erosion tracer, *Science of the Total Environment*, 566, 1489-1499, 10.1016/j.scitotenv.2016.06.035, 2016.
- Montanarella, L., Pennock, D. J., McKenzie, N., Badraoui, M., Chude, V., Baptista, I., Mamo, T., Yemefack, M., Aulakh, M. S., Yagi, K., Hong, S. Y., Vijarnsorn, P., Zhang, G. L., Arrouays, D., Black, H., Krasilnikov, P., Sobocka, J., Alegre, J., Henriquez, C. R., Mendonca-Santos, M. D., Taboada, M., Espinosa-Victoria, D., AlShankiti, A., AlaviPanah, S. K., Elsheikh, E. A. E., Hempel, J., Arbestain, M. C., Nachtergaele, F., and Vargas, R.: World's soils are under threat, *Soil*, 2, 79-82, 10.5194/soil-2-79-2016, 2016.
- 520 Nunes, J. P., Wainwright, J., Biellers, C. L., Darboux, F., Fiener, P., Finger, D., and Turnbull, L.: Better models are more effectively connected models, *Earth Surf. Process. Landf.*, 43, 1355-1360, <https://doi.org/10.1002/esp.4323>, 2018.
- Nyesheja, E. M., Chen, X., El-Tantawi, A. M., Karamage, F., Mupenzi, C., and Nsengiyumva, J. B.: Soil erosion assessment using RUSLE model in the Congo Nile Ridge region of Rwanda, *Physical Geography*, 40, 339-360, 10.1080/02723646.2018.1541706, 2019.
- 525 Nyssen, J., Poesen, J., Moeyersons, J., Deckers, J., Haile, M., and Lang, A.: Human impact on the environment in the Ethiopian and Eritrean highlands - a state of the art, *Earth-Sci. Rev.*, 64, 273-320, 10.1016/s0012-8252(03)00078-3, 2004.
- 530 Poesen, J. W. A., Verstraeten, G., Soenens, R., and Seynaeve, L.: Soil losses due to harvesting of chicory roots and sugar beet: an underrated geomorphic process?, *Catena*, 43, 35-47, 2001.
- Porto, P., and Walling, D. E.: Using plot experiments to test the validity of mass balance models employed to estimate soil redistribution rates from ^{137}Cs and ^{210}Pb -ex measurements, *Appl. Radiat. Isot.*, 70, 2451-2459, 10.1016/j.apradiso.2012.06.012, 2012.
- 535

- R: A Language and Environment for Statistical Computing: <https://www.R-project.org/>, access: 27.10.2020, 2019.
- Reichenbach, M., Fiener, P., Garland, G., Six, S., and Doetterl, S.: The role of geochemistry in organic carbon stabilization in tropical rainforest soils, *SOIL Discuss.*, 2021.
- 540 Ruyschaert, G., Poesen, J., Auerswald, K., Verstraeten, G., and Govers, G.: Soil losses due to potato harvesting at the regional scale in Belgium, *Soil Use Manag.*, 23, 156-161, 2007.
- Ryan, J. N., Illangasekare, T. H., Litaor, M. I., and Shannon, R.: Particle and plutonium mobilization in macroporous soils during rainfall simulations, *Environmental Science & Policy*, 32, 476-482, 1998.
- 545 van Ittersum, M. K., van Bussel, L. G. J., Wolf, J., Grassini, P., van Wart, J., Guilpart, N., Claessens, L., de Groot, H., Wiebe, K., Mason-D'Croz, D., Yang, H., Boogaard, H., van Oort, P. A. J., van Loon, M. P., Saito, K., Adimo, O., Adjei-Nsiah, S., Agali, A., Bala, A., Chikowo, R., Kaizzi, K., Kouressy, M., Makoi, J. H. J. R., Ouattara, K., Tesfaye, K., and Cassman, K. G.: Can sub-Saharan Africa feed itself?, *Proceedings of the National Academy of Sciences*, 113, 14964-14969, 10.1073/pnas.1610359113, 2016.
- Vanmaercke, M., Poesen, J., Broeckx, J., and Nyssen, J.: Sediment yield in Africa, *Earth-Sci. Rev.*, 136, 350-368, 10.1016/j.earscirev.2014.06.004, 2014.
- 550 Wallbrink, P. J., and Murray, A. S.: Use of radionuclides as indicators of erosion processes, *Hydrol. Processes*, 7, 297-304, 10.1002/hyp.3360070307, 1993.
- White, P. J., and Broadley, M. R.: Mechanisms of caesium uptake by plants, *New Phytol.*, 147, 241-256, 10.1046/j.1469-8137.2000.00704.x, 2000.
- 555 Wilcke, W., Yasin, S., Abramowski, U., Valarezo, C., and Zech, W.: Nutrient storage and turnover in organic layers under tropical montane rain forest in Ecuador, *Eur. J. Soil Sci.*, 53, 15-27, 10.1046/j.1365-2389.2002.00411.x, 2002.
- World Population Review: <https://worldpopulationreview.com>, access: 13.12.2020, 2020.
- WRB: World reference base for soil resources 2006, FAO, Rome, 1-145, 2006.
- 560 Wynants, M., Kelly, C., Mtei, K., Munishi, L., Patrick, A., Rabinovich, A., Nasser, M., Gilvear, D., Roberts, N., Boeckx, P., Wilson, G., Blake, W. H., and Ndakidemi, P.: Drivers of increased soil erosion in East Africa's agro-pastoral systems: changing interactions between the social, economic and natural domains, *Regional Environmental Change*, 19, 1909-1921, 10.1007/s10113-019-01520-9, 2019.
- Xiong, M. Q., Sun, R. H., and Chen, L. D.: A global comparison of soil erosion associated with land use and climate type, *Geoderma*, 343, 31-39, 10.1016/j.geoderma.2019.02.013, 2019.
- 565 Zhang, H., Aldana-Jague, E., Clapuyt, F., Wilken, F., Vanacker, V., and Van Oost, K.: Evaluating the potential of post-processing kinematic (PPK) georeferencing for UAV-based structure-from-motion (SfM) photogrammetry and surface change detection, *Earth Surf. Dynam.*, 7, 807-827, 10.5194/esurf-7-807-2019, 2019.
- Zhu, Y. G., and Smolders, E.: Plant uptake of radiocaesium: a review of mechanisms, regulation and application, *J. Exp. Bot.*, 51, 1635-1645, 10.1093/jexbot/51.351.1635, 2000.

Table 1: Coordinates and main characteristics of investigated study sites. Mean annual rainfall and temperature data is based on Fick and Hijmans (2017).

Study site	Location [Lat., Long., Elevation]	Max. slope [°]	Mean annual rainfall [mm]	Mean annual temperature [°C]	Mean field size [m ²]
DRC forest	2°19'S, 28°44'E, 2208-2248 m	60	1925	15.2	-
Uganda forest	0°28'N, 30°22'E, 1271-1306 m	55	1277	20.7	-
Rwanda forest	2°28'S, 29°06'E, 1271-1306 m	60	1689	17.3	-
DRC cropland	2°35'S, 28°43'E, 1711-1747 m	17	1586	17.7	662
Uganda cropland	0°34'N, 30°20'E, 1470-1510 m	20	1459	19.5	336

Table 2: Numbers of samples taken at three forest (DR Congo, Uganda, Rwanda) and two cropland (DR Congo, Uganda) study sites. While the O horizon depth is measured individually for each sample, the L horizon depth is assumed to be 1 cm as it was not possible to be accurately measured. L = L horizon; O = O horizon; M = mineral layer 1: 0-60 cm and 2: 60-120 in forest and 60-100 cm in cropland.

Forest			
	Plateau	Slope	Foot-slope
L, O, M1	10	6	6
M2	0	0	6
Cropland			
	Plateau*	Slope	Foot-slope
M1	12	27	12
M2	0	0	12

*Three depth increments 0 - 20 cm, 20 - 40 cm, 40 - 60 cm

580

Figures




Figure 1: Study area, locations of forest and cropland sites, topography and land use in the White Nile-Congo rift

585 region (land use data is based on: ESA Climate Change Initiative - Land Cover project 2017).




Figure 2: Distribution of $^{239+240}\text{Pu}$ inventories in three forests and two cropland study sites. In (a) the $^{239+240}\text{Pu}$ histogram for three forest sites is given, while (b) and (c) represent the cropland distribution at the slope and foot-slope topographic positions. The vertical lines represent the mean values serving as forest reference (Ref_{for}) and cropland plateau reference (Ref_{pit}). Red section at the left of the graphs illustrates the detection limit.

590




Figure 3: Catenae related depth profiles of $^{239+240}\text{Pu}$ activity and standard deviations (whisker) within three pristine forests (DRC: DR Congo, UGA: Uganda, RWA: Rwanda) and two cropland study sites. Please note that the L and O horizons are for illustrational purposes shown thicker than they naturally appear.




Figure 4: DR Congo (DRC) and Uganda (UGA) cropland soil erosion results of the mass balance model. A scenario
 600 assessment was carried out to understand $^{239+240}\text{Pu}$ inventory reduction until detection limit is reached after the
 simulation period: (a) soil erosion magnitude and corresponding (b) soil erosion frequency according to the mean
 forest (Ref_{for}) and mean cropland plateau (Ref_{plt}) assumption. (a) Represents the minimum quantity of soil erosion
 to pass the detection limit with different reference $^{239+240}\text{Pu}$ inventories. All displayed curves show the
 corresponding scenario runs of minimum soil erosion to cause a $^{239+240}\text{Pu}$ inventory reduction that falls below the
 605 detection limit. (b) Comparison between a continuous 55 years soil erosion rate vs. 5 erosive years that cause the
 same quantity of soil erosion. Soil erosion from 1 to 100 cm (55 yrs)⁻¹ is simulated. The coloured area describes
 the difference between the two scenarios, where the lower line is the 5 year scenario.




Figure 5: Soil redistribution derived from $^{239+240}\text{Pu}$ measurements using a mass balance model. Two different reference inventories (upper: Ref_{for} = mean forest reference; lower: Ref_{plt} mean cropland plateau reference) in two cropland areas of the White Nile-Congo rift region (left: DR Congo, right: Uganda) were used for calculation.

610

Appendix A

615 Figure A1: Overview photos and orthomosaics of the DR Congo (left) and Uganda (right) cropland sites. Image acquisition dates: DR Congo (left panel) overview photo Sep. 1st, 2018, orthomosaic Sep. 14th, 2019; Uganda (right panel) overview photo Oct. 12th, 2019, orthomosaic Jun. 12th, 2019.

