

1 **Aluminous clay and pedogenic Fe oxides modulate aggregation and**
2 **related carbon contents in soils of the humid tropics**

3

4 Maximilian Kirsten*¹, Robert Mikutta², Didas N. Kimaro³, Karl-Heinz Feger¹, Karsten Kalbitz¹

5

6

7 ¹Technische Universität Dresden, Institute of Soil Science and Site Ecology, Tharandt, Germany

8 ²Martin-Luther-Universität Halle-Wittenberg, Soil Science and Soil Protection, Halle/Saale Germany

9 ³Mwenge Catholic University, Directorate of Research Innovations and Consultancy, Moshi, Tanzania

10

11

12

13

14

15

16

17

18

19

20

21

22 **Keywords:** tropical soils, aggregate size distribution, aggregate stability, soil mineralogy, kaolinite,
23 gibbsite, goethite, hematite, land-use change

24

25 * maximilian.kirsten@tu-dresden.de

26 **Abstract**

27 Aggregation affects a wide range of physical and biogeochemical soil properties with positive effects on
28 soil carbon storage. For weathered tropical soils, aluminous clays (kaolinite and gibbsite) and pedogenic Fe
29 (oxyhydr)oxides (goethite and hematite; termed 'Fe oxides') have been suggested as important building
30 units for aggregates. However, as aluminosilicates, aluminum hydroxides, and Fe oxides are part of the clay-
31 sized fraction it is hard to separate, how certain mineral phases modulate aggregation. In addition, it is not
32 known what consequences this will have for organic carbon (OC) persistence after land-use change. We
33 selected topsoils with unique mineralogical compositions in the East Usambara Mountains of Tanzania
34 under forest and cropland land uses, varying in contents of aluminous clay and Fe oxides. Across the
35 mineralogical combinations, we determined the aggregate size distribution, aggregate stability, OC contents
36 of aggregate size fractions as well as changes in aggregation and OC contents under forest and cropland
37 land use. Patterns in soil aggregation were rather similar across the different mineralogical combinations
38 (high level of macroaggregation and high aggregate stability). Nevertheless, we found some statistically
39 significant effects of aluminous clay and pedogenic Fe oxides on aggregation and OC storage. An aluminous
40 clay content $> 250 \text{ g kg}^{-1}$ in combination with pedogenic Fe contents $< 60 \text{ g kg}^{-1}$ significantly promoted the
41 formation of large macroaggregates $> 4 \text{ mm}$. In contrast, a pedogenic Fe content $> 60 \text{ g kg}^{-1}$ in combination
42 with aluminous clay content of $< 250 \text{ g kg}^{-1}$ promoted OC storage and persistence even under agricultural
43 use. The combination with low aluminous clay and high pedogenic Fe contents displayed the highest OC
44 persistence, despite conversion of forest to cropland caused substantial disaggregation. Our data indicate
45 that aggregation in these tropical soils is modulated by the mineralogical regime, causing moderate but
46 significant differences in aggregate size distribution. Nevertheless, aggregation was little decisive for overall
47 OC persistence in these highly weathered soils, where OC storage is more regulated by direct mineral-
48 organic interactions.

49

50 **1. Introduction**

51 Many functions of soils such as food production, water purification as well as climate regulation are tightly
52 linked to soil structure (*Bronick and Lal, 2005; FAO, 2015; Six et al., 2004*). Aggregates are the structural
53 backbone of soil and changes in aggregation impacts various processes such as root development, soil
54 erosion, and soil organic carbon (OC) accumulation (*Chaplot et al., 2010; Le Bissonnais et al., 2018*). Based
55 on their size, soil aggregates are typically classified into small microaggregates ($< 20 \mu\text{m}$), large
56 microaggregates (20–250 μm), and macroaggregates ($> 0.25 \text{ mm}$) (*Tisdall and Oades, 1982*). Cementing
57 agents such as clay minerals, metal (oxyhydr)oxides, as well as organic matter (OM) are considered as
58 primary building units of microaggregates (*Totsche et al., 2018*), which provide the basis for the formation
59 of larger soil structural units (*Asano and Wagai, 2014*). The study by *Six et al. (2002)* points to the special
60 role of inorganic compounds such as clay minerals and pedogenic metal oxides in the formation of
61 aggregates in the tropics. Pedogenic iron (Fe_d) (oxyhydr)oxides (summarized as ‘Fe oxides’) have been
62 reported to facilitate macroaggregation (*Peng et al., 2015*) and aggregate stability (*Duiker et al., 2003*).
63 Under the acidic conditions of weathered tropical soils, Fe oxides provide positively charged surfaces
64 capable of reacting with negatively charged inorganic constituents, like clay minerals or OM (*Kaiser and*
65 *Guggenberger, 2003; Kleber et al., 2015; Six et al., 2004; Totsche et al., 2018*). Aggregation might be
66 ascribed to inorganic or organic cementing agents with no consensus about the relevance of each individual
67 agent. Understanding the effects of individual cementing agents for aggregation is needed to disentangle
68 their potential contribution to soil aggregation. For example, the extent of aggregation has been either
69 positively related to the contents of clay and OC (*Chaplot and Cooper, 2015; Paul et al., 2008; Spaccini et*
70 *al., 2001*), or to differences in the clay mineral composition (*Fernández-Ugalde et al., 2013*). Furthermore,
71 *Barthès et al. (2008)* showed that texture had no effect on macroaggregation over a range of tropical soils
72 characterized by low-activity clay minerals. Such kind of uncertainty may derive from the fact that the clay
73 size particle fraction ($< 2\text{-}\mu\text{m}$) not only contains OM and different types of clay minerals, but also variable
74 contents of pedogenic Fe and aluminum (Al) oxides (*Barré et al. 2014; Fernández-Ugalde et al. 2013;*
75 *Wagai and Mayer 2007*). *Denef et al. (2004)* showed that significant differences in the amount of

76 microaggregates encased in macroaggregates can be related to the clay mineral composition (2:1, mixed
77 layer, 1:1 clays). They assume that interactions of 1:1 clay minerals with Fe oxides cause a higher aggregate
78 stability compared to those involving 2:1 clay minerals (*Denef et al., 2002, 2004*). Such mutual interactions
79 between typical aluminous clay-sized minerals (e.g. kaolinite, gibbsite) and pedogenic Fe oxides are thus
80 possible drivers of aggregation in weathered tropical soils (*Durn et al., 2019*).

81 Soil aggregation is considered to be an important process that increases OC persistence, because of
82 the physical separation of OM from microorganisms and their exoenzymes (*Six et al., 2004*). Thus, improved
83 aggregation could contribute to enhanced OC storage in soils (*Kravchenko et al., 2015; Marín-Spiotta et al.,*
84 *2008; Schmidt et al., 2011*). Managing aggregation, e.g., for climate change mitigation, requires profound
85 knowledge on the controls of aggregation and their effects on OC persistence (*Paul et al., 2008*). To the best
86 of our knowledge there are no studies available, which investigated the influence of changes in the content
87 of clay minerals with low activity and the content of pedogenic metal oxides on aggregation under
88 comparable mineralogical conditions for weathered tropical soils. Macroaggregates are particularly
89 susceptible to soil management (*Six et al., 2000a; Totsche et al., 2018*). Consequently, destruction of
90 macroaggregates upon changes from forests to cropland might account for OC losses that were observed in
91 tropical soils (*Don et al., 2011; Kirsten et al., 2019; Mujuru et al., 2013*). The stability of aggregates should
92 thus determine OC losses induced by land-use change and higher losses should be related to lower aggregate
93 stability (*Denef et al., 2002; Le Bissonnais et al., 2018; Six et al., 2000b*). We are currently not aware of any
94 studies that solve the puzzle to which extent the amount of aluminous clay and pedogenic Fe oxides controls
95 soil aggregation and OC storage in highly weathered soils of the humid tropics.

96 This study takes advantage of soils under natural forest and cropland in the East Usambara Mountains
97 of Tanzania. The mineralogical composition of the study soils is very homogeneous with kaolinite and
98 gibbsite as the main aluminous minerals of the clay fraction and goethite and hematite as dominant
99 pedogenic Fe oxides (*Kirsten et al., 2021*). Yet, the ratio of aluminous clays to Fe oxides differed strongly,
100 giving rise to unique mineralogical combinations under both land use types. Thus, the conversion of natural
101 forest to cropland in the study region enables us to evaluate the effect of land-use change under each

102 mineralogical combination on soil physical properties and related OC persistence. In the precursor study,
103 we found a positive relationship between the storage of mineral-associated OC and the ratio of pedogenic
104 Fe to aluminous clay under forest and cropland land use, suggesting that a larger share of Fe oxides is linked
105 to larger OC storage and persistency against land-use change (*Kirsten et al., 2021*). In the present study, we
106 test whether aggregation and its contribution to OC storage follow similar patterns, or are decoupled from
107 the individual contribution of main mineral constituents. In detail, our main research goal was to investigate
108 the individual role of aluminous clay and pedogenic Fe oxides for determining (i) the soil aggregate size
109 distribution, (ii) aggregate stability, (iii) the consequences for OC allocation into different aggregate size
110 fractions, and (iv) the consequences for OC persistence related to land-use change. We hypothesize that the
111 mineralogical combination resulting in the largest aggregate stability also results in the largest OC
112 persistence. For this purpose, we determined the aggregate size distribution of soils under both land uses,
113 determined the OC contents of obtained aggregate fractions, and tested the stability of the two largest
114 aggregate size fractions (2–4 mm and > 4 mm). As a measure of OC persistence, the OC content of
115 aggregate size fractions was compared between the two land uses in the same mineralogical combination.
116 We generally focused on soil samples from 0–10 cm to test our current hypothesis since land-use induced
117 OC losses from soils of the study region largely occur in this depth increment (*Kirsten et al., 2019*).

118 **2. Material and methods**

119 **2.1 Study area and soil sampling**

120 The study was conducted in the Eastern Usambara Mountains of Tanzania close to the village Amani
121 (5°06'00"S; 38°38'00"E). The climate is humid monsoonal with a mean annual precipitation of 1,918 mm,
122 and a mean annual temperature of 20.6°C with low variability within the study area (*Hamilton and Bensted-*
123 *Smith*, 1989). The dominating Acrisols and Alisols, developed from Precambrian crystalline bedrock, are
124 deeply weathered and highly leached, with visible clay illuviation in the subsoil (*Kirsten et al.*, 2019).
125 Briefly, all soil samples were collected on mid-slope position. We sampled six plots under forest and three
126 under annual cropping. The site selection was done based on total clay amount determined in the field and
127 the associated total Fe amount measured with a portable XRF device (*Kirsten et al.*, 2021). We did not
128 observe systematic differences in vegetation composition of the forest sites and NMR spectra showed a
129 similar composition of litter for each of the two land uses investigated (*Kirsten et al.*, 2021). Furthermore,
130 several visits in the study region over the last decade (2012, 2013, 2015, and 2018) combined with personal
131 talks to farmers and local partners working in the region, enabled us to select cropland sites with similar
132 agricultural management (cultivation of cassava (*Manihot esculenta*), hand hoe tillage, biomass burning
133 before seed bed preparation). At each plot, mineral soil from three adjacent and randomly distributed soil
134 pits at mid-slope position was sampled at 0–5 and 5–10 cm depths. This procedure was chosen because we
135 identified two soil horizons at 0–5 and 5–10 cm depth based on differences in color and structure. To have
136 a consistent sampling design, we applied this distinction to the cropland sites, too. Living roots were
137 removed and aliquots of the soils were sieved to < 2 mm after drying at 40°C. For each depth increment,
138 three undisturbed soil cores (100 cm³) were collected for bulk density determination.

139

140 **2.2 Soil analyses**

141 *Basic soil properties and selected mineralogical combinations*

142 Bulk density was determined after drying the soil at 105°C and corrected for coarse fragments (*Carter and*
143 *Gregorich*, 2008). Soil pH was measured in 0.01 M CaCl₂ at a soil to solution ratio of 1 : 2.5. Extraction of

144 poorly crystalline Fe and Al phases as well as of Fe and Al complexed by OM was done with ammonium
145 oxalate according to *Schwertmann* (1964). Effective cation exchange capacity (CEC_{eff}) and base saturation
146 (BS) were determined following the procedure provided by *Trüby and Aldinger* (1989). Contents of OC and
147 total N were analyzed by high temperature combustion at 950°C and thermo-conductivity detection (Vario
148 EL III/Elementar, Heraeus, Langensfeld, Germany). A combined dithionite-citrate-bicarbonate extraction
149 and subsequent texture analysis was applied to determine the contents of aluminous clay and total pedogenic
150 Fe (Fe_d). Briefly, 5–6 g soil pre-treated with 30% H_2O_2 were extracted with 30 g sodium dithionite ($Na_2S_2O_4$)
151 and 1.35 L buffer solution (0.27 M trisodium citrate dihydrate ($C_6H_5Na_3O_7 \cdot 2H_2O$) + 0.11 M sodium
152 bicarbonate ($NaHCO_3$)) at 75°C in a water bath for 15 min (*Mehra and Jackson*, 1958). The Fe concentration
153 of the extracts were measured by inductively coupled plasma optical emission spectroscopy (ICP-OES)
154 using a CIROS-CCD instrument (Spectro, Kleve, Germany). The residues of the extraction were then
155 subjected to a texture analysis using the pipette method (*Gee and Bauder*, 1986). Details of the procedure
156 are described in *Kirsten et al.* (2021). Based on the respective content of aluminous clay and pedogenic Fe
157 oxide in the 5–10 cm depth increment, each sample was assigned to a certain mineralogical combination.
158 The threshold values for aluminous clay and pedogenic Fe to distinguish between “high” and “low” were
159 set to 250 g kg^{-1} and 60 g kg^{-1} , respectively. We differentiated four groups varying in contents of aluminous
160 clay and pedogenic Fe oxides under forest (i.e. ‘low clay–low Fe’, ‘low clay–high Fe’, ‘high clay–low Fe’,
161 ‘high clay–high Fe’), and three analogous groups under cropland (i.e. ‘low clay–low Fe’, ‘low clay–high
162 Fe’, ‘high clay–high Fe’). Using the threshold criterion for assigning the individual samples to a
163 mineralogical combination resulted in an unequal number of repetitions for combinations under forests
164 ($n = 3–7$) whereas those under cropland remained the same ($n = 3$).

165
166 *Aggregate size distribution, aggregate stability and carbon contents*
167 Aggregate size distribution was determined by dry sieving as it most closely resembles soil conditions at
168 the end of the long dry season. Undisturbed soil was dried at 40°C for 48 hours. Separation of aggregate
169 sizes was conducted with a sieving machine (AS 200 control “g”, Retsch, Hanau, Germany) combined with

170 a set of four sieves with meshes of 4, 2, 1, and 0.25 mm, respectively (Larney, 2008). The amplitude was
171 set to 1.51 mm (7.6 g-force), which was applied over a sieving duration of three minutes. Aggregate stability
172 was tested for the two largest aggregate size fractions (2–4 mm and > 4 mm). The fast wetting pretreatment
173 was applied to both fractions (Le Bissonnais, 1996) using a wet-sieving apparatus (Eijkelkamp, Giesbeek,
174 Netherlands) with sieve openings of 63 µm. This procedure simulates the transition of aggregates from dry
175 to rainy season. Sieving was conducted in ethanol for three minutes (stroke 1.3 cm, f = 34 min⁻¹). All
176 aggregates remaining on the sieve were dried at 105°C. Water-stable aggregates were subsequently
177 introduced to a sieving apparatus with a set of five sieves with mesh sizes of 4, 2, 1, 0.63, and 0.25 mm,
178 respectively (Larney, 2008). For each obtained aggregate fraction by dry sieving, OC contents analyzed by
179 high temperature combustion at 950°C and thermo-conductivity detection (Vario EL III/Elementar,
180 Heraeus, Langensfeld, Germany). The mass corrected OC content of a certain aggregate fraction was
181 calculated using equation 1 to resemble the contribution to total soil OC,

$$182 \text{ Mass - corrected } OC_{Aggregate} = \frac{m_i}{\sum_{i=0}^n m_i} \times OC_{Aggregate} \quad (\text{Eq. 1})$$

183 where m_i represents the mass of an aggregate size fraction (g), $\sum m_i$, the sum of masses of all size
184 fractions (g), and $OC_{Aggregate}$ the OC content of aggregate fraction "i".

185 The mean weight diameter (MWD) of aggregates was calculated using equation 2 for undisturbed soil to
186 describe the initial aggregate size distribution, and for the large aggregate size fractions after exposure to
187 the stability test to evaluate the effect of fast wetting on aggregate stability,

$$188 MWD = \sum_{i=0}^n \frac{m_i}{\sum m_i} \times d_i \quad (\text{Eq. 2})$$

189 where m_i represents the mass of an aggregate size fraction (g), $\sum m_i$, the sum of masses of all size
190 fractions (g), and d_i the mean mesh diameter of fraction "i" (mm). The MWD of the aggregate fraction
191 > 4 mm was estimated by doubling the largest sieve size diameter (Youker and McGuinness, 1957).

192

193 **2.3 Statistics and calculations**

194 The mean and standard deviation of data were calculated with the software package R (version 3.6.0). To
195 test for significant differences between mineralogical combinations, land uses, and depths, we applied the

196 linear model function [lm()] in combination with analysis of variance [aov(lm())]. The Tukey-HSD test was
197 used as a post-hoc comparison of means; the LSD-test was applied in the case of non-equality of variances.
198 Regression analysis was used to test for relationships between mineralogical properties and MWD, masses
199 of aggregate size fractions, aggregate stability, and OC losses due to land-use change. Statistical differences
200 are reported at a significance level of $p < 0.05$. Based on our selected threshold values for aluminous clay
201 and pedogenic Fe oxides, we were able to achieve the following number of replicates for the mineralogical
202 combinations: 'low clay–low Fe' under forest ($n = 4$), 'low clay–high Fe' under forest ($n = 4$), 'high clay–
203 low Fe' under forest ($n = 3$), 'high clay–high Fe' under forest ($n = 7$); all cropland combinations ($n = 3$).
204

205 **3. Results**

206 **3.1 Mineralogical composition and general soil properties**

207 The selected mineralogical combinations represent a broad spectrum of possible combinations mineral
208 aluminous clay and Fe oxide constituents. Amounts of aluminous clay varied between 149 and 438 g kg⁻¹,
209 and Fe_d between 21 and 101 g kg⁻¹ across all sites and land uses. Amorphous Fe and Al phases contributed
210 little to pedogenic oxides as indicated by low proportions of oxalate-extractable Fe and Al (Table 1). The
211 advanced weathering state of study soils was also reflected in low pH and CEC_{eff} values (Table 1).

212 **Table 1:** Basic properties of the two soil depth increments sampled along the mineralogical combinations with aluminous clay
 213 (clay), dithionite-citrate-bicarbonate-extractable Fe (Fe_d), total soil organic carbon content (OC), Fe_d to aluminous clay ratios
 214 ($Fe_d/clay$), effective cation exchange capacity (CEC_{eff}), hydrogen peroxide- and dithionite-citrate-bicarbonate-treated sand and silt
 215 contents, and oxalate-extractable Fe and Al content (Fe_o and Al_o). Aluminous clay represents the weight sum of kaolinite and
 216 gibbsite present in the < 2- μ m fraction after removal of OM and pedogenic Fe oxides. Lower case letters indicate significant
 217 differences within a certain land use as separated by depth, and capital letters denote significant differences between land uses.
 218 Sample numbers for the combinations are as follows: ‘low clay–low Fe’ under forest ($n = 4$), ‘low clay–high Fe’ under forest (n
 219 = 4), ‘high clay–low Fe’ under forest ($n = 3$), ‘high clay–high Fe’ under forest ($n = 7$); all cropland combinations ($n = 3$).

Land use	Mineralogical Combination	Depth (cm)	(g kg ⁻¹)				(g kg ⁻¹)				(0.01 M CaCl ₂)		CEC _{eff} (cmolc kg ⁻¹)
			Sand	Silt	Clay	Fe_d	$Fe_d/clay$	Fe_o	Al_o	OC	pH		
Forest	Low aluminous clay–	0–5	788 ^a (21)	63 ^c (24)	149 ^b (19)	21 ^d (4)	0.15 ^b (0.04)	1.4 ^a (0.3)	1.2 ^a (0.2)	76 ^{ab,A} (27)	3.5 ^b (0.1)	5.7 ^a (2.6)	
	Low pedogenic Fe oxides	5–10	712 ^a (46)	107 ^b (57)	181 ^b (19)	38 ^b (13)	0.21 ^{bc} (0.09)	1.8 ^a (0.3)	1.4 ^d (0.2)	34 ^{a,A} (6)	3.7 ^b (0.1)	2.9 ^a (0.1)	
Forest	Low aluminous clay–	0–5	617 ^b (36)	201 ^a (52)	182 ^b (38)	78 ^a (14)	0.45 ^a (0.12)	1.3 ^a (0.2)	1.5 ^a (0.2)	57 ^{b,A} (14)	3.8 ^a (0.2)	5.6 ^a (1.7)	
	High pedogenic Fe oxides	5–10	647 ^b (49)	179 ^a (26)	174 ^b (42)	77 ^a (4)	0.47 ^{ab} (0.13)	1.3 ^b (0.1)	1.6 ^a (0.3)	37 ^{b,A} (7)	3.8 ^{ab} (0.1)	3.2 ^a (0.9)	
Forest	High aluminous clay–	0–5	571 ^c (19)	131 ^b (32)	298 ^a (41)	36 ^c (5)	0.12 ^b (0.01)	0.9 ^b (0.0)	1.3 ^a (0.2)	43 ^b (6)	4.0 ^a (0.2)	5.2 ^a (1.1)	
	Low pedogenic Fe oxides	5–10	489 ^c (24)	137 ^{ab} (1)	374 ^a (24)	44 ^b (7)	0.12 ^c (0.02)	1.0 ^b (0.1)	1.5 ^a (0.3)	23 ^b (5)	3.9 ^{ab} (0.1)	3.0 ^a (0.4)	
Forest	High aluminous clay–	0–5	530 ^c (28)	152 ^b (24)	318 ^a (41)	67 ^b (5)	0.22 ^b (0.03)	1.2 ^{ab} (0.3)	1.9 ^a (0.8)	95 ^{a,A} (31)	4.1 ^a (0.2)	7.8 ^a (1.8)	
	High pedogenic Fe oxides	5–10	473 ^c (35)	178 ^a (45)	349 ^a (40)	81 ^a (6)	0.23 ^b (0.02)	1.3 ^b (0.1)	1.7 ^a (0.2)	35 ^{a,A} (5)	4.0 ^a (0.1)	4.9 ^a (4.0)	
Cropland	Low aluminous clay–	0–5	670 ^a (8)	103 ^c (4)	227 ^b (6)	30 ^c (2)	0.13 ^b (0.01)	0.6 ^c (0.0)	1.1 ^c (0.1)	19 ^{c,B} (0)	5.0 ^b (0.1)	5.1 ^b (0.2)	
	Low pedogenic Fe oxides	5–10	669 ^a (8)	118 ^b (28)	213 ^b (24)	29 ^c (4)	0.14 ^b (0.03)	0.6 ^c (0.0)	1.1 ^b (0.1)	19 ^{c,B} (1)	5.0 ^{ab} (0.1)	5.1 ^b (0.2)	
Cropland	Low aluminous clay–	0–5	602 ^b (17)	200 ^a (13)	198 ^b (29)	101 ^a (4)	0.51 ^a (0.06)	1.5 ^a (0.0)	4.1 ^a (0.2)	47 ^{a,A} (1)	4.9 ^c (0.1)	5.1 ^b (0.2)	

High pedogenic Fe oxides	5–10	579^b (19)	206^a (4)	215^b (23)	100^a (5)	0.47^m (0.07)	1.7^a (0.1)	4.3^a (0.6)	48^{aaA} (5)	4.8^b (0.1)	5.0^b (1.2)
Cropland High aluminous clay–	0–5	437^c (14)	129^b (12)	434^a (18)	63^b (3)	0.15^b (0.01)	1.2^b (0.0)	1.4^b (0.0)	34^{bbB} (1)	5.4^a (0.0)	9.4^a (0.5)
High pedogenic Fe oxides	5–10	399^c (18)	163^{ab} (35)	438^a (17)	66^b (4)	0.15^b (0.01)	1.2^b (0.1)	1.3^b (0.2)	30^{baA} (3)	5.2^a (0.1)	7.3^a (0.7)

221 **3.2 Influence of aluminous clay and pedogenic Fe on aggregate size distribution**


222 *Mean weight diameter*

223 The studied soils were highly aggregated and showed significant variation in their aggregate size distribution
224 across the mineralogical combinations (Figure 1a, Table 2). The low clay–low Fe combination under forest
225 displayed the significant smallest MWD (e.g., 2.9 mm in 0–5 cm depth; Table 2). In contrast, the low clay–
226 high Fe combination always had the largest MWD (e.g., 4.8 mm in 0–5 cm depth; Table 2) among the other
227 forest combinations. Our data suggest that the MWD under forest is significantly positively influenced by
228 the Fe_d content (e.g., $MWD_{Forest\ 0-5\ cm}: r^2 = 0.40, p < 0.001$; Table S1), whereas nearly no effect was observed
229 for aluminous clay. Contrary to the mineralogical combinations under forest, the significant smallest MWD
230 under cropland was within the low clay–high Fe combination (2.7 mm in both depths; Table 2). The low
231 clay–low Fe and high clay–high Fe cropland combinations showed no strong differences in their MWDs.
232 Nonetheless, a significant negative linear relationship existed between MWD and the pedogenic-Fe to
233 aluminous clay ratio ($MWD_{Cropland\ 0-5\ cm}: r^2 = 0.47, p = 0.03$; $MWD_{Cropland\ 5-10\ cm}: r^2 = 0.47, p = 0.02$) for the
234 mineralogical combinations under cropland (Table S1).

235

236 *Macroaggregates > 4 mm and 2–4 mm*

237 Corresponding to the smallest MWD, the low clay–low Fe forest combination contained the smallest
238 fraction of > 4 mm aggregates. The contribution of these large aggregates under forest increased in the order:
239 low clay–low Fe < low clay–high Fe = high clay–high Fe < high clay–low Fe (Figure 1a). For croplands,
240 the low clay–high Fe combination comprised the smallest amount of > 4 mm aggregates whereas the high
241 clay–high Fe combination exhibited the respective highest share (Figure 1a). The explained variance of
242 > 4 mm aggregate mass by aluminous clay and Fe_d was generally low, except for the cropland combinations
243 (positive effect of aluminous clay and negative effect of pedogenic Fe; Table S1).



244 **Figure 1:** Aggregate size distribution of the combined 0–5 and 5–10 cm depth increments (a), and relative
 245 mass-corrected OC contents (b) along the mineralogical combinations. Clay represents the weight sum of
 246 kaolinite and gibbsite present in the < 2- μ m fraction after removal of OM and pedogenic Fe oxides, and Fe
 247 denotes the content of pedogenic Fe oxides extracted with dithionite-citrate-bicarbonate. Sample numbers
 248 for the combinations are as follows: ‘low clay–low Fe’ under forest ($n = 4$), ‘low clay–high Fe’ under forest
 249 ($n = 4$), ‘high clay–low Fe’ under forest ($n = 3$), ‘high clay–high Fe’ under forest ($n = 7$); all cropland
 250 combinations ($n = 3$).
 251
 252

253 The mineralogical combinations affected the amounts of 2–4 mm aggregates differently than those
 254 of > 4 mm aggregates. The low clay–high Fe combination under forest and cropland contained slightly but
 255 significantly more 2–4 mm aggregates (Figure 1a), being associated with a significantly higher Fe_d to
 256 aluminous clay ratio (Table 1). In fact, in a multiple regression model for the entire data set (combined land
 257 uses and depths), we observed a positive relationship between the mass of 2–4 mm aggregates and Fe_d
 258 content, whereas the content of aluminous clay had a negative effect ($r^2 = 0.57$, $p < 0.001$; Table S1).
 259

260 *Microaggregates < 0.25 mm*

261 Across all mineralogical combinations, amounts of < 0.25 mm aggregates were principally similar, despite
 262 of significantly higher shares in the low clay–low Fe and high clay–high Fe combinations under forest. A
 263 significant larger amount of < 0.25 mm aggregates was observed in the low clay–high Fe combination under
 264 cropland. In this mineralogical combination, land-use change caused a quadrupling of < 0.25 mm aggregate

265 mass from about 30 to nearly 120 g kg⁻¹ (Table 2). In contrast to the macroaggregate fractions shown above,
266 there was no correlation between mineralogical parameters and the mass of < 0.25 mm aggregates, neither
267 for the entire data set (combined land uses and depths) nor when separated by soil depth (Table S1).

Table 2: Aggregate masses (mass) and OC content of aggregate size fractions (dry sieving) within different combinations of aluminous clay and pedogenic Fe oxides, OC change (Δ OC) between land uses within a certain mineralogical combination and depth, and related mean weight diameter (MWD). Aluminous clay represents the weight sum of kaolinite and gibbsite present in the $< 2\text{-}\mu\text{m}$ fraction after removal of OM and pedogenic Fe oxides. Lower case letters indicate significant differences within a certain land use separated by depth, and capital letters denote significant differences between land uses. Sample numbers for the combinations are as follows: 'low clay–low Fe' under forest ($n = 4$), 'low clay–high Fe' under forest ($n = 4$), 'high clay–low Fe' under forest ($n = 3$), 'high clay–high Fe' under forest ($n = 7$); all cropland combinations ($n = 3$).

Land use	Mineralogical Combination	Depth (cm)	> 4 mm (g kg ⁻¹)			2–4 mm (g kg ⁻¹)			1–2 mm (g kg ⁻¹)			0.25–1 mm (g kg ⁻¹)			< 0.25 mm (g kg ⁻¹)			
			mass	OC	Δ OC (%)	mass	OC	Δ OC (%)	mass	OC	Δ OC (%)	mass	OC	Δ OC (%)	mass	OC	Δ OC (%)	MWD (mm)
Forest	Low aluminous clay–pedogenic Fe oxides	0–5	249 ^{a,A} (33)	76 ^{a,A} (32)	na	144 ^{b,A} (21)	83 ^{a,A} (22)	na	191 ^{a,A} (4)	65 ^{a,A} (9)	na	345 ^{a,A} (40)	56 ^{ab,A} (18)	na	70 ^{a,A} (15)	125 ^{ab,A} (51)	na	2.9 ^{c,A} (0.3)
		5–10	343 ^{b,A} (61)	40 ^{a,A} (8)	na	176 ^{ab,A} (21)	39 ^{a,A} (10)	na	181 ^{a,A} (15)	27 ^{a,A} (9)	na	257 ^{a,A} (36)	28 ^{a,A} (5)	na	44 ^{a,B} (11)	51 ^{a,A} (17)	na	3.7 ^{a,A} (0.4)
Forest	Low aluminous clay–High pedogenic Fe oxides	0–5	493 ^{ab,A} (99)	68 ^{ab,A} (19)	na	210 ^{b,A} (20)	65 ^{a,A} (22)	na	115 ^{b,B} (38)	62 ^{a,A} (25)	na	150 ^{c,B} (42)	49 ^{b,A} (25)	na	33 ^{b,B} (14)	62 ^{b,A} (36)	na	4.8 ^{a,A} (0.7)
		5–10	451 ^{ab,A} (36)	40 ^{a,A} (11)	na	210 ^{b,A} (27)	36 ^{ab,B} (5)	na	139 ^{ab,B} (10)	29 ^{a,A} (7)	na	166 ^{b,B} (24)	31 ^{a,A} (11)	na	34 ^{a,B} (20)	44 ^{a,A} (18)	na	4.6 ^{a,A} (0.3)
Forest	High aluminous clay–Low pedogenic Fe oxides	0–5	604 ^a (84)	38 ^b (5)	na	140 ^b (21)	63 ^a (34)	na	100 ^b (21)	80 ^a (51)	na	125 ^c (31)	62 ^{ab} (28)	na	31 ^b (13)	101 ^{ab} (59)	na	4.3 ^{ab} (0.4)
		5–10	561 ^a (47)	26 ^a (14)	na	163 ^b (12)	28 ^b (7)	na	118 ^b (17)	22 ^a (3)	na	127 ^b (21)	25 ^a (6)	na	30 ^a (1)	43 ^a (18)	na	4.1 ^a (0.2)
Forest	High aluminous clay–High pedogenic Fe oxides	0–5	397 ^{b,A} (91)	86 ^{a,A} (21)	na	157 ^{b,A} (27)	89 ^{a,A} (32)	na	163 ^{a,A} (32)	99 ^{a,A} (50)	na	208 ^{b,B} (36)	91 ^{a,A} (38)	na	74 ^{a,A} (14)	133 ^{a,A} (47)	na	4.0 ^{b,A} (0.6)
		5–10	474 ^{ab,A} (139)	35 ^{a,A} (7)	na	156 ^{b,A} (27)	33 ^{ab,A} (4)	na	146 ^{ab,A} (41)	30 ^{a,A} (4)	na	172 ^{b,A} (61)	34 ^{a,A} (4)	na	52 ^{a,A} (26)	51 ^{a,A} (6)	na	4.6 ^{a,A} (1.0)
Cropland	Low aluminous clay–Low pedogenic Fe oxides	0–5	347 ^{a,A} (69)	20 ^{b,B} (3)	-73	147 ^{b,A} (13)	21 ^{c,B} (1)	-75	115 ^{b,B} (4)	17 ^{c,B} (1)	-74	318 ^{a,A} (52)	11 ^{c,B} (3)	-80	74 ^{b,A} (12)	24 ^{c,B} (1)	-81	3.6 ^{a,A} (0.5)
		5–10	368 ^{b,A} (28)	20 ^{b,B} (1)	-50	143 ^{b,A} (8)	22 ^{b,B} (5)	-44	113 ^{b,B} (10)	17 ^{b,A} (2)	-37	299 ^{a,A} (15)	11 ^{c,B} (2)	-61	77 ^{b,A} (1)	24 ^{c,A} (3)	-53	3.7 ^{b,A} (0.2)
Cropland	Low aluminous clay–	0–5	201 ^{b,B} (39)	47 ^{a,A} (7)	-30	212 ^{a,A} (12)	49 ^{a,A} (2)	-25	173 ^{a,A} (18)	42 ^{a,A} (3)	-32	296 ^{a,A} (33)	46 ^{a,A} (1)	-6	119 ^{a,A} (4)	62 ^{a,A} (2)	±0	2.7 ^{b,B} (0.3)

High pedogenic Fe oxides	5–10	194^{c,B} (11)	47^{a,A} (13)	+18	224^{a,A} (15)	49^{a,A} (4)	+36	177^{a,A} (1)	42^{a,A} (6)	+45	287^{a,A} (13)	45^{a,A} (3)	+45	118^{a,A} (29)	58^{a,A} (9)	+32	2.7^{c,B} (0.1)
Cropland High aluminous clay–	0–5	296^{b,A} (40)	26^{b,B} (6)	-71	159^{b,A} (8)	29^{b,B} (7)	-67	191^{a,A} (2)	28^{b,B} (4)	-71	278^{a,A} (25)	35^{b,A} (2)	-62	77^{b,A} (10)	41^{b,B} (1)	-69	3.3^{b,A} (0.3)
High pedogenic Fe oxides	5–10	593^{a,A} (95)	25^{b,A} (3)	-29	118^{b,A} (21)	26^{b,B} (2)	-21	107^{b,A} (29)	25^{b,A} (4)	-17	138^{b,A} (37)	32^{b,A} (3)	-6	43^{b,A} (10)	41^{b,B} (5)	-20	5.3^{a,A} (0.6)

na = not applicable.

274

275

276 In summary, mineralogical combinations and land use significantly affected the aggregate size distribution
277 of soils, despite quantitative relations to mineralogical proxies could not be observed for each aggregate
278 class. In undisturbed forest soils, higher pedogenic Fe contents resulted in increasing MWD especially in
279 0–5 cm depth and significantly larger amounts of > 2 mm aggregates. The conversion from forest to
280 croplands either decreased MWD, as particularly observed for the low clay–high Fe combination, or had no
281 effect (low clay–low Fe). Overall, the observed differences in aggregate masses and MWD were surprisingly
282 moderate, given the widely differing contents in aluminous clay and Fe oxides across the mineralogical
283 combinations.

284

285 **3.3 Aggregate stability**

286 In general, there was little variation of MWD values for > 4 mm aggregates over all mineralogical
287 combinations. In fact, the MWD of this fraction was always close to its calculated mean diameter (6 mm;
288 calculation was done after (*Youker and McGuinness, 1957*)), overall indicating a high stability.
289 Nevertheless, there were some minor differences in aggregate stability across mineralogical combinations.
290 The low clay–low Fe and high clay–low Fe combinations had a significantly lower aggregate stability in
291 comparison with the two other combinations under the two land uses (Table 3). The slightly higher
292 abundance of 2–4 mm aggregates in the low clay–high Fe combination under forest and cropland was
293 accompanied by a significantly higher aggregate stability under both land uses (Table 2 and 3). In summary,
294 all aggregates can be classified as stable with only minor differences imposed by the mineralogical
295 combinations. Slightly higher aggregate stability was associated with a larger amount of pedogenic Fe, and
296 increasing Fe_d to aluminous clay ratios, whereas differences in the amount of aluminous clay had almost no
297 effect on the aggregate stability (Table S2).

298

299

300

301

302 **Table 3:** Aggregate stability of selected aggregate size fractions after applying the fast wetting procedure
 303 along the different combinations of aluminous clay and pedogenic Fe oxides, indicated by the resulting
 304 mean weight diameter (MWD). Aluminous clay represents the weight sum of kaolinite and gibbsite present
 305 in the < 2- μ m fraction after removal of OM and pedogenic Fe oxides. Lower case letters indicate significant
 306 differences within a certain land use separated by depth, and capital letters denote significant differences
 307 between land uses. Sample numbers for the combinations are as follows: 'low clay–low Fe' under forest (n
 308 = 4), 'low clay–high Fe' under forest (n = 4), 'high clay–low Fe' under forest (n = 3), 'high clay–high Fe'
 309 under forest (n = 7); all cropland combinations (n = 3).

Land use	Mineralogical combination	Depth (cm)	MWD	
			Fast wetting > 4 mm (mm)	Fast wetting 2–4 mm (mm)
Forest	Low aluminous clay–	0–5	4.9 ^{b, A} (0.4)	2.6 ^{b, A} (0.1)
	Low pedogenic Fe oxides	5–10	5.1 ^{a, A} (0.3)	2.4 ^{b, A} (0.3)
Forest	Low aluminous clay–	0–5	5.6 ^{a, A} (0.2)	2.8 ^{a, A} (0.1)
	High pedogenic Fe oxides	5–10	4.9 ^{a, A} (0.9)	2.7 ^{a, A} (0.1)
Forest	High aluminous clay–	0–5	5.4 ^{ab} (0.4)	2.7 ^b (0.0)
	Low pedogenic Fe oxides	5–10	4.5 ^a (1.2)	2.4 ^b (0.3)
Forest	High aluminous clay–	0–5	5.5 ^{a, A} (0.2)	2.6 ^{b, A} (0.1)
	High pedogenic Fe oxides	5–10	5.2 ^{a, A} (0.4)	2.6 ^{ab, B} (0.1)
Cropland	Low aluminous clay–	0–5	4.4 ^{b, A} (0.1)	2.6 ^{c, A} (0.0)
	Low pedogenic Fe oxides	5–10	4.9 ^{b, A} (0.3)	2.4 ^{b, A} (0.1)
Cropland	Low aluminous clay–	0–5	5.2 ^{a, A} (0.2)	2.9 ^{a, A} (0.0)
	High pedogenic Fe oxides	5–10	5.3 ^{ab, A} (0.1)	2.8 ^{a, A} (0.0)
Cropland	High aluminous clay–	0–5	4.9 ^{a, B} (0.2)	2.7 ^{b, A} (0.1)
	High pedogenic Fe oxides	5–10	5.6 ^{a, A} (0.2)	2.8 ^{a, A} (0.0)

310

311 **3.4 Organic carbon in soils and aggregate size fractions**


312 In the entire data set, variation in mineral constituents caused pronounced differences in the OC content of

313 the soils between 19 to 95 g OC kg⁻¹ (Table 1). A significant proportion of the total OC content of all forest

314 soils was present in > 4 mm aggregates in both depth increments (low clay–low Fe: 33% < high clay–high

315 Fe: 43% < high clay–low Fe: 51% < low clay–high Fe: 52%; Figure 1b). Forest to cropland conversion
 316 caused OC losses from most aggregate size fractions (Figure 2). For the > 4 mm aggregates this was
 317 significantly modified by the mineralogical combinations at least at 0–5 cm depth, generally following the
 318 order: low clay–high Fe < high clay–high Fe < low clay–low Fe (Table S3). Losses of OC from aggregate
 319 size fractions were generally higher at 0–5 than at 5–10 cm depth (Figure 2). As mentioned above, no
 320 significant loss of total OC occurred for the low clay–high Fe combination, irrespective of the significant
 321 mass redistribution of the > 4 mm aggregate fraction into smaller aggregate fractions (Table 2). Although
 322 there were differences in OC losses among mineralogical combinations, there was little indication that
 323 coarser aggregate size fractions lost more OC than smaller ones (Table 2).

324



325 **Figure 2:** Mass-corrected OC contents of aggregate size fractions along the mineralogical combinations.
 326 Clay represents the weight sum of kaolinite and gibbsite present in the < 2- μ m fraction after removal of OM
 327 and pedogenic Fe oxides, and Fe denotes the content of pedogenic Fe oxides extracted with dithionite-
 328 citrate-bicarbonate. Sample numbers for the combinations are as follows: 'low clay–low Fe' under forest (*n*
 329

330 = 4), 'low clay–high Fe' under forest ($n = 4$), 'high clay–low Fe' under forest ($n = 3$), 'high clay–high Fe'
331 under forest ($n = 7$); all cropland combinations ($n = 3$).
332

333 **4. Discussion**

334 The aggregate size distribution of soils along the mineralogical combinations under both land uses were in
335 the range of values reported for African soils. For example, soils with strongly contrasting clay content (220
336 and 650 g kg⁻¹) but similar clay mineralogy (kaolinite) in the central highlands of Kenya displayed
337 macroaggregate contents of 245 and 636 g kg⁻¹ soil, respectively (*Gentile et al.*, 2010). In addition, for soils
338 from the catchment of the Riru river also located in the central highlands of Kenya it was shown that
339 macroaggregates (2–4.2 mm) displayed a large stability (*Kamamia et al.*, 2021). The reported MWD's after
340 application of the fast-wetting stability test were 2.5 mm for cropland and 3.2 mm for indigenous forest sites
341 (*Kamamia et al.*, 2021). These values are close to those observed in our study soils for 2–4 mm aggregates.
342 In contrast, soils in Brazil under native forest vegetation and similar mineral composition (kaolinite,
343 gibbsite, hematite) even subsumed over 90% of total aggregate mass in > 2 mm aggregates (*Maltoni et al.*,
344 2017). Nonetheless, reported data all point at a better soil structure and aggregate stability of tropical soils
345 dominated by low-activity clay minerals and well-crystalline Fe oxides, which is consistent with all
346 mineralogical combinations of this study.

347

348 **4.1 Aggregation and aggregate stability as controlled by aluminous clay and pedogenic Fe oxides**

349 Our data demonstrates relatively small differences in aggregation among the generally well-aggregated
350 study soils, being characterized by high aggregate stability despite of large variations in aluminous clay
351 (factor three) and pedogenic Fe (factor five) contents. Yet, we noticed some distinct modifications of the
352 aggregation size distribution and aggregate stability in both forest and cropland soils.

353

354 *Mineralogical control on the formation of large macroaggregates*

355 The low clay–low Fe soil under forest had a significantly smaller amount of > 4 mm and 2–4 mm aggregates
356 and a significantly lower MWD than all other mineralogical combinations. Notably, a combined increase in
357 aluminous clay and Fe oxides did not necessarily cause a shift towards larger aggregates and thus higher
358 MWD (see low clay–high Fe forest). Furthermore, the low clay–low Fe and high clay–high Fe combinations

359 under forest contained more < 0.25 mm aggregates. Thus, under undisturbed soil conditions it appears that
360 the formation of larger aggregates is promoted if one of the two aggregate-forming mineral fractions is more
361 abundant than the other (high clay–low Fe and low clay–high Fe combinations).

362 We assume that the positive effect of increasing aluminous clay content on the aggregate mass
363 > 4 mm is related to the hybrid electrostatic properties of kaolinite on edges (variable) and surfaces
364 (permanent negative), which enable the formation of characteristic cards-house structures (*Qafoku* and
365 *Sumner*, 2002). In addition to this increase in aggregation caused by the dominance in kaolinitic properties
366 (i.e. high clay–low Fe), we also expect that, similar to the study by *Dultz et al.* (2019), there are mixing
367 ratios between aluminous clay and pedogenic Fe minerals, which lead to improved aggregation (greater
368 MWD; i.e. low clay–high Fe). This effect is probably explained by changes in the electrostatic properties
369 of the mineralogical combinations, as was shown in the study by *Hou et al.* (2007) for kaolinite in different
370 relative combinations with goethite and hematite. Nevertheless, aluminous clay is the decisive control for
371 macroaggregation in these weathered tropical soils, confirming the often described promoting effect of
372 increasing clay content on aggregation (*Feller and Beare*, 1997). This is in line with results from two Oxisols
373 in Brazil (*Vrdoljak and Sposito*, 2002), showing kaolinite being the backbone of macroaggregates.

374 Consequently, the dominant role of pedogenic Fe oxides for macroaggregation under undisturbed
375 tropical soil conditions proposed by *Six et al.* (2002) cannot be confirmed in our study. This is also supported
376 by the low clay–high Fe forest soil, which contained a smaller amount of > 4 mm aggregates compared to
377 the high clay–low Fe forest soil in both depth increments. Furthermore, the high clay–low Fe and high clay–
378 high Fe combinations under forest also nicely demonstrate how nearly equal amounts of aluminous clay
379 plus pedogenic Fe oxides (i.e. similar clay contents) cause different amounts of > 4 mm aggregates.
380 Consequently, the connection between textural properties and aggregation can remain hidden (*Barthès et*
381 *al.*, 2008) without considering the mineralogical composition of the whole clay fraction (*Fernández-Ugalde*
382 *et al.*, 2013; *King et al.*, 2019; *West et al.*, 2004).

383 In contrast to the > 4 mm aggregates, 2–4 mm aggregates corresponded more clearly to the positive
384 effect of pedogenic Fe oxides on aggregation and aggregate stability as proposed for weathered tropical

385 soils (Igwe et al., 2013; Peng et al., 2015; Six et al., 2002). Both, the low clay–high Fe forest and low clay–
386 high Fe cropland soils contained somewhat but significantly more 2–4 mm aggregates than other mineral
387 combinations in concert with a higher aggregate stability of this particular fraction. This finding also
388 demonstrates that mineral interactions forming water-stable aggregates in tropical soils are differently
389 affected by a given mineralogical combination. Higher Fe_d to aluminous clay ratios (> 0.45) modulate
390 aggregate distribution towards aggregates 2–4 mm, whereas distinctly lower values (high clay–low Fe
391 forest: 0.12) shifted the maximum to > 4 mm aggregates.

392 Overall, the two macroaggregate fractions discussed above are differentially affected by the
393 mineralogical combinations, although the magnitude was less than expected, given the pronounced variation
394 in aluminous clay and Fe contents.

395

396 *Land use impact on aggregation within mineral combinations– implications for aggregate stability*

397 Land-use change had a distinct impact on aggregate distribution like indicated in other studies (Feller and
398 Beare, 1997; Six et al., 2002) and depended also on the mineralogical combinations, though croplands not
399 followed the trajectory observed under forest. A significantly lower MWD under low clay–high Fe rather
400 than low clay–low Fe can be mainly attributed to a reduced amount of > 4 mm aggregates. We assume that
401 differences in the ratio of pedogenic Fe to aluminous clay in the low clay–low Fe and high clay–high Fe
402 (0.13 to 0.15) in comparison with the low clay–high Fe combination (0.47 to 0.51) under cropland explains
403 the stability of ‘card-house’ structures like described for mineralogically similar Oxisols from Brazil and
404 India (Bartoli et al., 1992). Accordingly, a higher Fe_d to aluminous clay ratios seems to be disadvantageous
405 for the formation and stability of such structures, especially in > 4 mm aggregates. The different pH-
406 dependent charge characteristics of kaolinite and pedogenic Fe oxides (Kaiser and Guggenberger, 2003),
407 and their relative share can lead to altered charge properties of soils (Anda et al., 2008). We hypothesize,
408 that an increasing amount of Fe oxides adds more positive charge, thus possibly reducing structural integrity
409 and aggregate stability if not sufficiently compensated by OM or clay minerals. Furthermore, in the low

410 clay–high Fe cropland combination, land-use change caused a significant four-fold increase of < 0.25 mm
411 aggregates due to the breakdown of > 4 mm aggregates.

412 The less intense formation of > 4 mm aggregates in the low clay–high Fe forest combination was also
413 observed under cropland, whereas the low clay–low Fe and high clay–high Fe croplands showed either no
414 significant decrease or even an increase in > 4 mm aggregate mass. Thus, simultaneous abundance of large
415 amounts of aluminous clay and pedogenic Fe oxides preserved a higher aggregate stability than under
416 mineralogically imbalanced conditions, although no conclusions can be drawn for the high clay–low Fe
417 combination. Nonetheless, > 4 mm aggregates had a higher resistance to field operations in mineralogical
418 combinations with lower Fe_d to aluminous clay ratios (0.13 to 0.15). Nonetheless, our results show that
419 agricultural management does not necessarily decreases macroaggregation and related MWD's, like
420 reported in *Rabbi et al. (2015)*.

421
422 **4.2 Importance of aggregation for OC persistence – effects of aluminous clay and pedogenic Fe oxides**

423 Clay minerals and Fe oxides are considered as important mineral constituents fostering aggregation and
424 subsequent OC storage via physical protection (*Denef et al., 2004*). The overwhelming portion of OC in the
425 studied topsoils resided in mineral-organic associations (35–81%), whereas OC occluded in aggregates
426 amounted to 7–24%, with a lower share under cropland than forest as determined by density fractionation
427 (*Kirsten et al., 2021*). The low clay–high Fe cropland had an OC content more than twice larger than that of
428 the low clay–low Fe cropland, but comprised a significantly smaller MWD. Thus, a shift towards more
429 macroaggregation, indicated by a larger MWD in certain mineralogical combinations, did not result in
430 higher total OC storage, like shown for other tropical soils (*Barthès et al., 2008; Bartoli et al., 1991; Spaccini*
431 *et al., 2001*). The OC content of the > 4 mm aggregate and 2–4 mm aggregate fractions accounted for 42 to
432 73% of the total soil OC content (Figure 1b). This, however, does not *per se* indicate the relevance of
433 macroaggregation for OC storage in weathered tropical soils like proposed by others (*Feller and Beare,*
434 *1997; King et al., 2019; Six et al., 2002*). The high clay–low Fe forest with the highest share in > 4 mm and
435 2–4 mm aggregates had significant lower OC contents in these fractions than most other mineralogical

436 combinations. Comparing forest with cropland soils (Table 2), we observed significantly reduced OC
437 contents in the majority of macroaggregate fractions of the low clay–low Fe and high clay–high Fe
438 croplands, as reported in other studies (*Blanco-Canqui and Lal, 2004; Lobe et al., 2011*). In contrast, fewer
439 changes of aggregate-associated and total soil OC contents was observed in the low clay–high Fe
440 combination, despite it experienced the strongest disaggregation of the largest macroaggregates (Figure 1a
441 and Figure 2). We conclude that larger amounts of > 2 mm aggregates or higher stability during wet sieving
442 does not automatically translate into higher aggregate-associated OC contents, as reported for Ferralsols
443 (*Maltoni et al., 2017*). Given all these observations and the fact that occluded OM determined by density
444 fractionation was mostly of subordinate relevance, particularly in croplands, OC storage in study soils seems
445 rather disconnected from their aggregation status. Consequently, the loss of large aggregates and the mass
446 redistribution into smaller aggregate size fractions does not automatically imply a loss of soil OC, because
447 a substantial part of the OC in aggregate fractions is bound to minerals with a higher persistence against
448 land-use change (*Kirsten et al., 2021*). Here, density fractionation could shed more light on the nature and
449 quantity of OM located in certain aggregate size fractions.

450 Microaggregates contained the highest OC content per unit of mass for almost all mineralogical
451 combinations, depth increments, and land uses (Table 2). This is in line with the findings of *Chenu and*
452 *Plante (2006)* and *Lobe et al. (2011)* that microaggregates can significantly contribute to OC storage. As
453 aggregates were isolated by dry sieving, these microaggregates were not located inside larger aggregates,
454 rendering them principally better accessible for OC allocation. Particularly OC contained in the < 0.25 mm
455 aggregates of the low clay–high Fe combination revealed a strong persistence against land-use change,
456 which explains well the unaltered soil OC contents upon land-use change.

457

458 **5. Conclusions**

459 Classification of soils into mineralogical combinations of aluminous clay and pedogenic Fe oxides revealed
460 significant effects of mineral constituents on soil structure and related OC storage in weathered tropical
461 soils. Despite that, overall patterns across combinations were more similar than different, *i.e.*, always
462 comprising a high level of macroaggregation and aggregate stability. Aggregates > 4 mm of the low clay–
463 low Fe and high clay–high Fe combinations were less affected by land-use change, thus pedogenic Fe in a
464 certain relation with aluminous clay (0.13 to 0.23) seems beneficial to maintain the structural integrity of
465 macroaggregates. Despite the high physical stability, OC contents of macroaggregates declined substantially
466 in most mineralogical combinations if forest was compared with cropland land use. This highlights the fact
467 that structural integrity of macroaggregates during land-use change cannot be equated with OC persistence.
468 For the low clay–high Fe combination, substantial destruction of > 4 mm aggregates during land-use change
469 due to agricultural management was also not accompanied by higher OC losses. Hence, we must reject our
470 initial hypothesis that the mineralogical combination that results in the greatest aggregate stability best
471 preserves OC during the conversion from forest to cropland. Thus, the formation of macroaggregates cannot
472 be considered as a main stabilization process for OC in strongly weathered soils of the humid tropics. We
473 suggest that the formation of mineral-organic associations as part of the aggregate size fractions is the most
474 important process that preserves OC during land-use change in these soils.

475 **7. Author contribution**

476 KK, RM, MK, and KHF designed the project. MK, KK, RM, DNK, and KHF collected soil or data to
477 supported the sampling campaign. MK, KK, RM, and KHF evaluated data and all authors conducted a
478 thorough critical review of the manuscript. MK, KK, and RM wrote the manuscript with contribution of all
479 authors.

480

481 **8. Competing interests**

482 The authors declare that they have no conflict of interest.

483

484 **9. Acknowledgements**

485 We are grateful to the officials of Amani Nature Reserve who supported the field campaign in February
486 2018. Aloyce Mkongewa enthusiastically assisted fieldwork. We are also indebted to Gisela Ciesielski,
487 Manuela Unger, Mandy Meise, Tobias Krause, Thomas Klinger, Gudrun Nemson-von Koch, and Christine
488 Krenkewitz for laboratory support and analytical work. Finally, the authors would like to thank the two
489 reviewers for their comprehensive reading of the manuscript and constructive suggestions for its
490 improvement. This study was supported by grants of the Deutsche Forschungsgemeinschaft (DFG): FE
491 504/15-1, KA 1737/16-1, and MI 1377/11-1.

492 **10. References**

- 493 *Anda, M., Shamshuddin, J., Fauziah, C. I., Omar, S. S.* (2008): Mineralogy and factors controlling charge
494 development of three Oxisols developed from different parent materials. *Geoderma* 143, 153–167.
- 495 *Asano, M., Wagai, R.* (2014): Evidence of aggregate hierarchy at micro- to submicron scales in an allophanic
496 Andisol. *Geoderma* 216, 62–74.
- 497 *Barthès, B. G., Kouakoua, E., Larré-Larrouy, M.-C., Razafimbelo, T. M., de Luca, Edgar F., Azontonde, A.,*
498 *Neves, C. S.V.J., de Freitas, Pedro L., Feller, C. L.* (2008): Texture and sesquioxide effects on water-
499 stable aggregates and organic matter in some tropical soils. *Geoderma* 143, 14–25.
- 500 *Bartoli, F., Burtin, G., Herbillon, A. J.* (1991): Disaggregation and clay dispersion of Oxisols: Na resin, a
501 recommended methodology. *Geoderma* 49, 301–317.
- 502 *Bartoli, F., Philippy, R., Burlin, G.* (1992): Influence of organic matter on aggregation in Oxisols rich in
503 gibbsite or in goethite. I. Structures: the fractal approach. *Geoderma* 54, 231–257.
- 504 *Blanco-Canqui, H., Lal, R.* (2004): Mechanisms of Carbon Sequestration in Soil Aggregates. *CRC Crit.*
505 *Rev. Plant Sci.* 23, 481–504.
- 506 *Bronick, C. J., Lal, R.* (2005): Soil structure and management: a review. *Geoderma* 124, 3–22.
- 507 *Carter, M. R., Gregorich, E. G.* (Eds.) (2008): Soil sampling and methods of analysis. Canadian Society of
508 Soil Science; CRC Press, [Pinawa, Manitoba], Boca Raton, FL, USA, p. 1224.
- 509 *Chaplot, V., Bouahom, B., Valentin, C.* (2010): Soil organic carbon stocks in Laos: spatial variations and
510 controlling factors. *Glob. Change Biol.* 16, 1380–1393.
- 511 *Chaplot, V., Cooper, M.* (2015): Soil aggregate stability to predict organic carbon outputs from soils. *Ge-*
512 *oderma* 243-244, 205–213.
- 513 *Chenu, C., Plante, A. F.* (2006): Clay-sized organo-mineral complexes in a cultivation chronosequence:
514 revisiting the concept of the 'primary organo-mineral complex'. *Eur. J. Soil Sci.* 57, 596–607.
- 515 *Denef, K., Six, J., Merckx, R., Paustian, K.* (2002): Short-term effects of biological and physical forces on
516 aggregate formation in soils with different clay mineralogy. *Plant Soil* 246, 185–200.
- 517 *Denef, K., Six, J., Merckx, R., Paustian, K.* (2004): Carbon Sequestration in Microaggregates of No-Tillage
518 Soils with Different Clay Mineralogy. *Soil Sci. Soc. Am. J.* 68, 1935.
- 519 *Don, A., Schumacher, J., Freibauer, A.* (2011): Impact of tropical land-use change on soil organic carbon
520 stocks - a meta-analysis. *Glob. Change Biol.* 17, 1658–1670.
- 521 *Duiker, S. W., Rhoton, F. E., Torrent, J., Smeck, N. E., Lal, R.* (2003): Iron (Hydr)Oxide Crystallinity Effects
522 on Soil Aggregation. *Soil Sci. Soc. Am. J.* 67, 606.
- 523 *Dultz, S., Woche, S. K., Mikutta, R., Schrapel, M., Guggenberger, G.* (2019): Size and charge constraints in
524 microaggregation: Model experiments with mineral particle size fractions. *Applied Clay Science* 170, 29–
525 40.
- 526 *Durn, G., Škapin, S. D., Vdović, N., Rennert, T., Ottner, F., Ružičić, S., Cukrov, N., Sondi, I.* (2019): Impact
527 of iron oxides and soil organic matter on the surface physicochemical properties and aggregation of Terra
528 Rossa and Calcocambisol subsoil horizons from Istria (Croatia). *Catena* 183, 104184.
- 529 *FAO* (2015): Status of the World's Soil Resources: Main report. FAO; ITPS, Rome, XXXIX, 608 str.
- 530 *Feller, C., Beare, M. H.* (1997): Physical control of soil organic matter dynamics in the tropics. *Geoderma*
531 79, 69–116.

532 *Fernández-Ugalde, O., Barré, P., Hubert, F., Virto, I., Girardin, C., Ferrage, E., Caner, L., Chenu, C.*
533 (2013): Clay mineralogy differs qualitatively in aggregate-size classes: clay-mineral-based evidence for
534 aggregate hierarchy in temperate soils. *Eur. J. Soil Sci.* 64, 410–422.

535 *Gee, G.W., Bauder, J.W.* (1986): Particle-size analysis, in Klute, A., Page, A. L.: *Methods of soil analysis:*
536 *Part 1 Physical and mineralogical methods.* 2nd ed. no. 9. American Society of Agronomy; Soil Science
537 Society of America, Madison, pp. 383–412.

538 *Gentile, R., Vanlauwe, B., Kavoo, A., Chivenge, P., Six, J.* (2010): Residue quality and N fertilizer do not
539 influence aggregate stabilization of C and N in two tropical soils with contrasting texture. *Nutr. Cycling*
540 *Agroecosyst. (Nutrient Cycling in Agroecosystems)* 88, 121–131.

541 *Hamilton, A. C., Bensted-Smith, R.* (1989): *Forest Conservation in the East Usambara Mountains, Tanzania.*
542 *IUCN - The World Conservation Union, Gland, Switzerland.*

543 *Hou, T., Xu, R., Zhao, A.* (2007): Interaction between electric double layers of kaolinite and Fe/Al oxides in
544 suspensions. *Colloids and Surfaces A: Physicochemical and Engineering Aspects* 297, 91–94.

545 *Igwe, C. A., Zarei, M., Stahr, K.* (2013): Stability of aggregates of some weathered soils in south-eastern
546 Nigeria in relation to their geochemical properties. *J. Earth Syst. Sci.* 122, 1283–1294.

547 *Kaiser, K., Guggenberger, G.* (2003): Mineral surfaces and soil organic matter. *Eur. J. Soil Sci.* 54, 219–
548 236.

549 *Kamamia, A. W., Vogel, C., Mwangi, H. M., Feger, K.-H., Julich, S.* (2021): Mapping soil aggregate stability
550 using digital soil mapping: A case study of Ruiru reservoir catchment, Kenya. *Geoderma Regional* 24.

551 *King, A. E., Congreves, K. A., Deen, B., Dunfield, K. E., Voroney, R. P., Wagner-Riddle, C.* (2019): Quan-
552 tifying the relationships between soil fraction mass, fraction carbon, and total soil carbon to assess mech-
553 anisms of physical protection. *Soil Biol. Biochem.* 135, 95–107.

554 *Kirsten, M., Kimaro, D. N., Feger, K.-H., Kalbitz, K.* (2019): Impact of land use on soil organic carbon
555 stocks in the humid tropics of NE Tanzania. *J. Plant Nutr. Soil Sci.* 182, 625–636.

556 *Kirsten, M., Mikutta, R., Vogel, C., Thompson, A., Mueller, C. W., Kimaro, D. N., Bergsma, H. L. T., Feger,*
557 *K.-H., Kalbitz, K.* (2021): Iron oxides and aluminous clays selectively control soil carbon storage and
558 stability in the humid tropics. *Scientific Reports* 11.

559 *Kleber, M., Eusterhues, K., Keiluweit, M., Mikutta, C., Mikutta, R., Nico, P. S.* (2015): Mineral–Organic
560 Associations: Formation, Properties, and Relevance in Soil Environments. *Adv. Agron.* 130, 1–140.

561 *Kravchenko, A. N., Negassa, W. C., Guber, A. K., Rivers, M. L.* (2015): Protection of soil carbon within
562 macro-aggregates depends on intra-aggregate pore characteristics. *Scientific Reports* 5, 16261.

563 *Larney, F. J.* (2008): Dry-Aggregate Size Distribution, in Carter, M. R., Gregorich, E. G.: *Soil sampling*
564 *and methods of analysis.* 2nd ed. Canadian Society of Soil Science; CRC Press, [Pinawa, Manitoba], Boca
565 Raton, FL, USA, pp. 821–831.

566 *Le Bissonnais, Y.* (1996): Aggregate stability and assessment of soil crustability and erodibility: I. Theory
567 and methodology. *Eur. J. Soil Sci.* 47, 425–437.

568 *Le Bissonnais, Y., Prieto, I., Roumet, C., Nespoulous, J., Metayer, J., Huon, S., Villatoro, M., Stokes, A.*
569 (2018): Soil aggregate stability in Mediterranean and tropical agro-ecosystems: effect of plant roots and
570 soil characteristics. *Plant Soil* 424, 303–317.

571 *Lobe, I., Sandhage-Hofmann, A., Brodowski, S., du Preez, C. C., Amelung, W.* (2011): Aggregate dynamics
572 and associated soil organic matter contents as influenced by prolonged arable cropping in the South Afri-
573 can Highveld. *Geoderma* 162, 251–259.

574 *Maltoni, K. L., Mello, L. M. M. de, Dubbin, W. E.* (2017): The effect of Ferralsol mineralogy on the distri-
575 bution of organic C across aggregate size fractions under native vegetation and no-tillage agriculture. *Soil*
576 *Use Manag.* 33, 328–338.

577 *Marín-Spiotta, E., Swanston, C. W., Torn, M. S., Silver, W. L., Burton, S. D.* (2008): Chemical and mineral
578 control of soil carbon turnover in abandoned tropical pastures. *Geoderma* 143, 49–62.

579 *Mehra, O. P., Jackson, M. L.* (1958): Iron Oxide Removal from Soils and Clays by a Dithionite-Citrate
580 System Buffered with Sodium Bicarbonate. *Clays Clay Miner.* 7, 317–327.

581 *Mujuru, L., Mureva, A., Velthorst, E. J., Hoosbeek, M. R.* (2013): Land use and management effects on soil
582 organic matter fractions in Rhodic Ferralsols and Haplic Arenosols in Bindura and Shamva districts of
583 Zimbabwe. *Geoderma* 209-210, 262–272.

584 *Paul, S., Flessa, H., Veldkamp, E., López-Ulloa, M.* (2008): Stabilization of recent soil carbon in the humid
585 tropics following land use changes: Evidence from aggregate fractionation and stable isotope analyses.
586 *Biogeochemistry* 87, 247–263, www.jstor.org/stable/40343548.

587 *Peng, X., Yan, X., Zhou, H., Zhang, Y. Z., Sun, H.* (2015): Assessing the contributions of sesquioxides and
588 soil organic matter to aggregation in an Ultisol under long-term fertilization. *Soil Tillage Res.* 146, 89–
589 98.

590 *Qafoku, N. P., Sumner, M. E.* (2002): Adsorption and Desorption of Indifferent Ions in Variable Charge
591 Subsoils. *Soil Science Society of America Journal* 66, 1231–1239.

592 *Rabbi, S.M.F., Wilson, B. R., Lockwood, P. V., Daniel, H., Young, I. M.* (2015): Aggregate hierarchy and
593 carbon mineralization in two Oxisols of New South Wales, Australia. *Soil Tillage Res.* 146, 193–203.

594 *Schmidt, M. W. I., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. A., Kleber, M., Kögel-
595 Knabner, I., Lehmann, J., Manning, David A C, Nannipieri, P., Rasse, D. P., Weiner, S., Trumbore, S. E.*
596 (2011): Persistence of soil organic matter as an ecosystem property. *Nature* 478, 49–56.

597 *Schwertmann, U.* (1964): Differenzierung der Eisenoxide des Bodens durch Extraktion mit Ammoniumox-
598 alat-Lösung. *Z. Pflanzenernaehr. Dueng. Bodenk.* 105, 194–202.

599 *Six, J., Bossuyt, H., Degryze, S., Denef, K.* (2004): A history of research on the link between (micro)aggre-
600 gates, soil biota, and soil organic matter dynamics. *Soil Tillage Res.* 79, 7–31.

601 *Six, J., Elliott, E.T., Paustian, K.* (2000a): Soil macroaggregate turnover and microaggregate formation: A
602 mechanism for C sequestration under no-tillage agriculture. *Soil Biol. Biochem.* 32, 2099–2103.

603 *Six, J., Feller, C., Denef, K., Ogle, S. M., Moraes, J. C. de, Albrecht, A.* (2002): Soil organic matter, biota
604 and aggregation in temperate and tropical soils - Effects of no-tillage. *Agronomie* 22, 755–775.

605 *Six, J., Paustian, K., Elliott, E. T., Combrink, C.* (2000b): Soil Structure and Organic Matter. *Soil Sci. Soc.*
606 *Am. J.* 64, 681.

607 *Spaccini, R., Zena, A., Igwe, C. A., Mbagwu, J. S. C., Piccolo, A.* (2001): Carbohydrates in water-stable
608 aggregates and particle size fractions of forested and cultivated soils in two contrasting tropical ecosys-
609 tems. *Biogeochemistry* 53, 1–22, www.jstor.org/stable/1469625.

610 *Tisdall, J. M., Oades, J. M.* (1982): Organic matter and water-stable aggregates in soils. *J. Soil Sci.* 33, 141–
611 163.

- 612 Totsche, K. U., Amelung, W., Gerzabek, M. H., Guggenberger, G., Klumpp, E., Knief, C., Lehdorff, E.,
613 Mikutta, R., Peth, S., Prechtel, A., Ray, N., Kögel-Knabner, I. (2018): Microaggregates in soils. *J. Plant*
614 *Nutr. Soil Sci.* 181, 104–136.
- 615 Trüby, P., Aldinger, E. (1989): Eine Methode zur Bestimmung austauschbarer Kationen in Waldböden. *Z.*
616 *Pflanzenernaehr. Bodenk.* 152, 301–306.
- 617 Vrdoljak, G., Sposito, G. (2002): Soil aggregate hierarchy in a Brazilian oxisol, in *Soil Mineral-Organic*
618 *Matter-Microorganism Interactions and Ecosystem Health, Dynamics, Mobility and Transformation of*
619 *Pollutants and Nutrients. Developments in Soil Science. Elsevier*, pp. 197–217.
- 620 West, S. L., White, G. N., Deng, Y., McInnes, K. J., Juo, A. S. R., Dixon, J. B. (2004): Kaolinite, halloysite,
621 and iron oxide influence on physical behavior of formulated soils. *Soil Sci. Soc. Am. J.* 68, 1452–1460.
- 622 Youker, R. E., McGuinness, J. L. (1957): A short method of obtaining mean weight-diameter values of
623 aggregate analyses of soil. *J. Soil Sci.*, 291–294.