# Thermal signature and quantification of charcoal in soil by differential scanning calorimetry and BPCA markers

Brieuc Hardy<sup>1, 2</sup>, Nils Borchard<sup>3,4</sup> and Jens Leifeld<sup>5,6</sup>

5

<sup>1</sup> Sustainability, Systems & Prospectives Department – Soil, Water and Integrated Crop Production —Walloon Agricultural Research Center, Gembloux, Belgium

<sup>2</sup> Earth and Life Institute – Environmental Sciences, Université Catholique de Louvain, Croix du Sud 2, 1348 Louvain-la-Neuve, Belgium

<sup>3</sup> German Agricultural Society e.V., Eschborner Landstraße 122, 60489 Frankfurt, Germany

<sup>4</sup> Humboldt University Berlin, Faculty of Life Sciences, Invalidenstrasse 42, 10115 Berlin, Germany

<sup>5</sup> Agroscope, Climate and Agriculture Group, Reckenholzsstrasse 191, 8046 Zurich, Switzerland

<sup>6</sup> Environmental Geoscience, Universität Basel, Basel, Switzerland

Correspondence to: Brieuc Hardy (b.hardy@cra.wallonie.be)

- Abstract. Black carbon (BC) plays an important role in terrestrial carbon storage and can improve sustainably improve soil
  fertility. Nevertheless, the accurate quantification of BC remains a critical issue to fully unravel the functions and dynamics of BC in soil. In this work, we explored the potential of differential scanning calorimetry (DSC) to identify, characterize and quantify charcoal in the soil of pre-industrial charcoal kiln sites (CKS) from a diversity of forest and cropland soils in Belgium and Germany. Pre-industrial charcoals and uncharred soil organic matter (SOM) demonstrated a distinct thermal signature that allowed their discrimination, with charcoal being more thermally stable than SOM. The DSC pattern of charcoals is
  characterized by one to three specific exotherms varying in size and position depending on soil conditions. From our data, we assume that the thermal moieties within charcoal depend on the strength of chemical bonds of C atoms (increasing with the degree of aromatic condensation and decreasing with weathering) and on the activation energy required to onset-initiate the combustion. Despite the specific thermal features of charcoal, its decomposition spans over a wide range of temperatures that overlaps with the thermal signature of uncharred SOM. This stresses the challenge of BC quantification in soil and hinders the
- 25 use of cut-off temperatures to accurately quantify charcoal in soil. Therefore, charcoal-C content was estimated from the relative height of exotherms attributed either to the combustion of charcoal or SOM. For a selection of 45 soil samples, charcoal-C content estimated by DSC was compared to benzene polycarboxylic acids (BPCA) patternabundance, a widely used method to quantify BC in soil. The two methods correlated strongly ( $R^2 = 0.97$ ), with BPCA-C representing about one fifth of DSC-derived charcoal-C. This remind usreinforces the view that operationally-defined BC content has an absolute
- 30 quantitative value only if the recovery rate is controlled, which is very complicated for many case studies. Overall, our results demonstrate that dynamic thermal analysis is largely under-exploited despite providing rapidly and at low cost-quantitatively interpretable information all overacross the continuum of soil organic matter.

## 1 Introduction

Black carbon (BC) is the solid residue of incomplete combustion of biomass and fossil fuel, which comprises a wide range of

- 35 thermally altered materials from slightly charred biomass to highly recalcitrant condensates such as soot (Goldberg, 1985; Schmidt and Noack, 2000). A main fraction of terrestrial BC is stored in soil (Forbes et al., 2006; Leifeld et al., 2018; Preston and Schmidt, 2006; Reisser et al., 2016; Santín et al., 2016), where it has a longer residence time than uncharred organic matter (e.g. Wang et al., 2016; Wolf et al., 2013). The increased resistance of BC to (a)biotic degradation has been related to its fused aromatic ring structure (Solomon et al., 2007; Wang et al., 2016). BC has received much interest from soil scientists because
- 40 it plays an important role for in the provision of ecosystems services such as terrestrial carbon storage (Czimczik and Masiello, 2007; Knicker, 2011; Masiello, 2004; Preston and Schmidt, 2006; Schmidt and Noack, 2000) and sustainable soil fertility (Glaser et al., 2001, 2002; Glaser and Birk, 2012; Schmidt et al., 2021). Nevertheless, the accurate quantification of BC in environmental matrices is-remains as a critical issue that makes the role of BC unclear in geochemical processes (Mukherjee and Kumar, 2021).
- 45 According to the general definition reported earlier, BC comprises a wide range of materials with no clear-cut boundaries, which challenges quantification (Kappenberg et al., 2016; Roth et al., 2012; Schmidt et al., 2001). The various forms of BC cover a large molecular continuum (Hammes et al., 2007; Masiello, 2004) that reflects contrasting conditions of formation (Keiluweit et al., 2010; Wiedemeier et al., 2015a). Moreover, properties of chars produced at relatively low temperature overlap with that of uncharred organic compounds naturally present in soil. Consequently, quantification of BC relies on operational
- 50 definitions depending on specific objectives defined by researchers from very different fields in atmospheric, soil, sediment and paleo-environmental sciences (Hammes et al., 2007; Schmidt et al., 2001). The various methods record systematic differences because they recover a-different fractions of the BC continuum and do not all completely isolate BC from other carbon compounds (Hammes et al., 2007; Roth et al., 2012). As a result, Schmidt et al. (2001) reported that variation in the BC content estimated by four different methods for one individual sample was more than two orders of magnitude.
- 55 Typically, five categories of techniques of identification and quantification of BC in soil and sediments are distinguished: physical, thermal, chemical, spectroscopic and molecular markers techniques (Bird et al., 2015; Hammes et al., 2007). The principle of thermal and (thermo-)chemical separation techniques relies on the exposition of the sample to an oxidative treatment in standard conditions. Carbon surviving oxidation is determined by mass loss or elemental analysis and operationally defined as BC. <sup>13</sup>C Nuclear Magnetic Resonance (NMR) spectroscopy is sometimes used to identify BC in the
- 60 residue based on its aromatic signature. A molecular marker technique that is widely used by soil scientists consists in the quantification of benzene polycarboxylic acid (BPCA) markers liberated by digestion of BC in an acid medium (Brodowski et al., 2005; Glaser et al., 1998). This procedure has two main advantages: (i) it relies on the chemical decomposition of BC into markers that are specific to BC, related to its high aromaticity and (ii) it provides information on the degree of aromatic condensation erystallinity of BC according to the number of carboxyl groups on the edge of benzene of the BPCA markers;
- 65 which increases with the degree of aromatic condensation (Glaser et al., 1998). Nevertheless, this method has known

limitations, as soil type and soil organic matter content may affect both quantification of total BC and the pattern of BPCA markers (Brodowski et al., 2005; Kappenberg et al., 2016). ParticularlyIn particular, the recovery of BPCA-C may vary depending on the quality of BC (Hammes et al., 2007; Roth et al., 2012). The less condensed fraction of chars is suspected to be completely decomposed by the strong oxidative attack (Glaser et al., 1998) whereas the most refractory, graphitized forms

- of BC might resist digestion preceding BPCA analysis (Brodowski et al., 2005; Roth et al., 2012). As a result, BPCA-C was shown to correspond to a maximum of  $\frac{1}{2.27}$  of total C in charcoal in optimal conditions of recovery (Glaser et al., 1998). This factor of 2.27 is often used as a conservative conversion factor to estimate total BC content from BPCA-C extracted from soil<u>a</u> although it has been criticized for its validity (Glaser et al., 1998; Schneider et al., 2010).
- Among these techniques used for BC quantification, thermal methods are convenient because they are rapid, reproducible, inexpensive and require little sample preparation (Plante et al., 2009). Thermal resistance has been related to the biological availability of chars (Harvey et al., 2012; Plante et al., 2011) and to the residence time of BC in soil, as revealed by <sup>14</sup>C measurements (Leifeld et al., 2015; Plante et al., 2013). Static thermal methods rely on cut-off temperature values to distinguish between BC and non BC components (e.g. Gustafsson et al., 2001). Nevertheless, widely used chemo-thermal oxidation at 375°C is calibrated for soot (Elmquist et al., 2004) and was shown to recover only from 0 to 44 % of C in chars, with no
- 80 survival for those produced at < 850 °C (Nguyen et al., 2004). In contrast to static methods, dynamic thermal analysis techniques like thermogravimetry (TG) and differential scanning calorimetry (DSC) has have the potential to provide information for the entire continuum of materials that compose soil organic matter (SOM) by scanning a sample over a large range of temperatures (Leifeld, 2007; Plante et al., 2009). Leifeld (2007) highlighted that BC has a specific thermal signature that allows an unambiguous discrimination with from uncharred SOM, as BC is systematically more thermally stable. Among materials potentially interfering with the signature of BC, only bituminous coal had a thermal stability comparable to that of chars.</p>

In this work, we explored the analytical potential of DSC to identify and quantify charcoal-C in the soil of pre-industrial charcoal kiln sites, also referred to as relict charcoal hearths in the literature. Charcoal kiln sites are platforms of *in situ* charcoal production by the earthmound kiln method (Schenkel et al., 1998), which operated at a-maximal temperatures of 400–450 °C (Emrich, 1985). They are widespread in the former forested areas of westernhistorical forest areas in Europe as charcoal has long been the unique combustible used for smelting and steel-making (Hardy et al., 2016; Samojlik et al., 2013). These sites have received increasing attention in recent years, particularly as proxies for long-term soil amendment with biochar (Borchard et al., 2014; Burgeon et al., 2021; Dehkordi et al., 2020; Hardy et al., 2017a; Hirsch et al., 2018; Kerré et al., 2017; Lasota et al., 2021; Mastrolonardo et al., 2019; Pollet et al., 2022; Schneider et al., 2018; Zanutel et al., 2021), which is a promising technology to sequester carbon in soil while maintaining or improving soil fertility (Laird, 2008; Lehmann, 2007).

95 technology to sequester carbon in soil while maintaining or improving soil fertility (Laird, 2008; Lehmann, 2007). Three specific objectives were addressed here: we aimed (i) to characterize the DSC thermal signature of charcoal in soil relative to that of uncharred SOM; (ii) to quantify charcoal-C in soil based on specific charcoal features of the thermograms; and (iii) to compare the content of charcoal-C in soil estimated by DSC to the content of total BC estimated by the BPCA method. To meet these goals, we analyzed charcoal-rich soils sampled at various pre-industrial charcoal kiln sites of Wallonia in Belgium and of the Siegerland and the Eifel in Germany.

### 2 Material and methods

# 2.1 Soil samples

## 2.1.1 Soils of Belgium

- Two series of organo-mineral topsoil samples of kiln and adjacent reference soils from forest (sampled by (sub-)horizons of variable depth; N=38; described by Hardy *et al.*, 2016) and cropland (0-25 cm depth; N=34; described by Hardy et al., 2017a) were analyzed. Briefly, forest soils cover a wide range of textural classes, from sand to clay loam. Soil types of reference soils are Arenosols, Cambisols, Luvisols or Podzols according to the WRB 2014 classification (IUSS Working Group WRB, 2014). They are mainly strongly acidic, except for three Cambisols developed on calcareous parent rocks (limestone, dolostone and marl). In cropland, soil types of reference soils include mainly haplic Luvisols (15 sites) according to the WRB 2014
- 110 classification (IUSS Working Group WRB, 2014). One site was identified as an eutric Cambisol and one as a colluvic Regosol. The particle-size analysis of topsoil (USDA texture; IUSS Working Group WRB, 2014) indicated that soil texture is silt loam at 15 sites and loam at two sites. In Wallonia (Southern Belgium), the climate is oceanic and cold temperate, with mean annual temperature between 6.4 and 9.5 °C and rainfall from 750 to 1400 mm. Charcoal production virtually ceased in the early 19<sup>th</sup> century, when coke replaced charcoal as an industrial fuel in iron metallurgy, and had completely stopped by 1860 (Evrard,
- 115 1956). Therefore, we can reasonably assume that charcoal was deposited >150 years ago. To investigate the effect of the mineralogical background on the thermal signature of soil, we also selected five subsoil samples from relatively clay-rich argic horizons from forested and cultivated Luvisols. Subsoil samples were treated with 6 % H<sub>2</sub>O<sub>2</sub> at 70 °C during 30 days to oxidize SOM with a limited effect on soil mineralogy. The content of <u>soil organic carbon (SOC)</u> that survived oxidation ranged from 0.50 to 0.91 g kg<sup>-1</sup> according to elemental analysis.

### 120 2.1.2 Soils of Germany

For comparison of DSC results with BPCA biomarkers, we also analyzed 45 kiln and reference forest topsoil samples (0–5, 5–20 and occasionally 20-25 cm) from 10 different sites. These had been previously BPCA-characterized (Borchard et al., 2014) following the procedure of Brodowski et al. (2005). Five sites were located in the Siegerland region and five in the Eifel region of Germany. In the Siegerland, soils were Leptic Cambisols (IUSS Working Group WRB, 2014) developed on acidic

125 rock, whereas soils from the Eifel region were Haplic Luvisols, Mollic Leptosols and Leptic Cambisols (IUSS Working Group WRB, 2014) formed from the weathering of calcareous rock (Borchard et al., 2014). Soils from Siegerland were very acidic, with median pH values of 3.9 whereas soils from Eifel were base-rich and had a median pH value of 5.5 (Borchard et al. 2014). In both regions, the climate is cold temperate with mean annual temperatures of 8.9 and 7.7 °C and mean annual precipitations

of 946 and 717 mm in the Siegerland and the Eifel, respectively. The sites were abandoned > 60 years ago (Borchard et al., 130 2014).

# 2.2 Sample preparation and carbon analysis

All soils were air-dried at a maximum of 40 °C until constant weight was reached. They were then gently ground and sieved to 2 mm. Before DSC and BPCA analyzes, the < 2 mm fraction of each soil was ground to powder with an oscillating rings crusher (samples from Belgium) or a ball mill (samples from Germany). The inorganic C content was measured by the modified-pressure calcimeter method (Sherrod et al., 2002). The total content of C was determined by dry combustion, and corrected for inorganic C to obtain the total organic carbon (TOC) content, which includes uncharred SOC and charcoal-C.

Previously Prior to DSC analysis, samples exceeding 60 g kg<sup>-1</sup> of  $\frac{\text{SOC-TOC}}{\text{TOC}}$  were diluted with Al<sub>2</sub>O<sub>3</sub> and homogenized in a ball mill.

## 2.3 Charcoal pieces

- 140 In order to constrain the thermal signature of aged charcoal in soil, macrofragments of charcoal were extracted from kiln soil Ffor a selection of 20 sites from WalloniaBelgium, charcoal pieces were extracted from kiln soil in order to better constrain the thermal signature of charcoal in soil (Hardy et al., 2017b). By this approach, we assume that large charcoal particles have a thermal signature representative of charcoal residues in soil regardless of its size, provided that weathering over time didn't completely erase the aromatic character specific to chars. Charcoal particles > 1 mm were separated from about 2 kg of soil
- 145 by way of wet sieving. The residue, containing charcoal particles, was rinsed abundantly with demineralized water and air dried. Charcoal pieces were separated from inorganic material by flotation in water, and then rinsed again several times with demineralized water in a 500 ml beaker, until the water was clear. Plant residues were removed manually. Between 50 and > 400 charcoal pieces were collected for each site.

To compare thermal characteristics of aged charcoal to that of a charcoal that was never deposited in soil (referred to here as

150 "fresh" charcoal), we also analyzed a birch charcoal that was produced in a traditional mound kiln in <u>august August 2012</u> (Hardy et al., 2016). <u>Previously Prior</u> to analysis, charcoal particles were ground to a powder with an agate pestle and mortar. In total, 21 charcoal samples were analyzed.

# 2.4 Differential scanning calorimetry analysis

155

135

Soils and charcoals were analyzed by heat flux DSC with a DSC 100 (TA Instruments), which measures a-<u>the</u> temperature difference between the sample and an empty reference beside of it, subjected to the same heating program. A heat flow rate is calculated based on the voltage signal corresponding to the difference in temperature between the sample and the empty reference (Plante et al., 2009). Between 15 and 25 mg of soil ground to powder were weighed into an aluminium pan and scanned under a flow of 50 ml min<sup>-1</sup> synthetic air from room temperature to 600 °C, at a heating rate of 10 °C min<sup>-1</sup> (Leifeld, 2007).

- After subtraction of a linear baseline drawn between 150 °C and 600 °C, peak temperatures (°C), peak heights (W g<sup>-1</sup>), 160 temperature of at 50 % heat release (T50) and total heat of reaction (J g<sup>-1</sup>) were measured for each DSC thermogram with the Universal Analysis 2000 software (TA Instruments). Peak area (J g<sup>-1</sup>) was also measured for charcoal pieces only, by vertical drop after identification of identifying a the minima minimum between two peaks and splitting the peaks perpendicularly to the baseline (Figure 1). From repeated measurements (n=6) on one sample, we estimated a 95 % confidence interval for each DSC characteristic that was systematically < 2 % of the measured value.
- 165

The content of charcoal-C was estimated from DSC thermograms by the method described by (Hardy et al., 2017a), based on the relative height of the peaks derived from the combustion of charcoal and of that from the combustion of uncharred SOM (Leifeld, 2007). For acidic forest soils, only one peak was systematically discernable for charcoal and therefore used for quantification. ContrastinglyIn contrast, in cropland and Ca-rich forest soils, three peaks were clearly identifiable for charcoal

- 170
  - and used for quantification. The relationship between thermal signature of charcoal and soil environmental conditions was attributed to the presence of abundant  $Ca^{2+}$  adsorbed to carboxylate groups at the surface of aged charcoal under neutral pH. which decreases the thermal resistance of the O-rich, aged fraction of charcoal (Hardy et al., 2017b).

#### 2.4.1 **Sensibility** Sensitivity analysis

To test the influence of the pattern of heat fluxes related to the combustion of charcoal on the estimation of charcoal-C content, 175 we mathematically simulated soil-charcoal mixtures (n = 18) over a wide-representative range of charcoal-C concentrations (from 5 to 90 % of TOC) based on the DSC pattern of heat release from 9 pre-industrial charcoals from different kiln sites. Simulated mixtures were obtained by summing the thermogram of pure charcoals, at different doses, to the thermogram of a charcoal-free soil. By comparison of predicted values with calculated values, we obtained a root mean square error (RMSE) of 1.39 % of the amount of charcoal-C added.

#### 180 2.5 **BPCA** analyzes

Previously Prior to analysis, samples were dried at 40 °C to a constant weight and sieved to 2 mm. Benzene polycarboxylic acids were extracted from the 45 soils of Germany as specific markers for BC in soil, according to the procedure of Brodowski et al., (2005). An estimation of total BC content was obtained by multiplying total BPCA-C content by 2.27 (Brodowski et al., 2005).

#### 185 3 Results

#### 3.1 Thermal analysis of soils

Independently from soil conditions, reference soils had a characteristic DSC pattern, with a main maximum between 300 and 330 °C, at 310.3±10.0 (mean±s.d.) °C on average for forest reference soils and at 319.2±3.7°C for cropland reference soils (Figure 2). This peak <u>is-was</u> asymmetrical and spreads systematically towards higher temperatures. A smaller peak was sometimes visible in the range of 400–450 °C, particularly in cropland reference soils (Figure 2).

- Soils of from pre-industrial charcoal kiln sites had a more variable signature. In addition to the signal in the 300–330 °C range, they had from one to three additional exotherms of higher thermal stability (Figure 2). We observed two main types of DSC signature for kiln soils. In the first category, (very) acidic forest soils were pooled (Figure 2a-d). These soils showed a characteristic main exotherm at 391.8±14.7 °C, and a small peak of higher thermal stability at around 494.8±19.2 °C that was
- not always clearly visible. The second category comprised calcareous forest soils (Figure 2e, f) and cropland soils (Figure 2gi). The main difference with-to the thermal pattern of forest soils from the first group was the presence of multiple peaks, with an exotherm at 374.7±6.3°C and another at 422.6±2.7 °C in place of a unique peak at around 400 °C. Regardless of the presence of charcoal, most thermograms had a small, sharp endotherm at about 575 °C.

The deep Argic horizons that were H<sub>2</sub>O<sub>2</sub> treated and therefore contained almost no residual organic carbon (OC content < 0.91 g kg<sup>-1</sup>) recorded very limited heat fluxes by DSC analysis (Figure 3). In contrast to organo-mineral soils, no heat was released, and a small endotherm was even recorded between 350 and 600 °C, in addition to the same sharp endotherm at ~575°C as observed in most soils. This result suggests that, for the soils of this study, soil minerals have minor effects on the DSC signature. Accordingly, regression of total heat of reaction against <u>SOC-TOC</u> content provided a very high determination coefficient ( $R^2 \ge 0.97$ ), regardless of the dataset (Figure 4a-c), which highlighted the close relationship between heat released during DSC analysis and the combustion of soil organic materials. Small intercepts of the linear regressions might express

some influence of the mineralogy on heat fluxes.

190

# 3.2 Thermal analysis of charcoals

By superimposing the DSC thermogram of charcoal particles with that of the kiln soil from which particles were extracted, it appeared clearly appears that peaks recorded at temperature higher than 350 °C were related to the combustion of charcoal (Figure 5a). Nevertheless, as suggested by the variability of the thermal signature of kiln soils, the pattern of heat release of pre-industrial charcoals varied to some extent. From one to three main peaks were visible on the thermograms of pure charcoals (Figure 5b). The least stable peak (peak 1) showed the highest variability, with temperatures ranging from 360.5 to 415.6 °C and an average value of 378.1±15.6 °C (mean±sd). Temperature of the second and third peaks was less variable, with average values of 424.4±3.0 °C and 506.6±9.2 °C, respectively.

- The pattern of heat release of pre-industrial charcoals was also compared to that of one birch charcoal that was never aged in soil, produced in a traditional mound kiln\_(Figure 5c) in Dole (France), in august 2012. It is interesting to note that fresh charcoal had a thermal signature very different from that of pre-industrial charcoals (Figure 5c). Fresh charcoal had a main peak at 477 °C with a shoulder at 329 °C, which was lower in temperature than the temperature of the lowest maxima recorded in pre-industrial charcoals. Additionally, T50 of the fresh charcoal was 438°C as compared to that of pre-industrial charcoals 220 ranging from 388 to 418 °C with average values of 400.3±7.9 °C.
  - 7

# 3.3 Quantification of charcoal-C in soil with DSC and comparison with BPCA-C content

Data describing the selection of 45 soil samples from Germany used for the methodological comparison between the quantification of BC content by DSC and by BPCA molecular markers is presented in Appendix 1. By quantitative analysis of the DSC thermograms based on relative heights of peaks attributed to charcoal and uncharred SOM, we estimated a content of

- 225 charcoal-C of 13.5±7.8 g kg<sup>-1</sup> for kiln soils in cropland and 0.9±0.7 g kg<sup>-1</sup> for respective reference soils. In forest, charcoal contents estimated with DSC were much higher, as the sites were never diluted laterally by tillage. Forest kiln soil samples of from Wallonia-Belgium contained 68.7±35.3 g kg<sup>-1</sup> of charcoal-C on average and up to 173.2 g kg<sup>-1</sup> whereas in kiln soil samples from Germany, charcoal-C content is 79.7±68.9 g kg<sup>-1</sup> on average and up to 199.7 g kg<sup>-1</sup>.
- Thermal characteristics of soils from Germany were compared to their total BPCA-C content. We found a strong positive correlation (*r*=0.935) between T50 and the total amount of BPCA-C in soil (Figure 6). However, the BPCA-C content recovered for the forest kiln soils of Germany of 13.2±10.8 g kg<sup>-1</sup> was by far lower than the content of charcoal-C estimated by DSC of 79.7±68.9 g kg<sup>-1</sup>. After multiplication by the conversion factor of 2.27, total BC content estimated from BPCA was 30.3±24.4 g kg<sup>-1</sup>, which remains more than twice smaller thanless than half of the content of charcoal-C estimated by DSC. Nevertheless, both variables were strongly correlated. A significant linear relationship between DSC-estimated charcoal-C content (Figure 7a) or in absolute terms (Figure 7b), with determination-coefficients of determination of 0.89 and 0.97, respectively. The slope of the regression lines (Figure 7a, b) showed that total BPCA-C content underestimated the amount of charcoal-C predicted by DSC by a factor of around 5 the content charcoal C estimated by BPCA C differs by a factor of c. five as compared to the amount estimated by DSC. Accordingly, the ratio BPCA-C: charcoal-C was 0<sub>7</sub>.18±0<sub>7</sub>.03 on average.

#### 210

# 4 Discussion

#### 4.1 Thermal analysis of soils

For the soils of this study, the strong correlation between total heat release measured by DSC and SOCTOC content supports the idea that the mineral background of soil had little effect on the shape of exotherms from the combustion of SOM. Therefore,
in the present case exotherms can directly be related to reactions of combustion of organic components of soil. Nevertheless, in other soils the mineral backgroundmineralogy may interfere strongly with exotherms from SOM combustion. In addition to the inversion of quartz-α to quartz-β at 573 °C (visible on most thermograms of Figure 2), gibbsite, kaolinite and halloysite generate endotherms between 300 and 550 °C (Tan et al., 1986). These minerals, absent or present in relatively small amount in temperate soils of this study, are expected to be present in large amount in many clay-rich tropical soils (Uehara and Gillman, 1981). Nevertheless, the direct measurement of CO<sub>2</sub> emissions by evolved gas analysis (EGA)-rather than heat fluxes recorded

by DSC, alone or the combination with TG, have the potential to <u>get rid of remove most</u> interferences from soil minerals (Peltre et al., 2013) and to generalize the use of dynamic thermal analysis for the characterization and quantification of SOM pools. Thermal analysis of charcoal-rich kiln soils, adjacent reference soils and individual charcoals has highlighted that both charcoal and uncharred SOM are composed of a continuum of materials. Reference soils are dominated by thermally labile uncharred

- SOM compounds that degrade at around 300–330 °C. Spreading of the thermograms towards higher temperatures (Figure 2) suggests that more stable compounds are also present in smaller amount. The degradation of SOM in the 300–350 °C range has been related to the decomposition of aliphatic C molecules, such as carbohydrates (Dell'Abate et al., 2002; Kucerík et al., 2004) and in the range of 400–450°C to the combustion of aromatic C (Satoh, 1984). Accordingly, Li et al. (2002) reported that thermal stability of lignin is higher than that of cellulose but depends on its structure. More recently, Sanderman and
- Grandy (2020) confirmed, by means of combining thermal analysis and analytical pyrolysis of molecules released in different thermal windows, that polysaccharides and lipids are thermally more labile, whereas higher temperature volatiles comprise phenols, aromatics, and N-containing compounds. Therefore, the pattern of heat release of reference soils suggests that they are composed mainly of aliphatic-C from microbial residues as well as cellulose- and hemi-cellulose-derived SOM, mixed to a small amount of aromatic-C that decompose at higher temperature. Small differences in the shape of the thermograms from
- 265 cropland and forest reference soils probably result from a difference in the composition of SOM, possibly related to contrasting quality of organic matter inputs. Reference cropland soils systematically showed a small peak at ~400 °C, which corresponds to the temperature of the main peak attributed to charcoal in kiln soil. By visual inspection of bulk soils, black particles looking like charcoal were found in each of them. These particles might result from (i) from contamination of charcoal from the kiln site or (ii) burning that generally followed deforestation when land was converted from forest to agricultural land in the past (Hovois, 1953). Moreover, plant residues left on the field after harvest were commonly burnt until the 1980s (personal)
- communication of Joseph Dufey).

# 4.2 Thermal analysis of charcoals

Properties <u>The properties</u> and composition of charcoal are known to depend largely on conditions of production such as temperature and heating rate that control the degree of aromaticity and crystallinity of chars (Keiluweit et al., 2010).
 Accordingly, the high thermal stability of BC has been attributed to its polycondensed aromatic structure (De la Rosa et al., 2008). Consistently, the content of aromatic-C estimated by <sup>13</sup>C NMR spectroscopy correlates positively with the proportion of thermally refractory SOMA-positive correlation between the content of aromatic C estimated by <sup>14</sup>C NMR spectroscopy and the proportion of thermally refractory SOM was reported (Harvey et al., 2012; Leifeld, 2007). The thermal stability of aromatic compounds larger than that of aliphatic compounds or O- and H-rich C functionalities (Leifeld, 2007) is in line with the higher binding energy of C=C bonds (520 kJ/mol) as compared to higher than that of C-C, C-O or C-H bonds (350-412 kJ/mol) (Plante et al., 2009). Nevertheless, both degree of aromatic condensation and the presence of crystalline structures

9

(Wiedemeier et al., 2015b) <u>may the degree of crystallinity might</u> govern thermal resistance beyond aromaticity. Leifeld (2007) showed that hexane soot, charred wood and charred rice straw had the same aromaticity but different thermal stabilities. At

comparable aromaticity, thermal resistance depends mainly on the degree of aromatic condensation of char (Harvey et al.,

285 2012; Leifeld, 2007) and can be further influenced by other factors such as ash content (McBeath et al., 2015).

Leifeld (2007) The temperature of the thermally most stable peak was proposed as the most reliable feature to assess the thermal stability of charcoal (Leifeld, 2007) proposed temperature of the thermally most stable peak as the most reliable feature to compare thermal stability of charcoal to that of other organic materials. Leifeld (2007) He showed that the thermal stability 290 of pine wood charred under  $N_2$  increases with charring temperature, and recorded a complete loss of thermally labile compounds at 400 °C, in line with the process of aromatization of charcoal that occurs in the 280-400 °C range (Antal and Grønli, 2003; Bird et al., 2015). Leifeld (2007) also found that the most stable peak of charcoals and charred plant biomass had-occurred at a temperature > 500 °C. In contrast, fresh charcoal of this study had a main maximum at 477 °C and exhibited a signal in the low range of temperature (< 350 °C). These discrepancies might be explained by (i) differences in the quality 295 of charcoal and (ii) differences in the experimental parameters of the DSC analysis. Leifeld (2007) analyzed his samples with heating rate of 20°C min<sup>-1</sup> whereas in this study we used a heating rate of 10°C min<sup>-1</sup>. Temperature of heat release is decreased by slower heating rates (Fernández et al., 2010; Leifeld, 2007), which can explain for the smaller lower temperature of the most stable peak measured for our fresh charcoal. On the other hand, the survival of thermally labile compounds for charcoal produced in a traditional mound kiln suggests that temperature of charring was  $< 400^{\circ}$ C. This is not surprising, as wood 300 pyrolysis by the traditional mound kiln method is expected to reach a maximum of 400–450°C (Emrich, 1985), and that-the local maximum temperature of pyrolysis varies within the mound according to the distance from the hearth. In contrast, no signal was recorded in the lower range of temperature (< 350 °C) for pre-industrial charcoals. The presence of aliphatic C compounds, such as proteins and sugars, were detected in chars produced at low temperature and related to the biological accessibility of chars (Fabbri et al., 2012). T, with thehe amount of labile compounds decreased decreasing at higher with 305 temperature of pyrolysis (Fabbri et al., 2012). The disappearance of the thermally labile fraction for pre-industrial charcoals aged in soil suggests that this fraction was biologically reactive, probably made of residual incompletely transformed organic molecules dominated by aliphatic C. Therefore, it was more subject to (a)biotic decomposition than the thermally stable fraction of charcoal and was completely degraded since the time of charcoal production, > 150 years ago. Accordingly, the presence of a labile fraction in engineered biochars was found to contribute to early emissions of CO<sub>2</sub> after introduction to soil 310 (Sagrilo et al., 2014).

Interestingly, the thermal signature of fresh and aged charcoals is quite different. Aged charcoals are thermally less stable than fresh charcoals and generally have multiple exotherms in the 360–525 °C range. The main process in aging of charcoal is oxygenation starting from the surface and propagating to the core of the particle (Lehmann et al., 2005). Charcoal was produced at relatively low temperature at kiln sites (400–450 °C; Emrich, 1985) and therefore may contain aliphatic-C and amorphous

at relatively low temperature at kiln sites (400–450 °C; Emrich, 1985) and therefore may contain aliphatic-C and amorphous aromatic clusters of small size that would not be recovered as BC by the majority of existing BC quantification procedures.
 Moreover, physical, chemical and biological weathering occurring over time in soil are likely to have is supposed to decreased

the stability of charcoal (Ascough et al., 2011), <u>and creating-create</u> H- and O-rich C functionalities (Cheng et al., 2008; Hardy et al., 2017b; Lehmann et al., 2005) that have a decreased resistance to thermal oxidation. By relating thermal characteristics

- 320 with-to the elemental composition of charcoal, Hardy *et al.*, (2017b) highlighted that the O-rich fraction of charcoal had a specific thermal signature, corresponding to the peak of least thermal stability of charcoal (peak 1, Figure 5b). This is in line with the smaller binding energy of C-O bonds compared to C=C bonds of aromatic clusters in charcoal (Plante et al., 2009). As a result of weathering, the overall thermal stability of aged charcoals is smaller than that of fresh charcoal. Hardy *et al.*, (2017b) also found that the temperature of peak 1 was strongly negatively correlated to the content of Ca in charcoal. The most
- abundant O-rich functional groups in aged charcoals are carboxyl groups (Hardy et al., 2017b; Lehmann et al., 2005; Mao et al., 2012). These are known to have a strong affinity with Ca<sup>2+</sup> (Kalinichev and Kirkpatrick, 2007). The presence of Ca<sup>2+</sup> adsorbed to (poly-)carboxylate groups of charcoal might catalyze thermal decomposition by decreasing the binding energy of C-O bonds (e.g. Hu et al., 2018). ComparablySimilarly, the presence of Al and Fe in the form of trivalent cations complexed to humic compounds of Podzols was shown to alter thermal stability of SOM (Schnitzer et al., 1964). This highlights the importance of soil conditions on the thermal signature of BC, and, more generally, SOM.

Transmission electron micrographs of modern and fossil charcoals have provided evidence of organized and disorganized domains in the matrix of charcoal (Cohen-Ofri et al., 2006). The degree of organization, or aromatic condensation (McBeath and Smernik, 2009; Wiedemeier et al., 2015a) refers to the size and arrangement of aromatic clusters in BC, which increases
with temperature of pyrolysis > 350 °C, once the aromatization of BC is complete or close to (Keiluweit et al., 2010; McBeath et al., 2015). Bird et al. (2015) proposed the existence of three pools of different resistance (labile C, semi-labile aromatic C and stable aromatic polycyclic C) to explain the reactivity of BC. According to the conceptual model of Bird et al. (2015), labile C corresponds to the fraction of BC that is composed of minor pyrolysis products such as anhydrosugars and methoxylated phenols, mineralizable on the (very) short-term; semi-labile aromatic C corresponds to aromatic C with a low
degree of aromatic condensation; and stable aromatic polycyclic C to polycyclic aromatic clusters with a ring size > 7. Unfortunately, this view of charcoal stability made of three pools of distinct chemistry and reactivity cannot be directly translated to the presence of three exotherms measured for aged charcoals of this study. Indeed, fresh and aged charcoals of

345 thermal resistance of charcoal might rather be controlled by (1) the binding energy of C bonds, which is driven by both pyrolysis conditions (aromaticity and aromatic condensation) and the degree of weathering (oxygenation and hydrogenation) of charcoal; and (2) the accessibility to combustion of C moieties, or the activation energy necessary to initiate combustion. As the reaction is partially surface-controlled, it is expected that highly weathered outer surfaces of aged charcoal oxidize already-more readily at lower temperatures whereas the inner parts have no access to O<sub>2</sub> despite the temperature being high enough for reaction. In contrast, an onion-shaped soot particle can only be oxidized layer by layer, and probably requires a

this study were produced at a comparable temperature of about 400  $^{\circ}$ C (and are hence supposed to be both dominated by amorphous aromatic C). However, they have very contrasting DSC pattern of heat release. Therefore, we hypothesize that the

higher activation energy to onset combustion. According to this view, the DSC peaks of contrasting thermal resistance found

in aged charcoals might correspond to (i) C bonds that are weakened by the presence of O in their direct proximity (first peak, less thermally stable); (ii) aromatic C in clusters of small size that are partially weathered but that are not directly bound to oxygen (second peak); and (iii) unweathered aromatic C in clusters of large size (third peak, high thermal resistance). The

355 process of aging of BC in the environment is still incompletely understood and is of prime importance to unravel the role that BC plays in geochemical cycles. In that sense, dynamic thermal analysis has the potential to offer rapid, inexpensive continuous information on the complete BC continuum related to the binding energy of C bonds, which can bring precious-useful new information on the degree of weathering of BC aged in soil.

# 4.3 Quantification of charcoal-C by DSC: advantages and limitations.

- 360 Thermal analysis of charcoal-rich kiln soils, adjacent reference soils and individual charcoals has highlighted that both charcoal and uncharred SOM are composed of a continuum of materials. These two continuums largely overlap, which stresses the issue of BC quantification in soil by <u>static\_continuous</u>-thermal <u>analysismethods</u>: the choice of a cut-off temperature to discriminate quantitatively between charcoal-C and uncharred SOC is not reliable. Therefore, the relative height of peaks attributed either to charcoal or SOM were exploited here to quantify charcoal-C in the soil of pre-industrial kiln sites. For cropland soils, the DSC-derived content of charcoal-C stored in the topsoil of kiln sites correlated strongly (*r*=0.98) to the excess of OC (ΔOC) accumulated in the kiln soil relative to adjacent reference soil, as shown by Hardy et al. (2017a). The slope of the relationship between the two variables is -0.80. By a similar approach for forest soils from Germany, we obtained a slope of 1.0 for the regression line between ΔOC and charcoal-C content estimated by DSC (*r*=0.94). For forest soils from Belgium, sampling by soil sub-horizons of varying depth prevented from implementing the comparison between charcoal-C
- 370 content and  $\Delta OC$ . The consistency between estimates of charcoal-C content by DSC and  $\Delta OC$  brings confidence about the reliability of our DSC quantification procedure. Nevertheless, the slope of 1 for forest soils of Germany supports the view that  $\Delta OC$  is exclusively made of charcoal-C, whereas the slope of ~0.8 for cropland soils from Belgium indicates that only which indicates that ~80 % of  $\Delta OC$  is charcoal-C. This is in agreement with the assumption that the increase of OC at kiln sites is mainly related to charcoal enrichment, and supports the view result suggests that the presence of aged charcoal has
- 375 promoted the stabilization of a small amount of extra uncharred SOM in cropland soils of Belgium, which is consistent with other findings on similar sites (Burgeon et al., 2021; Hernandez-Soriano et al., 2015; Kerré et al., 2016; Liang et al., 2010). The fact that this increase in uncharred SOM is not measured in forests soils from Germany might originate from e. g. (i) the younger age (>60 years) of the sites; (ii) a contrasting effect of charcoal on biomass production or the dynamics of natural SOM depending on soil conditions; and (iii) the effect of repeated tillage over time in cropland that has diluted charcoal
- 380 <u>laterally (Hardy et al. 2017a) and might have accelerated the reconstitution of the natural SOM pool and its incorporation in soil. The consistency between estimates of charcoal-C content by DSC and △OC brings confidence about the reliability of our DSC quantification procedure.</u>

Although the relationship between BPCA-C and DSC-derived charcoal-C was strongly linear (Figure 7), the content of charcoal-C obtained by DSC was more than five times higher than the content of BPCA-C in soil. This result is in line with

- 385 the findings of Brodowski et al. (2005) who stated that BPCA-C may underestimate total charcoal-C content by a factor up to 4.5 or higher, either due to a complete digestion of the less condensed moieties of char or to an inaccessibility of the most refractory, condensed moieties (Brodowski et al., 2005; Glaser et al., 1998). This confirms, for CKS-charcoal kiln soils of this study, that the 2.27 multiplicative factor for the estimation of BC content from BPCA-C is overly-conservative. Overall, this result reminds us that the quantitative interpretation of operationally-defined BC measurements, regardless of the method, 390 must be done with great caution and have be presented as an absolute value only if the recovery rate is controlled, which is practically very complicated for most field case studies. Another source of uncertainty comes from DSC, particularly because of the variability in the shape of soil thermograms according to mineralogy and SOM quality. The shape of thermograms from pre-industrial charcoals did not seem to have much effect on the estimates. Indeed, we obtained very accurate estimates (RMSE = 1.39 %) of charcoal-C content with our peak index for soil-charcoal mixtures numerically simulated from the thermograms 395 of nine different pre-industrial charcoals with different shapes. Nevertheless, we were unable to test how the variability of thermal properties of uncharred SOM affects the accuracy of the estimation because of the difficulty to find soils completely free of BC. This point should be addressed in the future by setting up a strong calibration and validation dataset by adding known amounts of charcoal to a variety of soils initially free of charcoal (or to artificial mixtures of minerals and SOM) as conducted by e. g. (Hammes et al., (2007) for Roth et al., (2012) for other methods of BC quantification. To overcome the 400 issue raised by the variability in both SOM and BC depending on their composition and interactions with soil minerals, another way of improvement might come from peak decomposition of thermograms (Plante et al., 2005), as it is done for example with X-Ray photoelectron spectroscopy (XPS) spectra for atomic quantification. This approach would require to identify the identification of the thermal patterns associated to the different C moieties in SOM and BC and to model the shape of each
- 405 5 Conclusion

The main advantage of dynamic thermal analysis to characterize SOM comes from the fact that it provides a complete view of the continuum of organic materials present in soil. DSC analysis of the soil of pre-industrial charcoal kiln sites and adjacent charcoal-unaffected soils has stressed the complexity of BC quantification by highlighting the fact that the thermal properties of charcoal and uncharred SOM are variable and overlap to a large extent, invalidating the use of cut-off values for an accurate discrimination between aged charcoal and uncharred SOM. Thermal analysis by DSC turned out to be a precious-very useful tool to identify and characterize charcoal in the soil, though. Aged charcoal was shown to have a characteristic thermal signature, overall remaining more thermally resistant than uncharred SOM despite the decrease in thermal resistance-stability due to aging in soil. This thermal pattern was successfully used to quantify charcoal-C in cropland and forest soils. We found a strong linear relationship (R<sup>2</sup>=0.97) between DSC-derived charcoal-C content and BPCA-C content in the soils of this study, with BPCA-C representing about one fifth of DSC-derived charcoal-C. Despite the successful use of DSC to quantify charcoal-C in the soils of this study, our approach based on peak height and position has no general character because of the high

associated exotherms in order to decompose the thermograms on a rational basis.

13

variability in the pattern of heat release of BC and SOM depending on its composition. An approach by peak decomposition might help to overcome this issue, whereas the use of <u>evolved gas analysisEGA</u> rather than DSC has the potential to get rid of the interference of soil minerals with exotherms from SOM combustion. Overall, we believe that the potential of dynamic

420 thermal analysis to characterize and quantify soil organic materials is largely under-exploited despite providing rapidly and at low cost-information on the whole range of organic materials present in soil. In the perspective of amending soil with biochar at large scale to mitigate climate change, dynamic thermal analysis might be a precious-very useful tool to assess biochar stability prior to application and to trace-quantitatively and qualitatively trace its evolution in soil.

#### 6 Acknowledgements

425 Funds were provided by the General Directorate for Agriculture, Natural Resources and Environment - Public Service of Wallonia, Belgium (convention 11/13342) - and the Fonds Spéciaux de Recherche (FSR)\_-Université catholique de Louvain, Belgium (Grant 1C.31200.007). We thank Alain Plante from the University of Pennsylvania for TG, DSC and evolved gas analysisEGA preliminary tests made on a selection of samples and Robin Giger and Roman Hüppi from the Agroscope of Zürich for their indispensable help for DSC and elemental analyzes. BPCA analyzes were part of a study funded by the German Fadarel Institute for Constitute for

430 Federal Institute for Geosciences and Natural Resources (BGR).

# 7 Appendices

Appendix 1. <u>SOC-TOC</u> content, PLFA-C content and DSC-derived charcoal-C content for the selection of 45 soil samples from Germany.

# 8 Author contributions

435 BH characterized soil samples from Belgium and NB characterized soil samples from Germany. BH & JL ran DSC analyzes, made the calculations from the DSC graphs and interpreted DSC data. NB ran BPCA analyzes for the selection of samples from Germany. BH drafted the text revised by JL and NB.

## **9** Competing interests

Authors declare no conflict of interest

## 440 10 References

Antal, M. J. and Grønli, M.: The Art, Science, and Technology of Charcoal Production †, Ind. Eng. Chem. Res., 42(8), 1619–

1640, doi:10.1021/ie0207919, 2003.

Ascough, P. L., Bird, M. I., Francis, S. M., Thornton, B., Midwood, A. J., Scott, A. C. and Apperley, D.: Variability in oxidative degradation of charcoal: Influence of production conditions and environmental exposure, Geochim. Cosmochim. Acta, 75(9),

445 2361–2378, doi:10.1016/j.gca.2011.02.002, 2011.

Bird, M. I., Wynn, J. G., Saiz, G., Wurster, C. M. and McBeath, A.: The Pyrogenic Carbon Cycle, Annu. Rev. Earth Planet. Sci., 43(February), 273–298, doi:10.1146/annurev-earth-060614-105038, 2015.

Borchard, N., Ladd, B., Eschemann, S., Hegenberg, D., Möseler, B. M. and Amelung, W.: Black carbon and soil properties at historical charcoal production sites in Germany, Geoderma, 232–234, 236–242, doi:10.1016/j.geoderma.2014.05.007, 2014.

Brodowski, S., Rodionov, A., Haumaier, L., Glaser, B. and Amelung, W.: Revised black carbon assessment using benzene polycarboxylic acids, Org. Geochem., 36(9), 1299–1310, doi:10.1016/j.orggeochem.2005.03.011, 2005.
Burgeon, V., Fouché, J., Leifeld, J., Chenu, C. and Cornélis, J. T.: Organo-mineral associations largely contribute to the stabilization of century-old pyrogenic organic matter in cropland soils, Geoderma, 388(July 2020), doi:10.1016/j.geoderma.2020.114841, 2021.

Cheng, C.-H., Lehmann, J. and Engelhard, M. H.: Natural oxidation of black carbon in soils: Changes in molecular form and surface charge along a climosequence, Geochim. Cosmochim. Acta, 72(6), 1598–1610, doi:10.1016/j.gca.2008.01.010, 2008. Cohen-Ofri, I., Weiner, L., Boaretto, E., Mintz, G. and Weiner, S.: Modern and fossil charcoal: aspects of structure and diagenesis, J. Archaeol. Sci., 33(3), 428–439, doi:10.1016/j.jas.2005.08.008, 2006.

Czimczik, C. I. and Masiello, C. A.: Controls on black carbon storage in soils, Global Biogeochem. Cycles, 21, GB3005, doi:10.1029/2006GB002798, 2007.

Dehkordi, R. H., Burgeon, V., Fouche, J., Gomez, E. P., Cornelis, J. T., Nguyen, F., Denis, A. and Meersmans, J.: Using UAV collected RGB and multispectral images to evaluate winter wheat performance across a site characterized by century-old biochar patches in Belgium, Remote Sens., 12(15), doi:10.3390/RS12152504, 2020.

Dell'Abate, M. T., Benedetti, A., Trinchera, A. and Dazzi, C.: Humic substances along the profile of two Typic Haploxerert, Geoderma, 107(3–4), 281–296, doi:10.1016/S0016-7061(01)00153-7, 2002.

Elmquist, M., Gustafsson, O. and Andersson, P.: Quantification of sedimentary black carbon using the chemothermal oxidation method : an evaluation of ex situ pretreatments and standard additions approaches, Limnol. Oceanogr. Methods, 2, 417–427, 2004.

Emrich, W.: Handbook of Charcoal Making, D. Reidel Publishing Company, Dordrecht, The Netherlands., 1985.

470 Evrard, R.: Forges anciennes., Editions Solédi, Liège, Belgium., 1956.

475

Fabbri, D., Torri, C. and Spokas, K. A.: Analytical pyrolysis of synthetic chars derived from biomass with potential agronomic application (biochar). Relationships with impacts on microbial carbon dioxide production, J. Anal. Appl. Pyrolysis, 93, 77–84, doi:10.1016/j.jaap.2011.09.012, 2012.

Fernández, J. M., Plante, A. F., Leifeld, J. and Rasmussen, C.: Methodological considerations for using thermal analysis in the characterization of soil organic matter, J. Therm. Anal. Calorim., 104(1), 389–398, doi:10.1007/s10973-010-1145-6, 2010.

Forbes, M. S., Raison, R. J. and Skjemstad, J. O.: Formation, transformation and transport of black carbon (charcoal) in terrestrial and aquatic ecosystems, Sci. Total Environ., 370(1), 190–206, doi:10.1016/j.scitotenv.2006.06.007, 2006. Glaser, B. and Birk, J. J.: State of the scientific knowledge on properties and genesis of Anthropogenic Dark Earths in Central Amazonia (terra preta de Índio), Geochim. Cosmochim. Acta, 82, 39–51, doi:10.1016/j.gca.2010.11.029, 2012.

Glaser, B., Haumaier, L., Guggenberger, G. and Zech, W.: Black carbon in soils : the use of benzenecarboxylic acids as specific markers, Org. Geochem., 29(4), 811–819, 1998.
 Glaser, B., Haumaier, L., Guggenberger, G. and Zech, W.: The "Terra Preta" phenomenon: a model for sustainable agriculture

in the humid tropics, Naturwissenschaften, 88(1), 37–41, doi:10.1007/s001140000193, 2001.

Glaser, B., Lehmann, J. and Zech, W.: Ameliorating physical and chemical properties of highly weathered soils in the tropics
with charcoal - a review, Biol. Fertil. Soils, 35(4), 219–230, doi:10.1007/s00374-002-0466-4, 2002.

- Goldberg, E. D.: Black carbon in the environment, John Wiley & Sons, New York., 1985.
  Gustafsson, Ö., Bucheli, T. D., Kukulska, Z., Andersson, M., Largeau, C., Rouzaud, J. N., Reddy, C. M. and Eglinton, T. I.: Evaluation of a protocol for the quantification of black carbon in sediments., Global Biogeochem. Cycles, 15, 881–890, 2001.
  Hammes, K., Schmidt, M. W. I., Smernik, R. J., Currie, L. A., Ball, W. P., Nguyen, T. H., Louchouarn, P., Houel, S.,
- 490 Gustafsson, Ö., Elmquist, M., Cornelissen, G., Skjemstad, J. O., Masiello, C. a., Song, J., Peng, P., Mitra, S., Dunn, J. C., Hatcher, P. G., Hockaday, W. C., Smith, D. M., Hartkopf-Fröder, C., Böhmer, A., Lüer, B., Huebert, B. J., Amelung, W., Brodowski, S., Huang, L., Zhang, W., Gschwend, P. M., Flores-Cervantes, D. X., Largeau, C., Rouzaud, J.-N., Rumpel, C., Guggenberger, G., Kaiser, K., Rodionov, A., Gonzalez-Vila, F. J., Gonzalez-Perez, J. a., de la Rosa, J. M., Manning, D. a. C., López-Capél, E. and Ding, L.: Comparison of quantification methods to measure fire-derived (black/elemental) carbon in soils
- 495 and sediments using reference materials from soil, water, sediment and the atmosphere, Global Biogeochem. Cycles, 21(3), GB3016, doi:10.1029/2006GB002914, 2007.

Hardy, B., Cornelis, J.-T., Houben, D., Lambert, R. and Dufey, J. E.: The effect of pre-industrial charcoal kilns on chemical properties of forest soil of Wallonia, Belgium, Eur. J. Soil Sci., 67, 206–216, doi:10.1111/ejss.12324, 2016.

Hardy, B., Cornelis, J.-T., Houben, D., Leifeld, J., Lambert, R. and Dufey, J.: Evaluation of the long-term effect of biochar on
properties of temperate agricultural soil at pre-industrial charcoal kiln sites in Wallonia, Belgium, Eur. J. Soil Sci., 68, 80–89,
doi:10.1111/ejss.12395, 2017a.

Hardy, B., Leifeld, J., Knicker, H., Dufey, J. E., Deforce, K. and Cornélis, J. T.: Long term change in chemical properties of preindustrial charcoal particles aged in forest and agricultural temperate soil, Org. Geochem., 107, 33–45, doi:10.1016/j.orggeochem.2017.02.008, 2017b.

505 Harvey, O. R., Kuo, L.-J., Zimmerman, A. R., Louchouarn, P., Amonette, J. E. and Herbert, B. E.: An index-based approach to assessing recalcitrance and soil carbon sequestration potential of engineered black carbons (biochars)., Environ. Sci. Technol., 46(3), 1415–21, doi:10.1021/es2040398, 2012.

Hernandez-Soriano, M. C., Kerré, B., Goos, P., Hardy, B., Dufey, J. and Smolders, E.: Long-term effect of biochar on the stabilization of recent carbon: soils with historical inputs of charcoal, GCB Bioenergy, 1–11, doi:10.1111/gcbb.12250, 2015.

- 510 Hirsch, F., Schneider, A., Bauriegel, A., Raab, A. and Raab, T.: Formation, classification, and properties of soils at two relict charcoal hearth sites in Brandenburg, Germany, Front. Environ. Sci., 6(SEP), doi:10.3389/fenvs.2018.00094, 2018. Hoyois, G.: L'Ardenne et l'Ardennais. L'Evolution Economique et Sociale d'une région. Tome 2., Bruxelles-Paris: Éditions universitaires; Gembloux: J.Duculot, imprimeur-éditeur., 1953. Hu, Y., Wang, Z., Cheng, X., Liu, M. and Ma, C.: Effects of catalysts on combustion characteristics and kinetics of coal-char
- blends, IOP Conf. Ser. Earth Environ. Sci., 133(1), doi:10.1088/1755-1315/133/1/012023, 2018.
   IUSS Working Group WRB: World reference base for soil resources 2014. International soil classification system for naming soils and creating legends for soil maps. World Soil Resources Reports N° 106., Food and Agriculture Organization of the United Nations, Rome, Italy., 2014.

Kalinichev, A. G. and Kirkpatrick, R. J.: Molecular dynamics simulation of cationic complexation with natural organic matter, Eur. J. Soil Sci., 58(4), 909–917, doi:10.1111/j.1365-2389.2007.00929.x, 2007.

- Kappenberg, A., Bläsing, M., Lehndorff, E. and Amelung, W.: Black carbon assessment using benzene polycarboxylic acids: Limitations for organic-rich matrices, Org. Geochem., 94, 47–51, doi:doi:http://dx.doi.org/10.1016/j.orggeochem, 2016.
  Keiluweit, M., Nico, P. S., Johnson, M. G. and Kleber, M.: Dynamic molecular structure of plant biomass-derived black carbon (biochar)., Environ. Sci. Technol., 44(4), 1247–53, doi:10.1021/es9031419, 2010.
- 525 Kerré, B., Bravo, C. T., Leifeld, J., Cornelissen, G. and Smolders, E.: Historical soil amendment with charcoal increases sequestration of non-charcoal carbon : a comparison among methods of black carbon quantification, Eur. J. Soil Sci., 67, 324– 331, doi:10.1111/ejss.12338, 2016.

Kerré, B., Willaert, B. and Smolders, E.: Lower residue decomposition in historically charcoal-enriched soils is related to increased adsorption of organic matter, Soil Biol. Biochem., 104, 1–7, doi:10.1016/j.soilbio.2016.10.007, 2017.

Knicker, H.: Pyrogenic organic matter in soil: Its origin and occurrence, its chemistry and survival in soil environments, Quat. Int., 243(2), 251–263, doi:10.1016/j.quaint.2011.02.037, 2011.
 Kucerík, J., Kovár, J. and Pekar, M.: Thermoanalytical investigation of lignite humic acids fractions, J. Therm. Anal. Calorim.,

76(1), 55–65, doi:10.1023/B:JTAN.0000027803.24266.48, 2004.

520

535 Detection of Black Carbon in Soils by Py-GC/MS, Carbon-13 NMR Spectroscopy and Thermogravimetric Techniques, Soil Sci. Soc. Am. J., 72(1), 258, doi:10.2136/sssaj2007.0031, 2008.

Laird, D. A.: The charcoal vision: a win-win-win scenario for simultaneously producing bioenergy, permanently sequestering carbon, while improving soil and water quality, Agron. J., 100(1), 178–181, doi:10.2134/agronj2007.0161, 2008.

De la Rosa, J. M., Knicker, H., López-Capel, E., Manning, D. A. C., González-Perez, J. A. and González-Vila, F. J.: Direct

Lasota, J., Błońska, E., Babiak, T., Piaszczyk, W., Stępniewska, H., Jankowiak, R., Boroń, P. and Lenart-Boroń, A.: Effect of
Charcoal on the Properties, Enzyme Activities and Microbial Diversity of Temperate Pine Forest Soils, Forests, 12(11), 1488,
doi:10.3390/f12111488, 2021.

Lehmann, J.: Bio-energy in the black, Front. Ecol. Environ., 5(7), 381–387, 2007.

Lehmann, J., Liang, B., Solomon, D., Lerotic, M., Luizão, F., Kinyangi, J., Schäfer, T., Wirick, S. and Jacobsen, C.: Near-

edge X-ray absorption fine structure (NEXAFS) spectroscopy for mapping nano-scale distribution of organic carbon forms in

soil: Application to black carbon particles, Global Biogeochem. Cycles, 19, GB1013, doi:10.1029/2004GB002435, 2005.
Leifeld, J.: Thermal stability of black carbon characterised by oxidative differential scanning calorimetry, Org. Geochem., 38(1), 112–127, doi:10.1016/j.orggeochem.2006.08.004, 2007.
Leifeld, J., Heiling, M. and Hajdas, I.: Age and thermal stability of particulate organic matter fractions indicate the presence

of black carbon in soil, Radiocarbon, 57, 99–107, doi:10.2458/azu, 2015.

550 Leifeld, J., Alewell, C., Bader, C., Krüger, J. P., Mueller, C. W., Sommer, M., Steffens, M. and Szidat, S.: Pyrogenic Carbon Contributes Substantially to Carbon Storage in Intact and Degraded Northern Peatlands, L. Degrad. Dev., 29(7), 2082–2091, doi:10.1002/ldr.2812, 2018.

Li, J., Li, B. and Zhang, X. C.: Comparative studies of thermal degradation between Larch lignin and Manchurian Ash lignin., Polym. Degrad. Stab., 78, 279–285, 2002.

555 Liang, B., Lehmann, J., Sohi, S. P., Thies, J. E., O'Neill, B., Trujillo, L., Gaunt, J., Solomon, D., Grossman, J., Neves, E. G. and Luizão, F. J.: Black carbon affects the cycling of non-black carbon in soil, Org. Geochem., 41(2), 206–213, doi:10.1016/j.orggeochem.2009.09.007, 2010.

Mao, J.-D., Johnson, R. L., Lehmann, J., Olk, D. C., Neves, E. G., Thompson, M. L. and Schmidt-Rohr, K.: Abundant and stable char residues in soils: implications for soil fertility and carbon sequestration., Environ. Sci. Technol., 46(17), 9571–6,

560 doi:10.1021/es301107c, 2012.

570

Masiello, C. A.: New directions in black carbon organic geochemistry, Mar. Chem., 92(1-4), 201-213, doi:10.1016/j.marchem.2004.06.043, 2004.

Mastrolonardo, G., Calderaro, C., Cocozza, C., Hardy, B., Dufey, J. and Cornelis, J. T.: Long-term effect of charcoal accumulation in hearth soils on tree growth and nutrient cycling, Front. Environ. Sci., 7(APR), 1–15, doi:10.3389/fenvs.2019.00051, 2019.

McBeath, A. V. and Smernik, R. J.: Variation in the degree of aromatic condensation of chars, Org. Geochem., 40(12), 1161–1168, doi:10.1016/j.orggeochem.2009.09.006, 2009.

McBeath, A. V., Wurster, C. M. and Bird, M. I.: Influence of feedstock properties and pyrolysis conditions on biochar carbon stability as determined by hydrogen pyrolysis, Biomass and Bioenergy, 73, 155–173, doi:10.1016/j.biombioe.2014.12.022, 2015.

Mukherjee, S. and Kumar, M.: Cycling of black carbon and black nitrogen in the hydro-geosphere: Insights on the paradigm, pathway, and processes, Sci. Total Environ., 770, 144711, doi:https://doi.org/10.1016/j.scitotenv.2020.144711, 2021.

Nguyen, T. H., Brown, R. a and Ball, W. P.: An evaluation of thermal resistance as a measure of black carbon content in diesel soot, wood char, and sediment, Org. Geochem., 35(3), 217–234, doi:10.1016/j.orggeochem.2003.09.005, 2004.

575 Peltre, C., Fernández, J. M., Craine, J. M. and Plante, A. F.: Relationships between Biological and Thermal Indices of Soil Organic Matter Stability Differ with Soil Organic Carbon Level, Soil Sci. Soc. Am. J., 77(6), 2020, doi:10.2136/sssaj2013.02.0081, 2013. Plante, a. F., Pernes, M. and Chenu, C.: Changes in clay-associated organic matter quality in a C depletion sequence as measured by differential thermal analyses, Geoderma, 129(3–4), 186–199, doi:10.1016/j.geoderma.2004.12.043, 2005.

Plante, A. F., Fernández, J. M. and Leifeld, J.: Application of thermal analysis techniques in soil science, Geoderma, 153(1–2), 1–10, doi:10.1016/j.geoderma.2009.08.016, 2009.

Plante, A. F., Fernández, J. M., Haddix, M. L., Steinweg, J. M. and Conant, R. T.: Biological, chemical and thermal indices of soil organic matter stability in four grassland soils, Soil Biol. Biochem., 43(5), 1051–1058, doi:10.1016/j.soilbio.2011.01.024, 2011.

- Plante, A. F., Beaupré, S. R., Roberts, M. L. and Baisden, T.: Distribution of Radiocarbon Ages in Soil Organic Matter by Thermal Fractionation, Radiocarbon, 55(2), 1077–1083, doi:10.1017/s0033822200058215, 2013.
  Pollet, S., Chabert, A., Burgeon, V., Cornélis, J.-T., Fouché, J., Gers, C., Hardy, B. and Pey, B.: Limited effects of centuryold biochar on taxonomic and functional diversities of collembolan communities across land-uses, Soil Biol. Biochem., 164(June 2021), 108484, doi:10.1016/j.soilbio.2021.108484, 2022.
- 590 Preston, C. M. and Schmidt, M. W. I.: Black (pyrogenic) carbon: a synthesis of current knowledge and uncertainties with special consideration of boreal regions, Biogeosciences, 3(4), 397–420, doi:10.5194/bg-3-397-2006, 2006.
  Reisser, M., Purves, R. S., Schmidt, M. W. I. and Abiven, S.: Pyrogenic Carbon in Soils: A Literature-Based Inventory and a Global Estimation of Its Content in Soil Organic Carbon and Stocks, Front. Earth Sci., 4(August), 1–14, doi:10.3389/feart.2016.00080, 2016.
- 595 Roth, P. J., Lehndorff, E., Brodowski, S., Bornemann, L., Sanchez-García, L., Gustafsson, Ö. and Amelung, W.: Differentiation of charcoal, soot and diagenetic carbon in soil: Method comparison and perspectives, Org. Geochem., 46, 66– 75, doi:10.1016/j.orggeochem.2012.01.012, 2012.

Sagrilo, E., Jeffery, S., Hoffland, E. and Kuyper, T. W.: Emission of CO2 from biochar-amended soils and implications for soil organic carbon, GCB Bioenergy, 7, 1294–1304, doi:10.1111/gcbb.12234, 2014.

Samojlik, T., Jędrzejewska, B., Michniewicz, M., Krasnodębski, D., Dulinicz, M., Olczak, H., Karczewski, A. and Rotherham,
 I. D.: Tree species used for low-intensity production of charcoal and wood-tar in the 18th-century Białowieza Primeval Forest,
 Poland, Phytocoenologia, 43(1–2), 1–12, doi:10.1127/0340-269X/2013/0043-0511, 2013.
 Sanderman, J. and Grandy, A. S.: Ramped thermal analysis for isolating biologically meaningful soil organic matter fractions

with distinct residence times, Soil, 6(1), 131–144, doi:10.5194/soil-6-131-2020, 2020.

- 605 Santín, C., Doerr, S. H., Kane, E. S., Masiello, C. a., Ohlson, M., de la Rosa, J. M., Preston, C. M. and Dittmar, T.: Towards a global assessment of pyrogenic carbon from vegetation fires, Glob. Chang. Biol., 22(1), 76–91, doi:10.1111/gcb.12985, 2016. Schenkel, Y., Bertaux, P., Vanwijnsberghe, S. and Carre, J.: An evaluation of the mound kiln carbonization technique, Biomass and Bioenergy, 14(5/6), 505–516, 1998.
- Schmidt, H. P., Kammann, C., Hagemann, N., Leifeld, J., Bucheli, T. D., Sánchez Monedero, M. A. and Cayuela, M. L.:
  Biochar in agriculture A systematic review of 26 global meta-analyses, GCB Bioenergy, 13(11), 1708–1730, doi:10.1111/gcbb.12889, 2021.

Schmidt, M. W. I. and Noack, A. G.: Black carbon in soils and sediments: Analysis, distribution, implications, and current challenges, Global Biogeochem. Cycles, 14(3), 777–793, doi:10.1029/1999GB001208, 2000.

Schmidt, M. W. I., Skjemstad, J. O., Czimczik, C. I., Glaser, B., Prentice, K. M., Gelinas, Y. and Kuhlbusch, T. A. J.: 615 Comparative analysis of black carbon in soils, Global Biogeochem. Cycles, 15(1), 163–167, 2001.

Schneider, A., Hirsch, F., Raab, A. and Raab, T.: Dye tracer visualization of infiltration patterns in soils on relict charcoal hearths, Front. Environ. Sci., 6(NOV), 1–14, doi:10.3389/fenvs.2018.00143, 2018.

Schneider, M. P. W., Hilf, M., Vogt, U. F. and Schmidt, M. W. I.: The benzene polycarboxylic acid (BPCA) pattern of wood pyrolyzed between 200 °C and 1000 °C, Org. Geochem., 41(10), 1082–1088, doi:10.1016/j.orggeochem.2010.07.001, 2010.

620 Schnitzer, M., Turner, R. C. and Hoffman, I.: A thermo-gravimetric study of organic matter of representative Canadian podzol soils., Can. J. Soil Sci., 44, 7–13, 1964.

Sherrod, L. A., Dunn, G., Peterson, G. A. and Kolberg, R. L.: Inorganic Carbon Analysis by Modified Pressure-Calcimeter Method, Soil Sci. Soc. Am. J., 66, 299–305, 2002.

Solomon, D., Lehmann, J., Thies, J., Schäfer, T., Liang, B., Kinyangi, J., Neves, E., Petersen, J., Luizão, F. and Skjemstad, J.:

- 625 Molecular signature and sources of biochemical recalcitrance of organic C in Amazonian Dark Earths, Geochim. Cosmochim. Acta, 71(9), 2285–2298, doi:10.1016/j.gca.2007.02.014, 2007.
  - Tan, K. H., Hajek, B. F. and Barshad, I.: Thermal analysis techniques., in Methods of Soil Analysis, Part I. Physical and Mineralogical Methods. Agronomy Monograph, vol. 9., edited by A. Klute, pp. 151–183, American Society of Agronomy, Madison, WI Todor., 1986.
- 630 Uehara, G. and Gillman, P. G.: The Mineralogy, Chemistry, and Physics of Tropical Soils with Variable Charge Clays, edited by U. of Michigan, Westview Press, Ann Arbor, MI., 1981.
  Wang, J., Xiong, Z. and Kuzyakov, Y.: Biochar stability in soil: Meta-analysis of decomposition and priming effects, GCB Bioenergy, 8(3), 512–523, doi:10.1111/gcbb.12266, 2016.
  Wiedensite D. D. Altiene S., Herlerler, W. C. Krillerrich M. Klehen M. Merielle, C. A. McBardt, A. V. Nier, D. S.

Wiedemeier, D. B., Abiven, S., Hockaday, W. C., Keiluweit, M., Kleber, M., Masiello, C. A., McBeath, A. V., Nico, P. S.,

Pyle, L. A., Schneider, M. P. W., Smernik, R. J., Wiesenberg, G. L. B. and Schmidt, M. W. I.: Aromaticity and degree of aromatic condensation of char, Org. Geochem., 78, 135–143, doi:10.1016/j.orggeochem.2014.10.002, 2015a. Wiedemeier, D. B., Brodowski, S. and Wiesenberg, G. L. B.: Pyrogenic molecular markers: Linking PAH with BPCA analysis, Chemosphere, 119, 432–437, doi:10.1016/j.chemosphere.2014.06.046, 2015b.

Wolf, M., Lehndorff, E., Wiesenberg, G. L. B., Stockhausen, M., Schwark, L. and Amelung, W.: Towards reconstruction of
past fire regimes from geochemical analysis of charcoal, Org. Geochem., 55, 11–21, doi:10.1016/j.orggeochem.2012.11.002,

2013.

Zanutel, M., Garré, S. and Bielders, C. L.: Long-term effect of biochar on physical properties of agricultural soils with different textures at pre-industrial charcoal kiln sites in Wallonia (Belgium), Eur. J. Soil Sci., (April), 1–17, doi:10.1111/ejss.13157, 2021.

645



Figure 1. Measurement of characteristics from DSC thermograms (green line). After identification of peak maxima, peak height (W g-1) was measured as the maximum deviation from a linear baseline drawn between 150 and 600 °C, as illustrated for the main peak of the thermogram (0.9542 W g-1). Total heat release (527.6 J g-1) corresponds to the surface of the graph delimited by the linear baseline. Peak area (J g-1) was obtained by <u>identifying the minimum between two adjacent peaks and splitting the peaks</u> <u>perpendicularly to the baselinevertical drop between two adjacent peaks based on the position of the minimum between two peaks</u>. The black lines indicate the maximum rate of reaction associated to each maximum, and cross the baseline at the onset point

655 (Temperature at which oxidation of the material starts).





- 660
  - Figure 2. Differential scanning calorimetry thermograms of a representative selection of soils from pre-industrial charcoal kiln sites (black curves) and adjacent reference soils (grey curves). Four sites are located on (very) acidic forest soil (a, b, c, d), two sites located on calcareous forest soil (e, f) and three sites on cropland soils (g, h, i).



Figure 3. Differential scanning calorimetry thermograms of the subsoil (argic horizon) of a Haplic luvisol, H<sub>2</sub>O<sub>2</sub> treated.





Figure 4. Total heat released between 150 and 600 °C from kiln and reference soils as a function of <u>SOC\_TOC</u> content; a) forest soils <u>of from WalloniaBelgium</u>; b) cropland soils <u>of from WalloniaBelgium</u>; c) forest soils from Germany.





Figure 5. Differential scanning calorimetry thermograms of several charcoals; a) Thermogram of charcoal particles superimposed to the thermogram of the soil of the pre-industrial charcoal kiln site from which particles were extracted, and to the thermogram of adjacent reference soil (adapted from Hardy *et al.*, 2017a); b) Thermograms of charcoals extracted from an acidic forest soil and from a limed cropland soil. Numbers on the graph identify the maxima of exothermal peaks specific to the combustion of charcoal; c) Thermogram of a birch charcoal that was not aged in soil, produced *in august 2012* in a traditional mound kiln.



685 Figure 6. Temperature of 50 % heat release of bulk soils of <u>from</u> Germany from the study of Borchard et al. (2014) against the fraction of C from BPCA (BPCA-C/SOCTOC).



