

# Synergy between compost and cover crops in a Mediterranean row crop system leads to increased ~~subsurface soil~~ subsoil carbon storage

Daniel Rath<sup>1</sup>, Nathaniel Bogie<sup>2</sup>, Leonardo Deiss<sup>3</sup>, Sanjai J. Parikh<sup>1</sup>, Daoyuan Wang<sup>4</sup>, Samantha Ying<sup>5</sup>, Nicole Tautges<sup>6</sup>, Asmeret Asefaw Berhe<sup>7</sup>, Teamrat A. Ghezzehei<sup>7</sup>, Kate M. Scow<sup>1</sup>

5 <sup>1</sup>Department of Land, Air and Water Resources, University of California Davis, Davis, CA 95618, USA

<sup>2</sup>Department of Geology, San Jose State University, San Jose, CA 95192, USA

<sup>3</sup>School of Environment and Natural Resources, The Ohio State University, Wooster, OH 44691, USA

<sup>4</sup>Department of Environmental Science and Engineering, Shanghai University, Shanghai, OH 200444, CHN

<sup>5</sup>Department of Environmental Sciences, University of California Riverside, Riverside, CA 92521, USA

10 <sup>6</sup>Michael Fields Agricultural Institute, East Troy, WI 53120, USA

<sup>7</sup>Department of Life and Environmental Sciences, University of California Merced, Merced, CA 95342, USA

Correspondence to: Daniel Rath (darath@ucdavis.edu)

**Abstract.** ~~Subsurface~~Subsoil carbon stocks are a prime target for efforts to increase soil carbon storage for climate change mitigation ~~and improving soil health~~. However, ~~subsurfacesubsoil~~ carbon (C) dynamics are not well understood, especially in  
15 soils under long term intensive agricultural management. We compared ~~subsurface C dynamics~~subsoil C storage and soil organic matter (SOM) composition in tomato-corn rotations after 25 years of differing C and nutrient management in the California Central Valley: CONV (mineral fertilizer), CONV+WCC (mineral fertilizer + cover crops) and ORG (composted poultry manure + cover crops). The cover crop mix used in these systems are a mix of oat (*Avena sativa* L.), faba bean (*Vicia faba* L.) and hairy vetch (*Vicia villosa* Roth). Our results showed a ~19 Mg/ha increase in SOC stocks down to 1m under ORG  
20 systems, no significant SOC increases under CONV+WCC or CONV systems, and the ~~accumulation~~increased abundance of carboxyl rich C in the ~~subsurfacesubsoil~~ (60-100 cm) horizons of ~~all systems. Systems also had greater amounts of aromatic carbon in the order~~ ORG > and CONV+WCC > CONV. We identified a potential interaction between cover crops and compost, ~~theorizing that increased macropores from cover crop roots facilitate the transport of soluble C and nutrients into the subsurface, thereby increasing stocks. These systems.~~ Our results ~~demonstrateshow~~ the potential for ~~subsurface~~increased  
25 subsoil carbon storage with compost and cover crop amendments in tilled agricultural systems and ~~highlight~~identify potential ~~pathway~~pathways for increasing carbon transport and storage in ~~subsurface soils~~subsoil layers.

## 1 Introduction

Subsurface agricultural soils have the potential to store large amounts of carbon (Rumpel et al., 2012), with longer C residence times (Paul et al., 1997, 2001) relative to surface soils. The line between surface and subsurface soils is often drawn at the average tillage depth (30 cm) (Chen et al., 2018), but studies of microbial carbon processing and soil health indicators identify surface soils as shallow as 15 cm (Wordell-Dietrich et al., 2017). This top 15 cm, due to its high density of microbial biomass, is easy to sample, responds quickly to disturbance, and its outsized perceived role in ecosystem services and carbon processing may be one of the reasons that the average soil sampling depth has shrunk from 53 cm to 27 cm in studies published in the last 30 years (Yost and Hartemink, 2020). However, this shrinking sampling depth ignores decades of research into the role that subsurface soils play in increasing soil carbon stocks (Rapalee et al., 1998; Rumpel and Kögel Knabner, 2011), and is particularly problematic in agricultural studies given how practices such as cover crops, with rooting depths of up to 85 cm (Fan et al., 2016), can have drastically different effects on surface versus subsurface SOC accumulation (Bernal et al., 2016; Harrison et al., 2011; Tautges et al., 2019). To maximize C storage throughout the entire soil profile, we need to understand how land management influences carbon accumulation and storage in subsurface soils (Chenu et al., 2019).

Not recognizing the different potentials for carbon sequestration in surface and subsurface soils means losing the opportunity to take advantage of their unique biological, physical and chemical differences (Angst et al., 2018; Fierer et al., 2003b; Kautz et al., 2013). Subsurface soils generally have higher clay contents due to illuviation and eluviation processes (Jenny, 1941), as well as greater restrictions on carbon decomposition than in surface soils (Fierer et al., 2003a). The distribution of microbes and carbon are more decoupled in subsurface soils (Dungait et al., 2012) due to lower microbial biomass (Taylor et al., 2002) and more heterogeneous carbon distribution (Chabbi et al., 2009; Syswerda et al., 2011). Subsurface microbes are also less efficient at carbon assimilation than surface microbes, and thus more likely to mineralize SOC to CO<sub>2</sub> (Spohn et al., 2016), possibly due to energy and nutrient limitations (Salomé et al., 2010; Sanaullah et al., 2011). Carbon inputs to subsurface soils are primarily through dissolved organic carbon (DOC) movement (Kaiser and Kalbitz, 2012) and deeper rooting biomass and exudates (Sokol and Bradford, 2019). This contrasts with the greater variety of inputs to surface soils such as compost (White et al., 2020a), plant aboveground biomass/shoots (Liebmann et al., 2020), atmospheric deposition (Mladenov et al., 2012) and abundant shallower root biomass (Poeplau and Don, 2015).

Surface and subsurface SOC also respond differently to agricultural management practices that are concentrated at the soil's surface (Salomé et al., 2010; Syswerda et al., 2011). In a recent study after 19 years of management, cover cropping combined with mineral fertilizer application increased C stocks above 30 cm by ~3.5%, but decreased C over the entire 2 m profile by 7% (Tautges et al., 2019). In the same study, cover cropping combined with composting both increased C stocks above 30 cm by 5%, and increased C over the whole 2m profile by 12.6%. Estimates of whole profile carbon sequestration based solely on data from surface soils can lead to inaccurate estimates of carbon storage potential in agricultural systems (VandenBygaart et al., 2011).

Agricultural practices can increase or decrease subsurface SOC by modifying the processes that control soil carbon stabilization including occlusion in soil aggregates, sorption to soil minerals, and microbial processing of residues (Rumpel and Kögel Knabner, 2011). Crop root exudates can be efficiently transformed by microbes into stable soil carbon (Sokol and Bradford, 2019), but the same exudates can also destabilize aggregates and carbon mineral bonds that are key for long term stabilization (Keiluweit et al., 2015). Large inputs of dissolved organic carbon and nutrients can prime subsurface microbial biomass to decompose native SOC (Bernal et al., 2016; Kuzyakov, 2010), or can provide favorable nutrient stoichiometry for microbial carbon processing (Coonan et al., 2020; Kirkby et al., 2013). In addition, recent studies have highlighted that subsurface SOC may be vulnerable to loss under changing environmental conditions, including warming (Hicks Pries et al., 2018) and changes in soil moisture status (Min et al., 2020). Taken together, these studies demonstrate that to accurately predict whether a specific farming practice will increase or decrease subsurface SOC storage in a changing climate, it is necessary to perform studies that explicitly examine deeper soils.

70 Our overarching question asks how agricultural management changes carbon formation and storage processes in subsurface compared to surface soils. We determined the impacts of three land management systems on carbon storage and SOC-related indicators in the microbially active top 15 cm of soil, subsurface soils below 60 cm, and the intervening region (15-60 cm). Agricultural subsoils (>60cm) have the potential to store large amounts of carbon (Rumpel et al., 2012), for a longer period (Paul et al., 1997, 2001) relative to surface soils (<15cm). Surface soils are much easier to sample than subsoils, and respond  
75 more quickly to management, which makes them the focus of most studies of how soil organic carbon (SOC) is formed and stored. However, an increased focus on interrogating the surface soil to answer questions about processes in the entire soil profile exacerbates the risk of subsoils being treated merely as “more dilute surface soils” (Salomé et. al 2010) and ignores decades of research into the unique role that subsoils play in increasing soil carbon stocks (Rapalee et al., 1998; Rumpel and Kögel-Knabner, 2011). A focus on surface soils is particularly problematic in agricultural studies, given how practices such as  
80 cover cropping can have drastically different effects on surface versus subsoil SOC accumulation (Bernal et al., 2016; Harrison et al., 2011; Tautges et al., 2019) depending on the cover crop used. In addition, recent studies have highlighted that subsoil SOC may be vulnerable to loss under changing environmental conditions, such as warming (Hicks Pries et al., 2018) and drought (Min et al., 2020). To maximize C stored in the entire soil profile, we need to understand and capitalize on the numerous and interacting physical, chemical, and biological changes throughout the profile (Angst et al., 2018; Fierer et al.,  
85 2003b; Kautz et al., 2013).

The fact that the average soil sampling depth has decreased from 53 to 27 cm in studies published in the last 30 years (Yost and Hartemink, 2020) may be based on ease of sampling and a focus on surface microbiological processes, but also due to the lack of agreement on where surface soils end and subsoils begin. Depending on the goals of the study, the lower limit of surface soils may be anywhere between the top 7.5 and top 30 cm of the profile, while the upper limit of subsoils can range anywhere  
90 from 20-100cm (Soong et. al 2021, Chen et. al 2018, Lorenz and Lal 2005, Whitmore et. al 2015). There may also be an “intermediate” or “transition” zone that is operationally defined, often corresponding to the maximum tillage depth (Mobley et. al 2015). In this study, we have defined surface soils as the top 0-15 cm of the profile, and subsoils as the lower 60-100 cm, with the intervening 15-60 cm as an intermediate zone based on previous work carried out at our study site, and the relative lack of horizon formation in these young soils.

95 The unique role that subsoils play in storing SOC is due in part to the extensive, site-specific changes that happen across the soil profile. Often, there are changes in bulk density and mineralogy due to clay accumulation, but the exact magnitude and direction of this change varies depending on depositional environment and soil forming factors (Brady and Weill 2015, Soil Survey Staff 2014, Jenny 1941). Subsoils also experience much less disturbance than surface soils, with lower fluctuations in temperature and moisture content (Smitii 1932, Cole and Matthews 1939, Zeynoddin et. al 2019, de Quieroz et. al 2020) and  
100 less mechanical disturbance such as tillage (though it is important to note that tillage events deeper than 30 cm are not altogether rare in many systems). Inputs of oxygen, water, C, and nutrients are usually lower to subsoil than surface soils and mostly

occur via transport through the soil pore network constructed from intra-aggregate pore spaces, root channels, and cracks that form as the soil dries (Pagliai 2004, Sanderman and Amundson 2008). The types of C input are much less varied in subsoils, mostly coming from biomass and exudates of plants with deep roots (Sokol and Bradford, 2019) and transport of dissolved organic carbon (DOC). This downward transport of DOC is described by the “cascade theory” (Kaiser and Kalbitz, 2012), where subsoil DOC inputs undergo a series of successive sorption, desorption, microbial processing, and transport steps. This results in a gradual increase in the age of carbon as we move through the soil profile, with subsoil carbon molecules as old as  $10^3$ - $10^4$  years (Rumpel et. al 2012) compared to younger  $C_{14}$  ages of  $10^2$ - $10^3$  years in the top 30 cm. The transport of carbon into the subsoil via both roots and DOC movement leads to more heterogeneous carbon distribution (Chabbi et al., 2009; Syswerda et al., 2011) that is closely associated with the soil pore network.

Since the soil pore networks responsible for dissolved carbon transport are also hotspots of microbial activity (Banfield et. al 2017), carbon molecules found in the subsoil have often undergone extensive microbial transformation and processing. However, once that carbon does enter the subsoil, it is less likely to undergo further microbial processing due to the combination of heterogeneous carbon distribution, decoupled microbial-carbon presence (Dungait et al., 2012), greater metabolic and physical restrictions on carbon decomposition (Fierer et al., 2003a) and lower microbial biomass (Taylor et al., 2002). This leads to higher concentrations of simpler, microbially-derived carbohydrates, aliphatics and carboxylates in subsoils, in contrast to the more complex aromatic structures in cellulose and lignin present in surface soils (Roth et. al 2019). These simple microbial products may preferentially associate with mineral surfaces (Samson et. al 2020, Williams et. al 2018), driving the formation of mineral-associated organic matter and further rendering that carbon inaccessible to microbes. Subsoil microbial communities have adapted to this relative scarcity of carbon and nutrients (Salomé et al., 2010; Sanaullah et al., 2011) by increasing the proportion of Gram+ bacteria, whose thicker cell walls make them more resilient to adverse environmental conditions. These subsoil microbes may also optimize for survival rather than population growth, being less efficient at carbon assimilation than surface microbes (Spohn et al., 2016), and thus more likely to mineralize SOC to  $CO_2$ . Low carbon use efficiency would also be expected in soils with unfavourable carbon-nutrient stoichiometry for biomass production (Ng et. al 2014, Coonan et. al 2020).

Using existing methods to examine subsoils under different land management practices can help explain how the size and concentration of soil carbon stocks are related to types of carbon input and the status of the soil microbial community (Sradnick et. al 2014). While it is difficult to accurately estimate whether the carbon, nutrient and water status of a particular subsoil will promote or hamper microbial SOC decomposition (Soong et. al 2020); some insight can be obtained by looking at microbial stress levels. Phospholipid fatty acid analysis (PLFA) targets metabolically active cells in soil (Zhang et. al 2019) and is effective in measuring rapid changes in active microbial cell walls and membranes (Frostegård et. al 2010). Measurements of Gram negative: Gram positive ratios via PLFA agree with those obtained via the more recent 16s rRNA metabarcoding (Orwin et. al 2018) and are useful as an indicator of microbial nutrient limitation under different land management practices.

Understanding how these practices then affect the molecular composition of SOM is more difficult, as the most accurate method for quantifying specific C functional groups in soil (nuclear magnetic resonance, NMR) is sensitive to carbon concentrations, the presence of iron oxides, and requires extensive sample preparation if samples contain low C concentrations or a high abundance of paramagnetic species (Fe, Mn) (Bleam 1991, Smernik and Oades 2022). However, Fourier transform infrared spectroscopy (FTIR) presents a rapid, lower-cost method that allows pseudo-quantification of the relative abundance of certain carbon functional groups (Margenot et. al 2016). While FTIR may also be used as a high-throughput method to predict soil properties (Dangal et. al 2021, Deiss et. al 2020), it is particularly useful when comparing changes in SOM structure over time via spectral subtractions (Margenot et. al 2019).

Agricultural practices can increase or decrease subsoil SOC by modifying the physical, chemical, and biological processes that control microbial mineralization of soil carbon including occlusion in soil aggregates, sorption to soil minerals, microbial processing of residues and C transport into the subsoil (Rumpel and Kögel-Knabner, 2011). Crop root exudates can be efficiently transformed by microbes into stable soil carbon (Sokol and Bradford, 2019), but the same exudates can also destabilize aggregates and carbon-mineral bonds that are key for protecting carbon from mineralization (Keiluweit et al., 2015). Large inputs of dissolved organic carbon and nutrients can prime subsoil microbial biomass to decompose native SOC (Bernal et al., 2016; Kuzyakov, 2010), or provide the nutrients needed for microbes to process soil carbon (Coonan et al., 2020; Kirkby et al., 2013) and promote the formation of mineral-associated organic matter (MAOM) (Lavallee et. al 2020). Cover crops may not only increase soluble organic carbon inputs (Steenwerth and Belina, 2008 - rye) but can also influence carbon dynamics indirectly by increasing soil macroporosity and pore connectivity (Scott et al., 1994 - rye, vetch, and lupin, Haruna et al., 2018 - rye; Çerçioğlu et al., 2019 - rye, peas, vetch; Gulick et al., 1994 - bromegrass), as well as increasing topsoil disturbance due to the processes of planting, mowing, and incorporation. These indirect cover crop effects can lead to increases in both infiltration and hydraulic conductivity in fine textured soils and potential increases in soluble C transport, particularly over longer time scales. It is clear that to accurately predict whether a specific farming practice will increase or decrease subsoil SOC storage in a changing climate, it is necessary to perform studies that explicitly examine deeper soils.

Surface and subsoil SOC also respond differently to agricultural management practices that are concentrated at the soil's surface (Chenu et al., 2019; Syswerda et al., 2011). In a recent study at our field site, after 19 years of management cover cropping (oats, fava beans and vetch) combined with mineral fertilizer application increased C stocks above 30 cm by ~3.5%, but decreased C over the entire 2 m profile by 7% (Tautges et al., 2019). The same mix of cover crops combined with compost both increased C stocks above 30 cm by 5%, and increased C over the whole 2m profile by 12.6%. Estimates of whole-profile carbon sequestration based solely on data from surface soils can lead to inaccurate estimates of carbon storage potential in agricultural systems (VandenBygaart et al., 2011).

Our goal was to compare the magnitude of, and potential mechanism behind, changes in carbon stocks throughout a 1-m-deep soil profile after being managed for 25 years with different agricultural practices (as described in Tautges and Chiartas et al.

2019). Our experimental treatments were compost with cover crops (ORG), cover crop with mineral fertilizer (CONV+WCC), and mineral fertilizer application (CONV), and ~~indicators measured include aggregation, carbon chemistry, nutrient availability, and microbial community dynamics down to 1 m. Our specific hypotheses were: samples were taken from the microbially active top 15 cm of soil, subsoils below 60 cm, and the intervening region (15-60 cm). We hypothesized that~~ significant increases in subsoil carbon stocks were associated with the combination of high concentrations of soluble C and nutrients from compost and increased hydraulic transport associated with cover crops.

1. ~~SOC, DOC and nutrient stocks in surface (0-15 cm) and subsurface (60-100 cm) ORG systems will be higher relative to CONV and CONV+WCC stocks.~~
2. ~~Higher microbial biomass and evidence of microbially processed carbon is found in the subsurface of the ORG and CONV+WCC than of CONV systems.~~
3. ~~Soil aggregation and hydraulic conductivity is higher in ORG than CONV+WCC or CONV systems.~~

~~Given that small, cumulative subsurface management impacts may take decades to become detectable, the standard two to three year agronomic experiment is usually not sufficient to observe impacts of agricultural management practices on subsurface soils (Dick, 1992; Johnston and Poulton, 2018; Keel et al., 2019). We address this issue by asking our questions at the Century Experiment at the Russell Ranch Sustainable Agricultural Facility in Davis, CA, where inputs and management history have been tracked in detail over the last 25 years (Wolf et al., 2018). This unique opportunity allows us to draw conclusions relevant to row crops in California's Central Valley, one of the world's most productive agricultural regions (Pathak et al., 2018) and one quite susceptible to negative impacts of climate warming (Medellín-Azuara et al., 2011).~~

Given that small, cumulative subsoil management impacts may take decades to become detectable, impacts of agricultural management practices may not be detectable in the two-to-three-year focus of most agronomic field studies (Dick, 1992; Johnston and Poulton, 2018; Keel et al., 2019). We conducted our study at the Century Experiment at the Russell Ranch Sustainable Agricultural Facility in Davis, CA, where inputs and management history have been tracked over the last 25 years and are representative of row crop systems of the California Central valley (Wolf et al., 2018). This unique opportunity allows us to provide knowledge to make soil management recommendations for farming systems in California's Central Valley, one of the world's most productive agricultural regions (Pathak et al., 2018) and one quite susceptible to negative impacts of climate warming (Medellín-Azuara et al., 2011).

## 2 Methods

### 2.1 Field Site and Historical Management

The experiment was conducted at the Century Experiment at the Russell Ranch Sustainable Agricultural Facility in Davis, CA, in the southern region of the Sacramento Valley at an elevation of 16 m. A detailed description of management history at the Century Experiment is provided in Tautges and Chirtas et al (2019) and is described here only briefly. Davis experiences hot summers and cool winters, with a 2018-2019 average temperature of 16°C from November to March when cover cropping occurs, and 29°C during the normal vegetable production period of April to September. Average annual rainfall for the 2018-



200 2019 year was 812 mm, most of which fell between December - April in keeping with the xeric moisture regime in this area.  
(Supplementary Figure A5) (<http://atm.ucdavis.edu/weather/uc-davis-weather-climate-station/>).

The site has two soil types: (a) Yolo silt loam (Fine-silty, mixed, superactive, nonacid, thermic Mollic Xerofluvent) and (b) Rincon silty clay loam (fine, smectitic, thermic Mollic Haploxeralf). Detailed soil horizon information (classification, texture and depths) can be found in Supplementary Table A3 and the Century Experiment published dataset in Wolf et al.  
205 (~~2018~~-2018). Abbreviations used in this paper (CONV, CONV+WCC, ORG) correspond to the abbreviations used in Wolf  
et. al 2018 (CMT, LMT, OMT), and are identical to those used in Tautges & Chiartas et. al 2019 for ease of comparison.

The experimental design is a randomized complete block design (RCBD) with three blocks and nine systems. Two blocks are placed on the Rincon silty clay loam, and the third block is on the Yolo silt loam. Experimental plots were 64 m x 64 m (0.4 ha). Only three systems of the nine described in Tautges and Chiartas et al. (2019) were measured in the current paper: CONV  
210 (mineral fertilizer), CONV+WCC (mineral fertilizer + cover cropped) and ORG (composted poultry manure + cover cropped).  
All plots are in a two-year maize-tomato rotation, with three replicate plots of each crop in any given year. Each treatment  
sampled in this manuscript consisted of 3 plots under tomato and 3 plots under corn, to give a total of 9 corn plots and 9 tomato  
plots in total. All plots were irrigated with subsurface drip at the time of sampling, having converted from furrow irrigation to  
subsurface drip in 2014. While the lack of a compost-only treatment at Russell Ranch precludes conclusions about the impact  
215 of compost application alone, comparing the CONV+WCC treatment to the ORG treatment allows us to highlight how adding  
compost to a cover cropped plot impacts surface and subsoil carbon stocks, and provides insight into why these impacts occur.

## **2.2 Historic Carbon, Nutrient and Bulk Density Values**

Historical cover crop shoot, compost, and crop residue inputs were calculated based on the Century Experiment published  
dataset in Wolf et al. (2018). Total C and N of composted manure, aboveground cover crop biomass, and crop residues were  
220 determined on a CS 4010 Costech Elemental Analyzer (Costech Analytical Technologies). Total C and N incorporated was  
calculated by multiplying percent C and N of residues by total harvest biomass. Due to compost nutrient analysis not being  
performed every year, estimates from 1993-2000 used %C, N, P and S values averaged for that 7-year period, while estimates  
from 2000-2018 used %C, N, P and S values averaged for that 18-year period. Total aboveground C, N, P and S inputs were  
calculated by summing above ground crop residue, WCC, mineral fertilizer and compost inputs per plot per year. Calculated  
225 N inputs represent the total N content of the aboveground added WCC and crop residue biomass, and do not differentiate  
between fixed N and N uptake from the soil in the case of cover crop legumes.

Soil % carbon and nitrogen values for 0-15, 15-30, 30-60 and 60-100 cm in 1993 and 2012 were taken from Tautges & Chiartas  
et. al (2019), while values for the same depths in 2003 were taken from the Century Experiment published dataset in Wolf et.  
al (2018). Carbon and nitrogen analyses used in this paper were all performed using the same methods (Tautges and Chiartas  
230 et. al 2019, Wolf et, al 2018) on ball-milled, air-dried samples in a CS 4010 Costech Elemental Analyzer (Costech Analytical

Technologies). Total carbon and nitrogen values for 15-60 cm in 1993 were calculated by performing a weighted average of C and N % values from 15-30 and 30-60 cm.

Bulk density values used in this paper were sampled using a Giddings hydraulic probe to 2m in 1993, 2007 and 2012 (2007 values taken from Wolf et. al 2018 and 1993, 2012 values taken from Tautges et. al 2019). In 1993, bulk density was collected in 0–25, 25–50, 50–100, and 100–200 cm depth layers with an 8.25 cm diameter probe. In 2007 and 2012, bulk density was collected in 0–15, 15–30, 30–60, and 60–100 cm depth layers, with a 4.7 cm diameter probe. In 1993, 2007 and 2012, cores were collected from four random locations within each plot. Bulk densities were determined using mass of oven-dried soil (105°C, 24 hr.) and total volume of the core averaged for each depth increment (Blake and Hartge, 1986). Bulk density depths from 1993, 2007 and 2012 were adjusted to 2018 depths through the calculation of weighted averages using adjacent depth layers for comparison. Historical carbon stocks from 0-100 cm for 1993, 2003 and 2012 were calculated via depth weighted sum (Tautges and Chiartas et. al 2019) using bulk density values taken in 1993, 2007 and 2012 respectively. Depth-adjusted 2012 bulk density values were then used to calculate 2018 carbon and nutrient stocks due to the lack of more recent bulk density measurements for all plots. Bulk density values below 30 cm were assumed to have not undergone large changes between 2012-2018 (Tautges and Chiartas et. al 2019), while bulk density sampling from 0-20 cm in select Century Experiment plots indicated a limited difference in bulk density (less than 3%) from 2012-2019 (Wang, unpublished data).

### 2.3 Field Management–2018 Operations

Cover crop planting and incorporation in ORG and CONV+WCC systems in 2017-2018 followed the trend of previous years, being planted onto 15 cm raised beds 1.5 m apart with a mixture of oat (*Avena sativa* L., 42.0 %C, 2.5 %N), faba bean (*Vicia faba* L., 44.1 %C, 3.5 %N) and hairy vetch (*Vicia villosa* Roth, 44.5 %C, 5.2 %N), and terminated by mowing plus 2-3 disking passes in March. Cover crop biomass was sampled by cutting aboveground biomass from one 4.5 m<sup>2</sup> area in each plot prior to termination. Corn and tomato biomass residues were measured by cutting aboveground biomass at two 1.5 m<sup>2</sup> locations per plot after harvest. Biomass samples were oven dried at 65 °C for 4 days and ground to 2 mm prior to total C and N analysis. Fertilization during the 2017-2018 growing season was also similar to like previous years, with CONV and CONV+WCC plots receiving 325 kg/ha 8-24-6 (26 kg N/ha, 78 Kg P/ha, 19.5 kg K/ha) starter fertilizer at the time of planting. Tomato CONV plots also received ammonium sulfate at a total rate of 200 kg N/ha, while maize CONV plots received ammonium sulfate at a total rate of 235 kg N/ha.

From 1993-2018, ORG plots normally received a spring application (February 2018) of composted poultry manure at a rate of 3.6 Mg/ha (24.9 % C, 3.5 % N, 1.6 %P, 1.47 %S). However, during the 2018 season, these plots switched from spring to fall compost application, resulting in an additional application of 3.6 Mg/ha compost in September 2018.



## 2.3 Historic Carbon and Nutrient Inputs

Historical cover crop shoot, compost, and crop residue inputs were calculated based on the Century Experiment published dataset in Wolf et al. (2018). Total C and N of composted manure, aboveground cover crop biomass, and crop residues were determined on a CS 4010 Costech Elemental Analyzer (Costech Analytical Technologies). Total C and N incorporated was calculated by multiplying percent C and N of residues by total harvest biomass. Due to compost nutrient analysis not being performed every year, estimates from 1993–2000 used %C, N, P and S values averaged for that 7-year period, while estimates from 2000–2018 used %C, N, P and S values averaged for that 18-year period. Total aboveground C, N, P and S inputs were calculated by summing aboveground crop residue, WCC, mineral fertilizer and compost inputs per plot per year. Calculated N inputs represent the total N content of the aboveground added WCC and crop residue biomass, and do not differentiate between fixed N and N uptake from the soil in the case of cover crop legumes.

## 2.4 Soil Sampling

Soil sample collection took place in the 2018–2019 growing season. Plots were sampled at 4 timepoints: February 2018 (Pre-CC Incorporation), June 2018 (Mid-Season), September/October 2018 (Post-Harvest), and February 2019 (Pre-CC Incorporation). A substantial amount of variation in both EOC and mineral N measurements can occur during the growing season (Li et al. 2018). Our sampling regime at multiple timepoints was meant to account for that variation in both winter and summer months to give a more accurate snapshot of carbon and nutrient availability during the growing season. All sampling took place in the raised beds between furrows. Samples in February 2018, September/October 2018 and February 2019 were taken using a tractor-mounted Giddings probe with a diameter of 3 cm from all replicate plots of each system (n = 6 plots per treatment). Samples taken in June 2018 were taken using an auger to 100 cm and were only taken in the experimental plots planted with tomato (n = 3 plots per treatment). Three replicate cores were taken per plot, sectioned into 0–15, 15–60 and 60–100 cm depths, composited, and then subsampled. Aliquots of each soil were frozen at -20 °C for PLFA analysis within 48 hours of sampling, while the remaining samples were sieved to 8 mm and stored at 4 °C until analyzed.

Bulk density values used in this paper were sampled using a Giddings hydraulic probe to 2m in 1993 and 2012. In 1993, bulk density was collected in 0–25, 25–50, 50–100, and 100–200 cm depth layers with an 8.25 cm diameter probe. In 2012, bulk density was collected in 0–15, 15–30, 30–60, 60–100, and 100–200 cm depth layers, with a 4.7 cm diameter probe. In both 1993 and 2012, cores were collected from four random locations within each plot. Bulk densities were determined using mass of oven-dried soil (105°C, 24 hr.) and total volume of the core averaged for each depth increment (Blake and Hartge, 1986). Bulk density depths from 1993 and 2012 were adjusted to 2018 depths through the calculation of weighted averages using adjacent depth layers.

## 2.5 Carbon, Nutrient and Aggregation Analysis

Dissolved organic carbon was determined using a 0.5 M potassium sulfate extraction. 6 g of soil were extracted with 0.5 M K<sub>2</sub>SO<sub>4</sub> in a 1:5 ratio, shaken for one hour, filtered through Q5 filter paper and analyzed within 48 hours on a Shimadzu TOC-L Total Organic Carbon analyzer according to Jones and Willett (2006). Aliquots of the K<sub>2</sub>SO<sub>4</sub> extract were immediately

frozen at -20°C and later analyzed for nitrate by reacting with vanadium(III) chloride according to Doane and Horwath (2003); and ammonium via the Berthelot reaction as laid out in Rhine et al. (1998). Available calcium, phosphorus and sulfur were measured on 2 mm sieved air-dried samples using the Mehlich-3 soil test (Mehlich, 1984). Aggregation measurements were carried out using the method outlined in Wang et al. (2017), adapted from the wet sieving method outlined in Elliott (1986). Soils were gently passed through an 8mm sieve, and a 50g representative sample was submerged in room temperature water on top of a 2 mm sieve. This sieve was moved up and down for 2 min (50 submersions per minute) using an audio metronome to keep track of the number of submersions. The soil and water passed through the 2mm sieve were gently transferred by rinsing onto a 250 µm sieve and submerged again. The process was repeated using a 53 µm sieve to generate 4 aggregate size fractions (8 mm-2 mm, 2 mm-250 µm, 250 µm-50 µm, >50 µm) which were rinsed into pre-weighed aluminum pans, oven-dried at 60 °C, and weighed. Mean weight diameter of the aggregate fractions was calculated as the weighted average of the four aggregate size fractions (van Bavel, 1950). All analyses described below were carried out on samples taken during the 2018-2019 growing season. Extractable organic carbon was determined using a 0.5 M potassium sulfate extraction within 48 hours of sampling. 6 g of soil were extracted with 0.5 M K<sub>2</sub>SO<sub>4</sub> in a 1:5 ratio, shaken for one hour, filtered through Q5 filter paper and analyzed within 48 hours on a Shimadzu TOC-L Total Organic Carbon analyzer according to Jones and Willett (2006). Due to the moisture limited conditions present during summer at our study site, we chose an EOC extraction method as opposed to DOC sampling via tension lysimeters to compare soluble C measurements at different timepoints and soil water contents. Measurements of EOC are commonly used to estimate soluble carbon (Slessarev et. al 2020, Matlou et. al 2007) and may be more sensitive to recent carbon and litter inputs, making them more suitable for answering questions on the impacts of carbon input, nitrogen amendment and tillage (Li et. al 2018). Aliquots of the K<sub>2</sub>SO<sub>4</sub> extract were immediately frozen at -20°C and later analyzed for nitrate by reacting with vanadium (III) chloride according to Doane and Horwath (2003); and ammonium via the Berthelot reaction as laid out in Rhine et al. (1998). Available calcium, phosphorus and sulfur were measured on 2 mm sieved air-dried samples using the Mehlich-3 soil test (Mehlich, 1984). Total soil carbon and nitrogen values were measured on a CS 4010 Costech Elemental Analyzer (Costech Analytical Technologies) using air-dried, ball milled samples. 2018 carbon and nutrient stocks were calculated using depth-weighted sums (Tautges et. al 2019) with bulk density values from 2012. Aggregation measurements were carried out using the method outlined in Wang et al. (2017), adapted from the wet-sieving method outlined in Elliott (1986). Soils were gently passed through an 8mm sieve, and a 50g representative sample was submerged in room temperature water on top of a 2 mm sieve. This sieve was moved up and down for 2 min (50 submersions per minute) using an audio metronome to keep track of the number of submersions. The soil and water passed through the 2mm sieve were gently transferred by rinsing onto a 250 µm sieve and submerged again. The process was repeated using a 53 µm sieve to generate 4 aggregate size fractions (8 mm-2 mm, 2 mm-250 µm, 250 µm-50 µm, >50 µm) which were rinsed into pre-weighed aluminum pans, oven-dried at 60 °C, and weighed. Mean weight diameter of the aggregate fractions was calculated as the weighted average of the four aggregate size fractions (van Bavel, 1950).

## 2.6 Phospholipid Fatty Acid (PLFA) Analysis

PLFA analysis was carried out on 2018 samples using the high-throughput PLFA analysis method outlined in Buyer and Sasser (2012). Briefly, freeze-dried aliquots were extracted using Bligh-Dyer extractant. Phospholipid fractions were separated from the neutral lipid and glycolipid fractions using solid phase extraction columns. Phospholipids were then dried under N<sub>2</sub> gas, transesterified, and methylated. After methylation, the samples were dried again with N<sub>2</sub> gas and redissolved in hexane containing a known concentration of an internal standard (19:0) (Microbial ID, Newark, DE, USA). PLFAs were identified using the Sherlock software from Microbial Identification Systems and quantified using a gas chromatograph equipped with a flame ionization detector. A total of 56 different PLFAs were identified. PLFAs were assigned to Gram-positive, Gram negative, Cyclopropyl precursors, Saturated and Monounsaturated groups as outlined in Bossio and Scow (1998) (Supplementary Table [+A1](#)).

## 2.7 Hydraulic Conductivity and Moisture Content

Three 20 cm<sup>3</sup> cores were collected in September 2018 for saturated hydraulic conductivity from each plot that had been under tomato in 2017-2018 (9 plots out of a total of 189 plots). Cores were taken from a depth of 35 cm. Unfortunately, two cores were damaged during measurement, giving a total of 25 cores measured from the three treatments. Care was taken to transport the cores in foam holders to avoid creating compaction or preferential flow paths in transit. Cores were stored at 5 °C until measurement. A KSAT device was used to measure the cores with a falling head technique per the manufacturers manual and conductivity data was normalized to 20 °C using the Ksat software from the manufacturer (Meter Group, Pullman, Washington USA).

Soil moisture content was measured with a multi-depth profile capacitance probe in carbon fiber access tubes that were installed according to the manufacturer's recommendations with great care taken to avoid air gaps along the tube (PR 2/6, Delta-T Devices, Cambridge, UK). The factory calibration of the profile probe was used with an accuracy of  $\pm 0.04 \text{ m}^3 \text{ m}^{-3}$ . Volumetric soil moisture was measured at six depths (10, 20, 30, 40, 60, 100cm) (PR 2/6, Delta-T Devices, Cambridge, UK). Access tubes were installed in the field with a custom auger taking care to make the holes smooth and straight according to the manufacturer's recommendations. A total of 27 tubes were installed, with 3 tubes per subplot for a total of  $n = 9$  per treatment (ORG, CONV+WCC, CONV). The measurements were made on 8 dates between January 12 - March 1, 2019. Data was processed using R, and soil moisture depth from 10-100 cm was calculated using trapezoidal integration.

## 2.8 Fourier transform infrared Spectroscopy

Fourier transform infrared (FTIR) spectra of soil samples from 1993 and 2018 were collected using diffuse reflectance infrared Fourier transform spectroscopy (DRIFT; PIKE Technologies EasiDiff) with soil (air dried) diluted to 10% with KBr (~~Deiss et~~

al., 2020). All FTIR (Deiss et al., 2020). Spectra from 1993 samples were collected from air-dried, homogenized, archived soils from the Century Experiment Archive, while 2018 spectra were collected from air-dried, homogenized samples taken in 2018. 1993 spectra from 15-30 cm and 30-60 cm were combined into a single 15-60 cm spectra via weighted average for comparison with 2018 samples. The variation between these averaged 15-30 and 30-60 cm soils was found to be negligible for all three systems (Figure A6). All DRIFT spectra were collected using a Thermo Nicolet 6700 FTIR spectrometer (Thermo Scientific) using 256 scans,  $4\text{ cm}^{-1}$  resolution, and a DTGS detector. Three replicate samples were used, and average spectra were created for analysis. ~~Spectral subtractions were performed using Omnic 9.8.286 (Thermo Fisher Scientific).~~ Peak intensity ratios of aromatic to carboxyl moieties [ $\nu(\text{C}=\text{C})$ :  $\nu(\text{COO}^-)$  ( $1662\text{ cm}^{-1}$ : $1631\text{ cm}^{-1}$ )] were calculated using peak areas.

While FTIR is not a strictly quantitative tool for identifying specific compounds in mixed samples, it can be used pseudo-quantitatively because the absorption of IR light by a specific molecular bond at a specific electromagnetic frequency follows the Beer-Lambert Law (Beer's Law) (e.g., Margenot et al 2016, Smith 2001). Therefore, the height and area of a spectral peak are proportional to the abundance of molecules in a sample (linear relationship) and comparing the presence and absence of peaks and the relative differences in spectral contributions from each peak in a subtraction can suggest differences in C chemistry. However, it is important to note that spectral reflectance can lead to some non-linearity in concentration and absorbance, and thus pseudo quantification. Previous studies with DRIFTS in both the near-infrared (Dalal and Henry, 1986) and mid-infrared regions (Demyan et al., 2012; Margenot et al., 2015; West et al., 2020; Deiss et al., 2021) have shown direct associations between soil organic carbon concentration and absorbance at specific frequencies (depicted as peak height or area of single peaks or peak ratios). Spectral subtractions were performed using Omnic 9.8.286 (Thermo Fisher Scientific) and corrected for non-linearity of concentration and absorbance by using the Kubelka-Munk (KM) function. Plots of FTIR spectra were made using Origin 2018b (OriginLab Corporation). Subtractions were performed in two ways: 1) mean spectra, for each treatment and depth, of the 1993 spectra were separately subtracted from the corresponding 2018 spectra to reveal C chemistry changes over this period; and 2) the 2018 mean spectra, for each depth, were subtracted (ORG-CONV, ORG-CONV+WCC, CONV+WCC-CONV) to show the difference in C chemistry by treatment.

## 2.9 Statistical Analysis

All data analysis and graph production were done using R v. 4.0.2, (R Core Team, 2020) using the tidyverse package (Wickham et al., 2019). ~~Analysis of variance (ANOVA) was conducted using a linear model to determine the effects of management system, depth, and time point. Statistically significant~~ 4.0.2, (R Core Team, 2020) using the tidyverse package (Wickham et al., 2019). Analysis of variance (ANOVA) was conducted using a linear model to determine the effects of management system, depth, and time point. We first checked for normality and assumptions of the linear model prior to ANOVA, then fit a mixed effect model with "block" as a random effect. Since "block" was not significant for any of the variables measured, we removed it from the model. Statistical differences between management systems were analyzed separately for each depth using paired

## 390 3 Results

### 3.1 Nutrient Inputs


The cumulative estimated aboveground carbon input over 25 years was 186 Mg ha<sup>-1</sup>, 123 Mg ha<sup>-1</sup> and 113 Mg ha<sup>-1</sup> for ORG, CONV+WCC and CONV systems respectively. ~~Due to the combination of compost input and slightly increased and~~ Averaged per year over 25 years, C inputs to each system were 7.44 Mg ha<sup>-1</sup>, 4.92 Mg ha<sup>-1</sup> and 4.52 Mg ha<sup>-1</sup> for ORG, CONV+WCC and CONV systems respectively (Supplementary Table A4). Due to the combination of compost input and cover crop input, residue and root inputs, ORG systems received approximately 1.5x more carbon than CONV+WCC. Although CONV+WCC produced similar amounts of tomato residue and less maize residue than CONV, the presence of cover crops meant that CONV+WCC systems received 1.1x more carbon than CONV systems.

## 400 3 Results

### 3.1 Nutrient Inputs

The cumulative estimated aboveground carbon input over 25 years was 186 Mg ha<sup>-1</sup>, 123 Mg ha<sup>-1</sup> and 113 Mg ha<sup>-1</sup> for ORG, CONV+WCC and CONV systems respectively. Averaged per year over 25 years, C inputs to each system were 7.44 Mg ha<sup>-1</sup>, 4.92 Mg ha<sup>-1</sup> and 4.52 Mg ha<sup>-1</sup> for ORG, CONV+WCC and CONV systems respectively (Supplementary Table A4). Due to the combination of compost and cover crop residue and root inputs, ORG systems received approximately 1.5x more carbon than CONV+WCC. Although CONV+WCC produced similar amounts of tomato residue and less maize residue than CONV, the presence of cover crops meant that CONV+WCC systems received 1.1x more carbon than CONV systems.

Due to combined N inputs from cover crop and compost, ORG systems received 1.4x as much external N inputs (7.5 Mg ha<sup>-1</sup>) as CONV+WCC systems (5.4 Mg ha<sup>-1</sup>), and 1.65x as much N as CONV systems (4.5 Mg ha<sup>-1</sup>). External N inputs to CONV+WCC systems were close to 1 Mg ha<sup>-1</sup> higher than CONV systems over 25 years, with 40% of the external N inputs to CONV+WCC systems coming from the decomposition of cover crop residue, and the other 60% from mineral fertilizer application, compared to 100% of total N inputs in the CONV coming from mineral fertilizer application. ORG systems received over 3x as much phosphorus via compost (3.23 Mg ha<sup>-1</sup>) as CONV+WCC (1.09 Mg ha<sup>-1</sup>) and CONV (0.99 Mg ha<sup>-1</sup>) did from P fertilizer. ORG systems also received 1.15 Mg ha<sup>-1</sup> of sulfur from compost, approximately 0.5x as much as CONV+WCC (2.19 Mg ha<sup>-1</sup>) or CONV (1.98 Mg ha<sup>-1</sup>) systems received.









Figure 1a-d. Total aboveground carbon, nitrogen, phosphorus, and sulfur added per plot to ORG, CONV+WCC and CONV systems between 1993-2018. All values are given on a mass basis (Megagrams/hectare).

### 3.2 Soil Carbon Content Changes over 25 years

Carbon stocks in the 1 m profile of ORG systems showed ~~a marginally significant~~ increase of ~19 megagrams/hectare from 1993-2018 ( $p=0.06$ ) (Figure 2). Most of this carbon gain was concentrated in the 0-15 cm ( $\sim 5 \text{ Mg ha}^{-1}$ ,  $p<0.01$ ) and 15-60 cm depths ( $\sim 10 \text{ Mg ha}^{-1}$ ,  $p=0.1$ ), ~~but~~. Due to the large amount of variation present in these observations and the limited number of replicates, it was difficult to spot strong trends in carbon stock changes. The bottom 60-100 cm also showed ~~a non-significant~~ increase ( $\sim 3 \text{ Mg ha}^{-1}$ ,  $p=0.26$ ). No significant changes in carbon stocks in the 1 m profile were noted in CONV or CONV+WCC systems from 1993-2018 ( $p=0.47$ ,  $p=0.51$ ). When depth intervals were considered separately, only CONV systems showed a decrease in C stocks ( $\sim -3 \text{ Mg ha}^{-1}$ ), at the 0-15 ( $p<0.01$ ) depth (Figure 3). CONV+WCC systems did not show ~~significant~~ a clear trend of C decreases ~~decrease~~ at any individual depth ~~with the significance testing used~~.







Figure 2. Carbon stocks of the 1m profiles of ORG, CONV+WCC and CONV systems from 1993 to 2018. Carbon stocks are given in  $\text{Mg ha}^{-1}$ . Error bars denote standard error. Please note that all systems transitioned from furrow to drip irrigation in 2014.








Figure 3. Change in carbon stocks of ORG, CONV+WCC and CONV systems from 1993-2018 by depth. Values were obtained by subtracting carbon stocks in 1993 from 2018 stocks for individual systems, and then averaging by management system. Error bars denote standard error. (\* = significantly different from 0, p-value <0.05, + = significantly different from 0, p-value 0.05<x<0.1)

### 3.3 Moisture Content, Hydraulic Conductivity, and Cover Crop Roots

Cover cropped systems (ORG and CONV+WCC) stored approximately 10% more water than non-cover cropped systems (CONV) in the upper 1m of the soil profile during the 2019 winter (Fig 4). There was no difference in moisture content between ORG and CONV+WCC systems. Averaged hydraulic conductivity measurements showed no significant differences among all three systems, but treatments with cover crops (ORG and CONV+WCC) had values that spanned 3 orders of magnitude compared to treatments without cover crops (CONV) (Fig. 5).



455

Figure 4. Sum of moisture content (in ~~mm~~mm) in the upper 1m of ORG, CONV+WCC and CONV profiles during the Jan -Mar 2019 winter season. Error bars represent standard error.







Figure 5. Saturated hydraulic conductivity (cm/day) in ORG, CONV+WCC and CONV systems taken in August 2018.

3.4 Soil Nutrient Content: ~~Dissolved~~Extractable Organic Carbon, Mineral Nitrogen, Phosphorus, Sulfur

Composted systems (ORG) had ~~significantly~~ higher amounts of ~~dissolved~~extractable organic carbon (~~DOCEOC~~) ( $p<0.01$ ), plant available phosphorus ( $p<0.01$ ) and sulfur ( $p<0.01$ ) in the 1m profile than non-composted systems (CONV+WCC and CONV) during the 2018-2019 year (Fig. 6). These differences were most pronounced in the upper 15 cm, where ORG systems had approximately 2x more ~~DOCEOC~~ ( $p<0.01$ ), 3x more phosphorus ( $p<0.01$ ) and 1.75x more sulfur ( $p<0.01$ ) than CONV+WCC or CONV systems (Supplementary Figure 1).







Figure 6a-d. Dissolved/Extractable organic carbon, mineral N (NO<sub>3</sub>+ NH<sub>4</sub><sup>+</sup>), phosphorus and sulfur in 0-100cm profiles of ORG, CONV+WCC and CONV systems over the Feb 2018- Feb 2019 season. All values are given on a mass basis (kilograms/hectare). Error bars represent standard error.

CONV+WCC systems had significantly more mineral nitrogen (NO<sub>3</sub>+NH<sub>4</sub>) than CONV systems during the June and August timepoints (p=0.04), with up to 3.5x more mineral N than CONV systems mid-season, and 1.6x more mineral N at harvest. ORG systems trended towards higher mineral nitrogen during the April - September growing season but were not significantly different from CONV the magnitude of this difference was small (p=0.17).

Nutrient values showed large seasonal variation, with the highest levels of carbon, and nitrogen and sulfur observed during the June timepoint. DOC and highest sulfur levels at the August timepoint. EOC, mineral nitrogen, and sulfur values were lowest during the winter (Nov - Feb), which coincided with the period of highest rainfall. Phosphorus levels increased slightly throughout the 2018-2019 year.

485 Differences among systems and seasonal variation were also noted at a depth of 60 cm. ORG systems had ~~significantly~~ more ~~DOCEOC~~ (p<0.01), phosphorus (p<0.001), and sulfur at 60-100 cm than CONV+WCC or CONV systems. Mineral N values ~~were~~~~did~~ not ~~significantly different~~~~show large differences~~ between any of the three systems at 60-100 cm, though ORG and CONV+WCC systems trended higher during the growing season (Figure 7).

490






Figure 7a-d. **DissolvedExtractable** organic carbon, mineral N, phosphorus, and sulfur stocks at 60-100 cm in ORG, CONV+WCC and CONV systems over the Feb 2018- Feb 2019 season. All values are given in kg/ha. Error bars represent standard error.

### 495 3.5 Aggregation

There was no significant difference in MWD of aggregates between all three systems at any depth (Supplementary Figure [3A2](#)).


### 3.6 ~~Soil Carbon~~SOM Composition via FTIR

500 Spectral subtractions of 1993 ~~and from~~ 2018 FTIR spectra revealed positive peaks (increased absorbance) from 1900 to 1200 cm<sup>-1</sup> in all systems, indicating an increase in C functional groups within this region (e.g., aromatic, carboxyl) (Fig. 8A). FTIR band assignments are presented in Supplementary Table ~~B-~~A2. All treatments showed positive peaks indicating an increase in carboxylate functional groups between 1993 and 2018, as denoted by bands at 1625 cm<sup>-1</sup> and 1400 cm<sup>-1</sup> (Fig 8a). However,

ORG and CONV+WCC showed these distinct peaks at 15-60 and 60-100 cm depths, while CONV systems showed distinct peaks only at the 0-15 and 15-60 cm depths. CONV systems also showed a lower aromatic:carboxylate peak intensity ratio at all depths than ORG and CONV+WCC systems from 1993-2018 (Table 1).

A distinct increase in aromaticity in ORG systems at all depths between 1993 and 2018 was denoted by bands at  $1666\text{ cm}^{-1}$  and at  $1602\text{ cm}^{-1}$  (Fig 8A). There is little noticeable difference in aromaticity in both the CONV+WCC and CONV spectra over this time period. All treatments and depths showed an increase in carboxylate functional groups between 1993 and 2018, denoted by bands at  $1625\text{ cm}^{-1}$ ,  $1400\text{ cm}^{-1}$  and  $1384\text{ cm}^{-1}$ .

Comparison of 2018 spectra showed higher amounts of aromaticity ( $1666$  and  $1602\text{ cm}^{-1}$ ) for the top 0-15 cm in the order  $\text{ORG} > \text{CONV+WCC} > \text{CONV}$  (Fig 8B). ORG systems also showed more aromaticity at 15-60 cm than CONV+WCC or CONV, but this trend did not apply to the bottom 60-100 cm. Instead, higher amounts of carboxylate ( $1631\text{ cm}^{-1}$ ) functional groups were noted from 60-100 cm in the order  $\text{ORG} > \text{CONV+WCC} > \text{CONV}$ .



ORG and CONV+WCC systems showed distinct positive peaks associated with carboxylate functional groups at the 60-100 cm depths in 2018 when compared with CONV systems (bands at  $1631\text{ cm}^{-1}$ ), and slightly higher peaks associated with aromatic functional groups from 0-15 cm for CONV+WCC and 15-60 cm for ORG (bands at  $1662\text{ cm}^{-1}$ ) (Figure 8b). ORG also showed positive peaks associated with aromatic functional groups from 0-15 and 15-60 cm when compared to CONV+WCC (bands at  $1662$  and  $1602\text{ cm}^{-1}$ ) in 2018. Aromatic:carboxylate ratios provide an indication of the intensity of carboxyl peaks relative to aromatic peaks, which can be related back to concentrations of these functional groups in the sample. A lower aromatic:carboxyl ratio can indicate either more carboxyl or less aromatic functional groups, while a higher ratio can



mean increased aromatic or decreased carboxyl groups. Aromatic:carboxylate peak intensity ratios decreased with depth for ORG and CONV systems when looking at changes from 1993-2018, but CONV+WCC ratios increased with depth (Table 1).





Figure 8a,b. **FTIRDRIFT** spectral subtractions for the 1900-1200 cm<sup>-1</sup> range comparing (A) 2018-1993 spectra for ORG, CONV+WCC, and CONV, and (B) ORG, CONV+WCC, and CONV spectra in 2018. Spectra are plotted with Kubelka-Munk units on a common y-axis scale and are offset from one another for ease of comparison.

	<u>Depth</u>	<u>Peak Intensity Ratio (1662 cm<sup>-1</sup> : 1631 cm<sup>-1</sup>)</u>		
		<u>ORG</u>	<u>CONV+WCC</u>	<u>CONV</u>
<u>Subtraction: 2018-1993</u>	<u>0-15 cm</u>	<u>1.38</u>	<u>1.22</u>	<u>0.45</u>
	<u>15-60 cm</u>	<u>1.21</u>	<u>1.40</u>	<u>0.77</u>
	<u>60-100 cm</u>	<u>1.17</u>	<u>2.50</u>	<u>0.014</u>
		<u>ORG-CONV</u>	<u>ORG-(CONV+WCC)</u>	<u>(CONV+WCC)-CONV</u>
<u>Subtraction by treatment: 2018</u>	<u>0-15 cm</u>	<u>1.39</u>	<u>1.18</u>	<u>0.46</u>
	<u>15-60 cm</u>	<u>1.20</u>	<u>1.42</u>	<u>0.47</u>
	<u>60-100 cm</u>	<u>0.38</u>	<u>0.64</u>	<u>0.45</u>

**Table 1. Peak Intensity Ratios for aromatic (1662 cm<sup>-1</sup>) to asymmetric carboxyl (1631 cm<sup>-1</sup>) groups in spectral subtractions.**

### 3.7 Microbial Biomass and Stress Indicators - July 2018

Microbial biomass decreased with depth in all systems. ORG and CONV+WCC systems had more microbial biomass at 0-15cm than CONV systems (p=0.04 & p=0.04 respectively), while ORG systems had more microbial biomass at the 15-60 cm depth than CONV+WCC or CONV systems (p=0.03 & p=0.06 respectively). Saturated: Unsaturated fatty acid ratio and Cyclopropyl 19: precursor ratio increased with depth, with CONV systems ~~havingshowing a weaker trend of higher-trending~~ Cy19: pre (p=0.12) and saturated: unsaturated fatty acid ratios (p=0.07) than ORG at 60-100 cm. Gram+: Gram- ratio also increased with depth, with CONV systems having a higher ratio than ORG or CONV+WCC systems at 60-100 cm (p=0.01 and p=0.01 ~~repsectivelyrespectively~~).






Figure 9a-d. Microbial biomass and PLFA stress indicators measured in ORG, CONV+WCC and CONV systems during mid-season (July 2018). Ratios are unitless, while microbial biomass is given in kg/ha.

#### 550 4 Discussion

The ~19 Mg/ha increase in SOC over the 1m ORG profile after 25 years was attributed to a synergistic effect between cover crops and compost, which resulted in the movement of mobile carbon and nutrients deeper into the soil profile. We believe that high concentrations of mobile C and essential nutrients for microbial activity provided by the compost, combined with the easier movement of water downward associated with a history of cover-cropping, helped transport the material needed to build

555 C in the subsurface soil.

#### 4.1 Cover crop roots ~~increased~~increase water storage and movement ~~in subsurface soils~~into subsoils

Cover crops ~~increase both infiltration and hydraulic conductivity by increasing soil macroporosity and pore connectivity, particularly over longer time scales (Scott et al., 1994). This increase occurs due to an improvement in soil structure through aggregate formation, and a reduction in compaction due to increased organic matter content and formation of root channels (Chen and Weil, 2010; Franzluebbers, 2002). We measured a higher moisture content in CONV+WCC and ORG than CONV systems during the winter growing season, likely due to the presence of more water-filled spaces (Figure 4). However, we found no significant difference in aggregate MWD among systems (Supplementary Figure 4), potentially ruling out increased pore space due to aggregation as the cause of increased water content. We did find, however, more variable hydraulic conductivity in the two systems with cover crops (ORG and CONV+WCC, Figure 5). Roots may increase macroporosity as they decay (Ghestem et al., 2011), and increased water movement through these macropores can result in hydraulic conductivity values that can range over three orders of magnitude (Øygarden et al., 1997) similar to what we observed. We believe the more variable hydraulic conductivity and increased moisture content in ORG and CONV+WCC than CONV systems are due to the deeper, more abundant root-derived macropores from cover crops.~~

~~Due to the fact that tillage in all systems would likely eliminate differences among them in the top 30 cm, we would expect any differences in macroporosity and infiltration among treatments to be most affected by those roots that extend below the 30 cm plow layer (Hangen et al., 2002). Indeed, the mixture of cover crops in ORG and CONV+WCC included oats (~65–85 cm), fava beans (~52–70 cm) and vetch (~30–85 cm), all of which have extensive rooting deeper than 30 cm (Fan et al., 2016).~~

~~Cover crops increase both infiltration and hydraulic conductivity in fine textured soils by increasing soil macroporosity and pore connectivity, particularly over longer time scales (Scott et al., 1994; Haruna et al., 2018; Çerçioğlu et al., 2019; Gulick et al., 1994). Impacts of cover crops have been noted in previous work done in Russell Ranch soils: a cover crop mix of purple vetch (*Vicia benghalensis* L.), common vetch (*Vicia sativa* L.) and oats (*Avena sativa* L.) increased soil moisture-holding capacity during saturated conditions (Joyce et al. 2002); a cover crop of common vetch produced no changes in bulk density after 10 years (Colla et al. 2000) and a wheat (*Triticum aestivum* L.) cover crop increased infiltration by 43% and decreased DOC export by 54% in a furrow irrigated system, causing the soil profile to become a DOC sink (Mailapalli et al., 2012). Proposed mechanisms are that cover crops increase infiltration and hydraulic conductivity by increasing soil structure through aggregate formation, reduced soil crusting, and reduced soil compaction due to increased organic matter content and formation of root channels (Chen and Weil, 2010; Franzluebbers, 2002).~~

~~We measured a higher moisture content in CONV+WCC and ORG than CONV systems during the winter growing season, likely due to the presence of more water-filled spaces (Figure 4). Other potential, albeit less likely, explanations for increased moisture content could include lateral subsurface flow (unlikely due to the <1% slope of this field) and differences in runoff and runoff (also unlikely due to low slope). However, we found no difference in aggregate mean weight diameter (MWD) among systems (Supplementary Figure A2), potentially indicating that increased intra-aggregate pores were not behind the increases in moisture content. Instead, these differences in moisture content may result from biopores created by roots (Hangen et al., 2002) as our cover crop mix is known to have extensive root networks that extend deeper than the 30 cm plow layer. Rooting depths are ~65–85 cm in oats, ~52–70 cm in fava beans, and ~30–85 cm in vetch (Fan et al., 2016).~~

Despite the presence of cover crops, we found no significant difference in aggregate MWD among systems. This may be because root-induced soil alterations, such as aggregation, are highly localized and dependent on the root architecture of the cover crops. Specifically, cover crops with prominent tap roots, such as fava bean, are effective at creating continuous biopores, while fibrous roots such as in oats and hairy vetch, are particularly effective at promoting soil aggregate formation (Ogilvie et al., 2021). Therefore, the mixture of cover crops planted at the site likely resulted in a large amount of variation in aggregation and pore connectivity and may have resulted in the non-significant aggregate MWD values.

While mean hydraulic conductivity values were also not significantly different between treatment systems, hydraulic conductivities were more variable in the two systems with than without cover crops (Figure 5). Roots may increase macroporosity by opening channels as they decay (Ghestem et al., 2011), and increased water movement through these macropores can result in hydraulic conductivity values that can range over three orders of magnitude (Øygarden et al., 1997) similar to what we observed. In addition, the sample size used for  $K_{sat}$  measurements (cross-sectional area of 250 cm<sup>2</sup>) may be too small to capture the effects of cover crop roots, whose impacts are likely to be detected at a larger scale (Ozelim and Cavalcante, 2017). It is well recognized that hydraulic conductivity measurements can vary widely across fields and landscapes (Rahmati et al. 2018) and often do not reflect the presence of macropores (Brooks et al. 2004). Though our measurements do not reveal statistically significant differences between treatments, the scattered high-permeability zones in the cover-crop treatments likely play a role in rapid moisture redistribution and may explain the elevated deep moisture contents in ORG and CONV+WCC plots compared to CONV plots. We therefore attributed the more variable hydraulic conductivity and increased moisture content in ORG and CONV+WCC than CONV systems to the deeper, more abundant root-derived macropores from cover crops.

#### 4.2 Compost + Cover Crops increased ~~DOC~~the amount of EOC and ~~aromatic carbon~~carboxylate functional groups in subsurface soils

Compost application is associated both with high levels of mobile DOC (Wright et al., 2008; Zmora-Nahum et al., 2005), as well as a large proportion of aromatic functional groups, attributed to high amounts of lignin and other biomolecules (Leifeld et al., 2002). Research into the transport of organic molecules, such as described by the “cascade theory”, has shown that “fresh” carbon inputs are preferentially retained within the top 30 cm (Liebmann et al., 2020) and that aromatic moieties are rapidly removed from the soil solution at the surface (Leinemann et al., 2018). This removal may be a function of the relatively low solubilities of non-polar aromatic functional groups (Maxin and Kogel-Knabner, 1995), as well as their tendency to partition into other non-polar, insoluble organic matter (Pignatello, 1999). When these aromatic functional groups are eventually oxidized by microbes, the higher solubility of carboxylate and other O-rich functional groups may allow for greater C transport. In the “cascade theory”, carbon is transported downwards via a series of successive sorption, desorption, processing, and transport steps resulting in highly processed, O-enriched carbon accumulating in the subsoil (Kaiser and Kalbitz, 2012). As this stepwise process is triggered by fresh DOC inputs, the regular application of DOC-rich compost, combined with increased hydraulic conductivity due to WCC roots, can accelerate this “cascade”, leading to greater subsurface C transport. Increased DOC levels (Figure 6a), more aromatic carbon in the top 60 cm (Figure 8a), and relatively more oxidized carboxylate carbon in the bottom 60–100 cm of ORG plots point to an accelerated “cascade” process in ORG systems relative to CONV+WCC and CONV systems.



630 CONV+WCC systems received more C than CONV systems, but the application of cover crops did not increase DOC or aromatic carbon content deeper than 15 cm when compared to CONV (Figure 7a). This may be because microbial processing of cover crop and crop residues need to occur at the surface before these residues can be transported deeper as DOC (Chantigny, 2003; White et al., 2020b). The increase in carboxylate carbon in CONV+WCC and CONV plots, particularly in the bottom 60–100 cm after 25 years, does point to a “trickle” of soluble carbon inputs over time, albeit less than in ORG plots. Though our results do not provide sufficient data to determine the stability of the carbon at 60–100 cm, carboxylate functional groups would be more likely to form mineral-associated organic matter through cation bridging (Aquino et al., 2011) or association with charged surfaces, which could lead to C stabilization and increased storage times (Cotrufo et al., 2013; Leinemann et al., 2018).

635 Compost application is associated both with elevated soluble C (Wright et al., 2008; Zmora-Nahum et al., 2005) and a large proportion of aromatic functional groups derived from lignin and other biomolecules (Leifeld et al., 2002), while cover crop residue can increase dissolved and water extractable organic carbon both at (<15 cm) and below (>15cm) the soil surface (Singh et. al 2021 - cereal rye and hairy vetch). Research into the transport of soluble organic molecules in soil has led to development of the cascade theory (Kaiser and Kalbitz 2012) which describes the stepwise sorption, desorption, transport and processing of carbon molecules as they move deeper into the soil profile. This stepwise transformation and transport of carbon means that fresh carbon inputs are preferentially sorbed within the top 30 cm (Liebmann et al., 2020) and aromatic moieties are rapidly removed from the soil solution at the surface (Leinemann et al., 2018). This removal may be a function of the relatively low solubilities of non-polar aromatic functional groups (Maxin and Kogel-Knabner, 1995), as well as their tendency to partition into other non-polar, insoluble organic matter (Pignatello, 1999). When these aromatic functional groups are eventually oxidized by microbes, the higher solubility of carboxylate and other O-rich functional groups may allow for greater C transport. As the carbon cascade is triggered by fresh C inputs, the regular application of soluble C-rich compost and WCC residue, combined with increased hydraulic conductivity due to WCC roots can accelerate the process leading to greater subsoil C transport.

650 Increased EOC levels (Figure 6a) in ORG plots and relatively more oxidized carboxylate carbon in the bottom 60–100 cm of ORG and CONV+WCC plots relative to CONV plots (Figure 8b) point to an accelerated cascade process in these systems relative to CONV systems. Though our results do not provide sufficient support to determine the stability of the increased carbon stocks at 60–100 cm, carboxylate functional groups’ ability to form mineral-associated organic matter through cation bridging (Aquino et al., 2011) or associate with charged surfaces would promote MAOM formation and increase C storage times (Cotrufo et al., 2013; Leinemann et al., 2018).

660 Cover crops are associated with elevated EOC levels and more aromatic functional groups in topsoil SOC (Ding et. al 2005, Zhou et. al 2012). Application of cover crops in CONV+WCC did not increase EOC content deeper than 15 cm compared to CONV (Figure 7a) but did have an impact on carboxylate functional group presence at the 60-cm depth (Figure 8b), possibly indicating that cover crop residues are associated with smaller inputs of soluble C that are not as easily detected as the larger soluble C inputs coming from the compost. Additionally, the trend of increasing aromatic:carboxylate ratio with depth in CONV+WCC systems (Table 1) indicates that CONV+WCC systems may be accumulating more aromatic carbon relative to

carboxylate carbon in deeper soil layers. This accumulation of aromatic carbon in CONV+WCC subsoils may be due to cover crop root residue introducing lignin and cellulose directly into the subsoil, while the carboxylate-C obtained from the decomposition of surface residues can be potentially mineralized before being transported deeper as DOC (Chantigny, 2003; White et al., 2020a).

#### 4.3 Compost + Cover crops increased nutrient availability and decreased microbial stress in ~~subsurface soils~~subsoils

The higher N, P and S values noted in ORG subsurface soils (Figure 7a-d) can be attributed both to the higher organic N, P, and S inputs associated with compost (Preusch et al., 2002), as well as the increased mobility of these inputs. Organic phosphorus is more mobile than mineral P (Laos et al., 2000; Sharpley and Moyer, 2000), and mineralization of organic N and S into their soluble forms of nitrate and sulfate could also facilitate their movement (Edwards, 1998; Vinten et al., 1994).

Greater C and nutrient input can also explain the lower microbial stress ratios observed in subsurface ORG soils (Fig 6b, d). Higher values for these ratios, such as those observed in CONV plots, have been associated with nutrient and energy limitation (Bossio and Seow, 1998; Petersen and Klug, 1994). While adding compost did not appear to increase microbial biomass in ORG plots at 60-100 cm, it did increase biomass at 0-15 and 15-60 cm depths, attributed to more favorable nutrient stoichiometry for biomass formation (Kirkby et al., 2011; Richardson et al., 2014). Increased surface microbial biomass is an important potential source of C and other nutrients to subsurface layers through cell lysis from predation and wet-dry cycles (Bonkowski, 2004; Xiang et al., 2008) and is associated with increased stabilized carbon storage through microbial necromass formation (Buchmann and Schaumann, 2018; Jilling et al., 2020), contributing to the SOC increase observed in ORG systems (Figure 2b).

#### 4.4 Cover crop carbon was not as effective as compost carbon in increasing subsurface SOC

The SOC increases noted under the ORG system after 25 years are attributed to the increased mobility of compost added carbon and nutrients, as well as more microbial processing at the soil surface. Larger SOC increases under compost + cover crops relative to cover crops alone have also been noted in other long-term experiments (Brennan and Acosta-Martinez, 2017), indicating that carbon input from cover crops alone may not play as large of a role in increasing subsurface C. While cover crop biomass does represent significant C and N input to surface soils, the “paths” their roots create for mobile nutrients (either organic or mineral) to move downwards may be as important as their C and N inputs to subsurface SOC dynamics.

There is evidence that root-associated C inputs appear to be more effective than aboveground biomass in building SOC (Sokol and Bradford, 2019). The lack of an observed C increase in the CONV+WCC system was attributed to small inputs of C and N over time from cover crop roots priming decomposition of native SOC, potentially by stimulating phosphatases and accelerating MAOM breakdown (Cui et al., 2020; Mise et al., 2020). Additionally, common root exudates such as oxalic acid may have destabilized organomineral complexes (Keiluweit et al., 2015), making that carbon more accessible for decomposition. While any priming and destabilization of SOC due to cover crop roots would also be occurring in the ORG systems, we believe this was counteracted by the higher DOC inputs and more favorable nutrient stoichiometry for microbial biomass provided by the compost.

The trend of a SOC decrease over 25 years in CONV+WCC profiles is consistent with the results observed in Tautges and Chiartas et al. (2019), but we observed a smaller decrease in SOC stocks from 1993-2018 below 60 cm. This difference may be attributed to the high variability of soil C measurements (Kravchenko and Robertson, 2011; Wuest, 2014), but may also be due to the switch from furrow to drip irrigation in 2014. The reduction in water inputs that occurs when switching from furrow to drip can cause changes in C and N cycling enzymes (Schmidt et al., 2018), potentially increasing the prevalence of complex SOM, and reducing microbial mineralization. The potential for complex interactions of tillage, cover cropping, compost application and irrigation strategies in our experimental system indicate that care should be taken when applying these results to different soil types and climates.

The higher N, P and S values noted in ORG subsoils (Figure 7a-d) can be attributed both to the higher organic N, P, and S inputs associated with compost (Preusch et al., 2002), as well as the increased mobility of these inputs. Organic phosphorus is more mobile than mineral P (Laos et al., 2000; Sharpley and Moyer, 2000), and mineralization of organic N and S into their soluble forms of nitrate and sulfate could also facilitate their movement (Edwards, 1998; Vinten et al., 1994). Although soil microbial communities are primarily water and carbon limited (Soong et. al 2020), the addition of N, P and S in ratios like that found in soil organic matter may increase transformation of C inputs into SOM by up to 52% by promoting microbial anabolism (Coonan et. al 2020).

Greater C and nutrient inputs were associated with the lower Gram +: Gram - ratios observed in subsoil ORG soils (Fig 6b, d). Higher values for these ratios, such as those observed in CONV plots, have been associated with nutrient and energy limitation (Bossio and Scow, 1998; Petersen and Klug, 1994). Increases in these ratios represent an overall shift away from the thinner, more permeable cell membranes associated with Gram - bacteria and monounsaturated fatty acids; towards more tightly packed, less permeable cell membranes associated with Gram+ bacteria and saturated fatty acids (Silhavy et. al 2010). An increase in the Gram+:Gram- ratio has been associated with a decrease in easily available water and carbon (Fanin et. al 2019, Fierer et. al 2003b, Bossio et. al 1998), while an increase in the saturated: unsaturated ratio and cy17:pre ratios has been associated with low water potentials (-1.3 to -0.9 MPa) and potential dehydration (Moore-Kucera et. al 2007). The stress indicator trends in our data support our observations of increased soluble carbon and water content in ORG systems.

Adding compost and cover crop residue increases microbial biomass at the 0-15 and 15-60 cm but not 60-100 cm depth relative to CONV systems. This greater biomass increase in ORG than CONV plots was attributed to compost providing a favorable nutrient stoichiometry for biomass formation (Kirkby et al., 2011; Richardson et al., 2014). Increased microbial biomass in surface layers of soil is an important potential source of C and other nutrients to subsoil layers through cell lysis from predation and wet-dry cycles (Bonkowski, 2004; Xiang et al., 2008). It is associated with increased carbon storage through microbial necromass formation (Buchmann and Schaumann, 2018; Jilling et al., 2020), and we hypothesize that transport of microbial products downwards through the profile could have contributed to the SOC increase in the subsoil observed in ORG systems (Figure 2b).

#### **4.4 Compost + Cover Crops increased subsoil carbon stocks after 25 years, but Cover Crops alone did not**

We attribute the SOC increases under the ORG system after 25 years (Figure 2) to the increased mobility of compost-added carbon and nutrients combined with increased infiltration due to cover crop roots. Larger SOC increases under yard waste compost + cover crops relative to cover crops alone have also been noted in other California long term experiments on a loamy sand soil (White et. al 2020a - rye, fava bean, pea, common vetch, purple vetch), indicating that carbon input from cover crops alone may not play a large role in increasing subsoil C. While cover crop biomass does represent significant C and N input to

surface soils, the channels their roots create for mobile nutrients (either organic or mineral) to move downwards may be as important as their C and N inputs to subsoil SOC dynamics.

735 We noted significant seasonal variation in EOC, mineral N, P and S levels throughout the 2018-2019 growing season, though  
ORG plots consistently had higher EOC and P than CONV+WCC or CONV plots at all timepoints (Figures 6a,c). These  
soluble C and nutrient inputs likely peaked during the growing season due to the influence of compost application, root  
exudates and fertigation. Since the months of April-September are the driest months of the year at the study site, these C and  
740 nutrient inputs may have depended on irrigation water to be transported into subsoil layers, highlighting the importance of  
irrigation amounts and types (drip, furrow) to understanding changes in subsoil C stocks. The shifts in soluble C:N:P:S ratios  
during the course of the year may also indicate that carbon:nutrient stoichiometry during the growing season is more suitable  
for microbial biomass growth in these row cropped plots.

In contrast to the ORG system, SOC stocks did not significantly increase in the CONV+WCC plots after 25 years. While our  
FTIR results suggest that cover crop residues have an impact on subsoil carbon by increasing the proportion of carboxylate-C  
745 relative to the CONV system (Figure 8b), they do not suggest a clear reason behind the lack of an increase in SOC stocks. A  
possible hypothesis is that small inputs of C and N over time from cover crop roots primed decomposition of native SOC,  
potentially by stimulating phosphatases and accelerating MAOM breakdown (Cui et al., 2020; Mise et al., 2020). Additionally,  
common root exudates such as oxalic acid may have dispersed organomineral complexes (Keiluweit et al., 2015), making that  
carbon more accessible for decomposition. While any priming of SOC due to cover crop root exudates would also be occurring  
750 in the ORG systems, we believe this was counteracted by the higher EOC inputs and more favorable nutrient stoichiometry  
for microbial biomass provided by the compost.

We also observed a continual decline in SOC in subsoil in the conventional with cover crops treatment (CONV + WCC) as  
observed in Tautges and Chiartas et. al (2019); however, the rate of decline was lower over the last 7 years than in the first 19  
years of study. This slower decline in subsoil C stocks from 2012-2019 may be due to the switch from furrow to drip irrigation  
755 in 2014. Lower water inputs with drip reduces microbial activity and C and N cycling enzyme activities (e.g., beta-glucosidase  
and N-acetyl-glucosaminidase) in a large part of the bed in the surface of these same plots (Schmidt et al., 2018). This reduction  
in microbial activity potentially increased the prevalence of complex SOM by reducing microbial mineralization and may have  
facilitated greater DOC transport during the winter rainy season.


## 5 Conclusion

760 ~~We believe that the~~The combination of growing cover crops and compost amendment created a unique set of conditions  
conductive to carbon transport and ~~carbon~~ accumulation in the ~~subsurface~~subsoils of a tilled row crop rotation. This was, in  
part, likely due to increased hydraulic conductivity facilitated by ~~increased soil macropores created by~~ cover crop roots leading

to higher rates of transport of soluble C and nutrients from the surface to ~~subsurface soils~~ subsoil. In turn, higher transport led to increased C stocks and reduced levels of microbial stress. The accumulation of oxygen-rich carboxylate carbon in ~~subsurfacesubsoil~~ horizons under all treatments, attributed to an increase in microbially--processed carbon, ~~and the accumulation of aromatic carbon under compost application, attributed to the lower solubility of aromatic functional groups,~~ provide~~s~~es support for the “cascade theory” of carbon transport. These results demonstrate the potential for ~~subsurfacesubsoil~~ carbon storage in tilled agricultural systems, and highlight a potential pathway for increasing carbon transport, storage, and sequestration in ~~subsurface soils~~ subsoil layers.

## 6 Appendices

### 6.1 Appendix A






Figure A1. ~~Dissolved~~Extractable organic carbon, mineral N, phosphorus, and sulfur stocks at 0-15 cm in ORG, CONV+WCC and CONV systems over the Feb 2018- Feb 2019 season. All values are given in kg/ha. Error bars represent standard error.




Figure A2. Mean weight diameter of aggregates obtained by wet sieving for 0-15, 15-60 and 60-100 cm depth intervals in ORG, CONV+WCC and CONV systems.








Figure A3a,b. FTIR spectral subtractions for the 4000-1200  $\text{cm}^{-1}$  range comparing (A) 2018 -1993 spectra for ORG, CONV+WCC and CONV, and (B) ORG, CONV+WCC and CONV spectra in 2018.

(A)

(B)





**Figure A4a,b.** FTIR spectral subtractions for the 1900-1200  $\text{cm}^{-1}$  range comparing (A) 2018 -1993 spectra for ORG, CONV+WCC and CONV, and (B) ORG, CONV+WCC and CONV spectra in 2018.




Figure A5. Rainfall and soil temperature at 50 cm at Russell Ranch from Oct 2018 to Oct 2019. Data taken from <http://atm.ucdavis.edu/weather/uc-davis-weather-climate-station>




Figure A6. Example DRIFT spectra showing three replicate spectra, the mean spectra and the residual spectra for 1993 and 2008 samples.

Gram Positive	15:1 iso w6c; 15:0 iso; 15:0 anteiso; 16:0 iso, 17:1 iso w10c, 17:1 iso w9c; 17:1 anteiso w9c; 17:0 iso, 17:0 anteiso; 18:0 iso
Gram Negative	16:1 w9c; 16:1 w7c; 17:1 w8c; 17:0 cyclo w7c; 18:1 w7c; 19:0 cyclo w7c; 20:1 w9c; 21:1 w3c
Saturated	12:0; 14:0; 15:0; 16:0; 17:0; 20:0
Monounsaturated	16:1 w5c; 16:1 w7c; 18:1 w9c; 18:1 w7c
Cyclopropyl Indicator	19:0 cyclo w7c / 18:1w7c

Table A1 – PLFA (Phospholipid Fatty Acid) Assignments taken from Bossio and Scow (1998).

Wavenumber (cm <sup>-1</sup> )	IR Assignment
2800-3100	aliphatic $\nu_s(\text{CH}_2)$ , $\nu_{as}(\text{CH}_2)$ , $\nu_s(\text{CH}_3)$ , $\nu_{as}(\text{CH}_2)$
1700-1765	$\nu(\text{C=O})$
1666	aromatic $\nu(\text{C=C})$
1620-1631	$\nu_{as}(\text{COO})$
1602	skeletal $\nu(\text{C=C})$
1546	aromatic $\nu(\text{C=C})$
1417	$\delta(\text{C-H})$
1400	$\nu_s(\text{COO})$
1384	$\nu(\text{C-O})$ vibration aromatic and $\delta(\text{C-H})$ vibrations in $\text{CH}_3$ and $\text{CH}_2$

**Table A2 - FTIR Peak Assignments\* used for analysis of spectra**

\*Assignments taken from

Baes, A.U., Bloom, P.R., 1989. Diffuse reflectance Fourier transform infrared (DRIFT) of humic and fulvic acids. Soil Sci. Soc. Am. J. 53, 695–700. doi:10.2136/sssaj1989.03615995005300030008x;

Hesse, M., Meier, H., & Zeeh, B. (2005). Spektroskopische Methoden in der Organischen Chemie. (In German.) Georg Thieme Verlag, Stuttgart. doi:10.1002/pauz.19960250417

Parikh, S.J., A.J. Margenot, F.N.D. Mukome, F. Calderon, and K.W. Goynes. 2014. Soil Chemical Insights Provided through Vibrational Spectroscopy. *Adv. Agron-Adv. Agron*. 126:1-148

Orlov, D.S., 1986. Humus acids of soil. Rotterdam: Balkema. doi:10.1002/jpln.19871500116

<u>Soil Series</u>	<u>Rincon</u>	<u>Yolo</u>
<u>Soil Taxonomic Class</u>	<u>Fine, smectitic, thermic Mollic Haploxeralfs</u>	<u>Fine-silty, mixed, superactive, nonacid, thermic Mollic Xerofluvents</u>
<u>Horizon Designation</u>	<u>AP</u>	<u>AP</u>
<u>Depth (cm)</u>	<u>0-10</u>	<u>0-5</u>
<u>Texture</u>	<u>SiCL</u>	<u>SiL</u>
<u>pH</u>	<u>6.5</u>	<u>6.7</u>
<u>Organic Matter Content (%)</u>	<u>2.4</u>	<u>2.4</u>
<u>Clay Content (%)</u>	<u>31</u>	<u>30</u>
<u>Horizon Designation</u>	<u>A12</u>	<u>AP2</u>
<u>Depth (cm)</u>	<u>10-41</u>	<u>5-20</u>
<u>Texture</u>	<u>SiCL</u>	<u>SiL</u>
<u>pH</u>	<u>6.5</u>	<u>7.1</u>
<u>Organic Matter Content (%)</u>	<u>2.4</u>	<u>2.4</u>
<u>Clay Content (%)</u>	<u>31</u>	<u>30</u>
<u>Horizon Designation</u>	<u>B21t</u>	<u>A1</u>
<u>Depth (cm)</u>	<u>41-64</u>	<u>20-48</u>
<u>Texture</u>	<u>SC</u>	<u>SiL</u>
<u>pH</u>	<u>7</u>	<u>7.2</u>
<u>Organic Matter Content (%)</u>	<u>0.75</u>	<u>1.8</u>
<u>Clay Content (%)</u>	<u>40</u>	<u>30</u>
<u>Horizon Designation</u>	<u>B22t</u>	<u>A2</u>

<u>Depth (cm)</u>	<u>64-79</u>	<u>48-66</u>
<u>Texture</u>	<u>SC</u>	<u>SiL</u>
<u>pH</u>	<u>7.9</u>	<u>7.3</u>
<u>Organic Matter Content (%)</u>	<u>0.75</u>	<u>1.3</u>
<u>Clay Content (%)</u>	<u>40</u>	<u>30</u>
<u>Horizon Designation</u>	<u>B3tca</u>	<u>C1</u>
<u>Depth (cm)</u>	<u>79-102</u>	<u>66-84</u>
<u>Texture</u>	<u>SCL</u>	<u>SiL</u>
<u>pH</u>	<u>8</u>	<u>7.4</u>
<u>Organic Matter Content (%)</u>	<u>0.75</u>	<u>1</u>
<u>Clay Content (%)</u>	<u>40</u>	<u>28</u>
<u>Horizon Designation</u>		<u>C2</u>
<u>Depth (cm)</u>		<u>84-104</u>
<u>Texture</u>		<u>SiL</u>
<u>pH</u>		<u>7.4</u>
<u>Organic Matter Content (%)</u>		<u>0.8</u>
<u>Clay Content (%)</u>		<u>25</u>

**Table A3. Texture, pH, OM, and clay content for the Rincon and Yolo soil series found at Russell Ranch.**

<u>Input (Mg/Ha/yr)</u>	<u>CONV</u>	<u>CONV+WCC</u>	<u>ORG</u>
<u>Tomato Residue</u>	<u>1.73</u>	<u>1.57</u>	<u>1.54</u>
<u>Corn Residue</u>	<u>2.43</u>	<u>1.96</u>	<u>2.25</u>
<u>Compost</u>	<u>0</u>	<u>0</u>	<u>1.9</u>
<u>WCC Residue</u>	<u>0</u>	<u>1.086</u>	<u>1.48</u>
<u>WCC Nitrogen</u>	<u>0</u>	<u>0.09</u>	<u>0.11</u>
<u>Compost Nitrogen</u>	<u>0</u>	<u>0</u>	<u>0.19</u>
<u>Mineral N</u>	<u>0.18</u>	<u>0.13</u>	<u>0</u>
<u>Mineral Phosphorus</u>	<u>0.04</u>	<u>0.04</u>	<u>0</u>
<u>Organic Phosphorus</u>	<u>0</u>	<u>0</u>	<u>0.13</u>
<u>Mineral Sulfur</u>	<u>0.08</u>	<u>0.09</u>	<u>0</u>
<u>Organic Sulfur</u>	<u>0</u>	<u>0</u>	<u>0.05</u>

830 **Table A4. Average annual inputs of Carbon, Nitrogen, Phosphorus and Sulfur to ORG, CONV+WCC and CONV plots between 1993-2018. Please note that Sudangrass and Wheat carbon inputs were excluded from this table, as they were only grown for a limited amount of time.**

<u>Practices - 2018-2019</u>	<u>CONV</u>	<u>CONV+WCC</u>	<u>ORG</u>
<u>Compost Application</u>	<u>NA</u>	<u>NA</u>	<u>Apr 2018 and Oct 2019</u>
<u>Corn Harvest</u>	<u>Sept 10th</u>	<u>Sept 10th</u>	<u>Sept 10th</u>
<u>Tomato Harvest</u>	<u>Aug 2nd</u>	<u>Aug 2nd</u>	<u>Aug 2nd</u>
<u>Total Amount of Irrigation (mm/hectare)</u>	<u>245.13</u>	<u>239.43</u>	<u>538.27</u>
<u>Number of tractor passes/yr (7.5cm deep)</u>	<u>4</u>	<u>13</u>	<u>16</u>
<u>Number of tractor passes/yr (20.5cm deep)</u>	<u>4</u>	<u>4</u>	<u>4</u>

835 **Table A5. Management summary for ORG, CONV+WCC and CONV plots for the 2018-2019 year.**

## 7 Code availability

The code for the graphs and analyses in this manuscript is available at [https://github.com/danrath/2018\\_RRCARBON\\_DEPTH](https://github.com/danrath/2018_RRCARBON_DEPTH) (DOI: <https://zenodo.org/badge/latestdoi/181972884>)

## 8 Data availability

840 The data included in this manuscript is part of the Russell Ranch long term dataset and is available at  
https://github.com/danrath/2018\_RRCARBON\_DEPTH. (DOI: https://zenodo.org/badge/latestdoi/181972884)

## 9 Author contribution

845 NB, AAB, SY, KS, SP, TG and SPDR acquired funding for this project. DR, NT, NB, KS and DW designed the experiment and DR, NB, DW, LD, SP, TG, NT collected the data. DR, NB, LD, and SP performed data analysis and interpretation with assistance from DW, NT, KS, SY, TG and AAB. DR prepared the manuscript with contributions from all co-authors.

## 10 Competing interests

The authors declare that they have no conflict of interest.

## 11 Acknowledgements

850 Funding for this research was made possible by the University of California Office of the President, and the USDA National Institute of Food and Agriculture (NIFA) through the UCOP Catalyst Grant, the USDA-NIFA AFRI ELI Predoctoral Fellowship, Hatch Formula Funding CA 2122-H and multistate regional project W-2082. All opinions, findings, conclusions, and recommendations expressed in this publication are those of the authors and do not necessarily reflect the view of the National Institute of Food and Agriculture (NIFA), nor the United States Department of Agriculture (USDA). Authors would also like to express gratitude to Israel Herrera and the Russell Ranch staff for field assistance during the project, and Danielle  
855 Gelardi and Oren Hoffman for offering comments on the draft version of the manuscript.

## 12 References

- 860 ~~Angst, G., Messinger, J., Greiner, M., Häusler, W., Hertel, D., Kirfel, K., Kögel-Knabner, I., Leusehner, C., Rethemeyer, J. and Mueller, C. W.: Soil organic carbon stocks in topsoil and subsoil controlled by parent material, carbon input in the rhizosphere, and microbial derived compounds, Soil Biol. Biochem., 122, 19–30, https://doi.org/10.1016/j.soilbio.2018.03.026, 2018.~~
- ~~Aquino, A. J. A., Tunega, D., Schaumann, G. E., Haberhauer, G., Gerzabek, M. H. and Lischka, H.: The functionality of cation bridges for binding polar groups in soil aggregates, Int. J. Quantum Chem., 111(7–8), 1531–1542, https://doi.org/10.1002/qua.22693, 2011.~~
- 865 ~~van Bavel, C. H. M.: Mean-Weight Diameter of Soil Aggregates as a Statistical Index of Aggregation, Soil Sci. Soc. Am. J., 14(C), 20–23, https://doi.org/10.2136/sssaj1950.036159950014000e0005x, 1950.~~
- ~~Bernal, B., McKinley, D. C., Hungate, B. A., White, P. M., Mozdzer, T. J. and Magonigal, J. P.: Limits to soil carbon stability; Deep, ancient soil carbon decomposition stimulated by new labile organic inputs, Soil Biol. Biochem., 98, 85–94, https://doi.org/10.1016/j.soilbio.2016.04.007, 2016.~~
- 870 ~~Blake, G. R. and Hartge, K. H.: Bulk Density. In: A. Klute, editor, Methods of Soil Analysis: Part 1—Physical and Mineralogical Methods, in SSSA Book Ser. 5.1. SSSA, ASA, Madison, WI., vol. sssabookse, pp. 363–375, Soil Science Society of America, American Society of Agronomy, https://doi.org/10.2136/sssabookser5.1.2ed.c13, , 1986.~~
- ~~Bonkowski, M.: Protozoa and plant growth: the microbial loop in soil revisited, New Phytol., 162(3), 617–631,~~



<https://doi.org/10.1111/j.1469-8137.2004.01066.x>, 2004.

- 875 Bossio, D. A. and Seow, K. M.: Impacts of carbon and flooding on soil microbial communities: Phospholipid fatty acid profiles and substrate utilization patterns, *Microb. Ecol.*, 35(3), 265–278, <https://doi.org/10.1007/s002489900082>, 1998.
- Brennan, E. B. and Acosta-Martinez, V.: Cover cropping frequency is the main driver of soil microbial changes during six years of organic vegetable production, *Soil Biol. Biochem.*, 109, 188–204, <https://doi.org/10.1016/j.soilbio.2017.01.014>, 2017.
- 880 Buchmann, C. and Schaumann, G. E.: The contribution of various organic matter fractions to soil–water interactions and structural stability of an agriculturally cultivated soil, *J. Plant Nutr. Soil Sci.*, 181(4), 586–599, <https://doi.org/10.1002/jpln.201700437>, 2018.
- Chabbi, A., Kögel-Knabner, I. and Rumpel, C.: Stabilised carbon in subsoil horizons is located in spatially distinct parts of the soil profile, *Soil Biol. Biochem.*, 41(2), 256–261, <https://doi.org/10.1016/j.soilbio.2008.10.033>, 2009.
- Chantigny, M. H.: Dissolved and water-extractable organic matter in soils: A review on the influence of land use and management practices, in *Geoderma*, vol. 113, pp. 357–380, Elsevier, [https://doi.org/10.1016/S0016-7061\(02\)00370-1](https://doi.org/10.1016/S0016-7061(02)00370-1), 2003.
- 885 Chen, G. and Weil, R. R.: Penetration of cover crop roots through compacted soils, *Plant Soil*, 331(1), 31–43, <https://doi.org/10.1007/s11104-009-0223-7>, 2010.
- Chen, S., Martin, M. P., Saby, N. P. A., Walter, C., Angers, D. A. and Arrouays, D.: Fine resolution map of top- and subsoil carbon sequestration potential in France, *Sci. Total Environ.*, 630, 389–400, <https://doi.org/10.1016/j.scitotenv.2018.02.209>, 2018.
- 890 Chenu, C., Angers, D. A., Barré, P., Derrien, D., Arrouays, D. and Balesdent, J.: Increasing organic stocks in agricultural soils: Knowledge gaps and potential innovations, *Soil Tillage Res.*, 188, 41–52, <https://doi.org/10.1016/j.still.2018.04.011>, 2019.
- Coonan, E. C., Kirkby, C. A., Kirkegaard, J. A., Amidy, M. R., Strong, C. L. and Richardson, A. E.: Microorganisms and nutrient stoichiometry as mediators of soil organic matter dynamics, *Nutr. Cycl. Agroecosystems*, 117(3), 273–298, <https://doi.org/10.1007/s10705-020-10076-8>, 2020.
- 895 Cotrufo, M. F., Wallenstein, M. D., Boot, C. M., Denef, K. and Paul, E.: The Microbial Efficiency Matrix Stabilization (MEMS) framework integrates plant litter decomposition with soil organic matter stabilization: Do labile plant inputs form stable soil organic matter?, *Glob. Chang. Biol.*, 19(4), 988–995, <https://doi.org/10.1111/geb.12113>, 2013.
- Cui, Y., Zhang, Y., Duan, C., Wang, X., Zhang, X., Ju, W., Chen, H., Yue, S., Wang, Y., Li, S. and Fang, L.: Ecoenzymatic stoichiometry reveals microbial-phosphorus limitation decreases the nitrogen-cycling potential of soils in semi-arid agricultural ecosystems, *Soil Tillage Res.*, 197, 104463, <https://doi.org/10.1016/j.still.2019.104463>, 2020.
- 900 Deiss, L., Margenot, A. J., Culman, S. W. and Demyan, M. S.: Optimizing acquisition parameters in diffuse reflectance infrared Fourier transform spectroscopy of soils, *Soil Sci. Soc. Am. J.*, 84(3), 930–948, <https://doi.org/10.1002/saj2.20028>, 2020.
- Dick, R. P.: A review: long-term effects of agricultural systems on soil biochemical and microbial parameters, *Agric. Ecosyst. Environ.*, 40(1–4), 25–36, [https://doi.org/10.1016/0167-8809\(92\)90081-L](https://doi.org/10.1016/0167-8809(92)90081-L), 1992.
- 905 Doane, T. A. and Horwath, W. R.: Spectrophotometric determination of nitrate with a single reagent, *Anal. Lett.*, 36(12), 2713–2722, 2003.
- Dungait, J. A. J., Hopkins, D. W., Gregory, A. S. and Whitmore, A. P.: Soil organic matter turnover is governed by accessibility not recalcitrance, *Glob. Chang. Biol.*, 18(6), 1781–1796, <https://doi.org/10.1111/j.1365-2486.2012.02665.x>, 2012.

- 910 Edwards, P.: Sulfur cycling, retention, and mobility in soils: a review. [http://www.fs.fed.us/ne/newtown\\_square/publications/technical\\_reports/pdfs/1998/gtrne250.pdf](http://www.fs.fed.us/ne/newtown_square/publications/technical_reports/pdfs/1998/gtrne250.pdf), last access: 10 April 2020, 1998.
- Elliott, E. T.: Aggregate Structure and Carbon, Nitrogen, and Phosphorus in Native and Cultivated Soils, *Soil Sci. Soc. Am. J.*, 50(3), 627–633, <https://doi.org/10.2136/sssaj1986.03615995005000030017x>, 1986.
- 915 Fan, J., McConkey, B., Wang, H. and Janzen, H.: Root distribution by depth for temperate agricultural crops, *F. Crop. Res.*, 189, 68–74, <https://doi.org/10.1016/j.fcr.2016.02.013>, 2016.
- Fierer, N., Allen, A. S., Schimel, J. P. and Holden, P. A.: Controls on microbial CO<sub>2</sub> production: A comparison of surface and subsurface soil horizons, *Glob. Chang. Biol.*, 9(9), 1322–1332, <https://doi.org/10.1046/j.1365-2486.2003.00663.x>, 2003a.
- Fierer, N., Schimel, J. P. and Holden, P. A.: Variations in microbial community composition through two soil depth profiles, *Soil Biol. Biochem.*, 35(1), 167–176, [https://doi.org/10.1016/S0038-0717\(02\)00251-1](https://doi.org/10.1016/S0038-0717(02)00251-1), 2003b.
- 920 Franzluebbers, A. J.: Water infiltration and soil structure related to organic matter and its stratification with depth, *Soil Tillage Res.*, 66(2), 197–205, [https://doi.org/10.1016/S0167-1987\(02\)00027-2](https://doi.org/10.1016/S0167-1987(02)00027-2), 2002.
- Ghestem, M., Sidle, R. C. and Stokes, A.: The influence of plant root systems on subsurface flow: Implications for slope stability, *BioScience*, 61(11), 869–879, <https://doi.org/10.1525/bio.2011.61.11.6>, 2011.
- 925 Hangen, E., Buczko, U., Bens, O., Brunotte, J. and Hüttl, R. F.: Infiltration patterns into two soils under conventional and conservation tillage: Influence of the spatial distribution of plant root structures and soil animal activity, *Soil Tillage Res.*, 63(3–4), 181–186, [https://doi.org/10.1016/S0167-1987\(01\)00234-3](https://doi.org/10.1016/S0167-1987(01)00234-3), 2002.
- Harrison, R. B., Footen, P. W. and Strahm, B. D.: Deep soil horizons: Contribution and importance to soil carbon pools and in assessing whole ecosystem response to management and global change, *For. Sci.*, 57(1), 67–76, <https://doi.org/10.1093/forestscience/57.1.67>, 2011.
- 930 Hicks-Pries, C. E., Sulman, B. N., West, C., O'Neill, C., Poppleton, E., Porras, R. C., Castanha, C., Zhu, B., Wiedemeier, D. B. and Torn, M. S.: Root litter decomposition slows with soil depth, *Soil Biol. Biochem.*, 125(Soil Biology and Biochemistry 41 2009), 103–114, <https://doi.org/10.1016/j.soilbio.2018.07.002>, 2018.
- Jenny, H.: Factors of Soil Formation: A System of Quantitative Pedology, Dover Publications, New York. <http://soilslab.cfr.washington.edu/publications/Jenny-1941.pdf>, last access: 7 April 2015, 1941.
- 935 Jilling, A., Kane, D., Williams, A., Yannarell, A. C., Davis, A., Jordan, N. R., Koide, R. T., Mortensen, D. A., Smith, R. G., Snapp, S. S., Spokas, K. A. and Stuart-Grandy, A.: Rapid and distinct responses of particulate and mineral-associated organic nitrogen to conservation tillage and cover crops, *Geoderma*, 359, 114001, <https://doi.org/10.1016/j.geoderma.2019.114001>, 2020.
- 940 Johnston, A. E. and Poulton, P. R.: The importance of long term experiments in agriculture: their management to ensure continued crop production and soil fertility; the Rothamsted experience, *Eur. J. Soil Sci.*, 69(1), 113–125, <https://doi.org/10.1111/ejss.12521>, 2018.
- Jones, D. L. and Willett, V. B.: Experimental evaluation of methods to quantify dissolved organic nitrogen (DON) and dissolved organic carbon (DOC) in soil, *Soil Biol. Biochem.*, 38(5), 991–999, <https://doi.org/10.1016/J.SOILBIO.2005.08.012>, 2006.
- 945 Kaiser, K. and Kalbitz, K.: Cycling downwards dissolved organic matter in soils, *Soil Biol. Biochem.*, 52, 29–32, <https://doi.org/10.1016/J.SOILBIO.2012.04.002>, 2012.

- 950 Kautz, T., Amelung, W., Ewert, F., Gaiser, T., Horn, R., Jahn, R., Javaux, M., Kemna, A., Kuzyakov, Y., Munch, J. C., Pätzold, S., Peth, S., Scherer, H. W., Schlöter, M., Schneider, H., Vanderborght, J., Vetterlein, D., Walter, A., Wiesenberger, G. L. B. and Köpke, U.: Nutrient acquisition from arable subsoils in temperate climates: A review, *Soil Biol. Biochem.*, 57, 1003–1022, <https://doi.org/10.1016/j.soilbio.2012.09.014>, 2013.
- Keel, S. G., Anken, T., Büchi, L., Chervet, A., Fließbach, A., Flisch, R., Huguenin-Elie, O., Mäder, P., Mayer, J., Sinaj, S., Sturny, W., Wüst-Galley, C., Zihlmann, U. and Leifeld, J.: Loss of soil organic carbon in Swiss long-term agricultural experiments over a wide range of management practices, *Agric. Ecosyst. Environ.*, 286, <https://doi.org/10.1016/j.agee.2019.106654>, 2019.
- 955 Keiluweit, M., Bougoure, J. J., Nico, P. S., Pett Ridge, J., Weber, P. K. and Kleber, M.: Mineral protection of soil carbon counteracted by root exudates, *Nat. Clim. Chang.*, 5(6), 588–595, <https://doi.org/10.1038/nclimate2580>, 2015.
- Kirkby, C. A., Kirkegaard, J. A., Richardson, A. E., Wade, L. J., Blanchard, C. and Batten, G.: Stable soil organic matter: A comparison of C:N:P:S ratios in Australian and other world soils, *Geoderma*, 163(3–4), 197–208, <https://doi.org/10.1016/j.geoderma.2011.04.010>, 2011.
- 960 Kirkby, C. A., Richardson, A. E., Wade, L. J., Batten, G. D., Blanchard, C. and Kirkegaard, J. A.: Carbon nutrient stoichiometry to increase soil carbon sequestration, *Soil Biol. Biochem.*, 60, 77–86, <https://doi.org/10.1016/j.soilbio.2013.01.011>, 2013.
- Kravchenko, A. N. and Robertson, G. P.: Whole Profile Soil Carbon Stocks: The Danger of Assuming Too Much from Analyses of Too Little, *Soil Sci. Soc. Am. J.*, 75(1), 235–240, <https://doi.org/10.2136/sssaj2010.0076>, 2011.
- 965 Kuzyakov, Y.: Priming effects: Interactions between living and dead organic matter, *Soil Biol. Biochem.*, 42(9), 1363–1371, <https://doi.org/10.1016/j.soilbio.2010.04.003>, 2010.
- Laos, F., Satti, P., Walter, I., Mazzarino, M. J. and Moyano, S.: Nutrient availability of composted and noncomposted residues in a Patagonian Xeric Mollisol, *Biol. Fertil. Soils*, 31(6), 462–469, <https://doi.org/10.1007/s003740000192>, 2000.
- 970 Leifeld, J., Siebert, S. and Kögel-Knabner, I.: Changes in the chemical composition of soil organic matter after application of compost, *Eur. J. Soil Sci.*, 53(2), 299–309, <https://doi.org/10.1046/j.1351-0754.2002.00453.x>, 2002.
- Leinemann, T., Preusser, S., Mikutta, R., Kalbitz, K., Cerli, C., Höschen, C., Mueller, C. W., Kandeler, E. and Guggenberger, G.: Multiple exchange processes on mineral surfaces control the transport of dissolved organic matter through soil profiles, *Soil Biol. Biochem.*, 118, 79–90, <https://doi.org/10.1016/j.soilbio.2017.12.006>, 2018.
- 975 Liebmann, P., Wordell-Dietrich, P., Kalbitz, K., Mikutta, R., Kalks, F., Don, A., Woche, S. K., Dsilva, L. R. and Guggenberger, G.: Relevance of aboveground litter for soil organic matter formation—A soil profile perspective, *Biogeosciences*, 17(12), 3099–3113, <https://doi.org/10.5194/bg-17-3099-2020>, 2020.
- Maxin, C. R. and Kögel-Knabner, I.: Partitioning of polycyclic aromatic hydrocarbons (PAH) to water-soluble soil organic matter, *Eur. J. Soil Sci.*, 46(2), 193–204, <https://doi.org/10.1111/j.1365-2389.1995.tb01827.x>, 1995.
- 980 Medellín-Azuara, J., Howitt, R. E., MacEwan, D. J. and Lund, J. R.: Economic impacts of climate-related changes to California agriculture, *Clim. Change*, 109(SUPPL. 1), 387–405, <https://doi.org/10.1007/s10584-011-0314-3>, 2011.
- Mehlich, A.: Mehlich 3 soil test extractant: A modification of Mehlich 2 extractant, *Commun. Soil Sci. Plant Anal.*, 15(12), 1409–1416, <https://doi.org/10.1080/00103628409367568>, 1984.
- Min, K., Berhe, A. A., Khoi, C. M., van Asperen, H., Gillabel, J. and Six, J.: Differential effects of wetting and drying on soil CO<sub>2</sub> concentration and flux in near surface vs. deep soil layers, *Biogeochemistry*, 148(3), 255–269,

- 985 <https://doi.org/10.1007/s10533-020-00658-7>, 2020.
- Mise, K., Maruyama, R., Miyabara, Y., Kunito, T., Senoo, K. and Otsuka, S.: Time series analysis of phosphorus depleted microbial communities in carbon/nitrogen amended soils, *Appl. Soil Ecol.*, 145, 103346, <https://doi.org/10.1016/j.apsoil.2019.08.008>, 2020.
- 990 Mladenov, N., Williams, M. W., Schmidt, S. K. and Cawley, K.: Atmospheric deposition as a source of carbon and nutrients to an alpine catchment of the Colorado Rocky Mountains, *Biogeosciences*, 9(8), 3337–3355, <https://doi.org/10.5194/bg-9-3337-2012>, 2012.
- Øygarden, L., Kværner, J. and Jenssen, P. D.: Soil erosion via preferential flow to drainage systems in clay soils, *Geoderma*, 76(1–2), 65–86, [https://doi.org/10.1016/S0016-7061\(96\)00099-7](https://doi.org/10.1016/S0016-7061(96)00099-7), 1997.
- 995 Pathak, T., Maskey, M., Dahlberg, J., Kearns, F., Bali, K. and Zaccaria, D.: Climate Change Trends and Impacts on California Agriculture: A Detailed Review, *Agronomy*, 8(3), 25, <https://doi.org/10.3390/agronomy8030025>, 2018.
- Paul, E. A., Follett, R. F., Leavitt, S. W., Halvorson, A., Peterson, G. A. and Lyon, D. J.: Radiocarbon Dating for Determination of Soil Organic Matter Pool Sizes and Dynamics, *Soil Sci. Soc. Am. J.*, 61(4), 1058–1067, <https://doi.org/10.2136/sssaj1997.03615995006100040011x>, 1997.
- 1000 Paul, E. A., Collins, H. P. and Leavitt, S. W.: Dynamics of resistant soil carbon of midwestern agricultural soils measured by naturally occurring  $^{14}\text{C}$  abundance, *Geoderma*, 104(3–4), 239–256, [https://doi.org/10.1016/S0016-7061\(01\)00083-0](https://doi.org/10.1016/S0016-7061(01)00083-0), 2001.
- Petersen, S. O. and Klug, M. J.: Effects of sieving, storage, and incubation temperature on the phospholipid fatty acid profile of a soil microbial community, *Appl. Environ. Microbiol.*, 60(7), 2421–2430, <https://doi.org/10.1128/AEM.60.7.2421-2430.1994>, 1994.
- 1005 Pignatello, J. J.: The Measurement and Interpretation of Sorption and Desorption Rates for Organic Compounds in Soil Media, *Adv. Agron.*, 69(C), 1–73, [https://doi.org/10.1016/S0065-2113\(08\)60946-3](https://doi.org/10.1016/S0065-2113(08)60946-3), 1999.
- Poepplau, C. and Don, A.: Carbon sequestration in agricultural soils via cultivation of cover crops—A meta-analysis, *Agric. Ecosyst. Environ.*, 200, 33–41, 2015.
- Preusch, P. L., Adler, P. R., Sikora, L. J. and Tworkoski, T. J.: Nitrogen and Phosphorus Availability in Composted and Uncomposted Poultry Litter, *J. Environ. Qual.*, 31(6), 2051–2057, <https://doi.org/10.2134/jeq2002.2051>, 2002.
- 1010 Rapalee, G., Trumbore, S. E., Davidson, E. A., Harden, J. W. and Veldhuis, H.: Soil Carbon stocks and their rates of accumulation and loss in a boreal forest landscape, *Global Biogeochem. Cycles*, 12(4), 687–701, <https://doi.org/10.1029/98GB02336>, 1998.
- Rhine, E. D., Mulvaney, R. L., Pratt, E. J. and Sims, G. K.: Improving the Berthelot reaction for determining ammonium in soil extracts and water, *Soil Sci. Soc. Am. J.*, 62(2), 473–480, 1998.
- 1015 Richardson, A. E., Kirkby, C. A., Banerjee, S. and Kirkegaard, J. A.: The inorganic nutrient cost of building soil carbon, *Carbon Manag.*, 5(3), 265–268, <https://doi.org/10.1080/17583004.2014.923226>, 2014.
- Rumpel, C. and Kögel-Knabner, I.: Deep soil organic matter—a key but poorly understood component of terrestrial C cycle, *Plant Soil*, 338(1–2), 143–158, 2011.
- 1020 Rumpel, C., Chabbi, A. and Marschner, B.: Carbon storage and sequestration in subsoil horizons: Knowledge, gaps and potentials, in *Recarbonization of the Biosphere: Ecosystems and the Global Carbon Cycle*, pp. 445–464, Springer Netherlands, [https://doi.org/10.1007/978-94-007-4159-1\\_20](https://doi.org/10.1007/978-94-007-4159-1_20), 2012.

- Salomé, C., Nunan, N., Pouteau, V., Lerch, T. Z. and Chenu, C.: Carbon dynamics in topsoil and in subsoil may be controlled by different regulatory mechanisms, *Glob. Chang. Biol.*, 16(1), 416–426, <https://doi.org/10.1111/j.1365-2486.2009.01884.x>, 2010.
- 1025 Sanaullah, M., Chabbi, A., Leifeld, J., Bardoux, G., Billou, D. and Rumpel, C.: Decomposition and stabilization of root litter in top and subsoil horizons: What is the difference?, *Plant Soil*, 338(1), 127–141, <https://doi.org/10.1007/s11104-010-0554-4>, 2011.
- Schmidt, J. E., Peterson, C., Wang, D., Seow, K. M. and Gaudin, A. C. M.: Agroecosystem tradeoffs associated with conversion to subsurface drip irrigation in organic systems, *Agric. Water Manag.*, 202, 1–8, <https://doi.org/10.1016/j.agwat.2018.02.005>, 2018.
- 1030 Scott, H. D., Waddle, B. A., Williams, W., Frans, R. E. and Keisling, T. C.: Winter cover crops influence on cotton yield and selected soil properties1, *Commun. Soil Sci. Plant Anal.*, 25(19–20), 3087–3100, <https://doi.org/10.1080/00103629409369250>, 1994.
- 1035 Sharpley, A. and Moyer, B.: Phosphorus Forms in Manure and Compost and Their Release during Simulated Rainfall, *J. Environ. Qual.*, 29(5), 1462–1469, <https://doi.org/10.2134/jeq2000.00472425002900050012x>, 2000.
- Sokol, N. W. and Bradford, M. A.: Microbial formation of stable soil carbon is more efficient from belowground than aboveground input, *Nat. Geosci.*, 12(1), 46–53, <https://doi.org/10.1038/s41561-018-0258-6>, 2019.
- 1040 Spohn, M., Klaus, K., Wanek, W. and Richter, A.: Microbial carbon use efficiency and biomass turnover times depending on soil depth—Implications for carbon cycling, *Soil Biol. Biochem.*, 96, 74–81, <https://doi.org/10.1016/j.soilbio.2016.01.016>, 2016.
- Syswerda, S. P., Corbin, A. T., Mokma, D. L., Kravchenko, A. N. and Robertson, G. P.: Agricultural Management and Soil Carbon Storage in Surface vs. Deep Layers, *Soil Sci. Soc. Am. J.*, 75(1), 92–101, <https://doi.org/10.2136/sssaj2009.0414>, 2011.
- 1045 Tautges, N. E., Chiartas, J. L., Gaudin, A. C. M., O’Geen, A. T., Herrera, I. and Seow, K. M.: Deep soil inventories reveal that impacts of cover crops and compost on soil carbon sequestration differ in surface and subsurface soils, *Glob. Chang. Biol.*, 25(11), 3753–3766, <https://doi.org/10.1111/geb.14762>, 2019.
- Taylor, J. P., Wilson, B., Mills, M. S. and Burns, R. G.: Comparison of microbial numbers and enzymatic activities in surface soils and subsoils using various techniques, *Soil Biol. Biochem.*, 34(3), 387–401, [https://doi.org/10.1016/S0038-0717\(01\)00199-7](https://doi.org/10.1016/S0038-0717(01)00199-7), 2002.
- 1050 VandenBygaart, A. J., Bremer, E., McConkey, B. G., Ellert, B. H., Janzen, H. H., Angers, D. A., Carter, M. R., Drury, C. F., Lafond, G. P. and McKenzie, R. H.: Impact of Sampling Depth on Differences in Soil Carbon Stocks in Long Term Agroecosystem Experiments, *Soil Sci. Soc. Am. J.*, 75(1), 226–234, <https://doi.org/10.2136/sssaj2010.0099>, 2011.
- 1055 Vinten, A. J. A., Vivian, B. J., Wright, F. and Howard, R. S.: A comparative study of nitrate leaching from soils of differing textures under similar climatic and cropping conditions, *J. Hydrol.*, 159(1–4), 197–213, [https://doi.org/10.1016/0022-1694\(94\)90256-9](https://doi.org/10.1016/0022-1694(94)90256-9), 1994.
- Wang, D., Fonte, S. J., Parikh, S. J., Six, J. and Seow, K. M.: Biochar additions can enhance soil structure and the physical stabilization of C in aggregates, *Geoderma*, 303, 110–117, <https://doi.org/10.1016/j.geoderma.2017.05.027>, 2017.
- 1060 White, K. E., Brennan, E. B., Cavigelli, M. A. and Smith, R. F.: Winter cover crops increase readily decomposable soil carbon, but compost drives total soil carbon during eight years of intensive, organic vegetable production in California, edited by S. S. Snapp, *PLoS One*, 15(2), e0228677, <https://doi.org/10.1371/journal.pone.0228677>, 2020a.

White, K. E., Brennan, E. B., Cavigelli, M. A. and Smith, R. F.: Winter cover crops increase readily decomposable soil carbon, but compost drives total soil carbon during eight years of intensive, organic vegetable production in California, edited by S. S. Snapp, *PLoS One*, 15(2), e0228677, <https://doi.org/10.1371/journal.pone.0228677>, 2020b.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Golemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K. and Yutani, H.: Welcome to the Tidyverse, *J. Open Source Softw.*, 4(43), 1686, <https://doi.org/10.21105/joss.01686>, 2019.

Wolf, K. M., Torbert, E. E., Bryant, D., Burger, M., Denison, R. F., Herrera, I., Hopmans, J., Horwath, W., Kaffka, S., Kong, A. Y. Y., Norris, R. F., Six, J., Tomich, T. P. and Seow, K. M.: The century experiment: the first twenty years of UC Davis' Mediterranean agroecological experiment, *Ecology*, 99(2), 503, <https://doi.org/10.1002/ecy.2105>, 2018.

Wordell Dietrich, P., Don, A. and Helfrich, M.: Controlling factors for the stability of subsoil carbon in a Dystric Cambisol, *Geoderma*, 304, 40–48, <https://doi.org/10.1016/j.geoderma.2016.08.023>, 2017.

Wright, A. L., Provin, T. L., Hons, F. M., Zuberer, D. A. and White, R. H.: Compost impacts on dissolved organic carbon and available nitrogen and phosphorus in turfgrass soil, *Waste Manag.*, 28(6), 1057–1063, <https://doi.org/10.1016/j.wasman.2007.04.003>, 2008.

Wuest, S.: Seasonal Variation in Soil Organic Carbon, *Soil Sci. Soc. Am. J.*, 78(4), 1442–1447, <https://doi.org/10.2136/sssaj2013.10.0447>, 2014.

Xiang, S. R., Doyle, A., Holden, P. A. and Schimel, J. P.: Drying and rewetting effects on C and N mineralization and microbial activity in surface and subsurface California grassland soils, *Soil Biol. Biochem.*, 40(9), 2281–2289, <https://doi.org/10.1016/j.soilbio.2008.05.004>, 2008.

Yost, J. L. and Hartemink, A. E.: How deep is the soil studied—an analysis of four soil science journals, *Plant Soil*, 452(1–2), 5–18, <https://doi.org/10.1007/s11104-020-04550-z>, 2020.

Zmora Nahum, S., Markovitch, O., Tarchitzky, J. and Chen, Y.: Dissolved organic carbon (DOC) as a parameter of compost maturity, *Soil Biol. Biochem.*, 37(11), 2109–2116, 2005.

Angst, G., Messinger, J., Greiner, M., Häusler, W., Hertel, D., Kirfel, K., Kögel-Knabner, I., Leuschner, C., Rethemeyer, J. and Mueller, C. W.: Soil organic carbon stocks in topsoil and subsoil controlled by parent material, carbon input in the rhizosphere, and microbial-derived compounds, *Soil Biol. Biochem.*, 122, 19–30, <https://doi.org/10.1016/j.soilbio.2018.03.026>, 2018.

Aquino, A. J. A., Tunega, D., Schaumann, G. E., Haberhauer, G., Gerzabek, M. H. and Lischka, H.: The functionality of cation bridges for binding polar groups in soil aggregates, *Int. J. Quantum Chem.*, 111(7–8), 1531–1542, <https://doi.org/10.1002/qua.22693>, 2011.

Banfield, C. C., Dippold, M. A., Pausch, J., Hoang, D. T. T. and Kuzyakov, Y.: Biopore history determines the microbial community composition in subsoil hotspots, *Biol. Fertil. Soils* 2017 535, 53(5), 573–588, <https://doi.org/10.1007/S00374-017-1201-5>, 2017.

van Bavel, C. H. M.: Mean Weight-Diameter of Soil Aggregates as a Statistical Index of Aggregation, *Soil Sci. Soc. Am. J.*, 14(C), 20–23, <https://doi.org/10.2136/sssaj1950.036159950014000c0005x>, 1950.

Bernal, B., McKinley, D. C., Hungate, B. A., White, P. M., Mozdzer, T. J. and Megonigal, J. P.: Limits to soil carbon stability: Deep, ancient soil carbon decomposition stimulated by new labile organic inputs, *Soil Biol. Biochem.*, 98, 85–94, <https://doi.org/10.1016/J.SOILBIO.2016.04.007>, 2016.



- 1100 [Blake, G. R. and Hartge, K. H.: Bulk Density. In: A. Klute, editor, Methods of Soil Analysis: Part 1—Physical and Mineralogical Methods, in SSSA Book Ser. 5.1. SSSA, ASA, Madison, WI., vol. sssabookse, pp. 363–375, Soil Science Society of America, American Society of Agronomy, <https://doi.org/10.2136/sssabookser5.1.2ed.c13>, , 1986.](#)
- [Bleam, W. F.: Soil Science Applications of Nuclear Magnetic Resonance Spectroscopy, Adv. Agron., 46\(C\), 91–155, \[https://doi.org/10.1016/S0065-2113\\(08\\)60579-9\]\(https://doi.org/10.1016/S0065-2113\(08\)60579-9\), 1991](#)
- 1105 [Bonkowski, M.: Protozoa and plant growth: the microbial loop in soil revisited, New Phytol., 162\(3\), 617–631, <https://doi.org/10.1111/j.1469-8137.2004.01066.x>, 2004.](#)
- [Bossio, D. A. and Scow, K. M.: Impacts of carbon and flooding on soil microbial communities: Phospholipid fatty acid profiles and substrate utilization patterns, Microb. Ecol., 35\(3\), 265–278, <https://doi.org/10.1007/s002489900082>, 1998.](#)
- [Brady, N. C. and Weil, R. R.: The Nature and Properties of Soils, 15th ed., Pearson Education., 2017.](#)
- 1110 [Brooks, E. S., Boll, J. and McDaniel, P. A.: A hillslope-scale experiment to measure lateral saturated hydraulic conductivity, Water Resour. Res., 40\(4\), 4208, <https://doi.org/10.1029/2003WR002858>, 2004.](#)
- [Buchmann, C. and Schaumann, G. E.: The contribution of various organic matter fractions to soil–water interactions and structural stability of an agriculturally cultivated soil, J. Plant Nutr. Soil Sci., 181\(4\), 586–599, <https://doi.org/10.1002/jpln.201700437>, 2018.](#)
- 1115 [Çerçioğlu, M., Anderson, S. H., Udawatta, R. P. and Alagele, S.: Effect of cover crop management on soil hydraulic properties, Geoderma, 343, 247–253, <https://doi.org/10.1016/j.geoderma.2019.02.027>, 2019.](#)
- [Chabbi, A., Kögel-Knabner, I. and Rumpel, C.: Stabilised carbon in subsoil horizons is located in spatially distinct parts of the soil profile, Soil Biol. Biochem., 41\(2\), 256–261, <https://doi.org/10.1016/j.soilbio.2008.10.033>, 2009.](#)
- 1120 [Chantigny, M. H.: Dissolved and water-extractable organic matter in soils: A review on the influence of land use and management practices, in Geoderma, vol. 113, pp. 357–380, Elsevier, \[https://doi.org/10.1016/S0016-7061\\(02\\)00370-1\]\(https://doi.org/10.1016/S0016-7061\(02\)00370-1\), , 2003.](#)
- [Chen, G. and Weil, R. R.: Penetration of cover crop roots through compacted soils, Plant Soil, 331\(1\), 31–43, <https://doi.org/10.1007/s11104-009-0223-7>, 2010.](#)
- 1125 [Chen, S., Martin, M. P., Saby, N. P. A., Walter, C., Angers, D. A. and Arrouays, D.: Fine resolution map of top- and subsoil carbon sequestration potential in France, Sci. Total Environ., 630, 389–400, <https://doi.org/10.1016/j.scitotenv.2018.02.209>, 2018.](#)
- [Chenu, C., Angers, D. A., Barré, P., Derrien, D., Arrouays, D. and Balesdent, J.: Increasing organic stocks in agricultural soils: Knowledge gaps and potential innovations, Soil Tillage Res., 188, 41–52, <https://doi.org/10.1016/j.still.2018.04.011>, 2019.](#)
- 1130 [Cole, John S., and Oscar Roland Mathews. Subsoil moisture under semiarid conditions. No. 1488-2016-123487. 1939](#)
- [Colla, G., Mitchell, J. P., Joyce, B. A., Huyck, L. M., Wallender, W. W., Temple, S. R., Hsiao, T. C. and Poudel, D. D.: Soil physical properties and tomato yield and quality in alternative cropping systems, in Agronomy Journal, vol. 92, pp. 924–932, American Society of Agronomy, <https://doi.org/10.2134/agronj2000.925924x>, , 2000.](#)
- 1135 [Coonan, E. C., Kirkby, C. A., Kirkegaard, J. A., Amidy, M. R., Strong, C. L. and Richardson, A. E.: Microorganisms and nutrient stoichiometry as mediators of soil organic matter dynamics, Nutr. Cycl. Agroecosystems, 117\(3\), 273–298, <https://doi.org/10.1007/s10705-020-10076-8>, 2020.](#)

Cotrufo, M. F., Wallenstein, M. D., Boot, C. M., Denef, K. and Paul, E.: The Microbial Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition with soil organic matter stabilization: Do labile plant inputs form stable soil organic matter?, *Glob. Chang. Biol.*, 19(4), 988–995, <https://doi.org/10.1111/gcb.12113>, 2013.

Cui, Y., Zhang, Y., Duan, C., Wang, X., Zhang, X., Ju, W., Chen, H., Yue, S., Wang, Y., Li, S. and Fang, L.: Ecoenzymatic stoichiometry reveals microbial phosphorus limitation decreases the nitrogen cycling potential of soils in semi-arid agricultural ecosystems, *Soil Tillage Res.*, 197, 104463, <https://doi.org/10.1016/j.still.2019.104463>, 2020.

Dalal, R.C. and Henry, R.J., Simultaneous Determination of Moisture, Organic Carbon, and Total Nitrogen by Near Infrared Reflectance Spectrophotometry *Soil Sci. Soc. Am. J.*, 50: 120-123, <https://doi.org/10.2136/sssaj1986.03615995005000010023x>, 1986

Dangal, S. R. S., Sanderman, J., Wills, S., & Ramirez-Lopez, L. Accurate and Precise Prediction of Soil Properties from a Large Mid-Infrared Spectral Library. *Soil Syst.* 2019, Vol. 3, Page 11, 3(1), 11. <https://doi.org/10.3390/SOILSYSTEMS3010011>, 2019

Deiss, L., Margenot, A. J., Culman, S. W. and Demyan, M. S.: Optimizing acquisition parameters in diffuse reflectance infrared Fourier transform spectroscopy of soils, *Soil Sci. Soc. Am. J.*, 84(3), 930–948, <https://doi.org/10.1002/saj2.20028>, 2020.

Deiss, L., Sall, A., Demyan, M.S., Culman, S.W. Does crop rotation affect soil organic matter stratification in tillage systems. *Soil & Tillage Research* 209, 104932, 2021.

Demyan, M.S., Rasche, F., Schulz, E., Breulmann, M., Müller, T., Cadisch, G. Use of specific peaks obtained by diffuse reflectance Fourier transform mid-infrared spectroscopy to study the composition of organic matter in a Haplic Chernozem. *Eur. J. Soil Sci.* 63, 189–199. doi:10.1111/j.1365-2389.2011.01420.x, 2012

Dick, R. P.: A review: long-term effects of agricultural systems on soil biochemical and microbial parameters, *Agric. Ecosyst. Environ.*, 40(1–4), 25–36, [https://doi.org/10.1016/0167-8809\(92\)90081-L](https://doi.org/10.1016/0167-8809(92)90081-L), 1992.

Ding, G., Liu, X., Herbert, S., Novak, J., Amarasiriwardena, D. and Xing, B.: Effect of cover crop management on soil organic matter, *Geoderma*, 130(3–4), 229–239, <https://doi.org/10.1016/J.GEODERMA.2005.01.019>, 2006.

Doane, T. A. and Horwáth, W. R.: Spectrophotometric determination of nitrate with a single reagent, *Anal. Lett.*, 36(12), 2713–2722, 2003.

Dungait, J. A. J., Hopkins, D. W., Gregory, A. S. and Whitmore, A. P.: Soil organic matter turnover is governed by accessibility not recalcitrance, *Glob. Chang. Biol.*, 18(6), 1781–1796, <https://doi.org/10.1111/j.1365-2486.2012.02665.x>, 2012.

Edwards, P.: Sulfur cycling, retention, and mobility in soils: a review. [http://www.fs.fed.us/ne/newtown\\_square/publications/technical\\_reports/pdfs/1998/gtrne250.pdf](http://www.fs.fed.us/ne/newtown_square/publications/technical_reports/pdfs/1998/gtrne250.pdf), last access: 10 April 2020, 1998.

Elliott, E. T.: Aggregate Structure and Carbon, Nitrogen, and Phosphorus in Native and Cultivated Soils, *Soil Sci. Soc. Am. J.*, 50(3), 627–633, <https://doi.org/10.2136/sssaj1986.03615995005000030017x>, 1986.

Fan, J., McConkey, B., Wang, H. and Janzen, H.: Root distribution by depth for temperate agricultural crops, *F. Crop. Res.*, 189, 68–74, <https://doi.org/10.1016/j.fcr.2016.02.013>, 2016.



- 1175 Fanin, N., Kardol, P., Farrell, M., Nilsson, M. C., Gundale, M. J. and Wardle, D. A.: The ratio of Gram-positive to Gram-negative bacterial PLFA markers as an indicator of carbon availability in organic soils, *Soil Biol. Biochem.*, 128, 111–114, <https://doi.org/10.1016/j.soilbio.2018.10.010>, 2019.
- Fierer, N., Allen, A. S., Schimel, J. P. and Holden, P. A.: Controls on microbial CO<sub>2</sub> production: A comparison of surface and subsurface soil horizons, *Glob. Chang. Biol.*, 9(9), 1322–1332, <https://doi.org/10.1046/j.1365-2486.2003.00663.x>, 2003a.
- 1180 Fierer, N., Schimel, J. P. and Holden, P. A.: Variations in microbial community composition through two soil depth profiles, *Soil Biol. Biochem.*, 35(1), 167–176, [https://doi.org/10.1016/S0038-0717\(02\)00251-1](https://doi.org/10.1016/S0038-0717(02)00251-1), 2003b.
- Folorunso, O., Rolston, D., Prichard, P. and Louie, D.: Cover crops lower soil surface strength, may improve soil permeability, *Calif. Agric.*, 46(6), 26–27 // [calag.ucanr.edu/archive/?article=ca.v046n06p26](http://calag.ucanr.edu/archive/?article=ca.v046n06p26), last access: 6 May 2021, 1992.
- 1185 Franzluebbers, A. J.: Water infiltration and soil structure related to organic matter and its stratification with depth, *Soil Tillage Res.*, 66(2), 197–205, [https://doi.org/10.1016/S0167-1987\(02\)00027-2](https://doi.org/10.1016/S0167-1987(02)00027-2), 2002.
- Frostegård, Å., Tunlid, A. and Bååth, E.: Use and misuse of PLFA measurements in soils, *Soil Biol. Biochem.*, 43(8), 1621–1625, <https://doi.org/10.1016/j.soilbio.2010.11.021>, 2011.
- 1190 Ghestem, M., Sidle, R. C. and Stokes, A.: The influence of plant root systems on subsurface flow: Implications for slope stability, *Bioscience*, 61(11), 869–879, <https://doi.org/10.1525/bio.2011.61.11.6>, 2011.
- Gulick, S. H., Grimes, D. W., Munk, D. S. and Goldhamer, D. A.: Cover-Crop-Enhanced Water Infiltration of a Slowly Permeable Fine Sandy Loam, *Soil Sci. Soc. Am. J.*, 58(5), 1539–1546, <https://doi.org/10.2136/sssaj1994.03615995005800050038x>, 1994.
- 1195 Hangen, E., Buczko, U., Bens, O., Brunotte, J. and Hüttl, R. F.: Infiltration patterns into two soils under conventional and conservation tillage: Influence of the spatial distribution of plant root structures and soil animal activity, *Soil Tillage Res.*, 63(3–4), 181–186, [https://doi.org/10.1016/S0167-1987\(01\)00234-3](https://doi.org/10.1016/S0167-1987(01)00234-3), 2002.
- Harrison, R. B., Footen, P. W. and Strahm, B. D.: Deep soil horizons: Contribution and importance to soil carbon pools and in assessing whole-ecosystem response to management and global change, *For. Sci.*, 57(1), 67–76, <https://doi.org/10.1093/forestscience/57.1.67>, 2011.
- 1200 Haruna, S. I., Nkongolo, N. V., Anderson, S. H., Eivazi, F. and Zaibon, S.: In situ infiltration as influenced by cover crop and tillage management, *J. Soil Water Conserv.*, 73(2), 164–172, <https://doi.org/10.2489/jswc.73.2.164>, 2018.
- Hicks Pries, C. E., Sulman, B. N., West, C., O'Neill, C., Poppleton, E., Porras, R. C., Castanha, C., Zhu, B., Wiedemeier, D. B. and Torn, M. S.: Root litter decomposition slows with soil depth, *Soil Biol. Biochem.*, 125(Soil Biology and Biochemistry 41 2009), 103–114, <https://doi.org/10.1016/j.soilbio.2018.07.002>, 2018.
- Jenny, H.: Factors of Soil Formation: A System of Quantitative Pedology, Dover Publications, New York, <http://soilslab.cfr.washington.edu/publications/Jenny-1941.pdf>, last access: 7 April 2015, 1941.
- 1210 Jilling, A., Kane, D., Williams, A., Yannarell, A. C., Davis, A., Jordan, N. R., Koide, R. T., Mortensen, D. A., Smith, R. G., Snapp, S. S., Spokas, K. A. and Stuart Grandy, A.: Rapid and distinct responses of particulate and mineral-associated organic nitrogen to conservation tillage and cover crops, *Geoderma*, 359, 114001, <https://doi.org/10.1016/j.geoderma.2019.114001>, 2020.
- Johnston, A. E. and Poulton, P. R.: The importance of long-term experiments in agriculture: their management to ensure

- continued crop production and soil fertility; the Rothamsted experience, *Eur. J. Soil Sci.*, 69(1), 113–125, <https://doi.org/10.1111/ejss.12521>, 2018.
- Jones, D. L. and Willett, V. B.: Experimental evaluation of methods to quantify dissolved organic nitrogen (DON) and dissolved organic carbon (DOC) in soil, *Soil Biol. Biochem.*, 38(5), 991–999, <https://doi.org/10.1016/J.SOILBIO.2005.08.012>, 2006.
- Joyce, B. A., Wallender, W. W., Mitchell, J. P., Huyck, L. M., Temple, S. R., Brostrom, P. N. and Hsiao, T. C.: INFILTRATION AND SOIL WATER STORAGE UNDER WINTER COVER CROPPING IN CALIFORNIA'S SACRAMENTO VALLEY, *Trans. ASAE*, 45(2), 315–326, 2002.
- Kaiser, K. and Kalbitz, K.: Cycling downwards – dissolved organic matter in soils, *Soil Biol. Biochem.*, 52, 29–32, <https://doi.org/10.1016/J.SOILBIO.2012.04.002>, 2012.
- Kautz, T., Amelung, W., Ewert, F., Gaiser, T., Horn, R., Jahn, R., Javaux, M., Kemna, A., Kuzyakov, Y., Munch, J. C., Pätzold, S., Peth, S., Scherer, H. W., Schlöter, M., Schneider, H., Vanderborght, J., Vetterlein, D., Walter, A., Wiesenberg, G. L. B. and Köpke, U.: Nutrient acquisition from arable subsoils in temperate climates: A review, *Soil Biol. Biochem.*, 57, 1003–1022, <https://doi.org/10.1016/j.soilbio.2012.09.014>, 2013.
- Keel, S. G., Anken, T., Büchi, L., Chervet, A., Fliessbach, A., Flisch, R., Huguenin-Elie, O., Mäder, P., Mayer, J., Sinaj, S., Sturny, W., Wüst-Galley, C., Zihlmann, U. and Leifeld, J.: Loss of soil organic carbon in Swiss long-term agricultural experiments over a wide range of management practices, *Agric. Ecosyst. Environ.*, 286, <https://doi.org/10.1016/j.agee.2019.106654>, 2019.
- Keiluweit, M., Bougoure, J. J., Nico, P. S., Pett-Ridge, J., Weber, P. K. and Kleber, M.: Mineral protection of soil carbon counteracted by root exudates, *Nat. Clim. Chang.*, 5(6), 588–595, <https://doi.org/10.1038/nclimate2580>, 2015.
- Kirkby, C. A., Kirkegaard, J. A., Richardson, A. E., Wade, L. J., Blanchard, C. and Batten, G.: Stable soil organic matter: A comparison of C:N:P:S ratios in Australian and other world soils, *Geoderma*, 163(3–4), 197–208, <https://doi.org/10.1016/j.geoderma.2011.04.010>, 2011.
- Kirkby, C. A., Richardson, A. E., Wade, L. J., Batten, G. D., Blanchard, C. and Kirkegaard, J. A.: Carbon-nutrient stoichiometry to increase soil carbon sequestration, *Soil Biol. Biochem.*, 60, 77–86, <https://doi.org/10.1016/j.soilbio.2013.01.011>, 2013.
- Kuzyakov, Y.: Priming effects: Interactions between living and dead organic matter, *Soil Biol. Biochem.*, 42(9), 1363–1371, <https://doi.org/10.1016/j.soilbio.2010.04.003>, 2010.
- Laos, F., Satti, P., Walter, I., Mazzarino, M. J. and Moyano, S.: Nutrient availability of composted and noncomposted residues in a Patagonian Xeric Mollisol, *Biol. Fertil. Soils*, 31(6), 462–469, <https://doi.org/10.1007/s003740000192>, 2000.
- Lavallee, J. M., Soong, J. L. and Cotrufo, M. F.: Conceptualizing soil organic matter into particulate and mineral-associated forms to address global change in the 21st century, *Glob. Chang. Biol.*, 26(1), 261–273, <https://doi.org/10.1111/gcb.14859>, 2020.
- Leifeld, J., Siebert, S. and Kögel-Knabner, I.: Changes in the chemical composition of soil organic matter after application of compost, *Eur. J. Soil Sci.*, 53(2), 299–309, <https://doi.org/10.1046/j.1351-0754.2002.00453.x>, 2002.
- Leinemann, T., Preusser, S., Mikutta, R., Kalbitz, K., Cerli, C., Höschen, C., Mueller, C. W., Kandeler, E. and Guggenberger, G.: Multiple exchange processes on mineral surfaces control the transport of dissolved organic matter through soil profiles, *Soil Biol. Biochem.*, 118, 79–90, <https://doi.org/10.1016/j.soilbio.2017.12.006>, 2018.

- Li, S., Zheng, X., Liu, C., Yao, Z., Zhang, W. and Han, S.: Influences of observation method, season, soil depth, land use and management practice on soil dissolvable organic carbon concentrations: A meta-analysis, *Sci. Total Environ.*, 631–632, 105–114, <https://doi.org/10.1016/j.scitotenv.2018.02.238>, 2018.
- Liebmann, P., Wordell-Dietrich, P., Kalbitz, K., Mikutta, R., Kalks, F., Don, A., Woche, S. K., Dsilva, L. R. and Guggenberger, G.: Relevance of aboveground litter for soil organic matter formation - A soil profile perspective, *Biogeosciences*, 17(12), 3099–3113, <https://doi.org/10.5194/bg-17-3099-2020>, 2020.
- Lorenz, K. and Lal, R.: The Depth Distribution of Soil Organic Carbon in Relation to Land Use and Management and the Potential of Carbon Sequestration in Subsoil Horizons, *Adv. Agron.*, 88, 35–66, [https://doi.org/10.1016/S0065-2113\(05\)88002-2](https://doi.org/10.1016/S0065-2113(05)88002-2), 2005.
- Mailapalli, D. R., Horwath, W. R., Wallender, W. W. and Burger, M.: Infiltration, Runoff, and Export of Dissolved Organic Carbon from Furrow-Irrigated Forage Fields under Cover Crop and No-Till Management in the Arid Climate of California, *J. Irrig. Drain. Eng.*, 138(1), 35–42, [https://doi.org/10.1061/\(asce\)ir.1943-4774.0000385](https://doi.org/10.1061/(asce)ir.1943-4774.0000385), 2012.
- Matlou, M. C. and Haynes, R. J.: Soluble organic matter and microbial biomass C and N in soils under pasture and arable management and the leaching of organic C, N and nitrate in a lysimeter study, *Appl. Soil Ecol.*, 34(2–3), 160–167, <https://doi.org/10.1016/j.apsoil.2006.02.005>, 2006.
- Margenot, A.J., Calderón, F.J., Bowles, T.M., Parikh, S.J., Jackson, L.E. Soil organic matter functional group composition in relation to organic carbon, nitrogen, and phosphorus fractions in organically managed tomato fields. *Soil Sci. Soc. Am. J.* 79:772-782. 2015
- Margenot, A.J., F.J. Calderon, and S.J. Parikh. Limitations and potential of spectral subtractions in Fourier-transform infrared spectroscopy of soil samples. *Soil Sci. Soc. Am. J.* 80:10-26. 2016
- Margenot, A. J., Parikh, S. J., & Calderón, F. J. Improving infrared spectroscopy characterization of soil organic matter with spectral subtractions. *J. Vis. Exp.*, (143), e57464. <https://doi.org/10.3791/57464>, 2019
- Maxin, C. R. and Kogel-Knabner, I.: Partitioning of polycyclic aromatic hydrocarbons (PAH) to water-soluble soil organic matter, *Eur. J. Soil Sci.*, 46(2), 193–204, <https://doi.org/10.1111/j.1365-2389.1995.tb01827.x>, 1995.
- Medellín-Azuara, J., Howitt, R. E., MacEwan, D. J. and Lund, J. R.: Economic impacts of climate-related changes to California agriculture, *Clim. Change*, 109(SUPPL. 1), 387–405, <https://doi.org/10.1007/s10584-011-0314-3>, 2011.
- Mehlich, A.: Mehlich 3 soil test extractant: A modification of Mehlich 2 extractant, *Commun. Soil Sci. Plant Anal.*, 15(12), 1409–1416, <https://doi.org/10.1080/00103628409367568>, 1984.
- Min, K., Berhe, A. A., Khoi, C. M., van Asperen, H., Gillabel, J. and Six, J.: Differential effects of wetting and drying on soil CO<sub>2</sub> concentration and flux in near-surface vs. deep soil layers, *Biogeochemistry*, 148(3), 255–269, <https://doi.org/10.1007/s10533-020-00658-7>, 2020.
- Mise, K., Maruyama, R., Miyabara, Y., Kunito, T., Senoo, K. and Otsuka, S.: Time-series analysis of phosphorus-depleted microbial communities in carbon/nitrogen-amended soils, *Appl. Soil Ecol.*, 145, 103346, <https://doi.org/10.1016/j.apsoil.2019.08.008>, 2020.

- 1290 Mobley, M. L., Lajtha, K., Kramer, M. G., Bacon, A. R., Heine, P. R. and Richter, D. D.: Surficial gains and subsoil losses of soil carbon and nitrogen during secondary forest development, *Glob. Chang. Biol.*, 21(2), 986–996, <https://doi.org/10.1111/GCB.12715>, 2015.
- Moore-Kucera, J. and Dick, R. P.: PLFA profiling of microbial community structure and seasonal shifts in soils of a Douglas-fir chronosequence, *Microb. Ecol.*, 55(3), 500–511, <https://doi.org/10.1007/s00248-007-9295-1>, 2008.
- 1295 Ng, E. L., Patti, A. F., Rose, M. T., Schefe, C. R., Wilkinson, K. and Cavagnaro, T. R.: Functional stoichiometry of soil microbial communities after amendment with stabilised organic matter, *Soil Biol. Biochem.*, 76, 170–178, <https://doi.org/10.1016/j.soilbio.2014.05.016>, 2014.
- Ogilvie, C. M.; Ashiq, W.; Vasava, H. B.; Biswas, A. Quantifying Root-Soil Interactions in Cover Crop Systems: A Review. *Agriculture* 2021, 11, 218. <https://doi.org/10.3390/agriculture11030218>
- 1300 Orwin, K. H., Dickie, I. A., Holdaway, R. and Wood, J. R.: A comparison of the ability of PLFA and 16S rRNA gene metabarcoding to resolve soil community change and predict ecosystem functions, *Soil Biol. Biochem.*, 117, 27–35, <https://doi.org/10.1016/j.soilbio.2017.10.036>, 2018.
- 1305 Øygarden, L., Kværner, J. and Jenssen, P. D.: Soil erosion via preferential flow to drainage systems in clay soils, *Geoderma*, 76(1–2), 65–86, [https://doi.org/10.1016/S0016-7061\(96\)00099-7](https://doi.org/10.1016/S0016-7061(96)00099-7), 1997.
- Ozelim, L. C. de S. M. and Cavalcante, A. L. B.: Representative Elementary Volume Determination for Permeability and Porosity Using Numerical Three-Dimensional Experiments in Microtomography Data, *Int. J. Geomech.*, 18(2), 04017154, [https://doi.org/10.1061/\(asce\)gm.1943-5622.0001060](https://doi.org/10.1061/(asce)gm.1943-5622.0001060), 2018.
- 1310 Ozelim, L. C. de S. M. and Cavalcante, A. L. B.: Representative Elementary Volume Determination for Permeability and Porosity Using Numerical Three-Dimensional Experiments in Microtomography Data, *Int. J. Geomech.*, 18(2), 04017154, [https://doi.org/10.1061/\(asce\)gm.1943-5622.0001060](https://doi.org/10.1061/(asce)gm.1943-5622.0001060), 2018.
- 1315 Pagliai, M., Vignozzi, N. and Pellegrini, S.: Soil structure and the effect of management practices, *Soil Tillage Res.*, 79(2), 131–143, <https://doi.org/10.1016/J.STILL.2004.07.002>, 2004.
- Pathak, T., Maskey, M., Dahlberg, J., Kearns, F., Bali, K. and Zaccaria, D.: Climate Change Trends and Impacts on California Agriculture: A Detailed Review, *Agronomy*, 8(3), 25, <https://doi.org/10.3390/agronomy8030025>, 2018.
- 1320 Paul, E. A., Follett, R. F., Leavitt, S. W., Halvorson, A., Peterson, G. A. and Lyon, D. J.: Radiocarbon Dating for Determination of Soil Organic Matter Pool Sizes and Dynamics, *Soil Sci. Soc. Am. J.*, 61(4), 1058–1067, <https://doi.org/10.2136/sssaj1997.03615995006100040011x>, 1997.
- Paul, E. A., Collins, H. P. and Leavitt, S. W.: Dynamics of resistant soil carbon of midwestern agricultural soils measured by naturally occurring <sup>14</sup>C abundance, *Geoderma*, 104(3–4), 239–256, [https://doi.org/10.1016/S0016-7061\(01\)00083-0](https://doi.org/10.1016/S0016-7061(01)00083-0), 2001.
- 1325 Petersen, S. O. and Klug, M. J.: Effects of sieving, storage, and incubation temperature on the phospholipid fatty acid profile of a soil microbial community, *Appl. Environ. Microbiol.*, 60(7), 2421–2430, <https://doi.org/10.1128/AEM.60.7.2421-2430.1994>, 1994.
- Pignatello, J. J.: The Measurement and Interpretation of Sorption and Desorption Rates for Organic Compounds in Soil Media, *Adv. Agron.*, 69(C), 1–73, [https://doi.org/10.1016/S0065-2113\(08\)60946-3](https://doi.org/10.1016/S0065-2113(08)60946-3), 1999.

Preusch, P. L., Adler, P. R., Sikora, L. J. and Tworowski, T. J.: Nitrogen and Phosphorus Availability in Composted and Uncomposted Poultry Litter, *J. Environ. Qual.*, 31(6), 2051–2057, <https://doi.org/10.2134/jeq2002.2051>, 2002.

de Queiroz, M. G., da Silva, T. G. F., Zolnier, S., Jardim, A. M. da R. F., de Souza, C. A. A., Araújo Júnior, G. do N., de Morais, J. E. F., & de Souza, L. S. B.: Spatial and temporal dynamics of soil moisture for surfaces with a change in land use in the semi-arid region of Brazil. *Catena*, 188, 104457. <https://doi.org/10.1016/J.CATENA.2020.104457>, 2020

Rahmati, M., Weihermüller, L., Vanderborght, J., Pachepsky, Y. A., Mao, L., Sadeghi, S. H., Moosavi, N., Kheirfam, H., Montzka, C., Van Looy, K., Toth, B., Hazbavi, Z., Al Yamani, W., Albalasmeh, A. A., Alghzawi, M. Z., Angulo-Jaramillo, R., Antonino, A. C. D., Arampatzis, G., Armindo, R. A., Asadi, H., Bamutaze, Y., Battle-Aguilar, J., Béchet, B., Becker, F., Blöschl, G., Bohne, K., Braud, I., Castellano, C., Cerdà, A., Chalhoub, M., Cichota, R., Císlarová, M., Clothier, B., Coquet, Y., Cornelis, W., Corradini, C., Coutinho, A. P., De Oliveira, M. B., De Macedo, J. R., Durães, M. F., Emami, H., Eskandari, I., Farajnia, A., Flammmini, A., Fodor, N., Gharaibeh, M., Ghavimippanah, M. H., Ghezzehei, T. A., Giertz, S., Hatzigiannakis, E. G., Horn, R., Jiménez, J. J., Jacques, D., Keesstra, S. D., Kelishadi, H., Kiani-Harchegani, M., Kouselou, M., Jha, M. K., Lassabatere, L., Li, X., Liebig, M. A., Lichner, L., López, M. V., Machiwal, D., Mallants, D., Mallmann, M. S., De Oliveira Marques, J. D., Marshall, M. R., Mertens, J., Meunier, F., Mohammadi, M. H., Mohanty, B. P., Pulido-Moncada, M., Montenegro, S., Morbidelli, R., Moret-Fernández, D., Moosavi, A. A., Mosaddeghi, M. R., Mousavi, S. B., Mozaffari, H., Nabiollahi, K., Neyshabouri, M. R., Ottoni, M. V., Ottoni Filho, T. B., Pahlavan-Rad, M. R., Panagopoulos, A., Peth, S., Peyneau, P. E., Picciafuoco, T., Poesen, J., Pulido, M., Reinert, D. J., Reinsch, S., Rezaei, M., Roberts, F. P., Robinson, D., Rodrigo-Comino, J., Rotunno Filho, O. C., Saito, T., et al.: Development and analysis of the Soil Water Infiltration Global database, *Earth Syst. Sci. Data*, 10(3), 1237–1263, <https://doi.org/10.5194/essd-10-1237-2018>, 2018.

Rapalee, G., Trumbore, S. E., Davidson, E. A., Harden, J. W. and Veldhuis, H.: Soil Carbon stocks and their rates of accumulation and loss in a boreal forest landscape, *Global Biogeochem. Cycles*, 12(4), 687–701, <https://doi.org/10.1029/98GB02336>, 1998.

Rhine, E. D., Mulvaney, R. L., Pratt, E. J. and Sims, G. K.: Improving the Berthelot reaction for determining ammonium in soil extracts and water, *Soil Sci. Soc. Am. J.*, 62(2), 473–480, 1998.

Richardson, A. E., Kirkby, C. A., Banerjee, S. and Kirkegaard, J. A.: The inorganic nutrient cost of building soil carbon, *Carbon Manag.*, 5(3), 265–268, <https://doi.org/10.1080/17583004.2014.923226>, 2014.

Roth, V. N., Lange, M., Simon, C., Hertkorn, N., Bucher, S., Goodall, T., Griffiths, R. I., Mellado-Vázquez, P. G., Mommer, L., Oram, N. J., Weigelt, A., Dittmar, T. and Gleixner, G.: Persistence of dissolved organic matter explained by molecular changes during its passage through soil, *Nat. Geosci.*, 12(9), 755–761, <https://doi.org/10.1038/s41561-019-0417-4>, 2019.

Rumpel, C. and Kögel-Knabner, I.: Deep soil organic matter—a key but poorly understood component of terrestrial C cycle, *Plant Soil*, 338(1–2), 143–158, 2011.

Rumpel, C., Chabbi, A. and Marschner, B.: Carbon storage and sequestration in subsoil horizons: Knowledge, gaps and potentials, in *Recarbonization of the Biosphere: Ecosystems and the Global Carbon Cycle*, pp. 445–464, Springer Netherlands, [https://doi.org/10.1007/978-94-007-4159-1\\_20](https://doi.org/10.1007/978-94-007-4159-1_20), 2012.

Salomé, C., Nunan, N., Pouteau, V., Lerch, T. Z. and Chenu, C.: Carbon dynamics in topsoil and in subsoil may be controlled by different regulatory mechanisms, *Glob. Chang. Biol.*, 16(1), 416–426, <https://doi.org/10.1111/j.1365-2486.2009.01884.x>, 2010.

Sanderman, J. and Amundson, R.: A comparative study of dissolved organic carbon transport and stabilization in California forest and grassland soils, *Biogeochemistry*, 89(3), 309–327, <https://doi.org/10.1007/s10533-008-9221-8>, 2008.

- 1370 Samson, M. E., Chantigny, M. H., Vanasse, A., Menasseri-Aubry, S., Royer, I., & Angers, D. A. Management practices differently affect particulate and mineral-associated organic matter and their precursors in arable soils. *Soil Biol Biochem.* 148, 107867. <https://doi.org/10.1016/j.soilbio.2020.107867>, 2020
- Sanaullah, M., Chabbi, A., Leifeld, J., Bardoux, G., Billou, D. and Rumpel, C.: Decomposition and stabilization of root litter in top- and subsoil horizons: What is the difference?, *Plant Soil*, 338(1), 127–141, <https://doi.org/10.1007/s11104-010-0554-4>, 2011.
- 1375 Schmidt, J. E., Peterson, C., Wang, D., Scow, K. M. and Gaudin, A. C. M.: Agroecosystem tradeoffs associated with conversion to subsurface drip irrigation in organic systems, *Agric. Water Manag.*, 202, 1–8, <https://doi.org/10.1016/j.agwat.2018.02.005>, 2018.
- Scott, H. D., Waddle, B. A., Williams, W., Frans, R. E. and Keisling, T. C.: Winter cover crops influence on cotton yield and selected soil properties<sup>1</sup>, *Commun. Soil Sci. Plant Anal.*, 25(19–20), 3087–3100, <https://doi.org/10.1080/00103629409369250>, 1994.
- 1380 Sharpley, A. and Moyer, B.: Phosphorus Forms in Manure and Compost and Their Release during Simulated Rainfall, *J. Environ. Qual.*, 29(5), 1462–1469, <https://doi.org/10.2134/jeq2000.00472425002900050012x>, 2000.
- Silhavy, T. J., Kahne, D. and Walker, S.: The bacterial cell envelope., *Cold Spring Harb. Perspect. Biol.*, 2(5), <https://doi.org/10.1101/cshperspect.a000414>, 2010.
- 1385 Singh, G., Kaur, G., Williard, K. W. J., & Schoonover, J. E.. Cover crops and tillage effects on carbon–nitrogen pools: A lysimeter study. *Vadose Zone J.*, 20(2), e20110. <https://doi.org/10.1002/VZJ2.20110>, 2021
- Slessarev, E. W., Lin, Y., Jiménez, B. Y., Homyak, P. M., Chadwick, O. A., D'Antonio, C. M. and Schimel, J. P.: Cellular and extracellular C contributions to respiration after wetting dry soil, *Biogeochemistry*, 147(3), 307–324, <https://doi.org/10.1007/s10533-020-00645-y>, 2020.
- 1390 Smernik, R. J. and Oades, J. M.: Paramagnetic Effects on Solid State Carbon-13 Nuclear Magnetic Resonance Spectra of Soil Organic Matter, *J. Environ. Qual.*, 31(2), 414–420, <https://doi.org/10.2134/JEQ2002.4140>, 2002.
- Smith, B.C., Fundamentals of Fourier Transform Infrared Spectroscopy. Second Edition. CRC Press, Taylor and Francis Group. 2011
- Smitii, Alfred. "Seasonal subsoil temperature variations." *Journal of Agricultural Research* 44 (1932): 421.
- 1395 Soil Survey Staff. Keys to Soil Taxonomy, 12th ed. USDA-Natural Resources Conservation Service, Washington, DC. 2014
- Sokol, N. W. and Bradford, M. A.: Microbial formation of stable soil carbon is more efficient from belowground than aboveground input, *Nat. Geosci.*, 12(1), 46–53, <https://doi.org/10.1038/s41561-018-0258-6>, 2019.
- 1400 Soong, J. L., Castanha, C., Pries, C. E. H., Ofiti, N., Porras, R. C., Riley, W. J., Schmidt, M. W. I. and Torn, M. S.: Five years of whole-soil warming led to loss of subsoil carbon stocks and increased CO<sub>2</sub> efflux, *Sci. Adv.*, 7(21), eabd1343, <https://doi.org/10.1126/SCIADV.ABD1343>, 2021.
- Soong, J. L., Fuchslueger, L., Marañon-Jimenez, S., Torn, M. S., Janssens, I. A., Penuelas, J. and Richter, A.: Microbial carbon limitation: The need for integrating microorganisms into our understanding of ecosystem carbon cycling, *Glob. Chang. Biol.*, 26(4), 1953–1961, <https://doi.org/10.1111/gcb.14962>, 2020.
- Spohn, M., Klaus, K., Wanek, W. and Richter, A.: Microbial carbon use efficiency and biomass turnover times depending on



- 1405 soil depth - Implications for carbon cycling, Soil Biol. Biochem., 96, 74–81, <https://doi.org/10.1016/j.soilbio.2016.01.016>, 2016.
- Sradnick, A., Oltmanns, M., Raupp, J. and Joergensen, R. G.: Microbial residue indices down the soil profile after long-term addition of farmyard manure and mineral fertilizer to a sandy soil, Geoderma, 226–227(1), 79–84, <https://doi.org/10.1016/j.geoderma.2014.03.005>, 2014.
- 1410 Steenwerth, K. and Belina, K. M.: Cover crops enhance soil organic matter, carbon dynamics and microbiological function in a vineyard agroecosystem, Appl. Soil Ecol., 40(2), 359–369, <https://doi.org/10.1016/j.apsoil.2008.06.006>, 2008.
- Syswerda, S. P., Corbin, A. T., Mokma, D. L., Kravchenko, A. N. and Robertson, G. P.: Agricultural Management and Soil Carbon Storage in Surface vs. Deep Layers, Soil Sci. Soc. Am. J., 75(1), 92–101, <https://doi.org/10.2136/sssaj2009.0414>, 2011.
- 1415 Tautges, N. E., Chiartas, J. L., Gaudin, A. C. M., O’Geen, A. T., Herrera, I. and Scow, K. M.: Deep soil inventories reveal that impacts of cover crops and compost on soil carbon sequestration differ in surface and subsurface soils, Glob. Chang. Biol., 25(11), 3753–3766, <https://doi.org/10.1111/gcb.14762>, 2019.
- Taylor, J. P., Wilson, B., Mills, M. S. and Burns, R. G.: Comparison of microbial numbers and enzymatic activities in surface soils and subsoils using various techniques, Soil Biol. Biochem., 34(3), 387–401, [https://doi.org/10.1016/S0038-0717\(01\)00199-7](https://doi.org/10.1016/S0038-0717(01)00199-7), 2002.
- 1420 van Bavel, C. H. M.: Mean Weight-Diameter of Soil Aggregates as a Statistical Index of Aggregation, Soil Sci. Soc. Am. J., 14(C), 20–23, <https://doi.org/10.2136/sssaj1950.036159950014000c0005x>, 1950.
- VandenBygaart, A. J., Bremer, E., McConkey, B. G., Ellert, B. H., Janzen, H. H., Angers, D. A., Carter, M. R., Drury, C. F., Lafond, G. P. and McKenzie, R. H.: Impact of Sampling Depth on Differences in Soil Carbon Stocks in Long-Term Agroecosystem Experiments, Soil Sci. Soc. Am. J., 75(1), 226–234, <https://doi.org/10.2136/sssaj2010.0099>, 2011.
- 1425 Vinten, A. J. A., Vivian, B. J., Wright, F. and Howard, R. S.: A comparative study of nitrate leaching from soils of differing textures under similar climatic and cropping conditions, J. Hydrol., 159(1–4), 197–213, [https://doi.org/10.1016/0022-1694\(94\)90256-9](https://doi.org/10.1016/0022-1694(94)90256-9), 1994.
- Wang, D., Fonte, S. J., Parikh, S. J., Six, J. and Scow, K. M.: Biochar additions can enhance soil structure and the physical stabilization of C in aggregates, Geoderma, 303, 110–117, <https://doi.org/10.1016/j.geoderma.2017.05.027>, 2017.
- 1430 West, J.R., Cates, A.M., Ruark, M.D., Deiss, L., Whitman, T., Rui, Y.. Winter rye does not increase microbial necromass contributions to soil organic carbon in continuous corn silage in North Central US. Soil Biol. Biochem. 107899. doi:10.1016/j.soilbio.2020.107899, 2020
- 1435 White, K. E., Brennan, E. B., Cavigelli, M. A. and Smith, R. F.: Winter cover crops increase readily decomposable soil carbon, but compost drives total soil carbon during eight years of intensive, organic vegetable production in California, edited by S. S. Snapp, PLoS One, 15(2), e0228677, <https://doi.org/10.1371/journal.pone.0228677>, 2020a.
- Whitmore, A. P., Kirk, G. J. D. and Rawlins, B. G.: Technologies for increasing carbon storage in soil to mitigate climate change, Soil Use Manag., 31, 62–71, <https://doi.org/10.1111/sum.12115>, 2015.
- 1440 Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Golemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K. and Yutani, H.: Welcome to the Tidyverse, J. Open Source Softw., 4(43), 1686, <https://doi.org/10.21105/joss.01686>, 2019.

- 1445 Williams, E. K., Fogel, M. L., Berhe, A. A., & Plante, A. F. Distinct bioenergetic signatures in particulate versus mineral-associated soil organic matter. *Geoderma*, 330, 107–116. <https://doi.org/10.1016/j.geoderma.2018.05.024>, 2018
- Wolf, K. M., Torbert, E. E., Bryant, D., Burger, M., Denison, R. F., Herrera, I., Hopmans, J., Horwath, W., Kaffka, S., Kong, A. Y. Y., Norris, R. F., Six, J., Tomich, T. P. and Scow, K. M.: The century experiment: the first twenty years of UC Davis' Mediterranean agroecological experiment, *Ecology*, 99(2), 503, <https://doi.org/10.1002/ecy.2105>, 2018.
- 1450 Wordell-Dietrich, P., Don, A. and Helfrich, M.: Controlling factors for the stability of subsoil carbon in a Dystric Cambisol, *Geoderma*, 304, 40–48, <https://doi.org/10.1016/j.geoderma.2016.08.023>, 2017.
- Wright, A. L., Provin, T. L., Hons, F. M., Zuberer, D. A. and White, R. H.: Compost impacts on dissolved organic carbon and available nitrogen and phosphorus in turfgrass soil, *Waste Manag.*, 28(6), 1057–1063, <https://doi.org/10.1016/j.wasman.2007.04.003>, 2008.
- 1455 Xiang, S. R., Doyle, A., Holden, P. A. and Schimel, J. P.: Drying and rewetting effects on C and N mineralization and microbial activity in surface and subsurface California grassland soils, *Soil Biol. Biochem.*, 40(9), 2281–2289, <https://doi.org/10.1016/j.soilbio.2008.05.004>, 2008.
- Yost, J. L. and Hartemink, A. E.: How deep is the soil studied – an analysis of four soil science journals, *Plant Soil*, 452(1–2), 5–18, <https://doi.org/10.1007/s11104-020-04550-z>, 2020.
- 1460 Zeynoddin, M., Bonakdari, H., Ebtehaj, I., Esmailbeiki, F., Gharabaghi, B., & Zare Haghi, D. A reliable linear stochastic daily soil temperature forecast model. *Soil Till Res*, 189, 73–87. <https://doi.org/10.1016/J.STILL.2018.12.023>, 2019
- Zhang, Y., Zheng, N., Wang, J., Yao, H., Qiu, Q. and Chapman, S. J.: High turnover rate of free phospholipids in soil confirms the classic hypothesis of PLFA methodology, *Soil Biol. Biochem.*, 135, 323–330, <https://doi.org/10.1016/j.soilbio.2019.05.023>, 2019.
- 1465 Zhou, X., Chen, C., Wu, H. and Xu, Z.: Dynamics of soil extractable carbon and nitrogen under different cover crop residues, *J. Soils Sediments* 2012 126, 12(6), 844–853, <https://doi.org/10.1007/S11368-012-0515-Z>, 2012.
- Zmora-Nahum, S., Markovitch, O., Tarchitzky, J. and Chen, Y.: Dissolved organic carbon (DOC) as a parameter of compost maturity, *Soil Biol. Biochem.*, 37(11), 2109–2116, 2005.