

**Changes in soil physicochemical properties and bacterial
communities among different soil depths after long-term straw
mulching under a no-till system**

Zijun Zhou^{1,4,†}, Zengqiang Li^{2,†}, Kun Chen^{1,4}, Zhaoming Chen³, Xiangzhong Zeng^{1,4},
Song Guo^{1,4}, Hua Yu^{1,4}, Yuxian Shangguan^{1,4}, Qingrui Chen^{1,4,*}, Hongzhu Fan^{1,4},
Shihua Tu^{1,4}, Mingjiang He^{1,4}, Yusheng Qin^{1,4,*}

¹ Soil and Fertilizer Institute, Sichuan Academy of Agricultural Sciences, Chengdu, China

² College of Resources and Environment, Qingdao Agricultural University, Qingdao, China

³ Institute of Environmental Resources and Soil Fertilizer, Zhejiang Academy of Agricultural Sciences, Hangzhou, China

⁴ Monitoring and Experimental Station of Plant Nutrition and Agro-Environment for Sloping Land in South Region, Ministry of Agriculture and Rural Affairs, Chengdu, China

Correspondence to: Yusheng Qin (shengyuq@126.com), Qingrui Chen (qingruichen@163.com)




Figure S1. General distribution of the 14 most abundant bacterial phyla at the different soil depths for the CK and SM treatments.

