

1 Soil $\delta^{15}\text{N}$ is a better indicator of ecosystem nitrogen cycling than
2 plant $\delta^{15}\text{N}$: A global meta-analysis

3 Kaihua Liao^{1,2*}, Xiaoming Lai^{1,2}, Qing Zhu^{1,2,3*}

4 ¹Key Laboratory of Watershed Geographic Sciences, Nanjing Institute of Geography
5 and Limnology, Chinese Academy of Sciences, Nanjing 210008, China

6 ²University of Chinese Academy of Sciences, Beijing 100049, China

7 ³Jiangsu Collaborative Innovation Center of Regional Modern Agriculture &
8 Environmental Protection, Huaiyin Normal University, Huaian 223001, China

9

10 Submitted to: *Soil*

11

* Corresponding author. Tel.: +86 25 86882139; fax: +86 25 57714759.
E-mail addresses: khiao@niglas.ac.cn (Kaihua Liao); qzhu@niglas.ac.cn (Qing Zhu)

12 **Abstract.** The nitrogen-15 (^{15}N) natural abundance composition ($\delta^{15}\text{N}$) in soils or
13 plants is a useful tool to indicate the openness of ecosystem N cycling. This study was
14 aimed to evaluate the influence of the experimental warming on soil and plant $\delta^{15}\text{N}$.
15 We applied a global meta-analysis method to synthesize 79 and 76 paired
16 observations of soil and plant $\delta^{15}\text{N}$ from 20 published studies, respectively. Results
17 showed that the mean effect sizes of the soil and plant $\delta^{15}\text{N}$ under experimental
18 warming were -0.524 (95% CI: -0.987 to -0.162) and 0.189 (95% CI: -0.210 to 0.569),
19 respectively. This indicated that soil $\delta^{15}\text{N}$ had negative response to warming at the
20 global scale, where warming had no significant effect on plant $\delta^{15}\text{N}$. Experimental
21 warming significantly ($p < 0.05$) decreased soil $\delta^{15}\text{N}$ in Alkali and medium-textured
22 soils, in grassland/meadow, under air warming, for 4-10 yr warming period and for an
23 increase of > 3 °C in temperature, whereas it significantly ($p < 0.05$) increased soil
24 $\delta^{15}\text{N}$ in neutral and fine-textured soils and for an increase of 1.5-3 °C in temperature.
25 Plant $\delta^{15}\text{N}$ significantly ($p < 0.05$) increased with increasing temperature in neutral
26 and fine-textured soils and significantly ($p < 0.05$) decreased in alkali soil. Latitude
27 did not affect the warming effects on both soil and plant $\delta^{15}\text{N}$. However, the warming
28 effect on soil $\delta^{15}\text{N}$ was positively controlled by the mean annual temperature, which is
29 related to the fact that the higher temperature can strengthen the activity of soil
30 microbes. The effect of warming on plant $\delta^{15}\text{N}$ had weaker relationships with
31 environmental variables compared with that on soil $\delta^{15}\text{N}$. This implied that soil $\delta^{15}\text{N}$
32 was more effective than plant $\delta^{15}\text{N}$ in indicating the openness of global ecosystem N
33 cycling.

34 **1 Introduction**

35 Nitrogen (N) is one of the most important nutrient elements for plant growth and the
36 key limiting factors for vegetation productivity (McLay et al., 2001; Zhu et al., 2018;
37 Lu et al., 2020). On the one hand, if the available N in the soil is insufficient, it will
38 damage and weaken the ecosystem service function, including the supply of primary
39 material products, water conservation, climate regulation, etc. (Averill and Waring,
40 2018). On the other hand, if the available N in the soil is over supplied, it will also
41 damage the structure and function of the ecosystem, resulting in a series of
42 environmental problems such as soil acidification and imbalance of ecosystem
43 nutrient (Schrijver et al., 2008). The intermediate products of the N cycling processes,
44 such as nitrate nitrogen ($\text{NO}_3^- - \text{N}$), nitrous oxide (N_2O) and nitric oxide (NO), may
45 also cause eco-environmental pollution such as eutrophication of water body and
46 aggravation of climate-related issues (Liao et al., 2019). Therefore, it is of great
47 significance to reveal the openness of the ecosystem N cycle process for
48 understanding the plant N fixation and long-term trend of N cycling and protecting
49 the eco-environment (Wang et al., 2014; Wu et al., 2019). Openness is a measure of
50 both N inputs and outputs relative to internal cycling and determines both the
51 potential rate of N accumulation in the ecosystem and the potential for N losses
52 following a disturbance (Rastetter et al., 2021).

53 The ^{15}N natural abundance composition ($\delta^{15}\text{N}$) in soils or plants (leaves, shoots,
54 fine roots and litter) is often used to indicate the openness of ecosystem N cycling
55 (Robinson, 2001). This is because the lighter isotope of ^{14}N is always preferentially

56 lost from the ecosystem. Thus, the isotopic fractionation effect results in gradual ^{15}N
57 enrichment in the ecosystem (Aranibar et al., 2004). The larger the $\delta^{15}\text{N}$ value, the
58 higher degree of openness of N cycling. In addition, soil $\delta^{15}\text{N}$ also appears to reflect
59 the degree of decomposition of the organic matter, showing that $\delta^{15}\text{N}$ increases with
60 processing (Craine et al., 2015). A large number of studies have confirmed that
61 climate was the main factor regulating the soil and plant $\delta^{15}\text{N}$ (Craine et al., 2015;
62 Soper et al., 2015). Previous studies have demonstrated that precipitation had a
63 negative effect on soil and plant $\delta^{15}\text{N}$ from in-situ evidences to cross-sites syntheses
64 (Swap et al., 2004; Soper et al., 2015). However, the influence of temperature on soil
65 and plant $\delta^{15}\text{N}$ remains controversial. Some studies have shown that soil and plant
66 $\delta^{15}\text{N}$ increased with temperature (Amundson et al., 2003; Craine et al., 2015), while
67 others have indicated that $\delta^{15}\text{N}$ decreased with temperature (Cheng et al., 2009; Sheng
68 et al., 2014) or that they were not correlated (Yang et al., 2013). The various studies
69 suggested that the responses of soil and plant $\delta^{15}\text{N}$ to warming were very complex and
70 not well understood. In addition to climate factor, soil and plant $\delta^{15}\text{N}$ are affected by a
71 variety of other environmental factors, such as vegetation type, topography, soil
72 properties and management practices (Gurmesa et al., 2017; Wang et al., 2019).
73 However, we know little about the influences of environmental factors on the
74 warming effect on ecosystem N cycling, in terms of soil and plant $\delta^{15}\text{N}$.

75 Soil and air warming experiments have often been conducted to study the effect
76 of warming on the ecosystem N cycling at site scale (Schindlbacher et al., 2009). At
77 present, the effect of experimental warming on soil and plant $\delta^{15}\text{N}$ has not been

78 studied on a global scale. The objectives of this study were to: (i) detect the effect of
79 experimental warming on the soil and plant $\delta^{15}\text{N}$ based on a global meta-analysis of
80 20 studies; and (ii) identify the main factors influencing the warming effect on the soil
81 and plant $\delta^{15}\text{N}$. In addition, previous studies (e.g., Liu and Wang, 2009; Wang et al.,
82 2014) have found that the correlation between soil $\delta^{15}\text{N}$ and environmental factors
83 was stronger than that for plant, which may be due to the fact that soil samples
84 represented a long-term average for a given location, while plant samples were
85 affected by the microenvironment or the short-term environmental fluctuations.
86 Therefore, we specifically hypothesized that soil $\delta^{15}\text{N}$ is a better indicator of
87 ecosystem N cycling than plant $\delta^{15}\text{N}$.

88 **2 Materials and methods**

89 **2.1 Source of data and selection criteria**

90 Peer-reviewed journal articles and dissertations related to soil and plant $\delta^{15}\text{N}$ under
91 experimental warming were searched using Web of Science and China National
92 Knowledge Infrastructure (CNKI, <http://www.cnki.net>) until March 31, 2020 (Tab. 1).
93 The keywords used for the literature search were related to: “nitrogen isotope
94 composition”, “experimental warming” and “ecosystems nitrogen cycling”.

95 Our criteria were as follows: at least one of the target variables was contained,
96 including soils (different fractions, e.g., sand, silt, clay, aggregate and bulk soil) and
97 plants (leaves, shoots, roots and litters) $\delta^{15}\text{N}$; studies with climate gradients
98 (space-time substitution) were excluded and only field warming experimental studies
99 were included; only data from control and warming treatments were applied for

100 multifactor experiments; means, standard deviations (SD) (or standard errors (SE))
101 and sample sizes were directly provided or could be calculated from the studies; if one
102 article contained soil or plant $\delta^{15}\text{N}$ in multiple years, only the latest results were
103 applied since the observations should be independent in the meta-analysis (Hedges et
104 al., 1999).

105 **2.2 Data extraction and statistical analysis**

106 In total, 20 published papers were selected from 54 published papers. The locations of
107 warming experiments were presented and their site information is listed in Tab. 1. For
108 each study, the means, the statistical variation (SE or SD) and the sample size values
109 for treatment and control groups were extracted for each response variable ($\delta^{15}\text{N}$). In
110 addition to $\delta^{15}\text{N}$, the latitude, longitude, altitude, soil pH, organic matter content,
111 vegetation type, mean annual precipitation (MAP) and mean annual temperature
112 (MAT) were also extracted if they were provided (Tab. 1). All data were extracted
113 from tables or digitized from graphs with the software GetData v2.2.4
114 (<http://www.getdata-graph-digitizer.com>). A total of 79 and 76 paired observations for
115 soil and plant $\delta^{15}\text{N}$ were obtained, respectively.

116 The METAWIN 2.1 software (Sinauer Associates Inc., Sunderland, MA, USA)
117 (Rosenberg et al., 2000) was used to perform meta-analysis in this study. The Hedges'
118 *d* value was used as the effect size (Hedges et al., 1999). The absolute *d* value
119 indicated the magnitude of the treatment impact. Positive or negative *d* values
120 represented an increase or decrease effect of the treatment, respectively. Zero meant
121 no difference between treatment and control groups. Resampling tests were

122 incorporated into our meta-analysis using the bootstrap method (999 random
123 replicates). The mean effect size (calculated from 999 iterations) and 95% bootstrap
124 confidence intervals (CI) were then generated. If the 95% CI values of d did not
125 overlap zero, the effects of experimental warming on $\delta^{15}\text{N}$ were considered significant
126 at $p < 0.05$. We used a random effects model to test whether warming had a significant
127 effect on $\delta^{15}\text{N}$. To examine whether experimental conditions alter the response
128 direction and magnitude of soil and plant $\delta^{15}\text{N}$, observations were further divided into
129 subgroups according to the soil acidity-alkalinity (acid ($\text{pH} < 6.5$), neutral ($6.5 < \text{pH} <$
130 7.5), and alkali ($\text{pH} > 7.5$)), vegetation types (forest/shrub, moss/lichen, and
131 grassland/meadow), warming treatments (soil warming, air warming, and both soil
132 and air warming), soil texture (fine-, medium-, and coarse-textured soil), length of
133 warming (< 4 yr, 4-10 yr, and > 10 yr), and increase in temperature (< 1.5 °C, 1.5-3 °C,
134 and > 3 °C). A random effects model with a grouping variable was used to compare
135 responses among different subgroups. Linear regression analyses were applied to
136 assess the relationships between the Hedges' d values and environmental factors (i.e.,
137 latitude, altitude, MAT and MAP).

138 **3 Results**

139 Across all sites, the mean effect sizes of the soil and plant $\delta^{15}\text{N}$ under experimental
140 warming were -0.524 (95% CI: -0.987 to -0.162) and 0.189 (95% CI: -0.210 to 0.569),
141 respectively (Fig. 1). Experimental warming significantly ($p < 0.05$) decreased soil
142 $\delta^{15}\text{N}$ in Alkali (mean effect size = -2.484; 95% CI: -2.931 to -2.060) and
143 medium-textured (mean effect size = -0.676; 95% CI: -1.153 to -0.249) soils, in

144 grassland/meadow (mean effect size = -0.609; 95% CI: -1.076 to -0.190), under air
145 warming (mean effect size = -0.652; 95% CI: -1.081 to -0.273), for 4-10 yr warming
146 period (mean effect size = -0.652; 95% CI: -1.081 to -0.273) and for an increase of >
147 3 °C in temperature (mean effect size = -0.652; 95% CI: -1.081 to -0.273). However, it
148 significantly ($p < 0.05$) increased soil $\delta^{15}\text{N}$ in neutral (mean effect size = 0.359; 95%
149 CI: 0.078 to 0.620) and fine-texture soils (mean effect size = 2.394; 95% CI: 1.770 to
150 3.735), and for an increase of 1.5-3 °C in temperature (mean effect size = 0.409; 95%
151 CI: 0.070 to 0.707) (Fig. 2). Experimental warming did not significantly ($p > 0.05$)
152 change soil $\delta^{15}\text{N}$ under other experimental conditions.

153 In addition, experimental warming significantly ($p < 0.05$) increased plant $\delta^{15}\text{N}$ in
154 neutral (mean effect size = 3.157; 95% CI: 1.529 to 6.967) and fine-textured soils
155 (mean effect size = 1.202; 95% CI: 1.042 to 1.360), whereas it significantly ($p < 0.05$)
156 decreased plant $\delta^{15}\text{N}$ in alkali soil (mean effect size = -1.930; 95% CI: -2.325 to
157 -1.573) (Fig. 2). Experimental warming did not significantly ($p > 0.05$) change plant
158 $\delta^{15}\text{N}$ under other experimental conditions.

159 For soil and plant $\delta^{15}\text{N}$, their responses to experimental warming did not correlate
160 well with latitude ($p = 0.268$ and $p = 0.160$, respectively) (Fig. 3ab). However, the
161 Hedges' d values of soil $\delta^{15}\text{N}$ decreased significantly with altitude ($p < 0.001$) (Fig. 3c)
162 and increased significantly with MAT ($p < 0.001$) and MAP ($p < 0.001$) (Fig. 3eg). In
163 addition, the Hedges' d values of plant $\delta^{15}\text{N}$ were also found to increase significantly
164 with MAP ($p < 0.001$) (Fig. 3h). However, the responses of plant $\delta^{15}\text{N}$ to experimental
165 warming did not correlate well with altitude ($p = 0.109$) and MAT ($p = 0.002$) (Fig.

166 3df).

167 **4 Discussion**

168 A significant decreasing trend in soil $\delta^{15}\text{N}$ and no significant trend in plant $\delta^{15}\text{N}$ were
169 found in this study. This is somewhat inconsistent with previous findings. Chang et al.
170 (2017) observed that soil and plant $\delta^{15}\text{N}$ values decreased under warming in the
171 Tibetan permafrost. However, Zhang et al. (2019) found that the warming treatment
172 significantly increased soil and plant $\delta^{15}\text{N}$ in a subtropical forest. The various studies
173 suggest that soil and plant $\delta^{15}\text{N}$ are controlled by interactive effects of N fixation and
174 mineralization. At the global scale, $\delta^{15}\text{N}$ of N input (~ 0) is generally lower than that
175 of soil, so greater N fixation or higher N input (deposition and fertilization) under
176 warming can result in a lower soil $\delta^{15}\text{N}$ (Sorensen and Michelsen, 2011; Rousk and
177 Michelsen, 2017; Wang et al., 2018).

178 Soil pH has an important influence on nitrification, denitrification and N_2O
179 emissions from soils (Kyveryga et al., 2004). The results in this study showed that
180 when the soil was alkaline, the mean effect sizes of soil and plant $\delta^{15}\text{N}$ under warming
181 were negative, while when the soil was neutral, they were positive (Fig. 2ab).
182 Compared with alkaline condition, the near neutral conditions are more suitable for
183 the biological activities of heterotrophic denitrifying bacteria (Simek and Cooper,
184 2002). Therefore, the denitrification activity is usually higher under neutral conditions,
185 resulting in an enrichment of soil and plant N pools with ^{15}N (Kyveryga et al., 2004).
186 Vegetation type had limited effects on $\delta^{15}\text{N}$ under warming, except for soil $\delta^{15}\text{N}$ in
187 grassland/meadow (Fig. 2cd). This may be related to the differences in altitude, MAP

188 and MAT among three vegetation types (Tab. 1). The type of warming treatment was
189 found to have a substantial effect on soil $\delta^{15}\text{N}$, showing that the mean effect size of
190 soil $\delta^{15}\text{N}$ under air warming was negative and less than that under soil warming (Fig.
191 2ef). Salmon et al. (2016) have found that soil warming can increase N availability by
192 stimulating mineralization of organic matter in the warmed active layer. In addition,
193 air warming directly impacts aboveground temperatures and has an indirect effect on
194 soil $\delta^{15}\text{N}$ (Pardo et al., 2006). From Fig. 2gh, the finer the soil texture, the more
195 significant the positive effect of warming on soil and plant $\delta^{15}\text{N}$. The possible reason
196 is that the finer the soil texture, the stronger the adsorption of various ions on the soil
197 and the smaller the leaching loss of the soil, resulting in greater residual amount of
198 ^{15}N in the soil (Webster et al., 1986). In addition, the longer warming period and the
199 greater increase in temperature resulted in the more negative effect of warming on soil
200 $\delta^{15}\text{N}$ (Fig. 2ik). Chang et al. (2017) deduced that N fixation was greater under
201 warming and consequently resulted in a lower soil $\delta^{15}\text{N}$.

202 In the study of Mayor et al. (2015), soil and plant $\delta^{15}\text{N}$ were significantly ($p <$
203 0.001) and negatively correlated with latitude at the global scale. However, the
204 Hedges' d values of soil and plant $\delta^{15}\text{N}$ had weak correlations with latitude in this
205 study (Fig. 3). The warming effect on soil $\delta^{15}\text{N}$ was significantly ($p < 0.001$)
206 influenced by altitude, MAT and MAP. Among these, the strongest correlation was
207 observed for MAT. It is possible that soil $\delta^{15}\text{N}$ increased with increasing MAT when
208 the MAT exceeded a certain threshold (e.g., 9.8 °C as proposed by Craine et al.
209 (2015)). In this case, the increase in MAT can enhance the positive effect of

210 experimental warming on soil $\delta^{15}\text{N}$. In addition, the MAT can also affect ecosystem N
211 cycle by influencing soil texture. Craine et al. (2015) reported that hot sites had
212 greater clay concentrations than cold sites. As depicted in Fig. 2g, the finer the texture
213 of the soil, the more significant the effect of experimental warming on soil $\delta^{15}\text{N}$. High
214 d values of soil $\delta^{15}\text{N}$ corresponded to MAT of about 20 °C, which was the most
215 suitable temperature for nitrification and denitrification. However, warming had a
216 substantial negative impact on soil $\delta^{15}\text{N}$ when MAT decreased to around -5 °C.
217 Recently, Rousk et al. (2018) also found that the increase of temperature in the Arctic
218 promoted the biological N fixation, which can decrease the soil $\delta^{15}\text{N}$. The decrease of
219 d values of soil $\delta^{15}\text{N}$ with increasing altitude and decreasing MAP in this study might
220 be caused by the positive response of d values to MAT.

221 The relationships between the d values and environmental variables for plant
222 $\delta^{15}\text{N}$ were weaker than those for soil $\delta^{15}\text{N}$ (Fig. 3). The possible reason is that several
223 other factors (e.g., plant N concentrations and species richness) might co-regulate
224 plant $\delta^{15}\text{N}$ (Wu et al., 2019). This is consistent with the study of Craine et al. (2009),
225 who found different inflection points in soil and plant $\delta^{15}\text{N}$ relationships with MAT. In
226 addition, plants are generally depleted in ^{15}N relative to soils. Above results implied
227 that soil $\delta^{15}\text{N}$ was more efficient in indicating the openness of ecosystem N cycling
228 than plant $\delta^{15}\text{N}$ at the global scale. Although the present study provided a global
229 meta-analysis of the responses of $\delta^{15}\text{N}$ to experimental warming, the magnitude of
230 these responses might be uncertain. For example, a small number of observations
231 were obtained in moss/lichen under soil warming and both soil and air warming

232 treatments, which would affect the results of meta-analysis. Future research should
233 take more experimental data into account in order to better investigate the warming
234 effects on $\delta^{15}\text{N}$.

235 **6 Conclusions**

236 Our global meta-analysis indicated a significant decreasing trend in soil $\delta^{15}\text{N}$ and no
237 significant trend in plant $\delta^{15}\text{N}$ under experimental warming. Latitude did not affect the
238 warming effects on $\delta^{15}\text{N}$. However, the warming effect on $\delta^{15}\text{N}$ was related to soil
239 acidity-alkalinity, texture, vegetation type, warming treatment and period, increase in
240 temperature, altitude, MAT and MAP. The effect of warming on soil $\delta^{15}\text{N}$ was better
241 correlated with environmental variables compared with that on plant $\delta^{15}\text{N}$. Our
242 findings should be useful for understanding the underlying mechanisms of the
243 response of ecosystem N cycling to global warming.

244 **Data availability.** The data that support the findings of this study are available from
245 the corresponding author upon request.

246 **Author contributions.** KL and QZ designed this study, KL and XL performed the
247 meta-analysis, KL and QZ obtained funding, and KL and XL wrote the paper with
248 contributions from QZ.

249 **Competing interests.** The authors declare that they have no conflict of interest.

250 **Acknowledgements.** We thank two anonymous reviewers and editor for their efforts
251 on this paper. Support for this research was provided by the National Natural Science
252 Foundation of China and by Chinese Academy of Sciences.

253 **Financial support.** This study was financially supported by the National Natural

254 Science Foundation of China (41771107 and 42171077), the Key Research Program
255 of Frontier Sciences, Chinese Academy of Sciences (QYZDB-SSW-DQC038), and
256 the Youth Innovation Promotion Association, Chinese Academy of Sciences
257 (2020317).

258 **Review statement.** This paper was reviewed by two anonymous referees.

259 **References**

- 260 Aerts, R., Callaghan, T. V., Dorrepaal, E., van Logtestijn, R. S. P., and Cornelissen, J.
261 H. C.: Blackwell Publishing Ltd Seasonal climate manipulations result in
262 species-specific changes in leaf nutrient levels and isotopic composition in a
263 sub-arctic bog. *Funct. Ecol.*, 23, 680–688,
264 <https://doi.org/10.1111/j.1365-2435.2009.01566.x>, 2009.
- 265 Amundson, R., Austin, A. T., Schuur, E. A. G., Yoo, K., Matzek, V., Kendall, C.,
266 Uebersax, A., Brenner, D., and Baisden, W.T.: Global patterns of the isotopic
267 composition of soil and plant nitrogen. *Global Biogeochem. Cy.*, 17, 1031,
268 <https://doi.org/10.1029/2002GB001903>, 2003.
- 269 Anadon-Rosell, A., Ninot, J. M., Palacio, S., Grau, O., Nogués, S., Navarro, E.,
270 Carmen Sancho, M., and Carrillo, E.: Four years of experimental warming do not
271 modify the interaction between subalpine shrub species. *Oecologia*, 183, 1167–
272 1181, <https://doi.org/10.1007/S00442-017-3830-7>, 2017.
- 273 Aranibar, J. N., Otter, L., Macko, S. A., Feral, C. J. W., Epstein, H. E., Dowty, P. R.,
274 Eckardt, F., Shugart, H. H., and Swap, R. J.: Nitrogen cycling in the soil-plant
275 system along a precipitation gradient in the Kalahari sands. *Global Change Biol.*,

276 10, 359–373, <https://doi.org/10.1111/j.1365-2486.2003.00698.x>, 2004.

277 Averill, C., and Waring, B.: Nitrogen limitation of decomposition and decay: How
278 can it occur? *Global Change Biol.*, 4, 1417–1427,
279 <https://doi.org/10.1111/gcb.13980>, 2018.

280 Bijoor, N. S., Czimczik, C. I., Pataki, D. E., and Billings, S. A.: Effects of temperature
281 and fertilization on nitrogen cycling and community composition of an urban
282 lawn. *Global Change Biol.*, 14, 2119–2131,
283 <https://doi.org/10.1111/j.1365-2486.2008.01617.x>, 2008.

284 Chang, R., Wang, G., Yang, Y., and Chen, X.: Experimental warming increased soil
285 nitrogen sink in the Tibetan permafrost. *J. Geophys. Res. Biogeosci.*, 122, 1870–
286 1879, <https://doi.org/10.1002/2017JG003827>, 2017.

287 Cheng, W. X., Chen, Q. S., Xu, Y. Q., Han, X. G., and Li, L. H.: Climate and
288 ecosystem ¹⁵N natural abundance along a transect of Inner Mongolian grasslands:
289 Contrasting regional patterns and global patterns. *Global Biogeochem. Cy.*, 23,
290 GB2005, <https://doi.org/10.1029/2008GB003315>, 2009.

291 Cheng, X., Luo, Y., Xu, X., Sherry, R., and Zhang, Q.: Soil organic matter dynamics
292 in a North America tallgrass prairie after 9 yr of experimental warming.
293 *Biogeosciences*, 8, 1487–1498, <https://doi.org/10.5194/bg-8-1487-2011>, 2011.

294 Craine, J. M. et al.: Convergence of soil nitrogen isotopes across global climate
295 gradients. *Sci. Rep.*, 5, 8280, <https://doi.org/10.1038/srep08280>, 2015.

296 Dawes, M. A., Schleppei, P., Hättenschwiler, S., Rixen, C., and Hagedorn, F.: Soil
297 warming opens the nitrogen cycle at the alpine treeline. *Global Change Biol.*, 23,

298 421–434, <https://doi.org/10.1111/gcb.13365>, 2017.

299 Deane-Coe, K. K., Mauritz, M., Celis, G., Salmon, V., Crummer, K. G., Natali, S. M.,
300 and Schuur, E. A. G.: Experimental warming alters productivity and isotopic
301 signatures of tundra mosses. *Ecosystems*, 18, 1070–1082,
302 <https://doi.org/10.1007/s00442-015-3427-y>, 2015.

303 Gonzalez-Meler, M. A., Silva, L. B. C., Dias-De-Oliveira, E., Flower, C. E., and
304 Martinez, C. A.: Experimental air warming of a *Stylosanthes capitata*, vogel
305 dominated tropical pasture affects soil respiration and nitrogen dynamics. *Front.*
306 *Plant sci.*, 8, 46, <https://doi.org/10.3389/fpls.2017.00046>, 2017.

307 Gurmesa, G. A., Lu, X. K., Gundersen, P., Fang, Y. T., Mao, Q. G., Hao, C., and Mo,
308 J. M.: Nitrogen input N-15 signatures are reflected in plant N-15 natural
309 abundances in subtropical forests in China. *Biogeosciences*, 14, 2359–2370,
310 <https://doi.org/10.5194/bg-14-2359-2017>, 2017.

311 Hedges, L. V., Gurevitch, J., and Curtis, P. S.: The meta-analysis of response ratios in
312 experimental ecology. *Ecology*, 80, 1150–1156,
313 [https://doi.org/10.1890/0012-9658\(1999\)080\[1150:TMAORR\]2.0.CO;2](https://doi.org/10.1890/0012-9658(1999)080[1150:TMAORR]2.0.CO;2), 1999.

314 Hudson, J. M. G., Henry, G. H. R., and Cornwell, W. K.: Taller and larger: shifts in
315 Arctic tundra leaf traits after 16 years of experimental warming. *Global Change*
316 *Biol.*, 17, 1013–1021, <https://doi.org/10.1111/j.1365-2486.2010.02294.x>, 2011.

317 Kyveryga, P. M., Blackmer, A. M., Ellsworth, J. W., and Isla, R.: Soil pH effects on
318 nitrification of fall-applied anhydrous ammonia. *Soil Sci. Soc. Am. J.*, 68, 545–
319 551, <https://doi.org/10.2136/sssaj2004.0545>, 2004.

320 Liao, K., Lai, X., Zhou, Z., Zeng, X., Xie, W., Castellano, M. J., and Zhu, Q.:
321 Whether the rock fragment content should be considered when investigating
322 nitrogen cycle in stony soils? *J. Geophys. Res. Biogeosci.*, 124, 521–536,
323 <https://doi.org/10.1029/2018JG004780>, 2019.

324 Lim, H., Oren, R., Näsholm, T., Strömngren, M., Lundmark, T., Grip, H., and Linder,
325 S.: Boreal forest biomass accumulation is not increased by two decades of soil
326 warming. *Nat. Clim. Change*, 9, 49–52,
327 <https://doi.org/10.1038/s41558-018-0373-9>, 2019.

328 Liu, W., and Wang, Z.: Nitrogen isotopic composition of plant-soil in the Loess
329 Plateau and its responding to environmental change. *Chin. Sci. Bull.*, 54, 272–
330 279, <https://doi.org/10.1007/s11434-008-0442-y>, 2009.

331 Lu, Y., Gao, Y., and Yang, T.: A review of mass flux monitoring and estimation
332 methods for biogeochemical interface processes in watersheds. *J. Geogr. Sci.*, 30,
333 881–907, <https://doi.org/10.1007/s11442-020-1760-5>, 2020.

334 Lv, C., Zhang, Q., Hao, Y., Chen, Y., and Yang, Y.: Influence of short-term warming
335 on the composition of stable carbon and nitrogen isotopes in *Cunninghamia*
336 *lanceolata* in subtropical region of China. *Forest Res.*, 31, 27–32,
337 <https://doi.org/10.13275/j.cnki.lykxyj.2018.05.004>, 2018. (in Chinese)

338 Mayor, J., Bahram, M., Henkel, T., Buegger, F., Pritsch, K., and Tedersoo, L.:
339 Ectomycorrhizal impacts on plant nitrogen nutrition: emerging isotopic patterns,
340 latitudinal variation and hidden mechanisms. *Ecol. Lett.*, 18, 96–107,
341 <https://doi.org/10.1111/ele.12377>, 2015.

342 McLay, C., Dragten, R., Sparling, G., and Selvarajah, N.: Predicting groundwater
343 nitrate concentrations in a region of mixed agricultural land use, a comparison of
344 three approaches. *Environ. Pollut.*, 115, 191–204,
345 [https://doi.org/10.1016/S0269-7491\(01\)00111-7](https://doi.org/10.1016/S0269-7491(01)00111-7), 2001.

346 Munir, T. M., Khadka, B., Xu, B., and Strack, M.: Mineral nitrogen and phosphorus
347 pools affected by water table lowering and warming in a boreal forested peatland.
348 *Ecology*, 10, e1893, <https://doi.org/10.1002/eco.1893>, 2017.

349 Natali, S. M., Schuur, E. A. G., and Rubin, R. L.: Increased plant productivity in
350 Alaskan tundra as a result of experimental warming of soil and permafrost. *J.*
351 *Ecol.*, 100, 488–498, <https://doi.org/10.1111/j.1365-2745.2011.01925.x>, 2012.

352 Pardo, L. H., et al.: Regional assessment of N saturation using foliar and root $\delta^{15}\text{N}$.
353 *Biogeochemistry*, 80, 143–171, <https://doi.org/10.1007/s10533-006-9015-9>,
354 2006.

355 Peng, A.: Effects of artificial warming on alpine meadow in the permafrost region of
356 Qinghai-Tibet Plateau. Master dissertation, The University of Chinese Academy
357 of Sciences, 2017. (in Chinese)

358 Rastetter, E. B., Kling, G. W., Shaver, G. R., Crump, B. C., Gough, L., and Griffin, K.
359 L.: Ecosystem recovery from disturbance is constrained by N cycle openness,
360 vegetation-soil N distribution, form of N losses, and the balance between
361 vegetation and soil-microbial processes. *Ecosystems*, 24, 667–685,
362 <https://doi.org/10.1007/s10021-020-00542-3>, 2021.

363 Robinson, D.: $\delta^{15}\text{N}$ as an integrator of the nitrogen cycle. *Trends Ecol. Evol. (Amst.)*,

364 16, 153–162, [https://doi.org/10.1016/S0169-5347\(00\)02098-X](https://doi.org/10.1016/S0169-5347(00)02098-X), 2001.

365 Rosenberg, M., Adams, D., and Gurevitch, J.: MetaWin: Statistical Software for
366 Meta-Analysis. Sinauer Associates, Sunderland, MA, USA, 2000.

367 Rousk, K., and Michelsen, A.: Ecosystem nitrogen fixation throughout the snow-free
368 period in subarctic tundra: Effects of willow and birch litter addition and
369 warming. *Global Change Biol.*, 23, 1552–1563,
370 <https://doi.org/10.1111/gcb.13418>, 2017.

371 Rousk, K., Sorensen, P. L., and Michelsen, A.: What drives biological nitrogen
372 fixation in high arctic tundra: Moisture or temperature? *Ecosphere*, 9, e02117,
373 <https://doi.org/10.1002/ecs2.2117>, 2018.

374 Rui, Y., Wang, S., Xu, Z., Wang, Y., Chen, C., Zhou, X., Kang, X., Lu, S., Hu, Y.,
375 Lin, Q., and Luo, C.: Warming and grazing affect soil labile carbon and nitrogen
376 pools differently in an alpine meadow of the Qinghai–Tibet Plateau in China. *J.*
377 *Soils Sediments*, 11, 903–914, <https://doi.org/10.1007/s11368-011-0388-6>, 2011.

378 Salmon, V. G., Soucy, P., Mauritz, M., Celis, G., Natali, S. M., Mack, M. C., and
379 Schuur, E. A. G.: Nitrogen availability increases in a tundra ecosystem during
380 five years of experimental permafrost thaw. *Global Change Biol.*, 22, 1927–1941,
381 <https://doi.org/10.1111/gcb.13204>, 2016.

382 Schaeffer, S. M., Sharp, E., Schimel, J. A. P., and Welker, J. M.: Soil–plant N
383 processes in a High Arctic ecosystem, NW Greenland are altered by long-term
384 experimental warming and higher rainfall. *Global Change Biol.*, 19, 3529–3539,
385 <https://doi.org/10.1111/gcb.12318>, 2013.

386 Schindlbacher, A., Zechmeister-Boltenstern, S., and Jandl, R.: Carbon losses due to
387 soil warming: do autotrophic and heterotrophic soil respiration respond equally?
388 *Global Change Biol.*, 15, 901–913,
389 <https://doi.org/10.1111/j.1365-2486.2008.01757.x>, 2009.

390 Schneckner, J., Borken, W., Schindlbacher, A., and Wanek, W.: Little effects on soil
391 organic matter chemistry of density fractions after seven years of forest soil
392 warming. *Soil Biol. Biochem.*, 103, 300–307,
393 <https://doi.org/10.1016/j.soilbio.2016.09.003>, 2016.

394 Schrijver, A. D., Verheyen, K., Mertens, J., Staelens, J., Wuyts, K., and Muys, B.:
395 Nitrogen saturation and net ecosystem production. *Nature*, 451, E1,
396 <https://doi.org/10.1038/nature06578>, 2008.

397 Sheng, W., Yu, G., Fang, H., Liu, Y., Wang, Q., Chen, Z., and Zhang, L.: Regional
398 patterns of ^{15}N natural abundance in forest ecosystems along a large transect in
399 eastern China. *Sci. Rep.*, 4, 4249, <https://doi.org/10.1038/srep04249>, 2014.

400 Simek, M., and Cooper, J. E.: The influence of soil pH on denitrification: progress
401 towards the understanding of this interaction over the last 50 years. *Eur. J. Soil
402 Sci.*, 53, 345–354, <https://doi.org/10.1046/j.1365-2389.2002.00461.x>, 2002.

403 Soper, F. M., Richards, A. E., Siddique, I., Aidar, M. P. M., Cook, G. D., Hutley, L.
404 B., Robinson, N., and Schmidt, S.: Natural abundance ($\delta^{15}\text{N}$) indicates shifts in
405 nitrogen relations of woody taxa along a savanna-woodland continental rainfall
406 gradient. *Oecologia*, 178, 297–308, <https://doi.org/10.1007/s00442-014-3176-3>,
407 2015.

408 Sorensen, P. L., and Michelsen, A.: Long-term warming and litter addition affects
409 nitrogen fixation in a subarctic heath. *Global Change Biol.*, 17, 528–537,
410 <https://doi.org/10.1111/j.1365-2486.2010.02234.x>, 2011.

411 Swap, R. J., Aranibar, J. N., Dowty, P. R., Gilhooly, W. P., and Macko, S. A.: Natural
412 abundance of ^{13}C and ^{15}N in C_3 and C_4 vegetation of southern Africa: patterns
413 and implications. *Global Change Biol.*, 10, 350–358,
414 <https://doi.org/10.1111/j.1365-2486.2003.00702.x>, 2004.

415 Wang, C., Wang, X., Liu, D., Wu, H., Lv, X., Fang, Y., Cheng, W., Luo, W., Jiang, P.,
416 Shi, J., Yin, H., Zhou, J., Han, X., and Bai, E.: Aridity threshold in controlling
417 ecosystem nitrogen cycling in arid and semi-arid grasslands. *Nat. Commun.*, 5,
418 4799, <https://doi.org/10.1038/ncomms5799>, 2014.

419 Wang, Q., Nian, J., Xie, X., Yu, H., Zhang, J., Bai, J., Dong, G., Hu, J., Bai, B., Chen,
420 L., Xie, Q., Feng, J., Yang, X., Peng, J., Chen, F., Qian, Q., Li, J., and Zuo, J.:
421 Genetic variations in *ARE1* mediate grain yield by modulating nitrogen
422 utilization in rice. *Nat. Commun.*, 9, 735,
423 <https://doi.org/10.1038/s41467-017-02781-w>, 2018.

424 Wang, X., Jiang, Y., Ren, H., Yu, F., and Li, M.: Leaf and soil $\delta^{15}\text{N}$ patterns along
425 elevational gradients at both treelines and shrublines in three different climate
426 zones. *Forests*, 10, 557, <https://doi.org/10.3390/f10070557>, 2019.

427 Webster, C.P., Belford, R.K., and Cannell, R.Q.: Crop uptake and leaching losses of
428 ^{15}N labelled fertilizer nitrogen in relation to waterlogging of clay and sandy
429 loam soils. *Plant Soil*, 92, 89–101, 1986.

430 Wu, J., Song, M., Ma, W., Zhang, X., Shen, Z., Tarolli, P., Wurst, S., Shi, P.,
431 Ratzmann, G., Feng, Y., Li, M., Wang, X., and Tietjen, B.: Plant and soil's $\delta^{15}\text{N}$
432 are regulated by climate, soil nutrients, and species diversity in alpine grasslands
433 on the northern Tibetan Plateau. *Agr. Ecosyst. Environ.*, 281, 111–123,
434 <https://doi.org/10.1016/j.agee.2019.05.011>, 2019.

435 Yang, Y. H., Ji, C. J., Robinson, D., Zhu, B., Fang, H. J., Shen, H. H., and Fang, J. Y.:
436 Vegetation and soil ^{15}N natural abundance in alpine grasslands on the tibetan
437 plateau: patterns and implications. *Ecosystems*, 16, 1013–1024,
438 <https://doi.org/10.1007/s10021-013-9664-1>, 2013.

439 Zhang, Q., Zhou, J., Li, X., Yang, Z., Zheng, Y., Wang, J., Lin, W., Xie, J., Chen, Y.,
440 and Yang, Y.: Are the combined effects of warming and drought on foliar
441 C:N:P:K stoichiometry in a subtropical forest greater than their individual effects?
442 *For. Ecol. Manage.*, 448, 256–266, <https://doi.org/10.1016/j.foreco.2019.06.021>,
443 2019.

444 Zhao, Y., Xu, L., Yao, B., Ma, Z., Zhang, C., Wang, F., and Zhou, H.: Influence of
445 simulated warming to the carbon, nitrogen and their stability isotope-($\delta^{13}\text{C}$, $\delta^{15}\text{N}$)
446 contents in alpine meadow plant leaves. *Acta Bot. Boreal Occident Sin.*, 36,
447 777–783, <https://doi.org/10.7606/j.issn.1000-4025.2016.04.0777>, 2016. (in
448 Chinese)

449 Zhu, Q., Castellano, M. J., and Yang, G. S.: Coupling soil water processes and the
450 nitrogen cycle across spatial scales: Potentials, bottlenecks and solutions.
451 *Earth-Sci. Rev.*, 187, 248–258, <https://doi.org/10.1016/j.earscirev.2018.10.005>,

452

2018.

Table 1: Site characteristics from a global meta-analysis of 20 studies.

References	Country/Region	Vegetation types	Soil pH	Soil type	OMC ^a (%)	Latitude	Longitude	Altitude (m a.s.l)	MAT ^b (°C)	MAP ^c (mm)
Anadon-Rosell et al. (2017)	Spain	Subalpine shrub	4.49~4.6 3	Mineral soil	13.15~14.04	41.39 N	2.17 E	2250	3	1146.4
Zhang et al. (2019)	China	<i>C. lanceolata</i> seedlings	5.07	Oxisol	-	26.32 N	117.6 E	300	19.1	1670
Lim et al. (2019)	Sweden	Boreal forests	5.92~6.4 4	A thin, podzolic, sandy soil	-	64.12 N	19.45 E	310	2.4	600
Deane-Coe et al. (2015)	USA	Tundra mosses	-	Gelisol	-	63.88 N	149.23 W	700	-2.7~-1	138~228
Bijoor et al. (2008)	USA	Turfgrass lawn	-	Alkaline alo clay	-	33.7 N	117.7 W	30	18.6	352
Chang et al. (2017)	China	Alpine meadow	9.1~9.3	Gelisols	5.5	34.73 N	92.89 E	4750	-5.3	269.7
Gonzalez-Meler et al. (2017)	Brazil	Grasslands	5.0	Dystrophic red latosols	-	21.17 S	47.86 W	578	21.5	1100
Natali et al. (2012)	USA	Shrubs, sedges and mosses	-	Gelisol	-	63.88 N	149.23 W	700	-1	178~250
Munir et al. (2017)	Canada	Shrubs, mosses and trees	-	-	-	55.27 N	112.47 W			
Salmon et al. (2016)	USA	<i>Eriophorum vaginatum</i>	-	Gelisols	-	63.88 N	149.23 W	700	-1.45	200
Rui et al. (2011)	China	Alpine	-	-	-	37.62 N	101.2 E	3200	-2	500

Aerts et al. (2009)	Sweden	meadow Shrubs, mosses and trees	-	-	-	68.35 N	18.82 E	340	0.5	303
Cheng et al. (2011)	USA	Tallgrass prairie	Neutral pH	Nash-Lucien complex	-	34.98 N	97.52 W		16	911.4
Dawes et al. (2017)	Switzerland	Alpine treeline	-	Sandy Ranker and Podzols	-	46.77 N	9.87 E	2180	9.2	444
Schaeffer et al. (2013)	Greenland	Prostrate dwarf-shrub herb tundra	-	Turbic cryosols	-	76 N	68 W		4~8	<200
Schnecker et al. (2016)	Austria	Spruce forest	Near neutral pH	A mosaic of shallow Chromic Cambisols and Rendzic Leptosols	8.55~14.96	47.58 N	11.64 E	910	6.9	1506
Hudson et al. (2011)	Canada	Heath, willow and meadow	-	-	-	78.88 N	75.78 W		8.6~10. 4	
Lv et al. (2018)	China	<i>Cunninghamia a lanceolata</i> juveniles	-	Red soil	2.21	26.32 N	118 E		19.1	1585
Zhao et al. (2016)	China	Alpine meadow	-	Alpine meadow soils	-	37.48 N	101.2 E	3200~3250	-1.7	600
Peng (2017)	China	Alpine meadow	-	Alpine meadow soils	-	34.73 N	92.89 E	3200~4800	-5.03	267.4~426.3

^aSoil organic matter content; ^bMean annual temperature; ^cMean annual precipitation. If the soil organic carbon content was provided in the literature, soil

organic matter content was determined by multiplying the organic carbon content by a coefficient of 1.724.

List of Figures:

Figure 1: Effect sizes of the experimental warming on soil and plant $\delta^{15}\text{N}$ from a global meta-analysis of 20 studies. The error bars indicate effect sizes and 95% bootstrap confidence intervals (CI). The warming effect was statistically significant if the 95% CI did not bracket zero. The sample size for each variable is shown next to the bar.

Figure 2: Factors influencing the effect sizes of the soil and plant $\delta^{15}\text{N}$ under experimental warming from a global meta-analysis of 20 studies, including (a-b) soil acidity-alkalinity, (c-d) vegetation types, (e-f) warming treatments, (g-h) soil texture, (i-j) length of warming and (k-l) increase in temperature. The error bars indicate effect sizes and 95% bootstrap confidence intervals (CI). The warming effect was statistically significant if the 95% CI did not bracket zero. The sample size for each variable is shown next to the bar.

Figure 3: Relationships between the Hedges' d values of soil and plant $\delta^{15}\text{N}$ with the latitude, altitude, mean annual temperature (MAT) and mean annual precipitation (MAP) under experimental warming.

Figure 1





Figure 2









Figure 3

