

# Spatial variations, origins, and risk assessments of polycyclic aromatic hydrocarbons in French soils

Claire Froger<sup>\*a</sup>, Nicolas P. A. Saby<sup>a</sup>, Claudy C. Jolivet<sup>a</sup>, Line Boulonne<sup>a</sup>, Giovanni Caria<sup>b</sup>, Xavier Freulon<sup>c</sup>,  
5 Chantal de Fouquet<sup>c</sup>, H el ene Roussel<sup>d</sup>, Franck Marot<sup>d</sup>, Antonio Bispo<sup>a,d</sup>

<sup>a</sup> INRAE, US1106 Unit e Infosol, Centre de Recherches d'Orl ans, CS 40001, Ardon, 45075 Orl ans Cedex 2, France

<sup>b</sup> INRAE, US0010 Laboratoire d'analyses des sols, 273 rue de Cambrai 62000 Arras, France

10 <sup>c</sup> Mines ParisTech, Centre de g eosciences. 35, rue saint-Honor e. 77305 Fontainebleau, France. 75006 Paris, France

<sup>d</sup> ADEME, 20 Ave Gresille, F-49004 Angers 01, France

*Corresponding author:* Claire Froger (claire.froger@inrae.fr)

15 **Abstract.** Polycyclic aromatic hydrocarbons (PAHs) are persistent organic pollutants produced by anthropogenic activities that contaminate all environmental spheres, including soils. This study focused on PAHs measured in 2154 soils in France, covering the entire territory based on a regular sampling grid. The quantified concentrations in the  $\Sigma 15\text{PAHs}$  ranged from 5.1 to 31200  $\mu\text{g}\cdot\text{kg}^{-1}$ , with a median value of 32.6  $\mu\text{g}\cdot\text{kg}^{-1}$ , and PAHs were detected in 70% of the soil samples. The map of  $\Sigma 15\text{PAHs}$  concentrations revealed strong spatial variations in soil  
20 contamination throughout France, with larger concentrations in soils of industrial regions and near major cities. PAHs molecular diagnostic ratios supports the historical origin of PAHs in the northern part of France being linked to the significant emissions of PAHs in Europe during the industrial period of 1850–1950 with in particular the contribution of coal/biomass combustion and iron-steel production. A health risk assessment conducted for the residential population resulted in a median value of  $1.07 \times 10^{-8}$  in total lifetime cancer risk, with only 20 sites  
25 above the limit of  $10^{-6}$  and one above the limit of  $10^{-5}$  adopted by the French government. These results reveal the need to conduct large-scale studies on soil contamination to determine the fate of PAHs and evaluate the risks induced by soil pollution at a country-level scale.

Keywords: PAHs; soil monitoring; geostatistics; molecular diagnostic ratios; health risk

## 1 Introduction

30 Polycyclic aromatic hydrocarbons (PAHs) have been targeted by worldwide environmental and public health agencies since 1970 due to their mutagenicity and carcinogenicity (Grimmer, 1985). These substances, which are emitted during the incomplete combustion and pyrolysis of organic matter, are mainly associated with anthropogenic activities, especially 'heavy' industries (e.g. steel and coal), residential heating, and road traffic (Pacyna et al., 2003). Due to more than 150 years of contaminant release via human activities, all environmental  
35 spheres are now affected by persistent organic contaminants, including PAHs. Consequently, European measures on pollution control have been implemented, such as the Water Framework Directive or the Convention on Long-Range Transboundary Air Pollution; however, no such directives exist for soil quality, despite the fact that soil

contamination is a key challenge in the worldwide management of contaminants to protect both human health and ecosystem quality (Carre et al., 2017). Indeed, many studies pointed out that soils can be considered as both a  
40 “sink” and a “source” of PAHs in the environment. The mass balance of PAHs at the scale of a rural catchment conducted by Gocht et al. (2007) in Germany concluded that 90% of PAHs inputs from atmospheric deposition accumulate in the top-soils. Similar work done by Gateuille et al., (2014b) in a small watershed in France showed that atmospheric inputs of PAHs were low compared to the PAHs legacy pollution in soils. In addition, soil erosion has been identified as the predominant source of PAHs at the scale of the Seine River Basin (78,650 km<sup>2</sup>), one of  
45 the largest catchment in France (Gateuille et al., 2020).

The various studies that can be found in the literature on PAH soil contamination mainly focus on local soil pollution (Liu et al., 2016; Yang et al., 2008) or on soils under a particular type of land use (Cachada et al., 2012a; Jensen et al., 2007; Wang et al., 2017a). To efficiently manage soil contamination and prevent the release of pollutants stored in soils, we need to work at the scale of countries to determine the spatial variations in these  
50 contaminants and evaluate their risks to both ecosystems and human populations.

To fully understand the spatial distribution of PAHs, especially at a national scale, we need to determine the origin of the contaminants to potentially enact mitigation techniques at their sources. Considering the widespread contamination of Western Europe by the PAHs released since 1850 (Fernández et al., 2000), it is necessary to determine whether the contamination of soils observed nowadays is the result of legacy pollution or the current  
55 release of these substances. Multiple tracing methods have been developed to identify PAH sources. Among these techniques, PAH molecular ratios have been widely used to discriminate PAHs according to their pyrogenic and petrogenic origins (Ravindra et al., 2008; Yunker et al., 2002); however, these PAH ratios appear to be sensitive to environmental processes such as oxidation and photo-oxidation, resulting in their modification during PAH transport due to the preferential degradation of some PAHs (Biache et al., 2014; Zhang et al., 2005). To prevent  
60 misinterpretations, the use of a combination of multiple PAH ratios is necessary to more accurately identify the origin of PAHs (Kavouras et al., 2001; Tobiszewski and Namieśnik, 2012).

Due to health issues that are attributed to PAHs, an assessment of risks posed by contaminated soil to humans needs to be conducted to proceed with land management decisions. The United States Environmental Protection Agency (US EPA) developed guidelines to conduct human health risk assessments prior to the remediation of  
65 contaminated sites (US EPA 1991), and they then created a manual for soil screening (US EPA 1996). Improvements were made to adapt the risk assessment to the various types of soil contaminant exposure, including residential, occupational, and recreational exposure (US EPA 1996, 2020). In addition, the parameters used in the risk calculations are regularly updated based on the results reported in the literature (IRIS et al., 2017). Therefore, a health risk assessment has been applied in multiple studies addressing PAHs in soils (Albanese et al., 2015;  
70 Cachada et al., 2016; Wang et al., 2017b) using a determinist point estimate approach to first evaluate the health risks posed by soils, as recommended by the US EPA (2001).

Only a handful of studies have examined PAHs in soils at national scales such as those in the United Kingdom (Heywood et al., 2006), China (Ma et al., 2015), Switzerland (Desaules et al., 2010), Germany (Aichner et al. 2013), and the Netherlands (Brus et al., 2009). Unfortunately, the selected sampling design (e.g. Brus et al. 2009),  
75 soil sampling resolution, or land-use specifications (Aichner et al., 2013) have limited the use of these relevant datasets for accurately mapping the PAH spatial distribution at national scales. Currently, only Villanneau et al. (2013) has attempted to map the distribution of a selection of PAHs in French soils based on a coarse sampling

grid corresponding to a subset of 549 soils collected within the framework of the French Soil Quality Monitoring Network (RMQS: Réseau de Mesure de la Qualité des Sols). Their paper provided a first look at the spatial distribution of PAHs at the national scale; however, only four compounds (Benzo(b)fluoranthene, Fluoranthene, Pyrene and Phenanthrene) met the criteria to be mapped using geostatistical technics. The four resulting maps showed a general trend of PAHs in French soils but only broad scale spatial variation were detected in PAHs contamination due to the low-resolution of the input data. As this first work gave limited insights either in the identification of PAHs origins and their distribution, working with the sum of all PAHs molecules is therefore necessary to look further into PAHs distribution over the territory and to identify their origins. Finally, of the large-scale studies of PAHs in soils that have been reported, the spatial distribution and contaminant levels have been the main focus, while health risk assessments and source identification have only been assessed at local scales (Morillo et al., 2008; Wang et al., 2015, 2013). Therefore, it is necessary to conduct complete studies at national scales, including the identification of major PAH sources to understand their spatial distribution and an evaluation of the associated risks to build effective national policies. The objectives of the current study are thus: i) to evaluate the contamination of soils by PAHs in France in comparison with international soil monitoring studies; ii) to produce a high resolution map of the spatial distribution of  $\Sigma 15$ PAHs in French soils; iii) to determine the main origins of PAHs in soils based on PAH molecular ratios; iv) to evaluate the potential health risks for the residential population associated with PAHs in French soils in order to have a first look at the risks induced by the background contamination of soils.

## 2 Materials and methods

### 2.1 Sampling sites

The sampling sites were selected based on a regular grid of 16 km  $\times$  16 km, and they were sampled at the centre of each cell or in a radius of less than 1 km from the theoretical centre when no soil was found at the predicted location (Figure S1). Approximately 2200 sites were sampled, which consisted mostly of agricultural lands (50%), grasslands (20%), and forests (25%), with 1.2% of the region being urbanised areas (including brownfields) and 2.7% of the region being natural areas.

### 2.2 Sampling procedure and analysis

The sampling procedure and PAH analyses have been detailed in Villanneau et al. (2009; 2013); the soils were sampled from June 2000 to May 2010 using a systematic procedure. At each site, in a 20 m  $\times$  20 m grid, 25 core samples of topsoil (approximately 0–30 cm) were taken. The core samples were bulked to obtain a composite sample, which was air-dried and sieved to 2 mm before analysis.

The concentrations of the 16 PAH compounds were measured in the topsoil composite samples: naphthalene (Naph), acenaphthylene (Acy), acenaphthene (Ace), fluorene (Flu), phenanthrene (Phe), anthracene (Ant), fluoranthene (Flh), pyrene (Pyr), benzo(a)anthracene (BaA), chrysene (Chry), benzo(b)fluoranthene (BbF), benzo(k)fluoranthene (BkF), benzo(a)pyrene (BaP), dibenzo(ah)anthracene (DahA), indeno(123,cd)pyrene (IndP), and benzo(ghi)perylene (BghiP). These analyses were conducted at the Laboratory of Soils Analysis of the National Research Institute of Agronomy (INRA, in French) in Arras, France; this laboratory is accredited for

115 PAH analysis in soils by the French accreditation committee (COFRAC). All the analyses were conducted between 2008 and 2012; consequently, volatilization of lighter PAHs such as Naph could have occurred during the long-term storage of soils collected at the beginning of the sampling campaign.

A subsample of ~20 g was taken from topsoil composites, and PAHs were extracted using a Dionex ASE 200 extractor to perform pressure liquid extraction at high pressure (103.4 bar) and temperature (150 °C) with a mixture  
120 of acetone/hexane/dichloromethane (50/25/25; v/v/v). After extraction, the samples were partially concentrated by rotary evaporation to a volume of 5 mL and then completely evaporated under a gentle flow of nitrogen. The dry residue was taken up with 2 mL of methanol and filtered through a 0.2 µm PTFE filter to remove solid particles, which were finally transferred into 2 mL vials. The extracts were then injected into a thermoelectric high-performance liquid chromatograph (HPLC) with a quaternary pump P4000, an autosampler AS3000, a commutator  
125 SN4000 coupled with two Thermolectron detectors mounted in series, a diode array ultraviolet (UV6000), and a spectro-fluorimeter (F3000). A 10 µL-volume aliquot was injected in the HPLC, and the PAHs were separated on a Hypersil PAH column (100 mm in length, 0.46 mm diameter, 5 µm particle size) of a C18 reversed phase. In-depth technical details of the chromatographic parameters used are provided by Villanneau et al. (2009).

External calibration was used to quantify PAHs with concentrations from 40 to 1,000 µg·L<sup>-1</sup> detected by  
130 fluorescence, except for acenaphtylene and indeno(123,cd)pyrene, which were detected by ultraviolet irradiation. The limits of quantification (LOQ) were 0.005 mg·kg<sup>-1</sup> for fluorene and anthracene; 0.01 mg·kg<sup>-1</sup> for acenaphthene, phenanthrene, fluoranthene, pyrene, benzo(a)anthracene, benzo(b)fluoranthene, benzo(a)pyrene, and indeno(123,cd)pyrene; 0.02 mg·kg<sup>-1</sup> for naphthalene and dibenzo(ah)anthracene; 0.03 mg·kg<sup>-1</sup> for acenaphtylene; and 0.05 mg·kg<sup>-1</sup> for chrysene, benzo(k)fluoranthene, and benzo(ghi)perylene. The complete method for PAH  
135 analysis is referenced in the European analytical standard (NF EN 16181; AFNOR, June 2018). The analyses were validated using a reference material (values reported in Table S1) which has been systematically added to the sequence of soil sample analyses to validate the results. One of the 16 PAHs measured, acenaphtylene could not be validated by the reference material (i.e. value of 0 mg·kg<sup>-1</sup> for the reference material see Table S1). Therefore compound Acy quantified only in 1% of the soil samples was excluded from the study considering the remaining  
140 15 PAHs.

Finally, the expanded uncertainty was estimated for every PAH through repeatability and reproducibility tests using a coverage factor of two, which gives a confidence level of 95% (Table S2). In addition, the loss of the most volatile compound Naph could have occurred due to the soil sampling technique using composite samples and due to the soil analysis occurring years after the original sample collection. In addition, to avoid overestimation of the  
145 total PAH content, values under the LOQ were considered to have no PAH content (i.e. 0 µg·kg<sup>-1</sup>).

### 2.3 Geostatistical modelling

A geostatistical modelling approach was used to map the Σ15PAHs and their probability of exceeding a selected threshold (Webster and Oliver, 2007). The observations include many null values (30%) that correspond to values under the LOQ. They also demonstrate extreme values due to locally distinct processes such as point pollution  
150 from industrial sites. For zero-inflated and heavy-tailed distributions, standard geostatistical methods are not the most appropriate because they often lead to poor estimates. In addition, these methods do not properly handle the mapping of risks. Non-linear methods such as indicator kriging, disjunctive kriging, or conditional expectation have been developed for such cases and have found wide acceptance in soil science (Chilès and Delfiner, 2012;

Emery, 2006; Rivoirard, 1994). In this study, the total PAHs were modelled as a transform of a normal Gaussian field,  $Z = \phi(Y)$ . The spatial structure of the Gaussian field is first determined, and then maps of either the total PAHs or the risk of exceeding a threshold are computed by conditional expectation.

In the case of continuous distribution of the original data, the anamorphosis function strictly increases and can be inverted to assign to each observed value its Gaussian counterpart. Then, a common method for variogram estimation can be used (e.g. to calculate the empirical/experimental variogram by the method of moments and then to fit a model to the empirical variogram via non-linear least squares). Here, the proportion of null values is  $p_0 = 30$ ; hence, the transformation function  $\phi$  is identically equal to 0 for all Gaussian values lower than the Gaussian quantile  $y_0 = G^{-1}(p_0)$ . It follows that the actual value of  $Y$  is unknown where  $Z = 0$  but is only constrained by the inequality  $Y < y_0$ .

In such a case, two preliminary steps must be performed: determining the covariance of  $Y$  and then assigning a Gaussian value to each data point at which  $Z$  is null. For the first step, the zero-inflated distribution of the original data is transformed into an inflated Gaussian distribution with an inflated value equal to the quantile of the proportion of observed null values,  $y_0$  (a normal score is applied, and then all values below  $y_0$  are set to this value). Then, to estimate the covariance function of the Gaussian field, the empirical variogram of the inflated Gaussian variable is computed, and the model of the Gaussian field (without inflated values) is fitted by non-linear least squares, taking into account the relationship between the covariances between the Gaussian variables and the covariances between the transformed variables. Once the structure of  $Y$  is determined, the second step consists of simulating the values of the Gaussian field where the  $\Sigma 15\text{PAH}$  value is null, honouring the spatial structure and the inequality constraint. This is classically performed using a Gibbs sampler (Petitgas et al., 2017).

The validity of the best-fitted geostatistical model was then assessed in terms of the standardized squared prediction error (SSPE) using the results of a k-fold cross-validation. If the fitted model was a valid representation of the spatial variation of the Gaussian variable, then the sum of the squares of the errors would have an approximate  $\chi^2$  distribution with a mean of 1 and a median of 0.455 (Lark, 2002).

As both the variogram model and the value of the Gaussian field at all data points are defined, the conditional expectation, also called the multi-Gaussian kriging, can be calculated to map the total PAHs. The simple kriging of the Gaussian variable is computed first, and then the conditional expectation of any transform of the Gaussian field is derived using the following Eq. (1):

$$\{\psi(Y_s)\}^{CE} = \int_{-\infty}^{+\infty} \psi(y_s^{SK} + \sigma_{SK}u)g(u)du, \quad (1)$$

where  $y_s^{SK}$  and  $\sigma_{SK}$  are respectively the kriging value at location  $s$  (the nodes of the grid) and the standard deviation of the kriging error. The transform function  $\psi$  can be either the initial anamorphosis to map the total PAHs or the indicator  $y \mapsto 1_{y \geq \phi^{-1}(z_c)}$  to map the risk that the variable exceeds the threshold  $z_c$ . However, these estimates rely on the Gaussian values being simulated when the original values are null, and these simulated values are used for the kriging computation. Thus, the final estimate is established by the total expectation formula, where the expectation over the Gaussian values constrained by the inequalities is computed by a Monte Carlo integration, i.e. averaging different outcomes produced by the Gibbs sampler.

The geostatistical analysis and the linear and non-linear predictors were performed using the R software 'RGeostats' package (MINES ParisTech/ARMINES 2020).

## 2.4 Health risk assessment

From a toxicological standpoint, PAHs exhibit threshold and non-threshold effects. The risks for residential populations were estimated based on lifetime exposure; for such a duration of exposure, the risks are mainly induced by non-threshold effects. For the ranges of the concentrations measured in soils, it has been verified that the threshold effects are never exceeded, even for shorter exposures, including for children. Thus, only the non-threshold effects of PAHs were considered in the context of this study. The estimation of cancer risk associated with the exposure of individuals to potentially carcinogenic pollutants in French soils was calculated based on the equations developed by the US EPA (USEPA, 1991) and used in several studies (Cachada et al., 2016; Wang et al., 2017a; Yang et al., 2014). The incremental lifetime cancer risks (ILCRs) were calculated for each of the three pathways: soil ingestion, dermal absorption, and soil particle inhalation. To perform the risk assessment, we converted the concentrations of PAHs measured in every soil sample into a BaP equivalent concentration (i.e.  $BaP_{eq,soil}$ ) using Eq. (2):

$$BaP_{eq,soil} = \sum_{i=1}^n PAH_i \times TEF_i, \quad (2)$$

with  $BaP_{eq,soil}$  being the BaP equivalent concentration in soil in  $mg \cdot kg^{-1}$ ;  $PAH_i$  is the concentration of the  $i^{th}$  PAH molecule in the soil ( $mg \cdot kg^{-1}$ );  $TEF_i$  is the toxic equivalent factor for the  $i^{th}$  PAH molecule (see Table S3 in the supplementary material).

We estimated the lifetime cancer risk for the residential population (children + adults) living in the area of the sampling site considering only their direct exposure to contaminated soil (without the ingestion of vegetables collected from the garden). We assumed the local homogeneity of PAHs in soils, considering the PAH concentrations in French soils measured in the current study as the ‘background’ exposure, and therefore focused on the long-term impacts on the population (i.e. non-threshold effects).

Table 1 presents the equations used to calculate the ILCRs for the residential population, and Table 2 details the values of parameters used in the calculations based on the risk assessment guidance of the US EPA (1991, 1996, 2020).

The total lifetime cancer risk (TLCR) was calculated by summing the three ILCRs for the residential population and outdoor workers.

**Table 1: Equations used for the health risk assessment of the residential population.**

Pathway	Residential population
Ingestion	$ILCR_{ing,res} = \frac{C_{soil} \times IF_{soil/adj}}{AT \times 10^6} \times CSF_0$
	where $IF_{soil/adj} = \left( \frac{EF_{c,res} \times ED_{c,res} \times IRS_{c,res}}{BW_c} + \frac{EF_{a,res} \times (ED_{a,res} - ED_{c,res}) \times IRS_{a,res}}{BW_a} \right)$
Dermal	$ILCR_{der,res} = \frac{C_{soil} \times DFS_{soil/adj} \times ABS_d}{AT \times 10^6} \times \frac{CSF_0}{GIABS}$
	where $DFS_{soil/adj} = \left( \frac{EF_{c,res} \times ED_{c,res} \times SA_{c,res} \times AF_{c,res}}{BW_c} + \frac{EF_{a,res} \times (ED_{a,res} - ED_{c,res}) \times SA_{a,res} \times AF_{a,res}}{BW_a} \right)$
Inhalation	$ILCR_{inh,res} = \frac{C_{soil} \times EF_{a,res} \times ED_{a,res} \times ET_{a,res} \times (1/PEF)}{AT} \times IUR$
Total	$TLCR_{res} = ILCR_{ing,res} + ILCR_{der,res} + ILCR_{inh,res}$

**Table 2: Detailed parameters used in the incremental lifetime cancer risk assessment based on the US EPA (2020).**

<b>Parameter</b>		<b>Units</b>	<b>Value</b>
<b>AT</b>	Averaging time	d	25550 (365×70)
<b>EF<sub>c,res</sub></b>	Exposure frequency (child)	d·y <sup>-1</sup>	350
<b>ED<sub>c,res</sub></b>	Exposure duration (child)	y	6
<b>IRS<sub>c,res</sub></b>	Soil ingestion rate (child)	mg·d <sup>-1</sup>	200
<b>BW<sub>c</sub></b>	Body weight (child)	kg	15
<b>EF<sub>a,res</sub></b>	Exposure frequency (adult)	d·y <sup>-1</sup>	350
<b>ED<sub>a,res</sub></b>	Exposure duration (adult)	y	70
<b>IRS<sub>a,res</sub></b>	Soil ingestion rate (adult)	mg·d <sup>-1</sup>	100
<b>BW<sub>a</sub></b>	Body weight (adult)	kg	60
<b>SA<sub>c,res</sub></b>	Surface area (child)	cm <sup>2</sup>	2373
<b>AF<sub>c,res</sub></b>	Adherence factor (child)	mg·cm <sup>-2</sup>	0.2
<b>SA<sub>a,res</sub></b>	Surface area (adult)	cm <sup>2</sup>	6032
<b>AF<sub>a,res</sub></b>	Adherence factor (adult)	mg·cm <sup>-2</sup>	0.07
<b>ABS<sub>d</sub></b>	Dermal absorption factor	-	0.13
<b>GIABS</b>	Gastrointestinal absorption factor	-	1
<b>PEF</b>	Particle emission factor	m <sup>3</sup> ·kg <sup>-1</sup>	1.32×10 <sup>9</sup>
<b>CSF<sub>0</sub></b>	Cancer oral slope factor	(mg·kg <sup>-1</sup> d <sup>-1</sup> ) <sup>-1</sup>	1
<b>IUR</b>	Inhalation unit risk	(mg·m <sup>-3</sup> ) <sup>-1</sup>	0.6

### 3 Results and discussion

#### 3.1 PAH contents and the influence of soil characteristics

##### 3.1.1 PAH concentrations and composition

225 Descriptive statistics of the dataset are presented in Table 3, showing summary statistics for each PAH molecule and for the sum of the 15 PAHs. Over the 2154 soils sampled, PAHs (at least one molecule) could be quantified in 1512 soils corresponding to 70% of the samples. Among the 1512 soils with measurable PAH concentrations, 732 contained more than 6 PAH molecules above the LOQ.

230 Fluoranthene and phenanthrene were quantified in 56% and 55% of the soil samples (Table 4), respectively, followed by benzo(b)fluoranthene (48%) and pyrene (46%). Conversely, acenaphthene was measured in less than 2% of the total soil samples.

**Table 3: Summary of descriptive statistics in  $\mu\text{g}\cdot\text{kg}^{-1}$ .**

PAH molecule	N > LOQ	N > LOQ (%)	SD/mean	Min	Q25	Q50	Q75	Max	Mean
Naphthalene	127	6	9.5	< LOQ	< LOQ	< LOQ	< LOQ	1030	2.7
Acenaphthene	38	2	13.1	< LOQ	< LOQ	< LOQ	< LOQ	160	0.3
Fluorene	231	11	6.1	< LOQ	< LOQ	< LOQ	< LOQ	250	1.4
Phenanthrene	1176	55	4.7	< LOQ	< LOQ	11.2	19.275	3470	19.0
Anthracene	174	8	10.4	< LOQ	< LOQ	< LOQ	< LOQ	555	1.9
Fluoranthene	1201	56	5.1	< LOQ	< LOQ	12.05	26	6080	30.3
Pyrene	996	46	5.1	< LOQ	< LOQ	< LOQ	19.575	4370	22.4
Benzo(a)anthracene	559	26	5.5	< LOQ	< LOQ	< LOQ	10.2	2180	11.3
Chrysene	180	8	9.0	< LOQ	< LOQ	< LOQ	< LOQ	4140	11.3
Benzo(b)fluoranthene	1039	48	3.6	< LOQ	< LOQ	< LOQ	18.5	2220	18.0
Benzo(k)fluoranthene	825	38	< 4.9	< LOQ	< LOQ	< LOQ	8.0675	1460	8.2
Benzo(a)pyrene	658	31	4.7	< LOQ	< LOQ	< LOQ	12.3	1730	12.8
Dibenzo(ah)anthracene	148	7	8.6	< LOQ	< LOQ	< LOQ	< LOQ	1130	3.5
Indeno(123cd)pyrene	655	30	4.7	< LOQ	< LOQ	< LOQ	12.1	1830	10.8
Benzo(ghi)perylene	135	6	6.8	< LOQ	< LOQ	< LOQ	< LOQ	1530	7.2
$\Sigma 15\text{PAHs}$	1512	70	5.1	< LOQ <sup>a</sup>	< LOQ <sup>a</sup>	32.6	121.8	31193	161.0
BaP <sub>eq</sub>	1512	70	5.2	< LOQ <sup>a</sup>	< LOQ <sup>a</sup>	1.13	17.5	3705	21.4
2–3-ring PAHs (%)				NA	NA	10.3	21.3	100	18.5
4-ring PAHs (%)				NA	NA	36.3	49.2	100	28.9
5–6-ring PAHs (%)				NA	NA	22.8	39.8	100	22.7

<sup>a</sup>: for the sum of the 15 PAHs, < LOQ means that no PAH molecules were quantified

235 The results highlight the large differences between the levels of soil contamination, with PAH concentrations varying from 5.1 to 31200  $\mu\text{g}\cdot\text{kg}^{-1}$  in the  $\Sigma 15\text{PAHs}$ . However, extreme values were not frequent, as 90% of the PAH concentrations measured were under 500  $\mu\text{g}\cdot\text{kg}^{-1}$  in the  $\Sigma 15\text{PAHs}$ , and only 4% of the concentrations were above 1000  $\mu\text{g}\cdot\text{kg}^{-1}$  in the  $\Sigma 15\text{PAHs}$ .

240 The concentrations of the  $\Sigma 15\text{PAHs}$  measured in the current study were in the same range as the PAH contents measured in rural soils from various locations in Europe and China (Table 4), which also demonstrated large variations in the  $\Sigma 15\text{PAH}$  concentrations. The highest concentrations reported in the literature were mostly attributed to local sources, such as urban and industrial areas. Results similar to those in this study were found in Poland with respect to the PAH distribution, with 75% of the soil samples being characterized by PAH content  $\leq 694 \mu\text{g}\cdot\text{kg}^{-1}$  (Maliszewska-Kordybach et al., 2008).



245 **Table 4: PAH concentrations in soils reported in the literature in  $\mu\text{g}\cdot\text{kg}^{-1}$ .**

	Location	Land-use type	Concentrations in $\mu\text{g}\cdot\text{kg}^{-1}$	Number of PAHs considered	
	This study		5.1–31193	15	
<b>France</b>	Orgeval	Rural (n = 25) and urban (n = 8)	60–5305	13	(Gateuille et al., 2014b)
	Rouen	Rural (forest)	450–5650	14	(Motelay-Massei et al., 2004)
	Paris conurbation	Rural (n = 12) and urban (n = 20)	150–55000	15	(Gaspéri et al., 2018a)
<b>Norway</b>	Transect to Oslo	Forest soils	10–2600	16	(Jensen et al., 2007)
<b>Poland</b>	Entire country	Rural (n = 217)	80–7264	16	(Maliszewska-Kordybach et al., 2008)
<b>Switzerland</b>	Entire country	Urban (n = 2), semi-rural (n = 13), rural (n = 7), remote (n = 1)	19–6870	16	(Gubler et al., 2015)
<b>Germany</b>	Entire country	Forest soils (n = 447)	105–14889	16	(Aichner et al., 2013)
<b>UK</b>	England, Scotland, Wales, and Northern Ireland	Rural (n = 122)	24–128000 (mean: 608)	16	(Bull and Collins, 2013)
	Entire country	Rural (n = 120) and urban (n = 28)	9.9–5910 (mean: 377)	16	(Ma et al., 2015)
<b>China</b>	Dongjiang River Basin	Rural (n = 30)	23.5–231 (mean: 116)	16	(Zheng et al., 2014)

### 3.1.2 Influence of soil physicochemical parameters

The PAHs measured in the samples were mostly 4- and 5–6-ring PAHs respectively representing median values of 36% and 23% of the total PAHs, with the 2–3-ring PAH contribution being around 10% (Table 4). These observations are in accordance with literature values showing a lower content of light PAHs in soils over time and a stable content of heavy PAHs (Cui et al., 2020; Gubler et al., 2015). The sorption of PAHs on organic matter due to their lipophilic properties is known to be a key factor in PAH evolution in soils (Maliszewska-Kordybach et al., 2008; Yu et al., 2018). As the soil organic matter structure is strongly influenced by pH, the fate of PAHs could also be influenced by the soil pH, especially in forest soils (Aichner et al., 2013; Wenzel et al., 2002). Nevertheless, in the current study, no clear influence of organic carbon, organic matter (when available), or pH on the PAH content was evident, except for in urban parks (see Figures S2, S3, and S4 in the supplementary material). The low or non-existent impact of soil characteristics on the PAH concentrations found in multiple studies (Aichner et al., 2013; Maliszewska-Kordybach et al., 2008) could be attributed to the stronger influence of other factors such as source proximity and meteorological parameters (Heywood et al., 2006; Lohmann et al., 2000). In soils receiving identical amounts of PAHs emitted by anthropogenic activities, local variations in the pH and total organic carbon might exert an influence on the fate of those contaminants in soils (Wenzel et al., 2002). This is supported by the correlation between the PAH content and soil characteristics for soils in urban parks, as urban areas have highly concentrated PAH emissions from sources such as road traffic, industries, and residential emissions (Cachada et al., 2012b; Keyte et al., 2016). However, considering that five soils in this study were collected from urban parks, a similar conclusion cannot be made based on our dataset. At the scale of the entire

country, the influence of soil characteristics on PAH content would be minimized by the proximity of major emission sources (Aichner et al., 2013).




270 In addition to soil physicochemical characteristics, relationships between the PAH content in soils and land-use type have been noted in some studies (Maliszewska-Kordybach 1999); however, no differences in PAH content were observed among forests, cultivated lands, and permanent pastures in this study (see Figure S4 in the supplementary material). Only soil located in natural areas seems to present a lower content of PAHs (Kruskal-Wallis test,  $p$ -value < 0.05), but these results should be considered with caution since the number of natural areas accounted for less than 5% of the 2154 sites (Section 2.1). Therefore, agricultural practices appear to have a minor influence on the PAH concentrations measured in French soils.

## 275 3.2 Spatial distribution of PAHs

### 3.2.1 PAH-level distribution over France

280 In this study, we provide the first national map of the  $\Sigma 15$ PAHs in Figure 2A using multi-Gaussian kriging, which is a standard and versatile method. Once the kriging of the Gaussian variable is computed, maps for various transform functions can be easily derived. Here, it is particularly difficult to properly handle the LOQ, so a Monte Carlo integration was added for this purpose.

No evident anisotropy can be seen on the sample variogram; the omnidirectional estimated variogram and fitted model are shown in Figure 2C. The variogram displays a very large nugget effect, showing the substantial short-range variability for the  $\Sigma 15$ PAHs induced by the contamination process. A nested model was used to combine a nugget and a spherical model (range = 488 km). The conditional expectation approach assumes a stationary Gaussian field model. To meet this assumption, we forced the sill of the variogram model to 1, which corresponds to the total variance of the normally scored transformed variable. Notably, the estimated variogram continuously increases after 400 km, which corresponds to the north-to-south gradient observed in the data; however, this issue could be easily solved via point-kriging with a moving 250 km radius neighbourhood and a maximum of 150 data points (Rivoirard and Romary, 2011) so that only the well-fitted part of the variogram intervenes. The results of 290 the 10-fold cross-validation gave a mean SSPE value of 0.92 and a median SSPE value of 0.447, with both values falling within the confidence interval of the theoretical values. Finally, the  $R^2$  in the transformed space equalled to 0.33.



295 **Figure 1: (a) The estimated spatial distribution of the total PAH ( $\Sigma 15\text{PAHs}$ ) concentrations in  $\mu\text{g}\cdot\text{kg}^{-1}$  by conditional expectation along with major cities in France; (b) map of the estimated probability of exceeding  $100 \mu\text{g}\cdot\text{kg}^{-1}$  in the  $\Sigma 15\text{PAHs}$ ; (c) estimated (in red) and fitted (in black) variograms of the normally scored transformed  $\Sigma 15\text{PAHs}$ .**

The spatial distribution of the PAH concentrations in French soils highlighted wide variations in the contamination throughout the French territory. The north-eastern part of France, known to be an industrialised region, showed the highest levels of PAHs in the soils, as noted by Villanneau et al. (2013) for benzo(b)fluoranthene, fluoranthene, pyrene, and phenanthrene. In addition, higher PAH concentrations could be observed in the region of the Rhône

300

Valley, which is also known to be industrialised, and around major harbour cities along the west coast (Figure 1B). Finally, hot spots of PAHs were found near major cities such as Paris, Marseille, Lyon, Nantes or Bordeaux (Figure 1A). Those late observations contradicts the results of Villanneau et al. (2013), showing mostly the North-East – South West gradient without any evidence of cities impacts on soil contamination by PAHs.

305 To evaluate the state of contamination of the French soils, PAH concentrations should be compared with international historical background values, as PAHs could be emitted by natural sources such as wild fires during the Jurassic period (Killops and Massoud, 1992). Some studies have suggested a lower limit of 100  $\mu\text{g}\cdot\text{kg}^{-1}$  in the total PAHs, as in Canada (Canadian Council of Ministers of the Environment, 2010) or in the Seine River basin (Gaspéri et al., 2018b; Gateuille et al., 2014c). Maliszewska-Kordybach (1996) suggested a classification of  
310 agricultural soils in Poland based on the  $\Sigma 15\text{PAHs}$  concentrations, as presented in Table 5.

**Table 5: The soil contamination classification suggested by Maliszewska-Kordybach (1996) and a design-based estimation of the proportion of associated French soils.**

Soil classification	PAH ranges ( $\mu\text{g}\cdot\text{kg}^{-1}$ )	Proportion of French soils (in %)
Non-contaminated	< 200	83
Weakly contaminated	200–600	12
Contaminated	600–1000	2.1
Heavily contaminated	> 1000	2.9

The values outlined in Table 5 are supported by studies of pre-industrial periods (i.e. before 1830) such as  
315 Fernández et al. (2000), which showed PAH concentrations of 20–100  $\mu\text{g}\cdot\text{kg}^{-1}$  in 1830 in lake sediment cores across Europe. Other studies reported PAH concentrations under 200  $\mu\text{g}\cdot\text{kg}^{-1}$  in sediments from 1400 to 1850 (Elmquist et al., 2007; Leorri et al., 2014).

Based on the classification of Maliszewska-Kordybach (1996), the proportions of French soils attributed to each soil class were calculated (Table 5), showing that 17% could be considered contaminated when using a threshold  
320 value of 200  $\mu\text{g}\cdot\text{kg}^{-1}$ . When applying the Canadian background value of 100  $\mu\text{g}\cdot\text{kg}^{-1}$  (Table 5), 30% of the French soils could be considered contaminated (i.e. above the threshold value). When considering the outputs of the geostatistical model (Figure 1A), 30% of the surface of France has PAH concentrations in soils higher than the background value of 100  $\mu\text{g}\cdot\text{kg}^{-1}$ . To better assess the risk of exceeding this background value, we used a  
325 geostatistical model to map the probability that the  $\Sigma 15\text{PAHs}$  concentration in soils exceeds 100  $\mu\text{g}\cdot\text{kg}^{-1}$  (Figure 1B). Industrialised regions in northern France exhibit probabilities of exceeding 100  $\mu\text{g}\cdot\text{kg}^{-1}$  above 0.8. High probabilities of soils exceeding the threshold value were also found in the vicinity of the biggest cities in France, such as Paris, Marseille, Lyon, Nantes, La Rochelle, and Bordeaux (Figure 1B).

### 3.2.2 Enhancing spatial predictions

To take into account the uncertainty of the local mean, the ordinary multi-Gaussian kriging (Emery, 2006) was  
330 tested, but there is no significant difference from the standard conditional expectation presented here, as the ordinary kriging is very close to the simple kriging due to regular sampling. The variogram model of the latent Gaussian variable presents a significant nugget effect, which may result from local contamination or measurement errors, as previously mentioned. This introduces the issue of the support considered for the measurement (20 m  $\times$

20 m) and for the interpolated grid (with a mesh of 2000 m × 2000 m). Taking into account that the support and measurement errors in the Gaussian model may improve the spatial definition (see Chilès and Delfiner 2012 for a review of models for support changes and example applications).

The standard empirical variogram averages the squared differences from different zones: the variogram model is isotropic and stationary, but final maps suggest a north/south (or north-west/south-east) contrast and local anisotropies, which may or may not be partially explained by external co-variates (e.g. topography, distance to the sea, and dominant wind direction). Hence, modelling local anisotropies (Fouedjio et al., 2016) or using a non-stationary SPDE model (Pereira, 2019) could improve the structured part of the variogram model and, hence, the maps.


### 3.3 Major origin of PAHs using molecular diagnostic ratios

To identify the origin of PAHs in French soils, the four widely used molecular diagnostic ratios—Flh/(Flh + Py), Ant/(Ant + Phe), BaA/(BaA + Chry), and IdP/(IdP + BghiP)—were investigated to distinguish among petrogenic (i.e. crude oil), pyrogenic (coal, wood, and biomass combustion), and fossil fuel combustion sources (Table 6). We computed these ratios and mapped the results using a simple point map (Figure 2) without any interpolation, and missing values corresponded to sites at which none of the PAHs used in the ratios were quantified.


**Table 6: Most commonly used molecular diagnostic ratios associated with PAH origins.**

Ratio	Petrogenic	Fossil fuel combustion	Pyrogenic	References
<b>Flh/(Flh + Py)</b>	0–0.4	0.4–0.5	0.5–1	
<b>Ant/(Ant + Phe)</b>	0–0.1		0.1–1	(Biache et al., 2014; Ravindra et al., 2008;
<b>BaA/(BaA + Chry)</b>	0–0.2	0.2–0.35	0.35–1	Tobiszewski and Namieśnik, 2012; Yunker et al., 2002; Zhang et al., 2005)
<b>IdP/(IdP + BghiP)</b>	0–0.2	0.2–0.5	0.5–1	


Flh/(Flh + Pyr)




Ant/(Ant + Phe)



BaA/(BaA + Chry)



IndP/(IndP + BghiP)



350

**Figure 2: Spatialized molecular diagnostic ratios in French soils. Missing values correspond to sites at which none of the PAHs used in the ratios were quantified.**

The spatial distribution of the four diagnostic ratios appears to be similar to the majority of calculated ratios in the northern part of France (Figure 2) but with different indicators with respect to the PAH origins. The values of the Flh/(Flh + Pyr), IdP/(IdP + BghiP), and BaA/(BaA + Chry) ratios are mainly associated with pyrogenic signatures attributed to coal, wood, and biomass burning (Figure 2A, C, D), whereas the Ant/(Ant + Phe) values indicate petrogenic signatures attributed to crude oil (Figure 2B). This difference in the PAH origins based on the Ant/(Ant + Phe) ratio could be explained by the sensitivity of this ratio to environmental processes such as oxidation or biodegradation (Katsoyiannis and Breivik, 2014; Tobiszewski and Namieśnik, 2012). Phenanthrene and anthracene are lighter PAHs susceptible to multiple processes during their transport, shifting the source signature of Ant/(Ant + Phe) (Biache et al., 2014; Zhang et al., 2005). The lower detection of anthracene (8%) compared to phenanthrene (55%), resulting in many instances of Ant/(Ant + Phe) being close to 0, probably reflects anthracene

355

360



depletion rather than PAH origins. Consequently,  $\text{Ant}/(\text{Ant} + \text{Phe})$  should be used with caution when trying to identify the origin of PAHs and was thus considered an unreliable source signature in this study.

As shown by the  $\text{Flh}/(\text{Flh} + \text{Py})$ ,  $\text{IdP}/(\text{IdP} + \text{BghiP})$ , and  $\text{BaA}/(\text{BaA} + \text{Chry})$  diagnostic ratios (Figure 2A, C, and D), pyrogenic combustion such as coal and wood burning might be the main origin of PAHs measured in the soil of the north-eastern part of France and the Rhône Valley in the south-east. In the literature,  $\text{Flh}/(\text{Flh} + \text{Py})$  and  $\text{IdP}/(\text{IdP} + \text{BghiP})$  have been identified as the most conservative ratios due to the lower sensitivity of Flh, Py, IdP, and BghiP to degradation (Brändli et al., 2008; Tobiszewski and Namieśnik, 2012). In the case of  $\text{BaA}/(\text{BaA} + \text{Chry})$ , shifts in the signature have been observed during atmospheric transport due to photo-degradation (Katsoyiannis and Breivik, 2014); however, the concordance of the latter ratio with the PAH origins identified using the  $\text{Flh}/(\text{Flh} + \text{Py})$  and  $\text{IdP}/(\text{IdP} + \text{BghiP})$  diagnostic ratios suggests that it is a reliable molecular diagnostic ratio in this study. The spatial trend exhibited by  $\text{Flh}/(\text{Flh} + \text{Py})$  ratio for the sites with non-missing value combined with  $\text{IdP}/(\text{IdP} + \text{BghiP})$  and  $\text{BaA}/(\text{BaA} + \text{Chry})$  ratios seems to confirm the pyrogenic origin of PAHs in the north of France despite the substantial number of missing values for  $\text{IdP}/(\text{IdP} + \text{BghiP})$  and  $\text{BaA}/(\text{BaA} + \text{Chry})$  due to a lower detection of the PAH molecule used in these ratios.

In addition to the indication of pyrogenic origins of the PAHs by molecular diagnostic ratios, the homogeneous spatial pattern of both PAH concentrations and the ratio in north-eastern France (Figure 1 and 2) suggest contamination linked to regional atmospheric emissions, the latter being the major source of soil PAHs (Tobiszewski and Namieśnik, 2012). Multiple studies reported high PAH deposition rates in Western Europe from 1900 to 1970 due to industrial development, especially from heavy industries (e.g. coal and steel), with a massive peak in 1960 before a decrease due to pollution control (Elmqvist et al., 2007; Fernández et al., 2000; Gabrieli et al., 2010; Leorri et al., 2014). An estimation of the BaP emissions in Europe by Pacyna et al. (2003) identified Germany, France, and the UK as the main contributors, emitting 22%, 8%, and 11% of the 1300 tons of BaP released in 1970, respectively. In addition, the Gabrieli et al. (2010) study of atmospheric PAH deposition beginning in 1700 showed that, during 1900–1960, the  $\text{Flh}/(\text{Flh} + \text{Py})$  ratio decreased from 0.65 to 0.55. This decreasing trend was explained by the increasing contribution of fossil fuel combustion (gasoline, diesel) to local atmospheric PAH deposition (Gabrieli et al., 2010). This value of 0.6 in  $\text{Flh}/(\text{Flh} + \text{Py})$  appears to be the main value in the north-eastern part of France (Figure 2A), therefore supporting historical industrial PAH emissions as a major source of PAHs in the soils of this region. This hypothesis of the coal industry's contribution to PAH loads in Northern France during the mid-20<sup>th</sup> century was also suggested by Lorgeoux et al. (2016) using organic pollutant records from a sediment core of the Seine estuary.

The persistence of PAHs over more than a century and the absence of changes in the signature could be explained by multiple factors. First, the amount of PAHs emitted at the beginning of the 20<sup>th</sup> century was 10 to 100 times higher than the current PAH depositions (Fernández et al., 2000; Gabrieli et al., 2010). The high rates of PAH deposition during those 30 to 60 years could have prevented any changes in the PAH signatures in soils associated with source modifications after 1960 due to the stock of PAHs already polluting soils (Froger et al., 2019; Gateuille et al., 2014b). Second, the process of PAH sequestration in the soil could explain their long-term persistence in French soils (Biache et al., 2011). This sequestration is caused by the sorption of PAHs on organic matter and their diffusion through micropores in soils, reducing their bioavailability for microorganisms (Bogan and Sullivan, 2003; Semple et al., 2003). Overall, the aging of soils could increase PAH interactions in soils, enhancing the slow

but irreversible sequestration of PAHs and reducing their potential degradation in soils (Anyanwu and Semple, 2015; Duan et al., 2015; Ma et al., 2012; Ouvrard et al., 2013).

405 Additionally, the hotspots of PAH concentrations around major cities (Figure 1) suggests local sources of contaminants, such as in urban or suburban environments, and multiple sources of PAHs could contribute to PAH soil contamination. Current sources of PAHs, such as road traffic, local industries, and residential emissions (household heating) (Cachada et al., 2012a; Gateuille et al., 2014a; Lohmann et al., 2000) as well as legacy contamination due to former industries (Bertrand et al., 2015; Pies et al., 2007) could contribute to soil PAH  
410 content. However, precise source identification and apportionment cannot be done based only on PAH molecular diagnostic ratios, which only provide information about pyrogenic, petrogenic, or mixed PAHs in soils (Brändli et al., 2008; Tobiszewski and Namieśnik, 2012; Yunker et al., 2014). Multivariate analysis associated with linear mixing models and positive matrix factorization could be performed to reveal PAH sources and their respective contributions to PAH contamination in soils (Harrison et al., 1996; Larsen and Baker, 2003; Wang et al., 2015).  
415 Additionally, each PAH molecule may present different spatial structures depending on its source. Minimum and maximum auto-correlation factors (Petitgas et al., 2018; Switzer and Green, 1984) deduced from different indicators (for example, the indicators of the  $1_{PAH_i} \geq LOC_{LOQ}$ ) could be used. These factors are the spatial counterpart of the standard PCA, and their mapping could help regionalize the origin of PAHs and estimate the risk using alternate non-linear methods such as discrete disjunctive kriging. Finally, other fingerprinting methods  
420 involving new analyses such as carbon isotopes (Kim et al., 2008; Petrišič et al., 2013; Zhang et al., 2020) or alkyl-PAHs (Morales-Caselles et al., 2017; Yuan et al., 2015) could be used to complete source identification and apportionment.

### 3.4 Health risk assessment

#### 3.4.1 Total lifetime cancer risks in France

425 The results of the health risk assessment evaluated for the residential population are presented in Table 7. According to the US Environmental Protection Agency, the elevated risk is attributed to a TLCR above  $10^{-4}$ , meaning that the probability of an individual developing cancer over a lifetime is more than one in 10000. A TLCR value under  $10^{-6}$  is considered to indicate virtual safety, meaning that the probability of an individual developing cancer is less than one in 1000000. Consequently, TLCRs between  $10^{-4}$  and  $10^{-6}$  were associated with a ‘moderate  
430 risk’. However, the limit of  $10^{-6}$  is commonly used by the US EPA as the target risk value to calculate screening levels in soils. In France, the ministerial circular on December 10<sup>th</sup>, 1999 chose a value of  $10^{-5}$  as the limit defining the sites needing remediation for polluted soils.

Among the 2154 soils collected throughout the French territory, 54% had a TLCR value (i.e. 1512 sites), indicating that those soils were contaminated by at least one of the seven carcinogenic PAHs. TLCR values for the residential  
435 population ranged from  $1.89 \times 10^{-11}$  to  $1.37 \times 10^{-5}$ , with 20 sites presenting TLCRs above the safety target value of  $10^{-6}$  (i.e. 1% of the sites). Only one site showed a TLCR value above  $10^{-5}$ , which was considered a polluted site according to French legislation. Ingestion of soil contributed to 68.8% of the total cancer risk compared to dermal contact (31.2%) and inhalation (0.01%), which is commonly observed in the literature (Tong et al., 2018; Wang et al., 2017b).

440 Considering that soil samples were collected mainly in locations distant from anthropogenic sources (both industries and urban areas), the current study could be considered an evaluation of the background level of PAHs




445 in French soils. Consequently, soils contaminated by PAHs showed low risks for residential populations, considering that 99% of the soils presented TLCR values under  $10^{-6}$  (Table 7). In the literature, higher TLCR values have been identified in southern Italy with mean ILCRs of  $4.77 \times 10^{-6}$  for adults (Qi et al., 2020) and in China with a mean TLCR value of  $1.86 \times 10^{-6}$  in Shanghai (Tong et al., 2018) and an ILCR value of  $4.4 \times 10^{-5}$  in the Yangtze River Delta (Wang et al., 2017b). However, those studies were conducted mainly on industrial and highly urbanised regions and reflect the risks induced by soils under high anthropogenic pressure and not the background contamination over a large territory.

**Table 7: Summary of the total lifetime cancer risk (TLCR) attributed to French soils for the residential population.**

<b>TLCR residential population</b>	
<b>min</b>	$1.89 \times 10^{-11}$
<b>max</b>	$1.37 \times 10^{-5}$
<b>mean</b>	$1.13 \times 10^{-7}$
<b>q25</b>	$3.82 \times 10^{-9}$
<b>q50</b>	$1.07 \times 10^{-8}$
<b>q75</b>	$9.63 \times 10^{-8}$
<b>Number of sites &lt; <math>10^{-7}</math></b>	1144
<b>Number of sites between <math>10^{-7}</math> and <math>10^{-6}</math></b>	348
<b>Number of sites &gt; <math>10^{-6}</math></b>	20

## Risk for residents




450

**Figure 3: Health risk assessment for the residential population showing the total lifetime cancer risk (TLCR) for French soils along with the location of major cities. Missing values correspond to sites at which no PAHs were quantified, so a TLCR below  $10^{-7}$  can be assumed.**

455 Figure 3 presents the residential maps of TLCRs and indicates that the north-eastern part of France and the Rhône Valley had the most sites with estimated TLCRs reflecting the national PAH concentration distribution (Figure 1A). These observations can be explained by the positive correlation between the BaP toxic equivalent and the  $\Sigma 15$ PAHs shown in Figure 4. These results indicate that the contamination of soils by PAHs is driven by the most

carcinogenic PAHs with the highest TEFs (i.e., BaA, Chr, BbF, BkF, BaP, IndP, and DahA), whose concentrations are proportional to the total PAHs measured.



460

**Figure 4: Correlation between the BaPeq in mg·kg<sup>-1</sup> and the concentration of the  $\Sigma$ 15PAHs in µg·kg<sup>-1</sup> in French soils.**

The soils with TLCRs between  $10^{-7}$  and  $10^{-6}$  are concentrated in the northern part of France and in the vicinity of the main cities (Figure 3). Among the 17 main cities shown in Figure 3, 11 are located in areas in which TLCRs could be calculated, and four are situated in areas with multiple sites with TLCRs between  $10^{-7}$  and  $10^{-6}$ . As multiple sources of PAHs are concentrated in urban areas, soils with higher contamination levels that are usually found in cities might pose higher risks for the population, as in Naples, where the ILCRs were above  $1 \times 10^{-5}$  (Albanese et al., 2015). In France, a database of 300 urban soils was created (ADEME, 2018), and the soils therein had a median value of  $1560 \mu\text{g}\cdot\text{kg}^{-1}$  in the  $\Sigma$ 16PAHs, which is 50 times higher than the  $32 \mu\text{g}\cdot\text{kg}^{-1}$  in the  $\Sigma$ 15PAHs in the current study. Therefore, the higher risks posed by PAHs in soils would be expected in urbanised regions.

465

In addition, the distribution of soils with higher TLCRs around cities (especially Paris, Lille, and Marseille) highlights the influence of densely urbanised areas on the surrounding soils over a distance of up to several dozen kilometres (Jensen et al., 2007). Finally, the 20 sites with the highest TLCRs above  $10^{-6}$  appeared to be randomly distributed throughout the territory, suggesting local sources of PAHs at those sites, such as heavy industries.

470

The first results of the background national health risks linked to PAHs show the spatial heterogeneity in the risks encountered by the population and the necessity of considering soils in risk evaluations and the national surveillance of contaminants in the environment.

475

### 3.4.2 Limits and perspectives

As we used a site-specific determinist method to predict the TLCR at each site, the variations in the population characteristics and model parameters were not integrated in the calculation. Therefore, the use of a probabilistic model with a sensitivity analysis would result in more accurate predictions of cancer risks based on French population statistics (US EPA 2001; Bruce et al. 2007; Tong et al. 2018). However, the use of the point estimate

480

approach is the first step recommended by the US EPA (2001) when conducting risk assessments. These results provided an overview of the health risks associated with background PAH concentrations in French soils and the need to further evaluate these risks.

485 Three pathways of human exposure to PAHs in the soils were considered (i.e. ingestion, dermal contact, and inhalation). To integrate all exposure routes, PAH transference from soil to food production (crops, vegetables, and fruits) should be taken into account as well as groundwater transfer. Many studies have demonstrated that exposure through food ingestion significantly contributes to the total risk, as in the Yangtze River Delta, where TLCRs rose from  $4.4 \times 10^{-5}$  to  $9.17 \times 10^{-4}$  (Wang et al., 2017b). Similarly, in southern Italy, the risk induced by  
490 food ingestion exceeded the risks of the other pathways by two orders of magnitude (Qi et al., 2020). Therefore, the potential risks posed by PAHs in French soils could be underestimated, especially since a significant part of the production of cereals (soft wheat and barley) and some vegetables (potato) are cultivated in northern France. Consequently, a more in-depth health risk study integrating land-use types (crops, grassland, and forest), produce cultivation, and a scenario in which the residential population raises their own vegetables would result in deeper  
495 insight into the risks induced by PAHs in soils.

The risk estimations were made based on the additional toxic effects of PAHs using toxic equivalent factors (TEFs). However, little is known about the potential synergistic, antagonist, or additive effects of PAH mixtures on the human body, which could lead to either underestimation or overestimation of ILCRs. In addition, the proportion of PAHs absorbed in the body resulting in harmful effects is considered to be 100%; however, Hack and Selenka (1996) identified the influence of food type on the mobilisation of PAHs from soil in the digestive tract. Therefore, the population's diet could also influence the potential risks associated with soil contaminants. Consequently, to better evaluate the risk induced by food ingestion, bioavailability tests should be conducted using the standard diet of the targeted population. Furthermore, as PAHs in French soils were identified as originating from historical contamination, the aging process might reduce PAH bioavailability (Bogan and Sullivan, 2003;  
500 Semple et al., 2003) as well as their potential toxicity to organisms (Duan et al., 2015; Ma et al., 2012). Nevertheless, Reeves et al. (2001) demonstrated the low impact of soil aging on PAH availability and ingestion by organisms (i.e. rats in the study).

Finally, the sequestration of PAHs in the soil could result in the biodegradation of some molecules into metabolites such as fluoranthene-2,3-dione(F23Q), which is known to be toxic, can accumulate in the environment, and  
510 inhibits the degradation of other PAHs (Kazunga et al., 2001). This issue was also reported by Schmidt et al. (2010), who demonstrated that fungal PAH-metabolites tend to accumulate in the soil solution due to their high water solubility and appear to mineralise more slowly than the parent PAHs potentially leaching into the groundwater. These results support the need to study the fate of contaminants in soils and their associated risks.

#### 4 Conclusion

515 This study explored the concentrations of PAHs in 2154 soils collected throughout France based on an extended dataset collected using a systematic sampling grid covering all land-use types. Seventy percent of the soil samples presented PAH concentrations over the LOQ, and 30% of those soils were above the background value of  $100 \mu\text{g kg}^{-1}$  in the  $\Sigma 15\text{PAHs}$ , with a probability of 0.8 to exceed this threshold value in northern France. The spatial distribution of the  $\Sigma 15\text{PAHs}$  throughout the country showed a particular pattern, with higher PAH content in soils  
520 situated in the north-eastern part of France, in the Rhône Valley, and near the main cities. These results

demonstrated large-scale contamination of French soils by PAHs, which was identified by PAH molecular ratios as legacy contamination from the emissions linked to the industrialisation of Europe that started in 1850. The health risk assessment conducted for the residential population showed that 99% of the French soils presented a risk lower than the target value of TLCR of  $10^{-6}$  for the residential population. Only one site exceeded the limit of  $10^{-5}$  chosen by the French government to consider soil as contaminated in residential areas. A higher density of TLCRs between  $10^{-7}$  and  $10^{-6}$  was found in the north-eastern part of France, the Rhône Valley, and in soils surrounding the main cities. These results demonstrate the need to investigate the risk associated with PAHs in soils using a probabilistic method with French population statistics that includes food ingestion as a transfer pattern. Finally, this study demonstrated the heterogeneous risks associated with PAHs in soils over France; highlighted the environmental inequalities with regional variations in population and ecosystem exposure to hazardous contaminants; and indicated the need to conduct large-scale studies to understand contaminant behaviours, their fates in the ecosystem, and their associated risks.

### **Credit Author Statement**

Claire Froger: Conceptualization, Data Curation, Writing – Original Draft. N. P. A. Saby: Conceptualization, Supervision, Methodology, Data Curation, Formal Analysis, Writing – Review & Editing. C. C. Jolivet: Supervision, Resources, Writing – Review & Editing. L. Boulonne: Supervision, Resources, Writing – Review & Editing. G. Caria: Resources, Methodology, Validation, Writing – Review & Editing. X. Freulon: Software, Formal Analysis, Validation, Writing – Review & Editing. C. de Fouquet: Software, Formal Analysis, Validation, Writing – Review & Editing. H. Roussel: Methodology, Validation, Writing – Review & Editing. F. Marot: Methodology, Validation, Writing – Review & Editing. A. Bispo: Supervision, Writing – Review & Editing.

### **Competing interests**

The authors declare that they have no conflict of interest.

### **Acknowledgements**

The soil sampling and the analyses of physicochemical properties of soils were supported by the French Scientific Group of Interest on soils: the “GIS Sol,” involving the French Ministry in charge of the Ecological Transition (MTE), the French Ministry in charge of the Agriculture and Food (MAA), the French Agency for Environment and Energy Management (ADEME), the French Biodiversity Agency (OFB), The National Institute of Geographic and Forest Information (IGN), the French National Research Institute for Sustainable Development and the National Institute for Agronomic and Environmental Research (INRAE). We thank all the soil surveyors and technical assistants involved in sampling the sites. The PAH analyses were supported by a grant from the French Observatory for the Pesticides Residues (ORP).

## References

- ADEME: Guide pour la détermination des valeurs de fonds dans les sols – Echelle d'un site., 2018.
- Aichner, B., Bussian, B., Lehnik-Habrink, P. and Hein, S.: Levels and spatial distribution of persistent organic pollutants in the environment: A case study of German forest soils, *Environ. Sci. Technol.*, 47(22), 12703–12714, doi:10.1021/es4019833, 2013.
- 560 Albanese, S., Fontaine, B., Chen, W., Lima, A., Cannatelli, C., Piccolo, A., Qi, S., Wang, M. and De Vivo, B.: Polycyclic aromatic hydrocarbons in the soils of a densely populated region and associated human health risks: the Campania Plain (Southern Italy) case study, *Environ. Geochem. Health*, 37(1), 1–20, doi:10.1007/s10653-014-9626-3, 2015.
- 565 Anyanwu, I. N. and Semple, K. T.: Fate and behaviour of nitrogen-containing polycyclic aromatic hydrocarbons in soil, *Environ. Technol. Innov.*, 3, 108–120, doi:10.1016/j.eti.2015.02.006, 2015.
- Bertrand, O., Mondamert, L., Grosbois, C., Dhivert, E., Bourrain, X., Labanowski, J. and Desmet, M.: Storage and source of polycyclic aromatic hydrocarbons in sediments downstream of a major coal district in France, *Environ. Pollut.*, 207, 329–340, doi:10.1016/j.envpol.2015.09.028, 2015.
- 570 Biache, C., Ghislain, T., Faure, P. and Mansuy-Huault, L.: Low temperature oxidation of a coking plant soil organic matter and its major constituents: An experimental approach to simulate a long term evolution, *J. Hazard. Mater.*, 188(1–3), 221–230, doi:10.1016/j.jhazmat.2011.01.102, 2011.
- Biache, C., Mansuy-Huault, L. and Faure, P.: Impact of oxidation and biodegradation on the most commonly used polycyclic aromatic hydrocarbon (PAH) diagnostic ratios: Implications for the source identifications, *J. Hazard. Mater.*, 267, 31–39, doi:10.1016/j.jhazmat.2013.12.036, 2014.
- 575 Bogan, B. W. and Sullivan, W. R.: Physicochemical soil parameters affecting sequestration and mycobacterial biodegradation of polycyclic aromatic hydrocarbons in soil, *Chemosphere*, 52(10), 1717–1726, doi:10.1016/S0045-6535(03)00455-7, 2003.
- Brändli, R. C., Bucheli, T. D., Ammann, S., Desaulles, A., Keller, A., Blum, F. and Stahel, W. A.: Critical evaluation of PAH source apportionment tools using data from the Swiss soil monitoring network, *J. Environ. Monit.*, 10(11), 1278–1286, doi:10.1039/b807319h, 2008.
- 580 Bruce, E. D., Abusalih, A. A., McDonald, T. J. and Autenrieth, R. L.: Comparing deterministic and probabilistic risk assessments for sites contaminated by polycyclic aromatic hydrocarbons (PAHs), *J. Environ. Sci. Heal. Part A*, 42(6), 697–706, doi:10.1080/10934520701304328, 2007.
- 585 Brus, D. J., Lamé, F. P. J. and Nieuwenhuis, R. H.: National baseline survey of soil quality in the Netherlands, *Environ. Pollut.*, doi:10.1016/j.envpol.2009.02.028, 2009.
- Bull, S. and Collins, C.: Promoting the use of BaP as a marker for PAH exposure in UK soils, *Environ. Geochem. Health*, 35(1), 101–109, doi:10.1007/s10653-012-9462-2, 2013.
- 590 Cachada, A., Pato, P., Rocha-Santos, T., da Silva, E. F. and Duarte, A. C.: Levels, sources and potential human health risks of organic pollutants in urban soils, *Sci. Total Environ.*, 430, 184–192, doi:10.1016/j.scitotenv.2012.04.075, 2012a.
- Cachada, A., Pereira, M. E., Ferreira da Silva, E. and Duarte, A. C.: Sources of potentially toxic elements and organic pollutants in an urban area subjected to an industrial impact, *Environ. Monit. Assess.*, 184(1), 15–32, doi:10.1007/s10661-011-1943-8, 2012b.
- 595 Cachada, A., Ferreira da Silva, E., Duarte, A. C. and Pereira, R.: Risk assessment of urban soils contamination:

- The particular case of polycyclic aromatic hydrocarbons, *Sci. Total Environ.*, 551–552, 271–284, doi:10.1016/j.scitotenv.2016.02.012, 2016.
- Canadian Council of Ministers of the Environment: Canadian Soil Quality Guidelines: Carcinogenic and Other PAHs (Environmental and Human Health Effects). Scientific Criteria Document (revised). [online] Available from: <http://www.ec.gc.ca/ceqg-rcqe>, 2010.
- 600 Carre, F., Caudeville, J., Bonnard, R., Bert, V., Boucard, P. and Ramel, M.: Soil Contamination and Human Health: A Major Challenge for Global Soil Security, in *GLOBAL SOIL SECURITY*, edited by Field, DJ and Morgan, CLS and McBratney, AB, pp. 275–295, SPRINGER INT PUBLISHING AG, GEWERBESTRASSE 11, CHAM, CH-6330, SWITZERLAND., 2017.
- 605 Chilès, J.-P. and Delfiner, P.: *Geostatistics: Modeling Spatial Uncertainty*, Second Edition, John Wiley & Sons, Inc., Hoboken, NJ, USA., 2012.
- Cui, S., Zhang, Z., Fu, Q., Hough, R., Yates, K., Osprey, M., Yakowa, G. and Coull, M.: Long-term spatial and temporal patterns of polycyclic aromatic hydrocarbons (PAHs) in Scottish soils over 20 years (1990–2009): A national picture, *Geoderma*, doi:10.1016/j.geoderma.2019.114135, 2020.
- 610 Desaules, A., Ammann, S., Blum, F., Brändli, R. and Bucheli, T. D.: Teneurs en HAP et en PCB dans les sols en Suisse. Résultats de l’Observatoire national des sols 1995/1999, *Obs. Natl. des sols*, 2010.
- Duan, L., Naidu, R., Thavamani, P., Meaklim, J. and Megharaj, M.: Managing long-term polycyclic aromatic hydrocarbon contaminated soils: a risk-based approach, *Environ. Sci. Pollut. Res.*, 22(12), 8927–8941, doi:10.1007/s11356-013-2270-0, 2015.
- 615 Elmquist, M., Zencak, Z. and Gustafsson, Ö.: A 700 year sediment record of black carbon and polycyclic aromatic hydrocarbons near the EMEP air monitoring station in Aspövreten, Sweden, *Environ. Sci. Technol.*, 41(20), 6926–6932, doi:10.1021/es070546m, 2007.
- Emery, X.: Ordinary multigaussian kriging for mapping conditional probabilities of soil properties, *Geoderma*, 132(1–2), 75–88, doi:10.1016/j.geoderma.2005.04.019, 2006.
- 620 Fernández, P., Vilanova, R. M., Martínez, C., Appleby, P. and Grimalt, J. O.: The Historical Record of Atmospheric Pyrolytic Pollution over Europe Registered in the Sedimentary PAH from Remote Mountain Lakes, *Environ. Sci. Technol.*, 34(10), 1906–1913, doi:10.1021/es9912271, 2000.
- Fouedjio, F., Desassis, N. and Rivoirard, J.: A generalized convolution model and estimation for non-stationary random functions, *Spat. Stat.*, 16, 35–52, doi:10.1016/j.spasta.2016.01.002, 2016.
- 625 Froger, C., Quantin, C., Gasperi, J., Caupos, E., Monvoisin, G., Evrard, O. and Ayrault, S.: Impact of urban pressure on the spatial and temporal dynamics of PAH fluxes in an urban tributary of the Seine River (France), *Chemosphere*, 219, 1002–1013, doi:10.1016/j.chemosphere.2018.12.088, 2019.
- Gabrieli, J., Vallelonga, P., Cozzi, G., Gabrielli, P., Gambaro, A., Sigl, M., Decet, F., Schwikowski, M., Gäggeler, H., Boutron, C., Cescon, P. and Barbante, C.: Post 17th-century changes of european pah emissions recorded in high-altitude alpine snow and ice, *Environ. Sci. Technol.*, 44(9), 3260–3266, doi:10.1021/es903365s, 2010.
- 630 Gaspéri, J., Ayrault, S., Moreau-Guigon, E., Alliot, F., Labadie, P., Budzinski, H., Blanchard, M., Muresan, B., Caupos, E., Cladière, M., Gateuille, D., Tassin, B., Bordier, L., Teil, M.-J., Bourges, C., Desportes, A., Chevreuil, M. and Moilleron, R.: Contamination of soils by metals and organic micropollutants: case study of the Parisian conurbation, *Environ. Sci. Pollut. Res.*, 25(24), 23559–23573, doi:10.1007/s11356-016-8005-2, 2018a.
- 635 Gaspéri, J., Ayrault, S., Moreau-Guigon, E., Alliot, F., Labadie, P., Budzinski, H., Blanchard, M., Muresan, B.,

- Caupos, E., Cladière, M., Gateuille, D., Tassin, B., Bordier, L., Teil, M. J., Bourges, C., Desportes, A., Chevreuil, M. and Moilleron, R.: Contamination of soils by metals and organic micropollutants: case study of the Parisian conurbation, *Environ. Sci. Pollut. Res.*, 25(24), 23559–23573, doi:10.1007/s11356-016-8005-2, 2018b.
- Gateuille, D., Evrard, O., Lefevre, I., Moreau-Guigon, E., Alliot, F., Chevreuil, M. and Mouchel, J. M.: Combining measurements and modelling to quantify the contribution of atmospheric fallout, local industry and road traffic to PAH stocks in contrasting catchments, *Environ. Pollut.*, 189, 152–160, doi:10.1016/j.envpol.2014.02.029, 2014a.
- Gateuille, D., Evrard, O., Lefevre, I., Moreau-Guigon, E., Alliot, F., Chevreuil, M. and Mouchel, J.-M.: Mass balance and decontamination times of Polycyclic Aromatic Hydrocarbons in rural nested catchments of an early industrialized region (Seine River basin, France), *Sci. Total Environ.*, 470–471, 608–617, doi:10.1016/j.scitotenv.2013.10.009, 2014b.
- Gateuille, D., Evrard, O., Lefevre, I., Moreau-Guigon, E., Alliot, F., Chevreuil, M. and Mouchel, J.-M.: Mass balance and decontamination times of Polycyclic Aromatic Hydrocarbons in rural nested catchments of an early industrialized region (Seine River basin, France), *Sci. Total Environ.*, 470–471, 608–617, doi:10.1016/j.scitotenv.2013.10.009, 2014c.
- Gateuille, D., Gasperi, J., Briand, C., Guigon, E., Alliot, F., Blanchard, M., Teil, M.-J., Chevreuil, M., Rocher, V., Azimi, S., Thevenot, D., Moilleron, R., Brignon, J.-M., Meybeck, M. and Mouchel, J.-M.: Mass Balance of PAHs at the Scale of the Seine River Basin, in *The Seine River Basin*, pp. 163–187., 2020.
- Gocht, T., Ligouis, B., Hinderer, M. and Grathwohl, P.: ACCUMULATION OF POLYCYCLIC AROMATIC HYDROCARBONS IN RURAL SOILS BASED ON MASS BALANCES AT THE CATCHMENT SCALE, *Environ. Toxicol. Chem.*, 26(4), 591, doi:10.1897/06-287R.1, 2007.
- Grimmer, G.: PAH—Their contribution to the carcinogenicity of various emissions†, *Toxicol. Environ. Chem.*, 10(3), 171–181, doi:10.1080/02772248509357101, 1985.
- Gubler, A., Wächter, D., Blum, F. and Bucheli, T. D.: Remarkably constant PAH concentrations in Swiss soils over the last 30 years, *Environ. Sci. Process. Impacts*, 17(10), 1816–1828, doi:10.1039/c5em00344j, 2015.
- Hack, A. and Selenka, F.: Mobilization of PAH and PCB from contaminated soil using a digestive tract model, *Toxicol. Lett.*, 88(1–3), 199–210, doi:10.1016/0378-4274(96)03738-1, 1996.
- Harrison, R. M., Smith, D. J. T. and Luhana, L.: Source apportionment of atmospheric polycyclic aromatic hydrocarbons collected from an urban location in Birmingham, U.K, *Environ. Sci. Technol.*, 30(3), 825–832 ST-Source apportionment of atmospheric, 1996.
- Heywood, E., Wright, J., Wienburg, C. L., Black, H. I. J., Long, S. M., Osborn, D. and Spurgeon, D. J.: Factors influencing the national distribution of polycyclic aromatic hydrocarbons and polychlorinated biphenyls in British soils, *Environ. Sci. Technol.*, 40(24), 7629–7635, doi:10.1021/es061296x, 2006.
- IRIS, NCEA and U.S. EPA: Toxicological Review of Benzo(a)pyrene Executive summary., 2017.
- Jensen, H., Reimann, C., Finne, T. E., Ottesen, R. T. and Arnoldussen, A.: PAH-concentrations and compositions in the top 2 cm of forest soils along a 120 km long transect through agricultural areas, forests and the city of Oslo, Norway, *Environ. Pollut.*, 145(3), 829–838, doi:10.1016/j.envpol.2006.05.008, 2007.
- Katsoyiannis, A. and Breivik, K.: Model-based evaluation of the use of polycyclic aromatic hydrocarbons molecular diagnostic ratios as a source identification tool, *Environ. Pollut.*, 184, 488–494, doi:10.1016/j.envpol.2013.09.028, 2014.
- Kavouras, I. G., Koutrakis, P., Tsapakis, M., Lagoudaki, E., Stephanou, E. G., Von Baer, D. and Oyola, P.: Source



- Apportionment of Urban Particulate Aliphatic and Polynuclear Aromatic Hydrocarbons (PAHs) Using Multivariate Methods, *Environ. Sci. Technol.*, 35(11), 2288–2294, doi:10.1021/es001540z, 2001.
- Kazunga, C., Aitken, M. D., Gold, A. and Sangaiah, R.: Fluoranthene-2,3- and -1,5-diones are novel products from the bacterial transformation of fluoranthene, *Environ. Sci. Technol.*, 35(5), 917–922, doi:10.1021/es001605y, 680 2001.
- Keyte, I. J., Albinet, A., Harrison, R. M. and Harrison, R. M.: On-road traffic emissions of polycyclic aromatic hydrocarbons and their oxy- and nitro- derivative compounds measured in road tunnel environments, *Sci. Total Environ.*, 566–567, 1131–1142, doi:10.1016/j.scitotenv.2016.05.152, 2016.
- Killops, S. D. and Massoud, M. S.: Polycyclic aromatic hydrocarbons of pyrolytic origin in ancient sediments: 685 evidence for Jurassic vegetation fires, *Org. Geochem.*, 18(1), 1–7, doi:10.1016/0146-6380(92)90137-M, 1992.
- Kim, M., Kennicutt, M. C. and Qian, Y.: Source characterization using compound composition and stable carbon isotope ratio of PAHs in sediments from lakes, harbor, and shipping waterway, *Sci. Total Environ.*, 389(2–3), 367–377, doi:10.1016/j.scitotenv.2007.08.045, 2008.
- Lark, R.: Modelling complex soil properties as contaminated regionalized variables., *Geoderma*, (106), 173–190, 690 2002.
- Larsen, R. K. and Baker, J. E.: Source Apportionment of Polycyclic Aromatic Hydrocarbons in the Urban Atmosphere: A Comparison of Three Methods, *Environ. Sci. Technol.*, 37(9), 1873–1881, doi:10.1021/es0206184, 2003.
- Leorri, E., Mitra, S., Irabien, M. J., Zimmerman, A. R., Blake, W. H. and Cearreta, A.: A 700year record of 695 combustion-derived pollution in northern Spain: Tools to identify the Holocene/Anthropocene transition in coastal environments, *Sci. Total Environ.*, doi:10.1016/j.scitotenv.2013.09.064, 2014.
- Liu, G., Niu, J., Guo, W., An, X. and Zhao, L.: Ecological and health risk-based characterization of agricultural soils contaminated with polycyclic aromatic hydrocarbons in the vicinity of a chemical plant in China, *Chemosphere*, 163, 461–470, doi:10.1016/j.chemosphere.2016.08.056, 2016.
- 700 Lohmann, R., Northcott, G. L. and Jones, K. C.: Assessing the contribution of diffuse domestic burning as a source of PCDD/Fs, PCBs, and PAHs to the U.K. Atmosphere, *Environ. Sci. Technol.*, 34(14), 2892–2899, doi:10.1021/es991183w, 2000.
- Lorgeoux, C., Moilleron, R., Gasperi, J., Ayrault, S., Bonté, P., Lefèvre, I., Tassin, B., Bonté, P., Lefèvre, I. and Tassin, B.: Temporal trends of persistent organic pollutants in dated sediment cores: Chemical fingerprinting of 705 the anthropogenic impacts in the Seine River basin, Paris, *Sci. Total Environ.*, 541, 1355–1363, doi:10.1016/j.scitotenv.2015.09.147, 2016.
- Ma, L., Zhang, J., Han, L., Li, W., Xu, L., Hu, F. and Li, H.: The effects of aging time on the fraction distribution and bioavailability of PAH, *Chemosphere*, 86(10), 1072–1078, doi:10.1016/j.chemosphere.2011.11.065, 2012.
- Ma, W. L., Liu, L. Y., Tian, C. G., Qi, H., Jia, H. L., Song, W. W. and Li, Y. F.: Polycyclic aromatic hydrocarbons 710 in Chinese surface soil: occurrence and distribution, *Environ. Sci. Pollut. Res.*, 22(6), 4190–4200, doi:10.1007/s11356-014-3648-3, 2015.
- Maliszewska-kordybach, B.: PERSISTENT ORGANIC CONTAMINANTS IN THE ENVIRONMENT: PAHs AS A CASE STUDY., *Bioavailab. Org. Xenobiotics Environ.*, (January 1999), 2–36, doi:10.1007/978-94-015-9235-2, 1999.
- 715 Maliszewska-Kordybach, B.: Polycyclic aromatic hydrocarbons in agricultural soils in Poland: preliminary

- proposals for criteria to evaluate the level of soil contamination, *Appl. Geochemistry*, 11(1–2), 121–127, doi:10.1016/0883-2927(95)00076-3, 1996.
- Maliszewska-Kordybach, B., Smreczak, B., Klimkowicz-Pawlas, A. and Terelak, H.: Monitoring of the total content of polycyclic aromatic hydrocarbons (PAHs) in arable soils in Poland, *Chemosphere*, 73(8), 1284–1291, doi:10.1016/j.chemosphere.2008.07.009, 2008.
- MINES ParisTech / ARMINES: RGeostats: The Geostatistical R Package, 2020.
- Morales-Caselles, C., Yunker, M. B. and Ross, P. S.: Identification of Spilled Oil from the MV Marathassa (Vancouver, Canada 2015) Using Alkyl PAH Isomer Ratios, *Arch. Environ. Contam. Toxicol.*, 73(1), 118–130, doi:10.1007/s00244-017-0390-0, 2017.
- Morillo, E., Romero, A. S., Madrid, L., Villaverde, J. and Maqueda, C.: Characterization and sources of PAHs and potentially toxic metals in urban environments of Sevilla (southern Spain), *Water. Air. Soil Pollut.*, 187(1–4), 41–51, doi:10.1007/s11270-007-9495-9, 2008.
- Motelay-Massei, A., Ollivon, D., Garban, B., Teil, M. J., Blanchard, M. and Chevreuil, M.: Distribution and spatial trends of PAHs and PCBs in soils in the Seine River basin, France, *Chemosphere*, 55(4), 555–565, doi:10.1016/j.chemosphere.2003.11.054, 2004.
- Ouvrard, S., Chenot, E.-D., Masfaraud, J.-F. and Schwartz, C.: Long-term assessment of natural attenuation: statistical approach on soils with aged PAH contamination, *Biodegradation*, 24(4), 539–548, doi:10.1007/s10532-013-9618-5, 2013.
- Pacyna, J. M., Breivik, K., Münch, J. and Fudala, J.: European atmospheric emissions of selected persistent organic pollutants, 1970–1995, *Atmos. Environ.*, 37(1), 119–131, doi:10.1016/S1352-2310(03)00240-1, 2003.
- Pereira, M.: Generalized random fields on Riemannian manifolds : theory and practice, PSL Research University. [online] Available from: <https://pastel.archives-ouvertes.fr/tel-02499376>, 2019.
- Petitgas, P., Woillez, M., Rivoirard, J., Renard, D. and Bez, N.: Handbook of geostatistics in R for fisheries and marine ecology. ICES Cooperative Report 338., 2017.
- Petitgas, P., Woillez, M., Doray, M. and Rivoirard, J.: Indicator-Based Geostatistical Models For Mapping Fish Survey Data, *Math. Geosci.*, 50(2), 187–208, doi:10.1007/s11004-018-9725-2, 2018.
- Petrišič, M. G., Muri, G. and Ogrinc, N.: Source identification and sedimentary record of polycyclic aromatic hydrocarbons in lake bled (NW Slovenia) using stable carbon isotopes, *Environ. Sci. Technol.*, 47(3), 1280–1286, doi:10.1021/es303832v, 2013.
- Pies, C., Yang, Y. and Hofmann, T.: Distribution of polycyclic aromatic hydrocarbons (PAHs) in floodplain soils of the Mosel and Saar River, *J. Soils Sediments*, 7(4), 216–222, doi:10.1065/jss2007.06.233, 2007.
- Qi, P., Qu, C., Albanese, S., Lima, A., Cicchella, D., Hope, D., Cerino, P., Pizzolante, A., Zheng, H., Li, J. and De Vivo, B.: Investigation of polycyclic aromatic hydrocarbons in soils from Caserta provincial territory, southern Italy: Spatial distribution, source apportionment, and risk assessment, *J. Hazard. Mater.*, 383(April 2019), 121158, doi:10.1016/j.jhazmat.2019.121158, 2020.
- Ravindra, K., Sokhi, R. and Vangrieken, R.: Atmospheric polycyclic aromatic hydrocarbons: Source attribution, emission factors and regulation, *Atmos. Environ.*, 42(13), 2895–2921, doi:10.1016/j.atmosenv.2007.12.010, 2008.
- Reeves, W. R., McDonald, T. J., Bordelon, N. R., George, S. E. and Donnelly, K. C.: Impacts of aging on in vivo and in vitro measurements of soil-bound polycyclic aromatic hydrocarbon availability, *Environ. Sci. Technol.*, 35(8), 1637–1643, doi:10.1021/es0017110, 2001.

- Rivoirard, J.: Introduction to disjunctive kriging and non-linear geostatistics, Clarendon Press., 1994.
- Rivoirard, J. and Romary, T.: Continuity for Kriging with Moving Neighborhood, *Math. Geosci.*, 43(4), 469–481, doi:10.1007/s11004-011-9330-0, 2011.
- Schmidt, S. N., Christensen, J. H. and Johnsen, A. R.: Fungal PAH-metabolites resist mineralization by soil microorganisms, *Environ. Sci. Technol.*, 44(5), 1677–1682, doi:10.1021/es903415t, 2010.
- 760 Semple, K. T., Morriss, a. W. J. and Paton, G. I.: Bioavailability of hydrophobic organic contaminants in soils: fundamental concepts and techniques for analysis, *Eur. J. Soil Sci.*, 54(December), 809–818, doi:10.1046/j.1365-2389.2003.00564.x, 2003.
- Switzer, P. and Green, A.: Min/max autocorrelation factors for multivariate spatial imagery. Technical report No. 6., 1984.
- 765 Tobiszewski, M. and Namieśnik, J.: PAH diagnostic ratios for the identification of pollution emission sources, *Environ. Pollut.*, 162, 110–119, doi:10.1016/j.envpol.2011.10.025, 2012.
- Tong, R., Yang, X., Su, H., Pan, Y., Zhang, Q., Wang, J. and Long, M.: Levels, sources and probabilistic health risks of polycyclic aromatic hydrocarbons in the agricultural soils from sites neighboring suburban industries in Shanghai, *Sci. Total Environ.*, doi:10.1016/j.scitotenv.2017.10.179, 2018.
- 770 USEPA: Human Health Evaluation Manual Part (B, Development of Risk-based Preliminary Remediation Goals)., 1991.
- USEPA: Soil Screening Guidance : User 's Guide., *Environ. Prot. Agency*, (July), 1–49, doi:EPA/540/R-96/018, 1996.
- 775 USEPA: Risk Assessment Guidance for Superfund (RAGS) Volume III - Part A: Process for Conducting Probabilistic Risk Assessment, Appendix B, *Risk Assess. Guid. Superfund*, III(December), 1–385 [online] Available from: [http://www.epa.gov/sites/production/files/2015-09/documents/rags3adt\\_complete.pdf](http://www.epa.gov/sites/production/files/2015-09/documents/rags3adt_complete.pdf), 2001.
- USEPA: Regional Screening Levels (RSLs), U-S. *Environ. Protection Agency* [online] Available from: <https://www.epa.gov/risk/regional-screening-levels-rsls-users-guide> (Accessed 1 October 2020), 2020.
- 780 Villanneau, E., Saby, N. P. A., Arrouays, D., Jolivet, C. C., Boulonne, L., Caria, G., Barriuso, E., Bispo, A. and Briand, O.: Spatial distribution of lindane in topsoil of Northern France, *Chemosphere*, 77(9), 1249–1255, doi:10.1016/j.chemosphere.2009.08.060, 2009.
- Villanneau, E. J., Saby, N. P. A., Orton, T. G., Jolivet, C. C., Boulonne, L., Caria, G., Barriuso, E., Bispo, A., Briand, O. and Arrouays, D.: First evidence of large-scale PAH trends in French soils, *Environ. Chem. Lett.*, 11(1), 99–104, doi:10.1007/s10311-013-0401-y, 2013a.
- 785 Villanneau, E. J., Saby, N. P. A., Orton, T. G., Jolivet, C. C., Boulonne, L., Caria, G., Barriuso, E., Bispo, A., Briand, O. and Arrouays, D.: First evidence of large-scale PAH trends in French soils, *Environ. Chem. Lett.*, 11(1), 99–104, doi:10.1007/s10311-013-0401-y, 2013b.
- Wang, C., Wu, S., Zhou, S., Wang, H., Li, B., Chen, H., Yu, Y. and Shi, Y.: Polycyclic aromatic hydrocarbons in soils from urban to rural areas in Nanjing: Concentration, source, spatial distribution, and potential human health risk, *Sci. Total Environ.*, 527–528, 375–383, doi:10.1016/j.scitotenv.2015.05.025, 2015.
- 790 Wang, C., Wu, S., Zhou, S., Shi, Y. and Song, J.: Characteristics and Source Identification of Polycyclic Aromatic Hydrocarbons (PAHs) in Urban Soils: A Review, *Pedosphere*, 27(1), 17–26, doi:10.1016/S1002-0160(17)60293-5, 2017a.
- 795 Wang, J., Zhang, X., Ling, W., Liu, R., Liu, J., Kang, F. and Gao, Y.: Contamination and health risk assessment

of PAHs in soils and crops in industrial areas of the Yangtze River Delta region, China, *Chemosphere*, doi:10.1016/j.chemosphere.2016.10.113, 2017b.

Wang, X.-T., Miao, Y., Zhang, Y., Li, Y.-C., Wu, M.-H. and Yu, G.: Polycyclic aromatic hydrocarbons (PAHs) in urban soils of the megacity Shanghai: Occurrence, source apportionment and potential human health risk, *Sci. Total Environ.*, 447, 80–89, doi:10.1016/j.scitotenv.2012.12.086, 2013.

Webster, R. and Oliver, M.: *Geostatistics for Environmental Scientists*, 2nd Editio., 2007.

Wenzel, K. D., Manz, M., Hubert, A. and Schüürmann, G.: Fate of POPs (DDX, HCHs, PCBs) in upper soil layers of pine forests, *Sci. Total Environ.*, 286(1–3), 143–154, doi:10.1016/S0048-9697(01)00972-X, 2002.

Yang, W., Lang, Y. and Li, G.: Cancer risk of polycyclic aromatic hydrocarbons (PAHs) in the soils from Jiaozhou Bay wetland, *Chemosphere*, 112, 289–295, doi:10.1016/j.chemosphere.2014.04.074, 2014.

Yang, Y., Ligouis, B., Pies, C., Achten, C. and Hofmann, T.: Identification of carbonaceous geosorbents for PAHs by organic petrography in river floodplain soils, *Chemosphere*, 71(11), 2158–2167, doi:10.1016/j.chemosphere.2008.01.010, 2008.

Yu, L., Duan, L., Naidu, R. and Semple, K. T.: Abiotic factors controlling bioavailability and bioaccessibility of polycyclic aromatic hydrocarbons in soil: Putting together a bigger picture, *Sci. Total Environ.*, 613–614, 1140–1153, doi:10.1016/j.scitotenv.2017.09.025, 2018.

Yuan, K., Wang, X., Lin, L., Zou, S., Li, Y., Yang, Q. and Luan, T.: Characterizing the parent and alkyl polycyclic aromatic hydrocarbons in the Pearl River Estuary, Daya Bay and northern South China Sea: Influence of riverine input, *Environ. Pollut.*, 199, 66–72, doi:10.1016/j.envpol.2015.01.017, 2015.

Yunker, M. B., Macdonald, R. W., Vingarzan, R., Mitchell, R. H., Goyette, D. and Sylvestre, S.: PAHs in the Fraser River basin: a critical appraisal of PAH ratios as indicators of PAH source and composition, *Org. Geochem.*, 33(4), 489–515, doi:10.1016/S0146-6380(02)00002-5, 2002.

Yunker, M. B., McLaughlin, F. A., Fowler, M. G. and Fowler, B. R.: Source apportionment of the hydrocarbon background in sediment cores from Hecate Strait, a pristine sea on the west coast of British Columbia, Canada, *Org. Geochem.*, 76, 235–258, doi:10.1016/j.orggeochem.2014.08.010, 2014.

Zhang, R., Li, T., Russell, J., Zhang, F., Xiao, X., Cheng, Y., Liu, Z., Guan, M. and Han, Q.: Source apportionment of polycyclic aromatic hydrocarbons in continental shelf of the East China Sea with dual compound-specific isotopes ( $\delta^{13}\text{C}$  and  $\delta^2\text{H}$ ), *Sci. Total Environ.*, 704, 135459, doi:10.1016/j.scitotenv.2019.135459, 2020.

Zhang, X. L., Tao, S., Liu, W. X., Yang, Y., Zuo, Q. and Liu, S. Z.: Source Diagnostics of Polycyclic Aromatic Hydrocarbons Based on Species Ratios: A Multimedia Approach, *Environ. Sci. Technol.*, 39(23), 9109–9114, doi:10.1021/es0513741, 2005.

Zheng, T., Ran, Y. and Chen, L.: Polycyclic aromatic hydrocarbons (PAHs) in rural soils of Dongjiang River Basin: Occurrence, source apportionment, and potential human health risk, *J. Soils Sediments*, 14(1), 110–120, doi:10.1007/s11368-013-0753-8, 2014.

830