

How well does **Predictive-Digital** Soil Mapping represent soil geography?

An investigation from the USA

David G. Rossiter^{1, 2}, Laura Poggio¹, Dylan Beaudette³, and Zamir Libohova⁴

¹ISRIC-World Soil Information, Postbus 353, Wageningen 6700 AJ, NL

²Section of Soil & Crop Sciences, New York State College of Agriculture and Life Sciences, 233 Emerson Hall, Cornell University, Ithaca NY 14853 USA

³USDA-NRCS Soil and Plant Science Division, 19777 Greenley Rd., Sonora, CA, 95370 USA

⁴USDA-ARS, Dale Bumpers Small Farms Research Center, 6883 South State Highway 23, Booneville, AR 72927 USA

Correspondence: David G. Rossiter (david.rossiter@isric.org)

Abstract. We present methods to evaluate the spatial patterns of the geographic distribution of soil properties in the USA, as shown in gridded maps produced by **Predictive-Digital** Soil Mapping (**PSMDSM**) at global (SoilGrids v2), national (Soil Properties and Class 100m Grids of the USA), and regional (POLARIS soil properties) scales, and compare them to spatial patterns known from detailed field surveys (gNATSGO and gSSURGO). The methods are illustrated with an example: topsoil pH for an area in central New York State. A companion report examines other areas, soil properties, and depth sheesintervals. A set of R Markdown scripts is referenced so that readers can apply the analysis for areas of their interest. For the test case we discover and discuss substantial discrepancies between **PSMDSM** products, as well as large differences between the **PSMDSM** products and legacy field surveys. These differences are in whole-map statistics, visually-identifiable landscape features, level of detail, range and strength of spatial autocorrelation, landscape metrics (Shannon diversity and evenness, shape, aggregation, mean fractal dimension, co-occurrence vectors), and spatial patterns of property maps classified by histogram equalization. Histograms and variogram analysis revealed the smoothing effect of machine-learning models. Property class maps made by histogram equalization were substantially different, but there was no consistent trend in their landscape metrics. The model using only national points and covariates was not better-than-substantially different from the global model, and in some cases introduced artefacts from a lithology covariate. Uncertainty (5–95% confidence intervals) provided by SoilGrids and POLARIS were unrealistically wide compared to gNATSGO/gSSURGO low and high estimated values and show substantially different spatial patterns. We discuss the potential use of the **PSMDSM** products as a (partial) replacement for field-based soil surveys. There is no substitute for actually examining and interpreting the soil-landscape relation, but despite the issues revealed in this study, DSM can be an important aid to the soil surveyor.

Contents

20 1 Introduction

3

	2	Data sources	5
	2.1	<u>General character of the products</u>	5
	2.2	Primary data	7
	2.2	Environmental Covariates and geographic scope	9
25	2.2	Mapping methods	9
	2.3	Resolution, depths and coordinate reference systems	10
	2.4	Uncertainty assessment	11
	3	Evaluation methods	12
	3.1	Qualitative methods	12
30	3.2	Numerical methods – whole map	12
	3.3	Numerical methods – spatial continuity	13
	3.4	Numerical methods – patterns	13
	3.4.1	V-measure	13
	3.4.2	Landscape metrics	14
35	3.5	Regional patterns	16
	3.6	Local patterns	17
	3.6.1	Visual method	17
	3.6.2	Quantitative method	17
	4	Example area and soil property	18
40	5	Regional spatial patterns	18
	5.1	<u>Visual method</u>	18
	5.2	Regional maps	19
	5.3	Uncertainty	23
	5.4	Local spatial autocorrelation	29
45	5.5	Classification	30
	5.6	V-measure	30
	5.7	Landscape metrics	33
	6	Local spatial patterns	33
	6.1	Qualitative assessment	34
50	6.2	Quantitative assessment	34
	6.2.1	Class maps	36
	6.2.2	Local spatial autocorrelation	38

55 1 Introduction

~~Predictive Soil Mapping (PSM), also commonly referred to as~~ Digital Soil Mapping (~~DSM~~), has been defined (under the earlier term “Predictive Soil Mapping”) as “the development of a numerical or statistical model of the relationship among environmental variables and soil properties, which is then applied to a geographic data base to create a predictive map” (Scully et al., 2003). Since the seminal paper of McBratney et al. (2003), recently reviewed by Minasny and McBratney (2016), ~~PSM~~

60 ~~DSM~~ has been widely-applied from the field to global levels.

This is in contrast to what we here call “traditional” soil survey, in which the soil surveyor develops a mental model of the soil geography (Hudson, 1992) by interpreting the landscape with the aid of airphotos, purposive transects, and detailed profile descriptions at locations thought to represent the central concepts of the soil classes present in the study area (Soil Survey Division Staff, 20

65 A principal attraction of ~~PSM-DSM~~ is that it produces consistent, geometrically-correct and reproducible gridded maps over large areas, given training data (“point” observations of soil classes, properties or conditions), a set of environmental covariates covering the entire area to be mapped at some fixed grid resolution, and a set of algorithms implemented in computer code. This removes the need for expertise in discovering and interpreting the soil-landscape relations, also known as the “paradigm” of soil survey (Hudson, 1992), which is vital for traditional soil survey and difficult to acquire and harmonize among surveyors.

70 However, expertise in soil-landscape relations is still needed to ensure that DSM outputs are reasonable, and to discover reasons for any discrepancies.

Further, it may be that fewer locations can be visited in order to develop reliable models, as compared to traditional survey techniques. If the relation with covariates is strong, and locations representative of the entire covariate feature space are included in the training set, ~~large areas can be mapped~~ it may be possible to map large areas from relatively few field observations. This corresponds to the “homosoil” concept (Mallavan et al., 2010): identical environmental conditions (as represented by covariates) should result in the same soils. Maps made by ~~PSM-DSM~~ can include areas that are not accessible to field mappers because of permissions or difficult access, if the available training data cover the covariate space of the inaccessible area. However, ~~PSM-DSM~~ requires sufficient sampling density to cover the full covariate space, since most ~~PSM-DSM~~ methods do not ~~extrapolate~~ interpolate or extrapolate in soil property space, and in any case ~~extrapolation is inadvisable~~ it is inadvisable

80 to predict “too far” from the coverage of the training observations. This has been studied by Meyer and Pebesma, who have developed a method for measuring the distance in both covariate (Meyer and Pebesma, 2020) and geographic (Meyer and Pebesma, 2022) space between prediction locations and the set of training points.

~~It is not claimed that PSM is inherently superior to traditional survey, but~~ DSM avoids some well-known problems of traditional survey ~~are avoided~~: multiple survey projects over time with inconsistent standards and mapping concepts, inconsistency

85 among mappers, difficulties in objectively identifying boundaries, and indeed the need to identify boundaries. However, tradi-

tional soil surveyors and users of their maps are often critical of ~~PSM-DSM~~ products, and may not understand how they were made and how they should be used (Arrouays et al., 2020). In the USA there is increasing awareness of, and interest in, ~~PSM-DSM~~ products. Here the most important point of contention has to do with ~~PSM-DSM~~ resolution (pixel size), which implies a mapping scale, compared to the scale at which differences can be reliably interpreted for user needs. Criticism of ~~PSM-DSM~~ products is proportional to the degree to which their implied spatial precision and accuracy is over-sold.

Another benefit of DSM methods is the quantification of uncertainty inherent in various geostatistical and machine-learning approaches (Sztamári and Pásztor, 2018). In traditional mapping, uncertainty is implicitly encoded via mapping scale (which determines the size of the minimum delineation), map unit purity specification (e.g., complex, association, consociation), and taxonomic precision (e.g., soil series vs. suborder) (Soil Survey Division Staff, 2017).

95 The success of ~~PSM-DSM~~ in reproducing known “point” observations (i.e., pedons described in the field and characterized in the laboratory) is typically reported by evaluation (“validation”) statistics based on data splitting (~~independent-evaluation~~) or by cross-validation. These evaluations are almost never based on random sampling (Brus et al., 2011), and since the source point datasets are almost always biased towards certain land uses, access constraints or landscape locations, these evaluations carry forward these biases and must be interpreted with caution.

100 A more serious issue is that point evaluations of ~~PSM-DSM~~ products do not consider the spatial pattern of predictions. By contrast, ~~soil-surveys-based-on-extensive-field-observation-of-the-soil-landscape-(sometimes-called-the-“traditional”-soil-survey-method)~~ traditional soil surveys produce polygon maps of ~~soil map units (SMU) composed of one or more soil type units (STU), e.g., soil series, which then provide modal or representative profiles with measurements or estimates of soil properties, usually by genetic horizon. These~~ relatively homogeneous soil bodies (represented as soil map units), with the boundary lines placed at inflection points of maximum change between them (Lagacherie et al., 1996). These maps explicitly show the ~~soil-landscape, as interpreted by the soil surveyor and surveyor’s interpretation of the soil landscape as managed by the land user~~ developed from a mental model of the soil-forming processes, and which when viewed as a whole show the pattern of the soil cover. It has long been recognized that the soil cover forms patterns at various scales (Fridland, 1974; Hole and Campbell, 1985), so that the traditional soil mapper attempts to find those patterns expressed at the map design scale. Since ~~PSM-maps-are-not derived from a paradigm such as the surveyor’s explicit stratification of the soil landscape, their spatial pattern depends on the input data (training points and covariates) and the PSM algorithm applied~~ DSM predictions are on a grid cell basis, most DSM models have no concept of relatively homogeneous natural soil bodies nor inflection points between them. However, it might be expected that if the values of the DSM covariates representing the soil-forming factors also cluster in a similar pattern to the soil cover, the DSM predictions would also cluster and approximate map units from traditional survey. Convolutional

105 neural networks (e.g., Taghizadeh-Mehrjardi et al., 2020), not represented in the methods compared in this paper, explicitly consider neighbourhoods of various size, but not explicitly connectivity. The question is thus to what degree ~~PSM-DSM~~ products represent the actual soil landscape spatial pattern and, more importantly, the underlying pedogenetic and geomorphic processes.

120 ~~PSM-DSM~~ maps are most commonly produced at grid cell resolutions from 1 km to 30 m, and even to ~~10~~ < 10 m for precision agriculture applications. Environmental covariates are available at these resolution, so that ~~PSM-DSM~~ products at

high resolutions can show fine details that can not be presented at the design scale of ~~conventional maps~~ polygon maps made by traditional methods. These have minimum legible delineations (MLD) of 0.25 cm² (Vink, 1975) or 0.40 cm² (Forbes et al., 1982) on the published map, multiplied by the scale factor. For example, a polygon map at 1:24 000, typical of USA ~~conventional~~ traditional soil survey, can represent spatial patterns of 1.44 (Vink) to 2.3 (Forbes) ha minimum-size polygons.

125 The Forbes et al. (1982) criteria have been incorporated into NRCS soil survey standards (Schoeneberger et al., 2012; Soil Survey Division
These correspond to single grid cell resolutions of 240 to 384 m, coarser than higher-resolution PSM-DSM products from
(30 to 100 m). But the question remains whether this implied fine detail represents true differences or artefacts of the mapping
process – in other words, should the PSM-DSM map unit trust the apparent differences between adjacent grid cells, or are some
or most of these differences due to artefacts (“noise”) of the PSM-DSM process? Further, there is the question of how well the
130 medium-resolution products (e.g., 250 m) represent the soil landscape at regional extent.

The objective of this study is to present methods with which to evaluate the landscape and detailed level spatial patterns of ~~PSM maps~~ These maps may ~~DSM maps~~ These maps have been developed for global, national, or regional spatial extents. These patterns are compared with digital soil maps based on polygon maps produced by ~~field survey~~ traditional soil survey, using field study and expert soil-landscape analysis. We chose the USA as a study area because of the availability of field-based

135 soil surveys at 1:12 000 to 1:24 000 design scale, linked to detailed descriptions of modal soil profiles, available as a seamless
digital product. These comparisons may be useful in the context of current plans (Thompson et al., 2020) for updating and
completing the USA soil survey using PSM-DSM methods and GlobalSoilMap (GSM) specifications (Arrouays et al., 2014).
They should also be useful for developing realistic expectations for what PSM-DSM can and cannot deliver (Arrouays et al.,
2020).

140 To evaluate ~~PSM-DSM~~ methods we apply them to selected test areas and soil properties, and comment on the results. This paper introduces the methods and data sources, and includes an illustrative example (one area, one soil property, one depth ~~slice~~), with a minimum of discussion of soil geography interval), in the context of the soil geography of the selected region. A companion ISRIC Report (Rossiter et al., 2021) presents four case studies in diverse soil geographic contexts, each with different soil properties and depth ~~slices~~ intervals. We encourage readers to apply the methods to their own study areas within

145 the USA and to their soil properties of interest, to evaluate the utility of the several PSM-DSM products. For this, we provide
our analysis scripts as R Markdown documents (R Studio, 2020); see “Code availability” at the end of this paper.

2 ~~Data sources~~ Products compared

The ~~source maps~~ products compared in this study differ in their primary data source (soil maps and point observations), their
geographic scope, the mapping methods used to make the digital product, their resolution, depths and coordinate reference
150 systems, and how they assess and present uncertainty. We summarize these below; see the journal articles describing each
source for details.

2.1 General character of the products

The products are of three kinds: (1) digital products based on field-traditional soil survey without any statistical modelling; (2) PSM-DSM products based on field-traditional soil survey products and enhanced by statistical modelling using environmental covariates; and (3) PSM-DSM products based on statistical modelling using training points and environmental covariates. This latter is the most common PSM-DSM method worldwide, especially for areas without extensive field-traditional soil surveys. ~~These differ in their primary data, their environmental covariates and geographic scope, their mapping methods, the resolution of their predictions, and their uncertainty assessment. These have implications for interpreting their relative success. We summarize these below; see the Supplementary Materials and corresponding papers for details.~~

160 The first ~~source-kind of product~~ is represented by the reference ~~product-products~~ from the Natural Resources Conservation Service (NRCS) of the United States Department of Agriculture (USDA), based on extensive field survey ~~and~~, airphoto interpretation, thematic maps, and expert evaluation of DEM-derivatives. This is considered to be the most accurate information, despite the occasional presence of artefacts from the overall mapping program-programme, as explained later in this section. There are two closely-related products from the NRCS.

165 At the national level, The National Soil Geographic Database gNATSGO (NRCS Soils, 2022b) is a composite of the Soil Survey Geographic Database (SSURGO, mostly 1:24,000 scale), and State Soil Geographic Database (STATSGO2, 1:250,000 scale), and the ~~very~~ detailed Raster Soil Survey Database (RSS), according to the most detailed product available for all areas of the USA (?). ~~For each State It is aimed at users who require multi-state or CONUS extent mapping. For each state~~ or equivalent political unit, the SSURGO ~~and-or~~ STATSGO2 polygon maps of SMU produced by field-traditional survey have been rasterized to a grid, each cell keyed to a ~~a~~ soil map unit (SMU) (NRCS Soils, 2020a). Grid cells link to the best available (i.e., greatest detail STATSGO-SSURGO-RSS) SMU. The digital products are delivered at 30 ~~m-resolution and 90 m resolutions~~ for the 48 ~~States of the continental USA and the contiguous states and the federal~~ District of Columbia of the USA (abbreviated CONUS); ~~i. e., excluding Alaska, Hawaii and the Island Territories.~~ ~~The SMU are mappable landscape elements, at the survey design scale. These usually have multiple component STU, with reported estimated proportion and sometimes the landscape relations within the SMU. However the locations of the STU within the SMU are not mapped due to the design scale. The STU are linked to database tables of representative or synthetic soil profiles, with field and laboratory measurements of multiple soil properties, as well as interpretations for soil use. To obtain values for soil properties in a grid cell, properties of the components of the corresponding SMU are combined by area-weighted averaging. To obtain values at coarser resolution, grid cells are averaged by upscaling.~~

175

180 The second source is represented by **POLARIS soil properties** (Chaney et al., 2019) (further **PSP**), the result of harmonizing diverse SSURGO and STATSGO2 polygon data with the DSMART algorithm (Odgers et al., 2014) to produce a probabilistic raster soil class or component map ~~(At the state level, gSSURGO is also available (NRCS Soils, 2022a). This has higher resolution (both 10 m and 30 grid resolution) and then extracting property information from gSSURGO, aggregated by component name. Despite the source data, this is not an NRCS product.~~

185 There are two products representing the third source, one for the world and one for the continental USA only. This allows us ~~to compare globally~~ m) to minimize degradation of the original polygon delineations, and nationally-consistent products. The global product is SoilGrids v2.0 (further SG2) (ISRIC – World Soil Information, 2020; Poggio et al., 2021), a further

development of SoilGrids1km (Hengl et al., 2014) and SoilGrids250m (Hengl et al., 2017) is a direct gridding of SSURGO polygons. This uses a global point dataset and environmental covariates that cover the entire world (except the high Arctic and Antarctica), and models that are globally consistent. It does not use STATSGO2 for infilling nor RSS if available. The continental product is It thus is a gridded version of the Soil Properties and Class 100m Grids of the United States (further SPCG) (Ramcharan et al., 2018), which followed the methodology of Hengl et al. (2017) with the addition of USA-specific covariates, notably parent material and drainage classes extracted from SSURGO or STATSGO2, and only used the CONUS extent of environmental covariates in model building familiar SSURGO product that is used for local applications.

195 2.2 Primary data

A major difference between products is the extent to which primary data relies on field soil survey. SSURGO is refreshed annually for those users who do not wish to mix STATSGO or the new raster soil surveys into their analysis.

At one extreme are the hand-drawn. The gridded gNATSGO and gSSURGO maps are derived from the polygons of SSURGO, the basis of the gridded products gSSURGO and gNATSGO. These polygons are available from the NRCS as vector GIS layers (Natural Resources Conservation Service, 2019), and in a convenient format on a geographic background as SoilWeb (California Soil Resource Lab, 2020). These are a a representation of those delineated by the field surveyors on stereo-pairs or ortho-photos and subsequently converted to vector digital format by manual digitization. Soil surveys conducted in the last 15 years were compiled using on-screen digitization in a GIS. At boundaries between survey areas, polygon lines at survey limits have been matched during digitizing (D'Avelo and McLeese, 1998). These polygons are organized in soil map units (SMU) with one or more components (soil taxonomic units, STU), usually named for a soil series but more specific than the parent soil series concept. Taxa above the soil series (family or subgroup) are commonly used in soil surveys of national forest land or wilderness areas. Soil series are the lowest level of Soil Taxonomy (Soil Survey Division Staff, 2014) and are described in the Official Series Descriptions (OSD), as modal profiles with a set of ranges for the observed morphology and laboratory measurements.

210 The component STU in a mapped SMU vary in the observed field properties from the OSD modal description, but usually fit within soil series range. The observed field properties of soil component units are utilized for developing a set of interpretations for SSURGO polygon map units. Further, since field. These polygons are available from the NRCS as vector GIS layers (Natural Resources Conservation Service, 2019), and in a convenient format on a geographic background as SoilWeb (California Soil Resource Lab, 2020).

215 The SMU of the source maps are mappable landscape elements, at the survey design scale. These almost always have multiple component STU, with reported estimated proportion and geomorphic arrangement within the SMU when possible. However the locations of the STU within the SMU are not mapped due to the design scale. The STU are linked to database tables of representative or synthetic soil profiles, with field and laboratory measurements of multiple soil properties, as well as interpretations for soil use. To obtain values for soil properties in a gNATSGO or gSSURGO grid cell, properties of the components of the corresponding SMU are combined by area-weighted averaging. To obtain values at coarser resolutions, weighted-average properties of groups of grid cells are upscaled by averaging.

There are inherent problems with this product. First, since traditional surveys were carried out over a long time period, series names and mapping concepts may differ between adjacent survey areas (a. k.a. survey “vintage”). ~~At boundaries between survey areas, polygon lines at survey limits have been matched during digitizing (D’Avelo and McLeese, 1998).~~ Thus SSURGO SMU delineations and linked tabular data represent a progressive data collection and correlation effort spanning nearly 100 years. ~~As such~~ Therefore, there exist many soil survey vintages, each a snapshot in time, tied to specific land-use assumptions and technological limitations. Systematic, continuous updates to the entire SSURGO database have been made since 2013.

~~However~~ 2013 and are ongoing. Second, the transfer from unrectified photos to topographic base and the edge matching between survey areas has not always been flawless, and in addition polygons may have been mis-drawn on the original survey (Supplementary Fig. 4S1). Thus we can not take ~~gSSURGO~~ these primary polygon maps as a completely reliable ~~reference~~ georeference.

~~PSP does not use any point observations; rather, it samples pseudo-points from gSSURGO and uses these as training points for the DSMART disaggregation algorithm (see below).~~ The second kind of product is represented by POLARIS soil properties (Chaney et al., 2019) (further **PSP**), the result of harmonizing diverse SSURGO and STATSGO2 polygon data with the DSMART algorithm (Odgers et al., 2014) to produce a probabilistic raster soil class or component map (30 m grid resolution) and then extracting property information from gSSURGO or gNATSGO grid cells (representing polygons), aggregated by component name. Despite the source data, this is not an NRCS product and was developed independently of the NRCS.

~~At the other extreme is SG2, which~~ There are two products representing the third kind of product, one for the world and one for the continental USA only. This allows us to compare globally- and nationally-consistent products. The global product is SoilGrids v2.0 (further SG2) (ISRIC - World Soil Information, 2020; Poggio et al., 2021), a further development of SoilGrids1km (Hengl et al., 2014) and SoilGrids250m (Hengl et al., 2017). This uses a global point dataset and environmental covariates that cover the entire world (except the high Arctic and Antarctica), and global models. It does not use any information derived from SSURGO or STATSGO map units. Its training points are extracted from the freely-shareable World Soil Information Service (WoSIS) point dataset from ISRIC-World Soil Information (Batjes et al., 2020). These include all profiles in the ~~NRCS pedon~~ NCSS Laboratory Characterization Database database. The freely-sharable WoSIS points are augmented by several datasets included in WoSIS that can not be published externally due to restrictions by the original data providers to ISRIC, but which can be used in mapping. In total $\approx 240\ 000$ profiles were used in model building.

The continental product is the Soil Properties and Class 100m Grids of the United States (further SPCG) (Ramcharan et al., 2018), which followed the methodology of Hengl et al. (2017) with the addition of USA-specific covariates, notably parent material and drainage classes extracted from SSURGO or STATSGO2 map units, and only used the CONUS extent of environmental covariates in model building. SPCG is similar to SG2 in that it is primarily based on point observations, but it has a richer source of these than SG2: the NCSS Laboratory Characterization Database (34 183 pedons comprising 213 499 horizons), the National Soil Information System (NASIS), and the Rapid Carbon Assessment (RaCA) dataset (31 215 pedons); this latter only for organic C, total N, and bulk density.

2.2 Environmental Covariates and geographic scope

The three PSM products (SG2, PSP, and SPCG) use a large number of gridded GIS coverages as environmental covariates in their predictive models. These represent soil-forming factors, and include climate, ecology, geology, land use/cover, terrain, vegetation and hydrography; see Supplementary Information for details.

260 PSP also uses coarse-resolution (≈ 2) estimates of U, Th, and K γ -ray decay products. The model is trained in overlapping tiles, thus each tile is derived from a local model.

SG2 only uses environmental covariates available over the whole world. The model is trained for the whole world, not per-country or region, thus it is a global model. This corresponds to the “homosoil” concept (Mallavan et al., 2010): identical environmental conditions anywhere in the world should result in the same soils. The obvious question is whether or not the
265 additional information from outside the CONUS leads to an improved model for this region.

SPCG also uses SSURGO and STATSGO2 polygons. It also uses SSURGO map units to derive parent material (87) and drainage (4) classes as environmental-CONUS-specific covariates. The model is trained for the CONUS, thus it is a regional model.

2.2 Mapping methods

270 ~~gSSURGO is based on field~~ gNATSGO and gSSURGO are based on traditional soil survey, mostly on unrectified airphoto bases until the late 1990’s. The many individual survey areas prior to this time have been partially homogenized during a process of digitization and recompilation onto topographic or orthophoto bases during the 1990’s (D’Avelo and McLeese, 1998) and are provided as the polygon SSURGO map. In the early 2000’s for new surveys and updates a transition was made to on-screen digitization over orthophotos. Field methods are described in successive editions of the Soil Survey Manual (Soil Survey
275 Division Staff, 2017) and the field book for describing and sampling soils (Schoeneberger et al., 2012). Mapping is based on conceptual models of soil-landscape relations developed in each survey area (Hudson, 1992), confirmed by purposive auger
~~observations and a small number of~~ and full profile descriptions to characterize map unit composition. Component concepts are refined with any available laboratory characterization data, with (limited) new laboratory characterization performed as needed. Thus SSURGO provides a local model of soil-landscape relations, developed in each area from the most significant
280 soil forming factors relevant to that area. SSURGO is progressively updated by field inspection and correlation, as problems are identified by soil surveyors or map users. ~~The SSURGO polygons are rasterized to gSSURGO.~~ Since SSURGO is compiled from diverse ~~field~~ surveys over many years, in some areas there are artefacts of that survey process (Supplementary Fig. 2S2).

~~PSP uses~~ The three DSM products (SG2, PSP, and SPCG) use a large number of gridded GIS coverages as environmental covariates in their predictive models. These represent soil-forming factors, and include climate, ecology, geology, land use/cover, terrain, vegetation and hydrography (Supplementary Information §4). PSP also uses coarse-resolution (≈ 2 km) estimates of
285 U, Th, and K γ -ray decay products to represent suspected variation in parent material kind and origin.

PSP (Chaney et al., 2019) uses the DSMART disaggregation algorithm (Odgers et al., 2014) to predict the most probable component (STU), along with their probability of occurrence, at each 30 m resolution grid cell, and from the modal soil

properties of the component, a probability-weighted aggregation. ~~In this context “disaggregation”~~ Disaggregation is the process
290 of ~~attempting to take examining~~ a coarser-resolution gridded or smaller-scale polygon product which is known to have multiple
STU, and identify the locations at a finer grid resolution where these components would be found, should the original survey
have been at larger scale. This depends on fine-scale covariates that, in theory, relate to the STU within an SMU. It attempts
to deal with the problems caused by multiple surveys over time, inconsistencies among mappers, and poor georeference of
SMU boundaries by sampling out of mapped SMU polygons according to declared proportions of map unit components
295 (STU) and using these as pseudo-observations to train ~~PSM-DSM~~ models of STU occurrence. PSP does not use any point
observations; rather, it samples pseudo-points from gSSURGO or gNATSGO and uses these as training points for the DSMART
disaggregation algorithm (see below). The model is trained in overlapping tiles, each containing some set of SSURGO primary
surveys, and using covariates covering just the tile. Thus each POLARIS tile is derived from a local model in two senses.
PSP provides a fine-scale map equivalent to $\approx 1:3\ 000$ design scale, i.e., from 16 to 64 times finer resolution than the original
300 1:12 000 to 1:24 000 surveys included in SSURGO. An obvious question is whether it is possible to map at this resolution
from the SSURGO source, even with the fine-resolution covariates used by DSMART. ~~See Chaney et al. (2019) for details.~~
because of the probabilistic nature of selecting pseudo-points to match with components (STU).

The other two methods are representative of the dominant ~~PSM-DSM~~ method as implemented, with some differences in
detail, in many countries and for many properties (e.g., Reddy et al., 2021; Liu et al., 2020; Araujo-Carrillo et al., 2021).

305 SG2 (Poggio et al., 2021) uses random forests implemented in the `ranger` R package, with prior covariate selection by
recursive feature elimination and model tuning by cross-validation of model hyperparameters (number of covariates at each
tree split, number of trees in the forest). ~~See Poggio et al. (2021) for details~~The model is trained for the whole world, not
per-country or region, thus it is a global model. SPCG~~This is based on the “homosoil” concept (Mallavan et al., 2010): identical
environmental conditions anywhere in the world should result in the same soils. Its use in DSM assumes that all soil forming
310 factors are fully specified (i.e., over their whole range and with all their possible interactions) in the model and training set.
Due to the uneven distribution of training points in covariate space, as well as portions of covariate space with no observations,
this ideal situation is not met. An obvious question is whether or not the additional information from outside the CONUS leads
to an improved model for this region.~~

SPCG (Ramcharan et al., 2018) is an extension of the original SoilGrids approach, but uses an ensemble of two tree-based
315 machine learning methods: random forests (as in the original SoilGrids) and gradient boosting. ~~See Rameharan et al. (2018)
for details.~~The model is trained for the CONUS, not per-region, thus it is a reduced version of the “homosoil” concept. It is a
global model in the sense of “use all information over a wide area”, although this is not the entire globe, as in SG2.

2.3 Resolution, depths and coordinate reference systems

About 90% of gSSURGO ~~and gNATSGO~~ is derived from polygon maps with a design scale (1:12 000 to 1:24 000, depending
320 on the original survey), which corresponds to MLD 1.44 to 2.3 ha (1:24 000) or 0.38 to 0.575 ha (1:12 000) polygons,
depending on the definition of MLD (see above). These correspond to single grid cell resolutions of 240 to 384 m (1:24 000)
or 60 to 96 m (1:12 000). ~~gSSURGO gNATSGO includes some areas surveyed at smaller scale (1:250 000).~~ gNATSGO

is delivered as gridded coverages at 30 m or 90 m horizontal resolution on an Albers Equal Area projection covering the CONUS, with standard parallels at 29.5° and 45.5° N and the central meridian at -96° E on the NAD83 datum, which uses the GRS80 ellipsoid. ~~gNATSGO is a 90-m-resolution generalization of this. We have used the 30 m resolution product. gSSURGO is delivered as gridded coverages at 10 m or 30 m horizontal resolution on the same CONUS projection. We have used the 30 m resolution product.~~ Property information is provided per horizon or layer, each with depth limits. Thus to produce a prediction for a depth ~~slice~~interval these must be aggregated by depth-weighted average by thickness across the depth ~~slice~~interval. PSP predicts at 1 arc-second of longitude and latitude resolution, i.e., 0.0002777778° on the WGS84 datum, equivalent to ≈ 32 m latitude, and proportionally smaller longitude depending on latitude. Depth ~~slice~~intervals are the standards specified by GlobalSoilMap. SPCG predicts at 100 m resolution for seven point depths (0, 5, 15, 30, 60, 100 and 200 cm) in the same projection as gNATSGO and gSSURGO. Predictions are means of a depth sliceinterval. SG2 predicts at 250 m resolution for the standard depth ~~slice~~intervals specified by GlobalSoilMap on an equal-area Interrupted Goode Homolosine (IGH) projection on the WGS84 datum (Moreira de Sousa et al., 2019). Depth ~~slice~~interval predictions are in fact point predictions at the centre of the depth ~~slice~~interval, considered to represent that interval. The Supplementary Information §3 explains how these products are accessed and made compatible for comparison at regional and local scales.

2.4 Uncertainty assessment

SG2 and PSP predict the 5% and 95% quantiles of the distribution of predictions ~~using~~. SG2 uses Quantile Regression Forests (QRF) (Meinshausen, 2006) ~~-, whereas PSP's uncertainty estimates are based on property data available for each STU predicted by POLARIS. The profile property data are used to create a depth-harmonized profile with uncertainty for each standard depth interval. These-~~

These uncertainty limits are specified by the GlobalSoilMap consortium (Arrouays et al., 2014), defined as “the 90% Prediction Interval (PI) which reports the range of values within which the true value is expected to occur 9 times out of 10 . . . there is no assumption that this prediction interval is necessarily symmetric around the predicted value.” (Science Committee, 2012).

~~gSSURGO provides-~~

gNATSGO and gSSURGO provide “representative”, “upper” and “lower” limit values of each property of a STU, per horizon or layer. The National Soil Survey Handbook, §618.2 (United States Department of Agriculture, Natural Resources Conservation Service, n.d.) explains that the representative value approximates the median, but that the quantiles corresponding to the low and high values can be adjusted to the percentiles which best show the spread of the property within an STU. If there are sufficient laboratory data of sampled profiles of the STU in the National Soil Information System (NASIS) (Natural Resources Conservation Service, n.d.), these are used as the basis for establishing the range. In all cases expert opinion is used to adjust these to represent the range that a map user can expect to find in the field. Thus these are not directly comparable to the results of QRF, but do give some idea of how the field mappers, supported by laboratory observations, conceive of the spread of a property. Note that none of these assessments imply a parametric probability distribution ~~-, only~~ ranges of selected quantiles. Libohova et al. (2014) discuss how these estimates can be derived for USA products following the GlobalSoilMap.net specifications.

As pointed out by Arrouays et al. (2020), “[t]he user community requires training in, and experience with, the new digital soil map products, especially about the use of uncertainties”. It would be hoped that the uncertainties computed by different methods would be similar, ~~but as will be seen here, that is not the case.~~

360 3 Evaluation methods

We compared ~~PSM-DSM~~ products at regional (nominal 250 m grid cells) and local (nominal 30 m grid cells) levels. We evaluated both qualitatively, i.e., by visual inspection followed by expert interpretation, and numerically, over a $1 \times 1^\circ$ tile, ~~selected based on its diverse soil-forming factors and environments and our familiarity with its soil geography. Spatial patterns were evaluated over a 0.2×0.2~~ For the pattern analysis within this area we selected a $0.20 \times 0.20^\circ$ subtile ~~and projected the maps to the UTM18N grid on the WGS84 datum (ESPG code 32618).~~

To compare maps at the regional resolution (250 m), the higher-resolution maps (gSSURGO, PSP, SPCG) were aggregated to the lower resolution by weighted averaging (resampling) of the high-resolution pixels within one low-resolution pixel. Thus there is smoothing inherent in the regional comparisons.

To compare maps at the local resolution, we only included the two products (gSSURGO and PSP) provided at that resolution,
370 along with the global product (SG2) as reference, this latter downscaled by increasing the grid resolution without any attempt to disaggregate within the larger grid cell, over a $0.15 \times 0.15^\circ$ subtile.

3.1 Qualitative methods

Qualitative methods for comparing maps rely on expert judgement to identify known soil-geographic patterns and evaluate to what extent they are represented on the gridded maps. The maps are displayed side-by-side along with a map of their pairwise
375 differences. Areas of disagreement are identified and discussed.

The DSM product can be evaluated at selected known points, typically from field observation of test areas: is the “correct” soil type or property predicted? and if not, is the error a reasonable approximation? More interesting are patterns in the DSM product. These can be compared to patterns used in the mental model of traditional soil survey, for example, toposquences and sequences of contrasting parent material.

In both cases (points and patterns) the evaluator may be able to infer which DSM covariates would be needed to improve the map.

3.2 Numerical methods – whole map

Numerical methods for comparing gridded maps as a whole include (1) MD: Mean difference (also known as the bias), i.e., the average disagreement between maps; (2) RMSD: root mean squared difference; (3) RMSD adjusted for MD, i.e., the RMSD
385 after ~~accounting for any bias~~ subtracting the bias from each prediction. These take the first-listed map as reference and the second as the map to evaluate. They can be normalized by the number of grid cells or total area. In addition, all maps can be compared by their Pearson (linear) correlations. These methods are of limited interpretive value ~~—they do characterize the entire~~

~~map, especially for overall. Their main use is to characterize the bias (MD), but do not explain where the spatial over the entire map; they do not reveal where any~~ discrepancies occur. For example, there can be no bias overall but a large difference in the amount and values of higher and lower differences. This will be reflected in the RMSD, although not shown on a map.

3.3 Numerical methods – spatial continuity

Soil properties are usually spatially-correlated: we expect similar values of properties in nearby grid cells. The degree of local spatial continuity can be assessed by the variogram computed over local neighbourhoods of the gridded map. We computed and modelled the variogram within a local neighbourhood and automatically fit with an exponential model, using the `fit.variogram` function of the `gstat` R package (Pebesma, 2004). Spatial structure is characterized by the range, proportional nugget and structural sill of the fitted variogram model. The range shows the radius over which the selected property has spatial correlation. The proportional nugget shows the ~~inherent variability at a point~~ variability at the prediction point at the centre of a grid cell, at a scale shorter than the grid spacing. The structural sill shows the overall variability within the range. These metrics show differences in spatial continuity (range), total variability (total sill) and short-range unexplained variability (proportional nugget) between maps.

3.4 Numerical methods – patterns

Numerical methods for comparing patterns include: (1) the “V-measure” method (§3.4.1) (Nowosad and Stepinski, 2018) implemented in the `sabre` “Spatial Association Between REgionalizations” R package (Nowosad, 2020); and (2) landscape-level metrics (§3.4.2) (Uuemaa et al., 2013) as used in ecology and derived from the FRAGSTATS computer program (McGarigal et al., 2012), implemented in the ~~landscapemetrics~~ landscapemetrics R package (Hesselbarth et al., 2019). These include Shannon diversity and evenness, landscape shape index, and fractal dimension. Although the ecological relevance of FRAGSTATS metrics have been criticized (Kupfer, 2012), here we use them to characterize spatial patterns of soil properties, not as inputs to landscape ecology models. Most of the metrics used here have also been used by Pindral et al. (2020) in a study of urban pedodiversity.

These methods must be applied to classified maps, so the continuous soil property maps must first be classified into ranges before analysis. ~~Clearly, different~~ Different choices of class limits and widths will result in different values of these measures. A somewhat objective method to choose classes is histogram equalization. The analyst determines the number of classes, and equal numbers of grid cells are in each class. To compare maps, the combined values of all maps are used to construct the histogram. For the “V-measure” ~~they must also~~ the gridded maps must be polygonized.

~~These metrics do not compare the values of the classes, only their spatial pattern.~~

3.4.1 V-measure

The V-measure ~~method evaluates the spatial association between two regionalizations, i. e., partitions of a map into classes, called~~ metrics compare different spatial partitions of the same domain, in this case, maps with classified soil properties. The

420 intent is to reveal how similar are these partitions. Two maps could have the same total areas of each class, and even the same number of polygons within each class, and even the same size distribution of these polygons, and yet be completely different in how they partition space into classes.

The polygons of a classified map are termed regions of a regionalization in the first (reference) map and zones of a partition in the second map. These are intersected to produce segments/segment polygons of the combined map. ~~The segments, which are~~ labelled with both zone and region ~~are classes. These polygons are then~~ used to compute two metrics ~~called of the map to~~ be evaluated (1) homogeneity and (2) completeness, both with respect to the ~~base (first-listed) regionalization~~ regionalization of the reference map.

The ~~Homogeneity~~ homogeneity of the second map is a measure of the variance of the regions within a zone, normalized by the variance of the regions in the entire domain of the first map. ~~If These variances are computed by the Shannon entropy based on areas of the segments. If the variance of the regions within the regions within zones have less variance than within the entire second map, the second map is to some extent~~ zones is small, the partition is relatively homogeneous with respect to the ~~first map regionalization~~. A perfectly homogeneous partition (~~second map~~) (with value 1) is when each zone of the ~~first map is entirely second map is~~ within a single region of the second reference map. In this case each zone has only one reference class. A perfectly inhomogeneous partition (with value 0) is when each zone has the same composition of regions as the entire domain of the first map, i.e., the second map's partition (to be ~~tested~~ evaluated) is essentially random with respect to the first map's regionalization.

Completeness of the second map is the inverse of homogeneity: it assesses the variance of the zones within a region, normalized by the variance of the zones in the entire domain of the second map. ~~This~~ It evaluates the homogeneity of regions with respect to zones, and shows how well the ~~partition of the second regionalization of the reference~~ regionalization ~~partition of the map to be evaluated. A perfectly complete regionalization is when each region of the reference map is entirely within a single zone of the map to be evaluated. In this case a polygon of the reference map will not be split among zones.~~

These two together are combined into a single measure, the V-measure, as the harmonic mean of homogeneity h and completeness c (Equation 1). This has a range between 0 (no spatial association between the maps) and 1 (perfect association). Obviously, we prefer high association between maps produced by ~~PSM-DSM~~ and a reference map. We can also assess the agreement of the patterns produced by different ~~PSM-methods~~ DSM methods, selecting one as a reference.

$$V = \frac{h \times c}{h + c} \quad (1)$$

3.4.2 Landscape metrics

The landscape metrics applicable to soil maps (as opposed to, e.g., ~~vegetation maps~~ maps of vegetation types) have diverse interpretations. We compare the metrics of two maps to see if they have a similar concept of the (~~classified~~) soil landscape. The landscapemetrics package can compute many ~~indices; we choose a few that will best~~ FRAGSTAT indices. We chose several that show the landscape-level difference between maps. We ~~do~~ did not consider metrics of individual patches, except as

they contribute to landscape-level metrics. The algorithms for these can be [read from the found in the package](#) code repository (Hesselbarth, 2021); [here](#) we present the formulas and [interpretations in the following text](#)[their interpretations](#).

- 455 – The **Shannon Diversity Index** sh_{di} (Equation 2), where p_i is the proportion of pixels of class $i = (1 \dots N)$, characterizing the landscape diversity [in two senses according to two factors: number of classes and their proportions. It is widely used as a summary diversity measure, although it does not distinguish between the two factors.](#) More classes and/or a more even distribution of proportions lead to a higher landscape diversity. This does not account for spatial contiguity, it just considers the class of each pixel, irrespective of position. [In this example the number of classes in each map will be similar, a maximum of eight \(the chosen histogram equalization classes, computed over the combined range of all](#)
460 [maps\), but some maps may lack representatives of the highest or lowest classes, and so will have only seven classes.](#)

$$D = - \sum_{i=1}^N p_i \ln p_i \quad (2)$$

- The **Shannon Evenness Index** sh_{ei} (Equation 3) is a normalization of Shannon Diversity by the maximum diversity possible for the given number of classes (N). It varies from 0 (completely uneven distribution - low landscape diversity) to 1 (all proportions are equal - high landscape diversity). It does not depend on the number of classes. [Again, this does not account for spatial contiguity, and thus isolates the effect of class proportion.](#)
465

$$E = \frac{D}{\ln N} \quad (3)$$

- The **Landscape Shape Index** ls_{i} (Equation 4), where A is the total area of the landscape and E' is the total length of edges, including the boundary, quantifies the internal boundary complexity of a landscape tile, with a value of 1 when the landscape consists of a single square patch, increasing without limit as the length of edges within the landscape increases. [This metric characterizes the degree of compactness of the contiguous areas of the classes.](#)
470

$$LSI = \frac{0.25E'}{\sqrt{A}} \quad (4)$$

- The **Landscape Aggregation Index** la_{i} (Equation 5), where g_{ii} is the number of like adjacencies, ~~max~~ g_{ii} (~~max~~ g_{ii}) is the classwise maximum possible number of like adjacencies of class i (i.e., if all pixels in the class were in one cluster), and P_i is the proportion of landscape comprised of class i , to weight the index by class prevalence. Thus la_{i} equals
475 the number of like adjacencies divided by the theoretical maximum possible number of like adjacencies, summed over

each class and over the entire landscape. It ranges from 0 for maximally disaggregated to 100 for maximally aggregated landscapes. This metric characterizes how dispersed are the classes.

$$AI = \left[\sum_{i=1}^m \left(\frac{g_{ii}}{max - g_{ii}} \right) P_i \right] (100) \quad (5)$$

480 – The **Mean Fractal Dimension** `frac_mn` characterizes the complexity of the landscape as the mean of the fractal dimension of all patches in the landscape. It approaches 1 if all patches are square, and 2 if all patches are irregular. It is scale-independent. The patch-level fractal dimensions (Equation 6) are computed from the patch perimeters p_{ij} in linear units and areas a_{ij} in square units; these are then averaged to obtain `frac_mn`.

$$FRAC = \frac{2 * \ln * (0.25 * p_{ij})}{\ln a_{ij}} \quad (6)$$

485 – The **Co-occurrence vector** `cove` proposed by Nowosad (2021) summarizes the entire adjacency structure of the map and can be used to compare map structures. This is a normalized form of the co-occurrence matrix, which counts all the pairs of the adjacent cells for each category in a local landscape, in the form of a cross-classification matrix. This vector can be considered as a probability vector for the co-occurrence of different classes. Co-occurrence vectors of different categorical maps can then be compared by computing the distance between them. Many distance measures are possible; we choose the Jensen-Shannon distance (Equation 7), which computes the entropy H of each probability vector v_i and entropy of their average, and from these the distance in entropy space between them. Increasing values indicate increasing dissimilarity in the adjacency patterns. The computation of `cove` is implemented in the `motif` R package, 490 and the Jensen-Shannon distance in the `philentropy` R package.

$$JSD(v_1, v_2) = H\left(\frac{v_1 + v_2}{2}\right) - \frac{1}{2}[H(v_1) + H(v_2)] \quad (7)$$

3.5 Regional patterns

495 Regional patterns are at the scale of regional trends such as lithologic units, elevation zones in mountains, and repeating patterns (e.g., basin-and-range, ridge-and-valley). ~~A “region” in this context is on the order of 1×1 to 5×5 . An appropriate resolution for this scale is $(250\text{ m})^2$, as used in SG2.~~

~~For this evaluation we produced difference maps of the PSM products (SG2, SPCG, PSP) vs. the gNATSGO gridded maps. All except SG2 were upsealed to the SG2 resolution (250 m).~~ The gNATSGO maps are ~~assumed to be the most correct~~ taken 500 as the reference, although we are well aware that they may not always correspond to ground truth. We then comment on the differences and speculate on the causes, based on our knowledge of the ~~PSM-DSM~~ procedures used to make each product and the nature of the soil landscape.

505 ~~We selected an example 1×1 test area, based on its well-known soil-forming factors and environments and our familiarity with the soil geography. For the pattern analysis within this area we selected a 0.20×0.20 subtile and projected the maps to the UTM18N grid on the WGS84 datum (ESPG code 32618).~~

3.6 Local patterns

Local patterns are at the scale of geomorphic features such as hillslope catenas, fluvial terraces, outwash fans, valley trains and drumlin fields. ~~“Local” in this context is on the order of 10×10 . An appropriate resolution for this scale is $\approx 30 \times 30$. Only gSSURGO and PSP map at this level; we included the other two products to see the effect of (unjustified) downscaling.~~

510 ~~For this evaluation we used the same test areas as [This evaluation within the test area](#) for the regional patterns, but examined smaller areas with [examining a smaller area with a distinctive soil-landscape relations pattern](#). The gSSURGO maps are taken as the reference. ~~These were~~ [This was](#) evaluated by two methods, [as follows](#).~~

3.6.1 Visual method

515 We produced ground overlays of key soil properties at selected depth ~~shies~~ [intervals](#) with corresponding KML specifications, and displayed these in Google Earth as semi-transparent overlays, using the original resolution of each product, projected into WGS84 geographic coordinates as required by Google Earth. These were then compared with ~~gSSURGO~~ [gNATSGO](#) maps streamed within Google Earth by SoilWeb Earth (California Soil Resource Lab, 2020). This shows the mapped polygons, labelled with their map unit, and linked to the map unit description, which in turn is linked to the Official Series Descriptions (OSD) (NRCS Soils, 2020b) with complete description of the soil properties modal values and ranges.

520 ~~Figure 3 shows SSURGO map units draped over a ground overlay of pH (0–5 cm) from SG2, produced by the SoilWeb streaming coverage in Google EarthPro, with a point query showing the SSURGO map unit composition. The map unit is described by its constituent soil series and their estimated proportions. Each series can then be queried for its Official Series Description (OSD) (NRCS Soils, 2020b), which gives a typical profile, a range of properties, and a link to lab data for the series. Clearly the pattern of properties follows the map unit delineations. This figure also shows the result of gridding the SSURGO polygons on a 30 m grid: grid cells overlap polygon boundaries.~~

~~SoilWeb view of SSURGO map units and a ground overlay of pH, 0–5 cm, predicted by SG2. Colours are low (red) to yellow (high) pH. Point inquiry at $-76^{\circ}38'05''$ E, $42^{\circ}19'53''$ N~~

3.6.2 Quantitative method

530 This follows the procedures of the regional assessment, except that V-measures are not computed, due to the very fine pattern of classified polygons.




Figure 1. Bedrock geology of Central New York State, transect from N-43° N (left) – 42°, centred on -76° 30' E. Orientation N (left) to S (right). Chronological and topographic sequence from Upper Silurian (N) through Upper Devonian (S) sedimentary rocks, notably the Onondaga limestone (green “Don”) and Tully limestone (crosshatched red, “Dt”). Source: (New York State Geological Survey, 1970)

4 Example area and soil property

To illustrate the method, we selected one area familiar to the first author, and an important soil property with strong spatial variability and pattern, namely pH in the 0–5 layer. We selected this property because in our experience this is often well-modelled by PSM-DSM methods. For example, SG2 had global cross-validation statistics of 0.78 pH median RMSD and a Model Efficiency Coefficient (MEC, the R^2 of the 1:1 line actual vs. observed) of 0.67 (Poggio et al., 2021). We select the topmost depth shies-interval because it is most represented by many environmental covariates, especially land cover as well as those derived from remote sensing. Thus the example shown here may be the best case, where we would hope that all mapping methods should provide similar results.

The example area is in central New York State, bounding box (-77 – -76° E), (42–43° N); the subtile for pattern evaluation was (-76.8 – -76.6° E), (42.2–42.4° N), centred at Cayuta NY. See the Supplementary Information for an explanation of the regional soil geomorphologyThe regional geomorphology is described by Bloom (2018). The underlying bedrock is a sedimentary sequence from Ordovician (north) to upper Devonian (south), with a wide variety of sedimentary facies. A strip of the bedrock geology map (New York State Geological Survey, 1970) covering part of the study area is shown in Fig. 1.

The entire area has been glaciated, the portion north of about 42° 15' (Valley Heads terminal moraines) somewhat more recently than the southern portion. A fragment of the surficial geology map (New York State Geological Survey, 1986) is shown in Fig. 2. This shows strongly-expressed features resulting from the most recent glaciation; these are well-known to the traditional soil surveyors. Many glacial features are present and relevant to soil geography: ground moraine, deep glacial troughs with proglacial lake sediments, beach lines, outwash valley trains, kame terraces and hanging deltas. Soil reaction in the northern half is largely controlled by limestone spread by the glacier from outcrops of the Onondaga and Tully limestones (Fig. 1), decreasing to the south.

5 Regional spatial patterns

5.1 Visual method




Figure 2. Surficial geology of Central New York State near Moravia NY. Legend: ground moraine (pink; if stippled shallow over bedrock), proglacial lakes (brown), organic swamps (dark green), bedrock or very thin soil cover (red), till moraine (purple), kame moraines (orange), lacustrine sand (light green), outwash sand and gravel (yellow). Source: (New York State Geological Survey, 1986)

Visual inspection of a DSM product over the landscape can be useful to identify anomalies and the degree to which the DSM product captures landscape features. These are over small areas where the soil-landscape relation is known to the evaluator.

555 This can not be part of a systematic evaluation, but can reveal areas of concern or agreement.

As an example, Figure 3 shows SSURGO map units, from a 1965 1:20k design scale soil survey, minimum legible delineation 1.6 ha (Cornell University Geospatial Information Repository (CUGIR), n.d.), draped over a ground overlay of pH (0–5 cm) from SG2, produced by the SoilWeb streaming coverage in Google EarthPro, with a point query showing the SSURGO map unit composition (Figure 4). The map unit is described by its constituent soil series and their estimated proportions. Each series can then be queried for its Official Series Description (OSD) (NRCS Soils, 2020b), which gives a typical profile, a range of properties, and a link to lab data for the series. In this case, the pattern of properties as predicted by SG2 somewhat follows the map unit delineations, but at a much coarser resolution. This is especially evident at the transition from the end moraine (map units beginning with H) and the steep slopes with thin till from local bedrock (map unit L₀F).

560

5.2 Regional maps

565 Table 1 shows the statistical differences between gNATSGO (reference) and the PSM-DSM products. All PSM-DSM products under-predict topsoil pH with respect to gNATSGO, by about 0.38–0.48 pH units. The RMSD is substantial also, on the order of 0.49–0.67 pH units, somewhat less than this when corrected for bias.




Figure 3. SoilWeb view of SSURGO map units and a ground overlay of pH, 0–5 cm, predicted by SG2. Colours are low (red) to yellow (high) pH. Centre at $-76^{\circ}38'04''$ E, $42^{\circ}20'07''$ N. Interactive view of SSURGO: <https://casoilresource.lawr.ucdavis.edu/gmap/?loc=42.33215,-76.63590,z15>

Figure 5 shows whole-map histograms. PSP has a bimodal distributionsdistribution, and predicts few pH values around pH 5.8. This is quite strange was unexpected, since this value is well-represented in the gNATSGO map. It may be an artifact of a covariate that is influential over a wider area than this tile and results in two regional distributions from contrasting elevation or climate zones. The other distributions are fairly symmetric, although SG2 and SPCG are more even than gNATSGO, which is strongly concentrated near pH 6.2. This shows the smoothing effect of the machine learning models.

Figure 6 shows the pairwise Pearson correlations between the products. The products are overall well-correlated. SG2 and SPCG are very closely correlated, since they use similar mapping methods, despite the additional covariates used by SPCG. PSP and gNATSGO are also closely-correlated. These correlations do not account for bias. They do however show that the maps are similar in their overall pattern as evaluated per-grid cell.

Figure 7 shows gNATSGO (reference) along with the predictions of pH of the PSP products. Figure 8 shows these as difference maps. These figures reveal substantial differences between products. The most obvious difference is in the detail of




Figure 4. SSURGO map unit HrC composition at $-76^{\circ}38'05''$ E, $42^{\circ}19'53''$ N. Link to interactive map unit summary: https://casoilresource.lawr.ucdavis.edu/soil_web/list_components.php?mukey=295620

the spatial pattern. Despite having been upscaled to regional resolution, gNATSGO shows finer detail than the other products, especially PSP.

These figures also show the spatial distribution of the bias compared to gNATSGO (as reference). Compared to gNATSGO, SG2 and SPCG underestimate under-predict pH in the higher hills in the NE portion of the map, and in the glacio-lacustrine sediments along the lakeshores. The disagreement along the lakeshores is because SG2 and SPCG do not use a surficial geology map, which would be especially useful in recently-glaciated areas such as this. The disagreement with in the higher hills seems to be a direct result of elevation. This is not because of extrapolation in feature space, because at these elevations SG2 also misses the soils derived from Onondaga limestone glacial till towards the southern end of the till plain. SG2 has no information on parent material and uses global models. SPCG has very similar differences, despite using SSURGO-derived parent material as a covariate.

PSP predictions are closer to gNATSGO than are those of SG2, which is not surprising since PSP also uses gSSURGO gNATSGO as its primary information source. This product has removed some of the fine variation of gNATSGO. However the disaggregation by DSMART results in quite some discrepancies with gNATSGO. In particular, the Homer-Tully outwash valley (northeast side of map) is under-predicted by one pH unit, and the surrounding hills over-predicted by almost as much.

Product	MD	RMSD	RMSD.Adjusted
SG2	3.796	6.111	4.789
PSP	3.843	4.908	3.052
SPCG	4.815	6.693	4.649

Table 1. Statistical differences between gNATSGO and PSM-DSM products, pHx10, 0–5 cm




Figure 5. Histograms of pHx10, 0–5 cm. Note the bimodal distribution of PSP and the flatter distributions of SG2 and SPCG compared to gNATSGO.




Figure 6. Pearson correlations between all products, pH, 0–5 cm. Strong correlations, especially between gNATSGO vs. PSP, and SG2 vs. SPCG.

Many of the valley trains (southern side of map, running towards the Susquehanna River) are under-predicted. This is likely due to PSP's soil series predictions, which are based on estimated map unit composition and random selection of series locations within map units for PSM-DSM calibration.

5.3 Uncertainty

The 5%, 50%, and 95% prediction quantile maps are shown in Fig. 9 (SG2) and 10 (PSP). The “low”, “representative” and “high” values from gNATSGO are shown in Fig. 11. Each figure has its own stretch. gNATSGO has narrower ranges than the two DSM products and by design does not include unrealistic values. SG2 and PSP have unrealistically wide ranges at all locations. In addition, PSP shows a curious feature: fine patterning at the two extremes that is not present at the median prediction.

Figure 12 shows the inter-quartile range 5–95% (IQR) for the two PSM-DSM products, along with the low-high range for gNATSGO.

- Quantiles of the prediction, SG2, pHx10, 0–5 cm
- Quantiles of the prediction, PSP, pHx10, 0–5 cm
- Low, representative, high values from gNATSGO, pHx10, 0–5 cm
- Inter-quartile ranges 0.05–0.95, pHx10, 0–5 cm




Figure 7. Topsoil (0–5 cm) pHx10, according to gNATSGO and [PSM-DSM](#) products. [See text for discussion.](#)




Figure 8. Difference between gNATSGO and PSM-DSM products, pHx10, 0–5 cm. See text for discussion.




Figure 9. Quantiles of the prediction, SG2, pHx10, 0–5 cm. Note the unrealistically wide range at all locations and consistent patterning among quantiles.




Figure 10. Quantiles of the prediction, PSP, pHx10, 0–5 cm. Note the unrealistically wide range at all locations and the fine patterning at the two extremes.




Figure 11. Low, representative, high values from gNATSGO, pHx10, 0–5 cm





Figure 12. Inter-quantile ranges 0.05–0.95, pHx10, 0–5 cm. SG2 IQR is fairly consistent from about 2.5 to 3.5 pH. PSP IQR has a wider range and more spatial patterning. gNATSGO “low-high” range is narrower.

SG2 has a fairly consistent IQR, mostly from about 2.5 to 3.5 pH, whereas PSP has a much wider range of uncertainties, mostly from about 1.5 to 4.5 pH, and shows much more spatial pattern. PSP has the widest ranges on the steep valley sides and especially in the Seneca Army Depot at the north inter-lake area, and the lowest on the broad till plains and through valleys. These are wide ranges, and although an honest reflection of the ~~PSM-DSM~~ models, should give pause to map users. This suggests that the GlobalSoilMap specifications for uncertainty (Arrouays et al., 2014) are unduly pessimistic. Sources for uncertainty assessment (SG2: training points and global covariates, PSP: mapped soil series and national covariates) and the different machine learning methods lead to greatly different estimates of prediction uncertainty. ~~gNATSGO has narrower ranges than the two PSM products and by design does not include unrealistic values.~~ The gNATSGO “low-high” range is narrower than the DSM IQR, but these are not comparable, because the expert-assigned range is not based on an estimate of a 5-95% IQR.

Figure 13 shows the differences between the IQR of the DSM products and the low-high range from gNATSGO. Both DSM products almost everywhere have substantially wider ranges than gNATSGO, however the pattern of differences is not similar. For example, the difference with SG250 is much larger in the north of the study area, whereas PSP has the larger differences in the southern hills.

IQR/range 5/95% vs. low/high, POLARIS-gNATSGO



IQR/range 5/95% vs. low/high, SG250-gNATSGO




Figure 13. IQR/range 5/95% vs. low/high, POLARIS-gNATSGO (left), SG250-gNATSGO (right), pHx10, 0-5 cm. Each figure has its own stretch

5.4 Local spatial autocorrelation

The local variograms and their fitted exponential models are shown in Fig. 14. Table 2 shows their statistics.




Figure 14. Fitted variograms, pH 0–5 cm. Semivariance units $(\text{pH} \times 10)^2$. Note the shorter range of gNATSGO and low sill of PSP.

Product	Effective range	Structural Sill	Proportional Nugget
gNATSGO	1938.00	10.32	0.00
SG2	3699.00	12.93	0.00
SPCG	6924.00	11.81	0.01
PSP	3918.00	6.50	0.02

Table 2. Fitted variogram parameters, pH 0–5 cm. Effective range in m; structural sill in $(\text{pH} \times 10)^2$, proportional nugget on $[0 \dots 1]$

gNATSGO has the shortest effective range. This ~~indicate~~ indicates fine-scale structure at 250 m resolution, which is of the same order as the minimum legible delineation (MLD) as a grid cell (see Introduction). The ~~PSM~~ mappers who defined the boundaries between soil classes (and thus representative property values) were able to divide the landscape at this high spatial frequency, if appropriate to the soil pattern. The DSM products have longer ranges, ~~showing that these models do not~~

625

PSM_products	V_measure	Homogeneity	Completeness
gNATSGO vs. SG2	0.0128	0.0143	0.0116
gNATSGO vs. SPCG	0.0258	0.0275	0.0243
gNATSGO vs. PSP	0.084	0.0897	0.079
SPCG vs. SG2	0.3342	0.3495	0.3201

Table 3. V-measure statistics, pHx10 0–5 cm

capture well fine-scale variation. These show a smoothing effect, likely due to the longer-range spatial continuity in many of the covariates. PSP has a long range and low sill longer range and lower sill than the gNATSGO from which it is derived, due to the harmonization from DSMART inherent in the DSMART algorithm. It has the highest proportional nugget, due to DSMART randomly assigning pixels within a gNATSGO map unit to its constituents, so that neighbouring pixels may be contrasting at the shortest separation. The very low proportional nuggets of the other products are due to the relatively coarse resolution.

5.5 Classification

Figure 15 shows the topsoil pH classified into eight histogram-equalized classes in the a 0.2 x 0.2° sub-tile. The higher pH are shown in green, the lower in red. Class limits are approximately 5.01, 5.14, 5.27, 5.40, 5.54, 5.71, and 6.02 pH, with the extreme values of 4.52 and 6.96 pH. The maps show obvious spatial differences in class distribution. gNATSGO shows more areas in the highest pH class than the PSP products DSM products, which is consistent with the results from continuous property maps. The products based on gSSURGO, i.e., gNATSGO and PSP, show a finer spatial pattern than the purely PSM products, i.e., SG2 and SPCG pattern of gNATSGO is the coarsest, because the classified values come from minimum-area polygons, whereas the DSM products predict per-grid cell. PSP shows the finest spatial pattern because of its disaggregation algorithm that randomly divides gNATSGO polygons according to component proportion. If these components are in different pH classes, there will be fine-scale pattern within the original polygon. But This is clearly the case in the large gNATSGO polygon in the northeast portion of the map (Connecticut Hill). In this example SPCG shows large homogeneous areas of the lowest pH class, covering the highest hills, whereas SG2 presents a more nuanced view.

5.6 V-measure

Table 3 shows the statistics from several V-measure comparisons, based on the histogram-equalized class maps. Only SG2 and SPCG have somewhat comparable patterns. gNATSGO is considerably different from all other products, due to its detailed spatial pattern based on field survey. the DSM products because of its derivation from minimum-area polygons.

Figure ?? shows the computed homogeneity and completeness 16 shows the inhomogeneity and incompleteness of the SG2 pH class map (the second map for the V-measure), with respect to the gNATSGO pH class map (the reference map). In the yellow These values are the inverse of the composite values of Table 3: the very low values in the table correspond to high values in the figure. In the homogeneity map, the blue polygons are the most homogeneous areas of the homogeneity map one, one






Figure 15. pH classes, 0–5 cm, central NY, detail. [Most areas of gNATSGO are in higher pH classes. PSP has the finest spatial pattern due to the DSMART disaggregation algorithm.](#)

Inhomogeneity -- SG2 vs. gNATSGO



Incompleteness -- SG2 vs. gNATSGO





Figure 16. Homogeneity (left) and completeness (right) measures of the SG2 pH class map, with respect to the reference gNATSGO pH class map, 0–5 cm. Values are the inhomogeneity of each zone (left) and incompleteness of each region (right).

Figure 17. Homogeneity (left) and ~~Completeness~~-completeness (right) measures of the SG2 pH class map, with respect to ~~gNATSGO~~-the SPCG pH class map, 0–5 cm. Values are the inhomogeneity of each zone (left) and incompleteness of each region (right).

655 ~~gNATSGO-predicted-class-is-contained-in~~ SG2 map, i.e., where an SG2 polygon has the most homogeneous set of gNATSGO classified values and thus comes closest to the reference. In the completeness map, the blue polygons are the most complete areas of the SG2 map, i.e., where the gNATSGO reference map has the most homogeneous set of SG2 classified values. The two maps have no areas with similar patterns.

660 A contrasting result is shown in Figure 17, which compares the SG2 ~~region; in the blue areas many are.~~ pH class map with respect to the SPCG map. These maps were made with similar methods and at the same resolution. ~~Overall the agreement is fairly good.~~ The inhomogeneity and incompleteness are much lower than for the previous map, showing that the pattern of these two classified maps are fairly similar.

product	ai	frac_mn	lsi	shdi	shei
gNATSGO	48.188	1.034	22.602	1.666	0.801
SG2	50.659	1.034	21.768	2.06	0.991
SPCG	58.483	1.041	18.557	1.887	0.907
PSP	47.025	1.04	23.232	1.898	0.913

Table 4. Landscape metrics statistics, pH 0–5 cm. frac_mn: Mean Fractal Dimension; lsi: Landscape Shape Index; shdi: Shannon Diversity; shei: Shannon Evenness; ai: Aggregation Index

~~Landscape metrics statistics, pH 0–5 cm. frac_mn: Mean Fractal Dimension; lsi: Landscape Shape Index; shdi: Shannon Diversity; shei: Shannon Evenness; ai: Aggregation Index~~

	gNATSGO	SG2	SPCG	PSP
gNATSGO	0.000	0.149	0.281	0.261
SG2	0.149	0.000	0.067	0.087
SPCG	0.281	0.067	0.000	0.111
PSP	0.261	0.087	0.111	0.000

Table 5. Jensen-Shannon distance between co-occurrence vectors

5.7 Landscape metrics

Table 4 shows the statistics from the landscape metrics calculations. The mean fractal dimensions are almost identical. There is quite some range of aggregations, with SPCG most aggregated, i.e., least complex. ~~Otherwise the results are inconsistent; all we can say is that the map patterns vary considerably among products~~PSP has the most complex landscape shape, due to its fine-scale disaggregation of gSSURGO polygons. The Shannon diversity indices are highest for SG2, indicating the most even areal division into classes. This may be an artefact of the histogram equalization.

Table 5 shows the Jensen-Shannon distance between co-occurrence vectors of the four products. The co-occurrence patterns of SG2 is quite similar to that of the other DSM products, whereas gNATSGO is quite different than PSP ~~and SPCG and somewhat different than SG2. This shows that, given this histogram equalization and for the selected property and depth interval, none of the DSM products well-match the pattern from traditional soil survey.~~

6 Local spatial patterns

The ~~main~~ interest here is to see how well ~~PSM-DSM~~ methods at relatively fine resolution reproduce known relations at the local geomorphic level, e.g., hillslopes, transects across valleys with multiple terrace levels, and within farms. It has been claimed that ~~PSM-DSM~~ at 30 m resolution is sufficient for management of, or even within, individual farm fields. ~~This-PSM~~ is the only ~~PSM-DSM~~ product which predicts at this resolution.

We examine this first qualitatively, i.e., by visual inspection, and then quantitatively, mostly following the methods of the regional assessment.

6.1 Qualitative assessment

Here we use silt concentration, as it reveals stronger qualitative discrepancies than pH in this test area. Figure 18 shows the silt concentration of the 0–5 cm layer for (top) the gridded SSURGO overlain on the original polygons from which it was derived, and (bottom) the disaggregated PSP grid cells in a hilly landscape near Caroline, NY. ~~Red colours are low silt, in this window alluvial fans (the C* map units). Pale grey colours are organic soils (the Hk, Hl map units). Light colours are high-silt surface soils (the L*, V*, B*, M* map units), from thin glacial till developed on shale and mudstone bedrock.~~

The gSSURGO product follows the SSURGO lines exactly. Some of the sharp boundary lines do correspond with abrupt transitions on the ground, for example where the steep hillsides are buried by fan alluvium. But others are not, for example on the hilltops. These differences are because the predicted silt concentrations are taken from the official series descriptions. PSP follows the map unit lines fairly well, but is much finer-grained; each 30 m pixel is separately predicted. This results in some smoothing of the abrupt boundary lines from gSSURGO on the hilltops. However within some SSURGO map units PSP predicts quite some differences in topsoil silt concentration. These are map units with contrasting components, which PSP attempts to disaggregate according to their correlation with covariates. For the most part these do not seem to be related to terrain or land use.

For example, Fig. 19 shows detail of the Holly-Papakting map unit within this PSP window. This map unit has two contrasting soils in similar proportions: a mineral alluvial soil (Holly series) and an organic soil (Papakting series); the second has much lower silt concentration.

It is difficult to see the reason for the pattern within this map unit. PSP has placed the component series in their proper proportions but not according to any ~~landscape feature~~apparent landscape feature or covariate.

Another example from this same area is shown in Supplementary Information §5.

6.2 Quantitative assessment

To see the fine differences at this high resolution, we consider a $0.15 \times 0.15^\circ$ subtile with lower-right corner $-76.30^\circ\text{E}, 42.45^\circ\text{N}$ and evaluate pH, as in the regional assessment (§5).

Table 6 shows the statistical differences between gSSURGO (reference) and the ~~PSM-DSM~~ products, along with the predictions of pH. Figure 20 shows the pairwise Pearson correlations between the maps. These results are comparable to those for the full tile at regional resolution: both SG2 and PSP under-predict pH by about 0.35–0.45 pH. Correlations are fairly strong between PSP and gSSURGO, and between SG2 and PSP, but weak between SG2 and gSSURGO.

Figure 21 shows gSSURGO (reference) along with the predictions of pH by the PSP products. Figure 22 shows these as difference maps. Clearly, gSSURGO has overall higher values than the other two products, and despite the fine resolution, has in general large areas of identical values. The differentiation between map units follows sharp boundaries even within a single landscape (e.g., the plateau towards the S of the map), and this is likely an artefact of relying on the representative profiles in




Figure 18. Ground overlay from gSSURGO (top) and PSP (bottom), silt % 0–5 cm, with SSURGO polygons from SoilWeb. Centre of image $-76^{\circ}16'25''$ E, $42^{\circ}22'53''$ N; view azimuth 247° . Red colours are low silt, in this window alluvial fans (the C* map units). Pale grey colours are organic soils (the Hk, H1 map units). Light colours are high-silt surface soils (the L*, V*, B*, M* map units), from thin glacial till developed on shale and mudstone bedrock. gNATSGO polygons have only one value, PSP disaggregates these, hence the pixelated pattern and somewhat smoothed boundaries.




Figure 19. Ground overlay from PSP in the Holly-Papaking map unit, silt % 0–5 cm. Centre of image $-76^{\circ}16'03''\text{E}, 42^{\circ}22'30''\text{N}$. Disaggregation appears to be random and not related to covariates.

PSM_product	MD	RMSD	RMSD.Adjusted
SG2	4.436	6.758	5.097
PSP	3.462	5.625	4.433

Table 6. Statistical differences between gSSURGO and PSM-DSM products, pHx10, 0–5 cm. Centre of map $-76^{\circ}30'30''\text{E}, 42^{\circ}52'30''\text{N}$

the official series descriptions for property values. PSP has a finer pattern, due to disaggregation, and shows a **more realistic** local pattern by smoothing out smoother local pattern, without the sharp boundaries between map units within a landscape. PSP shows large areas of low pH. SG2 does not follow well the landscape lines, especially the sharp boundaries between uplands and valleys, and predicts very low pH (≈ 4.5) on the plateau. It is difficult to recognize local landscape units in this global product.

6.2.1 Class maps

715 Figure 23 shows the topsoil pH classified into eight histogram-equalized classes. Class limits in this area are approximately 5.30, 5.44, 5.55, 5.61, 5.74, 5.89, and 6.15 pH, with the extreme values of 4.44 and 7.00 pH. SG2 clearly is less detailed than the other two products. PSP shows a fine pattern, not closely related to the fine pattern of gSSURGO. As previously noted, gSSURGO is consistently about one pH class higher than the other products.




Figure 20. Pearson correlations between local products, pH, 0–5 cm. [These moderate but weak for gSSURGO vs. SG2](#)




Figure 21. Topsoil (0–5 cm) pHx10, according to gSSURGO and [PSM-DSM](#) products. [See text for discussion.](#)




Figure 22. Difference between gSSURGO and PSM-DSM products, pHx10, 0–5 cm. [See text for discussion.](#)




Figure 23. pH classes, 0–5 cm. Coordinates are UTM 18N meters.

6.2.2 Local spatial autocorrelation

720 The local variograms and their fitted exponential models are shown in Fig. 24. Table 7 shows their statistics. gSSURGO has the shortest effective range and highest sill. PSP has a longer range and low sill, due to the harmonization from DSMART that removes some of the overall variability. SG2 has no nugget variance, a low sill, and long range, consistent with its regional scale.

6.2.3 Landscape metrics

725 Table 8 shows the statistics from the landscape metrics calculations. The mean fractal dimensions are almost identical. SG2 is much more aggregated, i.e., least complex, than gSSURGO or PSP. PSP has a higher landscape shape and Shannon diversity




Figure 24. Fitted variograms, pH 0–5 cm. Semivariance units $(\text{pH} \times 10)^2$. Note the short range of gSSURGO and low sill of SG2 and PSP.

Product	Effective range	Structural Sill	Proportional Nugget
gSSURGO	774.00	13.67	0.12
SG2	2550.00	7.34	0.00
PSP	1455.00	6.36	0.22

Table 7. Fitted variogram parameters, pH 0–5 cm. Effective range in m; structural sill in $(\text{pH} \times 10)^2$, proportional nugget on $[0 \dots 1]$

product	ai	frac_mn	lsi	shdi	shei
gSSURGO	73.658	1.049	71.395	1.845	0.887
SG2	87.647	1.106	34.978	1.941	0.934
PSP	56.376	1.045	116.476	2.006	0.965

Table 8. Landscape metrics statistics (local), pH 0–5 cm. *frac_mn*: Mean Fractal Dimension; *lsi*: Landscape Shape Index; *shdi*: Shannon Diversity; *shei*: Shannon Evenness; *ai*: Aggregation Index

	gSSURGO	SG2	PSP
gSSURGO	0.000	0.218	0.168
SG2	0.218	0.000	0.112
PSP	0.168	0.112	0.000

Table 9. Jensen-Shannon distance between co-occurrence vectors (local)

than the other products. Table 5-9 shows the Jensen-Shannon distance between co-occurrence vectors of the four products. The co-occurrence patterns of SG2 is somewhat similar to that PSP but quite different from gSSURGO.

7 Conclusions

730 The presented methods are well-able to expose differences between PSM in maps produced by different DSM mapping methods, and between these and field-traditional soil survey. There are also well-documented differences between maps based produced by traditional survey methods. For example, Bie and Beckett (1973) compared four independent surveys of a 19 km² area in Cyprus, and found that the maps differed considerably in their map unit purity and their proportions of interclass and intraclass variability. Thus the use of the NRCS products as reference should be seen as a basis for comparison with DSM products, not as the “truth”. However, it is the best available representation the of soil landscape at the given design scale and legend.

Our methods for comparing maps have two limitations due to the decision that they be applicable, using the supplied computer code, to any area within the USA. The first is the use of histogram equalization for the class maps which are then evaluated for the class pattern. For specific areas and properties it would be preferable to use established class limits relevant for land use, for example limits from soil survey interpretation tables. The second is the choice of the exponential model for automatic variogram fitting, as well as the somewhat arbitrary choice of empirical variogram cutoff and bin width. For each area, property and depth interval variograms could be computed and fit according to the analysts’ prior knowledge.

740 A variety of metrics to compare DSM products among themselves and to the reference map are proposed in this paper. This raises several questions about their utility and possible redundancy. Since we only consider one example case here, our conclusions are tentative. Comparing the results here with those in the companion case studies report, we know that these are context-dependent, and no general conclusions can be drawn. All the metrics provide useful information based on different summaries of the maps, so none are redundant.

A first question is which metrics best reflect the visual differences in patterns, e.g., for the regional patterns of Fig. 7. For the continuous maps, the whole-map histograms (Fig. 5) reveal whether the feature-space distribution of the property known from gNATSGO has been distorted by the DSM method. <<<< HEAD In the example case PSP produced a strongly bimodal distribution, so that its map shows few values near pH 5.8. ===== In the example case PSP produced a strongly bimodal distribution, so its map showed few values near pH 5.8. >>>> 56de475b489470b570c995fd75098582716df8e9 Much of the patterning in the strongly acid soil region is homogenized towards lower values, and there is a sharper boundary between the strongly and moderately acid areas. By contrast, both SG2 and SPCG reduced the peak modal value pH 6 and made more predictions towards the two tails of the univariate distribution. This can be seen in the resulting maps by more areas with the colours towards the two ends of the colour ramp. The whole-map variograms (Fig. 14) reveal the longer-range spatial continuity of the DSM products, compared to gNATSGO. This can be seen in the maps as less fine detail and larger areas with similar values.

755 For the classified maps (e.g., Fig. 15), the large discrepancies between them is due to the slicing from histogram equalization. Each landscape metric (Table 4) reveals a different aspect of the maps. For example, the aggregation index a_i shows that SPCG contains much larger one-class areas, on average, than the other products, and this is clear in the figure. Consistent with this, the landscape shape index l_{s_i} shows that SPCG has a simpler overall shape.

765 A second question is which metrics best discriminate the different DSM products. The whole-map histograms and variograms clearly show which products are more similar. In the example case, SG2 and SPCG are quite close and PSP is substantially different by both these metrics. The Jensen-Shannon distance between co-occurrence vectors (Table 5) clearly shows dissimilarity in the adjacency patterns of classes. In the example case again SG2 and SPCG are quite close, but here PSP is not too different. The landscape metrics were inconsistent in this case.

770 It is clear from the “best case” example presented in this paper that different PSM-DSM methods, with different training points, covariates, and algorithms, can produce quite different predictive soil maps. Comparing-Thus comparing maps with point-wise evaluation from (almost always biased) field observations gives an incomplete picture of how the different methods represent the soil landscape, which is after all what dictates how the soil is used and managed.

The main findings from the example case are:

- 775 1. Although the regional products (250 m resolution) are well-correlated, the PSM-DSM products are biased, under-predicting topsoil pH by about 0.38–0.48 pH units. They also differ substantially, with a RMSD adjusted for bias on the order of 0.31–0.48 pH. This is based on representative pH values of the mapped soil-seriesSTU, not on measured values.
- 780 2. The PSM-DSM products differ substantially among themselves and with the reference product in their local spatial pattern, as revealed by empirical variograms. gNATSGO has a short effective range, but this is smoothed to a range 2 to 3.5 times as long by PSM-DSM.
- 785 3. Classification by histogram equalization reveals major differences in the spatial patterns of the produced class maps, as evaluated both by visual inspection and landscape metrics.
- 790 4. Despite using USA-specific covariates (parent material, drainage classes) derived from gSSURGO-gNATSGO and covariates limited in geographic scope to the USA, the predictive map made by SPCG is not substantially different from that made by SG2, likely due to the similar modelling method.
- 795 5. The estimates of uncertainty provided by SG2 and PSP are substantially different, both in width of the uncertainty interval (much narrower in SG2) and in spatial pattern. This could be in part because SG2 is a global model, whereas PSP is based on local soil surveys and covariates restricted to one tile. The confidence intervals seem unrealistically wide compared to the expert-derived high-low value range provided by gSSURGOgNATSGO.
6. At the local level (30 m resolution) the disaggregation provided by PSP does not appear to correspond to landscape positions associated with STU components. PSP obscures the fine-scale details of the local spatial pattern, and SG is substantially more general, due to its resolution.

These results will differ in different soil geographic regions, for different soil properties, and for different depth slieesintervals, as shown in the companion Case Studies report.

795 Our methods have two limitations due to-Why are the results from these DSM examples so poor? Why do they not better approximate traditional surveys? We present some possible reasons:

1. The dominant DSM methods do not explicitly consider spatial continuity or pattern. Experiments have been started with convolutional neural networks and other methods with varying window sizes of covariates.
2. Environmental covariates to represent past soil-forming conditions (the “time” factor) are only available since the satellite remote sensing age, very short in terms of soil formation.
- 800 3. Environmental covariates to represent soil parent material (e.g., surficial geology) are not available globally, and even for the USA the decision that they be applicable, using the supplied computer code, to any area within the USA proxy of using parent material derived from SSURGO in SPCG was not of sufficient precision to improve the predictive models. The first is the use of histogram equalization for the class maps which are then evaluated for the class pattern.
- 805 4. Point observations were mostly placed by the soil surveyor at “typical” or “representative” locations in order to characterize map units, and do not capture the full range of variability along toposequences. For specific areas and properties it would be preferable to use established class limits relevant for land use, for example limits from soil survey interpretation tables. The second is the choice of the exponential model for automatic variogram fitting.
5. Poor georeference of legacy point observations, many from the pre-GPS era, leads to poor correlation with environmental covariates, hence to poor models, hence to much noise in the DSM product, which can obscure patterns.
- 810 6. Traditional soil survey uses is also a predictive activity. The surveyors uses as “covariates” (i.e., non-soil environmental information related to soil geography) what can be inferred from airphotos, as well as the somewhat arbitrary choice of empirical variogram cutoff and bin width. For each area, property and depth slice variograms could be computed and fit according to the analysts’ prior knowledge. direct landscape observation (terrain, vegetation, land use etc.). These give a more detailed and nuanced view of than possible at the resolutions used in practical DSM at regional scale, i.e., 100 m or coarser.
- 815

~~Despite the discrepancies between PSM-~~

Despite the discrepancies between DSM products and field survey, PSM-DSM can be a valuable tool for soil survey. Because of the expense and difficulty of field survey, in practice DSM is likely to be the most used method of making or updating soil maps in areas with no or poorly-resources soil survey organizations. For unsurveyed areas DSM can provide a useful pre-map

820 for planning sampling and field survey, thereby optimizing scarce resources for field work. It has the advantage of being reproducible and objective, given a set of training points, relevant environmental covariates, and a PSM-machine-learning method. ~~Therefore, its output should be examined and compared with field-based maps to identify possible improvements, especially in areas with difficult field access or complex, difficult to interpret soil-landscape patterns~~ Many of its problematic results are due to a set of training points, often with imprecise georeference, that do not properly occupy the covariate feature

825 space and, and to the lack of covariates to represent some aspects of pedogenesis over time. For unsurveyed areas PSM provides a useful pre-map for planning sampling and field survey.

830 In the USA (our study area) and in other countries with active soil survey programmes, DSM will be an important but not dominant tool in the overall survey. Soil survey as practiced by the NRCS uses methods from DSM, applied statistical modeling, and numerical ecology, along with an active and focused field programme. For example, supervised classification of terrain derivatives and satellite imagery has been successfully used to check internal consistency of map unit concepts and assist with the placement of delineations. The aim is to blend the most applicable tools from traditional field survey and applied statistical methods, supported by pedologic theory and regional land use considerations.

835 Of course, soil survey must be based on a proper examination of the soil itself. There is no substitute for actually examining the soil and landscape, ~~but despite the issues revealed in this study, PSM can be an important aid to the soil surveyor.~~ for either traditional soil survey, or as a reliable basis for DSM.

840 *Code availability.* Source code as R Markdown documents are available at <https://github.com/ncss-tech/compare-psm>. These can be used to (1) import all products to compare, as well as some others not considered in this study; (2) create ground overlays and corresponding KML files for display in Google Earth; (3) compare SG2 and PSP for $1 \times 1^\circ$ tiles; (4) compare SG2 with SPCG and gNATSGO for any rectangular tile; (5) compute landscape metrics and compare them between products for any subtile of these; (6) evaluate the success of PSP in disaggregating at 30 m resolution.

Author contributions. DGR conceptualized the approach, did most of the writing, wrote the R Markdown documents and performed the example case study. LP provided DSM expertise and detailed knowledge of SG2. DB and ZL provided USA-specific expertise, in particular about the NRCS and its products and services. All authors collaborated on the motivation, methods and conclusions.

Competing interests. There are no competing interests.

845 *Acknowledgements.* The contribution of Zamir Libohova was mostly accomplished during his tenure at USDA-NRCS-National Soil Survey Center, 100 Centennial Mall North, Room 152, Lincoln, NE 68508-3866 USA.

His contribution was partly supported by the LE STUDIUM Loire Valley Institute for Advanced Studies through its LE STUDIUM Research Consortium Programme.

References

- 850 Araujo-Carrillo, G. A., Varón-Ramírez, V. M., Jaramillo-Barrios, C. I., Estupiñan-Casallas, J. M., Silva-Arero, E. A., Gómez-Latorre, D. A., and Martínez-Maldonado, F. E.: IRAKA: The First Colombian Soil Information System with Digital Soil Mapping Products, *CATENA*, 196, 104940, <https://doi.org/10.1016/j.catena.2020.104940>, 2021.
- Arrouays, D., Grundy, M. G., Hartemink, A. E., Hempel, J. W., Heuvelink, G. B., Hong, S. Y., Lagacherie, P., Lelyk, G., McBratney, A. B., McKenzie, N. J., d.L. Mendonca-Santos, M., Minasny, B., Montanarella, L., Odeh, I. O., Sanchez, P. A., Thompson, J. A., and Zhang, G.-L.: GlobalSoilMap: Towards a Fine-Resolution Global Grid of Soil Properties, *Advances in Agronomy*, 125, 93–134, 2014.
- 855 Arrouays, D., McBratney, A., Bouma, J., Libohova, Z., Richer-de-Forges, A. C., Morgan, C. L., Roudier, P., Poggio, L., and Mulder, V. L.: Impressions of Digital Soil Maps: The Good, the Not so Good, and Making Them Ever Better, *Geoderma Regional*, 20, e00255, <https://doi.org/10.1016/j.geodrs.2020.e00255>, 2020.
- Batjes, N. H., Ribeiro, E., and van Oostrum, A.: Standardised Soil Profile Data to Support Global Mapping and Modelling (WoSIS Snapshot 2019), *Earth System Science Data*, 12, 299–320, <https://doi.org/10.5194/essd-12-299-2020>, 2020.
- 860 Bie, S. W. and Beckett, P. H. T.: Comparison of Four Independent Soil Surveys by Air-Photo Interpretation, Paphos Area (Cyprus), *Photogrammetria*, 29, 189–202, 1973.
- Bloom, A. L.: *Gorges History: Landscapes and Geology of the Finger Lakes Region*, Paleontological Research Institution, Ithaca, New York, 2018.
- 865 Brus, D., Kempen, B., and Heuvelink, G.: Sampling for Validation of Digital Soil Maps, *European Journal of Soil Science*, 62, 394–407, <https://doi.org/10.1111/j.1365-2389.2011.01364.x>, 2011.
- California Soil Resource Lab: SoilWeb Apps, <https://casoilresource.lawr.ucdavis.edu/soilweb-apps/>, 2020.
- Chaney, N., Minasny, B., Herman, J., Nauman, T., Brungard, C., Morgan, C., McBratney, A., Wood, E., and Yimam, Y.: POLARIS Soil Properties: 30-m Probabilistic Maps of Soil Properties over the Contiguous United States, *Water Resources Research*, 55, 2916–2938, 870 <https://doi.org/10.1029/2018WR022797>, 2019.
- Cornell University Geospatial Information Repository (CUGIR): Soil Survey, Tompkins County NY, 1965 (FGDC Metadata), <https://cugir-data.s3.amazonaws.com/00/74/98/fgdc.html>, n.d.
- D’Avelo, T. P. and McLeese, R. L.: Why Are Those Lines Placed Where They Are?: An Investigation of Soil Map Recompilation Methods, *Soil Survey Horizons*, 39, 119–126, <https://doi.org/10.2136/sh1998.4.0119>, 1998.
- 875 Forbes, T., Rossiter, D., and Van Wambeke, A.: *Guidelines for Evaluating the Adequacy of Soil Resource Inventories*, Cornell University Department of Agronomy, Ithaca, NY, 1982.
- Fridland, V. M.: Structure of the Soil Mantle, *Geoderma*, 12, 35–42, [https://doi.org/10.1016/0016-7061\(74\)90036-6](https://doi.org/10.1016/0016-7061(74)90036-6), 1974.
- Hengl, T., de Jesus, J. M., MacMillan, R. A., Batjes, N. H., Heuvelink, G. B. M., Ribeiro, E., Samuel-Rosa, A., Kempen, B., Leenaars, J. G. B., Walsh, M. G., and Gonzalez, M. R.: SoilGrids1km — Global Soil Information Based on Automated Mapping, *PLOS ONE*, 9, 880 e105992, <https://doi.org/10.1371/journal.pone.0105992>, 2014.
- Hengl, T., de Jesus, J. M., Heuvelink, G. B. M., Gonzalez, M. R., Kilibarda, M., Blagotić, A., Shangguan, W., Wright, M. N., Geng, X., Bauer-Marschallinger, B., Guevara, M. A., Vargas, R., MacMillan, R. A., Batjes, N. H., Leenaars, J. G. B., Ribeiro, E., Wheeler, I., Mantel, S., and Kempen, B.: SoilGrids250m: Global Gridded Soil Information Based on Machine Learning, *PLOS ONE*, 12, e0169748, <https://doi.org/10.1371/journal.pone.0169748>, 2017.
- 885 Hesselbarth, M. H.: *R-Spatialecology/Landscapemetrics*, *r-spatialecology*, <https://github.com/r-spatialecology/landscapemetrics>, 2021.

- Hesselbarth, M. H., Sciaini, M., With, K. A., Wiegand, K., and Nowosad, J.: *Landscapemetrics: An Open-Source R Tool to Calculate Landscape Metrics*, *Ecography*, 42, 1648–1657, <https://doi.org/10.1111/ecog.04617>, 2019.
- Hole, F. and Campbell, J.: *Soil Landscape Analysis*, Rowman & Allanheld, Totowa, NJ, 1985.
- Hudson, B. D.: *The Soil Survey as Paradigm-Based Science*, *Soil Science Society of America Journal*, 56, 836–841, <https://doi.org/10.2136/sssaj1992.03615995005600030027x>, 1992.
- 890 ISRIC - World Soil Information: *SoilGrids – Global Gridded Soil Information*, <https://www.isric.org/explore/soilgrids>, 2020.
- Kupfer, J. A.: *Landscape Ecology and Biogeography: Rethinking Landscape Metrics in a Post-FRAGSTATS Landscape*, *Progress in Physical Geography-Earth and Environment*, 36, 400–420, <https://doi.org/10.1177/0309133312439594>, 2012.
- Lagacherie, P., Andrieux, P., and Bouzigues, R.: *Fuzziness and Uncertainty of Soil Boundaries: From Reality to Coding in GIS*, in: *Geographic Objects with Indeterminate Boundaries*, edited by Burrough, P. A., Frank, A. U., and Salgé, F., *GISDATA 2*, pp. 275–286, Taylor & Francis, London, 1996.
- 895 Libohova, Z., Wills, S., and Odgers, N. P.: *Legacy data quality and uncertainty estimation for United States GlobalSoilMap products*, in: *GlobalSoilMap: Basis of the Global Spatial Soil Information System*, edited by Arrouays, D., McKenzie, N., Hempel, J., DeForges, A. C. R., and McBratney, A., pp. 63–68, Crc Press-Taylor & Francis Group, Boca Raton, <http://www.webofscience.com/wos/woscc/full-record/WOS:000341268700011>, wOS:000341268700011, 2014.
- 900 Liu, F., Rossiter, D. G., Zhang, G.-L., and Li, D.-C.: *A Soil Colour Map of China*, *Geoderma*, 379, 114–156, <https://doi.org/10.1016/j.geoderma.2020.114556>, 2020.
- Mallavan, B., Minasny, B., and McBratney, A.: *Homosoil, a Methodology for Quantitative Extrapolation of Soil Information Across the Globe*, in: *Digital Soil Mapping*, edited by Boettinger, J. L., Howell, D. W., Moore, A. C., Hartemink, A. E., and Kienast-Brown, S., pp. 137–150, Springer Netherlands, Dordrecht, <http://www.springerlink.com/content/g227720377103484/>, 2010.
- 905 McBratney, A. B., Mendonça Santos, M. L., and Minasny, B.: *On Digital Soil Mapping*, *Geoderma*, 117, 3–52, [https://doi.org/10.1016/S0016-7061\(03\)00223-4](https://doi.org/10.1016/S0016-7061(03)00223-4), 2003.
- McGarigal, K., Cushman, S. A., and Ene, E.: *FRAGSTATS v4: Spatial Pattern Analysis Program for Categorical and Continuous Maps*, Tech. rep., University of Massachusetts, Amherst, MA, <http://www.umass.edu/landeco/research/fragstats/fragstats.html>, 2012.
- 910 Meinshausen, N.: *Quantile Regression Forests*, *Journal of Machine Learning Research*, 7, 983–999, 2006.
- Meyer, H. and Pebesma, E.: *Predicting into Unknown Space? Estimating the Area of Applicability of Spatial Prediction Models*, arXiv:2005.07939 [cs, stat], <http://arxiv.org/abs/2005.07939>, 2020.
- Meyer, H. and Pebesma, E.: *Machine Learning-Based Global Maps of Ecological Variables and the Challenge of Assessing Them*, *Nature Communications*, 13, 2208, <https://doi.org/10.1038/s41467-022-29838-9>, 2022.
- 915 Minasny, B. and McBratney, A. B.: *Digital Soil Mapping: A Brief History and Some Lessons*, *Geoderma*, 264, Part B, 301–311, <https://doi.org/10.1016/j.geoderma.2015.07.017>, 2016.
- Moreira de Sousa, L., Poggio, L., and Kempen, B.: *Comparison of FOSS4G Supported Equal-Area Projections Using Discrete Distortion Indicatrices*, *ISPRS International Journal of Geo-Information*, 8, 351, <https://doi.org/10.3390/ijgi8080351>, 2019.
- Natural Resources Conservation Service: *Web Soil Survey*, <https://websoilsurvey.nrcs.usda.gov/>, 2019.
- 920 Natural Resources Conservation Service: *National Soil Information System (NASIS)*, https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/tools/?cid=nrcs142p2_053552, n.d.
- New York State Geological Survey: *Geologic Map of New York*, New York State Geological Survey, Albany, NY, <http://www.nysm.nysed.gov/research-collections/geology/gis>, 1970.

- New York State Geological Survey: Surficial Geologic Map of New York, New York State Geological Survey, Albany, NY, <http://www.nysm.nysed.gov/research-collections/geology/gis>, 1986.
- 925
- Nowosad, J.: *sabre*: Spatial Association Between Regionalizations, <https://nowosad.github.io/sabre/>, 2020.
- Nowosad, J.: *Motif*: An Open-Source R Tool for Pattern-Based Spatial Analysis, *Landscape Ecology*, 36, 29–43, <https://doi.org/10.1007/s10980-020-01135-0>, 2021.
- Nowosad, J. and Stepinski, T. F.: Spatial Association between Regionalizations Using the Information-Theoretical V-Measure, *International Journal of Geographical Information Science*, 32, 2386–2401, <https://doi.org/10.1080/13658816.2018.1511794>, 2018.
- 930
- NRCS Soils: Soils, <https://nrcs.app.box.com/v/soils>, 2020a.
- NRCS Soils: Official Soil Series Descriptions, https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/class/?cid=nrcs142p2_053587, 2020b.
- NRCS Soils: Description of Gridded Soil Survey Geographic (gSSURGO) Database, https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2_053628, 2022a.
- 935
- NRCS Soils: Gridded National Soil Survey Geographic Database (gNATSGO), <https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/geo/?cid=nrcseprd1464625>, 2022b.
- Odgers, N. P., McBratney, A. B., Minasny, B., Sun, W., and Clifford, D.: DSMART: An Algorithm to Spatially Disaggregate Soil Map Units, in: *GlobalSoilMap: Basis of the Global Spatial Soil Information System*, edited by Arrouays, D., McKenzie, N., Hempel, J., DeForges, A. C. R., and McBratney, A., pp. 261–266, CRC Press-Taylor & Francis Group, Boca Raton, 2014.
- 940
- Pebesma, E. J.: Multivariable Geostatistics in S: The Gstat Package, *Computers & Geosciences*, 30, 683–691, <https://doi.org/10.1016/j.cageo.2004.03.012>, 2004.
- Pindral, S., Kot, R., Hulisz, P., and Charzyński, P.: Landscape Metrics as a Tool for Analysis of Urban Pedodiversity, *Land Degradation & Development*, 31, 2281–2294, <https://doi.org/10.1002/ldr.3601>, 2020.
- 945
- Poggio, L., de Sousa, L. M., Batjes, N. H., Heuvelink, G. B. M., Kempen, B., Ribeiro, E., and Rossiter, D.: SoilGrids 2.0: Producing Soil Information for the Globe with Quantified Spatial Uncertainty, *SOIL*, 7, 217–240, <https://doi.org/10.5194/soil-7-217-2021>, 2021.
- R Studio: R Markdown, <https://rmarkdown.rstudio.com/>, 2020.
- Ramcharan, A., Hengl, T., Nauman, T., Brungard, C., Waltman, S., Wills, S., and Thompson, J.: Soil Property and Class Maps of the Conterminous United States at 100-Meter Spatial Resolution, *Soil Science Society of America Journal*, 82, 186–201, <https://doi.org/10.2136/sssaj2017.04.0122>, 2018.
- 950
- Reddy, N. N., Chakraborty, P., Roy, S., Singh, K., Minasny, B., McBratney, A. B., Biswas, A., and Das, B. S.: Legacy Data-Based National-Scale Digital Mapping of Key Soil Properties in India, *Geoderma*, 381, 114 684, <https://doi.org/10.1016/j.geoderma.2020.114684>, 2021.
- Rossiter, D. G., Poggio, L., Beaudette, D., and Libohova, Z.: How Well Does Predictive Soil Mapping Represent Soil Geography? An Investigation from the USA. Case Studies., ISRIC Report 2016-004, ISRIC-World Soil Information, ISRIC-World Soil Information, <https://doi.org/10.17027/isric-wdcsoils.20160004>, 2021.
- 955
- Schoeneberger, P. J., Wysocki, D. A., Benham, E. C., and Soil Survey Staff: *Field Book for Describing and Sampling Soils*, USDA Natural Resources Conservation Service, Lincoln, NE, 3.0 edn., <http://soils.usda.gov/technical/fieldbook/>, 2012.
- Science Committee: Specifications: Tiered GlobalSoilMap.Net Products; Release 2.3, Tech. rep., GlobalSoilMap.net, <http://www.ozdsm.com.au/resources/GlobalSoilMap%20specs%20version%20point3.pdf>, 2012.
- 960
- Scull, P., Franklin, J., Chadwick, O., and McArthur, D.: Predictive Soil Mapping: A Review, *Progress in Physical Geography*, 27, 171–197, <https://doi.org/10.1191/0309133303pp366ra>, 2003.

- Soil Survey Division Staff: Keys to Soil Taxonomy, US Government Printing Office, Washington, DC, twelfth edn., www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/class/, 2014.
- 965 Soil Survey Division Staff: Soil Survey Manual, no. 18 in USDA Handbook, Government Printing Office, Washington, DC, http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/planners/?cid=nrcs142p2_054262, 2017.
- Szatmári, G. and Pásztor, L.: Comparison of Various Uncertainty Modelling Approaches Based on Geostatistics and Machine Learning Algorithms, *Geoderma*, <https://doi.org/10.1016/j.geoderma.2018.09.008>, 2018.
- 970 Taghizadeh-Mehrjardi, R., Mahdianpari, M., Mohammadimanesh, F., Behrens, T., Toomanian, N., Scholten, T., and Schmidt, K.: Multi-Task Convolutional Neural Networks Outperformed Random Forest for Mapping Soil Particle Size Fractions in Central Iran, *Geoderma*, 376, 114 552, <https://doi.org/10.1016/j.geoderma.2020.114552>, 2020.
- Thompson, J. A., Kienast-Brown, S., D’Avello, T., Philippe, J., and Brungard, C.: Soils2026 and Digital Soil Mapping – A Foundation for the Future of Soils Information in the United States, *Geoderma Regional*, 22, e00 294, <https://doi.org/10.1016/j.geodrs.2020.e00294>, 2020.
- United States Department of Agriculture, Natural Resources Conservation Service: National Soil Survey Handbook, United States Department of Agriculture, Natural Resources Conservation Service, Washington, D.C., https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2_054242, n.d.
- 975 Uuemaa, E., Mander, U., and Marja, R.: Trends in the Use of Landscape Spatial Metrics as Landscape Indicators: A Review, *Ecological Indicators*, 28, 100–106, <https://doi.org/10.1016/j.ecolind.2012.07.018>, 2013.
- Vink, A.: Land Use in Advancing Agriculture, no. 1 in *Advanced Series in Agricultural Sciences.*, Springer-Verlag, New York, 1975.