

30 that yield-invigorating soils exhibited higher contents of organic matter than yield-debilitating soils and
31 harboured unique bacterial communities. Greater alpha diversity and higher relative abundances of
32 [Planctomycetes](#)[Planctomycetota](#) and [Chloroflexi](#)[Chloroflexota](#) were observed in yield-debilitating soils.
33 Co-occurrence network analysis revealed that yield-invigorating soils displayed a greater number of
34 [functionally interrelated modules \(meta-modules\)](#) and a higher proportion of negative links to positive
35 links. [Chloroflexi](#)[Chloroflexota](#) was recognized as a keystone taxon in manipulating the interaction of
36 bacterial communities in yield-invigorating soils. Structural equation modelling showed that soil organic
37 matter, beta diversity of bacterial community, and network connector ([Chloroflexi](#)[Chloroflexota](#)) were
38 [identified as potential](#) key factors [supporting in explaining the](#) high-yield pear production. Altogether,
39 we provide evidence that yield-invigorating soils across a range of locations appear to share common
40 features, including accumulation of soil organic matter, higher microbial diversity, enrichment of key
41 taxa like [Chloroflexi](#)[Chloroflexota](#), and maintaining a competitive network. These findings have
42 implications for science-based guidance for sustainable food production.

43

44 **Keywords:** Soil organic matter, Microbial diversity, Random forest prediction, Co-occurrence network,
45 Keystone taxa

46

47 **1 Introduction**

48 Soils are essential to human wellbeing due to their great contributions to the production of food, fiber,
49 feed, and medicine (Raaijmakers and Mazzola, 2016). Soil organisms play critical roles in maintaining
50 these ecosystem services, such as driving nutrient cycling, maintaining soil fertility, improving plant
51 productivity and suppressing plant diseases (Bender et al., 2016; Barrios, 2007). Microorganisms
52 participate in nearly all soil biological processes, and the microbial abundance, community composition
53 and activity primarily determine the sustainable productivity of agricultural lands (Philippot et al., 2013).
54 [Fungi participate in decomposition of organic matter and deliver nutrients for plant growth \(Frac et al.,](#)
55 [2018\), however, Given that considering that](#) bacteria are the most diverse and abundant group of
56 microorganisms in soil, bacterial communities and their functions can be pivotal indicators for crop
57 production in agroecosystems (van der Heijden et al., 2008).

58 In general, an increase in microbial diversity is linked to a high-yielding crop production mainly
59 through improving the host resilience to physical or chemical disturbances, modifying plant competition,

60 and facilitating plant access to nutrients (Chaer et al., 2009; Kennedy and Smith, 1995). Since individual
61 organisms do not live in isolation but rather form a complex system of inter-species interactions in soil,
62 interactions among community members were found to be related to crop production in the [potato](#)
63 monoculture system (Lu et al., 2013). Enrichments of key functional microbes in soil were deemed to
64 serve specific soil system functions, such as suppressing soil-borne pathogens and maintaining
65 sustainable crop production (Banerjee et al., 2018). However, the relative contributions of microbial
66 diversity, interactions among community members, or enrichment of key taxa to crop production remain
67 largely unknown. Therefore, it is highly desirable to identify pivotal indicators of bacterial community
68 composition in response to high-yielding crop production.

69 Changes in composition of soil bacterial communities across space are often strongly correlated
70 with soil pH (Fierer and Jackson, 2006). Soil pH has been recognized as a key driver in determining the
71 assembly of bacterial community in arable soils by field or microcosm experiments (Rousk et al., 2010).
72 However, recent studies have demonstrated that compositions of soil bacterial communities were driven
73 by a myriad of soil abiotic traits, such as organic matter contents, ~~nutrient contents and forms~~ [and contents](#)
74 [of soil nutrient](#) (Tian et al., 2018; Wang et al., 2018). For example, soil bacterial community composition,
75 which determines the ability of soil to suppress soil-borne pathogens, was found to be strongly correlated
76 with soil organic matter (Shen et al., 2018). An imbalanced ratio of soil nutrients, *i.e.*, ratio of nitrogen
77 to phosphorus or potassium could be a driving force altering the bacterial community composition in
78 long-term fertilized soils (Eo and Park, 2016). ~~Therefore, k~~Key soil chemical properties identified in
79 controlling the distribution and abundance of bacterial community is largely depending on [the](#)
80 [geographical distributions of soils](#) ~~soil sampling scale or treatments~~. As a consequence, a better
81 understanding of the relationship between soil edaphic properties and bacterial community composition
82 is critical to develop targeted manipulation options to increase soil service provisions.

83 ~~Pear (*Pyrus*) is the third most important temperate fruit species second only to grape and apple,~~
84 ~~belonging to the subfamily *Pomoideae* in the family *Rosaceae*. As a popular fruit in the world market,~~
85 ~~pear has been cultivated globally, and China is the biggest pear producer (FAOSTAT, 2019). 'Sucui No.~~
86 ~~1' pear, an early-maturing cultivated variety bred by the Jiangsu Academy of Agricultural Sciences,~~
87 ~~China, has displayed distinct advantages over other cultivates in Eastern and Central China, because this~~
88 ~~variety is easy to produce, adaptable to the environment, and has good quality and high economic~~
89 ~~benefits.~~ ["Sucui No. 1' pear, is an early-maturing variety bred by the Jiangsu Academy of Agricultural](#)

90 [Sciences, China, and has been popularly cultivated in Eastern and Central China, due to the advantages](#)
91 [including easy to produce, adaptable to the environment, and has good quality and high economic](#)
92 [benefits](#) –(Lin et al., 2013). With the increasing demand in China, sustainable production of high-quality
93 pear is becoming increasingly important. Manipulation of soil microbiomes has shown to be an effective
94 way to increase soil productivity (Chaparro et al., 2012). Considering that large-scale surveys could
95 exhibit the diversity of soil microbial communities exceeds what is found in host-associated communities
96 (~~Zorz-Toju~~ et al., ~~2019~~2018), it is necessary to explore the general microbial characteristics of multiple
97 yield-invigorating soils and identify key environmental drivers in assembling bacterial community.

98 In this study, ~~compared to local average yield, orchard showing higher pear yield production was~~
99 ~~recognized as yield-invigorating (YI) orchard while orchard displaying lower pear yield production was~~
100 ~~regard as yield-debilitating orchard (YD). After field surveys accomplished in 2019, six yield-~~
101 ~~invigorating and adjacent yield-debilitating pear orchards in total, which were identified through field~~
102 ~~surveys, were selected for further analysis of soil chemical properties and microbiome.~~ We hypothesized
103 that ~~high input of organic fertilizer could improve soil structure and modify chemical properties, which~~
104 ~~leading to YI yield-invigorating pear orchard~~ soils harbor unique bacterial communities ~~that maintains~~
105 ~~the high-yielding pear production, which are manipulated by key soil abiotic factors.~~ To address this, soil
106 bacterial communities and edaphic properties of ~~the study sites were compared six yield-invigorating~~
107 ~~and adjacent yield-debilitating pear orchards were compared~~ to (1) decipher the differences of taxonomic
108 diversity, and composition of the bacterial community, and (2) determine the contributions of
109 environmental variables to the changes in the structure of bacterial communities.

110 **2 Methods**

111 2.1 Study sites and experimental design

112 From July - August 2019, a field production survey of orchards cultivated with ‘Suci No. 1’ pear was
113 performed after pear fruits harvest to compare the differences of soil nutrients and microbiota between
114 yield-invigorating (~~YI~~) with yield-debilitating (~~YD~~) orchards. The locations, planting density, cropping
115 years, soil type and total yield were recorded. To minimize the effects of microclimate at each site, only
116 pair-located pear orchards with invigorating and debilitating yield and at similar growth stage were
117 selected for this research. In total, six pair-located yield-invigorating and -debilitating pear orchards
118 distributed in four cities of Jiangsu province, China, were selected in the main pear production areas ([Fig.](#)
119 [1, Table S1](#)). The yield per tree was obtained by dividing the total yield per hectare by plant density;

120 (~~Fig. 1A, Table S1~~).

121 **Fig. 1 here**

122
123 Paired yield-invigorating and yield-debilitating orchards from Fengxian (FX), Suining (SN) and
124 Tongshan (TS) were maintained in the Xuzhou city under the warm temperate sub-humid monsoon
125 climate. This site has a mean annual temperature (MAT) of 14.5 °C and mean annual precipitation (MAP)
126 of 847 mm. Orchards from Taixing (TX) were located in the Taizhou city under the humid southern
127 subtropical climate with a MAT of 15.3 °C and MAP of 1055 mm. Orchards from Gaochun (GC) were
128 located in the Nanjing city under the humid subtropical monsoon climate with a MAT of 15.4 °C and
129 MAP of 1106 mm. Orchards from Zhangjiagang (ZJ) were located in the Suzhou city under the humid
130 subtropical monsoon climate with a MAT of 15.7 °C and MAP of 1094 mm. For paired yield-invigorating
131 and-debilitating orchards, the irrigation and pesticide management practices were similar according to
132 farm records. However, yield-invigorating orchard was usually amended with more organic fertilizer
133 under integrated nutrients management whereas the co-located yield-debilitating orchard received more
134 chemical fertilizer under intensive management. ~~The yield per tree was obtained by dividing the total~~
135 ~~yield per hectare by plant density.~~Detailed information about **fertilization regimes for** each orchard is
136 shown in Table ~~S1~~ **S2**.

137 2.2 Soil sample collection and chemical properties determination

138 Along with the field survey, soil sampling campaigns were performed from July - August 2019 after pear
139 fruits harvest. For each yield-invigorating or -debilitating orchard, four subplots with three pear trees in
140 each subplot were randomly selected for soil sampling. Subsequently three soil cores (0-20 cm) under
141 the trunk base for each tree were collected using a 25 mm soil auger. In total, nine soil cores for each
142 subplot were pooled as a composite sample and finally four composite soil samples for each orchard
143 were collected and promptly transported on ice to the laboratory. After sifting through a 2 mm sieve and
144 thoroughly mixing, one portion of each soil sample was air-dried for chemical property analyses while
145 the remainder was stored at -70 °C for DNA extraction.

146 Soil chemical properties including soil pH, content of organic matter (OM), total nitrogen (TN),
147 available phosphorus (AP), available potassium (AK), alkali-hydrolyzale nitrogen (N), exchangeable
148 calcium (Ca), effective magnesium (Mg), effective iron (Fe), effective manganese (Mn), effective copper
149 (Cu) and effective zinc (Zn), were measured according to methods described by Shen et al. (2018) and

150 Huang et al. (2019). Briefly, soil pH was determined using a glass electrode meter in a suspension with
151 a 1:5 soil/water ratio (w/v). Soil OM was determined using the potassium dichromate external heating
152 method. TN was determined using a dry combustion method on an Element Analyzer (Vario EL,
153 Germany). AP and AK were determined using the molybdenum blue method after soil was extracted with
154 sodium bicarbonate and flame photometry after soil was extracted with ammonium acetate, respectively.
155 Soil alkaline hydrolysable nitrogen (N) was measured by the alkaline hydrolysable diffusion method.
156 Contents of soil Ca, Mg, Fe, Mn, Cu and Zn were determined by the atomic absorption spectroscopy
157 method using ICE 3300 AAS Atomic Absorption Spectrometer (ThermoScientific, USA) after acid
158 hydrolysis.

159 2.3 Soil DNA extraction and bacterial abundance quantification

160 Genomic DNA from 0.25 g soil for each sample was extracted by using the DNeasy® PowerSoil® Kit
161 (QIAGEN GmbH, Germany) according to the manufacturer's instructions. The abundances of soil
162 bacteria were determined with Eub338F/Eub518R primer using a 7500 Real Time PCR System (Applied
163 Biosystems, USA). Standard curves were generated by using 10-fold serial dilutions of a plasmid
164 containing a full-length copy of the 16S rRNA gene from *Escherichia coli*. Quantitative PCR analysis
165 was performed in 96-well plates with a 20- μ l mixture for each reaction using SYBR®Premix Ex Taq™
166 (TaKaRa, Japan). Thermal cycling was conducted according to a standard procedure with three replicates,
167 and the results were expressed as log copy numbers g⁻¹ dry soil.

168 2.4 Sequencing library construction and sequencing

169 The gene-specific primers 515F/806R with 12 bp barcode were used to amplify the V4 region of bacterial
170 16S rRNA gene on the BioRad S1000 (Bio-Rad Laboratory, CA) ~~roughly~~ according to the protocols
171 described by Caporaso et al. (2011). All constructed libraries were sequenced using the Illumina
172 NovaSeq 6000 at the Guangdong Magigene Biotechnology Co., Ltd. (Guangzhou, China).

173 2.5 Sequence processing

174 Quality filtering of the paired-end raw reads was performed to obtain the high-quality clean reads
175 according to the Trimmomatic ~~V0.33~~ (V0.33) (Bolger et al., 2014) quality control process. Sequences
176 were assigned to each sample based on their unique barcode, after which the barcodes and primers were
177 removed. Paired-end clean reads were merged using FLASH (V1.2.11 (Magoč and Salzberg, 2011)). Raw
178 tags were processed to generate the final ASV (Amplicon Sequence Variant) table file at 97% pairwise
179 identity according to the QIIME2 pipeline (Bolyen et al., 2019). The nonbacterial and mitochondrial

180 ASVs and extremely low frequency ASVs (relative abundance < 0.01%) were removed. A representative
181 sequence for each ASV was selected and classified using the RDP classifier [against the RDP Bacterial](#)
182 [16S database](#) (Wang et al., 2007) ~~against the RDP Bacterial 16S database.~~

183 2.6 Statistical analyses

184 Statistical analyses were performed using the software SPSS 20.0 ([SPSS Technologies, Armonk, NY,](#)
185 [USA](#)) and R (<http://www.R-project.org/>). Non-normal data were square-root or log transformed when
186 necessary. The significance of soil properties or microbial taxa in yield-invigorating or-debilitating
187 orchards was determined based on paired Wilcoxon rank sum test, and adjusted *P* values (< 0.05) were
188 obtained by the FDR method. Mantel tests were used to identify the correlations between microbial
189 community composition and pear yield, and soil chemical properties using the ‘vegan’ package ([Oksanen](#)
190 [et al., 2013](#)) in R. The linear regression analyses relating yield to selected microbial taxa or soil chemical
191 properties were conducted using the ‘basicTrendline’ package ([Mei et al., 2018](#)) in R.

192 Principal Coordinate Analysis (PCoA) based on the Bray-Curtis distance was performed in
193 MOTHUR (V1.38.1) (Schloss et al., 2009) and visualized by the ‘ggplot2’ package ([Wickham and Chang,](#)
194 [2015](#)) in R to explore the differences in microbial community composition. Permutational multivariate
195 analysis of variance (PERMANOVA) was performed to evaluate the significant differences of microbial
196 community composition according to sample locations and orchard yield using the ‘vegan’ package in R.
197 Microbial alpha diversity indexes (Chao, Shannon) were calculated based on randomly resampled ASV
198 abundance matrices at the same depth (23,800 sequences) in MOTHUR. A Venn diagram was generated
199 based on the final ASVs to compare microbial community composition between yield-invigorating and
200 -debilitating orchard soils. The affiliations of unique and shared ASVs in yield-invigorating and -
201 debilitating soils were compared to evaluate the differences of the bacterial community composition and
202 plotted using the ‘pheatmap’ package (<https://cran.r-project.org/web/packages/pheatmap/>) in R. Fold
203 changes (log₂ transformed) of shared ASVs across yield-invigorating and -debilitating soils were
204 calculated. The ASVs with fold change ratios > 2 and unique ASVs in yield-invigorating soils were
205 recognized as potential responders to yield promotion. In addition, to better understand the bacterial
206 community composition, relative abundances at the genus level were compared by STAMP software
207 v2.1.3 (Parks et al., 2014).

208 [The phylogenetic molecular ecological networks \(pMEN\) were constructed using the random](#)
209 [matrix theory-based approach to explore the organization of bacterial communities in yield-invigorating](#)

210 [\(YI\) or yield-debilitating \(YD\) soil samples](#). Potential ecological interactions among bacteria were
211 determined by modeling the microbial community using Molecular Ecological Network Analysis
212 (<http://ieg2.ou.edu/MENA>) based on pear yield ([Deng et al., 2012](#)). [Given the large number of rare taxa](#)
213 [that are specific to certain locations, ASVs that occurred in less than half of soil samples and lower than](#)
214 [0.01% were filtered, which resulting in 591 and 485 ASVs for YI and YD samples respectively, before](#)
215 [networks constructed. After removal of ASVs whose abundances were lower than 0.01%, the ASVs table](#)
216 [appeared in at least half soil samples were merged for phylogenetic molecular ecological network \(pMEN\)](#)
217 [construction \(Deng et al., 2012\)](#). The microbial network was constructed using random matrix theory-
218 based at 0.94 similarity threshold and visualized using Cytoscape 2.8.3 software ~~(V3.5.1,~~
219 ~~<http://cytoscape.org/>)~~ [Module \(Smoot et al., 2011\)](#). [Module](#) clustering and composition in yield-
220 invigorating and -debilitating networks were compared and plotted in R using the ‘pheatmap’ and
221 ‘ggplot2’ packages. [And the threshold values of Zi and Pi was 0.62 2.5 and 2.50.62, respectively for](#)
222 [topology analysis of the network](#). Redundancy analysis (RDA) was performed in the R ‘vegan’ package
223 to examine the relationship among frequencies of ASVs, samples and soil variables, which were selected
224 using ‘stepAIC’ in R. Variance partitioning analysis (VPA) was used to determine the contributions of
225 soil properties, sample location, and yield, as well as interactions among the variation in a microbial
226 community with hellinger-transformed data. The predictors of selected soil properties for explaining the
227 pear ~~yield-yield~~ were identified by random forest regression analysis (Boulesteix, et al., 2012). The
228 significance of each predictor in the response variables was assessed with the ‘rfPermute’ package ([Liaw](#)
229 [and Wiener, 2002](#)) with 1000 permutations based on 1000 trees. Structural equation modelling was
230 applied to evaluate relative contributions of soil chemical properties and bacterial community to pear
231 yield ([Grace et al., 2006](#); [Schermelleh-Engel et al., 2003](#)). The conceptual SEM fitness was examined on
232 the basis of a non-significant chi-square test ($P > 0.05$) and the goodness-of-fit index (GFI). Model was
233 fitted using the ‘lavaan’ package in R (Rosseel, 2012).

234 **3 Results**

235 3.1 Overview of sequencing data

236 In total, 1,622,858 16S rRNA sequences were retained after quality control and a total of 9,394 ASVs
237 were obtained for the 16S rRNA gene sequences based on 97% similarity. Among the total 16S rRNA
238 gene sequences, 159 ASVs with 74,372 sequences were classified as Archaea while 9,235 ASVs with
239 1,548,486 sequences were identified as Bacteria. Among Bacteria, [AcidobacteriaAcidobacteriota](#),

240 [ProteobacteriaPseudomonadota](#), [ChloroflexiChloroflexota](#), [PlanctomycetesPlanctomycetota](#) and
241 [ActinobacteriaActinomycetota](#) were the most abundant phyla (Fig. S1).

242 3.2 Soil chemical properties

243 Soil chemical properties differed significantly among the locations and orchard yield types (Table S3).
244 However, when taken all sites together, only a higher relative abundance of soil organic matter (OM) on
245 average was observed in yield-invigorating orchards compared to that in yield-debilitating orchards
246 based on the Wilcoxon test.

247

248 3.3 Bacterial abundances and community compositions

249 Yield-invigorating orchards together displayed significantly higher abundances of total bacteria than that
250 in co-located yield-debilitating orchards based on real time PCR result (Fig. ~~2B2A~~). Meanwhile, bacterial
251 community compositions at the ASV level were significantly correlated to pear yield ($r = 0.460$, $p =$
252 0.001) (Fig. ~~1C2B~~).

253

Fig. ~~12~~ here

254 PCoA based on Bray-Curtis distance matrices clearly revealed ~~treatment~~location-based differences
255 in bacterial community compositions (Fig. ~~2A3A~~). Six distinct groups representing samples from
256 different locations (FX, GC, SN, TS, TX and ZJ) were obviously separated and confirmed by
257 PERMANOVA test ($F = 14.9$, $P = 0.001$). At each location, soil bacterial community composition in
258 yield-invigorating orchards was significantly separated from that in co-located yield-debilitating
259 orchards, which was also confirmed by PERMANOVA test ($F = 3.6$, $P = 0.001$). Although only the
260 Shannon diversity in yield-invigorating orchards from GC and ZJ was significantly higher than that in
261 co-located yield-debilitating orchards (Fig. S2), the mean alpha diversity indices of Chao and Shannon
262 in all yield-invigorating orchards were significantly higher than those in all yield-debilitating orchards
263 based on the paired Wilcoxon test (Fig. ~~2B3B~~).

264

Fig. ~~23~~ here

265 The Venn diagram showed that ~~4,540~~ ASVs occupying over 90% of total sequences were shared
266 between yield-invigorating and -debilitating orchards (Fig. ~~2C3C~~). Among these shared ASVs, the fold
267 changes larger than 2 of ASVs in yield-invigorating compared to yield-debilitating orchards were
268 ~~recognized as potentially-potential responders linked-linking~~ to yield improvement. Surprisingly, none
269 of these ASVs potentially linked to yield improvement were shared among six separated collocated

270 orchards (Fig. S3). A total of 2,546 unique ASVs with 53,222 sequences were found in all yield-
271 invigorating orchards ~~and while~~ 2,308 unique ASVs with 44,389 sequences were observed in all yield-
272 invigorating orchards, ~~among~~ Among these unique ASVs, ~~which~~ almost 70% of ~~total~~ ASVs were
273 shared ~~among these unique ASVs~~ between yield-invigorating orchards and -debilitating orchards.
274 However, no shared unique ASVs were found among six separately located orchards. The affiliation of
275 unique and shared ASVs at the phylum level exhibited that the ProteobacteriaPseudomonadota,
276 PlanctomycetesPlanctomycetota, ChloroflexiChloroflexota, AcidobacteriaAcidobacteriota and
277 ActinobacteriaActinomycetota were the top five phyla (Fig. ~~2D3D~~).

278 At the phylum level, the relative abundances of bacterial dominant phyla varied across the location
279 and orchard yield condition ~~(Fig. 3A)~~. ProteobacteriaPseudomonadota, AcidobacteriaAcidobacteriota,
280 ActinobacteriaActinomycetota, ChloroflexiChloroflexota and PlanctomycetesPlanctomycetota were the
281 top five abundant phyla (Fig. 4). The mean abundance of ChloroflexiChloroflexota and
282 PlanctomycetesPlanctomycetota was significantly higher while Firmicutes was significantly lower in
283 yield-invigorating orchards compared to yield-debilitating orchards based on Wilcoxon test (Fig. 3A, Fig.
284 S4).

285 At a finer resolution, 967 genera were observed for all soil samples, among which 299 genera
286 appeared in more than half of soil samples in yield-invigorating or -debilitating orchards. However, only
287 34 genera displayed significant differences between yield-invigorating or -debilitating orchard soils
288 based on STAMP analysis (Fig. ~~3BS5~~). Interestingly, *Ornatilinea*, *Ktedonobacter*, *Longilinea*, belonging
289 to ChloroflexiChloroflexota, were significantly enriched in yield-invigorating orchard soils. *Gimesia* in
290 PlanctomycetesPlanctomycetota and *Arenimonas* in ProteobacteriaPseudomonadota showed
291 significantly higher relative abundances in yield-invigorating orchard soils than in yield-debilitating
292 orchard soils.

293 **Fig. ~~34~~ here**

294 3.4 Co-occurrence patterns of bacterial community

295 ~~The phylogenetic molecular ecological networks were constructed using the random matrix theory based~~
296 ~~approach to explore the organization of bacterial communities in yield-invigorating (YI) or yield-~~
297 ~~debilitating (YD) soil samples. After filtering ASVs that occurred in less than half of soil samples, 591~~
298 ~~ASVs for yield-invigorating samples and 485 ASVs for yield-debilitating samples were used to construct~~
299 ~~the networks.~~ The yield-invigoratingYI network contained 302 nodes, 448 edges, and 11 larger modules

300 (> 5 nodes), with an average connectivity (avgK) of 2.967, average path distance of 5.494 and clustering
301 coefficient (avgCC) of 0.152, while the values in the [yield-debilitating-YD](#) network were 235, 334, 9,
302 2.843, 6.232 and 0.131, respectively (Fig. [4A5A](#), Table [S2S4](#)). The module eigengene network analysis
303 revealed a difference in the higher-order organization between the two networks. Notably, the node
304 composition was substantially different between the two networks as the relative abundances of dominant
305 phyla were obviously different among different modules (Fig. [4B-5B](#) and C). A higher proportion of
306 nodes in the module of [yield-invigorating-YI](#) network was unique. ASVs affiliated to
307 [AcidobacteriaAcidobacteriota](#), [ChloroflexiChloroflexota](#), [ProteobacteriaPseudomonadota](#),
308 [ActinobacteriaActinomycetota](#), and [PlanctomycetesPlanctomycetota](#) within the unique modules (M9,
309 M10 and M11) were observed in the [yield-invigorating-YI](#) versus [yield-debilitating-YD](#) network.

310 Analysis using the threshold values of Z_i ([within-module connectivity](#)) and P_i ([among-module](#)
311 [connectivity](#)) showed that majority of nodes from both networks were categorized as peripherals that had
312 only a few links and almost always linked to the nodes within their own modules (Fig. [4D5D](#)). Although
313 only three nodes affiliated to [AcidobacteriaAcidobacteriota](#) were categorized as module hubs in the [yield-](#)
314 [invigorating-YI](#) network, seven nodes belonging to [AcidobacteriaAcidobacteriota](#),
315 [ActinobacteriaActinomycetota](#) and [ProteobacteriaPseudomonadota](#) were categorized as module hubs in
316 the [yield-debilitating-YD](#) network. Interestingly, four nodes including *Longilinea* species from
317 [ChloroflexiChloroflexota](#) in the [yield-invigorating-YI](#) network whereas only one node in the [yield-](#)
318 [debilitating-YD](#) network was categorized as module connectors (Table [S3S5](#)).

319 **Fig. 45 here**

320 3.5 Relationships between soil chemical properties and microbial community composition

321 ~~Soil chemical properties differed significantly among the locations and orchard yield types (Table S4).~~
322 ~~Together, yield invigorating orchards exhibited a significantly higher content of soil organic matter (OM)~~
323 ~~compared to yield debilitating orchards based on the Wilcoxon test.~~ Soil chemical properties were
324 significantly correlated to the bacterial community compositions (Mantel: $r = 0.803$, $p = 0.001$). Soil
325 chemical properties, location, and orchard explained 44.9% of the observed variation, leaving 55.1% of
326 the variation unexplained for bacterial community composition based on VPA [analysis-result](#) (Fig. [5A6A](#)).
327 Variation in the community composition was largely explained by soil properties (42.3%), and was also
328 influenced by locations and orchard yield types.

329 After forward stepwise selection, the module including soil OM, TN, alkaline N, AP and AK,

330 available calcium (Ca), copper (Cu) and manganese (Mn) explained the majority of the variation in
331 bacterial community composition (Fig. ~~5B6B~~). As evidenced by the RDA vectors, OM within the module
332 was ~~identified as among~~ the most-top important soil ~~properties~~ property that determines the composition
333 ~~in of shaping~~ bacterial community ~~composition~~. Random forest analysis showed that contents of soil Mn,
334 OM and Ca were the top parameters for predicting the orchard yield (Fig. ~~5C6C~~). Furthermore, soil OM
335 was also significantly correlated with bacterial communities as revealed by Mantel test (Fig. ~~5D6D~~, Table
336 ~~5556~~).

337 **Fig. ~~5-6~~ here**

338 3.6 Relationships of soil chemical and microbial indicators with orchard yield

339 Soil OM as potentially key soil chemical properties and bacterial alpha diversity, beta diversity and
340 relative abundance of ~~Chloroflexi~~ Chloroflexota and ~~Planctomyces~~ Planctomycetota as potentially key
341 microbial indicators ~~in determining associated with~~ pear yield were used to construct a model to explain
342 yield improvement. Final structural equation modelling (path analysis) (Fig. ~~67~~ and ~~S6~~) showed that the
343 strongest driver explaining yield improvement was beta diversity of bacterial community (PCoA) ($r =$
344 0.959 , $P < 0.001$), which was positively affected by content of soil OM ($r = 0.843$, $P < 0.001$). Alpha
345 diversity (Chao) of bacterial community also determined yield improvement to a large extent ($r = 0.542$,
346 $P = 0.009$). However, alpha diversity was not significantly correlated with content of soil OM.

347 **Fig. ~~6-7~~ here**

348 **4 Discussion**

349 Although pear is among the most important fruits worldwide, soil microbial communities in pear
350 orchards have been largely under-investigated (Huang et al., 2019). The present study attempts to
351 decipher the bacterial community linked to high-yield production of pear. Our results based on Mantel
352 analysis suggested ~~a directly~~ significant correlations ~~between among~~ bacterial community, soil chemical
353 properties and pear yield. Microbial characteristics responding to yield promotion have repeatedly been
354 observed on several crops depending on single experimental site (Zhong et al., 2020; Qiao et al., 2019;
355 Shen et al., 2013). It remained unclear, however, whether these distinctions are ubiquitous at a large-
356 scale. By comparing multiple co-located yield-invigorating and -debilitating orchards, we demonstrate
357 that high-yielding pear production soils exhibited high organic matter contents and harbored ~~shared~~
358 bacterial communities with high diversity, significantly enriched indigenous microbes and more
359 interactivewell-organized-interaction network, which was triggered by high-inputs of soil organic

360 ~~matter~~fertilizer. Here we discussed these main results and potential mechanisms in detail.

361 Microbial diversity is critical to soil ecosystems in maintaining the integrity, function and long-term
362 sustainability (Kennedy and Smith, 1995). Higher soil biodiversity is considered to be linked to a more
363 stable system and enhance the combination of vital microbial functions and processes (Cardinale et al.,
364 2006; Bell et al., 2005). In line with a previous report that crop yield was correlated to the soil bacterial
365 diversity (Zhao et al., 2014), greater diversity of bacterial community in yield-invigorating soils was
366 observed in the present study. ~~Hence we~~ ~~Our results indicate~~ ~~infer~~ that higher microbial diversity may
367 result in a more ~~productive stable~~ agroecosystem, contributing to sustainable pear production.

368 In this study, we found that ~~Proteobacteria~~Pseudomonadota, ~~Acidobacteria~~Acidobacteriota,
369 ~~Actinobacteria~~Actinomycetota, ~~Planctomyetes~~Planctomycetota and ~~Chloroflexi~~Chloroflexota were the
370 top abundant phyla. This result roughly agreed with previous studies showing that
371 ~~Proteobacteria~~Pseudomonadota, ~~Acidobacteria~~Acidobacteriota and ~~Actinobacteria~~Actinomycetota are
372 usually dominant bacterial taxa in agricultural soils (Xun et al., 2019; Dai et al., 2018), while
373 ~~Planctomyetes~~Planctomycetota and ~~Chloroflexi~~Chloroflexota exhibit an unexpectedly high relative
374 abundance in rice cropped soil (Edwards et al., 2015) and sandy loam soil (Pathan et al., 2021). The
375 highest relative abundance of ~~Proteobacteria~~Pseudomonadota was probably explained by the fact that
376 ~~Proteobacteria~~Pseudomonadota are considered as copiotrophic bacteria and ~~always~~ flourish in soils with
377 large amounts of available nutrients (Fierer et al., 2007).

378 Moreover, a significantly higher abundance of ~~Planctomyetes~~Planctomycetota and
379 ~~Chloroflexi~~Chloroflexota was observed in yield-invigorating orchards, indicating that
380 ~~Planctomyetes~~Planctomycetota and ~~Chloroflexi~~Chloroflexota may be ~~responsible~~ ~~associated with~~ ~~for~~
381 pear yield-improvement. There is no direct evidence showing that ~~Planctomyetes~~Planctomycetota could
382 improve plant growth. ~~However~~, ~~Planctomyetes~~Planctomycetota has been reported to be involved in
383 ~~the~~ ~~many~~ soil biological processes such as ammoxidation, carbohydrate and polysaccharide
384 ~~metabolism~~metabolie (Fuerst, 2017). This implies that ~~Planctomyetes~~Planctomycetota may promote
385 plant production through improving soil ~~biological~~ fertility. ~~Chloroflexi~~Chloroflexota is a facultative
386 anaerobic phylum including autotrophic, heterotrophic and mixotrophic taxa (Speirs et al., 2019).
387 Considering that soil amended with organic fertilizer ~~could~~ ~~may~~ enhance the soil water holding capacity,
388 the yield-invigorating soils with more organic material input have a higher soil moisture content,
389 especially after irrigation, probably leading to the enrichment of ~~Chloroflexi~~Chloroflexota in soil.

390 ~~Furthermore, it has been well documented that ChloroflexiChloroflexota could grow well in drought~~
391 ~~conditions (Ullah et al., 2019), implying that yield invigorating soils with a higher relative abundance of~~
392 ~~ChloroflexiChloroflexota may exhibit excellent resistance to environmental stress to support sustainable~~
393 ~~crop production.~~

394 Network analysis is a systems-level method to explore interactions within an ecosystem that cannot
395 be directly observed through co-occurrence analysis (Fath et al., 2007). Similar to the food web network
396 analyses in macro ecosystems, microorganisms also form complex interactions with other species (Faust
397 and Raes, 2012) and have been widely investigated to explore the linkage of microbial network with soil
398 function, such as nutrient supply (Fan et al., 2021) and disease suppression (Lu et al., 2013). Overall, in
399 line with previous findings (Hu et al., 2020), the topological properties of the constructed networks,
400 including connectivity, average clustering coefficients, average degree distance, and modularity indicate
401 that these networks are scale-free, modular and “small world”. In short, a scale-free network represents
402 that a network whose connectivity follows a power law, and most of nodes have only a few connections
403 with other nodes. Meanwhile, a small-world network is the network in which most nodes are not
404 neighbors of one another, but most nodes can be reached by a few paths. Modularity is a fundamental
405 characteristic of biological network as a module in the network is a group of nodes that are highly
406 connected within the group, but very few connections outside the group (Deng et al., 2012). Our
407 comparative network analysis indicated that microbial co-occurrence patterns in soils links to- different
408 ~~were correlated to~~ pear production. As a meta-module is usually considered as a group of modules
409 functionally interrelated (Langfelder and Horvath, 2007), a greater number of meta-modules were
410 identified in the network constructed from yield-invigorating soils, suggesting that a greater number of
411 network nodes in the yield-invigorating soils were functionally interrelated than those in the yield-
412 debilitating soils. A majority of nodes in the meta-modules were not shared between yield-invigorating
413 and -debilitating networks, indicating basal shifts in network architecture during pear production with
414 contrasting yield performance.

415 Furthermore, a higher proportion of negative interactions to positive interactions were identified in
416 the network constructed from yield-invigorating network than the yield-debilitating network. Our results
417 indicated ~~that~~ stronger resource competitions in yield-invigorating soils, which means that the soil co-
418 occurrence network was more stable to maintain soil ecosystem function (Coyte et al., 2015). In this our
419 study, three module connectors and three module hubs were identified as potentially key taxa in the yield-

420 invigorating network. Interestingly, among those key species, ASV357 affiliated to *Longilinea*,
421 belonging to the ~~Chloroflexi~~Chloroflexota, was recognized as a key phylum in improving pear yield.
422 Similarly, ~~Chloroflexi~~Chloroflexota was reported to be key-stone taxa in the constructed network from
423 agricultural soils with 40-years fertilization (Fan et al., 2021). ~~Chloroflexi-Chloroflexota play key roles~~
424 ~~in connecting network nodes of soil microbiome probably due to that Chloroflexota could participate in~~
425 ~~degrading plant compounds to create more nutrients via pathways for the degradation of starch, cellulose,~~
426 ~~and longchain sugars, as it is positively correlated with genes for amino sugars, sugar alcohols and simple~~
427 ~~carbohydrate metabolic pathways (Hug et al., 2013).~~ play key roles in manipulating soil microbiome
428 probably due to that Chloroflexi could participate in degrading plant compounds to create more niches
429 via pathways for the degradation of starch, pyrogallol, cellulose, and longchain sugars, as it is positively
430 correlated with genes for amino sugars, sugar alcohols and simple carbohydrate metabolic pathways
431 (Hug et al., 2013).

432 Soil pH is generally recognized as the main driver in the assembly of bacterial community;
433 especially in the studies related to geographic distribution of microorganisms (Fierer and Jackson, 2006).
434 Soil pH varying across a wide range allows insights into the relationships between pH and soil bacterial
435 communities in those researches. Therefore, we speculated that there are important factors other than pH
436 in shaping soil bacterial communities in our study, given that soil pH only ranged within two units. In
437 this study, a significantly higher content of soil organic matter was observed in yield-invigorating
438 orchards, demonstrating that soil organic matter ~~can~~ could also drive the assembly of bacterial community.
439 Consensus is emerging that microbial ~~residues~~materials are an important constituent of soil organic
440 matter (Kallenbach et al., 2016), ~~a higher content of soil organic matter usually supports a more diverse~~
441 ~~microbial community~~the quantity and quality of soil organic matter determines the diversity of microbial
442 ~~community~~, which participate in almost all soil biological processes (Fierer, 2017). ~~Despite the quality~~
443 ~~of soil organic matter was not evaluated in this study, the quality of soil organic matter was associated~~
444 ~~with the diversity of microbial community (Ding et al., 2015), which implies more attentions should be~~
445 ~~paid to illustrate the relationship between the quality of soil organic matter and microbial community in~~
446 ~~our future work.~~

447 ~~Structural equation modelling approach has been widely used to decipher keystone indicators~~
448 ~~associated with soil function and crop production in agroecosystems (Jiang et al., 2020; Chen et al., 2019).~~
449 ~~In the present study, we observed that soil organic matter, beta diversity of bacterial community, and~~

450 network connector were key indicators in supporting high-yield pear production based on the structural
451 equation modelling results. Worth to mention, soil organic matter was not directly linked to the yield in
452 the constructed model, indicating that soil organic matter maintain the high-yielding pear production
453 probably via the indirect ways. Therefore, we proposed that yield-invigorating soils harbour unique
454 bacterial communities that may improve soil biological fertility, which could be driven by soil organic
455 matter and manipulated by keystone species (Chloroflexota) through altering the bacterial interactions.

456
457 ~~Structural equation modelling approach has been widely used to decipher keystone indicators~~
458 ~~associated with soil function and crop production in agroecosystems (Jiang et al., 2020; Chen et al., 2019).~~
459 ~~In the present study, we observed that soil organic matter, beta diversity of bacterial community, and~~
460 ~~network connector were key indicators in supporting high yield pear production based on the structural~~
461 ~~equation modelling results. Therefore, we proposed that yield invigorating soils harbour unique bacterial~~
462 ~~communities that could improve soil biological fertility, which could be driven by soil organic matter~~
463 ~~and manipulated by keystone species (ChloroflexiChloroflexota) through altering the bacterial~~
464 ~~interactions.~~

465 75 **Conclusions**

466 In conclusion, ~~by comparing six paired located orchards, our results demonstrated that~~ yield-invigorating
467 soils ~~showed~~ displayed a higher content of organic matter and harboured unique bacterial community
468 with greater diversity than yield-debilitating soils. ~~We further~~ Further highlight that
469 ~~ChloroflexiChloroflexota~~ was significantly enriched and ~~served-identified~~ as a potential keystone taxon
470 in manipulating the interaction of bacterial community in yield-invigorating soils. These findings ~~help~~
471 ~~elucidate~~ indicated that soil organic matter triggered the assembly of soil microbiome. – the role of soil
472 microbiome which both participated in maintaining crop production ~~and factors controlling the assembly~~
473 ~~of soil microbiome~~. Such knowledge is a first step toward harnessing soil microbiome in support of
474 sustainable agroecosystems.

475
476 **Data availability.** Raw amplicon sequencing data for each sample used in this study was deposited at
477 the National Center for Biotechnology Information (NCBI) in the FASTQ format and is available under
478 the accession number PRJNA749397. Other data that support the findings of this study are available on
479 request from the corresponding author (Xiaomei Ye).

480

481 **Authors' contributions**

482 L. Wang: performed all experiments; L. Wang, X. Ye, and Z. Shen: designed the study, and wrote the
483 majority of the manuscript; L. Wang and Z. Shen and C. Tao: analyzed the data; H. Hu, J. Du, Y. Xi, J.
484 Lin, and D. Chen: participated in the design of the study, provided comments and edited the manuscript.
485 The authors read and approved the final manuscript.

486

487 **Competing interests.** The authors declare that they have no conflict of interest.

488

489 **Disclaimer.** Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional
490 claims in published maps and institutional affiliations.

491

492 **Acknowledgements.** We sincerely thank all those who have assisted me with any part of this paper and
493 all pear orchards owners for providing access to the soil sampling.

494

495 **Financial support.** This research was supported by Jiangsu Agricultural Science and Technology
496 Innovation Fund (CX(19)3094) and National Natural Science Foundation of China (31801842 and
497 42090065).

498

499 **Appendix A. Supplementary data**

500 Supplementary figures and tables to this article can be found in the supplemental material.

501

502 **References**

503 Banerjee, S., Schlaeppli, K., van der Heijden, M.G.J: Keystone taxa as drivers of microbiome structure
504 and functioning. *Nat. Rev. Microbiol.*, 16, 567-576, 2018.

505 Barrios, E.: Soil biota, ecosystem services and land productivity. *Ecological Eco.*, 64, 269-285, 2007.

506 Bell, T., Newman, J.A., Silverman, B.W., Turner, S.L., Lilley, A.K.: The contribution of species richness
507 and composition to bacterial services. *Nature*, 436, 1157-1160, 2005.

508 Bender, S.F., Wagg, C., van der Heijden, M.G.A.: An underground revolution: biodiversity and soil
509 ecological engineering for agricultural sustainability. *Trends Ecol Evol.*, 31, 440-452, 2016.

510 [Bolger, A.M., Lohse, M., Usadel, M.: Trimmomatic: a flexible trimmer for Illumina sequence data.](#)
511 [Bioinformatics, 30, 2114-2120, 2014.](#)

512 Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., Alexander, H.,
513 Alm, E.J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J.E., Bittinger, K., Brejnrod, A., Brislawn,
514 C.J., Brown, C.T., Callahan, B.J., Caraballo-Rodríguez, A.M., Chase, J., Cope, E.K., et al.:
515 Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat.*
516 *Biotechnol.*, 37, 852-857, 2019.

517 Boulesteix, A.L., Janitza, S., Kruppa, J., König, I.R.: Overview of random forest methodology and
518 practical guidance with emphasis on computational biology and bioinformatics. *Wires Data Min.*
519 *Knowl.*, 2, 493-507, 2012.

520 Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A., Turnbaugh, P.J., Fierer,
521 N., Knight, R.: Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample.
522 *P. Nat. Acad. Sci. USA*, 108, 4516-4522, 2011.

523 Cardinale, B.J., Srivastava, D.S., Duffy, J.E., Wright, J.P., Downing, A.L., Sankaran, M., Jouseau, C.:
524 Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature*, 443, 989-992,
525 2006.

526 Chaer, G., Fernandes, M., Myrold, D., Bottomley, P.: Comparative resistance and resilience of soil
527 microbial communities and enzyme activities in adjacent native forest and agricultural soils. *Microb*
528 *Ecol.*, 58, 414-424, 2009.

529 Chaparro, J.M., Sheflin, A.M., Manter, D.K., Vivanco, J.M.: Manipulating the soil microbiome to
530 increase soil health and plant fertility. *Biol. Fert. Soils*, 48, 489-499, 2012.

531 Chen, L., Jiang, Y., Liang, C., Luo, Y., Xu, Q., Han, C., Zhao, Q., Sun, B.: Competitive interaction with
532 keystone taxa induced negative priming under biochar amendments. *Microbiome*, 7, 77, 2019.

533 Coyte, K.Z., Schluter, J., Foster, K.R.: The ecology of the microbiome: networks, competition, and
534 stability. *Science*, 350, 663-666, 2015.

535 Dai, Z., Su, W., Chen, H., Barberán, A., Zhao, H., Yu, M., Yu, L., Brookes, P.C., Schadt, C.W., Chang,
536 S.X., Xu, J.: Long-term nitrogen fertilization decreases bacterial diversity and favors the growth of
537 Actinobacteria and Proteobacteria in agro-ecosystems across the globe. *Global Change Biol.*, 24,
538 3452-3461, 2018.

539 Deng, Y., Jiang, Y., Yang, Y., He, Z., Luo, F., Zhou, J.: Molecular ecological network analyses. *BMC*

540 Bioinformatics, 13, 1-20, 2012.

541 [Ding, J., Zhang, Y., Wang, M., Sun, X., Cong, J., Deng, Y., Lu, H., Yuan, T., van Nostrand, J.D., Li, D.,](#)

542 [Zhou, J., Yang, Y.: Soil organic matter quantity and quality shape microbial community](#)

543 [compositions of subtropical broadleaved forests. Mol. Ecol., 24, 5175-5185, 2015.](#)

544 Edwards, J., Johnson, C., Santos-Medellín, C., Lurie, E., Podishetty, N.K., Bhatnagar, S., Eisen, J.A.,

545 Sundaresan, V.: Structure, variation, and assembly of the root-associated microbiomes of rice. P.

546 Nat. Acad. Sci. USA, 112, E911-E920, 2015.

547 Eo, J., Park, K.: Long-term effects of imbalanced fertilization on the composition and diversity of soil

548 bacterial community. Agr. Ecosyst. Environ., 231, 176-182, 2016.

549

550 Fan, K., Delgado-Baquerizo, M., Guo, X., Wang, D., Zhu, Y., Chu, H.: Biodiversity of key-stone

551 phylotypes determines crop production in a 4-decade fertilization experiment. ISME J., 15, 550-

552 561, 2021.

553 Fath, B.D., Scharler, U.M., Ulanowicz, R.E., Hannon, B.: Ecological network analysis: network

554 construction. Ecol. Model., 208, 49-55, 2007.

555 Faust, K., Raes, J.: Microbial interactions: from networks to models. Nat. Rev. Microbiol., 10, 538-550,

556 2012.

557 Fierer, N., Bradford, M.A., Jackson, R.B.J.: Toward an ecological classification of soil bacteria. Ecology,

558 88, 1354-1364, 2007.

559 Fierer, N., Jackson, R.B.J.: The diversity and biogeography of soil bacterial communities. P. Nat. Acad.

560 Sci. USA, 103, 626-631, 2006.

561 Fierer, N.: Embracing the unknown: disentangling the complexities of the soil microbiome. Nat. Rev.

562 Microbiol., 15, 579-590, 2017.

563 [FAO: FAOSTAT Database Resources. Food and Agriculture Organization of the United Nations, Data of](#)

564 [crop production \(2021-08-01\) \[2021-03-18\]. <http://www.fao.org/faostat/en/#data/QC>. 2019.](#)

565 [Frac, M., Hannula, S.E., Bełka, M., Jędrzycka, M.: Fungal biodiversity and their role in soil health. Front.](#)

566 [Microbiol., 9, 707, 2018.](#)

567 Fuerst, J.A.: Planctomycetes—new models for microbial cells and activities. I. Kurtböke (Ed.), Microbial

568 Resources, Elsevier/Academic Press, 1-27, 2017.

569 [Grace, J., José J.S., Meir, P., Miranda, H.S., Monteset R.A.: Productivity and carbon fluxes of tropical](#)

570 [savannas. J Biogeogr., 33, 387-400, 2006.](#)

571 Hu, Q., Tan, L., Gu, S., Xiao, Y., Xiong, X., Zeng, W., Feng, K., Wei, Z., Deng, Y.: Network analysis
572 infers the wilt pathogen invasion associated with non-detrimental bacteria. *npj Biofilms Microb.*, 6,
573 1-8, 2020.

574 Huang, Z., Zhao, F., Wang, M., Qi, K., Wu, J., Zhang, S.: Soil chemical properties and geographical
575 distance exerted effects on arbuscular mycorrhizal fungal community composition in pear orchards
576 in Jiangsu Province, China. *Appl. Soil Ecol.*, 142, 18-24, 2019.

577 Hug, L.A., Castelle, C.J., Wrighton, K.C., Thomas, B.C., Sharon, I., Frischkorn, K.R., Williams, K.H.,
578 Tringe, S.G., Banfield, J.F.: Community genomic analyses constrain the distribution of metabolic
579 traits across the Chloroflexi phylum and indicate roles in sediment carbon cycling. *Microbiome*, 1,
580 1-17, 2013.

581 Jiang, Y., Luan, L., Hu, K., Liu, M., Chen, Z., Geisen, S., Chen, X., Li, H., Xu, Q., Bonkowski, M., Sun,
582 B.: Trophic interactions as determinants of the arbuscular mycorrhizal fungal community with
583 cascading plant-promoting consequences. *Microbiome*, 8, 1-14, 2020.

584 Kallenbach, C.M., Frey, S.D., Grandy, A.S.J.: Direct evidence for microbial-derived soil organic matter
585 formation and its ecophysiological controls. *Nat. Commun.*, 7, 1-10, 2016.

586 Kennedy, A.C., Smith, K.J.: Soil microbial diversity and the sustainability of agricultural soils. *Plant Soil*,
587 170, 75-86, 1995.

588 [Kolde, R., Kolde, M. Package 'pheatmap'. 2018, \[https://cran.r-\]\(https://cran.r-project.org/web/packages/pheatmap/index.html\)](#)
589 [project.org/web/packages/pheatmap/index.html](https://cran.r-project.org/web/packages/pheatmap/index.html).

590 Langfelder, P., Horvath, S.J.: Eigengene networks for studying the relationships between co-expression
591 modules. *BMC Syst. Biol.*, 1, 1-17, 2007.

592 [Liaw, A., Wiener, M.: Classification and regression by randomForest. Rnews, 2, 18-22, 2002.](#)

593 Lin, J., Sheng, B., Li, X., Yang, Q., Wang, Z., Li, H., Wang, H., Cheng, Y.: A new *pyrus pyrifolia* cultivar
594 'Sucui 1'. *Acta Hort. Sin.*, 40, 1849-1850, 2013 (In Chinese)

595 Lu, L., Yin, S., Liu, X., Zhang, W., Gua, T., Shen, Q., Qiu, H.: Fungal networks in yield-invigorating
596 and-debilitating soils induced by prolonged potato monoculture. *Soil Biol. Biochem.*, 65, 186-194,
597 2013.

598 [Magoč, T., Salzberg, S. L.: FLASH: fast length adjustment of short reads to improve genome assemblies.](#)
599 [Bioinformatics, 27, 2957-2963, 2011.](#)

600 [Mei, W., Yu, G., Lai, J., Rao, Q., Umezawa, Y.: basicTrendline: add trendline and confidence interval of](http://cran.r-project.org/package=basicTrendline)
601 [basic regression models to plot. R package version 2.0.3, 2018, http://cran.r-](http://cran.r-project.org/package=basicTrendline)
602 [project.org/package=basicTrendline.](http://cran.r-project.org/package=basicTrendline)

603 [Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P.R., O'Hara,](http://CRAN.R-project.org/package=vegan)
604 [R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H.: Package 'vegan'.](http://CRAN.R-project.org/package=vegan)
605 [Community ecology package, version 2, 2013, http://CRAN.R-project.org/package=vegan.](http://CRAN.R-project.org/package=vegan)

606 Parks, D.H., Tyson, G.W., Hugenholtz, P., Beiko, R.G.: STAMP: statistical analysis of taxonomic and
607 functional profiles. *Bioinformatics*, 30, 3123-3124, 2014.

608 Pathan, S.I., Roccotelli, A., Petrovičová, B., Romeo, M., Badagliacca, G., Monti, M., Gelsomino, A..
609 Temporal dynamics of total and active prokaryotic communities in two Mediterranean orchard soils
610 treated with solid anaerobic digestate or managed under no-tillage. *Biol. Fert. Soils.*, 57, 837-861,
611 2021.

612 Philippot, L., Raaijmakers, J.M., Lemanceau, P., van der Putten, W.H.: Going back to the roots: the
613 microbial ecology of the rhizosphere. *Nat. Rev. Microbiol.*, 11, 789-799, 2013.

614 Qiao, C., Penton, C.R., Xiong, W., Liu, C., Wang, R., Liu, Z., Xu, X., Li, R., Shen, Q.: Reshaping the
615 rhizosphere microbiome by bio-organic amendment to enhance crop yield in a maize-cabbage
616 rotation system. *Appl. Soil Ecol.*, 142, 136-146, 2019.

617 Raaijmakers, J.M., Mazzola, M.J.: Soil immune responses. *Science*, 352, 1392-1393, 2016.

618 Rosseel, Y.: Lavaan: An R package for structural equation modeling and more. *J. Stat. Softw.*, 48, 1-36,
619 2012.

620 Rousk, J., Bååth, E., Brookes, P.C., Lauber, C.L., Lozupone, C., Caporaso, J.G., Knight, R., Fierer, N.:
621 Soil bacterial and fungal communities across a pH gradient in an arable soil. *ISME J.*, 4, 1340-1351,
622 2010.

623 Schermelleh-Engel, K., Moosbrugger, H., Müller, H.J.: Evaluating the fit of structural equation models:
624 Tests of significance and descriptive goodness-of-fit measures. *Methods Psychol. Res.*, 8, 23-74,
625 2003.

626 Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., Lesniewski, R.A.,
627 Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B., Thallinger, G.G., Van Horn, D.J.,
628 Weber, C.F.: Introducing mothur: open-source, platform-independent, community-supported
629 software for describing and comparing microbial communities. *Appl. Environ. Microbiol.*, 75,

630 7537-7541, 2009.

631 Shen, Z., Penton, C.R., Lv, N., Xue, C., Yuan, X., Ruan, Y., Li, R., Shen, Q.: Banana Fusarium wilt
632 disease incidence is influenced by shifts of soil microbial communities under different monoculture
633 spans. *Microb. Ecol.*, 75, 739-750, 2018.

634 Shen, Z., Zhong, S., Wang, Y., Wang, B., Mei, X., Li, R., Ruan, Y., Shen, Q.: Induced soil microbial
635 suppression of banana fusarium wilt disease using compost and biofertilizers to improve yield and
636 quality. *Eur. J. Soil Biol.*, 57, 1-8, 2013.

637 [Smoot, M.E., Ono, K., Ruscheinski, J., Wang, P.L., Ideker, T.: Cytoscape 2.8: new features for data
638 integration and network visualization. *Bioinformatics*, 27, 431-432, 2011.](#)

639 Speirs, L.B.M., Rice, D.T.F., Petrovski, S., Seviour, R.J.: The phylogeny, biodiversity, and ecology of the
640 *Chloroflexi* in activated sludge. *Front Microbiol.*, 10, 2015, 2019.

641 Tian, J., He, N., Hale, L., Niu, S., Yu, G., Liu, Y., Blagodatskaya, E., Kuzyakov, Y., Gao, Q., Zhou, J.:
642 Soil organic matter availability and climate drive latitudinal patterns in bacterial diversity from
643 tropical to cold temperate forests. *Funct. Ecol.*, 32, 61-70, 2018.

644 [Toju, H., Peay, K.G., Yamamichi, M. Narisawa, K., Hiruma K., Naito K., Fukuda S., Ushio M., Nakaoka
645 S., Onoda Y., Yoshida K., Schlaeppi K., Bai Y., Sugiura R., Ichihashi Y., Minamisawa K., Kiers &
646 E.T.: Core microbiomes for sustainable agroecosystems. *Nat Plants*, 4, 247–257, 2018.](#)

647 ~~Ullah, A., Nisar, M., Ali, H., Hazrat, A., Hayat, K., Keerio, A.A., Ihsan, M., Laiq, M., Ullah, S., Fahad,
648 S., Kan A., Khan, A.H., Akbar, A., Yang, X.: Drought tolerance improvement in plants: an
649 endophytic bacterial approach. *Appl. Environ. Microbiol.*, 103, 7385-7397, 2019.~~

650 van der Heijden, M.G., Bardgett, R.D., van Straalen, N.M.J.: The unseen majority: soil microbes as
651 drivers of plant diversity and productivity in terrestrial ecosystems. *Ecol. Lett.*, 11, 296-310, 2008.

652 Wang, C., Liu, D., Bai, E.: Decreasing soil microbial diversity is associated with decreasing microbial
653 biomass under nitrogen addition. *Soil Biol. Biochem.*, 120, 126-133, 2018.

654 Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R.: Naive Bayesian classifier for rapid assignment of rRNA
655 sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.*, 73, 5261-5267, 2007.

656

657 Xun, W., Li, W., Xiong, W., Ren, Y., Liu, Y., Miao, Y., Xu, Z., Zhang, N., Shen, Q., Zhang, R.: Diversity-
658 triggered deterministic bacterial assembly constrains community functions. *Nat. Commun.*, 10, 1-
659 10, 2019.

660 Zhao, J., Ni, T., Li, Y., Xiong, W., Ran, W., Shen, B., Shen, Q., Zhang, R.: Responses of bacterial
661 communities in arable soils in a rice-wheat cropping system to different fertilizer regimes and
662 sampling times. *PloS One*, 9, e85301, 2014.


663 Zhong, Y., Hu, J., Xi, Q., Zhang, S., Li, X., Pan, X., Zhao, R., Wang, R., Yang, W., Shangguan, Z., Hu,
664 F., Yang, C., Wang, W.. Soil microbial mechanisms promoting ultrahigh rice yield. *Soil Biol.*
665 *Biochem.*, 143: 107741, 2020.

666 ~~Zorz, J.K., Sharp, C., Kleiner, M., Gordon, P.M.K., Pon, R.T., Dong, X., Strous, M.: A shared core~~
667 ~~microbiome in soda lakes separated by large distances. *Nat. Commun.*, 10, 4230, 2019.~~


668

669 **Figure legends**

670 **Fig. 1 Distribution of studied field sites, ~~quantitation of the abundance of bacteria population, and~~**
671 **~~linkage of microbial composition to pear yield.~~ (A) Map showing the sites of six pair-located orchards**
672 **sampled in this study. (B) Violin plot showing the abundance of total bacteria for all selected orchards. ^{*}**
673 **~~indicates a significant difference between yield invigorating (YI) and yield debilitating (YD) orchards~~**
674 **~~based on Wilcoxon tests ($p < 0.05$).~~ (C) Correlation plot showing the relationship of microbial**
675 **~~composition and yield based on braycurtis distances calculated by Mantel test.~~**



676



677

678


679 **Fig. 2 Quantitation of the abundance of bacteria population, and linkage of microbial composition**

680 **to pear yield.** (A) Violin plot showing the abundance of total bacteria for all selected orchards. * indicates


681 a significant difference between yield-invigorating (YI) and yield-debilitating (YD) orchards based on

682 Wilcoxon tests ($p < 0.05$). (B) Correlation plot showing the relationship of microbial composition and

683 yield based on braycurtis distances calculated by Mantel test.




684







685

|686
687

688 **Fig. _23 Overview of bacterial composition and alpha diversity.** (A) Principal Coordinates Analysis
 689 (PCoA) plot displaying the bacterial community composition calculated based on braycurtis distances.
 690 (B) Violin plot showing the alpha diversity indices (Chao and Shannon) for all selected orchards. *
 691 indicates a significant difference between yield-invigorating (YI) and yield-debilitating (YD) orchards
 692 based on Wilcoxon tests ($p < 0.05$). (C) Venn plot depicting the unique and shared bacterial ASVs
 693 between yield-invigorating (YI) and yield-debilitating (YD) orchards at ASV and sequence insights.
 694 Uni. YI and Uni. YD represent unique ASVs or sequences in the YI or YD soils while Shared represent
 695 shared ASVs or sequences between the YI and YD soils. (D) Heatmap displaying the composition of
 696 unique and shared ASVs at phylum level in YI and YD soils. Numbers in the cell represent the number
 697 of ASVs affiliated to that phylum.




698




699

700

701 **Fig. 3.4** Key taxonomic groups in distinguishing yield-invigorating (YI) and yield-debilitating (YD) orchards.
 702 **orchards.** (A)-Stacked bar chart (left panel) showing dominant phyla affiliation in YI and YD soils for
 703 six pair-located sites while horizontal histogram (right panel) depicting relative changes of dominant
 704 phyla in YI soils compared to those in YD soils. (B) Genus level taxonomic analysis of bacterial
 705 sequences obtained from yield-invigorating (YI) and yield-debilitating (YD) orchards using the STAMP
 706 software. Cyan bars represent the YI soils and pink bars represent the YD soils.






708


709

710

711 **Fig. 4-5** Co-occurrence networks of bacterial community and identified keystone taxa in
 712 **distinguishing yield-invigorating (YI) and yield-debilitating (YD) orchards.** (A) An overview of
 713 microbial phylogenetic molecular ecological networks constructed from YI and YD soils. Line with blue
 714 color indicates positive correlations whereas lines with red color signifies negative correlations in each
 715 network. Modules containing larger than five nodes in the networks are labeled with corresponding letter
 716 followed by a number. Circular node colors indicate different bacterial phyla. (B) Stacked figure showing
 717 Bubble graph showing the relative abundance of nodes in each module within each network at the phylum
 718 level. (C) Venn plot depicting the unique and shared bacterial ASVs between two networks construed
 719 from YI and YD soils. Left panel is plotted based on the original nodes used in building network while
 720 right panel is plotted based on the nodes from modules. Uni. YI and Uni. YD represent unique ASVs in
 721 the YI or YD networks while Shared represent shared ASVs between the YI and YD networks. (D) Zi-Pi
 722 plot showing the distribution of nodes based on their topological roles. The threshold values of Zi and Pi
 723 for categorizing OTUs were 2.5 and 0.62 respectively. Node colors indicate different bacterial phyla and
 724 node size represent the relative abundance in each network.




725







726

727

728 **Fig. 5-6 Relationships among bacterial community, soil edaphic factors and pear yield.** (A)
729 Variance partitioning analysis (VPA) map of the effects of soil edaphic properties, sample locations,
730 pear yield and interactions of these factors on the bacterial community. (B) Redundancy analysis
731 (RDA) plot showing the relationships among all assigned bacterial ASVs and measured soil edaphic
732 properties for all soils after stepwise selection. (C) Random forest mean predictor importance of
733 selected soil edaphic properties used ~~in the~~ as drivers in predicting the pear yield. Red bar indicates that
734 the given predictor is significant while black bar indicates that the given predictor is non-significant.
735 The %IncMSE in the up panel means the increase in mean squared error while IncNodePurity indicates
736 the increase in node purity. The values of these two indices represent the importance of each variable to
737 predict the module. A larger value indicates that the variable is more important. (D) Correlation plot
738 showing the relationship of microbial composition and soil organic matter based on braycurtis
739 distances calculated by Mantel test.




740




741

742

743 **Fig. 6-7** Structural equation modeling (SEM) describing the biotic and abiotic factors in affecting
744 **the crop production.** Structural equation model was built incorporating soil organic matter, microbial
745 biomass, beta diversity (PCoA), key taxa, network hubs including module hubs and network
746 connectors, and yield. The path analysis numbers adjacent to arrows indicate the relationship's effect
747 size and the associated bootstrap *P*-value. Cyan and red arrows indicate positive and negative
748 relationships, respectively. Paths with non-significant coefficients are presented as gray lines.



749



750

751