

30 invigorating soils exhibited higher contents of organic matter than yield-debilitating soils and harboured
31 unique bacterial communities. Greater alpha diversity and higher relative abundances of Planctomycetota
32 and Chloroflexota were observed in yield-debilitating soils. Co-occurrence network analysis revealed
33 that yield-invigorating soils displayed a greater number of functionally interrelated modules (meta-
34 modules) and a higher proportion of negative links to positive links. Chloroflexota was recognized as a
35 keystone taxon in manipulating the interaction of bacterial communities in yield-invigorating soils.
36 ~~Structural equation modelling showed that soil organic matter, beta diversity of bacterial community, and~~
37 ~~network connector (Chloroflexota) were identified as potential key factors in explaining the high yield~~
38 ~~pear production.~~ Altogether, we provide evidence that yield-invigorating soils across a range of locations
39 appear to share common features, including accumulation of soil organic matter, higher microbial
40 diversity, enrichment of key taxa like Chloroflexota, and maintaining a competitive network. These
41 findings have implications for science-based guidance for sustainable food production.

42

43 **Keywords:** Soil organic matter, Microbial diversity, Random forest prediction, Co-occurrence network,
44 Keystone taxa

45

46 **1 Introduction**

47 Soils are essential to human wellbeing due to their great contributions to the production of food, fiber,
48 feed, and medicine (Raaijmakers and Mazzola, 2016). Soil organisms play critical roles in maintaining
49 these ecosystem services, such as driving nutrient cycling, maintaining soil fertility, improving plant
50 productivity and suppressing plant diseases (Bender et al., 2016; Barrios, 2007). Microorganisms
51 participate in nearly all soil biological processes, and the microbial abundance, community composition
52 and activity primarily determine the sustainable productivity of agricultural lands (Philippot et al., 2013).
53 Fungi participate in decomposition of organic matter and deliver nutrients for plant growth (Frąc et al.,
54 2018), however, considering that bacteria are the most diverse and abundant group of microorganisms in
55 soil, bacterial communities and their functions can be pivotal indicators for crop production in
56 agroecosystems (van der Heijden et al., 2008).

57 In general, an increase in microbial diversity is linked to a high-yielding crop production mainly
58 through improving the host resilience to physical or chemical disturbances, modifying plant competition,
59 and facilitating plant access to nutrients (Chaer et al., 2009; Kennedy and Smith, 1995). Since individual

60 organisms do not live in isolation but rather form a complex system of inter-species interactions in soil,
61 interactions among community members were found to be related to crop production in the potato
62 monoculture system (Lu et al., 2013). Enrichments of key functional microbes in soil were deemed to
63 serve specific soil system functions, such as suppressing soil-borne pathogens and maintaining
64 sustainable crop production (Banerjee et al., 2018). However, the relative contributions of microbial
65 diversity, interactions among community members, or enrichment of key taxa to crop production remain
66 largely unknown. Therefore, it is highly desirable to identify pivotal indicators of bacterial community
67 composition in response to high-yielding crop production.

68 Changes in composition of soil bacterial communities across space are often strongly correlated
69 with soil pH (Fierer and Jackson, 2006). Soil pH has been recognized as a key driver in determining the
70 assembly of bacterial community in arable soils by field ~~and microcosms~~ experiments (Rousk et al.,
71 2010). However, recent studies have demonstrated that compositions of soil bacterial communities were
72 driven by a myriad of soil abiotic traits, such as organic matter contents, forms and contents of soil
73 nutrient (Tian et al., 2018; Wang et al., 2018). For example, soil bacterial community composition, which
74 determines the ability of soil to suppress soil-borne pathogens, was found to be strongly correlated with
75 soil organic matter (Shen et al., 2018). An imbalanced ratio of soil nutrients, *i.e.*, ratio of nitrogen to
76 phosphorus or potassium could be a driving force altering the bacterial community composition in long-
77 term fertilized soils (Eo and Park, 2016). Key soil chemical properties identified in controlling the
78 distribution and abundance of bacterial community is largely depending on the geographical distributions
79 of soils. As a consequence, a better understanding of the relationship between soil edaphic properties and
80 bacterial community composition is critical to develop targeted manipulation options to increase soil
81 service provisions.

82 'Sucui No. 1' pear, is an early-maturing variety bred by the Jiangsu Academy of Agricultural
83 Sciences, China, and has been popularly cultivated in Eastern and Central China, due to the advantages
84 including easy to produce, adaptable to the environment, and has good quality and high economic
85 benefits (Lin et al., 2013). With the increasing demand in China, sustainable production of high-quality
86 pear is becoming increasingly important. Manipulation of soil microbiomes has shown to be an effective
87 way to increase soil productivity (Chaparro et al., 2012). Considering that large-scale surveys could
88 exhibit the diversity of soil microbial communities exceeds what is found in host-associated communities
89 (Toju et al., 2018), it is necessary to explore the general microbial characteristics of multiple yield-

90 invigorating soils and identify key environmental drivers in assembling bacterial community.

91 In this study, ~~compared to local average yield~~, orchard ~~showing with~~ higher pear yield production
92 ~~compared to local average yield~~ was recognized as yield-invigorating (YI) orchard while orchard
93 ~~displaying having~~ lower pear yield production ~~in comparison with local average yield~~ was regarded as
94 yield-debilitating orchard (YD). After field surveys accomplished in 2019, six ~~separately located~~ yield-
95 invigorating ~~pear orchards~~ and adjacent yield-debilitating pear orchards ~~in total~~ were selected for further
96 analysis of soil chemical properties and microbiome. We hypothesized that high input of organic fertilizer
97 could improve soil structure and modify ~~soil~~ chemical properties, which ~~leading leads~~ to YI soils
98 harbor~~ing~~ unique bacterial communities ~~that maintains associated with the~~ high-yielding pear production.
99 To address this, soil bacterial communities and edaphic properties of the study sites were compared to
100 (1) decipher the differences of taxonomic diversity, and composition of the bacterial community, and (2)
101 determine the contributions of environmental variables to the changes in the structure of bacterial
102 communities.

103 2 Methods

104 2.1 Study sites and experimental design

105 From July - August 2019, a field production survey of orchards cultivated with ‘Suci No. 1’ pear was
106 performed after pear fruits harvest to compare the differences of soil nutrients and microbiota between
107 yield-invigorating with yield-debilitating orchards. The locations, planting density, cropping years, soil
108 type and total yield were recorded. To minimize the effects of microclimate at each site, only pair-located
109 pear orchards with invigorating and debilitating yield and at similar growth stage were selected for this
110 research. In total, six pair-located yield-invigorating and -debilitating pear orchards distributed in four
111 cities of Jiangsu province, China, were selected in the main pear production areas (Fig. 1, Table S1). The
112 yield per tree was obtained by dividing the total yield per hectare by plant density.

113 **Fig. 1 here**

114 Paired yield-invigorating and yield-debilitating orchards from Fengxian (FX), Suining (SN) and
115 Tongshan (TS) were maintained in the Xuzhou city under the warm temperate sub-humid monsoon
116 climate. This site has a mean annual temperature (MAT) of 14.5 °C and mean annual precipitation (MAP)
117 of 847 mm. Orchards from Taixing (TX) were located in the Taizhou city under the humid southern
118 subtropical climate with a MAT of 15.3 °C and MAP of 1055 mm. Orchards from Gaochun (GC) were
119 located in the Nanjing city under the humid subtropical monsoon climate with a MAT of 15.4 °C and

120 MAP of 1106 mm. Orchards from Zhangjiagang (ZJ) were located in the Suzhou city under the humid
121 subtropical monsoon climate with a MAT of 15.7 °C and MAP of 1094 mm. For paired yield-invigorating
122 and-debilitating orchards, the irrigation and pesticide management practices were similar according to
123 farm records. However, yield-invigorating orchard was usually amended with more organic fertilizer
124 under integrated nutrients management whereas the co-located yield-debilitating orchard received more
125 chemical fertilizer under intensive management. Detailed information about fertilization regimes for each
126 orchard is shown in Table S2.

127 2.2 Soil sample collection and chemical properties determination

128 Along with the field survey, soil sampling campaigns were performed from July - August 2019 after pear
129 fruits harvest. For each yield-invigorating or -debilitating orchard, four subplots with three pear trees in
130 each subplot were randomly selected for soil sampling. Subsequently three soil cores (0-20 cm) under
131 the trunk base for each tree were collected using a 25 mm soil auger. In total, nine soil cores for each
132 subplot were pooled as a composite sample and finally four composite soil samples for each orchard
133 were collected and promptly transported on ice to the laboratory. After sifting through a 2 mm sieve and
134 thoroughly mixing, one portion of each soil sample was air-dried for chemical property analyses while
135 the remainder was stored at -70 °C for DNA extraction.

136 Soil chemical properties including soil pH, content of organic matter (OM), total nitrogen (TN),
137 available phosphorus (AP), available potassium (AK), alkali-hydrolyzable nitrogen (N), exchangeable
138 calcium (Ca), effective magnesium (Mg), effective iron (Fe), effective manganese (Mn), effective copper
139 (Cu) and effective zinc (Zn), were measured according to methods described by Shen et al. (2018) and
140 Huang et al. (2019). Briefly, soil pH was determined using a glass electrode meter in a suspension with
141 a 1:5 soil/water ratio (w/v). Soil OM was determined using the potassium dichromate external heating
142 method. TN was determined using a dry combustion method on an Element Analyzer (Vario EL,
143 Germany). AP and AK were determined using the molybdenum blue method after soil was extracted with
144 sodium bicarbonate and flame photometry after soil was extracted with ammonium acetate, respectively.
145 Soil alkaline hydrolysable nitrogen (N) was measured by the alkaline hydrolysable diffusion method.
146 Contents of soil Ca, Mg, Fe, Mn, Cu and Zn were determined by the atomic absorption spectroscopy
147 method using ICE 3300 AAS Atomic Absorption Spectrometer (ThermoScientific, USA) after acid
148 hydrolysis.

149 2.3 Soil DNA extraction and bacterial abundance quantification

150 Genomic DNA from 0.25 g soil for each sample was extracted by using the DNeasy® PowerSoil® Kit
151 (QIAGEN GmbH, Germany) according to the manufacturer's instructions. The abundances of soil
152 bacteria were determined with Eub338F/Eub518R primer using a 7500 Real Time PCR System (Applied
153 Biosystems, USA). Standard curves were generated by using 10-fold serial dilutions of a plasmid
154 containing a full-length copy of the 16S rRNA gene from *Escherichia coli*. Quantitative PCR analysis
155 was performed in 96-well plates with a 20- μ l mixture for each reaction using SYBR®Premix Ex Taq™
156 (TaKaRa, Japan). Thermal cycling was conducted according to a standard procedure with three replicates,
157 and the results were expressed as log copy numbers g⁻¹ dry soil.

158 2.4 Sequencing library construction and sequencing

159 The gene-specific primers 515F/806R with 12 bp barcode were used to amplify the V4 region of bacterial
160 16S rRNA gene on the BioRad S1000 (Bio-Rad Laboratory, CA) according to the protocols described
161 by Caporaso et al. (2011). All constructed libraries were sequenced using the Illumina NovaSeq 6000 at
162 the Guangdong Magigene Biotechnology Co., Ltd. (Guangzhou, China).

163 2.5 Sequence processing

164 Quality filtering of the paired-end raw reads was performed to obtain the high-quality clean reads
165 according to the Trimmomatic V0.33 (Bolger et al., 2014) quality control process. Sequences were
166 assigned to each sample based on their unique barcode, after which the barcodes and primers were
167 removed. Paired-end clean reads were merged using FLASH V1.2.11 (Magoč and Salzberg, 2011). Raw
168 tags were processed to generate the final ~~ASV~~ (Amplicon Sequence Variant (ASV)) table file at 97%
169 pairwise identity according to the QIIME2 pipeline (Bolyen et al., 2019). The nonbacterial and
170 mitochondrial ASVs and extremely low frequency ASVs (relative abundance < 0.01%) were removed.
171 A representative sequence for each ASV was selected and classified using the RDP classifier against the
172 RDP Bacterial 16S database (Wang et al., 2007).

173 2.6 Statistical analyses

174 Statistical analyses were performed using the software SPSS 20.0 (SPSS Technologies, Armonk, NY,
175 USA) and R (<http://www.R-project.org/>). Non-normal data were square-root or log transformed when
176 necessary. The significance of soil properties or microbial taxa in yield-invigorating or-debilitating
177 orchards was determined based on paired Wilcoxon rank sum test, and adjusted *P* values (< 0.05) were
178 obtained by the FDR method. Mantel tests were used to identify the correlations between microbial
179 community composition and pear yield, and soil chemical properties using the 'vegan' package (Oksanen

180 et al., 2013) in R. The linear regression analyses relating yield to selected microbial taxa or soil chemical
181 properties were conducted using the ‘basicTrendline’ package (Mei et al., 2018) in R.

182 Principal Coordinate Analysis (PCoA) based on the Bray-Curtis distance was performed in
183 MOTHUR V1.38.1 (Schloss et al., 2009) and visualized by the ‘ggplot2’ package (Wickham and Chang,
184 2015) in R to explore the differences in microbial community composition. Permutational multivariate
185 analysis of variance (PERMANOVA) was performed to evaluate the significant differences of microbial
186 community composition according to sample locations and orchard yield using the ‘vegan’ package in R.
187 Microbial alpha diversity indexes (Chao, Shannon) were calculated based on randomly resampled ASV
188 abundance matrices at the same depth (23,800 sequences) in MOTHUR. A Venn diagram was generated
189 based on the final ASVs to compare microbial community composition between yield-invigorating and
190 -debilitating orchard soils. The affiliations of unique and shared ASVs in yield-invigorating and -
191 debilitating soils were compared to evaluate the differences of the bacterial community composition and
192 plotted using the ‘pheatmap’ package (<https://cran.r-project.org/web/packages/pheatmap>) in R. Fold
193 changes (log₂ transformed) of shared ASVs across yield-invigorating and -debilitating soils were
194 calculated. The ASVs with fold change ratios > 2 and unique ASVs in yield-invigorating soils were
195 recognized as potential responders to yield promotion. In addition, to better understand the bacterial
196 community composition, relative abundances at the genus level were compared by STAMP software
197 v2.1.3 (Parks et al., 2014).

198 The phylogenetic molecular ecological networks (pMEN) were constructed using the random
199 matrix theory-based approach to explore the organization of bacterial communities in yield-invigorating
200 (YI) or yield-debilitating (YD) soil samples. Potential ecological interactions among bacteria were
201 determined by modeling the microbial community using Molecular Ecological Network Analysis
202 (<http://ieg2.ou.edu/MENA>) based on pear yield (Deng et al., 2012). Given the large number of rare taxa
203 that are specific to certain locations, ASVs that occurred in less than half of soil samples and lower than
204 0.01% were filtered, which resulting in 591 and 485 ASVs for YI and YD samples respectively, before
205 networks constructed. The microbial network was constructed using random matrix theory-based at
206 0.94 similarity threshold and visualized using Cytoscape 2.8.3 software (Smoot et al., 2011). Module
207 clustering and composition in yield-invigorating and -debilitating networks were compared and plotted
208 in R using the ‘pheatmap’ and ‘ggplot2’ packages. And the threshold values of Z_i and P_i was 2.5 and
209 0.62, respectively for topology analysis of the network. Redundancy analysis (RDA) was performed in

210 the R ‘vegan’ package to examine the relationship among frequencies of ASVs, samples and soil
211 variables, which were selected using ‘stepAIC’ in R. Variance partitioning analysis (VPA) was used to
212 determine the contributions of soil properties, sample location, and yield, as well as interactions among
213 the variation in a microbial community with hellinger-transformed data. The predictors of selected soil
214 properties for explaining the pear yield were identified by random forest regression analysis (Boulesteix,
215 et al., 2012). The significance of each predictor in the response variables was assessed with the
216 ‘rfPermute’ package (Liaw and Wiener, 2002) with 1000 permutations based on 1000 trees. ~~Structural~~
217 ~~equation modelling was applied to evaluate relative contributions of soil chemical properties and~~
218 ~~bacterial community to pear yield (Grace et al., 2006; Schermelleh-Engel et al., 2003). The conceptual~~
219 ~~SEM fitness was examined on the basis of a non-significant chi-square test ($P > 0.05$) and the goodness-~~
220 ~~of fit index (GFI). Model was fitted using the ‘lavaan’ package in R (Rosseel, 2012).~~

221 **3 Results**

222 3.1 Overview of sequencing data

223 In total, 1,622,858 16S rRNA sequences were retained after quality control and a total of 9,394 ASVs
224 were obtained for the 16S rRNA gene sequences based on 97% similarity. Among the total 16S rRNA
225 gene sequences, 159 ASVs with 74,372 sequences were classified as Archaea while 9,235 ASVs with
226 1,548,486 sequences were identified as Bacteria. Among Bacteria, Acidobacteriota, Pseudomonadota,
227 Chloroflexota, Planctomycetota and Actinomycetota were the most abundant phyla (Fig. S1).

228 3.2 Soil chemical properties

229 Soil chemical properties differed ~~significantly~~ among the locations and orchard yield types (Table S3).
230 On average, yield-invigorating orchards showed obviously higher contents of OM, AP, Mg and Fe, and
231 a lower content of Mn in comparison with those in yield-debilitating orchards. However, when taken all
232 sites together, only a higher relative abundance of ~~soil organic matter (OM)~~ on average was observed in
233 yield-invigorating orchards compared to that in yield-debilitating orchards based on the Wilcoxon test
234 ($P < 0.05$).

235 3.3 Bacterial abundances and community compositions

236 Yield-invigorating orchards together displayed significantly higher abundances of total bacteria than that
237 in co-located yield-debilitating orchards based on real time PCR result (Fig. 2A). Meanwhile, bacterial
238 community compositions at the ASV level were significantly correlated to pear yield ($r = 0.460$, $p =$
239 0.001) (Fig. 2B).

240

Fig. 2 here

241 PCoA based on Bray-Curtis distance matrices clearly revealed location-based differences in
242 bacterial community compositions (Fig. 3A). Six distinct groups representing samples from different
243 locations (FX, GC, SN, TS, TX and ZJ) were obviously separated and confirmed by PERMANOVA test
244 ($F = 14.9$, $P = 0.001$). At each location, soil bacterial community composition in yield-invigorating
245 orchards was significantly separated from that in co-located yield-debilitating orchards, which was also
246 confirmed by PERMANOVA test ($F = 3.6$, $P = 0.001$). Although only the Shannon diversity in yield-
247 invigorating orchards from GC and ZJ was significantly higher than that in co-located yield-debilitating
248 orchards (Fig. S2), the mean alpha diversity indices of Chao and Shannon in all yield-invigorating
249 orchards were significantly higher than those in all yield-debilitating orchards based on the paired
250 Wilcoxon test (Fig. 3B).

251

Fig. 3 here

252 The Venn diagram showed that 4,540 ASVs occupying over 90% of total sequences were shared
253 between yield-invigorating and -debilitating orchards (Fig. 3C). Among these shared ASVs, the fold
254 changes larger than 2 of ASVs in yield-invigorating compared to yield-debilitating orchards were
255 recognized as potential responders linking to yield improvement. Surprisingly, none of these ASVs
256 potentially linked to yield improvement were shared among six separated collocated orchards (Fig. S3).
257 A total of 2,546 unique ASVs with 53,222 sequences were found in all yield-invigorating orchards while
258 2,308 unique ASVs with 44,389 sequences were observed in all yield-debilitating orchards. Among
259 these unique ASVs, almost 70% of ASVs were shared between yield-invigorating orchards and -
260 debilitating orchards. However, no shared unique ASVs were found among six separately located
261 orchards. The affiliation of unique and shared ASVs at the phylum level exhibited that the
262 Pseudomonadota, Planctomycetota, Chloroflexota, Acidobacteriota and Actinomycetota were the top
263 five phyla (Fig. 3D).

264

265 At the phylum level, the relative abundances of bacterial dominant phyla varied across the location
266 and orchard yield condition. Pseudomonadota, Acidobacteriota, Actinomycetota, Chloroflexota and
267 Planctomycetota were the top five abundant phyla (Fig. 4). The mean abundance of Chloroflexota and
268 Planctomycetota was significantly higher while Firmicutes was significantly lower in yield-invigorating
269 orchards compared to yield-debilitating orchards based on Wilcoxon test (Fig. S4).

269

At a finer resolution, 967 genera were observed for all soil samples, among which 299 genera

270 appeared in more than half of soil samples in yield-invigorating or -debilitating orchards. However, only
271 34 genera displayed significant differences between yield-invigorating or -debilitating orchard soils
272 based on STAMP analysis (Fig. S5). Interestingly, *Ornatilinea*, *Ktedonobacter*, *Longilinea*, belonging
273 to Chloroflexota, were significantly enriched in yield-invigorating orchard soils. *Gimesia* in
274 Planctomycetota and *Arenimonas* in Pseudomonadota showed significantly higher relative abundances
275 in yield-invigorating orchard soils than in yield-debilitating orchard soils.

276 **Fig. 4 here**

277 3.4 Co-occurrence patterns of bacterial community

278 The yield-invigorating network contained 302 nodes, 448 edges, and 11 larger modules (> 5 nodes), with
279 an average connectivity (avgK) of 2.967, average path distance of 5.494 and clustering coefficient
280 (avgCC) of 0.152, while the values in the yield-debilitating network were 235, 334, 9, 2.843, 6.232 and
281 0.131, respectively (Fig. 5A, Table S4). The module eigengene network analysis revealed a difference in
282 the higher-order organization between the two networks. Notably, the node composition was substantially
283 different between the two networks as the relative abundances of dominant phyla were obviously
284 different among different modules (Fig. 5B and C). A higher proportion of nodes in the module of yield-
285 invigorating network was unique. ASVs affiliated to Acidobacteriota, Chloroflexota, Pseudomonadota,
286 Actinomycetota, and Planctomycetota within the unique modules (M9, M10 and M11) were observed in
287 the yield-invigorating versus yield-debilitating network.

288 Analysis using the threshold values of Z_i (within-module connectivity) and P_i (among-module
289 connectivity) showed that majority of nodes from both networks were categorized as peripherals that had
290 only a few links and almost always linked to the nodes within their own modules (Fig. 5D). Although
291 only three nodes affiliated to Acidobacteriota were categorized as module hubs in the yield-invigorating
292 network, seven nodes belonging to Acidobacteriota, Actinomycetota and Pseudomonadota were
293 categorized as module hubs in the yield-debilitating network. Interestingly, four nodes including
294 *Longilinea* species from Chloroflexota in the yield-invigorating network whereas only one node in the
295 yield-debilitating network was categorized as module connectors (Table S5).

296 **Fig. 5 here**

297 3.5 Relationships between soil chemical properties and microbial community composition

298 Soil chemical properties were significantly correlated to the bacterial community compositions (Mantel:
299 $r = 0.803$, $p = 0.001$). Soil chemical properties, location, and orchard explained 44.9% of the observed

300 variation, leaving 55.1% of the variation unexplained for bacterial community composition based on
301 VPA result (Fig. 6A). Variation in the community composition was largely explained by soil properties
302 (42.3%), and was also influenced by locations and orchard yield types.

303 After forward stepwise selection, the module including soil OM, TN, alkaline N, AP and AK,
304 available calcium (Ca), copper (Cu) and manganese (Mn) explained the majority of the variation in
305 bacterial community composition (Fig. 6B). As evidenced by the RDA vectors, OM within the module
306 was identified as the top important soil property that determines the composition of bacterial community.
307 Random forest analysis showed that contents of soil Mn, OM and Ca were the top parameters for
308 predicting the orchard yield (Fig. 6C). Furthermore, soil OM was also significantly correlated with
309 bacterial communities as revealed by Mantel test (Fig. 6D, Table S6).

310 **Fig. 6 here**

311 3.6 Relationships of soil chemical and microbial indicators with orchard yield

312 Soil OM as potentially key soil chemical properties and bacterial alpha diversity, beta diversity and
313 relative abundance of Chloroflexota and Planctomycetota as potentially key microbial indicators
314 associated with pear yield were used to construct a model to explain yield improvement. Final structural
315 equation modelling (path analysis) (Fig. 7 and S6) showed that the strongest driver explaining yield
316 improvement was beta diversity of bacterial community (PCoA) ($r = 0.959$, $P < 0.001$), which was
317 positively affected by content of soil OM ($r = 0.843$, $P < 0.001$). Alpha diversity (Chao) of bacterial
318 community also determined yield improvement to a large extent ($r = 0.542$, $P = 0.009$). However, alpha
319 diversity was not significantly correlated with content of soil OM.

320 **Fig. 7 here**

321 **4 Discussion**

322 Although pear is among the most important fruits worldwide, soil microbial communities in pear
323 orchards have been largely under-investigated (Huang et al., 2019). The present study attempts to
324 decipher the bacterial community linked to high-yield production of pear. Our results based on Mantel
325 analysis suggested significant correlations among bacterial community, soil chemical properties and pear
326 yield. Microbial characteristics responding to yield promotion have repeatedly been observed on several
327 crops depending on single experimental site (Zhong et al., 2020; Qiao et al., 2019; Shen et al., 2013). It
328 remained unclear, however, whether these distinctions are ubiquitous at a large-scale. By comparing
329 multiple co-located yield-invigorating and -debilitating orchards, we demonstrate that high-yielding pear

330 production soils exhibited high organic matter contents and harbored bacterial communities with high
331 diversity, significantly enriched indigenous microbes and more interactive network, which was triggered
332 by high-inputs of soil organic fertilizer. Here we discussed these main results and potential mechanisms
333 in detail.

334 Microbial diversity is critical to soil ecosystems in maintaining the integrity, function and long-term
335 sustainability (Kennedy and Smith, 1995). Higher soil biodiversity is considered to be linked to a more
336 stable system and enhance the combination of vital microbial functions and processes (Cardinale et al.,
337 2006; Bell et al., 2005). In line with a previous report that crop yield was correlated to the soil bacterial
338 diversity (Zhao et al., 2014), greater diversity of bacterial community in yield-invigorating soils was
339 observed in the present study. Hence we infer that higher microbial diversity may result in a more
340 productive agroecosystem, contributing to sustainable pear production.

341 In this study, we found that Pseudomonadota, Acidobacteriota, Actinomycetota, Planctomycetota
342 and Chloroflexota were the top abundant phyla. This result roughly agreed with previous studies showing
343 that Pseudomonadota, Acidobacteriota and Actinomycetota are usually dominant bacterial taxa in
344 agricultural soils (Xun et al., 2019; Dai et al., 2018), while Planctomycetota and Chloroflexota exhibit
345 an unexpectedly high relative abundance in rice cropped soil (Edwards et al., 2015) and sandy loam soil
346 (Pathan et al., 2021). The highest relative abundance of Pseudomonadota was probably explained by the
347 fact that Pseudomonadota are considered as copiotrophic bacteria and flourish in soils with large amounts
348 of available nutrients (Fierer et al., 2007).

349 Moreover, a significantly higher abundance of Planctomycetota and Chloroflexota was observed in
350 yield-invigorating orchards, indicating that Planctomycetota and Chloroflexota may be associated with
351 pear yield-improvement. There is no direct evidence showing that Planctomycetota could improve plant
352 growth. However, Planctomycetota has been reported to be involved in many soil biological processes
353 such as ammonification, carbohydrate and polysaccharide metabolism (Fuerst, 2017). This implies that
354 Planctomycetota may promote plant production through improving soil fertility. Chloroflexota is a
355 facultative anaerobic phylum including autotrophic, heterotrophic and mixotrophic taxa (Speirs et al.,
356 2019). Considering that soil amended with organic fertilizer may enhance the soil water holding capacity,
357 the yield-invigorating soils with more organic material input have a higher soil moisture content,
358 especially after irrigation, probably leading to the enrichment of Chloroflexota in soil.

359 Network analysis is a systems-level method to explore interactions within an ecosystem that cannot

360 be directly observed through co-occurrence analysis (Fath et al., 2007). Similar to the food web network
361 analyses in macro ecosystems, microorganisms also form complex interactions with other species (Faust
362 and Raes, 2012) and have been widely investigated to explore the linkage of microbial network with soil
363 function, such as nutrient supply (Fan et al., 2021) and disease suppression (Lu et al., 2013). Overall, in
364 line with previous findings (Hu et al., 2020), the topological properties of the constructed networks,
365 including connectivity, average clustering coefficients, average degree distance, and modularity indicate
366 that these networks are scale-free, modular and small world. In short, a scale-free network represents that
367 a network whose connectivity follows a power law, and most of nodes have only a few connections with
368 other nodes. Meanwhile, a small-world network is the network in which most nodes are not neighbors of
369 one another, but most nodes can be reached by a few paths. Modularity is a fundamental characteristic
370 of biological network as a module in the network is a group of nodes that are highly connected within
371 the group, but very few connections outside the group (Deng et al., 2012). Our comparative network
372 analysis indicated that microbial co-occurrence patterns in soils links to different pear production. As a
373 meta-module is usually considered as a group of modules functionally interrelated (Langfelder and
374 Horvath, 2007), a greater number of meta-modules were identified in the network constructed from yield-
375 invigorating soils, suggesting that a greater number of network nodes in the yield-invigorating soils were
376 functionally interrelated than those in the yield-debilitating soils. A majority of nodes in the meta-
377 modules were not shared between yield-invigorating and -debilitating networks, indicating basal shifts
378 in network architecture during pear production with contrasting yield performance.

379 Furthermore, a higher proportion of negative interactions to positive interactions were identified in
380 the network constructed from yield-invigorating network than the yield-debilitating network. Our results
381 indicated stronger resource competitions in yield-invigorating soils, which means that the soil co-
382 occurrence network was more stable to maintain soil ecosystem function (Coyte et al., 2015). In our
383 study, three module connectors and three module hubs were identified as potentially key taxa in the yield-
384 invigorating network. Interestingly, among those key species, ASV357 affiliated to *Longilinea*,
385 belonging to the Chloroflexota, was recognized as a key phylum associated with higher pear yields in
386 improving pear yield. Similarly, Chloroflexota was reported to be key-stone taxa in the constructed
387 network from agricultural soils with 40-years fertilization (Fan et al., 2021). Chloroflexota play key roles
388 in connecting network nodes of soil microbiome probably due to that Chloroflexota could participate in
389 degrading plant compounds to create more nutrients via pathways for the degradation of starch, cellulose,

390 and longchain sugars, as it is positively correlated with genes for amino sugars, sugar alcohols and simple
391 carbohydrate metabolic pathways (Hug et al., 2013).

392 In this study, a significantly higher content of soil organic matter was observed in yield-invigorating
393 orchards, demonstrating that soil organic matter could drive the assembly of bacterial community.
394 Consensus is emerging that microbial residues are an important constituent of soil organic matter
395 (Kallenbach et al., 2016),, which participate in almost all soil biological processes (Fierer, 2017). Despite
396 the quality of soil organic matter was not evaluated in this study, the quality of soil organic matter was
397 associated with the diversity of microbial community (Ding et al., 2015), which implies more attentions
398 should be paid to illustrate the relationship between the quality of soil organic matter and microbial
399 community in our future work.

400 Structural equation modelling approach has been widely used to decipher keystone indicators
401 associated with soil function and crop production in agroecosystems (Jiang et al., 2020; Chen et al., 2019).
402 In the present study, we observed that soil organic matter, beta diversity of bacterial community, and
403 network connector were key indicators in supporting high-yield pear production based on the structural
404 equation modelling results. Worth to mention, soil organic matter was not directly linked to the yield in
405 the constructed model, indicating that soil organic matter maintain the high-yielding pear production
406 probably via the indirect ways. ~~Therefore, we proposed that yield-invigorating soils harbour unique~~
407 ~~bacterial communities that may improve soil biological fertility, which could be driven by soil organic~~
408 ~~matter and manipulated by keystone species (Chloroflexota) through altering the bacterial interactions.~~

409 **5 Conclusions**

410 In conclusion, yield-invigorating soils displayed a higher content of organic matter and harboured unique
411 bacterial community with greater diversity than yield-debilitating soils. Further Chloroflexota was
412 significantly enriched and identified as a potential keystone taxon in manipulating the interaction of
413 bacterial community in yield-invigorating soils. These findings indicated that soil organic matter
414 triggered the assembly of soil microbiome, which both participated in maintaining crop production . Such
415 knowledge is a first step toward harnessing soil microbiome in support of sustainable agroecosystems.

416

417 **Data availability.** Raw amplicon sequencing data for each sample used in this study was deposited at
418 the National Center for Biotechnology Information (NCBI) in the FASTQ format and is available under
419 the accession number PRJNA749397. Other data that support the findings of this study are available on

420 request from the corresponding author (Xiaomei Ye).

421

422 **Authors' contributions**

423 L. Wang: performed all experiments; L. Wang, X. Ye, and Z. Shen: designed the study, and wrote the
424 majority of the manuscript; L. Wang and Z. Shen and C. Tao: analyzed the data; H. Hu, J. Du, Y. Xi, J.
425 Lin, and D. Chen: participated in the design of the study, provided comments and edited the manuscript.
426 The authors read and approved the final manuscript.

427

428 **Competing interests.** The authors declare that they have no conflict of interest.

429

430 **Disclaimer.** Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional
431 claims in published maps and institutional affiliations.

432

433 **Acknowledgements.** We sincerely thank all those who have assisted me with any part of this paper and
434 all pear orchards owners for providing access to the soil sampling.

435

436 **Financial support.** This research was supported by Jiangsu Agricultural Science and Technology
437 Innovation Fund (CX(19)3094) and National Natural Science Foundation of China (31801842 and
438 42090065).

439

440 **Appendix A. Supplementary data**

441 Supplementary figures and tables to this article can be found in the supplemental material.

442

443 **References**

444 Banerjee, S., Schlaeppli, K., van der Heijden, M.G.J: Keystone taxa as drivers of microbiome structure
445 and functioning. *Nat. Rev. Microbiol.*, 16, 567-576, 2018.

446 Barrios, E.: Soil biota, ecosystem services and land productivity. *Ecological Eco.*, 64, 269-285, 2007.

447 Bell, T., Newman, J.A., Silverman, B.W., Turner, S.L., Lilley, A.K.: The contribution of species richness
448 and composition to bacterial services. *Nature*, 436, 1157-1160, 2005.

449 Bender, S.F., Wagg, C., van der Heijden, M.G.A.: An underground revolution: biodiversity and soil

450 ecological engineering for agricultural sustainability. *Trends Ecol Evol.*, 31, 440-452, 2016.

451 Bolger, A.M., Lohse, M., Usadel, M.: Trimmomatic: a flexible trimmer for Illumina sequence data.
452 *Bioinformatics*, 30, 2114-2120, 2014.

453 Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., Alexander, H.,
454 Alm, E.J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J.E., Bittinger, K., Brejnrod, A., Brislawn,
455 C.J., Brown, C.T., Callahan, B.J., Caraballo-Rodríguez, A.M., Chase, J., Cope, E.K., et al.:
456 Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat.*
457 *Biotechnol.*, 37, 852-857, 2019.

458 Boulesteix, A.L., Janitza, S., Kruppa, J., König, I.R.: Overview of random forest methodology and
459 practical guidance with emphasis on computational biology and bioinformatics. *Wires Data Min.*
460 *Knowl.*, 2, 493-507, 2012.

461 Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A., Turnbaugh, P.J., Fierer,
462 N., Knight, R.: Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample.
463 *P. Nat. Acad. Sci. USA*, 108, 4516-4522, 2011.

464 Cardinale, B.J., Srivastava, D.S., Duffy, J.E., Wright, J.P., Downing, A.L., Sankaran, M., Jouseau, C.:
465 Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature*, 443, 989-992,
466 2006.

467 Chaer, G., Fernandes, M., Myrold, D., Bottomley, P.: Comparative resistance and resilience of soil
468 microbial communities and enzyme activities in adjacent native forest and agricultural soils. *Microb*
469 *Ecol.*, 58, 414-424, 2009.

470 Chaparro, J.M., Sheflin, A.M., Manter, D.K., Vivanco, J.M.: Manipulating the soil microbiome to
471 increase soil health and plant fertility. *Biol. Fert. Soils*, 48, 489-499, 2012.

472 Chen, L., Jiang, Y., Liang, C., Luo, Y., Xu, Q., Han, C., Zhao, Q., Sun, B.: Competitive interaction with
473 keystone taxa induced negative priming under biochar amendments. *Microbiome*, 7, 77, 2019.

474 Coyte, K.Z., Schluter, J., Foster, K.R.: The ecology of the microbiome: networks, competition, and
475 stability. *Science*, 350, 663-666, 2015.

476 Dai, Z., Su, W., Chen, H., Barberán, A., Zhao, H., Yu, M., Yu, L., Brookes, P.C., Schadt, C.W., Chang,
477 S.X., Xu, J.: Long-term nitrogen fertilization decreases bacterial diversity and favors the growth of
478 Actinobacteria and Proteobacteria in agro-ecosystems across the globe. *Global Change Biol.*, 24,
479 3452-3461, 2018.

480 Deng, Y., Jiang, Y., Yang, Y., He, Z., Luo, F., Zhou, J.: Molecular ecological network analyses. *BMC*
481 *Bioinformatics*, 13, 1-20, 2012.

482 Ding, J., Zhang, Y., Wang, M., Sun, X., Cong, J., Deng, Y., Lu, H., Yuan, T., van Nostrand, J.D., Li, D.,
483 Zhou, J., Yang, Y.: Soil organic matter quantity and quality shape microbial community
484 compositions of subtropical broadleaved forests. *Mol. Ecol.*, 24, 5175-5185, 2015.

485 Edwards, J., Johnson, C., Santos-Medellín, C., Lurie, E., Podishetty, N.K., Bhatnagar, S., Eisen, J.A.,
486 Sundaresan, V.: Structure, variation, and assembly of the root-associated microbiomes of rice. *P.*
487 *Nat. Acad. Sci. USA*, 112, E911-E920, 2015.

488 Eo, J., Park, K.: Long-term effects of imbalanced fertilization on the composition and diversity of soil
489 bacterial community. *Agr. Ecosyst. Environ.*, 231, 176-182, 2016.

490 Fan, K., Delgado-Baquerizo, M., Guo, X., Wang, D., Zhu, Y., Chu, H.: Biodiversity of key-stone
491 phylotypes determines crop production in a 4-decade fertilization experiment. *ISME J.*, 15, 550-
492 561, 2021.

493 Fath, B.D., Scharler, U.M., Ulanowicz, R.E., Hannon, B.: Ecological network analysis: network
494 construction. *Ecol. Model.*, 208, 49-55, 2007.

495 Faust, K., Raes, J.: Microbial interactions: from networks to models. *Nat. Rev. Microbiol.*, 10, 538-550,
496 2012.

497 Fierer, N., Bradford, M.A., Jackson, R.B.J.: Toward an ecological classification of soil bacteria. *Ecology*,
498 88, 1354-1364, 2007.

499 Fierer, N., Jackson, R.B.J.: The diversity and biogeography of soil bacterial communities. *P. Nat. Acad.*
500 *Sci. USA*, 103, 626-631, 2006.

501 Fierer, N.: Embracing the unknown: disentangling the complexities of the soil microbiome. *Nat. Rev.*
502 *Microbiol.*, 15, 579-590, 2017.

503 Frąc, M., Hannula, S.E., Bełka, M., Jędrzycka, M.: Fungal biodiversity and their role in soil health. *Front.*
504 *Microbiol.*, 9, 707, 2018.

505 Fuerst, J.A.: Planctomycetes—new models for microbial cells and activities. I. Kurtböke (Ed.), *Microbial*
506 *Resources*, Elsevier/Academic Press, 1-27, 2017.

507 ~~Grace, J., José J.S., Meir, P., Miranda, H.S., Monteset R.A.: Productivity and carbon fluxes of tropical~~
508 ~~savannas. *J Biogeogr.*, 33, 387-400, 2006.~~ Grace, J.B. (Ed): *Structural Equation Modeling and*
509 *Natural Systems*, Cambridge University Press, Cambridge, United Kingdom, 2006.

510 Hu, Q., Tan, L., Gu, S., Xiao, Y., Xiong, X., Zeng, W., Feng, K., Wei, Z., Deng, Y.: Network analysis
511 infers the wilt pathogen invasion associated with non-detrimental bacteria. *npj Biofilms Microb.*, 6,
512 1-8, 2020.

513 Huang, Z., Zhao, F., Wang, M., Qi, K., Wu, J., Zhang, S.: Soil chemical properties and geographical
514 distance exerted effects on arbuscular mycorrhizal fungal community composition in pear orchards
515 in Jiangsu Province, China. *Appl. Soil Ecol.*, 142, 18-24, 2019.

516 Hug, L.A., Castelle, C.J., Wrighton, K.C., Thomas, B.C., Sharon, I., Frischkorn, K.R., Williams, K.H.,
517 Tringe, S.G., Banfield, J.F.: Community genomic analyses constrain the distribution of metabolic
518 traits across the Chloroflexi phylum and indicate roles in sediment carbon cycling. *Microbiome*, 1,
519 1-17, 2013.

520 Jiang, Y., Luan, L., Hu, K., Liu, M., Chen, Z., Geisen, S., Chen, X., Li, H., Xu, Q., Bonkowski, M., Sun,
521 B.: Trophic interactions as determinants of the arbuscular mycorrhizal fungal community with
522 cascading plant-promoting consequences. *Microbiome*, 8, 1-14, 2020.

523 Kallenbach, C.M., Frey, S.D., Grandy, A.S.J.: Direct evidence for microbial-derived soil organic matter
524 formation and its ecophysiological controls. *Nat. Commun.*, 7, 1-10, 2016.

525 Kennedy, A.C., Smith, K.J.: Soil microbial diversity and the sustainability of agricultural soils. *Plant Soil*,
526 170, 75-86, 1995.

527 Kolde, R., Kolde, M. Package 'pheatmap'. 2018, [https://cran.r-](https://cran.r-project.org/web/packages/pheatmap/index.html)
528 [project.org/web/packages/pheatmap/index.html](https://cran.r-project.org/web/packages/pheatmap/index.html).

529 Langfelder, P., Horvath, S.J.: Eigengene networks for studying the relationships between co-expression
530 modules. *BMC Syst. Biol.*, 1, 1-17, 2007.

531 Liaw, A., Wiener, M.: Classification and regression by randomForest. *Rnews*, 2, 18-22, 2002.

532 Lin, J., Sheng, B., Li, X., Yang, Q., Wang, Z., Li, H., Wang, H., Cheng, Y.: A new *pyrus pyrifolia* cultivar
533 'Sucui 1'. *Acta Hort. Sin.*, 40, 1849-1850, 2013 (In Chinese)

534 Lu, L., Yin, S., Liu, X., Zhang, W., Gao, T., Shen, Q., Qiu, H.: Fungal networks in yield-invigorating
535 and-debilitating soils induced by prolonged potato monoculture. *Soil Biol. Biochem.*, 65, 186-194,
536 2013.

537 Magoč, T., Salzberg, S. L.: FLASH: fast length adjustment of short reads to improve genome assemblies.
538 *Bioinformatics*, 27, 2957-2963, 2011.

539 Mei, W., Yu, G., Lai, J., Rao, Q., Umezawa, Y.: basicTrendline: add trendline and confidence interval of

540 basic regression models to plot. R package version 2.0.3, 2018, [http://cran.r-](http://cran.r-project.org/package=basicTrendline)
541 [project.org/package=basicTrendline](http://cran.r-project.org/package=basicTrendline).

542 Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'Hara,
543 R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H.: Package 'vegan'.
544 Community ecology package, version 2, 2013, <http://CRAN.R-project.org/package=vegan>.

545 Parks, D.H., Tyson, G.W., Hugenholtz, P., Beiko, R.G.: STAMP: statistical analysis of taxonomic and
546 functional profiles. *Bioinformatics*, 30, 3123-3124, 2014.

547 Pathan, S.I., Roccotelli, A., Petrovičová, B., Romeo, M., Badagliacca, G., Monti, M., Gelsomino, A..
548 Temporal dynamics of total and active prokaryotic communities in two Mediterranean orchard soils
549 treated with solid anaerobic digestate or managed under no-tillage. *Biol. Fert. Soils.*, 57, 837-861,
550 2021.

551 Philippot, L., Raaijmakers, J.M., Lemanceau, P., van der Putten, W.H.: Going back to the roots: the
552 microbial ecology of the rhizosphere. *Nat. Rev. Microbiol.*, 11, 789-799, 2013.

553 Qiao, C., Penton, C.R., Xiong, W., Liu, C., Wang, R., Liu, Z., Xu, X., Li, R., Shen, Q.: Reshaping the
554 rhizosphere microbiome by bio-organic amendment to enhance crop yield in a maize-cabbage
555 rotation system. *Appl. Soil Ecol.*, 142, 136-146, 2019.

556 Raaijmakers, J.M., Mazzola, M.J.: Soil immune responses. *Science*, 352, 1392-1393, 2016.

557 Rosseel, Y.: Lavaan: An R package for structural equation modeling and more. *J. Stat. Softw.*, 48, 1-36,
558 2012.

559 Rousk, J., Bååth, E., Brookes, P.C., Lauber, C.L., Lozupone, C., Caporaso, J.G., Knight, R., Fierer, N.:
560 Soil bacterial and fungal communities across a pH gradient in an arable soil. *ISME J.*, 4, 1340-1351,
561 2010.

562 Schermelleh-Engel, K., Moosbrugger, H., Müller, H.J.: Evaluating the fit of structural equation models:
563 Tests of significance and descriptive goodness-of-fit measures. *Methods Psychol. Res.*, 8, 23-74,
564 2003.

565 Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., Lesniewski, R.A.,
566 Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B., Thallinger, G.G., Van Horn, D.J.,
567 Weber, C.F.: Introducing mothur: open-source, platform-independent, community-supported
568 software for describing and comparing microbial communities. *Appl. Environ. Microbiol.*, 75,
569 7537-7541, 2009.

570 Shen, Z., Penton, C.R., Lv, N., Xue, C., Yuan, X., Ruan, Y., Li, R., Shen, Q.: Banana Fusarium wilt
571 disease incidence is influenced by shifts of soil microbial communities under different monoculture
572 spans. *Microb. Ecol.*, 75, 739-750, 2018.

573 Shen, Z., Zhong, S., Wang, Y., Wang, B., Mei, X., Li, R., Ruan, Y., Shen, Q.: Induced soil microbial
574 suppression of banana fusarium wilt disease using compost and biofertilizers to improve yield and
575 quality. *Eur. J. Soil Biol.*, 57, 1-8, 2013.

576 Smoot, M.E., Ono, K., Ruscheinski, J., Wang, P.L., Ideker, T.: Cytoscape 2.8: new features for data
577 integration and network visualization. *Bioinformatics*, 27, 431-432, 2011.

578 Speirs, L.B.M., Rice, D.T.F., Petrovski, S., Seviour, R.J.: The phylogeny, biodiversity, and ecology of the
579 *Chloroflexi* in activated sludge. *Front Microbiol.*, 10, 2015, 2019.

580 Tian, J., He, N., Hale, L., Niu, S., Yu, G., Liu, Y., Blagodatskaya, E., Kuzyakov, Y., Gao, Q., Zhou, J.:
581 Soil organic matter availability and climate drive latitudinal patterns in bacterial diversity from
582 tropical to cold temperate forests. *Funct. Ecol.*, 32, 61-70, 2018.

583 Toju, H., Peay, K.G., Yamamichi, M. Narisawa, K., Hiruma K., Naito K., Fukuda S., Ushio M., Nakaoka
584 S., Onoda Y., Yoshida K., Schlaeppi K., Bai Y., Sugiura R., Ichihashi Y., Minamisawa K., Kiers &
585 E.T.: Core microbiomes for sustainable agroecosystems. *Nat Plants*, 4, 247–257, 2018.

586 van der Heijden, M.G., Bardgett, R.D., van Straalen, N.M.J.: The unseen majority: soil microbes as
587 drivers of plant diversity and productivity in terrestrial ecosystems. *Ecol. Lett.*, 11, 296-310, 2008.

588 Wang, C., Liu, D., Bai, E.: Decreasing soil microbial diversity is associated with decreasing microbial
589 biomass under nitrogen addition. *Soil Biol. Biochem.*, 120, 126-133, 2018.

590 Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R.: Naive Bayesian classifier for rapid assignment of rRNA
591 sequences into the new bacterial taxonomy. *Appl. Environ. Microbiol.*, 73, 5261-5267, 2007.

592 Xun, W., Li, W., Xiong, W., Ren, Y., Liu, Y., Miao, Y., Xu, Z., Zhang, N., Shen, Q., Zhang, R.: Diversity-
593 triggered deterministic bacterial assembly constrains community functions. *Nat. Commun.*, 10, 1-
594 10, 2019.

595 Zhao, J., Ni, T., Li, Y., Xiong, W., Ran, W., Shen, B., Shen, Q., Zhang, R.: Responses of bacterial
596 communities in arable soils in a rice-wheat cropping system to different fertilizer regimes and
597 sampling times. *PloS One*, 9, e85301, 2014.


598 Zhong, Y., Hu, J., Xi, Q., Zhang, S., Li, X., Pan, X., Zhao, R., Wang, R., Yang, W., Shangguan, Z., Hu,
599 F., Yang, C., Wang, W.. Soil microbial mechanisms promoting ultrahigh rice yield. *Soil Biol.*

600 Biochem., 143: 107741, 2020.

601

602 **Figure legends**


603 **Fig. 1 Distribution of studied field sites.** Map showing the sites of six pair-located orchards sampled in
604 this study.
605



606

607


608 **Fig. 2 Quantitation of the abundance of bacteria population, and linkage of microbial composition**
609 **to pear yield.** (A) Violin plot showing the abundance of total bacteria for all selected orchards. * indicates
610 a significant difference between yield-invigorating (YI) and yield-debilitating (YD) orchards based on
611 Wilcoxon tests ($p < 0.05$). (B) Correlation plot showing the relationship of microbial composition and
612 yield based on braycurtis distances calculated by Mantel test.



613


614

615 **Fig. 3 Overview of bacterial composition and alpha diversity.** (A) Principal Coordinates Analysis
 616 (PCoA) plot displaying the bacterial community composition calculated based on braycurtis distances.
 617 (B) Violin plot showing the alpha diversity indices (Chao and Shannon) for all selected orchards. *
 618 indicates a significant difference between yield-invigorating (YI) and yield-debilitating (YD) orchards
 619 based on Wilcoxon tests ($p < 0.05$). (C) Venn plot depicting the unique and shared bacterial ASVs
 620 between yield-invigorating (YI) and yield-debilitating (YD) orchards at ASV and sequence insights.
 621 Uni. YI and Uni. YD represent unique ASVs or sequences in the YI or YD soils while Shared represent
 622 shared ASVs or sequences between the YI and YD soils. (D) Heatmap displaying the composition of
 623 unique and shared ASVs at phylum level in YI and YD soils. Numbers in the cell represent the number
 624 of ASVs affiliated to that phylum.



625
 626
 627
 628

629 **Fig. 4 Key taxonomic groups in distinguishing yield-invigorating (YI) and yield-debilitating (YD)**
 630 **orchards.** Stacked bar chart (left panel) showing dominant phyla affiliation in YI and YD soils for six
 631 pair-located sites while horizontal histogram (right panel) depicting relative changes of dominant phyla
 632 in YI soils compared to those in YD soils.




633

634

635

636


637 **Fig. 5 Co-occurrence networks of bacterial community and identified keystone taxa in**
638 **distinguishing yield-invigorating (YI) and yield-debilitating (YD) orchards.** (A) An overview of
639 microbial phylogenetic molecular ecological networks constructed from YI and YD soils. Line with blue
640 color indicates positive correlations whereas lines with red color signifies negative correlations in each
641 network. Modules containing larger than five nodes in the networks are labeled with corresponding letter
642 followed by a number. Circular node colors indicate different bacterial phyla. (B) Stacked figure showing
643 the relative abundance of nodes in each module within each network at the phylum level. (C) Venn plot
644 depicting the unique and shared bacterial ASVs between two networks construed from YI and YD soils.
645 Left panel is plotted based on the original nodes used in building network while right panel is plotted
646 based on the nodes from modules. Uni. YI and Uni. YD represent unique ASVs in the YI or YD networks
647 while Shared represent shared ASVs between the YI and YD networks. (D) Zi-Pi plot showing the
648 distribution of nodes based on their topological roles. The threshold values of Zi and Pi for categorizing
649 OTUs were 2.5 and 0.62 respectively. Node colors indicate different bacterial phyla and node size
650 represent the relative abundance in each network.



651


652

653 **Fig. 6 Relationships among bacterial community, soil edaphic factors and pear yield.** (A) Variance
 654 partitioning analysis (VPA) map of the effects of soil edaphic properties, sample locations, pear yield
 655 and interactions of these factors on the bacterial community. (B) Redundancy analysis (RDA) plot
 656 showing the relationships among all assigned bacterial ASVs and measured soil edaphic properties for
 657 all soils after stepwise selection. (C) Random forest mean predictor importance of selected soil edaphic
 658 properties used as drivers in predicting the pear yield. Red bar indicates that the given predictor is
 659 significant while black bar indicates that the given predictor is non-significant. The %IncMSE in the up
 660 panel means the increase in mean squared error while IncNodePurity indicates the increase in node
 661 purity. The values of these two indices represent the importance of each variable to predict the module.
 662 A larger value indicates that the variable is more important. (D) Correlation plot showing the
 663 relationship of microbial composition and soil organic matter based on braycurtis distances calculated
 664 by Mantel test.



665
 666
 667

668 **Fig. 7 Structural equation modeling (SEM) describing the biotic and abiotic factors in affecting**
 669 **the crop production.** Structural equation model was built incorporating soil organic matter, microbial
 670 biomass, beta diversity (PCoA), key taxa, network hubs including module hubs and network
 671 connectors, and yield. The path analysis numbers adjacent to arrows indicate the relationship's effect
 672 size and the associated bootstrap *P* value. Cyan and red arrows indicate positive and negative
 673 relationships, respectively. Paths with non-significant coefficients are presented as gray lines.



674
 675
 676