

1 **Soil bacterial community triggered by organic matter inputs**  
2 **associates with a high-yielding pear production**

3  
4 Li Wang<sup>1</sup>, Xiaomei Ye<sup>1✉</sup>, Hangwei Hu<sup>2</sup>, Jing Du<sup>1</sup>, Yonglan Xi<sup>1</sup>, Zongzhuan Shen<sup>3✉</sup>, Jing Lin<sup>4</sup>, Deli  
5 Chen<sup>2</sup>  
6

7 **Affiliations**

8 <sup>1</sup> Institute of Animal Science, Jiangsu Academy of Agricultural Sciences, Key Laboratory of Crop and  
9 Livestock Integrated Farming, Ministry of Agriculture and Rural Affairs, Nanjing 210014, China

10 <sup>2</sup> Faculty of Veterinary and Agricultural Sciences, University of Melbourne, Melbourne, Victoria 3010,  
11 Australia

12 <sup>3</sup> Jiangsu Provincial Key Lab of Solid Organic Waste Utilization, Jiangsu Collaborative Innovation  
13 Center of Solid Organic Wastes, Educational Ministry Engineering Center of Resource-saving fertilizers,  
14 The Key Laboratory of Plant Immunity, Nanjing Agricultural University, Nanjing, 210095, China

15 <sup>4</sup> Institute of Pomology, Jiangsu Academy of Agricultural Sciences, Nanjing 210014, China

16  
17 ✉ Corresponding author: Xiaomei Ye, Institute of Animal Science, Jiangsu Academy of Agricultural  
18 Sciences, Key Laboratory of Crop and Livestock Integrated Farming, Ministry of Agriculture and Rural  
19 Affairs, Nanjing 210014, China. Zognzhuan Shen, College of Resources and Environmental Sciences,  
20 Nanjing Agricultural University, 210095, Nanjing, China. E-mail: shenzongz@njau.edu.cn. Tel:  
21 (86)02584395521.  
22

23 **Abstract** The roles of microorganisms in enhancing crop production have been demonstrated for a range  
24 of cropping systems. Most studies to date, however, have been confined to a limited number of locations,  
25 making it difficult to identify general soil biotic and abiotic characteristics underpinning the yield-  
26 promotion across various locations. This knowledge gap limits our capacity to harness soil microbiome  
27 to improve crop production. Here we used high-throughput amplicon sequencing to investigate the  
28 common features of bacterial community composition, ecological networks and physicochemical  
29 properties in six yield-invigorating and adjacent yield-debilitating orchards. We found that yield-

30 invigorating soils exhibited higher contents of organic matter than yield-debilitating soils and harboured  
31 unique bacterial communities. Greater alpha diversity and higher relative abundances of Planctomycetota  
32 and Chloroflexota were observed in yield-debilitating soils. Co-occurrence network analysis revealed  
33 that yield-invigorating soils displayed a greater number of functionally interrelated modules (meta-  
34 modules) and a higher proportion of negative links to positive links. Chloroflexota was recognized as a  
35 keystone taxon in manipulating the interaction of bacterial communities in yield-invigorating soils.  
36 Structural equation modelling showed that soil organic matter, beta diversity of bacterial community, and  
37 network connector (Chloroflexota) were identified as potential key factors in explaining the high-yield  
38 pear production. Altogether, we provide evidence that yield-invigorating soils across a range of locations  
39 appear to share common features, including accumulation of soil organic matter, higher microbial  
40 diversity, enrichment of key taxa like Chloroflexota, and maintaining a competitive network. These  
41 findings have implications for science-based guidance for sustainable food production.

42

43 **Keywords:** Soil organic matter, Microbial diversity, Random forest prediction, Co-occurrence network,  
44 Keystone taxa

45

## 46 **1 Introduction**

47 Soils are essential to human wellbeing due to their great contributions to the production of food, fiber,  
48 feed, and medicine (Raaijmakers and Mazzola, 2016). Soil organisms play critical roles in maintaining  
49 these ecosystem services, such as driving nutrient cycling, maintaining soil fertility, improving plant  
50 productivity and suppressing plant diseases (Bender et al., 2016; Barrios, 2007). Microorganisms  
51 participate in nearly all soil biological processes, and the microbial abundance, community composition  
52 and activity primarily determine the sustainable productivity of agricultural lands (Philippot et al., 2013).  
53 Fungi participate in decomposition of organic matter and deliver nutrients for plant growth (Fr ac et al.,  
54 2018), however, considering that bacteria are the most diverse and abundant group of microorganisms in  
55 soil, bacterial communities and their functions can be pivotal indicators for crop production in  
56 agroecosystems (van der Heijden et al., 2008).

57 In general, an increase in microbial diversity is linked to a high-yielding crop production mainly  
58 through improving the host resilience to physical or chemical disturbances, modifying plant competition,  
59 and facilitating plant access to nutrients (Chaer et al., 2009; Kennedy and Smith, 1995). Since individual

60 organisms do not live in isolation but rather form a complex system of inter-species interactions in soil,  
61 interactions among community members were found to be related to crop production in the potato  
62 monoculture system (Lu et al., 2013). Enrichments of key functional microbes in soil were deemed to  
63 serve specific soil system functions, such as suppressing soil-borne pathogens and maintaining  
64 sustainable crop production (Banerjee et al., 2018). However, the relative contributions of microbial  
65 diversity, interactions among community members, or enrichment of key taxa to crop production remain  
66 largely unknown. Therefore, it is highly desirable to identify pivotal indicators of bacterial community  
67 composition in response to high-yielding crop production.

68 Changes in composition of soil bacterial communities across space are often strongly correlated  
69 with soil pH (Fierer and Jackson, 2006). Soil pH has been recognized as a key driver in determining the  
70 assembly of bacterial community in arable soils by field or microcosm experiments (Rousk et al., 2010).  
71 However, recent studies have demonstrated that compositions of soil bacterial communities were driven  
72 by a myriad of soil abiotic traits, such as organic matter contents, forms and contents of soil nutrient  
73 (Tian et al., 2018; Wang et al., 2018). For example, soil bacterial community composition, which  
74 determines the ability of soil to suppress soil-borne pathogens, was found to be strongly correlated with  
75 soil organic matter (Shen et al., 2018). An imbalanced ratio of soil nutrients, *i.e.*, ratio of nitrogen to  
76 phosphorus or potassium could be a driving force altering the bacterial community composition in long-  
77 term fertilized soils (Eo and Park, 2016). Key soil chemical properties identified in controlling the  
78 distribution and abundance of bacterial community is largely depending on the geographical distributions  
79 of soils. As a consequence, a better understanding of the relationship between soil edaphic properties and  
80 bacterial community composition is critical to develop targeted manipulation options to increase soil  
81 service provisions.

82 'Sucui No. 1' pear, is an early-maturing variety bred by the Jiangsu Academy of Agricultural  
83 Sciences, China, and has been popularly cultivated in Eastern and Central China, due to the advantages  
84 including easy to produce, adaptable to the environment, and has good quality and high economic  
85 benefits (Lin et al., 2013). With the increasing demand in China, sustainable production of high-quality  
86 pear is becoming increasingly important. Manipulation of soil microbiomes has shown to be an effective  
87 way to increase soil productivity (Chaparro et al., 2012). Considering that large-scale surveys could  
88 exhibit the diversity of soil microbial communities exceeds what is found in host-associated communities  
89 (Toju et al., 2018), it is necessary to explore the general microbial characteristics of multiple yield-

90 invigorating soils and identify key environmental drivers in assembling bacterial community.

91 In this study, compared to local average yield, orchard showing higher pear yield production was  
92 recognized as yield-invigorating (YI) orchard while orchard displaying lower pear yield production was  
93 regard as yield-debilitating orchard (YD). After field surveys accomplished in 2019, six yield-  
94 invigorating and adjacent yield-debilitating pear orchards in total were selected for further analysis of  
95 soil chemical properties and microbiome. We hypothesized that high input of organic fertilizer could  
96 improve soil structure and modify chemical properties, which leading to YI soils harbor unique bacterial  
97 communities that maintains the high-yielding pear production. To address this, soil bacterial communities  
98 and edaphic properties of the study sites were compared to (1) decipher the differences of taxonomic  
99 diversity, and composition of the bacterial community, and (2) determine the contributions of  
100 environmental variables to the changes in the structure of bacterial communities.

## 101 **2 Methods**

### 102 2.1 Study sites and experimental design

103 From July - August 2019, a field production survey of orchards cultivated with ‘Suci No. 1’ pear was  
104 performed after pear fruits harvest to compare the differences of soil nutrients and microbiota between  
105 yield-invigorating with yield-debilitating orchards. The locations, planting density, cropping years, soil  
106 type and total yield were recorded. To minimize the effects of microclimate at each site, only pair-located  
107 pear orchards with invigorating and debilitating yield and at similar growth stage were selected for this  
108 research. In total, six pair-located yield-invigorating and -debilitating pear orchards distributed in four  
109 cities of Jiangsu province, China, were selected in the main pear production areas (Fig. 1, Table S1). The  
110 yield per tree was obtained by dividing the total yield per hectare by plant density.

111 **Fig. 1 here**

112 Paired yield-invigorating and yield-debilitating orchards from Fengxian (FX), Suining (SN) and  
113 Tongshan (TS) were maintained in the Xuzhou city under the warm temperate sub-humid monsoon  
114 climate. This site has a mean annual temperature (MAT) of 14.5 °C and mean annual precipitation (MAP)  
115 of 847 mm. Orchards from Taixing (TX) were located in the Taizhou city under the humid southern  
116 subtropical climate with a MAT of 15.3 °C and MAP of 1055 mm. Orchards from Gaochun (GC) were  
117 located in the Nanjing city under the humid subtropical monsoon climate with a MAT of 15.4 °C and  
118 MAP of 1106 mm. Orchards from Zhangjiagang (ZJ) were located in the Suzhou city under the humid  
119 subtropical monsoon climate with a MAT of 15.7 °C and MAP of 1094 mm. For paired yield-invigorating

120 and-debilitating orchards, the irrigation and pesticide management practices were similar according to  
121 farm records. However, yield-invigorating orchard was usually amended with more organic fertilizer  
122 under integrated nutrients management whereas the co-located yield-debilitating orchard received more  
123 chemical fertilizer under intensive management. Detailed information about fertilization regimes for each  
124 orchard is shown in Table S2.

## 125 2.2 Soil sample collection and chemical properties determination

126 Along with the field survey, soil sampling campaigns were performed from July - August 2019 after pear  
127 fruits harvest. For each yield-invigorating or -debilitating orchard, four subplots with three pear trees in  
128 each subplot were randomly selected for soil sampling. Subsequently three soil cores (0-20 cm) under  
129 the trunk base for each tree were collected using a 25 mm soil auger. In total, nine soil cores for each  
130 subplot were pooled as a composite sample and finally four composite soil samples for each orchard  
131 were collected and promptly transported on ice to the laboratory. After sifting through a 2 mm sieve and  
132 thoroughly mixing, one portion of each soil sample was air-dried for chemical property analyses while  
133 the remainder was stored at -70 °C for DNA extraction.

134 Soil chemical properties including soil pH, content of organic matter (OM), total nitrogen (TN),  
135 available phosphorus (AP), available potassium (AK), alkali-hydrolyzable nitrogen (N), exchangeable  
136 calcium (Ca), effective magnesium (Mg), effective iron (Fe), effective manganese (Mn), effective copper  
137 (Cu) and effective zinc (Zn), were measured according to methods described by Shen et al. (2018) and  
138 Huang et al. (2019). Briefly, soil pH was determined using a glass electrode meter in a suspension with  
139 a 1:5 soil/water ratio (w/v). Soil OM was determined using the potassium dichromate external heating  
140 method. TN was determined using a dry combustion method on an Element Analyzer (Vario EL,  
141 Germany). AP and AK were determined using the molybdenum blue method after soil was extracted with  
142 sodium bicarbonate and flame photometry after soil was extracted with ammonium acetate, respectively.  
143 Soil alkaline hydrolysable nitrogen (N) was measured by the alkaline hydrolysable diffusion method.  
144 Contents of soil Ca, Mg, Fe, Mn, Cu and Zn were determined by the atomic absorption spectroscopy  
145 method using ICE 3300 AAS Atomic Absorption Spectrometer (ThermoScientific, USA) after acid  
146 hydrolysis.

## 147 2.3 Soil DNA extraction and bacterial abundance quantification

148 Genomic DNA from 0.25 g soil for each sample was extracted by using the DNeasy® PowerSoil® Kit  
149 (QIAGEN GmbH, Germany) according to the manufacturer's instructions. The abundances of soil

150 bacteria were determined with Eub338F/Eub518R primer using a 7500 Real Time PCR System (Applied  
151 Biosystems, USA). Standard curves were generated by using 10-fold serial dilutions of a plasmid  
152 containing a full-length copy of the 16S rRNA gene from *Escherichia coli*. Quantitative PCR analysis  
153 was performed in 96-well plates with a 20- $\mu$ l mixture for each reaction using SYBR<sup>®</sup>Premix Ex Taq<sup>™</sup>  
154 (TaKaRa, Japan). Thermal cycling was conducted according to a standard procedure with three replicates,  
155 and the results were expressed as log copy numbers g<sup>-1</sup> dry soil.

#### 156 2.4 Sequencing library construction and sequencing

157 The gene-specific primers 515F/806R with 12 bp barcode were used to amplify the V4 region of bacterial  
158 16S rRNA gene on the BioRad S1000 (Bio-Rad Laboratory, CA) according to the protocols described  
159 by Caporaso et al. (2011). All constructed libraries were sequenced using the Illumina NovaSeq 6000 at  
160 the Guangdong Magigene Biotechnology Co., Ltd. (Guangzhou, China).

#### 161 2.5 Sequence processing

162 Quality filtering of the paired-end raw reads was performed to obtain the high-quality clean reads  
163 according to the Trimmomatic V0.33 (Bolger et al., 2014) quality control process. Sequences were  
164 assigned to each sample based on their unique barcode, after which the barcodes and primers were  
165 removed. Paired-end clean reads were merged using FLASH V1.2.11 (Magoč and Salzberg, 2011). Raw  
166 tags were processed to generate the final ASV (Amplicon Sequence Variant) table file at 97% pairwise  
167 identity according to the QIIME2 pipeline (Bolyen et al., 2019). The nonbacterial and mitochondrial  
168 ASVs and extremely low frequency ASVs (relative abundance < 0.01%) were removed. A representative  
169 sequence for each ASV was selected and classified using the RDP classifier against the RDP Bacterial  
170 16S database (Wang et al., 2007).

#### 171 2.6 Statistical analyses

172 Statistical analyses were performed using the software SPSS 20.0 (SPSS Technologies, Armonk, NY,  
173 USA) and R (<http://www.R-project.org/>). Non-normal data were square-root or log transformed when  
174 necessary. The significance of soil properties or microbial taxa in yield-invigorating or-debilitating  
175 orchards was determined based on paired Wilcoxon rank sum test, and adjusted *P* values (< 0.05) were  
176 obtained by the FDR method. Mantel tests were used to identify the correlations between microbial  
177 community composition and pear yield, and soil chemical properties using the ‘vegan’ package (Oksanen  
178 et al., 2013) in R. The linear regression analyses relating yield to selected microbial taxa or soil chemical  
179 properties were conducted using the ‘basicTrendline’ package (Mei et al., 2018) in R.

180 Principal Coordinate Analysis (PCoA) based on the Bray-Curtis distance was performed in  
181 MOTHUR V1.38.1 (Schloss et al., 2009) and visualized by the ‘ggplot2’ package (Wickham and Chang,  
182 2015) in R to explore the differences in microbial community composition. Permutational multivariate  
183 analysis of variance (PERMANOVA) was performed to evaluate the significant differences of microbial  
184 community composition according to sample locations and orchard yield using the ‘vegan’ package in R.  
185 Microbial alpha diversity indexes (Chao, Shannon) were calculated based on randomly resampled ASV  
186 abundance matrices at the same depth (23,800 sequences) in MOTHUR. A Venn diagram was generated  
187 based on the final ASVs to compare microbial community composition between yield-invigorating and  
188 -debilitating orchard soils. The affiliations of unique and shared ASVs in yield-invigorating and -  
189 debilitating soils were compared to evaluate the differences of the bacterial community composition and  
190 plotted using the ‘pheatmap’ package (<https://cran.r-project.org/web/packages/pheatmap>) in R. Fold  
191 changes (log<sub>2</sub> transformed) of shared ASVs across yield-invigorating and -debilitating soils were  
192 calculated. The ASVs with fold change ratios > 2 and unique ASVs in yield-invigorating soils were  
193 recognized as potential responders to yield promotion. In addition, to better understand the bacterial  
194 community composition, relative abundances at the genus level were compared by STAMP software  
195 v2.1.3 (Parks et al., 2014).

196 The phylogenetic molecular ecological networks (pMEN) were constructed using the random  
197 matrix theory-based approach to explore the organization of bacterial communities in yield-invigorating  
198 (YI) or yield-debilitating (YD) soil samples. Potential ecological interactions among bacteria were  
199 determined by modeling the microbial community using Molecular Ecological Network Analysis  
200 (<http://ieg2.ou.edu/MENA>) based on pear yield (Deng et al., 2012). Given the large number of rare taxa  
201 that are specific to certain locations, ASVs that occurred in less than half of soil samples and lower than  
202 0.01% were filtered, which resulting in 591 and 485 ASVs for YI and YD samples respectively, before  
203 networks constructed. The microbial network was constructed using random matrix theory-based at  
204 0.94 similarity threshold and visualized using Cytoscape 2.8.3 software (Smoot et al., 2011). Module  
205 clustering and composition in yield-invigorating and -debilitating networks were compared and plotted  
206 in R using the ‘pheatmap’ and ‘ggplot2’ packages. And the threshold values of  $Z_i$  and  $P_i$  was 2.5 and  
207 0.62, respectively for topology analysis of the network. Redundancy analysis (RDA) was performed in  
208 the R ‘vegan’ package to examine the relationship among frequencies of ASVs, samples and soil  
209 variables, which were selected using ‘stepAIC’ in R. Variance partitioning analysis (VPA) was used to

210 determine the contributions of soil properties, sample location, and yield, as well as interactions among  
211 the variation in a microbial community with hellinger-transformed data. The predictors of selected soil  
212 properties for explaining the pear yield were identified by random forest regression analysis (Boulesteix,  
213 et al., 2012). The significance of each predictor in the response variables was assessed with the  
214 ‘rfPermute’ package (Liaw and Wiener, 2002) with 1000 permutations based on 1000 trees. Structural  
215 equation modelling was applied to evaluate relative contributions of soil chemical properties and  
216 bacterial community to pear yield (Grace et al., 2006; Schermelleh-Engel et al., 2003). The conceptual  
217 SEM fitness was examined on the basis of a non-significant chi-square test ( $P > 0.05$ ) and the goodness-  
218 of-fit index (GFI). Model was fitted using the ‘lavaan’ package in R (Rosseel, 2012).

### 219 **3 Results**

#### 220 3.1 Overview of sequencing data

221 In total, 1,622,858 16S rRNA sequences were retained after quality control and a total of 9,394 ASVs  
222 were obtained for the 16S rRNA gene sequences based on 97% similarity. Among the total 16S rRNA  
223 gene sequences, 159 ASVs with 74,372 sequences were classified as Archaea while 9,235 ASVs with  
224 1,548,486 sequences were identified as Bacteria. Among Bacteria, Acidobacteriota, Pseudomonadota,  
225 Chloroflexota, Planctomycetota and Actinomycetota were the most abundant phyla (Fig. S1).

#### 226 3.2 Soil chemical properties

227 Soil chemical properties differed significantly among the locations and orchard yield types (Table S3).  
228 However, when taken all sites together, only a higher relative abundance of soil organic matter (OM) on  
229 average was observed in yield-invigorating orchards compared to that in yield-debilitating orchards  
230 based on the Wilcoxon test.

#### 231 3.3 Bacterial abundances and community compositions

232 Yield-invigorating orchards together displayed significantly higher abundances of total bacteria than that  
233 in co-located yield-debilitating orchards based on real time PCR result (Fig. 2A). Meanwhile, bacterial  
234 community compositions at the ASV level were significantly correlated to pear yield ( $r = 0.460$ ,  $p =$   
235  $0.001$ ) (Fig. 2B).

### 236 **Fig. 2 here**

237 PCoA based on Bray-Curtis distance matrices clearly revealed location-based differences in  
238 bacterial community compositions (Fig. 3A). Six distinct groups representing samples from different  
239 locations (FX, GC, SN, TS, TX and ZJ) were obviously separated and confirmed by PERMANOVA test



240 ( $F = 14.9$ ,  $P = 0.001$ ). At each location, soil bacterial community composition in yield-invigorating  
241 orchards was significantly separated from that in co-located yield-debilitating orchards, which was also  
242 confirmed by PERMANOVA test ( $F = 3.6$ ,  $P = 0.001$ ). Although only the Shannon diversity in yield-  
243 invigorating orchards from GC and ZJ was significantly higher than that in co-located yield-debilitating  
244 orchards (Fig. S2), the mean alpha diversity indices of Chao and Shannon in all yield-invigorating  
245 orchards were significantly higher than those in all yield-debilitating orchards based on the paired  
246 Wilcoxon test (Fig. 3B).

247 **Fig. 3 here**

248 The Venn diagram showed that 4,540 ASVs occupying over 90% of total sequences were shared  
249 between yield-invigorating and -debilitating orchards (Fig. 3C). Among these shared ASVs, the fold  
250 changes larger than 2 of ASVs in yield-invigorating compared to yield-debilitating orchards were  
251 recognized as potential responders linking to yield improvement. Surprisingly, none of these ASVs  
252 potentially linked to yield improvement were shared among six separated collocated orchards (Fig. S3).  
253 A total of 2,546 unique ASVs with 53,222 sequences were found in all yield-invigorating orchards while  
254 2,308 unique ASVs with 44,389 sequences were observed in all yield-debilitating orchards. Among  
255 these unique ASVs, almost 70% of ASVs were shared between yield-invigorating orchards and -  
256 debilitating orchards. However, no shared unique ASVs were found among six separately located  
257 orchards. The affiliation of unique and shared ASVs at the phylum level exhibited that the  
258 Pseudomonadota, Planctomycetota, Chloroflexota, Acidobacteriota and Actinomycetota were the top  
259 five phyla (Fig. 3D).

260 At the phylum level, the relative abundances of bacterial dominant phyla varied across the location  
261 and orchard yield condition. Pseudomonadota, Acidobacteriota, Actinomycetota, Chloroflexota and  
262 Planctomycetota were the top five abundant phyla (Fig. 4). The mean abundance of Chloroflexota and  
263 Planctomycetota was significantly higher while Firmicutes was significantly lower in yield-invigorating  
264 orchards compared to yield-debilitating orchards based on Wilcoxon test (Fig. S4).

265 At a finer resolution, 967 genera were observed for all soil samples, among which 299 genera  
266 appeared in more than half of soil samples in yield-invigorating or -debilitating orchards. However, only  
267 34 genera displayed significant differences between yield-invigorating or -debilitating orchard soils  
268 based on STAMP analysis (Fig. S5). Interestingly, *Ornatilinea*, *Ktedonobacter*, *Longilinea*, belonging  
269 to Chloroflexota, were significantly enriched in yield-invigorating orchard soils. *Gimesia* in

270 Planctomycetota and *Arenimonas* in Pseudomonadota showed significantly higher relative abundances  
271 in yield-invigorating orchard soils than in yield-debilitating orchard soils.

272 **Fig. 4 here**

### 273 3.4 Co-occurrence patterns of bacterial community

274 The yield-invigorating network contained 302 nodes, 448 edges, and 11 larger modules (> 5 nodes), with  
275 an average connectivity (avgK) of 2.967, average path distance of 5.494 and clustering coefficient  
276 (avgCC) of 0.152, while the values in the yield-debilitating network were 235, 334, 9, 2.843, 6.232 and  
277 0.131, respectively (Fig. 5A, Table S4). The module eigengene network analysis revealed a difference in  
278 the higher-order organization between the two networks. Notably, the node composition was substantially  
279 different between the two networks as the relative abundances of dominant phyla were obviously  
280 different among different modules (Fig. 5B and C). A higher proportion of nodes in the module of yield-  
281 invigorating network was unique. ASVs affiliated to Acidobacteriota, Chloroflexota, Pseudomonadota,  
282 Actinomycetota, and Planctomycetota within the unique modules (M9, M10 and M11) were observed in  
283 the yield-invigorating versus yield-debilitating network.

284 Analysis using the threshold values of  $Z_i$  (within-module connectivity) and  $P_i$  (among-module  
285 connectivity) showed that majority of nodes from both networks were categorized as peripherals that had  
286 only a few links and almost always linked to the nodes within their own modules (Fig. 5D). Although  
287 only three nodes affiliated to Acidobacteriota were categorized as module hubs in the yield-invigorating  
288 network, seven nodes belonging to Acidobacteriota, Actinomycetota and Pseudomonadota were  
289 categorized as module hubs in the yield-debilitating network. Interestingly, four nodes including  
290 *Longilinea* species from Chloroflexota in the yield-invigorating network whereas only one node in the  
291 yield-debilitating network was categorized as module connectors (Table S5).

292 **Fig. 5 here**

### 293 3.5 Relationships between soil chemical properties and microbial community composition

294 Soil chemical properties were significantly correlated to the bacterial community compositions (Mantel:  
295  $r = 0.803$ ,  $p = 0.001$ ). Soil chemical properties, location, and orchard explained 44.9% of the observed  
296 variation, leaving 55.1% of the variation unexplained for bacterial community composition based on  
297 VPA result (Fig. 6A). Variation in the community composition was largely explained by soil properties  
298 (42.3%), and was also influenced by locations and orchard yield types.

299 After forward stepwise selection, the module including soil OM, TN, alkaline N, AP and AK,

300 available calcium (Ca), copper (Cu) and manganese (Mn) explained the majority of the variation in  
301 bacterial community composition (Fig. 6B). As evidenced by the RDA vectors, OM within the module  
302 was identified as the top important soil property that determines the composition of bacterial community.  
303 Random forest analysis showed that contents of soil Mn, OM and Ca were the top parameters for  
304 predicting the orchard yield (Fig. 6C). Furthermore, soil OM was also significantly correlated with  
305 bacterial communities as revealed by Mantel test (Fig. 6D, Table S6).

306 **Fig. 6 here**

307 3.6 Relationships of soil chemical and microbial indicators with orchard yield

308 Soil OM as potentially key soil chemical properties and bacterial alpha diversity, beta diversity and  
309 relative abundance of Chloroflexota and Planctomycetota as potentially key microbial indicators  
310 associated with pear yield were used to construct a model to explain yield improvement. Final structural  
311 equation modelling (path analysis) (Fig. 7 and S6) showed that the strongest driver explaining yield  
312 improvement was beta diversity of bacterial community (PCoA) ( $r = 0.959$ ,  $P < 0.001$ ), which was  
313 positively affected by content of soil OM ( $r = 0.843$ ,  $P < 0.001$ ). Alpha diversity (Chao) of bacterial  
314 community also determined yield improvement to a large extent ( $r = 0.542$ ,  $P = 0.009$ ). However, alpha  
315 diversity was not significantly correlated with content of soil OM.

316 **Fig. 7 here**

#### 317 **4 Discussion**

318 Although pear is among the most important fruits worldwide, soil microbial communities in pear  
319 orchards have been largely under-investigated (Huang et al., 2019). The present study attempts to  
320 decipher the bacterial community linked to high-yield production of pear. Our results based on Mantel  
321 analysis suggested significant correlations among bacterial community, soil chemical properties and pear  
322 yield. Microbial characteristics responding to yield promotion have repeatedly been observed on several  
323 crops depending on single experimental site (Zhong et al., 2020; Qiao et al., 2019; Shen et al., 2013). It  
324 remained unclear, however, whether these distinctions are ubiquitous at a large-scale. By comparing  
325 multiple co-located yield-invigorating and -debilitating orchards, we demonstrate that high-yielding pear  
326 production soils exhibited high organic matter contents and harbored bacterial communities with high  
327 diversity, significantly enriched indigenous microbes and more interactive network, which was triggered  
328 by high-inputs of soil organic fertilizer. Here we discussed these main results and potential mechanisms  
329 in detail.

330 Microbial diversity is critical to soil ecosystems in maintaining the integrity, function and long-term  
331 sustainability (Kennedy and Smith, 1995). Higher soil biodiversity is considered to be linked to a more  
332 stable system and enhance the combination of vital microbial functions and processes (Cardinale et al.,  
333 2006; Bell et al., 2005). In line with a previous report that crop yield was correlated to the soil bacterial  
334 diversity (Zhao et al., 2014), greater diversity of bacterial community in yield-invigorating soils was  
335 observed in the present study. Hence we infer that higher microbial diversity may result in a more  
336 productive agroecosystem, contributing to sustainable pear production.

337 In this study, we found that Pseudomonadota, Acidobacteriota, Actinomycetota, Planctomycetota  
338 and Chloroflexota were the top abundant phyla. This result roughly agreed with previous studies showing  
339 that Pseudomonadota, Acidobacteriota and Actinomycetota are usually dominant bacterial taxa in  
340 agricultural soils (Xun et al., 2019; Dai et al., 2018), while Planctomycetota and Chloroflexota exhibit  
341 an unexpectedly high relative abundance in rice cropped soil (Edwards et al., 2015) and sandy loam soil  
342 (Pathan et al., 2021). The highest relative abundance of Pseudomonadota was probably explained by the  
343 fact that Pseudomonadota are considered as copiotrophic bacteria and flourish in soils with large amounts  
344 of available nutrients (Fierer et al., 2007).

345 Moreover, a significantly higher abundance of Planctomycetota and Chloroflexota was observed in  
346 yield-invigorating orchards, indicating that Planctomycetota and Chloroflexota may be associated with  
347 pear yield-improvement. There is no direct evidence showing that Planctomycetota could improve plant  
348 growth. However, Planctomycetota has been reported to be involved in many soil biological processes  
349 such as ammonification, carbohydrate and polysaccharide metabolism (Fuerst, 2017). This implies that  
350 Planctomycetota may promote plant production through improving soil fertility. Chloroflexota is a  
351 facultative anaerobic phylum including autotrophic, heterotrophic and mixotrophic taxa (Speirs et al.,  
352 2019). Considering that soil amended with organic fertilizer may enhance the soil water holding capacity,  
353 the yield-invigorating soils with more organic material input have a higher soil moisture content,  
354 especially after irrigation, probably leading to the enrichment of Chloroflexota in soil.

355 Network analysis is a systems-level method to explore interactions within an ecosystem that cannot  
356 be directly observed through co-occurrence analysis (Fath et al., 2007). Similar to the food web network  
357 analyses in macro ecosystems, microorganisms also form complex interactions with other species (Faust  
358 and Raes, 2012) and have been widely investigated to explore the linkage of microbial network with soil  
359 function, such as nutrient supply (Fan et al., 2021) and disease suppression (Lu et al., 2013). Overall, in

360 line with previous findings (Hu et al., 2020), the topological properties of the constructed networks,  
361 including connectivity, average clustering coefficients, average degree distance, and modularity indicate  
362 that these networks are scale-free, modular and small world. In short, a scale-free network represents that  
363 a network whose connectivity follows a power law, and most of nodes have only a few connections with  
364 other nodes. Meanwhile, a small-world network is the network in which most nodes are not neighbors of  
365 one another, but most nodes can be reached by a few paths. Modularity is a fundamental characteristic  
366 of biological network as a module in the network is a group of nodes that are highly connected within  
367 the group, but very few connections outside the group (Deng et al., 2012). Our comparative network  
368 analysis indicated that microbial co-occurrence patterns in soils links to different pear production. As a  
369 meta-module is usually considered as a group of modules functionally interrelated (Langfelder and  
370 Horvath, 2007), a greater number of meta-modules were identified in the network constructed from yield-  
371 invigorating soils, suggesting that a greater number of network nodes in the yield-invigorating soils were  
372 functionally interrelated than those in the yield-debilitating soils. A majority of nodes in the meta-  
373 modules were not shared between yield-invigorating and -debilitating networks, indicating basal shifts  
374 in network architecture during pear production with contrasting yield performance.

375 Furthermore, a higher proportion of negative interactions to positive interactions were identified in  
376 the network constructed from yield-invigorating network than the yield-debilitating network. Our results  
377 indicated stronger resource competitions in yield-invigorating soils, which means that the soil co-  
378 occurrence network was more stable to maintain soil ecosystem function (Coyte et al., 2015). In our  
379 study, three module connectors and three module hubs were identified as potentially key taxa in the yield-  
380 invigorating network. Interestingly, among those key species, ASV357 affiliated to *Longilinea*,  
381 belonging to the Chloroflexota, was recognized as a key phylum in improving pear yield. Similarly,  
382 Chloroflexota was reported to be key-stone taxa in the constructed network from agricultural soils with  
383 40-years fertilization (Fan et al., 2021). Chloroflexota play key roles in connecting network nodes of soil  
384 microbiome probably due to that Chloroflexota could participate in degrading plant compounds to create  
385 more nutrients via pathways for the degradation of starch, cellulose, and longchain sugars, as it is  
386 positively correlated with genes for amino sugars, sugar alcohols and simple carbohydrate metabolic  
387 pathways (Hug et al., 2013).

388 In this study, a significantly higher content of soil organic matter was observed in yield-invigorating  
389 orchards, demonstrating that soil organic matter could drive the assembly of bacterial community.

390 Consensus is emerging that microbial residues are an important constituent of soil organic matter  
391 (Kallenbach et al., 2016), , which participate in almost all soil biological processes (Fierer, 2017). Despite  
392 the quality of soil organic matter was not evaluated in this study, the quality of soil organic matter was  
393 associated with the diversity of microbial community (Ding et al., 2015), which implies more attentions  
394 should be paid to illustrate the relationship between the quality of soil organic matter and microbial  
395 community in our future work.

396 Structural equation modelling approach has been widely used to decipher keystone indicators  
397 associated with soil function and crop production in agroecosystems (Jiang et al., 2020; Chen et al., 2019).  
398 In the present study, we observed that soil organic matter, beta diversity of bacterial community, and  
399 network connector were key indicators in supporting high-yield pear production based on the structural  
400 equation modelling results. Worth to mention, soil organic matter was not directly linked to the yield in  
401 the constructed model, indicating that soil organic matter maintain the high-yielding pear production  
402 probably via the indirect ways. Therefore, we proposed that yield-invigorating soils harbour unique  
403 bacterial communities that may improve soil biological fertility, which could be driven by soil organic  
404 matter and manipulated by keystone species (Chloroflexota) through altering the bacterial interactions.

## 405 **5 Conclusions**

406 In conclusion, yield-invigorating soils displayed a higher content of organic matter and harboured unique  
407 bacterial community with greater diversity than yield-debilitating soils. Further Chloroflexota was  
408 significantly enriched and identified as a potential keystone taxon in manipulating the interaction of  
409 bacterial community in yield-invigorating soils. These findings indicated that soil organic matter  
410 triggered the assembly of soil microbiome, which both participated in maintaining crop production . Such  
411 knowledge is a first step toward harnessing soil microbiome in support of sustainable agroecosystems.

412

413 **Data availability.** Raw amplicon sequencing data for each sample used in this study was deposited at  
414 the National Center for Biotechnology Information (NCBI) in the FASTQ format and is available under  
415 the accession number PRJNA749397. Other data that support the findings of this study are available on  
416 request from the corresponding author (Xiaomei Ye).

417

## 418 **Authors' contributions**

419 L. Wang: performed all experiments; L. Wang, X. Ye, and Z. Shen: designed the study, and wrote the

420 majority of the manuscript; L. Wang and Z. Shen and C. Tao: analyzed the data; H. Hu, J. Du, Y. Xi, J.  
421 Lin, and D. Chen: participated in the design of the study, provided comments and edited the manuscript.  
422 The authors read and approved the final manuscript.

423

424 **Competing interests.** The authors declare that they have no conflict of interest.

425

426 **Disclaimer.** Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional  
427 claims in published maps and institutional affiliations.

428

429 **Acknowledgements.** We sincerely thank all those who have assisted me with any part of this paper and  
430 all pear orchards owners for providing access to the soil sampling.

431

432 **Financial support.** This research was supported by Jiangsu Agricultural Science and Technology  
433 Innovation Fund (CX(19)3094) and National Natural Science Foundation of China (31801842 and  
434 42090065).

435

#### 436 **Appendix A. Supplementary data**

437 Supplementary figures and tables to this article can be found in the supplemental material.

438

#### 439 **References**

440 Banerjee, S., Schlaeppli, K., van der Heijden, M.G.J: Keystone taxa as drivers of microbiome structure  
441 and functioning. *Nat. Rev. Microbiol.*, 16, 567-576, 2018.

442 Barrios, E.: Soil biota, ecosystem services and land productivity. *Ecological Eco.*, 64, 269-285, 2007.

443 Bell, T., Newman, J.A., Silverman, B.W., Turner, S.L., Lilley, A.K.: The contribution of species richness  
444 and composition to bacterial services. *Nature*, 436, 1157-1160, 2005.

445 Bender, S.F., Wagg, C., van der Heijden, M.G.A.: An underground revolution: biodiversity and soil  
446 ecological engineering for agricultural sustainability. *Trends Ecol Evol.*, 31, 440-452, 2016.

447 Bolger, A.M., Lohse, M., Usadel, M.: Trimmomatic: a flexible trimmer for Illumina sequence data.  
448 *Bioinformatics*, 30, 2114-2120, 2014.

449 Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., Alexander, H.,

450 Alm, E.J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J.E., Bittinger, K., Brejnrod, A., Brislawn,  
451 C.J., Brown, C.T., Callahan, B.J., Caraballo-Rodríguez, A.M., Chase, J., Cope, E.K., et al.:  
452 Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat.*  
453 *Biotechnol.*, 37, 852-857, 2019.

454 Boulesteix, A.L., Janitza, S., Kruppa, J., König, I.R.: Overview of random forest methodology and  
455 practical guidance with emphasis on computational biology and bioinformatics. *Wires Data Min.*  
456 *Knowl.*, 2, 493-507, 2012.

457 Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A., Turnbaugh, P.J., Fierer,  
458 N., Knight, R.: Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample.  
459 *P. Nat. Acad. Sci. USA*, 108, 4516-4522, 2011.

460 Cardinale, B.J., Srivastava, D.S., Duffy, J.E., Wright, J.P., Downing, A.L., Sankaran, M., Jouseau, C.:  
461 Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature*, 443, 989-992,  
462 2006.

463 Chaer, G., Fernandes, M., Myrold, D., Bottomley, P.: Comparative resistance and resilience of soil  
464 microbial communities and enzyme activities in adjacent native forest and agricultural soils. *Microb*  
465 *Ecol.*, 58, 414-424, 2009.

466 Chaparro, J.M., Sheflin, A.M., Manter, D.K., Vivanco, J.M.: Manipulating the soil microbiome to  
467 increase soil health and plant fertility. *Biol. Fert. Soils*, 48, 489-499, 2012.

468 Chen, L., Jiang, Y., Liang, C., Luo, Y., Xu, Q., Han, C., Zhao, Q., Sun, B.: Competitive interaction with  
469 keystone taxa induced negative priming under biochar amendments. *Microbiome*, 7, 77, 2019.

470 Coyte, K.Z., Schluter, J., Foster, K.R.: The ecology of the microbiome: networks, competition, and  
471 stability. *Science*, 350, 663-666, 2015.

472 Dai, Z., Su, W., Chen, H., Barberán, A., Zhao, H., Yu, M., Yu, L., Brookes, P.C., Schadt, C.W., Chang,  
473 S.X., Xu, J.: Long-term nitrogen fertilization decreases bacterial diversity and favors the growth of  
474 Actinobacteria and Proteobacteria in agro-ecosystems across the globe. *Global Change Biol.*, 24,  
475 3452-3461, 2018.

476 Deng, Y., Jiang, Y., Yang, Y., He, Z., Luo, F., Zhou, J.: Molecular ecological network analyses. *BMC*  
477 *Bioinformatics*, 13, 1-20, 2012.

478 Ding, J., Zhang, Y., Wang, M., Sun, X., Cong, J., Deng, Y., Lu, H., Yuan, T., van Nostrand, J.D., Li, D.,  
479 Zhou, J., Yang, Y.: Soil organic matter quantity and quality shape microbial community



480 compositions of subtropical broadleaved forests. *Mol. Ecol.*, 24, 5175-5185, 2015.

481 Edwards, J., Johnson, C., Santos-Medellín, C., Lurie, E., Podishetty, N.K., Bhatnagar, S., Eisen, J.A.,  
482 Sundaresan, V.: Structure, variation, and assembly of the root-associated microbiomes of rice. P.  
483 *Nat. Acad. Sci. USA*, 112, E911-E920, 2015.

484 Eo, J., Park, K.: Long-term effects of imbalanced fertilization on the composition and diversity of soil  
485 bacterial community. *Agr. Ecosyst. Environ.*, 231, 176-182, 2016.

486 Fan, K., Delgado-Baquerizo, M., Guo, X., Wang, D., Zhu, Y., Chu, H.: Biodiversity of key-stone  
487 phylotypes determines crop production in a 4-decade fertilization experiment. *ISME J.*, 15, 550-  
488 561, 2021.

489 Fath, B.D., Scharler, U.M., Ulanowicz, R.E., Hannon, B.: Ecological network analysis: network  
490 construction. *Ecol. Model.*, 208, 49-55, 2007.

491 Faust, K., Raes, J.: Microbial interactions: from networks to models. *Nat. Rev. Microbiol.*, 10, 538-550,  
492 2012.

493 Fierer, N., Bradford, M.A., Jackson, R.B.J.: Toward an ecological classification of soil bacteria. *Ecology*,  
494 88, 1354-1364, 2007.

495 Fierer, N., Jackson, R.B.J.: The diversity and biogeography of soil bacterial communities. P. *Nat. Acad.*  
496 *Sci. USA*, 103, 626-631, 2006.

497 Fierer, N.: Embracing the unknown: disentangling the complexities of the soil microbiome. *Nat. Rev.*  
498 *Microbiol.*, 15, 579-590, 2017.

499 Frąc, M., Hannula, S.E., Bełka, M., Jędrzycka, M.: Fungal biodiversity and their role in soil health. *Front.*  
500 *Microbiol.*, 9, 707, 2018.

501 Fuerst, J.A.: Planctomycetes—new models for microbial cells and activities. I. Kurtböke (Ed.), *Microbial*  
502 *Resources*, Elsevier/Academic Press, 1-27, 2017.

503 Grace, J., José J.S., Meir, P., Miranda, H.S., Monteset R.A.: Productivity and carbon fluxes of tropical  
504 savannas. *J Biogeogr.*, 33, 387-400, 2006.

505 Hu, Q., Tan, L., Gu, S., Xiao, Y., Xiong, X., Zeng, W., Feng, K., Wei, Z., Deng, Y.: Network analysis  
506 infers the wilt pathogen invasion associated with non-detrimental bacteria. *npj Biofilms Microb.*, 6,  
507 1-8, 2020.

508 Huang, Z., Zhao, F., Wang, M., Qi, K., Wu, J., Zhang, S.: Soil chemical properties and geographical  
509 distance exerted effects on arbuscular mycorrhizal fungal community composition in pear orchards

510 in Jiangsu Province, China. *Appl. Soil Ecol.*, 142, 18-24, 2019.

511 Hug, L.A., Castelle, C.J., Wrighton, K.C., Thomas, B.C., Sharon, I., Frischkorn, K.R., Williams, K.H.,  
512 Tringe, S.G., Banfield, J.F.: Community genomic analyses constrain the distribution of metabolic  
513 traits across the Chloroflexi phylum and indicate roles in sediment carbon cycling. *Microbiome*, 1,  
514 1-17, 2013.

515 Jiang, Y., Luan, L., Hu, K., Liu, M., Chen, Z., Geisen, S., Chen, X., Li, H., Xu, Q., Bonkowski, M., Sun,  
516 B.: Trophic interactions as determinants of the arbuscular mycorrhizal fungal community with  
517 cascading plant-promoting consequences. *Microbiome*, 8, 1-14, 2020.

518 Kallenbach, C.M., Frey, S.D., Grandy, A.S.J.: Direct evidence for microbial-derived soil organic matter  
519 formation and its ecophysiological controls. *Nat. Commun.*, 7, 1-10, 2016.

520 Kennedy, A.C., Smith, K.J.: Soil microbial diversity and the sustainability of agricultural soils. *Plant Soil*,  
521 170, 75-86, 1995.

522 Kolde, R., Kolde, M. Package 'pheatmap'. 2018, [https://cran.r-](https://cran.r-project.org/web/packages/pheatmap/index.html)  
523 [project.org/web/packages/pheatmap/index.html](https://cran.r-project.org/web/packages/pheatmap/index.html).

524 Langfelder, P., Horvath, S.J.: Eigengene networks for studying the relationships between co-expression  
525 modules. *BMC Syst. Biol.*, 1, 1-17, 2007.

526 Liaw, A., Wiener, M.: Classification and regression by randomForest. *Rnews*, 2, 18-22, 2002.

527 Lin, J., Sheng, B., Li, X., Yang, Q., Wang, Z., Li, H., Wang, H., Cheng, Y.: A new pyrus pyrifolia cultivar  
528 'Sucui 1'. *Acta Hort. Sin.*, 40, 1849-1850, 2013 (In Chinese)

529 Lu, L., Yin, S., Liu, X., Zhang, W., Gua, T., Shen, Q., Qiu, H.: Fungal networks in yield-invigorating  
530 and-debilitating soils induced by prolonged potato monoculture. *Soil Biol. Biochem.*, 65, 186-194,  
531 2013.

532 Magoč, T., Salzberg, S. L.: FLASH: fast length adjustment of short reads to improve genome assemblies.  
533 *Bioinformatics*, 27, 2957-2963, 2011.

534 Mei, W., Yu, G., Lai, J., Rao, Q., Umezawa, Y.: basicTrendline: add trendline and confidence interval of  
535 basic regression models to plot. R package version 2.0.3, 2018, [http://cran.r-](http://cran.r-project.org/package=basicTrendline)  
536 [project.org/package=basicTrendline](http://cran.r-project.org/package=basicTrendline).

537 Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlenn, D., Minchin, P.R., O'Hara,  
538 R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H.: Package 'vegan'.  
539 Community ecology package, version 2, 2013, <http://CRAN.R-project.org/package=vegan>.

540 Parks, D.H., Tyson, G.W., Hugenholtz, P., Beiko, R.G.: STAMP: statistical analysis of taxonomic and  
541 functional profiles. *Bioinformatics*, 30, 3123-3124, 2014.

542 Pathan, S.I., Roccotelli, A., Petrovičová, B., Romeo, M., Badagliacca, G., Monti, M., Gelsomino, A..  
543 Temporal dynamics of total and active prokaryotic communities in two Mediterranean orchard soils  
544 treated with solid anaerobic digestate or managed under no-tillage. *Biol. Fert. Soils.*, 57, 837-861,  
545 2021.

546 Philippot, L., Raaijmakers, J.M., Lemanceau, P., van der Putten, W.H.: Going back to the roots: the  
547 microbial ecology of the rhizosphere. *Nat. Rev. Microbiol.*, 11, 789-799, 2013.

548 Qiao, C., Penton, C.R., Xiong, W., Liu, C., Wang, R., Liu, Z., Xu, X., Li, R., Shen, Q.: Reshaping the  
549 rhizosphere microbiome by bio-organic amendment to enhance crop yield in a maize-cabbage  
550 rotation system. *Appl. Soil Ecol.*, 142, 136-146, 2019.

551 Raaijmakers, J.M., Mazzola, M.J.: Soil immune responses. *Science*, 352, 1392-1393, 2016.

552 Rosseel, Y.: Lavaan: An R package for structural equation modeling and more. *J. Stat. Softw.*, 48, 1-36,  
553 2012.

554 Rousk, J., Bååth, E., Brookes, P.C., Lauber, C.L., Lozupone, C., Caporaso, J.G., Knight, R., Fierer, N.:  
555 Soil bacterial and fungal communities across a pH gradient in an arable soil. *ISME J.*, 4, 1340-1351,  
556 2010.

557 Schermelleh-Engel, K., Moosbrugger, H., Müller, H.J.: Evaluating the fit of structural equation models:  
558 Tests of significance and descriptive goodness-of-fit measures. *Methods Psychol. Res.*, 8, 23-74,  
559 2003.

560 Schloss, P.D., Westcott, S.L., Ryabin, T., Hall, J.R., Hartmann, M., Hollister, E.B., Lesniewski, R.A.,  
561 Oakley, B.B., Parks, D.H., Robinson, C.J., Sahl, J.W., Stres, B., Thallinger, G.G., Van Horn, D.J.,  
562 Weber, C.F.: Introducing mothur: open-source, platform-independent, community-supported  
563 software for describing and comparing microbial communities. *Appl. Environ. Microbiol.*, 75,  
564 7537-7541, 2009.

565 Shen, Z., Penton, C.R., Lv, N., Xue, C., Yuan, X., Ruan, Y., Li, R., Shen, Q.: Banana Fusarium wilt  
566 disease incidence is influenced by shifts of soil microbial communities under different monoculture  
567 spans. *Microb. Ecol.*, 75, 739-750, 2018.

568 Shen, Z., Zhong, S., Wang, Y., Wang, B., Mei, X., Li, R., Ruan, Y., Shen, Q.: Induced soil microbial  
569 suppression of banana fusarium wilt disease using compost and biofertilizers to improve yield and

570 quality. Eur. J. Soil Biol., 57, 1-8, 2013.

571 Smoot, M.E., Ono, K., Ruscheinski, J., Wang, P.L., Ideker, T.: Cytoscape 2.8: new features for data  
572 integration and network visualization. Bioinformatics, 27, 431-432, 2011.

573 Speirs, L.B.M., Rice, D.T.F., Petrovski, S., Seviour, R.J.: The phylogeny, biodiversity, and ecology of the  
574 *Chloroflexi* in activated sludge. Front Microbiol., 10, 2015, 2019.

575 Tian, J., He, N., Hale, L., Niu, S., Yu, G., Liu, Y., Blagodatskaya, E., Kuzyakov, Y., Gao, Q., Zhou, J.:  
576 Soil organic matter availability and climate drive latitudinal patterns in bacterial diversity from  
577 tropical to cold temperate forests. Funct. Ecol., 32, 61-70, 2018.

578 Toju, H., Peay, K.G., Yamamichi, M. Narisawa, K., Hiruma K., Naito K., Fukuda S., Ushio M., Nakaoka  
579 S., Onoda Y., Yoshida K., Schlaeppi K., Bai Y., Sugiura R., Ichihashi Y., Minamisawa K., Kiers &  
580 E.T.: Core microbiomes for sustainable agroecosystems. Nat Plants, 4, 247–257, 2018.

581 van der Heijden, M.G., Bardgett, R.D., van Straalen, N.M.J.: The unseen majority: soil microbes as  
582 drivers of plant diversity and productivity in terrestrial ecosystems. Ecol. Lett., 11, 296-310, 2008.

583 Wang, C., Liu, D., Bai, E.: Decreasing soil microbial diversity is associated with decreasing microbial  
584 biomass under nitrogen addition. Soil Biol. Biochem., 120, 126-133, 2018.

585 Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R.: Naive Bayesian classifier for rapid assignment of rRNA  
586 sequences into the new bacterial taxonomy. Appl. Environ. Microbiol., 73, 5261-5267, 2007.

587 Xun, W., Li, W., Xiong, W., Ren, Y., Liu, Y., Miao, Y., Xu, Z., Zhang, N., Shen, Q., Zhang, R.: Diversity-  
588 triggered deterministic bacterial assembly constrains community functions. Nat. Commun., 10, 1-  
589 10, 2019.

590 Zhao, J., Ni, T., Li, Y., Xiong, W., Ran, W., Shen, B., Shen, Q., Zhang, R.: Responses of bacterial  
591 communities in arable soils in a rice-wheat cropping system to different fertilizer regimes and  
592 sampling times. PloS One, 9, e85301, 2014.


593 Zhong, Y., Hu, J., Xi, Q., Zhang, S., Li, X., Pan, X., Zhao, R., Wang, R., Yang, W., Shangguan, Z., Hu,  
594 F., Yang, C., Wang, W.. Soil microbial mechanisms promoting ultrahigh rice yield. Soil Biol.  
595 Biochem., 143: 107741, 2020.

596

597 **Figure legends**

598 **Fig. 1 Distribution of studied field sites.** Map showing the sites of six pair-located orchards sampled in  
599 this study.

600



601

602


603 **Fig. 2 Quantitation of the abundance of bacteria population, and linkage of microbial composition**

604 **to pear yield.** (A) Violin plot showing the abundance of total bacteria for all selected orchards. \* indicates

605 a significant difference between yield-invigorating (YI) and yield-debilitating (YD) orchards based on

606 Wilcoxon tests ( $p < 0.05$ ). (B) Correlation plot showing the relationship of microbial composition and


607 yield based on braycurtis distances calculated by Mantel test.



608


609

610 **Fig. 3 Overview of bacterial composition and alpha diversity.** (A) Principal Coordinates Analysis  
 611 (PCoA) plot displaying the bacterial community composition calculated based on braycurtis distances.  
 612 (B) Violin plot showing the alpha diversity indices (Chao and Shannon) for all selected orchards. \*  
 613 indicates a significant difference between yield-invigorating (YI) and yield-debilitating (YD) orchards  
 614 based on Wilcoxon tests ( $p < 0.05$ ). (C) Venn plot depicting the unique and shared bacterial ASVs  
 615 between yield-invigorating (YI) and yield-debilitating (YD) orchards at ASV and sequence insights.  
 616 Uni. YI and Uni. YD represent unique ASVs or sequences in the YI or YD soils while Shared represent  
 617 shared ASVs or sequences between the YI and YD soils. (D) Heatmap displaying the composition of  
 618 unique and shared ASVs at phylum level in YI and YD soils. Numbers in the cell represent the number  
 619 of ASVs affiliated to that phylum.



620  
 621  
 622  
 623

624 **Fig. 4 Key taxonomic groups in distinguishing yield-invigorating (YI) and yield-debilitating (YD)**  
 625 **orchards.** Stacked bar chart (left panel) showing dominant phyla affiliation in YI and YD soils for six  
 626 pair-located sites while horizontal histogram (right panel) depicting relative changes of dominant phyla  
 627 in YI soils compared to those in YD soils.




628

629

630

631

632 **Fig. 5 Co-occurrence networks of bacterial community and identified keystone taxa in**  
633 **distinguishing yield-invigorating (YI) and yield-debilitating (YD) orchards.** (A) An overview of  
634 microbial phylogenetic molecular ecological networks constructed from YI and YD soils. Line with blue  
635 color indicates positive correlations whereas lines with red color signifies negative correlations in each  
636 network. Modules containing larger than five nodes in the networks are labeled with corresponding letter  
637 followed by a number. Circular node colors indicate different bacterial phyla. (B) Stacked figure showing  
638 the relative abundance of nodes in each module within each network at the phylum level. (C) Venn plot  
639 depicting the unique and shared bacterial ASVs between two networks construed from YI and YD soils.  
640 Left panel is plotted based on the original nodes used in building network while right panel is plotted  
641 based on the nodes from modules. Uni. YI and Uni. YD represent unique ASVs in the YI or YD networks  
642 while Shared represent shared ASVs between the YI and YD networks. (D) Zi-Pi plot showing the  
643 distribution of nodes based on their topological roles. The threshold values of Zi and Pi for categorizing  
644 OTUs were 2.5 and 0.62 respectively. Node colors indicate different bacterial phyla and node size  
645 represent the relative abundance in each network.




646

647




648 **Fig. 6 Relationships among bacterial community, soil edaphic factors and pear yield.** (A) Variance  
 649 partitioning analysis (VPA) map of the effects of soil edaphic properties, sample locations, pear yield  
 650 and interactions of these factors on the bacterial community. (B) Redundancy analysis (RDA) plot  
 651 showing the relationships among all assigned bacterial ASVs and measured soil edaphic properties for  
 652 all soils after stepwise selection. (C) Random forest mean predictor importance of selected soil edaphic  
 653 properties used as drivers in predicting the pear yield. Red bar indicates that the given predictor is  
 654 significant while black bar indicates that the given predictor is non-significant. The %IncMSE in the up  
 655 panel means the increase in mean squared error while IncNodePurity indicates the increase in node  
 656 purity. The values of these two indices represent the importance of each variable to predict the module.  
 657 A larger value indicates that the variable is more important. (D) Correlation plot showing the  
 658 relationship of microbial composition and soil organic matter based on braycurtis distances calculated  
 659 by Mantel test.



660  
 661  
 662

663 **Fig. 7 Structural equation modeling (SEM) describing the biotic and abiotic factors in affecting**  
 664 **the crop production.** Structural equation model was built incorporating soil organic matter, microbial  
 665 biomass, beta diversity (PCoA), key taxa, network hubs including module hubs and network  
 666 connectors, and yield. The path analysis numbers adjacent to arrows indicate the relationship's effect  
 667 size and the associated bootstrap *P*-value. Cyan and red arrows indicate positive and negative  
 668 relationships, respectively. Paths with non-significant coefficients are presented as gray lines.



669  
 670  
 671