

# Tropical Andosol organic carbon quality and degradability in relation to soil geochemistry as affected by land use

Sastrika Anindita<sup>1</sup>, Peter Finke<sup>1</sup>, Steven Sleutel<sup>1</sup>

<sup>1</sup>Department of Environment, Ghent University, Ghent, 9000, Belgium

5 *Correspondence to:* Sastrika Anindita (sastrika.anindita@ugent.be)

**Abstract.** Land use is recognized to impact soil geochemistry on the centennial to millennial time scale, with implications for the distribution and stability of soil organic carbon (SOC). Juvenile volcanic soils in tropical areas are subject to much faster pedogenesis, with then also possibly a significant mediation by land use on much shorter centennial or even decadal scale. Very scarce observational evidence exists and so such indirect implications of land use on SOC cycling are largely unknown.

10 We here investigated SOC fractions, substrate specific mineralization (SOC or added plant residue), and net priming of SOC in function of forest or agricultural land use on Indonesian volcanic soils. The content of oxalate-extracted Al ( $Al_o$ ) and exchangeable Ca correlated well with OC associated with sand-sized aggregates. The concomitant ~~near doubling~~more than tripling of the proportion of SOC in sand-sized 400 J ml<sup>-1</sup> ultrasonication resistant aggregates compared to forest and likewise contrasts in  $Al_o$  ~~and Ca~~ suggest ~~an that~~ enhanced formation of Al-(hydr)oxides ~~and liming~~ promoted aggregation and physical  
15 occlusion of OC. This was importantly also consistent with a relatively lesser degradability of SOC in the agricultural sites, though the net priming of SOC and degradability of added <sup>13</sup>C-labelled ryegrass was not found to depend on land use. We expected that ~~-amorphous Al~~ content under agricultural land use would mainly have promoted mineral-association of SOC compared to under pine forest but found no indications for this. Improved small scale aggregation of tropical Andosols caused by conversion to agriculture and high carbon input via organic fertilizer may thus partially counter the otherwise expectable  
20 decline of SOC stocks following cultivation. Such indirect land use effects on the SOC balance appeared relevant for correct interpretation and prediction of the long-term C-balance of (agro)ecosystems with soil subject to intense development, like the here studied tropical Andosols.

## 1 Introduction

Storage of soil organic carbon (SOC) in terrestrial ecosystems can improve ecosystem services such as soil health, agricultural  
25 productivity, and climate change mitigation (Baldock, 2007; Lehmann and Kleber, 2015). The storage of SOC is influenced by the interaction of ecological processes with net primary productivity and heterotrophic respiration usually being most important terms of the SOC balance. Land use determines the net-C input to soil and in doing so often bears an overriding control on the SOC balance, compared to other drivers of the C-cycle at ecosystem level. Forest ecosystems usually have relatively high net primary productivity compared to agricultural land use (Smith et al., 2014). The net result of cultivation

30 then typically leads to a decline in SOC stock and increased human-induced CO<sub>2</sub> emission (Don et al., 2011; Wei et al., 2014).  
The long-term SOC balance of ecosystems also depends on the capacity of soils to stabilize newly entering OC against  
microbial decomposition, resulting from heterotrophic respiration. Physical occlusion of OC at the microaggregate scale as  
well as binding of OC onto reactive mineral surfaces constitute two major mechanisms for stabilization of SOC (Matus et al.,  
2014). Soil characteristics such as content of poorly crystalline minerals, specific surface area, clay content and degree of soil  
35 micro-aggregation have all been linked to the capacity of soils to accumulate SOC (Hernández et al., 2012; Kleber et al., 2015;  
Mikutta et al., 2006, 2009; Poirier et al., 2020). While soil mineralogy itself is thus a well-recognized controller of SOC  
stabilization, surprisingly little attention has gone into understanding how land use history, through its impact on soil  
geochemistry might in turn indirectly control stability of SOC and organic matter (OM) entering soil.

40 The geochemistry of the reactive mineral phase of a soil is the resultant of both the composition of its parent material and soil  
weathering status (Mikutta et al., 2009). Weathering status is crucially driven by the time since the onset of weathering, local  
climate and hydrologic conditions, resilience of minerals to weathering and vegetation cover (Doetterl et al., 2015). Land use,  
through its control on vegetation, can impact the weathering process by modifying the pH, influencing soil biological activity  
and nutrient levels, and releasing organic complexing compounds (Van Breemen et al., 1983; Cronan, 2018). On agricultural  
45 lands, moreover fertilizer addition and soil disturbance caused by tillage have also been reported to significantly enhance  
weathering (Churchman and Lowe, 2012; Li et al., 2021; Taylor et al., 2016). While an extensive and still growing number of  
studies considered land use impacts on SOC stock, soil OM quality and its degradability (Covaleda et al., 2011; Cusack et al.,  
2013; Huygens et al., 2005), just very few of have considered the potential indirect impact of land use on SOC stability through  
its effect on the soil mineral phase. Moreover, the coverage remains limited for several soil groups, including tropical Andosols.

50 A recent study on the Gálápagos islands demonstrated that even ~~after maximum of fifteen years within a few centuries to just  
several decades~~ conversion of native forest to cultivated land strikingly accelerated soil weathering (Gerzabek et al., 2019). In  
particular, formation of secondary minerals should impact SOC retention in volcanic soils as their high capacity to do so is  
acknowledged to emerge from abundantly present poorly crystalline Al and Fe oxides. Indeed, Asano and Wagai (2014)  
concluded the importance of short-range order minerals or organo-Al complexes for SOC stabilization in Andosols from  
55 correlations with OC stock (Miyazawa et al., 2013), mean residence time (reviewed by Parfitt (2009)), and by chemical  
characterization of organo-mineral associations (Basile-Doelsch et al., 2007; Mikutta et al., 2009). Next to a large specific  
surface area, also microaggregation and occlusion therein of intra-microaggregate particulate OM (iPOM) is known to grant a  
degree of protection against its microbial decay (Six et al., 2000a), particularly in Andosols (Asano et al., 2018; Asano and  
Wagai, 2014). Introduction of tillage and removal of permanent vegetative coverage usually adversely affect iPOM-induced  
60 physical protection. However, differences between untilled native vegetation or secondary forest on the one hand and tilled  
cropland might not be significant for volcanic soils (Dörner et al., 2012; Huygens et al., 2005; Linlin et al., 2016) as abundant  
nano sized organo-mineral composites overridingly act as binding agents in microaggregates in Andosols (Asano and Wagai,  
2015).

65 In Indonesia, the Sunda volcanic arc region (Mt. Tangkuban Perahu and Mt. Burangrang) in West Java, was covered by native forest until only about two centuries back but has since then been largely replaced by secondary pine forest or agricultural land use. With differently aged volcanic deposits and co-occurrence of native forest, secondary forest and cultivated land this region forms a useful means to understand how Andosol development would impact SOC storage and stability. A first objective was to further investigate how cultivation history in this region would have resulted in differential soil aggregation and levels of  
70 iPOM, and mineral associated OM vs. free relatively unprotected OM. We did so by assessing soil OM fractions for a set of 6 Indonesian volcanic soil (Cambisols and Andosols) with native forest, secondary pine forest or agricultural land use. A second goal was to infer how OM degradability in the topsoil of these tropical volcanic soils itself differs in function of forest vs. agricultural land use. Assessing the indirect effect of land use on SOC degradability through its mediation of soil mineralogy is, however, complicated by the fact that quality of native SOC itself is also function of land use. We therefore assessed the  
75 degradability of a single model exogenous OM source (<sup>13</sup>C-labelled ryegrass residues) to see how land use changes affect stability of OC in soil. We hypothesize that the enhanced formation of pedogenic poorly crystalline Al under agricultural land use, as we previously confirmed (Anindita et al., 2022)(~~Anindita et al., under review~~), results in a relative stabilization of the exogenous OM as compared to native forest or secondary forest. In doing so we take account for the well-known phenomenon that labile OC-inputs can accelerate the decomposition of native SOC, a process called positive priming (Chen et al., 2014),  
80 or conversely restrain decomposition of SOC and induce negative priming (Blagodatskaya et al., 2014; Qiao et al., 2014). By using a C-isotope mixing model, mineralization of the exogenous OM (i.e. ryegrass) and native SOC are distinguished in the soil incubation experiments.

## 2 Materials and methods

### 2.1 Site description and soil sampling

85 Our study covered soils from the Mount Tangkuban Perahu and Mount Burangrang regions that are part of the Sunda Volcanic complex in West Java, Indonesia. Six sites were selected to represent the dominant land use types in the area, viz. primary forest (NF-y), pine forest (PF-i1 and PF-o2) and horticulture (AG-y1, AG-i2, and AG-o3). The mean annual temperature in this region study sites is 19–21°C and mean annual precipitation ranges around 2000–~~3000000~~ mm per year. ~~Six sites were selected to represent the dominant land use types in the area, viz. primary forest (NF), pine forest (PF1 and PF2) and horticulture (AG1, AG2, and AG3).~~ Soils at sites NF-y and PF-i1 are andic Cambisols and at the PF-o2, AG-y1, AG-i2, AG-o3 sites aluandic Andosols (Anindita et al., 2022)(~~Anindita et al., under review~~). All soils contain silicate minerals (i.e. quartz, cristobalite, tridymite), gibbsite, albite, hornblende, 1:1 clay minerals, and non-crystalline materials (Anindita et al., 2022)(~~Anindita et al., under review~~). The proportion of primary minerals is higher in the younger NF-y soil as it is closer (within 1.5 km) to the crater of Mt. Tangkuban Perahu and received ash more recently. The pine forest and agricultural sites  
90 were originally all under the primary forest vegetation. At PF-i1 and PF-o2, secondary forests were planted in 1962 for

restoration purposes, following earlier deforestation. Agricultural land use at AG-y1, AG-i2, and AG-o3 dates back to just 30–50 years ago with mainly diverse horticultural crop rotations with crops potatoes (*Solanum tuberosum*), cabbage (*Brassica oleracea*), green bean (*Phaseolus vulgaris*), tomatoes (*Solanum lycopersicum*), and chayote (*Sechium edule*). Detailed information about land use history, and land management are given in (Anindita et al., 2022) Anindita et al. (under review).

100

At every site, soil samples were taken at 0–20, 20–40, and 60–80 cm depth. At each depth, the bulk samples were mixed into composite ones, which were then dried to the air, homogenised and sieved (< 2 mm) until further analysis at the Dept. of Environment of Ghent University (Belgium). The pH H<sub>2</sub>O was measured using a glass electrode pH meter in a 1:5 soil solution slurry. The estimated clay content was determined using the pipet method after full soil dispersion with Na<sup>+</sup>-resin (Bartoli et al., 1991), after testing several other procedures for soil texture analysis that proved incompatible with the investigated set of volcanic soils. The concentrations of total C and N were determined by dry combustion at 1100°C using a LECO 928 series CN-analyser. Aluminum extracted by 0.1 M pyrophosphate solution (Al<sub>p</sub>) was taken as measure of complexed Al to OM (McKeague, 1967). Iron and aluminum extractable by ammonium oxalate (Al<sub>o</sub>, Fe<sub>o</sub>) were used as estimates of X-ray amorphous Al and Fe. We previously quantified crystalline minerals using X-ray diffraction analysis. The amount of amorphous materials undetectable by X-ray diffraction was semi-quantitatively estimated from the difference between the obtained and the real amount of internal standard (Zincite 20%) after semi-quantification of crystalline minerals using BGMN Rietveld and Profex as user interface. Soil specific surface area (SSA) was assessed from adsorption-desorption of N<sub>2</sub> at 77°C following Brunauer–Emmett–Teller (BET) approach. Detailed methods and the results of geochemical analyses of the samples were documented in (Anindita et al., 2022) Anindita et al. (under review). Selected physicochemical soil properties are given in Table 1. More detailed mineralogical information of geochemical soil properties can be found in Supplementary material 1.

115

**Table 1. Physicochemical soil properties of 6 sampled soil profiles in the Sunda Volcanic complex in West Java, Indonesia (partly taken from Anindita et al. (2022) under review)**

Depth (cm)	pH H <sub>2</sub> O	BD <sup>a</sup> (g cm <sup>-3</sup> )	OC <u>stock<sup>a</sup></u> Mg ha <sup>-1</sup> g kg <sup>-1</sup>	δ <sup>13</sup> C ‰	C/N ratio	Clay %	Exch. Ca cmol kg <sup>-1</sup>	NH <sub>4</sub> -oxalate extractable		Al <sub>p</sub> <sup>a</sup> mg g <sup>-1</sup>	SSA <sup>a</sup> m <sup>2</sup> g <sup>-1</sup>	Amorphous materials g kg <sup>-1</sup>	Land use <u>&amp; estimated annual C-inputs<sup>b</sup></u>
								Al <sub>o</sub> <sup>a</sup>	Fe <sub>o</sub> <sup>a</sup>				
Primary forest (NF- <u>y</u> )													
0–20	4.1	0.77	<u>56.937.1</u>	-27.0	15.1	18.70	0.2	3.5	15.5	3.4	9	288	Natural vegetation ( <u>estimated</u>
20–40	4.4	0.78	<u>43.126.0</u>		20.2	19.81	0.1	2.3	25.4	3.5	27	254	<u>plant C input: 6.7 – 12.2 Mg ha<sup>-1</sup></u>
60–80	4.3	0.65	<u>37.428.6</u>		20.0	29.51	0.1	4.8	52.4	2.8	69	378	<u>yr<sup>-1</sup></u> )

Pine forest (PF- <u>i1</u> )													
0–20	4.5	0.64	<del>30.423.6</del>	-24.3	10.8	67.27	1.8	8.1	11.9	41.0	91	471	Pine forest
20–40	4.4	0.61	<del>15.913.0</del>		10.2	71.56	0.5	10.4	12.2	32.2	117	429	(estimated plant C input: 3.6 –
70–90	4.4	0.69	<del>8.15.9</del>		10.0	64.35	0.2	13.7	21.1	13.7	124	481	9.1 Mg ha <sup>-1</sup> yr <sup>-1</sup> )
Pine forest (PF- <u>o2</u> )													
0–20	4.7	0.50	<del>33.03.0</del>	-24.2	12.6	55.48	0.7	28.5	17.5	10.1	107	497	Pine forest
20–40	4.9	0.56	<del>20.418.2</del>		12.2	46.43	0.4	27.7	17.2	9.8	145	506	(estimated plant C input: 3.6 –
60–80	4.4	0.60	<del>18.85.7</del>		11.9	48.82	0.4	25.9	17.5	9.4	127	447	9.1 Mg ha <sup>-1</sup> yr <sup>-1</sup> )
Agriculture (AG- <u>y1</u> )													
0–20	5.1	0.67	<del>40.229.9</del>	-13.6	18.2	68.21	7.7	14.4	21.1	26.3	114	548	Intensive horticulture ± 50 years;
20–40	4.9	0.66	<del>17.313.0</del>		11.9	63.06	5.7	16.5	13.0	19.7	125	523	(estimated plant C input = 1.8
60–80	4.8	0.64	<del>14.411.2</del>		10.8	61.43	4.7	22.3	18.1	12.9	134	547	Mg ha <sup>-1</sup> yr <sup>-1</sup> , organic fertilizer C- input = 11.5 Mg ha <sup>-1</sup> yr <sup>-1</sup> ) organic fertilizer ± 76 Mg ha <sup>-1</sup> year <sup>-1</sup> )
Agriculture (AG- <u>i2</u> )													
0–20	5.4	0.66	<del>71.454.4</del>	-15.4	14.7	27.24	6.6	31.4	12.3	7.6	48	623	Intensive horticulture ± 40 years;
20–40	5.8	0.47	<del>39.642.6</del>		17.9	35.62	10.0	33.2	12.1	5.8	128	645	(estimated plant C input = 1.6
50–70	5.9	0.52	<del>28.126.9</del>		16.9	47.07	7.0	36.4	11.6	5.1	185	650	Mg ha <sup>-1</sup> yr <sup>-1</sup> , organic fertilizer C- input = 6.42 Mg ha <sup>-1</sup> yr <sup>-1</sup> ) organic fertilizer ± 20 Mg ha <sup>-1</sup> year <sup>-1</sup> )
Agriculture (AG- <u>o3</u> )													
0–20	5.4	0.65	<del>61.147.0</del>	-24.0	10.5	54.91	10.5	17.1	9.7	9.0	74	531	Horticulture < 30 years;
20–40	5.6	0.61	<del>35.028.8</del>		9.8	50.02	7.0	15.6	9.5	8.6	91	502	(estimated plant C input = 0.98
60–80	5.8	0.62	<del>9.77.8</del>		8.4	62.81	6.0	9.1	8.0	10.7	108	460	Mg ha <sup>-1</sup> yr <sup>-1</sup> , organic fertilizer C- input = 6.8 Mg ha <sup>-1</sup> yr <sup>-1</sup> ) organic fertilizer ± 18 Mg ha <sup>-1</sup> year <sup>-1</sup> )

120 <sup>a</sup>BD = bulk density; OC = organic carbon; Al<sub>o</sub> = aluminum extracted by NH<sub>4</sub>-oxalate; Fe<sub>o</sub> = iron extracted by NH<sub>4</sub>-oxalate; Al<sub>p</sub> = aluminum extracted by Na-pyrophosphate; SSA = specific surface area

<sup>b</sup> C-inputs from vegetation (forest or annual crops) and organic amendments. Forest C-inputs estimated from studies on Indonesian pine forest ((Bruijnzeel, 1985) ~~x~~) and Indonesian tropical primary forest ((Guillaume et al., 2018; Hertel et al., 2009) ~~y~~). Crop C-inputs based on farmer interviews and SSB & UGhent (2008). C-inputs from organic amendments were based on farmer interviews and lab-analyses.

## 125 2.2 Soil organic matter fractionation

Soils were fractionated using size, density and chemical separation steps according to [a modified version of a fractionation scheme proposed by Zimmermann et al. \(2007\)](#) (Supplementary 2 Fig. 2, Fig. S1). This procedure separates SOC into five

130 pools, namely into i) OC contained in soil aggregates and associated with sand (S+A), ii) in free particulate organic matter (POM), iii) in water dissolvable OC (DOC), iv) in oxidizable OC associated with the silt and clay (s+c – rSOC) fraction, and  
135 v) in a chemically resistant SOC (rSOC) pool. Briefly, 30 g of equivalent dry soil (< 2 mm) was initially dispersed in water by a calibrated ultrasonic probe with an output energy of 22 J ml<sup>-1</sup>. The resulting slurries were wet-sieved over a 63 µm sieve and rinsed with deionized water until the rinsing water became clear. In the original Zimmermann et al. (2007) method the > 63 µm fraction is further separated into S+A and POM based on density differences using sodium polytungstate at 1.8 g cm<sup>-3</sup>. However, several samples could not be well separated into free OM on the one hand and predominantly mineral material of  
140 the S+A fraction on the other. After pre-tests we instead used a 1.5 g cm<sup>-3</sup> sodium polytungstate solution to separate free POM from the S+A soil fraction, as per Cerli et al. (2012)'s advise to per soil optimize density cut-off for best separation of free and occluded POM. A subsample of the < 63 µm suspension (± 10 ml) derived from the wet sieving step was filtered through a 0.45 µm nylon membrane filter. The filtrate was analyzed for its dissolved OC concentration using a HT-I Formacs TOC-analyzer (Skalar, The Netherlands). The DOC pool was calculated by multiplying this OC concentration with the volume of  
145 flush water used in the preceding wet sieving step. The < 63 µm suspension was dried at 40°C and weighted to constitute the overall silt+clay (s+c) fraction. A chemically inert share of the SOC (rSOC) was isolated by subjecting the s+c fraction to oxidation by 6 % NaOCl. To this end, one gram of s+c material was weighted inside a 65 ml Nalgene centrifuge tube and 50 ml 6 % NaOCl adjusted to pH 8 was added and allowed to react for 18 h at 25°C. After centrifugation, remnant NaOCl was discarded by decantation. This procedure was repeated three times, the pellet was washed three more times with deionized  
150 water, transferred to a pre-weighted aluminum cup, dried at 40°C and weighted. The obtained S+A fraction remains composite as it contains both silt+clay associated OC next to occluded POM. In addition to the original scheme proposed by Zimmermann et al. (2007) and Poeplau et al. (2013) a further fractionation of the S+A into s+c and POM and sand-associated OM was included: A 3 5g subsample of the obtained S+A fraction was further dispersed by ultrasonication at 400 J ml<sup>-1</sup> and then separated into two size fractions by passing the obtained slurries over a 63µm sieve. The obtained <63µm fraction contains s+c OM and the >63µm fraction primarily occluded POM and sand-bound OM contained in S+A as well as OM contained in >63µm aggregates resistant to dispersal at 400 J ml<sup>-1</sup>. The S+A s+c fraction was further on considered as to constitute a single joint OM fraction with s+c obtained after the first 22 J ml<sup>-1</sup> and 63µm sieving step. All fractions were analyzed for their C concentration using a LECO 928 series CN-analyzer. ~~Further details are given in Zimmermann et al. (2007) and Poeplau et al. (2013).~~

### 155 2.3 Soil incubation experiment and isotopic signature measurements

Soils were incubated in a standardized way to compare degradability of a model plant-C substrate and SOC in the topsoil (0–20cm) of the 6 considered sites. Decomposition of either substrate was derived by regular measurement of the soil CO<sub>2</sub> efflux and inference of its δ<sup>13</sup>C signature. Soil mesocosms were prepared by repacking approximately 150 g of soil (depending on bulk density) into PVC tubes (diameter: 6.8 cm, height: 7 cm) to reach a height of 6.2 cm and bulk density as encountered at  
160 the field sites. Soil moisture content was set to 50 % water-filled pore space by addition of deionized water. A <sup>13</sup>C-labelled

plant-substrate, i.e. pulse-labelled ryegrass (*Lolium perenne*) with  $\delta^{13}\text{C}$  of +53.7 ‰, was applied at a dose of 1 g kg<sup>-1</sup> of dry matter. The contrast in  $\delta^{13}\text{C}$  with SOC (Table 1) allowed to distinguish emitted CO<sub>2</sub> into parts stemming from either native SOC or grass-C mineralization. The  $\delta^{13}\text{C}$  of topsoil (expressed as  $\delta^{13}\text{C}$  value (‰) vs. the international Vienna Pee Dee Belemnite standard) was measured using a PDZ Europe ANCA-GSL elemental analyser, interfaced with a Sercon 20–22 IRMS  
 165 with SysCon electronics (SetCon, Ceshire, UK).

Soils were incubated at 20°C for 120 days. On days 1, 3, 5, 8, 12, 17, 24, 32, 38, 47, 59, 73, 87, 101, 111, 120 soil CO<sub>2</sub> emission was inferred by measuring CO<sub>2</sub> build-up in a cylindrical closed-chamber attached consecutively on top of each PVC tube for at least 10 min. The evolution of the headspace CO<sub>2</sub> concentration and its  $\delta^{13}\text{C}$  was measured every 4 sec by connecting the  
 170 closed-chamber with a cavity ring-down spectrometer (G2201-*i* CRDS isotopic CO<sub>2</sub> analyser, Picarro, USA) in a loop via Teflon tubing. Soil CO<sub>2</sub> efflux rate in mg C kg<sup>-1</sup> h<sup>-1</sup> was calculated from the slope of the accumulating CO<sub>2</sub> concentration in function of time using the ideal gas law. The  $\delta^{13}\text{C}$  of emitted CO<sub>2</sub> was estimated from the y-axis intercept of derived Keeling plots (Keeling, 1958). The fraction of CO<sub>2</sub> derived from grass was calculated using the following equations introduced by Werth and Kuzyakov (2010):

$$175 \quad f_{\text{grass-CO}_2} = \frac{\delta^{13}\text{C}_{\text{total-CO}_2} - \delta^{13}\text{C}_{\text{SOC-CO}_2}}{\delta^{13}\text{C}_{\text{grass-CO}_2} - \delta^{13}\text{C}_{\text{SOC-CO}_2}} \quad (1)$$

$$\delta^{13}\text{C}_{\text{grass-CO}_2} = \delta^{13}\text{C}_{\text{grass}} - \epsilon_{\text{grass}} \quad (2)$$

$$\delta^{13}\text{C}_{\text{SOC-CO}_2} = \delta^{13}\text{C}_{\text{SOC}} - \epsilon_{\text{SOC}} \quad (3)$$

with  $\epsilon_{\text{SOC}}$  (in ‰) the net C-isotopic fractionation resulting from native SOC mineralization and diffusive CO<sub>2</sub> transport in soil until its efflux into the headspace air. The  $\epsilon_{\text{SOC}}$  was obtained from CO<sub>2</sub>-efflux monitoring parallel triplicate sets of control soils  
 180 without grass added. A third parallel set of soils were amended with a very high dose of grass (viz. 6 g kg<sup>-1</sup> added grass, i.e. about 4 t C ha<sup>-1</sup>) to estimate  $\epsilon_{\text{grass}}$ . This value was operationally calculated by subtracting  $\delta^{13}\text{C}_{\text{grass}}$  from the peak CO<sub>2</sub> emission  $\delta^{13}\text{C}$ -CO<sub>2</sub> of soil with 6 g kg<sup>-1</sup> grass under the assumption that virtually any emitted CO<sub>2</sub> was then derived from grass mineralization.

185 Using equations 1, 2 and 3 the fraction of emitted CO<sub>2</sub> derived from grass-C mineralization was calculated per soil and per point in time. The rate of grass-C mineralization ( $C_{\text{grass-min}}$ , in mg C kg<sup>-1</sup> h<sup>-1</sup>) for each measurement day was calculated by multiplying the  $f_{\text{grass}}$  value with the total CO<sub>2</sub> emission. The rate of SOC mineralization ( $C_{\text{SOC min}}$ , in mg C kg<sup>-1</sup> h<sup>-1</sup>) was calculated by subtracting the rate of  $C_{\text{grass-min}}$  from the total CO<sub>2</sub>-C emission rate. Cumulative CO<sub>2</sub> emissions derived from native SOC-C and grass-C were obtained by consecutively summing up  $C_{\text{SOC-CO}_2}$  and  $C_{\text{grass-CO}_2}$  respectively per time increment  
 190 between flux measurements. The % of mineralized  $C_{\text{SOC}}$  was then derived from dividing  $C_{\text{SOC-CO}_2}$  by  $C_{\text{SOC}}$ .



## 2.5 Statistics

Independent samples t-tests were used to compare both land uses (native and pine forest vs. agriculture, or alternatively agriculture vs. pine forests only) for the distribution of OC over soil fractions and for the cumulative amounts of grass-C or native SOC mineralized. Pearson correlation coefficients were used to investigate relations between soil carbon fraction proportions and geochemical properties. The non-parametric test was used if the data were not normally distributed or if the condition of homoscedasticity was unmet. Analyses were completed in IBM SPSS Statistics 27.

## 3 Results

### 3.1 Organic carbon in soil fractions

Across the studied soils a greater part of soil material ( $\pm 70.55\%$ ) was found as s+cS+A fraction (Table 2). The highest There were particularly large proportions of s+cS+A fraction occurred mainly in the PF-i, PF-o, and AG-o soils in the AG1, AG2, and topsoil PF1 soils (> 86.0 %). Conversely, in the PF2 and AG3-AG-i2 subsoils, the s+cS+A fraction was largest. The OC concentration of soil fractions decreased in the order: of POM > s+c (including rSOC) > S+A > DOC and was generally higher in the topsoil than subsoil. The OC concentration of the S+A fraction was higher under agricultural than under forest land use ( $p < 0.01$ ), particularly in topsoils of the sites under agricultural land use AG1 and AG3 ( $27.5-50.5$  g C kg<sup>-1</sup> fraction). No such difference in OC concentration existed between SOC in the s+c, POM, DOC and rSOC fractions under agriculture compared to forest (i.e., including primary forest). However, the OC concentration of s+c, s+c - rSOC, and rSOC fractions was higher ( $p < 0.01$ ) in the agricultural than when compared to the pine forests soils solely sites. The size of the DOC fraction was negligibly small ( $\pm 0.002$  g C kg<sup>-1</sup> fraction) and is further disregarded. The Total SOC content was not significantly different under forest ( $29.625.0 \pm 15.112.4$  g C kg<sup>-1</sup> soil) and agricultural ( $29.630.6 \pm 15.116.4$  g C kg<sup>-1</sup> soil) land use, but it was higher under agriculture than pine forest ( $16.318.4 \pm 8.97.7$  g C kg<sup>-1</sup> soil) ( $P < 0.015$ ).

Overall, the bulk of SOC was present in the s+cS+A ( $\pm 79.47\%$ ) with higher proportion in fraction, then followed by the rSOC ( $\pm 44.25\%$ ) than in the s+c - rSOC, s+c - rSOC ( $\pm 35.24\%$ ) fraction, and POM followed by S+A ( $+ 19. < 5\%$ ) and POM ( $+ 1\%$ ) fractions (Fig. 1). The rSOC constituted 34-67% of OC in s+c. The distribution of SOC across soil fractions varied to some extent between both land uses. The amount of OC in the S+A fraction was higher in the agricultural soils ( $11.9 \pm 10.13$  g C kg<sup>-1</sup> soil) than under forest ( $2.8 \pm 1.4$  g C kg<sup>-1</sup> soil;  $p < 0.01$ ) soils. Likewise, its relative contribution to SOC was larger in the agricultural soils (32 % of SOC), both when compared to all three forest sites (8% of SOC) (P < 0.01) or to just the two pine forest sites (9% of SOC) (P < 0.01). Particularly in topsoil of AG-y1 ( $\pm 13.23$  g C kg<sup>-1</sup> soil) and AG-i2 ( $3.17$  g C kg<sup>-1</sup> soil) a larger share of the SOC was in the S+A fraction. In contrast, more C was contained in free POM and in the s+c fraction under forest than under agricultural land use (p < 0.05). The s+c followed an opposite trend with a larger SOC proportion under forest (all three (89.5 % of SOC) or just the two pine forests (89.8 % of SOC)) than under agricultural land-use (68 % of SOC)



( $P < 0.05$ ). Out of the s+c ~~the~~ resistant SOC (rSOC) fraction accounted for approximately ~~a third to half~~ ~~half one quarter~~ of SOC and its ~~amount (in g C kg<sup>-1</sup> soil) and~~ proportion (44 % of SOC) was ~~likewise~~ higher under forest than ~~under~~ agricultural land use ( $p < 0.05$ ). Free POM ~~contributed on average less than 5 % of SOC and~~ constituted a ~~significantly~~ larger share of SOC in topsoil compared to both subsoil depth layers ( $p < 0.05$ ), ~~although this pool contributed on average less than 5 % of SOC. The occasional low POM C-concentrations (Table 2) indicate that some mineral matter was indicate that some mineral matter was left in the isolated POM fraction. However, given its very small mass proportion (median 0.1%) this artefact has negligible further impact on the overall distribution of SOC across soil fractions. The share of SOC proportion of in free POM (% of SOC) was also higher in under forest than agricultural land use ( $P < 0.015$ ).~~ -


230

**Table 2.** Mass proportion, C concentration and total content of SOC contained in soil fractions isolated by a modified version of the Zimmermann et al. (2007) method

Depth (cm)	Mass proportion (%)			C concentration (g kg <sup>-1</sup> fraction)					Amount of C (g kg <sup>-1</sup> soil)				
	POM	S+A	s+c	POM	S+A	s+c	DOC	POM	S+A	s+c	s+c - rSOC	rSOC	
Primary forest (NF-y)													
0–20	0.2 ± 0.1	37.6 ± 1.9	62.2 ± 2.0	412.5 ± 33.5	4.6 ± 3.6	31.1 ± 9.4	30.5 ± 3.3	1.0 ± 0.5	0.9 ± 0.3	1.7 ± 1.3	19.4 ± 6.3	18.9 ± 2.0	
20–40	1.1 ± 0.3	31.0 ± 1.1	67.9 ± 1.4	393.0 ± 0.0	2.9 ± 1.3	12.9 ± 2.4	24.9 ± 2.2	0.7 ± 0.7	4.4 ± 1.2	0.9 ± 0.4	8.8 ± 1.6	16.9 ± 1.7	
60–80	0.1 ± 0.1	27.3 ± 2.0	72.6 ± 2.0	367.5 ± 88.5	12.9 ± 4.8	16.0 ± 8.1	36.3 ± 1.5	0.8 ± 0.5	0.5 ± 0.4	3.6 ± 1.6	11.5 ± 5.9	26.3 ± 1.3	
Pine forest (PF-i)													
0–20	0.1 ± 0.1	9.2 ± 1.4	90.7 ± 1.5	227.0 ± 6.0	11.5 ± 0.4	12.2 ± 1.3	9.8 ± 1.0	1.9 ± 0.5	0.3 ± 0.2	1.1 ± 0.1	11.1 ± 1.3	8.9 ± 1.0	
20–40	0.1 ± 0.0	16.5 ± 2.4	83.5 ± 2.4	162.0 ± 9.5	11.6 ± 0.2	5.6 ± 0.5	7.1 ± 0.3	1.0 ± 0.4	0.1 ± 0.1	1.9 ± 0.3	4.7 ± 0.4	6.0 ± 0.4	
70–90	0.0 ± 0.0	7.1 ± 1.9	92.9 ± 1.9	103.6 ± 7.5	3.6 ± 0.1	2.9 ± 1.5	4.1 ± 0.2	0.4 ± 0.2	0.1 ± 0.0	0.3 ± 0.1	2.7 ± 1.4	3.8 ± 0.2	
Pine forest (PF-o)													
0–20	0.4 ± 0.1	16.3 ± 1.6	83.3 ± 1.4	188.3 ± 43.2	19.0 ± 1.5	12.9 ± 3.4	17.6 ± 1.3	1.7 ± 0.6	0.8 ± 0.1	3.1 ± 0.4	10.8 ± 2.9	14.6 ± 0.8	
20–40	0.2 ± 0.1	15.5 ± 3.7	84.3 ± 3.8	148.7 ± 47.3	10.3 ± 1.3	7.4 ± 1.1	10.4 ± 0.1	1.5 ± 0.4	0.2 ± 0.1	1.6 ± 0.6	6.2 ± 0.7	8.8 ± 0.3	
60–80	0.1 ± 0.1	11.5 ± 2.2	88.4 ± 2.2	135.6 ± 48.9	9.6 ± 0.8	3.3 ± 0.9	8.0 ± 0.3	1.1 ± 0.0	0.1 ± 0.0	1.1 ± 0.2	2.9 ± 0.8	7.1 ± 0.4	
Agriculture (AG-y)													
0–20	0.1 ± 0.0	39.6 ± 6.3	60.3 ± 6.3	206.0 ± 4.0	32.2 ± 0.7	12.7 ± 2.0	17.8 ± 0.1	1.5 ± 0.9	0.1 ± 0.5	12.8 ± 2.1	7.7 ± 1.6	10.7 ± 1.2	
20–40	0.0 ± 0.0	34.8 ± 1.5	65.2 ± 1.5	63.7 ± 0.9	15.5 ± 0.4	6.6 ± 0.5	9.5 ± 0.3	0.3 ± 0.1	0.02 ± 0.0	5.4 ± 0.2	4.3 ± 0.5	6.2 ± 0.1	
60–80	0.0 ± 0.0	30.5 ± 3.5	69.4 ± 3.5	15.6 ± 2.4	12.2 ± 0.3	5.8 ± 0.5	7.6 ± 0.2	0.4 ± 0.1	0.01 ± 0.0	3.7 ± 0.4	4.0 ± 0.5	5.2 ± 0.4	
Agriculture (AG-i)													
0–20	0.1 ± 0.0	45.7 ± 1.0	54.2 ± 1.1	350.0 ± 52.0	48.7 ± 2.4	26.7 ± 3.6	36.1 ± 3.2	0.6 ± 0.1	0.2 ± 0.1	22.3 ± 1.3	14.5 ± 2.1	19.6 ± 1.6	
20–40	0.0 ± 0.0	66.3 ± 1.4	33.7 ± 1.4	401.0 ± 88.0	46.2 ± 0.3	16.1 ± 2.3	31.4 ± 1.8	1.1 ± 0.7	0.04 ± 0.0	30.6 ± 0.8	5.5 ± 1.0	10.6 ± 2.6	
50–70	0.0 ± 0.0	64.1 ± 1.8	35.8 ± 1.8	587.5 ± 240	26.3 ± 0.1	10.4 ± 0.9	19.7 ± 0.7	0.8 ± 0.4	0.1 ± 0.0	16.9 ± 0.4	3.7 ± 0.3	7.1 ± 0.4	
Agriculture (AG-o)													

0-20	0.2 ± 0.0	9.2 ± 0.3	90.6 ± 0.3	420.7 ± 88.3	29.6 ± 2.6	21.7 ± 5.4	17.2 ± 0.8	1.8 ± 0.4	0.8 ± 0.3	2.7 ± 0.3	19.7 ± 5.0	15.6 ± 0.7
20-40	0.2 ± 0.0	8.2 ± 0.5	91.6 ± 0.5	311.3 ± 39.9	19.0 ± 0.9	14.4 ± 0.8	12.5 ± 1.1	1.3 ± 0.4	0.5 ± 0.1	1.6 ± 0.1	13.2 ± 0.8	11.5 ± 1.0
60-80	0.0 ± 0.0	4.6 ± 0.7	95.4 ± 0.7	92.6 ± 19.2	7.1 ± 1.2	7.0 ± 1.3	3.0 ± 0.3	1.2 ± 0.9	0.1 ± 0.0	0.3 ± 0.1	6.7 ± 1.3	2.8 ± 0.3

235



**Figure 1.** The distribution of OC over soil fractions obtained by a modified version of the Zimmerman et al. (2007) procedure for 6 sampled (0-60cm) soil profiles in the Sunda Volcanic complex in West Java, Indonesia. Vertical bars indicate standard deviations for three lab-replicates (n = 3)

240 **3.2 C:N ratio, amorphous aluminum and iron in the sand-aggregates (S+A) and silt-clay (s+c) fractions**

The ratio of C:N ranged from 6 – 20 in S+A and s+c fractions (Table 3) and their mean values did not differ between C:N ratio in S+A and s+c fractions. The C:N ratio in S+A fraction C:N-ratio was higher in the agriculture than in the forest soils ( $p < 0.05$ ), whereas this ratio was similar (non-significant) in s+c fraction. The content of  $Al_0$  and  $Fe_0$  varied between 1.5 to 57  $g\ kg^{-1}$  and 3.27-6 to 4774  $g\ kg^{-1}$  in the S+A and s+c fractions, respectively, whereas  $Fe_0$  ranged from 9.6 to 30  $g\ kg^{-1}$  and 7.6 to 74  $g\ kg^{-1}$  in the S+A and s+c fractions, respectively. A much lower  $Al_0$  content in the NF-y site s+c and S+A fractions compared to all other soils was in line with its distinctive total elemental composition previously reported by Anindita et al., (2022). The S+A  $Al_0$  content was higher under agriculture ( $32 + 12\ mg\ g^{-1}$ ) than pine forest ( $20\ mg\ g^{-1}$ ) (p

245

< 0.05) use. The s+c fraction Fe<sub>o</sub> content was higher under pine forest (21.6 + 4.7 mg g<sup>-1</sup>) than under agricultural land-use (17 + 7.2 mg g<sup>-1</sup>) (p < 0.01).

250

**Table 3.** Ratio of C:N, oxalate extractable--aluminiumaluminum (Al<sub>o</sub>) and iron (Fe<sub>o</sub>) in S+A and s+c fractions


Depth	S+A fraction			s+c fraction		
	C:N ratio	Al <sub>o</sub> mg kg <sup>-1</sup>	Fe <sub>o</sub>	C:N ratio	Al <sub>o</sub> mg kg <sup>-1</sup>	Fe <sub>o</sub>
<u>Primary forest (NF-y)</u>						
0-20	6.5 + 2.9	1.7 + 0.5	17.5 + 1.6	15.5 + 0.4	4.2 + 0.6	26.6 + 0.3
20-40	6.6 + 1.6	1.7 + 0.3	28.1 + 4.0	19.8 + 0.5	3.2 + 0.6	39.3 + 5.7
60-80	15 + 2.2	1.5 + 0.5	20.0 + 7.6	20.1 + 0.6	7.6 + 0.4	74.2 + 3.6
<u>Pine forest (PF-i)</u>						
0-20	9.5 + 0.2	11.0 + 0.6	19.9 + 1.3	9.3 + 0.7	11.1 + 0.6	15.8 + 0.7
20-40	9.1 + 0.6	15.3 + 0.2	21.1 + 0.9	8.9 + 0.2	15.3 + 1.9	20.3 + 5.2
70-90	7.3 + 0.5	10.2 + 0.2	28.5 + 5.0	9.7 + 2.8	16.7 + 0.3	29.8 + 1.1
<u>Pine forest (PF-o)</u>						
0-20	12.0 + 0.5	34.6 + 3.6	24.9 + 3.5	11.2 + 0.2	37.1 + 0.8	21.6 + 0.6
20-40	10.5 + 0.3	28.5 + 2.6	27.9 + 1.5	10.7 + 1.2	34.9 + 1.1	20.2 + 0.5
60-80	10.6 + 0.3	10.6 + 0.3	26.1 + 1.9	10 + 0.6	28.1 + 2.8	21.9 + 2.1
<u>Agriculture (AG-y)</u>						
0-20	18.7 + 0.7	19.9 + 0.3	30.1 + 3.5	17.0 + 0.8	18.6 + 2.1	31.9 + 4.2
20-40	12.1 + 0.6	25.7 + 1.5	24.0 + 1.9	11.1 + 0.6	19.3 + 0.4	16.5 + 1.5
60-80	11.0 + 0.8	29.4 + 1.0	22.1 + 2.8	10.6 + 0.3	26.5 + 1.0	20.5 + 2.5
<u>Agriculture (AG-i)</u>						
0-20	19.6 + 0.3	44.8 + 1.2	11.8 + 1.1	11.7 + 0.2	44.5 + 1.5	13.7 + 0.3
20-40	19.8 + 0.4	53.0 + 3.6	11.0 + 1.8	15.6 + 0.3	45.7 + 0.5	10.2 + 0.2
50-70	18.0 + 0.9	56.6 + 1.2	9.6 + 0.5	16.2 + 0.4	46.6 + 2.2	7.6 + 0.7
<u>Agriculture (AG-o)</u>						
0-20	12.6 + 0.5	21.7 + 2.5	27.4 + 1.5	9.4 + 0.9	22.0 + 0.4	14.0 + 0.9
20-40	9.5 + 0.5	23.1 + 1.6	34.3 + 5.0	8.1 + 0.2	21.6 + 0.8	15.3 + 1.7
60-80	7.5 + 0.04	16.1 + 1.0	33.5 + 2.9	7.5 + 0.6	14.5 + 0.5	22.7 + 1.2

### 3.32 Correlations between SOC fractions and selected chemical soil properties

The proportion of several of the isolated SOC fractions correlated to some of the geochemical soil properties (Fig. 2). Across all land uses and depths, the SOC proportion of the S+A fraction was positively correlated with Al<sub>o</sub> (i.e., Al<sub>o</sub> (in bulk soil) and as

255

well as in the S+A fraction),  $Al_o+1/2Fe_o$ , pH  $H_2O$ , exchangeable Ca, C:N ratio of S+A fraction, and content of amorphous materials ( $p < 0.01$ ). These soil properties were also positively correlated with the amount of OC in S+A fraction ( $g\ C\ kg^{-1}$  soil) and OC concentration in S+A fraction ( $g\ C\ kg^{-1}$  fraction). When considering only the pine forest and agricultural soils, SOC proportion of the S+A fraction was still positively correlated to those soil properties:  $Al_o$  content ( $r = 0.80$ ;  $P < 0.01$ ). The SOC proportion in s+c fraction was positively correlated with exchangeable Al and negatively correlated with pH  $H_2O$  and C:N ratio in s+c fraction ( $p < 0.01$ ). There existed no significant correlations between the proportion of the other SOC fractions and any of the analyzed soil properties for the full dataset. But when considering only pine forest and agriculture, positive correlations existed between SOC proportion in s+c fraction with the amount of  $Fe_o$  s+c fraction ( $p < 0.01$ ). OC concentrations of the s+c and rSOC fractions ( $g\ C\ kg^{-1}$  fraction) with  $Al_o$  and  $Al_o+1/2Fe_o$ . The proportion of rSOC in s+c fraction (i.e.  $rSOC / s+c \times 100$ ) s+c rSOC correlated positively ( $P < 0.05$ ) to  $Al_o$  content ( $r = 0.67$ ), SSA ( $r = 0.54$ ), the micropore volume ( $r = 0.66$ ) and the  $Fe_o$  content ( $r = 0.52$ ).



270

**Figure 2.** Scatter plots of the SOC proportion of the S+A fraction and selected geochemical soil properties, with Pearson correlation coefficients and corresponding statistical significance level indicated.— Sites under agriculture and forest are indicated by  $\Delta$  and  $\bullet$  respectively.

### 3.4.3 Carbon mineralization experiment


#### 3.4.3.1 Gross soil carbon mineralization

275

Across all sites, C-mineralization (C-min) rates were highest for AG-i2 and AG-o3 and peaked already during the first day of incubation, while for the other soils rates peaked at day 3 or 5 only. After day 38, the C-min rates stabilized across time. The cumulative total CO<sub>2</sub>-C emission and derived amount of C-mineralized (SOC + ryegrass) generally followed a sequence NF-y < AG-y1 < AG-i2 < PF-i1 < AG-o3 < PF-o2. Pine forest soils displayed higher cumulative C mineralization than agricultural soils ( $p < 0.05$ ). Overall, the 120-day cumulative C mineralization was 194–419 mg kg<sup>-1</sup> higher in soil with grass added than

280

without (Fig. 3).




285

**Figure 3.** Gross cumulative amount of C mineralized in soil with and without 1 g C kg<sup>-1</sup> ryegrass added. Vertical bars indicate standard deviations of three lab replicates ( $n = 3$ ) and are presented for the grass amended soils only. Sites under forest and agriculture are indicated by  $\bullet$  and  $\Delta$ , respectively.

### 3.43.2 Grass C-mineralization

The  $C_{\text{grass-min}}$  rates peaked within 12 days after the start of the incubation, and they dropped thereafter to just half to one third of the initial rates. The  $C_{\text{grass}}$  mineralization rate was conspicuously high on the first day of incubation in soils AG-i2 and AG-o3, whereas for the other soils mineralization peaked on day 5 only. From day 38 onwards, the  $C_{\text{grass-min}}$  rate had become more or less similar in all six sites. By the end of the 120-days incubations, 22-38 % of the added ryegrass had been mineralized. The highest cumulative  $C_{\text{grass-min}}$  was found for PF-o2 ( $389 \text{ mg kg}^{-1}$ ), followed by AG-i2 > AG-o3  $\approx$  PF-i1  $\approx$  AG-y1 > NF-y. There were no significant differences in  $C_{\text{grass-min}}$  between the sets of agricultural and forest soils.




**Figure 4.** The percentage of  $C_{\text{grass}}$  (left) and  $C_{\text{SOC}}$  (right) mineralized during 120 days of incubation in soils with  $1 \text{ g kg}^{-1}$  grass or no grass added, respectively. Vertical bars indicate standard deviations of triplicate lab repetitions. Sites under forest and agriculture are indicated by  $\bullet$  and  $\Delta$ , respectively.

### 3.43.3 Native SOC C-mineralization

Mineralization of SOC ( $C_{\text{SOC-min}}$ ) contributed more to gross soil C mineralization than  $C_{\text{grass-min}}$  did. There were large differences in  $C_{\text{SOC-min}}$  between the 0-20 cm soil layers of the six sites. By the end of the incubation, the highest cumulative amount of SOC mineralized was found for PF-o2 ( $953 \pm 13 \text{ mg kg}^{-1}$ ) while a fourfold lower amount was mineralized in case of NF-y ( $195 \pm 17 \text{ mg kg}^{-1}$ ). Significantly less SOC was mineralized in the agricultural soils ( $517 \pm 90 \text{ mg kg}^{-1}$ ) than in the pine forest soils ( $813 \pm 156 \text{ mg kg}^{-1}$ ) ( $p < 0.01$ ). Relatively, 0.5–2.9 % of the SOC had been mineralized within the 120 days incubations (Fig. 4). A larger % of SOC was mineralized in the pine forest soils ( $2.9 \% \pm 0.1$ ) than in the agricultural soils ( $1.2 \% \pm 0.3$ ) ( $p < 0.01$ ).

Priming of native SOC mineralization was assessed by comparing  $C_{SOC-min}$  between the soils with and without  $1 \text{ g kg}^{-1}$  grass added (Fig. 5). During the first week, grass addition tended to increase the  $C_{SOC-min}$  rate in most soils, i.e. there was positive priming of native SOC. Along the 120-days experiment, however, the direction and magnitude of the net cumulative priming effect varied strongly among soils. Little or no net priming of  $C_{SOC-min}$  occurred in the NF-y soil, as a result of initial positive priming until day 17 but negative priming afterwards. In the pine forest soils likewise periods of positive and negative priming alternated, with at the end of the 120-days period an alike limited net positive priming of SOC at both PF sites. Priming of SOC was diverse among the three agricultural soils. While strong positive priming throughout the 120-days period in the AG-o soil, while the opposite was true for AG-y and no net priming occurred in case of AG-i. On average, the net 120-days cumulative priming of SOC in forest ( $19.3 \text{ mg CO}_2\text{-C kg}^{-1} \text{ day}^{-1}$ ) was lower than agricultural ( $31.2 \text{ mg CO}_2\text{-C kg}^{-1} \text{ day}^{-1}$ ) land use, but due to wide variation between replicate sites this difference proved insignificant and moreover net priming rates were not significantly different from zero.



**Figure 5.** Net effect of grass addition ( $1 \text{ g kg}^{-1}$ ) on cumulative native SOC mineralization. Positive and negative figures denote net positive or net negative cumulative priming of SOC mineralization. Dotted lines present the temporal evolution of mean net cumulative priming effect for either forest or agricultural land use, with respectively only upper or lower error bars representing the standard deviations on the data.

## 4 Discussion

### 4.1 Land use effects on free and aggregate protected OC

Overall, the portion of S+A fraction material (i.e.,  $> 63 \mu\text{m}$  400 J ml<sup>-1</sup> ultrasonication resistant aggregates and sand particles) was higher-nearly double in the agricultural ( $364 \pm 220 \%$ ) than forest sites ( $1948 \pm 104 \%$ ) ( $p < 0.05$ ). This indicated




stronger soil aggregation and aggregate stability under agriculture. In line, ~~even a double-fourfold~~ share of SOC resided in the S+A fraction under agriculture (~~61-32 ± 21-22~~ %) than forest (~~34-7.7 ± 15-4~~ %) ( $p < 0.05$ ). The important role of soil microaggregates for SOC storage has been displayed by numerous studies (Dungait et al., 2012; Rabbi et al., 2016; Six et al., 2002). We expected a lesser share of SOC to be occluded inside aggregates in the tilled agricultural soils, as absence of tillage results in less disruption of macroaggregates with increased formation of microaggregates and occlusion of POM (Six et al., 2000b; Zheng et al., 2018). However, tillage is not the only determinant of soil aggregate formation and breakdown, ~~as~~ aggregate stability also depends on presence of binding agents that cluster mineral and organic particles into aggregates. Repeated addition of labile C via manure and compost amendments was shown to result in greater formation of macroaggregate (Du et al., 2014; Mikha and Rice, 2004; Yu et al., 2012) by increased production of microbial-derived binding agents. Manure and compost amendments to the AG sites were vast (Table 1), and so the enhanced addition of labile C as compared to the forest sites may have boosted aggregate formation with more SOC storage in the S+A fraction under agricultural land use. Next to organic binding agents, Wagai et al. (2018) also found reactive metal phases to be important binding agents in Allophanic Andisols which was in line with the higher amount of  $Al_o$  S+A in agricultural than forest soils ( $p < 0.05$ ). With high to very high levels of  $Al_o$  (up to 4% of the soil mass), it seems very likely that amorphous to poorly crystalline  $Al$ -containing minerals would have been prominent binding agents in our set of soils. The bulk soil and S+A fraction contents of  $Al_o$  were also larger in the agricultural than in the forest soils ( $p < 0.05$ ). Moreover, ~~the~~ the SOC proportion of the S+A fraction and its OC concentration ( $g\ C\ kg^{-1}$  S+A fraction) correlated positively with contents of  $Al_o$  and  $Al_o+1/2Fe_o$ , and with the % of X-ray amorphous materials. This indicates that higher levels of  $Al_o$  under agriculture than forest would have enhanced aggregation, and in doing so occluded a larger share of SOC inside these aggregates. When comparing the agricultural sites and only the pine forest sites likewise differences existed. In other words the results were not biased by the deviating mineralogical composition and texture of the NF site. Other than aluminium oxides, the content of exchangeable Ca also correlated positively to S+A fraction mass proportion ( $r = 0.405$ ) and the SOC proportion of the S+A fraction ( $r = 0.5961$ ); ( $p < 0.01$ ). Soil Ca is also well known for its positive effects on aggregation via formation of clay-polyvalent cation-OM complexes (de Kerchove and Elimelech, 2007; Six et al., 2004). This mechanism might have co-promoted aggregate occlusion of OC and its accumulation (Bertrand et al., 2007; Paul et al., 2003), but given the high  $Al_o$  levels, likely  $Ca^{2+}$  would have played a smaller role. The higher levels of exchangeable Ca under agriculture compared to forest ( $p < 0.05$ ) evidently result from liming which increases the pH, ~~thus having higher capacity to bind  $Ca^{2+}$~~ . The concomitant difference in  $Al_o$ , exchangeable  $Ca^{2+}$  and S+A OC content between agricultural and forest sites makes it impossible to derive the relative importance of these various binding agents in stimulation of aggregate OC occlusion under agricultural land use. Notwithstanding, when considering only pine forest and agricultural sites, the S+A SOC proportion sole best correlated to the S+A  $Al_o$  content ( $r = 0.80$ ;  $p < 0.01$ ), strengthening the view that to a substantial extent enhanced  $Al_o$  content under agriculture promoted storage of SOC inside stable aggregates. By expanding the Zimmerman et al. (2007) fractionation procedure with a stronger ( $400\ J\ ml^{-1}$ ) ultrasonic dispersion step we attempted to further isolate s+c associated OC from POM contained in the S+A fraction. The intermediate C:N ratio (mean:  $12 + 4.7$ ) of the final  $400\ J\ ml^{-1}$  S+A fraction indicates that next to POM it still also contained

360 ~~other, likely mineral associated, OM, complicating further interpretation of mechanisms for S+A SOC storage in the~~  
~~agricultural sites. Nevertheless, the higher S+A fraction C:N ratio of the agricultural sites vs. the forests ( $P < 0.05$ ) demonstrates~~  
~~that at least to some extent enhanced SOC storage in aggregates under agricultural land-use resulted from extra occluded POM~~  
~~storage. Particularly for the 0 – 20 cm layer the contrast in the S+A C:N ratio (agriculture: 17.0 vs. forest: 9.3) supports this~~  
~~view.-~~ Lastly, less free POM was found in the agricultural than forest soils ( $p < 0.015$ ), likely as a logical consequence of  
365 more POM occlusion under agriculture. However, we cannot exclude that also a better degradability of crop-derived residues  
vs. pine or native forest litter co-explains these trends. It is well established that litter rich in cutin, waxes and lignin such as  
derived from pine forests is less degradable than plant litter richer in carbohydrates and organic N like most crop residues  
(Berg and McLaugherty, 2014).

#### 4.2 Effect of land use on SOC in the silt and clay fraction

370 Interaction of SOC with silt and clay provides long-term stabilization to SOC against decomposition and therefore forms a key  
mechanism for SOC sequestration (De Clercq et al., 2015). SOC sorbs to phyllosilicate clays, Al-, Fe- and Mn-oxides, poorly  
crystalline minerals, as well as polyvalent cations that form a bridge between minerals and organic constituents (Blanco-Canqui  
and Lal, 2004). However, for our soils we found that despite higher concentrations of  $Al_0$  under agriculture, a lesser share of  
the SOC was present in the s+c fraction (as % of SOC) compared to the forest soils ( $p < 0.05$ ). ~~The content of s+c OC (in g C~~  
375 ~~kg<sup>-1</sup> soil) displayed likewise trends. Particularly, the all depth layers at the NF site had high s+c content and proportion of~~  
~~SOC, but this was in fact accompanied with a larger contribution of younger less-weathered minerals present, low  $Al_0$  level~~  
~~and smaller share of amorphous minerals (Anindita et al., under review). Hence, this association of OC with  $Al_0$ -containing~~  
minerals could not explain the higher SOC proportion ~~and amount~~ of OC in s+c under forest. ~~As the amount of s+c OC (in g~~  
380 ~~C kg<sup>-1</sup> soil) was in fact similar between both land-uses it appears that a lower s+c SOC contribution in the agricultural soils~~  
~~was instead simply mirroring trends in the S+A fraction. The s+c fraction proportion to SOC correlated negatively to soil pH~~  
~~and positively to exchangeable Al content but likewise these correlations probably do not point to a causal relationship but~~  
~~rather result from the higher SOC proportion of S+A OC under agriculture alongside liming-caused lifted soil pH and lowered~~  
~~exchangeable Al. The weak positive correlations with between the s+c SOC proportion SOC in s+c fraction may contribute~~  
385 ~~to larger SOC proportion in s+c fraction under pine forest and  $Fe_0$  in s+c ( $r = 0.45$ ;  $p < 0.01$ ) could indicate that, but we did~~  
~~not find good correlation between  $Fe_0$  s+c with OC in s+c fraction association with poorly crystalline Fe stabilizes OM in~~  
~~the silt and clay fraction. But, again as s+c SOC content did not differ between the agricultural and the forest sites while  $Fe_0$~~   
~~did, such land-use control on s+c  $Fe_0$  associated SOC seems unlikely. Instead, most likely a larger s+c proportion of SOC~~  
under forest simply mirrors trends in the S+A fraction, although it cannot be ruled out that other factors such as a lower soil  
nutrient availability, lower soil pH, and likely wetter microclimate and more biochemically inert OC inputs would have caused  
390 relative storage of OC in the s+c fraction under forest. Our experimental design did not allow to infer unambiguously if such  
soil and environmental factors alternatively caused the larger SOC proportions in s+c.

The s+c fraction was per the Zimmerman et al. (2007) fractionation procedure subdivided into oxidizable and resistant parts, expected to coincide roughly with SOC stabilized by mineral-association as opposed to its own biochemical inertness, respectively. Significant positive correlations existed between the share of s+c OC that was oxidizable by 6 % NaOCl (i.e., the  $r_{SOC} / s+c \times 100$ ) with the s+c fraction contents of  $Al_0$ , % of amorphous materials and the bulk soil SSA, and micropore volume content of amorphous materials (Fig. 6) (not shown), especially in subsoils where such the correlations with  $Al_0$  were strong (Fig. 6). These correlations suggest that the oxidizable s+c OC-fraction indeed at least partially coincides with mineral-associated SOC, in spite that on the opposite Association of SOC with Fe and Al (hydr)oxides has indeed just as well before been confirmed found to limit oxidation of SOC by 6 % NaOCl (Mikutta et al., 2006). Since we also found that agricultural land use had higher  $Al_0$  by 14% compared to pine forest (Table 1), these results at first sight might suggest that the enhanced mineral weathering under agricultural land use lowered the chemical oxidizability of s+c OC. However, by comparing pine forest and agriculture, the difference in the As neither chemical inertness of the s+c OC (agriculture: 53 % of s+c vs. forest: 59.49 % of s+c), nor the s+c  $Al_0$ , bulk soil SSA and micropore volume appeared insignificant between differed between both land uses it appears that land-use history did not impact the degree of mineral association of OC and its content in these Andosols. (agriculture: 53 % of s+c vs. pine forest: 49 % of s+c). This outcome then further at least contradicts that more s+c OC would accumulate under forest because of a more biochemical inertness of the OC inputs compared to agriculture.



**Figure. 6** Relation between the proportion of rSOC of the s+c fraction in the pine forest and agricultural soils with the soil  $Al_0$  (all dots jointly,  $n = 30$ ) and for the set of subsoil observations (in blue) specifically; and likewise relations with the soil specific surface area and the micropore volume % of amorphous materials, and the soil specific surface area ( $n = 15$ , SSA and micropore volume X-ray diffraction not replicated). Sites under agriculture and pine forest are indicated by  $\Delta$  and  $\bullet$ , respectively. Pearson correlation coefficients in black and blue represent correlations for the total dataset and for the set of subsoils, respectively.

### 4.3 Land use effects on native SOC degradability

420 The percentage of SOC mineralized after 120 days of incubation was less than half in the agricultural soils (1.2 %) than in the pine forest soils (2.9 %) ( $p < 0.01$ ). This result was rather unexpected given the lower soil pH by 0.8 under pine forest, as acidic conditions are well known to retard SOC decomposition of SOC (Högberg et al., 2007; Malik et al., 2018) in line with Anda and Dahlgren (2020). To understand the apparent relative stabilization of SOC in the agricultural soils, we again considered the distribution of SOC across soil fractions. We (Anindita et al., 2022) ~~(Anindita et al., under review)~~ previously hypothesized that the nearly doubled  $\text{Al}_o$  content of the agricultural soils ( $21.8 \text{ g kg}^{-1}$  on average) compared to the pine forest soils ( $13.9 \text{ g kg}^{-1}$ ) would partially explain the relatively elevated SOC stocks under agriculture for our study area due to enhanced organo–mineral association. Basile-Doelsch et al. (2005) likewise found that volcanic ash soil horizons containing much poorly crystalline material (proto–imogolite and proto–imogolite allophane) store large amounts of organic matter which turns over very slowly. But as no negative correlations existed between the amount or proportion of SOC mineralized and  $\text{Al}_o$  content, this hypothesis was not confirmed. As mentioned above, we did find a higher % of SOC to be present in the rSOC fraction under forest but its content ( $\text{g kg}^{-1}$  soil) was not higher than under agricultural land use and again no correlation existed with SOC mineralization. Mikutta et al. (2006) indeed demonstrated that a larger chemical recalcitrance of SOC does not really translate into a larger biological stability. ~~Moreover, the share of SOC in rSOC was actually lower (but not significantly) under forest than under agriculture and so as argued above, Once more, it appears that~~ the larger SOC proportion of rSOC seems to simply result from less accumulation of SOC in the S+A fraction under forest.

The proportion and concentration of the S+A fraction OC were higher under agriculture, so enhanced soil aggregation and occlusion of SOC might have significantly physically protected SOC against microbially mediated decomposition compared to pine forest land use. A study of Andosols in Colombia by Gijsman and Sanz (1998) also reported that aggregation holds considerable control on SOC decomposability as these authors observed a significant increase of  $\text{CO}_2$  mineralization after crushing large and small ~~-~~macroaggregates ( $> 53 \mu\text{m}$ ). Indeed, we found that the amount of OC ( $\text{g C kg}^{-1}$  soil) in the S+A fraction negatively correlated to % SOC mineralized for the set of soils with agricultural and pine forest land use ( $r = -0.785$ ;  $p < 0.01$ ). It then appears that enhanced soil aggregation and occlusion of SOC under agricultural land use in part limited degradability of SOC, rather than enhanced association of SOC with soil minerals, including poorly crystalline or amorphous  $\text{Al}$ -containing minerals. As argued in 4.2, we can add to this that Fformation of amorphous materials and  $\text{Al}_o$  in the agricultural soils most likely promoted soil aggregation. Hence, we ~~provisionally conclude~~hypothesize that enhanced pedogenesis forms a relevant indirect mechanism via which conversion to agricultural land use impacts the SOC balance of these relatively young tropical volcanic soils. ~~Overall~~However, since large amounts of exogenous OM are annually being applied to the in this study investigated agricultural fields (Table 1), we cannot unequivocally exclude that these instead explain the high SOC stocks and

450 contribution of aggregate associated OC here and more observations on an expanded set of sites will be required to test this hypothesis.

#### 4.4 Land use effects of exogenous organic matter decomposition and net priming of SOC

In contrast to native SOC mineralization, added grass C mineralization did not differ between forest and agricultural soils. Therefore, apparently protection of ryegrass OC either by mineral association or occlusion inside aggregates must have been similar between both land uses and, in any case, did not impact its degradability in soil. Laboratory incubations with disturbed soils do not necessarily adequately reproduce field conditions in terms of soil structure, microclimate, and food webs. The observed similar degradability of ryegrass-C does not suggest a likewise enhanced storage of freshly added OM like for native SOC under agriculture than under forest. Possibly, such impact only emerges on the longer term when the added OM has been subject to sufficient diminution into smaller POM that could be occluded in microaggregates. Six et al. (2002) summarized that particularly 50–250µm scaled microaggregates grant sizable physical protection to POM, while the macroaggregate (> 250 µm) structure exerts little control on POM decomposability. Adding ryegrass to the 0–20 cm soils initially seemed to impact mineralization of native SOC with a general positive priming effect in first week of incubation, in line with many other studies, e.g. Liu et al. (2017). Thereafter, however, in four of the six studied soils net negative priming countered this initial stimulation of SOC mineralization. Such adverse impact of adding fresh OM on SOC mineralization could be due to a preferential utilization of the added substrate by microorganisms (Derrien et al., 2014). Regardless, the net result was that after 120 days SOC mineralization was not significantly stimulated (on average by 1.6 %). There was also no land use effect on the net priming of SOC after 120 days. Hence, in spite of differences in SOC quality under both land uses, as inferred from different SOC proportions of S+A and free POM, no differential stimulation of SOC mineralization by addition of a relatively labile plant-C substrate existed between agricultural or forest land use. These observations imply that no adverse effect is to be expected of the addition of generally relatively labile C (above ground crop residues, animal manure) on stability of the native SOC present under agricultural land use.

#### 5 Conclusions

475 Following the fractionation of SOC by Zimmermann et al., (2007), the silt- and clay associated OC forms the dominant SOC pool in tropical volcanic soils in Indonesia, regardless of land-use. TheBut land-use strongly impacts SOC contained within 400 J m<sup>-1</sup> ultrasonic dispersion resistant soil sand-aggregates-fraction, as its contribution to SOC rose from 9% of SOC in forest to as much as 32% of SOC in several decades after conversion to agricultural use than pine forest soils, with alongside also more OC stored in this fraction. This fraction in general was a dominant SOC pool in our soils, accounting for 47% of total SOC in all soil fractions, and S+A SOC proportion was positively assoeiated-related to Al<sub>o</sub>, Al<sub>o</sub>+1/2Fe<sub>o</sub>, exchangeable Ca,

480 pH, and the % of amorphous materials. Our study thus indicates ~~that agricultural land use to stimulated higher level~~the more  
~~amount development~~ of ~~amorphous materials, particularly~~ poorly crystalline Al<sub>o</sub>, ~~more amorphous materials and under~~  
~~agricultural land use, which in turn stimulate~~promoted soil aggregation and physical occlusion of OC compared to forest land  
use. ~~The~~A negative correlation between relative decomposability of the SOC and the portion of SOC that is physically  
occluded ~~on the one hand and to lower relative decomposability of the SOC on the other, further~~ suggests that enhanced soil  
485 aggregation under cropland ~~indeed~~ effectively stabilizes part of the SOC compared to under forest land use. Contrary to our  
hypothesis, however, we found no proof that stimulated formation of Al<sub>o</sub> and amorphous minerals would have increased  
association of SOC with soil minerals under agriculture. Based on the present study we postulate that the enhanced formation  
of amorphous minerals and Al<sub>o</sub> under agriculture with high OM inputs promoted development of stable soil aggregates  
and OC occlusion therein and this would in part counter otherwise expected losses of SOC compared to primary and secondary  
490 forest. However, the contribution of large OM inputs vs. land-use conversion per sé could not be elucidated here and this will  
require study of other tropical Andosol forest-agricultural land use pairs with detailed inventory of OC inputs.~~Based on the~~  
~~present study we do postulate that the stable aggregates via formation of amorphous materials Al<sub>o</sub> faster pedogenesis under~~  
~~agriculture and high C input via organic fertilizer would counter otherwise expected loss of SOC compared to secondary forest,~~  
~~but further study of more tropical Andosol land use pairs with detailed inventory of OC inputs is now needed to affirm this.~~  
495 To the least, our study points at the overall need to account for potential indirect land use effects on stability of SOC via its  
control on pedogenesis. Especially, this is so for pedoclimatic combinations where weathering can be very fast, like in tropical  
Andosols that have only relatively recently been cultivated.

## Appendix

No Appendix

## 500 Data availability

The data generated in this study are available from the corresponding author upon reasonable request

## Author contributions

SS, PF, and SA conceptualized the study. SA conducted soil sampling and performed the experiment. SA prepared the manuscript with contributions from all the co-authors.

## 505 Competing interest

The authors declare that they have no conflict of interest

## Acknowledgments

We thank Indonesia Endowment Fund for Education (LPDP) for the internal support of this research, Orly Mendoza, Haichao Li, and Heleen Dero from Soil Fertility Research Group (Ghent University) for laboratory assistance.

## 510 Financial support

This research has been supported by Indonesia Endowment Fund for Education (LPDP).

## References

- Anda, M. and Dahlgren, R. A.: Long-term response of tropical Andisol properties to conversion from rainforest to agriculture, *Catena*, 194, 104679, doi:10.1016/j.catena.2020.104679, 2020.
- 515 [Anindita, S., Sleutel, S., Vandenberghe, D., Grave, J. De, Vandenhende, V. and Finke, P.: Land use impacts on weathering, soil properties, and carbon storage in wet Andosols, Indonesia, \*Geoderma\*, 423\(May\), 115963, doi:10.1016/j.geoderma.2022.115963, 2022.](#)
- ~~Anindita, S., Sleutel, S., Vandenberghe, D., De Grave, J., Vandenhende, V. and Finke, P.: Land use impacts on weathering and carbon sequestration in Sunda Tangkuban Perahu volcanic region, Indonesia, n.d.~~
- 520 Asano, M. and Wagai, R.: Evidence of aggregate hierarchy at micro- to submicron scales in an allophanic andisol, *Geoderma*, 216, 62–74, doi:10.1016/j.geoderma.2013.10.005, 2014.
- Asano, M. and Wagai, R.: Distinctive organic matter pools among particle-size fractions detected by solid-state  $^{13}\text{C}$ -NMR,  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  analyses only after strong dispersion in an allophanic andisol, *Soil Sci. Plant Nutr.*, 61(2), 242–248, doi:10.1080/00380768.2014.982492, 2015.
- 525 Asano, M., Wagai, R., Yamaguchi, N., Takeichi, Y., Maeda, M., Suga, H. and Takahashi, Y.: In Search of a Binding Agent: Nano-Scale Evidence of Preferential Carbon Associations with Poorly-Crystalline Mineral Phases in Physically-Stable, Clay-Sized Aggregates, *Soil Syst.*, 2(2), 32, doi:10.3390/soilsystems2020032, 2018.
- Baldock, J. A.: Composition and Cycling of Organic Carbon in Soil, in *Nutrient Cycling in Terrestrial Ecosystems*, edited by P. Marschner and Z. Rengel, pp. 1–35, Springer Berlin, Heidelberg., 2007.
- 530 Bartoli, F., Burtin, G. and Herbillon, A. J.: Disaggregation and clay dispersion of Oxisols: Na resin, a recommended methodology, *Geoderma*, 49(3–4), 301–317, doi:10.1016/0016-7061(91)90082-5, 1991.
- Basile-Doelsch, I., Amundson, R., Stone, W. E. E., Borschneck, D., Bottero, J. Y., Moustier, S., Masin, F. and Colin, F.: Mineral control of carbon pools in a volcanic soil horizon, *Geoderma*, 137(3–4), 477–489, doi:10.1016/j.geoderma.2006.10.006, 2007.
- 535 Berg, B. and McLaugherty, C.: *Plant litter: decomposition, humus formation, carbon sequestration*, Third., Springer Berlin Heidelberg, New York., 2014.



- Bertrand, I., Delfosse, O. and Mary, B.: Carbon and nitrogen mineralization in acidic, limed and calcareous agricultural soils: Apparent and actual effects, *Soil Biol. Biochem.*, 39(1), 276–288, doi:10.1016/j.soilbio.2006.07.016, 2007.
- Blagodatskaya, E., Khomyakov, N., Myachina, O., Bogomolova, I., Blagodatsky, S. and Kuzyakov, Y.: Microbial interactions affect sources of priming induced by cellulose, *Soil Biol. Biochem.*, 74, 39–49, doi:10.1016/j.soilbio.2014.02.017, 2014.
- Blanco-Canqui, H. and Lal, R.: Mechanisms of carbon sequestration in soil aggregates, *CRC. Crit. Rev. Plant Sci.*, 23(6), 481–504, doi:10.1080/07352680490886842, 2004.
- Van Breemen, N., Mulder, J. and Driscoll, C.: Acidification and alkalization of soils, *Plant Soil*, 75(3), 283–308 [online] Available from: <http://www.jstor.com/stable/42934465>, 1983.
- Bruijnzeel, L. A.: Nutrient Content of Litterfall in Coniferous Forest Plantations in Central Java, Indonesia, *J. Trop. Ecol.*, 1(4), 353–372 [online] Available from: <http://www.jstor.org/stable/2559453>, 1985.
- Cerli, C., Celi, L., Kalbitz, K., Guggenberger, G. and Kaiser, K.: Separation of light and heavy organic matter fractions in soil - Testing for proper density cut-off and dispersion level, *Geoderma*, 170, 403–416, doi:10.1016/j.geoderma.2011.10.009, 2012.
- Chen, R., Senbayram, M., Blagodatsky, S., Myachina, O., Dittert, K., Lin, X., Blagodatskaya, E. and Kuzyakov, Y.: Soil C and N availability determine the priming effect: Microbial N mining and stoichiometric decomposition theories, *Glob. Chang. Biol.*, 20(7), 2356–2367, doi:10.1111/gcb.12475, 2014.
- Churchman, G. J. and Lowe, D. J.: Alteration, formation, and occurrence of minerals in soils, in *Handbook of Soil Sciences*, vol. 1, edited by P. M. Huang, Y. Li, and M. E. Sumner, pp. 20.1-20.72, CRC Press (Taylor & Francis), Boca Raton, FL., 2012.
- De Clercq, T., Heiling, M., Dercon, G., Resch, C., Aigner, M., Mayer, L., Mao, Y., Elsen, A., Steier, P., Leifeld, J. and Merckx, R.: Predicting soil organic matter stability in agricultural fields through carbon and nitrogen stable isotopes, *Soil Biol. Biochem.*, 88, 29–38, doi:10.1016/j.soilbio.2015.05.011, 2015.
- Covaleda, S., Gallardo, J. F., García-Oliva, F., Kirchmann, H., Prat, C., Bravo, M. and Etchevers, J. D.: Land-use effects on the distribution of soil organic carbon within particle-size fractions of volcanic soils in the Transmexican Volcanic Belt (Mexico), *Soil Use Manag.*, 27(2), 186–194, doi:10.1111/j.1475-2743.2011.00341.x, 2011.
- Cronan, C. S.: *Ecosystem Biogeochemistry*, Springer. [online] Available from: <http://link.springer.com/10.1007/978-3-319-66444-6>, 2018.
- Cusack, D. F., Chadwick, O. A., Ladefoged, T. and Vitousek, P. M.: Long-term effects of agriculture on soil carbon pools and carbon chemistry along a Hawaiian environmental gradient, *Biogeochemistry*, 112(1–3), 229–243, doi:10.1007/s10533-012-9718-z, 2013.
- Derrien, D., Plain, C., Courty, P. E., Gelhaye, L., Moerdijk-Poortvliet, T. C. W., Thomas, F., Versini, A., Zeller, B., Koutika, L. S., Boschker, H. T. S. and Epron, D.: Does the addition of labile substrate destabilise old soil organic matter?, *Soil Biol. Biochem.*, 76(November 2017), 149–160, doi:10.1016/j.soilbio.2014.04.030, 2014.
- Doetterl, S., Stevens, A., Six, J., Merckx, R., Van Oost, K., Casanova Pinto, M., Casanova-Katny, A., Muñoz, C., Boudin, M., Zagal Venegas, E. and Boeckx, P.: Soil carbon storage controlled by interactions between geochemistry and climate, *Nat.*

- Geosci., 8(10), 780–783, doi:10.1038/ngeo2516, 2015.
- Don, A., Schumacher, J. and Freibauer, A.: Impact of tropical land-use change on soil organic carbon stocks - a meta-analysis, *Glob. Chang. Biol.*, 17(4), 1658–1670, doi:10.1111/j.1365-2486.2010.02336.x, 2011.
- 575 Dörner, J., Dec, D., Feest, E., Vásquez, N. and Díaz, M.: Dynamics of soil structure and pore functions of a volcanic ash soil under tillage, *Soil Tillage Res.*, 125, 52–60, doi:10.1016/j.still.2012.05.019, 2012.
- Du, Z. liu, Wu, W. liang, Zhang, Q. zhong, Guo, Y. bin and Meng, F. qiao: Long-Term Manure Amendments Enhance Soil Aggregation and Carbon Saturation of Stable Pools in North China Plain, *J. Integr. Agric.*, 13(10), 2276–2285, doi:10.1016/S2095-3119(14)60823-6, 2014.
- 580 Dungait, J. A. J., Hopkins, D. W., Gregory, A. S. and Whitmore, A. P.: Soil organic matter turnover is governed by accessibility not recalcitrance, *Glob. Chang. Biol.*, 18(6), 1781–1796, doi:10.1111/j.1365-2486.2012.02665.x, 2012.
- Gerzabek, M. H., Bajraktarevic, A., Keiblinger, K., Mentler, A., Rechberger, M., Tintner, J., Wriessnig, K., Gartner, M., Valenzuela, X. S., Troya, A., Couenberg, P. M., Jäger, H., Carrión, J. E. and Zehetner, F.: Agriculture changes soil properties on the Galápagos Islands-two case studies, *Soil Res.*, 57(3), 201–214, doi:10.1071/SR18331, 2019.
- 585 Gijsman, A. J. and Sanz, J. I.: Soil organic matter pools in a volcanic-ash soil under fallow or cultivation with applied chicken manure, *Eur. J. Soil Sci.*, 49(3), 427–436, doi:10.1046/j.1365-2389.1998.4930427.x, 1998.
- Guillaume, T., Kotowska, M. M., Hertel, D., Knohl, A., Krashevskaya, V., Murtillaksono, K., Scheu, S. and Kuzyakov, Y.: Carbon costs and benefits of Indonesian rainforest conversion to plantations, *Nat. Commun.*, 9(1), doi:10.1038/s41467-018-04755-y, 2018.
- 590 Hernández, Z., Almendros, G., Carral, P., Álvarez, A., Knicker, H. and Pérez-Trujillo, J. P.: Influence of non-crystalline minerals in the total amount, resilience and molecular composition of the organic matter in volcanic ash soils (Tenerife Island, Spain), *Eur. J. Soil Sci.*, 63(5), 603–615, doi:10.1111/j.1365-2389.2012.01497.x, 2012.
- Hertel, D., Moser, G., Culmsee, H., Erasmí, S., Horna, V., Schuldt, B. and Leuschner, C.: Below- and above-ground biomass and net primary production in a paleotropical natural forest (Sulawesi, Indonesia) as compared to neotropical forests, *For. Ecol. Manage.*, 258(9), 1904–1912, doi:10.1016/j.foreco.2009.07.019, 2009.
- 595 Högberg, M. N., Högberg, P. and Myrold, D. D.: Is microbial community composition in boreal forest soils determined by pH, C-to-N ratio, the trees, or all three?, *Oecologia*, 150(4), 590–601, doi:10.1007/s00442-006-0562-5, 2007.
- Huygens, D., Boeckx, P., Van Cleemput, O., Godoy, R. and Oyarzun, C.: Aggregate structure and stability linked to carbon dynamics in a south Chilean Andisol, *Biogeosciences Discuss.*, 2(1), 203–238, doi:10.5194/bgd-2-203-2005, 2005.
- 600 Keeling, D.: The concentration and isotopic abundances of atmospheric carbon dioxide in rural areas, *Geochim. Cosmochim. Acta*, 13, 322–334, doi:https://doi.org/10.1016/0016-7037(58)90033-4, 1958.
- de Kerchove, A. J. and Elimelech, M.: Formation of polysaccharide gel layers in the presence of Ca<sup>2+</sup> and K<sup>+</sup> ions: Measurements and mechanisms, *Biomacromolecules*, 8(1), 113–121, doi:10.1021/bm060670i, 2007.
- Kleber, M., Eusterhues, K., Keilweit, M., Mikutta, C., Mikutta, R. and Nico, P. S.: Mineral-Organic Associations: Formation, Properties, and Relevance in Soil Environments, in *Advances in Agronomy*, vol. 130, edited by D. . Sparks, pp. 1–140, Elsevier

- 605 Ltd., 2015.
- Lehmann, J. and Kleber, M.: The contentious nature of soil organic matter, *Nature*, 528(7580), 60–68, doi:10.1038/nature16069, 2015.
- Li, J., Du, J., Zhong, S., Ci, E. and Wei, C.: Changes in the profile properties and chemical weathering characteristics of cultivated soils affected by anthropic activities, *Sci. Rep.*, 11(1), 20822, doi:10.1038/s41598-021-00302-w, 2021.
- 610 Linlin, G., Taku, N., Hiromi, I. and Zhigang, S.: Carbon Mineralization Associated with Soil Aggregates as Affected by Short-Term Tillage, *J. Resour. Ecol.*, 7(2), 101–106, doi:10.5814/j.issn.1674-764x.2016.02.004, 2016.
- Liu, X. J. A., Sun, J., Mau, R. L., Finley, B. K., Compson, Z. G., van Gestel, N., Brown, J. R., Schwartz, E., Dijkstra, P. and Hungate, B. A.: Labile carbon input determines the direction and magnitude of the priming effect, *Appl. Soil Ecol.*, 109, 7–13, doi:10.1016/j.apsoil.2016.10.002, 2017.
- 615 Malik, A. A., Puissant, J., Buckeridge, K. M., Goodall, T., Jehmlich, N., Chowdhury, S., Gweon, H. S., Peyton, J. M., Mason, K. E., van Agtmaal, M., Bland, A., Clark, I. M., Whitaker, J., Pywell, R. F., Ostle, N., Gleixner, G. and Griffiths, R. I.: Land use driven change in soil pH affects microbial carbon cycling processes, *Nat. Commun.*, 9(1), 1–10, doi:10.1038/s41467-018-05980-1, 2018.
- Matus, F., Rumpel, C., Neculman, R., Panichini, M. and Mora, M. L.: Soil carbon storage and stabilisation in andic soils: A  
620 review, *Catena*, 120, 102–110, doi:10.1016/j.catena.2014.04.008, 2014.
- McKeague, J. A.: An evaluation of 0.1 M pyrophosphate and pyrophosphate-dithionite in comparison with oxalate as extractants of the accumulation products in podzols and some other soils, *Can. J. Soil Sci.*, 47(2), 95–99, doi:10.4141/cjss67-017, 1967.
- Mikha, M. M. and Rice, C. W.: Tillage and Manure Effects on Soil and Aggregate-Associated Carbon and Nitrogen, *Soil Sci. Soc. Am. J.*, 68(3), 809–816, doi:10.2136/sssaj2004.8090, 2004.
- 625 Mikutta, R., Kleber, M., Torn, M. S. and Jahn, R.: Stabilization of soil organic matter: Association with minerals or chemical recalcitrance?, *Biogeochemistry*, 77(1), 25–56, doi:10.1007/s10533-005-0712-6, 2006.
- Mikutta, R., Schaumann, G. E., Gildemeister, D., Bonneville, S., Kramer, M. G., Chorover, J., Chadwick, O. A. and Guggenberger, G.: Biogeochemistry of mineral-organic associations across a long-term mineralogical soil gradient (0.3-4100  
630 kyr), Hawaiian Islands, *Geochim. Cosmochim. Acta*, 73(7), 2034–2060, doi:10.1016/j.gca.2008.12.028, 2009.
- Miyazawa, M., Takahashi, T., Sato, T., Kanno, H. and Nanzyo, M.: Factors controlling accumulation and decomposition of organic carbon in humus horizons of Andosols: A case study for distinctive non-allophanic Andosols in northeastern Japan, *Biol. Fertil. Soils*, 49(7), 929–938, doi:10.1007/s00374-013-0792-8, 2013.
- Parfitt, R. L.: Allophane and imogolite: role in soil biogeochemical processes, *Clay Miner.*, 44(1), 135–155,  
635 doi:10.1180/claymin.2009.044.1.135, 2009.
- Paul, E. A., Morris, S. J., Six, J., Paustian, K. and Gregorich, E. G.: Interpretation of Soil Carbon and Nitrogen Dynamics in Agricultural and Afforested Soils, *Soil Sci. Soc. Am. J.*, 67(5), 1620–1628, doi:10.2136/sssaj2003.1620, 2003.
- Poepflau, C., Don, A., Dondini, M., Leifeld, J., Nemo, R., Schumacher, J., Senapati, N. and Wiesmeier, M.: Reproducibility of

- a soil organic carbon fractionation method to derive RothC carbon pools, *Eur. J. Soil Sci.*, 64(6), 735–746, doi:10.1111/ejss.12088, 2013.
- 640 Poirier, V., Basile-Doelsch, I., Balesdent, J., Borschneck, D., Whalen, J. K. and Angers, D. A.: Organo-Mineral Interactions Are More Important for Organic Matter Retention in Subsoil Than Topsoil, *Soil Syst.*, 4(1), 4, doi:10.3390/soilsystems4010004, 2020.
- Qiao, N., Schaefer, D., Blagodatskaya, E., Zou, X., Xu, X. and Kuzyakov, Y.: Labile carbon retention compensates for CO<sub>2</sub> released by priming in forest soils, *Glob. Chang. Biol.*, 20(6), 1943–1954, doi:10.1111/gcb.12458, 2014.
- 645 Rabbi, S. M. F., Daniel, H., Lockwood, P. V., Macdonald, C., Pereg, L., Tighe, M., Wilson, B. R. and Young, I. M.: Physical soil architectural traits are functionally linked to carbon decomposition and bacterial diversity, *Sci. Rep.*, 6(September), doi:10.1038/srep33012, 2016.
- Six, J., Elliott, E. T. and Paustian, K.: Soil macroaggregate turnover and microaggregate formation: A mechanism for C sequestration under no-tillage agriculture, *Soil Biol. Biochem.*, 32(14), 2099–2103, doi:10.1016/S0038-0717(00)00179-6, 2000a.
- 650 Six, J., Paustian, K., Elliott, E. T. and Combrink, C.: Soil Structure and Organic Matter I. Distribution of Aggregate-Size Classes and Aggregate-Associated Carbon, *Soil Sci. Soc. Am. J.*, 64(2), 681–689, doi:10.2136/sssaj2000.642681x, 2000b.
- Six, J., Conant, R. T., Paul, E. A. and Paustian, K.: Stabilization mechanisms of soil organic matter: Implications for C-saturation of soils, *Plant Soil*, 241(2), 155–176, doi:10.1023/A:1016125726789, 2002.
- 655 Six, J., Bossuyt, H., Degryze, S. and Denef, K.: A history of research on the link between (micro)aggregates, soil biota, and soil organic matter dynamics, *Soil Tillage Res.*, 79(1), 7–31, doi:10.1016/j.still.2004.03.008, 2004.
- Smith, W. ., Cleveland, C. ., Reed, S. . and Running, S. .: Agricultural conversion without external water and nutrient inputs reduces terrestrial vegetation productivity, *Geophys. Prospect.*, (April), 6298–6305, doi:10.1002/2013GL058857. **Received**, 660 2014.
- [SSB & UGhent: Study LA BOD/STUD 2006 01 04 for the Flemish Government - Development of an expertsystem for C-management advise for cropland soils: part 1 literature study & part 2 development of an advice system \(in Dutch\). Dept. Environment, Nature and Energy of the Flemish Government. Brussels, https://publicaties.vlaanderen.be/view-file/20207, 2008.](https://publicaties.vlaanderen.be/view-file/20207,2008)
- 665 Taylor, M. D., Lowe, D. J., Hardi, P., Smidt, G. A. and Schnug, E.: Comparing volcanic glass shards in unfertilised and fertilised Andisols derived from rhyolitic tephra, New Zealand: Evidence for accelerated weathering and implications for land management, *Geoderma*, 271(November 2018), 91–98, doi:10.1016/j.geoderma.2016.01.035, 2016.
- Wagai, R., Kajiura, M., Uchida, M. and Asano, M.: Distinctive roles of two aggregate binding agents in allophanic andisols: Young carbon and poorly-crystalline metal phases with old carbon, *Soil Syst.*, 2(2), 1–23, doi:10.3390/soilsystems2020029, 670 2018.
- Wei, X., Shao, M., Gale, W. and Li, L.: Global pattern of soil carbon losses due to the conversion of forests to agricultural land, *Sci. Rep.*, 4, 6–11, doi:10.1038/srep04062, 2014.

- Werth, M. and Kuzyakov, Y.:  $^{13}\text{C}$  fractionation at the root-microorganisms-soil interface: A review and outlook for partitioning studies, *Soil Biol. Biochem.*, 42(9), 1372–1384, doi:10.1016/j.soilbio.2010.04.009, 2010.
- 675 Yu, H., Ding, W., Luo, J., Geng, R., Ghani, A. and Cai, Z.: Effects of long-term compost and fertilizer application on stability of aggregate-associated organic carbon in an intensively cultivated sandy loam soil, *Biol. Fertil. Soils*, 48(3), 325–336, doi:10.1007/s00374-011-0629-2, 2012.
- Zheng, H., Liu, W., Zheng, J., Luo, Y., Li, R., Wang, H. and Qi, H.: Effect of long-term tillage on soil aggregates and aggregate-associated carbon in black soil of northeast China, *PLoS One*, 13(6), 1–18, doi:10.1371/journal.pone.0199523, 2018.
- 680 Zimmermann, M., Leifeld, J., Schmidt, M. W. I., Smith, P. and Fuhrer, J.: Measured soil organic matter fractions can be related to pools in the RothC model, *Eur. J. Soil Sci.*, 58(3), 658–667, doi:10.1111/j.1365-2389.2006.00855.x, 2007.