

Impact of agricultural management on soil aggregates and associated organic carbon fractions: Analysis of long-term experiments in Europe

Ioanna S. Panagea¹, Antonios Apostolakis², Antonio Berti³, Jenny Bussell⁴, Pavel Čermak⁵, Jan Diels¹, Annemie Elsen⁶, Helena Kusa⁵, Ilaria Piccoli³, Jean Poesen^{1,7}, Chris Stoate⁴, Mia Tits⁶, Zoltan Toth⁸, and Guido Wyseure¹

¹Department of Earth and Environmental Sciences, KU Leuven, Leuven, 3001, Belgium

²Max-Planck-Institute for Biogeochemistry, Jena, 07745, Germany

³DAFNAE Department, University of Padova, Legnaro PD, 35020, Italy

⁴Game & Wildlife Conservation Trust, Allerton Project, Loddington, Leicester, LE7 9XE, UK

⁵Crop Research Institute, Prague 6-Ruzyně, 161 06, Czech Republic

⁶Soil Service of Belgium (BDB), Heverlee, 3001, Belgium

⁷Institute of Earth and Environmental Sciences, UMCS, Lublin, 20-718, Poland

⁸Institute of Agronomy, Georgikon Campus, Keszthely, Hungarian University of Agriculture and Life Sciences, Gödöllő, 2100, Hungary

Correspondence: Ioanna S. Panagea (ioanna.panagea@kuleuven.be)

Abstract. Inversion tillage is a commonly applied soil cultivation practice in Europe, which often has been blamed for deteriorating topsoil stability and organic carbon (OC) content. In this study, the potential to reverse these negative effects in the topsoil by alternative agricultural management practices are evaluated in seven long-term experiments (running from 8 to 54 years the moment of sampling) in five European countries (Belgium, Czech Republic, Hungary, Italy and UK). Topsoil samples (0–15 cm) were collected and analysed to evaluate the effects of conservation tillage (reduced and no-tillage) and increased organic inputs of different origin (farmyard manure, compost, crop residues) combined with inversion tillage on topsoil stability, soil aggregates and within these, OC distribution using wet sieving after slaking. Effects from the treatments on the two main components of organic matter i.e., particulate (POM) and mineral associated (MAOM), were also evaluated using dispersion and size fractionation. Reduced and no-tillage practices, as well as the additions of manure or compost, increased the aggregates mean weight diameter (MWD) (up to 49% in the Belgian study-site) and topsoil OC (up to 51% in the Belgian study-site), as well as the OC corresponding to the different aggregate size fractions. The incorporation of crop residues had a positive impact on the MWD but a less profound effect on OC content both on total OC and on OC associated with the different aggregates. A negative relationship between the mass and the OC content of the microaggregates (53–250 μm) was identified in all experiments. There was no effect on the mass of the macroaggregates and the occluded microaggregates (mM) within these, while the corresponding OC contents increased with less tillage and more organic inputs. Inversion tillage led to less POM within the mM, whereas the different organic inputs did not affect it. In all experiments where the total POM increased, the total soil organic carbon (SOC) was also affected positively. We concluded that the negative effects of inversion

tillage on topsoil can be mitigated by reducing the tillage intensity or by adding organic materials, optimally combined with non-inversion tillage methods.

20 1 Introduction

Soil tillage by mouldboard ploughing which inverts and mixes the soil is a classical conventional agricultural practice in temperate agricultural systems, including most of Europe. This soil cultivation technique is used to prepare the soil for seeding, to reduce soil compaction, to control weeds and crop residues and to improve water and nutrients availability for plants (Cannell, 1985; Peigné et al., 2018; Schneider et al., 2017; Hobson et al., 2021). Inverting the topsoil especially by mouldboard ploughing is blamed for deteriorating the stability and arrangement of the soil aggregates (soil structure) with subsequent negative effects on soil organic matter (SOM) preservation and sequestration potential (Six et al., 2000a; Six and Paustian, 2014). For this reason, introducing agricultural practices that counter the adverse effects of mouldboard tillage on soil structure and SOM will contribute to sustainable agricultural management and climate change mitigation (Cooper et al., 2021).

The protection of SOM in the soil aggregates has been proposed as an important stabilization mechanism. According to Oades (1984), macroaggregates are initially formed around fresh plant material (i.e. particulate organic matter (POM)) entering the soil due to physical, chemical, and biological processes. As the organic matter within macroaggregates ($> 250 \mu\text{m}$) decomposes, an internal structure develops consisting of microaggregates ($53\text{--}250 \mu\text{m}$) within which the SOM is better protected (i.e., occluded SOM (mM-C)) as compared to non-occluded SOM (Ananyeva et al., 2013; De Clercq et al., 2015; Denef et al., 2001; King et al., 2019). When macroaggregates break down, for instance due to wetting-drying cycles, the microaggregates are released (Six et al., 2000a; Denef et al., 2001). However, agricultural practices such as inversion tillage may break down soil macroaggregates prematurely, which results in the release of unstable microaggregates and leads to increased SOM accessibility for decomposition (Kotronakis et al., 2017; Six et al., 2000a). For this reason, the fraction of microaggregates held within the macroaggregates (mM) has been proposed as an indicator to monitor present and potential soil structural and SOM changes caused by management practices (Denef et al., 2004, 2007; Kong et al., 2005; Six et al., 2000a; Six and Paustian, 2014).

Tillage effects on soil structure and especially topsoil SOM are a point of controversy in the literature. Several studies observed an increase in macroaggregates under non-inversion / zero till as compared to conventional inversion tillage agriculture (Jat et al., 2019; Paustian et al., 1997; Six et al., 2000b). Reduced soil disturbance is often accompanied by an increase in SOC in the topsoil (Apostolakis et al., 2017; Fuentes et al., 2012; Giannakis et al., 2014; Karlen et al., 2013; Virto et al., 2012) or even the whole soil profile (Cooper et al., 2021; Varvel and Wilhelm, 2011) especially when it is combined with increased inputs from cover crops or intercrops (Boddey et al., 2010; Fuentes et al., 2009; Minasny et al., 2017). This increase is mainly explained by differences among the practices in terms of organic input accumulation in the topsoil (Andruschkewitsch et al., 2014; Virto et al., 2012) and reduced mineralization rates. Other studies, reported limited or no effects of reduced or no tillage on SOM or even negative effects in some cases, especially when the entire soil profile was considered (Blanco-Canqui and Lal, 2008; Camarotto et al., 2020; Du et al., 2017; Haddaway et al., 2017; Piccoli et al., 2016; Meurer et al., 2018). This was

explained mainly by the concentration of organic materials in the top/surface layer due to no redistribution through inversion tillage in the deeper soil layers. Reduced and zero tillage have also been claimed to cause other agricultural risks like increased soil compaction, impaired root development, increased weed and disease pressure leading to more pesticide use and, possibly, lower crop yields and poor crop establishment (Berner et al., 2008; Ogle et al., 2012; Peigné et al., 2018; Piccoli et al., 2021; 55 Van den Putte et al., 2010). Overall, while tillage reduction or cessation seems to alleviate the negative effects on soil structure and topsoil SOM, it also removes some of its beneficial effects especially in the short term. Kay and VandenBygaart (2002) and Sartori et al. (2021) speculated that in the short-term, soil compaction would be expected as a result of tillage absence and traffic load, while crop yield and soil structure stabilization are expected in the mid- and long-term (>15years) (Rusinamhodzi et al., 2011), as a result of greater biological activity and SOM redistribution.

60 While soil tillage increases the turnover rate of OM, a supply of organic materials should compensate for the losses. Under conventional agriculture, the addition of exogenous organic materials, like compost or manure, could offset the negative effects of inversion-tillage on soil structure (Williams et al., 2017) while maintaining the beneficial ones. Gross and Glaser (2021) concluded that regardless of tillage intensity manure application can increase the SOC stocks and that animal manure led to a greater SOC increase compared to green manure or plant-derived organic amendments (e.g., straw). While compost and 65 animal manure additions are found to improve soil aggregation and increase both the total and the aggregate associated OC in croplands receiving conventional tillage, the inorganic fertilization did not result in such improved soil fertility (De Clercq et al., 2016; Kotronakis et al., 2017; Lin et al., 2019; Yin et al., 2016). Minasny et al. (2017) reviewed several studies and found that organic amendments (such as compost or manure) lead to a SOC accumulation of on average $0.5 \text{ Mg C ha}^{-1} \text{ year}^{-1}$ while residue incorporation leads to accumulation of on average $0.35 \text{ Mg C ha}^{-1} \text{ year}^{-1}$. Incorporation of crop residues increased total 70 SOC from 2.7 up to 18.2 % (Bolinder et al., 2020) when compared to residue removal as well as increased the SOC within the different aggregate fractions (Zhao et al., 2018) and improved aggregation (Zhang et al., 2014). The effects of different residue management practices are highly connected with the tillage methods that are combined with it, thus leading to controversial findings when the latter is overlooked. Paul et al. (2013) concluded that tillage or residue management alone does not affect SOC, but when the residues are incorporated into the soil in the 15 – 30 cm layer, the SOC content increases whereas the 75 aggregate stability is not improved. On the other hand, Li et al. (2020) identified non-inversion tillage (breaking the upper soil layers without any mixing or inverting) coupled with residue retention (when compared to conventional inversion or zero tillage with or without residue retention) as the optimum system to increase SOC stocks in the 0 – 30 cm soil layer.

Management effects on SOC and soil aggregation vary in the literature possibly due to differences in the climatic conditions and/or soil properties between studied regions (Pan et al., 2021), as well as, variations in the application and incorporation 80 rates of the organic materials. Methodological differences between studies during sampling or analysis processes (Poeplau et al., 2018) may also be responsible for the controversial responses reported. For this reason, studies that investigate the impact of multiple and combined agricultural practices on several regions under a common methodological framework are needed. Such studies are in general scarce due to the logistical constraints associated with field experiments, and available studies in different countries are difficult to compare due to differences in objectives and analyses methods of heterogeneous

85 field experiments. Meurer et al. (2018) identified the lack of documented required data for meta-analysis of such studies as another limitation factor.

SOM is a complex mixture that can be separated into multiple pools with distinct functional properties (Schrumpf et al., 2013; Trumbore, 2009). The separation of SOM into particulate organic matter (POM) that consists of plant-derived, relatively undecomposed light fragments, and mineral-associated organic matter (MAOM) that consists of mostly simple molecules, 90 has been proposed for many decades (Cambardella and Elliott, 1992; Poepflau et al., 2018). Lavalley et al. (2020) suggested the applicability of SOM separation into POM and MAOM for the investigation of management effects on carbon cycling. Coupling the separation of SOM in OM occluded in different aggregation fractions and in POM and MAOM within each aggregation fraction can help elucidating the effects of tillage intensity and soil amendments on SOM stability.

In this study, we investigated the effects of three agricultural practice categories on soil structure and SOC distribution in four 95 aggregate size fractions and one subfraction (i.e. occluded microaggregates (mM)). The same field and laboratory protocols used for the analysis of seven long-term experiments in five European countries. The agricultural practice categories were: i) soil cultivation including zero, minimum and conventional tillage, ii) crop residue incorporation or removal, iii) addition of exogenous organic or inorganic material including manure, compost and NPK fertilizers.

Our hypotheses were that in the topsoil:

- 100 – Inversion tillage reduces the mass of water-stable (large) macroaggregates and of SOC stabilized in macroaggregates and microaggregates, and this effect increases with increasing tillage intensity (from zero to conventional mouldboard ploughing).
- Matured exogenous organic fertilization, (i.e., manure and compost application), increases the mass of water-stable macroaggregates and increases SOC in all aggregate fractions, even under inversion tillage.
- 105 – Incorporation of the previous crop residues increases the mass of water-stable macroaggregates and occluded POM due to higher additions of fresh plant-derived organic matter, and thus leads to higher SOC stabilized in both occluded and free microaggregate, compared to the practice of crop residue removal.
- Increased organic inputs and lower soil disturbances lead to more water-stable mM and corresponding SOC (mM-C), (i.e., the mass of mM and occluded OC exhibit a linear relationship).

110 Our objectives were therefore: i) to evaluate the effects of the different treatments/practices on the topsoil structural stability and SOC, ii) to determine the soil mass and OC distribution in four water-stable aggregate size fractions (i.e., large macroaggregates: $LM > 2000 \mu\text{m}$; macroaggregates: $2000 < M < 250 \mu\text{m}$; microaggregates: $250 < \text{micro} < 53 \mu\text{m}$; clay and silt domains: $s\&c < 53 \mu\text{m}$), iii) to determine the effects of the practices on the occluded microaggregates (mM) and the corresponding OC (mM-C) and iv) to determine the OC in POM and MAOM in the bulk soil and the aggregation fractions.

2.1 Study site description

For our objectives, we took the option to combine several long-term experiments and to have a wide range of soil management practices and combinations rather than just concentrating on only one practice. Logistically, to handle seven experiments from five countries by the same methods, sampling within under identical conditions and laboratory analysis in the same laboratory we had to concentrate on the 0 – 15 cm topsoil. Thus, topsoil samples (upper 15 cm) were collected from different long-term agricultural experiments (8 to 54 years) with different treatments (the towns, countries, coordinates, main soil type and climate of the sites are given in Table A1 taken from Panagea et al. (2021)). The long-term experiments were set up independently from one another with different objectives and under different environmental conditions. Nevertheless, they offer the possibility to explore a wide range of representative management practices and pedo-climatological conditions across Europe as they cover a wide gradient both west-east and north-south (Fig.1). For each country, the field experiment included management practices (and intensities) that were adapted to the local conditions and were commonly applied by local farmers, but also intensities that were considered extreme, to evaluate the maximum effects (e.g., 45 Mg ha⁻¹ compost annually in BE).

For this research, a subset of the experimental treatments was selected to include various treatment categories to test our hypotheses (Table 1). This resulted in 79 experimental plots under 26 treatments. The first category includes primarily treatments with different soil cultivation intensities (CZ, HU_2, UK), the second category focuses on the addition of different types of exogenous organic materials (BE, IT_1c, IT_1p, HU_1), in cropping systems that are conventionally tilled, and the third category deals with the incorporation of crop residues (HU_1, IT2_c, IT_2l), also in systems under conventional inversion tillage. The experiments in Italy are conducted on two different soil types each and, in this study, were analysed as separate experiments: a clay and an initially peaty soil for IT_1 (i.e., IT_1c and IT_1p) and a clay and a loamy soil for experiment IT_2 (i.e., IT_2c and IT_2l). The selected treatments per experiment are presented in Table 1. At the five study sites, an identical sampling and analysis procedure was performed for determining the soil aggregates size fraction distribution, the OC content in each size fraction, and the separation of SOM into POM and MAOM (Figure 2) . For consistency reasons, all laboratory analyses were conducted in the same laboratory with the same equipment, by the same person. This is particularly important for the analysis of the aggregates.

2.2 Aggregate separation

Field-moist topsoil samples were taken with a sharp shovel in a Z-shape sampling design within each experimental plot from the upper 15 cm of soil, mixed and directly broken to pass a <8 mm sieve. Soil samples were stored in plastic containers to avoid compaction during transportation and then stored in the refrigerator until air-drying. All samples were air-dried and stored in a dark and dry place at room temperature.

Aggregate separation was done by wet sieving, as prescribed from Elliott (1986) and described in Six et al. (2000b). A 100 g subsample of 8 mm-sieved and air-dried soil was submerged in demineralized water for 5 min on a 2000 μ m sieve. Aggregates were separated by moving the sieve up and down mechanically 50 times in two minutes. The > 2000 μ m fraction




Figure 1. Map with the location of the long-term experiments

was backwashed in a glass jar and the $<2000 \mu\text{m}$ fraction was transferred in the next sieve ($250 \mu\text{m}$) and sieved with the same methodology. This procedure was repeated with a $53 \mu\text{m}$ sieve. All aggregate fractions were oven-dried (50°C) until dried, 150 weighted and stored for further fractionation and analysis. The aggregates were distinguished into large macroaggregates (>2 mm) (LM), macroaggregates ($250 \mu\text{m}$ – 2 mm) (M), microaggregates ($53 \mu\text{m}$ – $250 \mu\text{m}$) (micro) and silt and clay size particles ($<53 \mu\text{m}$) (s&c). The procedure was replicated three times for each sample from each experimental plot and the averaged values used for the following analysis.

Table 1: Details of the treatments in the various experiments.

Code	Treatments	Tillage system	Sampling month, year	Years applied	Repl.(#), design	Main crop type
CZ	<p>Conventional: Conventional ploughing (Turning of stubble—furrow opener at 10 cm, Mouldboard plough at 22 cm) (control)</p> <p>Minimum: Minimum tillage (Turning of stubble—furrow opener at 10 cm, 30 % of crop residues remain on the soil surface)</p> <p>Zero: Zero tillage (all residues remain in the soil surface)</p>	Depending on the experimental treatment	Nov 2018	23	4, †	Oil rapeseed, winter wheat, peas

Table 1 – continued from previous page

Code	Treatments	Tillage system	Sampling month, year	Years applied	Repl.(#), design	Main crop type
HU_2	<p>Conventional: Deep winter ploughing (27–28 cm) + secondary tillage (control)</p> <p>Minimum: Disking just before drilling (<15 cm)</p> <p>Shallow: Shallow winter disking (<15 cm) +secondary tillage</p>	Depending on the experimental treatment	Nov 2018	46	4, †	Winter wheat, maize
UK	<p>Conventional: Ploughing at 25 cm (control)</p> <p>Direct drilling: Direct drilling of the seeds into previous crop residues</p>	Depending on the experimental treatment	Apr 2019	8	3, ‡	Winter wheat, wheat, oat
BE	<p>No organic: No organic fertilization (control)</p> <p>45tn3-yearly: 45 Mg ha⁻¹ compost* applied every three years</p> <p>15tnyearly: 15 Mg ha⁻¹ compost* applied yearly</p> <p>45tnyearly: 45 Mg ha⁻¹ compost* applied yearly</p> <p>*C/N ≈ 12</p>	Conventional tillage up to 23 cm and bed preparation according to the crop type	Oct 2019	22	4, ‡	Winter wheat, carrots, sugar beet, potatoes
HU_1	<p>NPK: Only mineral fertilization/ removal of straw(control)</p> <p>NPK+FYM: 35Mg ha⁻¹ 0.5% N, farmyard manure application every 3 years/removal of straw</p> <p>NPK+STR: Straw and stalk incorporation completed with 10 kg N*Mg straw ha⁻¹</p>	Conventional tillage: shallow stubble tillage, ploughing (27 cm) in Autumn, secondary tillage and seedbed preparation	Nov 2018	35	3, †	Maize, winter wheat, winter barley
IT_1c	<p>Unfertilized: No organic or mineral fertilization (control)</p> <p>Manure L1: 20 Mg ha⁻¹ manure applied annually*</p> <p>Manure L2: 40 Mg ha⁻¹ manure applied annually*</p>	Shovelling-inversion tillage (0-20 cm) each autumn after the removal of crop residues.	Nov 2018	54	2, ‡	Maize, winter wheat, potato, tillage radish (winter cover crop), ryegrass, silage maize
IT_1p	<p>Unfertilized: No organic or mineral fertilization (control)</p> <p>Manure L1: 20 Mg ha⁻¹ manure applied annually*</p> <p>Manure L2: 40 Mg ha⁻¹ manure applied annually*</p> <p>*Farmyard manure from dairy cows (20% dry matter, 0.5 % N, 0.25 % P₂O₅, 0.7 % K₂O)</p>					
IT_2c	<p>Residue Remov.: Removal of the previous crop residues (control)</p> <p>Residue incorp.: Burial of the previous crop residues</p>	Shovelling-inversion tillage (0-20 cm) each autumn	Nov 2018	48	3, ‡	Maize, winter wheat, potato, ryegrass, silage maize, tillage radish

Table 1 – continued from previous page

Code	Treatments	Tillage system	Sampling month, year	Years applied	Repl.(#), design	Main crop type
IT_2I	Residue Remov.: Removal of the previous crop residues (control)					
	Residue incorp.: Burial of the previous crop residues					

‡ Randomized complete block design (RCBD), † Split plot randomized complete block design (Split Plot-RCBD).

155

2.3 Determination of occluded microaggregates

The isolation of the microaggregates held within the macroaggregates was done by a microaggregate isolator using the method proposed by Six et al. (2000a). A 10 g subsample of macroaggregates was placed in the microaggregate isolator on top of a 250 μm mesh sieve, immersed in demineralized water and shaken with 50 glass beads (4 mm diameter). Steady and continuous water flow helped the microaggregates to flush on a 53 μm sieve. When only water was flushed in the sieve, the material on the 53 μm sieve was sieved to ensure that the isolated microaggregates were water-stable, backwashed in a glass jar, oven-dried (50 °C) weighed and stored. The material on the 250 μm sieve was also backwashed in a glass jar, oven-dried (50 °C) weighed and stored for further analysis.

160

2.4 Size fractionation of SOM to MAOM and POM and sand correction

A size fractionation approach was used to distinguish SOM to its components, i.e., MAOM and POM as described in Cambardella and Elliott (1992), and to determine the sand content of all aggregate fractions $>53 \mu\text{m}$ using the method of Elliott et al. (1991). Specifically, the average upper size limit specification from MAOM at 53 μm was used which is also considered as the lower limit for the sand fraction. An aliquot of 5 g from each aggregation fraction was dispersed in 20 ml sodium hexametaphosphate (5 g L⁻¹), (0.5 %) for 18 hours at 180 rpm, and sieved through a 2 mm sieve to remove possible small stones and a 53 μm sieve to determine sand content. The fraction remaining in the latter sieve was backwashed in a glass jar, oven-dried (105 °C) until constant mass and weighed. The procedure was replicated twice for each sample and the average value was used for the following analysis. By dispersing and sieving the fractions through the 53 μm sieve, the fraction remaining in the sieve was considered sand and the POM component of the SOM and the fraction that passed through the sieve was taken as the MAOM component and s&c fraction.

170

The relevant aggregate weight percentages were corrected for their sand content to allow comparisons between soils with different sand contents. The sand corrected fraction proportions were calculated as proposed in Six et al. (2002).

175

The soil stability was assessed using the Mean Weight Diameter (MWD) (Eq. 1) which is the most widely used index, defined as the sum of the weighted mean diameter of the aggregate size fractions where the weighting factor of each fraction is the proportion of the total sample weight for each fraction (Nimmo, 2013).




Figure 2. Aggregate and SOM separation scheme used in this study. For each fraction the different soil particles and relevant organic matter components as considered in this research are presented. This separation scheme designed following a combination of methods and concepts described in Cambardella and Elliott (1992); Elliott et al. (1991); Six et al. (2000a)

$$180 \quad MWD(\mu m) = \frac{8000\mu m + 2000\mu m}{2} LM\% + \frac{2000\mu m + 250\mu m}{2} M\% + \frac{250\mu m + 53\mu m}{2} micro\% + \frac{53\mu m}{2} s\&c\% \quad (1)$$

2.5 Soil Organic Carbon determination

Soil organic carbon content (SOC) which is the largest and easiest component of SOM to quantify, was used in this study. The SOC content was determined by dry combustion and mass spectrometry elemental analysis (Carlo-Erba EA 1110, Thermo Scientific). A representative subsample of each fraction was taken with a soil sample splitter, ground and weighed into an Ag capsule. To determine only the carbon present in organic form, carbonates were removed with the addition of HCl (35 %). After drying at 40 °C for 24 hours, the soil samples were loaded into the autosampler for combustion with oxygen with the presence of chromium trioxide (catalyst). The mass percentage was determined after quantification by infrared absorption spectroscopy of the organic carbon (OC) that reacted to carbon dioxide (CO₂).

2.6 Statistical analysis

One-way Analysis of Variance (ANOVA) (Webster, 2007) was carried out using R-Studio with R version 3.6.1 (R Core Team, 2019; RStudio Team, 2016) to test for differences between treatments. Estimated marginal means by factors were computed by the least square method using the package “emmeans” (Lenth, 2020). All graphs were produced with the package “ggplot2” (Wickham, 2016). The statistical significance was checked at $p < 0.05$. The assumptions of normality and homoscedasticity of the residuals were assessed by visual inspection of the Q-Q plots and plots of the normalized residuals against the fitted values.

3 Results

3.1 Soil stability

The soil structure stability of the sand corrected aggregates represented by the MWD (Figure 3) was a sensitive index reflecting management changes. In CZ, the zero-tillage treatment had almost double MWD in comparison to the minimum or conventional tillage. The same applied to the HU_2 experiment between the shallow and conventional tillage. The UK experiment presented the highest MWD values of all the experiments monitored up to 3000 μm for the direct drilling (zero tillage) treatment, which was significantly higher than the conventional tillage.

The addition of animal manure or compost led to higher MWD but differences among the treatments were only statistically significant in the BE experiment. Adding 45 Mg ha⁻¹ of VFG compost annually for 22 years, led to a MWD of about 1200 μm which was significantly higher compared to the addition of 45 Mg ha⁻¹ tri-annually (one third of the previous treatment) and from no addition of compost. In the IT_1c, IT_1p and HU_1 experiments, even after 54, and 35 years respectively, there were no significant differences among the treatments but the visual trends were consistent i.e., adding manure increased the MWD. In the HU_1 experiment the incorporation of the crop residues led to significantly higher MWD compared to the NPK-only treatment. Similarly, in the IT_2c and IT_2l experiments, the incorporation of residues resulted in higher values of MWD, but the differences were statistically significant only in the IT_2l.




Figure 3. Mean weight diameter (MWD) of sand free aggregates in the topsoil (0–15 cm) for each study site (see also Table 1 for the codes and a description of the treatments). Error bars represent the standard error. Within an experiment, bars with a different letter differ significantly according to Tukey’s test ($p < 0.05$).

210 3.2 Soil Organic Carbon

The total SOC (Fig. A1) expressed in its two main fractions (i.e., POM and MAOM) (Figure 4), ranged between 10 g C kg⁻¹ soil and 70 g C kg⁻¹ soil (1 % –7 %) among all experiments. In all cases when significant increases in the total POM were identified as a result of management, the total SOC was also increased. The treatments with the minimum soil disturbance in CZ and HU_2 and the treatments where manure or compost were applied in BE, IT_1c, presented significantly higher values of total POM and MAOM in the topsoil, than the treatments where conventional inversion tillage took place or when they were unamended. The retention of the crop residues did not lead to an increase of the topsoil MAOM when compared to their removal while the incorporation of straw significantly increased the POM in HU_1.




Figure 4. Total topsoil (0–15 cm) SOC separated in its components for each study site (see Table 1 for the codes and a description of the treatments). MAOM stands for mineral associated organic carbon and POM for particulate organic carbon. Error bars represent the standard error. Within an experiment and fraction, bars with a different letter differ significantly according to Tukey’s test ($p < 0.05$).

3.3 Aggregate weight and organic carbon distribution

Reduced tillage intensities caused statistically significant differences in the mass distribution of water-stable large macroaggregates and microaggregates, with those causing less soil disturbance resulting in more water-stable large macroaggregates but fewer microaggregates (Figure 5). The macroaggregates between 250 μm and 2 mm did not present significant differences in the CZ and HU_2 experiments, but they differed significantly in the UK experiment in which the conventional tillage resulted in more water-stable macroaggregates than the direct drilling treatment. Different rates of manure or compost applications caused significant differences (in case of BE and IT_1p) or an increasing trend when the organic input level was higher in the large macroaggregates. In the BE experiment, an abundance of water-stable microaggregates was also observed in the treatment without addition of compost compared to those with addition. In the HU_1 experiment, the application of FYM resulted in more macroaggregates and fewer microaggregates than when only mineral fertilizer was applied. In addition, when the straw

was incorporated in the soil, we found significantly more water-stable macroaggregates and less microaggregates than when FYM and mineral fertilizer or only mineral fertilizer was applied. The residue retention in IT_2c and IT_2l did not result in significant differences (except the microaggregates fraction in IT_2l) in the aggregate size fractions.




Figure 5. Weight distribution of sand free aggregate fractions for each study site (Codes and description of the practices are presented in Table 1). The error bars represent the standard error. Within an experiment and fraction, bars with a different letter differ significantly according to Tukey's test ($p < 0.05$).

The SOC content in each aggregate fraction after the sand correction is illustrated in Figure 6. In the CZ experiment, zero tillage resulted in higher OC contents than conventional tillage in every aggregate fraction except the large macroaggregates, although this fraction exhibited the highest OC content up to 35 g C kg^{-1} soil. Interestingly, the minimum tillage resulted in the same OC content with the zero tillage in the macroaggregates fraction. In the microaggregates fraction, minimum tillage presented significantly lower values than the zero tillage and significantly higher OC content than the conventional tillage. In general, the HU_2 experiment followed the same trend as the CZ. Shallow disking resulted in statistically higher OC than conventional tillage in all fractions apart from the large macroaggregates. In the UK experiment, significant differences between

the two treatments were only present in the microaggregates and s&c fractions, with the direct drilling presenting higher OC content (more than 40 g C kg⁻¹ soil in the microaggregates fraction).




Figure 6. SOC content in each aggregate fraction expressed in a sand-free soil for each study (Codes and description of the practices are presented in Table 1). The error bars represent the standard error. Within an experiment and fraction, bars with a different letter differ significantly according to Tukey's test ($p < 0.05$).

240 In the compost or manure experiments, i.e. BE, IT_1c and IT_1p, the OC content in the aggregate fractions (Figure 6) responded as the total OC (Figure 4). The higher application rates led to higher OC contents in the different fractions. In the BE experiment, the annual application of 45 Mg ha⁻¹ compost led to significantly higher OC in the macroaggregates and microaggregates fractions compared to every other treatment. In the IT_1c experiment, addition of manure led to significantly higher OC among all the treatments in the macroaggregates fraction. In the microaggregates and large macroaggregates fractions, there was large variability between the experimental plots and no significant differences were found. Similarly, in the IT_1p experiment, where even though the OC content within the different fractions went up to about 80 g C kg⁻¹ soil, the in-between experimental plots variability did not allow for safe comparisons among the treatments. In the experiments where the treat-

245

ments dealt with the residues management, i.e., HU_1, IT_2c and IT_2l, we did not observe significant differences among the treatments even after 48 years of application.

250 3.4 Mass and OC dynamics in the occluded microaggregates

Based on our observations, we found that tillage intensity and OM additions i) influenced the quality of macroaggregates in terms of OC content, but not their mass, and ii) led to a negative relationship between OC content and aggregate mass in the microaggregates fraction. We therefore isolated the microaggregates within macroaggregates (mM) to understand better the underlying processes. The analysis did not show significant differences among the treatments in all categories on the amount
255 of mM (data shown in Fig. A2), but it did show significant differences in their OC content (mM-C) (Figure 7). Over all study sites, their OC content ranged from 10 to 47 g C kg⁻¹ mM. In the CZ experiment significantly more mM-C was observed in the zero tillage in comparison with the conventional, while in the HU_1 there was significantly more occluded OC in the shallow disking+secondary tillage compared to the other two treatments (i.e., shallow disking alone and conventional deep ploughing). In the UK experiment, OC content in the mM fraction did not differ among the treatments. In the BE experiment,
260 only the annual addition of 45 Mg ha⁻¹ of compost led to increased OC content in the mM fraction compared to the other application rates. Both in the IT_1c and IT_1p experiments, the occluded OC was higher when manure was added than when it was not, but the difference was statistically significant only for IT_1c. The residue retention led to more mM-C only in the HU_1 experiment.

When comparing the differences in each OC component among the treatments (Figure 8), the same trends with the total
265 OC were followed/observed. In the BE experiment when 45 Mg ha⁻¹ of compost were added the MAOM was double compared to the treatments with no addition of compost and statistically significantly higher than the other two treatments with lower amounts of compost addition. In the IT_1c experiment when animal manure was added (both levels) the MAOM was significantly higher than in the unfertilized treatment. The same results were observed in the HU_1 experiment. Comparing the treatment that included only application of mineral fertilizers with those that got either manure or residues, we observed that
270 the MAOM was significantly higher in the case of farmyard manure addition and retention of the residues.

4 Discussion

Improving the topsoil structure and increasing the SOM through improved soil management while ensuring agricultural production and farmers' income, constitutes an important challenge in the agricultural sector (Dignac et al., 2017). Restoring degraded agricultural soils via natural attenuation (i.e. set-aside) demands long periods of fallow, especially in regions with
275 low primary production (Apostolakis et al., 2017), which is not compatible with ensuring farmers' income or societal objectives for food production. This is especially important for regions with stagnating yields or with yields that are considerably less than their potential (Wiesmeier et al., 2015; Schils et al., 2018) due to land mismanagement and poor soil quality, and for regions with low primary production. For this reason, additional measures are needed.




Figure 7. SOC content in the microaggregates held within the macroaggregate fraction (mM-C) (Codes and description of the practices are presented in Table 1). The OC content is expressed in g C per kg of occluded microaggregates. The error bars represent the standard error. Within an experiment, bars with a different letter differ significantly according to Tukey's test ($p < 0.05$).

In this study, we explored the potential to reverse the negative effects of inversion tillage in the topsoil structure and SOM
 280 by applying different management practices that are potentially soil improving. To test our hypotheses, we exploited a network
 of ongoing long-term experiments running from 8 to 54 years the moment of sampling. We used common sampling and
 samples analyses protocols as, these factors can contribute to the strong controversies in the effects of practices demonstrated
 in literature (Blanco-Canqui and Lal, 2008; Haddaway et al., 2017; Karlen et al., 2013; Sheehy et al., 2015; Meurer et al.,
 285 2018). The practices evaluated, included the addition of soil amendments such as compost or manure and the management of
 crop residues combined with inversion tillage, and the reduction of tillage intensities (from minimal to zero). These practices
 control both the spatial and temporal distribution of OM inputs into the soil and regulate the sensitivity of OM to mineralization,
 affecting the OM stocks (Jastrow et al., 2007). Such practices can i) reduce OM mineralization by minimizing disturbances, ii)
 promote aggregation and iii) increase the SOC content and aggregate stability, but on the other hand they might also enhance




Figure 8. OC content in each SOM component in the mM size fraction in each study site (Codes and description of the practices are presented in Table 1). The OC content is expressed in g C per kg of occluded microaggregates. The error bars represent the standard error. Within an experiment and fraction, bars with a different letter differ significantly according to Tukey's test ($p < 0.05$).

the decomposition of old OC due to the addition of fresh OM inputs (e.g. residues), i.e., the priming effect (Fontaine et al., 2007; Liu et al., 2020).

Generally, it was observed that inversion tillage in CZ, HU_2 and UK and less organic inputs in BE, IT_1, HU_1 and IT_2, deteriorated the topsoil aggregate stability and decreased the topsoil SOC. Moreover, inversion tillage and less organic inputs led to fewer large macroaggregates with no profound differences in the OC content within these, and mainly no differences in the macroaggregates quantity but a significantly lower OC content in these macroaggregates. Finally, these practices resulted in more microaggregates, but carbon depleted, and a small increase in the s&c fraction weight but with reduced OC content.

The results obtained after soil separation into aggregates and the determination of their OC content confirm our first hypothesis that inversion tillage reduces the mass of water-stable large macroaggregates (Figure 5) and the SOC stabilized in macroaggregates and microaggregates (Figure 6) including the mM (Figure 7). However, as also observed in some other studies (Modak et al., 2020; Piazza et al., 2020), we did not observe a decrease in the mass of macroaggregates under inversion tillage as initially hypothesised. The tillage systems that reduced soil disturbances to a minimum or zero level and avoided excessive soil inversion, (e.g., CZ, UK and HU_2), improved the topsoil stability, increased the mass of large macroaggregates, did not affect the mass of the macroaggregates and reduced the mass of all other fractions. They finally increased the OC in all fractions in the 0–15 cm soil layer. The mass decrease of the large macroaggregates and the reduction of the SOC in the aggregates fractions in conventionally tilled soils has been presented in several studies (Devine et al., 2014; Jat et al., 2019; Mikha and Rice, 2004; Sheehy et al., 2015) but most of them also found a reduction of the macroaggregates mass (Jat et al., 2019; Mikha and Rice, 2004; Mondal and Chakraborty, 2022; Plaza-Bonilla et al., 2010; Song et al., 2019; Zheng et al., 2018). This can be linked to the seasonal soil aggregate stability variation depending on the soil management, climatic conditions, root stage development, microbial activity and freezing-thawing processes (Dimoyiannis, 2009; Yang and Wander, 1998; Batista et al., 2022; Jirků et al., 2010), which favoured the formation and preservation of large macroaggregates at expense of macroaggregates.

Our second hypothesis dealt with the addition of exogenous material which already underwent decomposition before the application on the soil, in fields in BE, IT_1c, IT_1p and HU_1 where conventional inversion tillage was the main soil cultivation practice. The initial hypothesis was that matured exogenous organic material will increase the mass of water-stable macroaggregates and SOC in all aggregates fractions even if will combined with inversion tillage. Indeed, under most conditions, adding matured exogenous organic material such as compost or manure increased substantially the OC content within all aggregate fractions as well as the total OC in the topsoil, even if combined with inversion tillage (De Clercq et al., 2016; Lin et al., 2019; Mikha and Rice, 2004; Zhang et al., 2021). Interestingly, we did not observe an increased mass of water-stable macroaggregates as initially hypothesized and demonstrated in previous studies (Mikha and Rice, 2004; Wen et al., 2021; Wortmann and Shapiro, 2008). Results similar to ours, were obtained by Lin et al. (2019) and Zhang et al. (2021) who even observed a reduction of the macroaggregates and an increase of the large macroaggregates after manure fertilization. Nevertheless, addition of exogenous organic material increased the mass of the large macroaggregates as also observed by Mikha and Rice (2004) which, despite being a small fraction, is important for maintaining a good soil structure. The macroaggregates presence controls the distribution of macro-pores that influence the water flow particularly near the soil surface. It is also associated with better pore connectivity, optimum movement and storage of gases, water, heat and nutrients, enhanced biological activity and root penetration.

The addition of exogenous organic sources had a pronounced impact on soil structure and SOM regardless of the tillage intensity as also observed by Gross and Glaser (2021). On the other hand, the effects of crop residue left on the soil or incorporated into the soil varied among the different study sites. Based on our findings in all experiments sampled in our study that dealt with residue management, our third hypothesis that incorporation of the crop residues will lead to an increase in the mass of macroaggregates and the occluded POM cannot be supported. Even if the stability of the soil improved, with several

experiments to present statistically significant increase in the MWD as also observed by Modak et al. (2020) and mass of macroaggregates, the corresponding SOC contents and the total SOC were not affected positively by the incorporation of fresh organic materials as had been reported by Powlson et al. (2011). Similar results were obtained by Lin et al. (2019) while other studies demonstrated a parallel increase of aggregates mass (large macroaggregates, macroaggregates and microaggregates) and respective OC contents (Fuentes et al., 2012; Karlen et al., 1994; Zhao et al., 2018). This may be because the exogenous organic materials (compost in BE or manure in IT_1) were at least partially decomposed and so in a more stable form (more resistant soil organic matter) (Berti et al., 2016) and of a higher quality (low C/N ratio) (Castellano et al., 2015). In contrast, the crop residues were not previously decomposed, and they decomposed fast when applied to the soil (Struijk et al., 2020).

In addition, there are concerns and evidence (Dignac et al., 2017; Fontaine et al., 2007; Wang et al., 2015) that the addition and incorporation of fresh organic material combined with conventional deep ploughing may trigger the microbial demand for carbon. This may lead to increased mineralization and consumption of the old existing OC in the soil (i.e., priming effect). Nevertheless, the priming effect, input quality and decomposition rate seem to play a role only in the short term (days to year) as in the long-term (decades) there is evidence from Cardinael et al. (2015) and Thomsen et al. (2013) that do not show differences in the SOC retention, regardless of the initial organic material quality. The combination of residue retention with zero (Pu et al., 2019), minimum tillage, or shallow ploughing could be more effective for SOC storage increase even in the short term (<6years) (Li et al., 2020) compared to conventional tillage practices (Dal Ferro et al., 2020; Fuentes et al., 2012; Luo et al., 2010; Xu et al., 2019).

Reduced or no-tillage keeps fresh organic material concentrated on the soil surface or in the topsoil (Andruschkewitsch et al., 2014; Virto et al., 2012) and keeps the roots intact increasing the POM concentration. Commonly, the roots contribute more to SOC storage than the above-ground inputs, due to an approximately double efficiency in conversion into stable SOC compared to above-ground biomass (e.g., crop residue) (Berti et al., 2016; Kätterer et al., 2011; Rumpel and Kögel-Knabner, 2011). Reduced topsoil disturbances promote aggregation and subsequently provide adequate time for the added fresh POM to become protected within large aggregates and further to become stable in the occluded microaggregates (Six et al., 2000a). These processes lead to reduced mineralization not only in the zero tillage systems but also under reduced tillage in several cases (Chen et al., 2019; Ghimire et al., 2017).

An important aspect is to unravel the soil processes responsible for long-term OC storage and, more importantly in the agricultural sector, the SOC components that are sensitive to management changes. This allows identification of the changes of management effects, which could be either positive or negative towards a desired soil composition and structure. The physically protected OC within the microaggregates fraction has been identified as the second most protected after the OC chemically bonded to the soil mineral particles. A linear negative relationship between the microaggregates fraction weight percentage and their carbon content was detected in this study (F-stat: 15.61, p-value: 0.000171), as also found in agricultural or native soils (Six et al., 2000b) under different levels of disturbances and organic material addition. This points out that the increasing cultivation intensity and the lack of organic materials lead to a loss of carbon rich macroaggregates and to an increase of microaggregates which are though, C-depleted.

365 In most of the investigated experiments, the mass of both the macroaggregates (except for UK and HU_1) and occluded
microaggregates were not affected by the different management practices or their intensities. On the other hand, the total
SOC (g C kg^{-1} soil), macroaggregate C and mM-C (g C kg^{-1} mM) were in most cases positively affected when the tillage
was reduced, or when more organic materials were added. Therefore, we can partially reject our fourth hypothesis stating
that increased organic inputs and reduced soil disturbances will lead to more mM and corresponding SOC (mM-C). Increased
370 SOC inputs and lower soil disturbances led indeed to more mM-C but not necessarily to more water-stable mM. The parallel
change of the occluded microaggregates with the corresponding OC content (i.e., more mM-C when the mM are increasing)
proposed by Six et al. (2000a) and supported by several other studies (King et al., 2019; Modak et al., 2020; Piazza et al.,
2020; Zhao et al., 2018). However, our findings do not support this hypothesis. Similarly, Andruschkewitsch et al. (2014) and
Deneff et al. (2007) also concluded that there is no concomitant change in specific cases. Thus, the relationship between mass
375 of occluded microaggregates and corresponding OC as well as the applicability of this concept in different soils and under
various management practices should be further evaluated.

In the experiments with significant differences in the mM-C (g C kg^{-1} mM) (Figure 7), the differences were reflected by the
different occluded carbon components, i.e., mM-MAOM (g MAOC kg^{-1} mM) and mM-POM (g POC kg^{-1} mM) (Figure 8), as
well as, the total SOC content (g C kg^{-1} soil) (Figure 4). In the treatments in which the soil was less disturbed compared to those
380 that were under conventional tillage, both the MAOM and POM were higher, as also found by Modak et al. (2020). Taking
into account that the quality of the macroaggregates and the occluded microaggregates depends on the POM (Six et al., 1998),
which is trapped initially within the macroaggregates, together with the above-mentioned results on their mass we suggest that
the total soil OC content is independent of the macroaggregates quantity but rather on their quality in terms of POM which
is trapped within them. In the experiments which included addition of organic materials like manure or compost, an increase
385 was found only in the MAOM component. In contrast, in the experiments that incorporated the fresh residues in the soil no
changes in the MAOM nor in the POM were identified. POM is mainly plant derived and has been proven in other studies to
increase when the residues are retained or incorporated in the soil (Modak et al., 2020) or if manure is applied (Wen et al.,
2021). In our research these practises were combined with inversion conventional tillage, and thus we suggest that the time that
the macroaggregates remain intact is crucial for the labile POM to stabilize within the occluded microaggregates.

390 An important difference found between exogenous (matured) material and fresh crop residue in our results is that the first
can possibly counteract the inversion tillage effects on topsoil OC while the latter cannot. In the case of matured exogenous
organic material, as in BE there is an increasing trend of SOC with increasing compost doses under conventional tillage, which
is visible throughout all treatments but is only statistically significant at the highest dose. This does not mean that there is no
effect at lower doses. However, because of the great field variability, statistical differences are often difficult to demonstrate.
395 That is the reason that the extremely high dose for Belgium of 45 Mg ha^{-1} compost to the plots every year has been included
even if from a practical point of view, it is not feasible to be implemented in large scale agricultural fields. Apart from the
impracticability at field scale though, the application of high doses of organic fertilisers is not always preferred. It may lead to
several other negative effects such as, leaching or accumulation of nutrients with subsequent negative effects on plants, soil,
water, microorganisms and climate change mitigation (Song et al., 2017; Lazcano et al., 2021). In the case of IT_1 both levels

400 of manure application in the clay soil increased the total soil OC when compared with the unfertilized field (no Nitrogen of any nature). Again, the option of not fertilizing the fields is neither preferred nor common in the current agricultural systems. It should be stressed that, even if fields under mineral fertilization (NPK fertilization but no manure) had been compared with fields that received manure, differences in the OC content probably will not be noticeable, as in Italy the mineral fertilization alone seems not able to maintain the SOC levels (Lugato et al., 2010).

405 Our results coming from the analysis of diverse cropping systems across five European countries give strong indications about the effects of the common factor i.e., the tillage method in the topsoil structure and its dynamics. Nevertheless, our analysis focused only in the top 15 cm soil layer, and this is a limitation that should not be overlooked in the interpretation of the results. The concentration of the residues in the topsoil and the lack of mixing of these residues because of reduced or no tillage throughout the deeper soil horizons have been linked to constraints regarding the maintenance of OC level in the deeper
410 soil horizons. Some authors claim that the SOC balance over the whole soil profile remains unchanged (Blanco-Canqui and Lal, 2008; Chen et al., 2019; Haddaway et al., 2017) or even decreases (Meurer et al., 2018). Nevertheless, there is recent evidence that the adoption of zero or reduced tillage systems also maintains better the existing SOC levels in the deeper horizons (Cooper et al., 2021; Varvel and Wilhelm, 2011), so, increasing the levels of SOC in the topsoil and keeping the subsoil SOC content unchanged, lead to a net SOC increase in the system (Cooper et al., 2021).

415 Based on the above-mentioned results, we agree partly with the suggestion of Chenu et al. (2019) that the best option to increase SOC is to increase the inputs compared to decreasing the outputs through mineralization. However, we need to stress that a combination of increased organic inputs coupled with less intensive tillage methods that will stop or reduce the further release of SOC seems more optimal and realistic. Adding the parameter of reduced tillage intensity, the time required for the microaggregates to be formed properly within the macroaggregates and the OC to get stabilized within these is considered. In
420 addition, most probably by reducing the depth and intensity of tillage, the priming effect will be weakened in systems with addition of fresh OM, leading to higher SOC stocks. This proposal should be validated with further studies including factorial treatments of several tillage intensities and different organic materials, ideally over broad environmental conditions.

5 Conclusions

We analysed seven long-term experiments in five European countries to evaluate soil management practices that could inverse
425 the negative effects of inversion conventional tillage on topsoil structural stability and consequently the SOC content. Following testing several hypotheses and considering the limitation of the sampling depth we conclude that:

- Inversion tillage reduces the mass of topsoil water-stable large macroaggregates, and the OC stabilized in the macroaggregates and microaggregates and this effect increases with increasing tillage intensity, but it does not affect the mass of macroaggregates.
- 430 – Addition of matured exogenous organic materials increases in most cases the topsoil soil OC content in all aggregate fractions even under inversion tillage, but it does not increase the mass of water-stable macroaggregates.

– Incorporation of the previous crops residues into the soil does not increase the mass of the topsoil water-stable macroaggregate nor the occluded POM and does not influence in a consistent way the OC content in the occluded or in free microaggregates.

435 – Increased organic material inputs and reduced soil disturbances do not lead to more water-stable occluded microaggregates but they do lead to increased OC within these (mM-C).

We propose that conventional inversion soil tillage affects topsoil structure in a two-fold way: i) mainly by disrupting the aggregation cycle, thus limiting the time for the development of a stable structure and, subsequently, the time for SOM stabilization within the occluded microaggregates and ii) by increasing the decomposition of OM, and thus limiting the OM that
440 is available to promote aggregation. Therefore, we suggest that there are two ways to protect and/or mitigate soil degradation caused by conventional- inversion tillage in the topsoil: i) to reduce the tillage intensity and allow for the soil structure to build and to stabilize SOM and ii) to increase the OM inputs, combined with non-inversion tillage practices, in order to increase the SOC stocks and diminish the decomposition of old OC. By these ways, the topsoil structural stability, as well as the quality of the topsoil in terms of OC content improves with a subsequent positive effect on the topsoil functions.

Table A1. Description of the study sites and experiments. Table taken from Panagea et al. (2021).

Code	Town, Country	Coordinates (Deci- mal De- grees)	Agro- Climate Zone (Ceglar et al., 2019)	Start of Ex- peri- ment	Soil Type	Name of Experiment	Reference
CZ	Prague- Ruzyně, CZ	50.0880 14.2980	Continental	1995	Silt Loam	Tillage trial	Mühlbachová et al. (2015)
HU_2	Keszthely, HU	46.7346 17.2302	Pannonian	1972	Silt Loam	Soil tillage systems in wheat and maize bi culture	Hoffmann and Kismányoky (2001)
UK	Loddington, UK	52.6089 0.83257	Maritime North	2011	Clay loam	Soil Biology and Soil Health	-
BE	Bierbeek, BE	50.8244 4.79605	Maritime North	1997	Silt Loam	VFG Compost trial	Tits et al. (2014)
HU_1	Keszthely, HU	46.7332 17.2295	Pannonian	1983	Silt Loam	Organic & inorganic fertilization trial- IOSDV	Kismányoky and Tóth (2013)
IT_1c	Legnaro, IT	45.3506 11.9497	Maritime South	1964	Silty Clay Loam	Organic & mineral fertilization trial	Giardini (2004)
IT_1p	Legnaro, IT	45.3506 11.9497	Maritime South	1964	Peat 18% OC initially	Organic & mineral fertilization trial	Giardini (2004)
IT_2c	Legnaro, IT	45.3507 11.9498	Maritime South	1970	Silty Clay Loam	Nitrogen fertilization and crop residue trial	Giardini (2004)
IT_2l	Legnaro, IT	45.3507 11.9498	Maritime South	1970	Silt Loam	Nitrogen fertilization and crop residue trial	Giardini (2004)




Figure A1. Topsoil (0–15 cm) organic carbon expressed in sand free soil for each study site. The OC content is expressed in g C per kg of sand free soil. The error bars represent the standard error. Within an experiment, bars with a different letter differ significantly according to Tukey's test (p<0.05).




Figure A2. Mass percentage of the occluded within the macroaggregates, microaggregates. The mM mass is expressed as percentage of macroaggregates. The error bars represent the standard error. Within an experiment, bars with a different letter differ significantly according to Tukey's test ($p < 0.05$).

Appendix B: Crop yield results




Figure B1. Crop yield in each study site. The error bars represent the standard error. Within an experiment and year, bars with a different letter differ significantly according to Tukey's test ($p<0.05$).

The different treatment categories did not present specific patterns or consistent differences when it came to crop yield (Fig. B1). In the tillage experiments, only the treatments in the CZ experiment caused statistically significant differences. Specifically, both in 2019 and 2020 the minimum tillage yielded more than the conventionally tilled field. Zero tillage did not cause a yield loss when compared with conventional tillage. Addition of manure caused a statistically significant increase in the IT_1c and the IT_1p (only in 2019) when compared to the unfertilized plots. Addition of compost in BE did not cause significant differences in yields. Finally, the incorporation of residues created only a small increase in individual cases such during 2019 in HU_1 and 2020 in the IT_2l experiments.

None of the practices evaluated in this research caused a yield reduction opposing the results of Song et al. (2019), and this offers opportunities for a possible adoption from farmers, but our results also showed that there is a small potential for adopting practices that improve the topsoil SOC and stability, to increase the crop yields. Individual similar field experiments

have shown a positive relationship between SOC and crop yield (D'Hose et al., 2014), however similar to our results Vonk et al. (2020) found a poor association between yield and SOC for common crops in Europe and highlighted the need for a diversified strategy when it comes to motivating farmers to increase their field SOM.

460 *Data availability.* All raw data will be available in the SoilCare community in Zenodo repository (<https://zenodo.org/communities/soilcare>),
after the acceptance of this paper.

Author contributions. Conceptualization, I.S.P; methodology and soil sampling and analysis I.S.P.; data analysis, I.S.P.; visualization, I.S.P.;
writing—original draft preparation, I.S.P.; A.A. and G.W.; writing—review and editing: J.P.; J.D.; M.T., A.E., I.P., A.B., C.S., Z.T., H.K.,
J.B. and P.Č.; supervision, G.W.; J.D.; J.P.; project coordination, G.W.; funding acquisition, G.W.; All authors have read and agreed to the
465 published version of the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We thank all the study sites owners for permitting the use of the experimental fields where this research was done and
providing all types of help during the sampling campaign. We thank the laboratory assistant of the Department of Earth and Environmental
Sciences of the KU Leuven Lore Fondu and the postdoctoral researcher Dr. Zita Kelemen for providing valuable guidance and help during
470 the soil samples analysis.

Financial support. This research was funded by the project “SoilCare: Soil Care for profitable and sustainable crop production in Europe”
from the H2020 Programme under grant agreement no 677407.

References

- Ananyeva, K., Wang, W., Smucker, A. J., Rivers, M. L., and Kravchenko, A. N.: Can intra-aggregate pore structures affect the aggregate's effectiveness in protecting carbon?, *Soil Biology and Biochemistry*, 57, 868–875, <https://doi.org/10.1016/j.soilbio.2012.10.019>, 2013.
- 475 Andruschkewitsch, R., Koch, H. J., and Ludwig, B.: Effect of long-term tillage treatments on the temporal dynamics of water-stable aggregates and on macro-aggregate turnover at three German sites, *Geoderma*, 217–218, 57–64, <https://doi.org/10.1016/j.geoderma.2013.10.022>, 2014.
- Apostolakis, A., Panakoulia, S., Nikolaidis, N. P., and Paranychianakis, N. V.: Shifts in soil structure and soil organic matter in a chronosequence of set-aside fields, *Soil and Tillage Research*, 174, 113–119, <https://doi.org/10.1016/j.still.2017.07.004>, 2017.
- 480 Batista, A., Nunes, M., Pessoa, T., and Libardi, P. L.: Seasonal Variation of the Rhizosphere Soil Aggregation in an Oxisol., SSRN [preprint], <http://dx.doi.org/10.2139/ssrn.4074493>, 2022.
- Berner, A., Hildermann, I., Fließbach, A., Pfiffner, L., Niggli, U., and Mäder, P.: Crop yield and soil fertility response to reduced tillage under organic management, *Soil and Tillage Research*, 101, 89–96, <https://doi.org/10.1016/j.still.2008.07.012>, 2008.
- 485 Berti, A., Morari, F., Dal Ferro, N., Simonetti, G., and Polese, R.: Organic input quality is more important than its quantity: C turnover coefficients in different cropping systems, *European Journal of Agronomy*, 77, 138–145, <https://doi.org/10.1016/j.eja.2016.03.005>, 2016.
- Blanco-Canqui, H. and Lal, R.: No-Tillage and Soil-Profile Carbon Sequestration: An On-Farm Assessment, *Soil Science Society of America Journal*, 72, 693–701, <https://doi.org/10.2136/sssaj2007.0233>, 2008.
- Boddey, R. M., Jantalia, C. P., Conceição, P. C., Zanatta, J. A., Bayer, C., Mielniczuk, J., Dieckow, J., Dos Santos, H. P., Denardin, J. E., Aita, C., Giacomini, S. J., Alves, B. J., and Urquiaga, S.: Carbon accumulation at depth in Ferralsols under zero-till subtropical agriculture, *Global Change Biology*, 16, 784–795, <https://doi.org/10.1111/j.1365-2486.2009.02020.x>, 2010.
- 490 Bolinder, M. A., Crotty, F., Elsen, A., Frac, M., Kismányoky, T., Lipiec, J., Tits, M., Tóth, Z., and Kätterer, T.: The effect of crop residues, cover crops, manures and nitrogen fertilization on soil organic carbon changes in agroecosystems: a synthesis of reviews, *Mitigation and Adaptation Strategies for Global Change*, 25, 929–952, <https://doi.org/10.1007/s11027-020-09916-3>, 2020.
- 495 Camarotto, C., Piccoli, I., Dal Ferro, N., Polese, R., Chiarini, F., Furlan, L., and Morari, F.: Have we reached the turning point? Looking for evidence of SOC increase under conservation agriculture and cover crop practices, *European Journal of Soil Science*, 71, 1050–1063, <https://doi.org/10.1111/ejss.12953>, 2020.
- Cambardella, C. A. and Elliott, E. T.: Particulate Soil Organic-Matter Changes across a Grassland Cultivation Sequence, *Soil Science Society of America Journal*, 56, 777–783, <https://doi.org/10.2136/sssaj1992.03615995005600030017x>, 1992.
- 500 Cannell, R. Q.: Reduced tillage in north-west Europe-A review, *Soil and Tillage Research*, 5, 129–177, [https://doi.org/10.1016/0167-1987\(85\)90028-5](https://doi.org/10.1016/0167-1987(85)90028-5), 1985.
- Cardinael, R., Eglin, T., Guenet, B., Neill, C., Houot, S., and Chenu, C.: Is priming effect a significant process for long-term SOC dynamics? Analysis of a 52-years old experiment, *Biogeochemistry*, 123, 203–219, <https://doi.org/10.1007/s10533-014-0063-2>, 2015.
- Castellano, M. J., Mueller, K. E., Olk, D. C., Sawyer, J. E., and Six, J.: Integrating plant litter quality, soil organic matter stabilization, and the carbon saturation concept, *Global Change Biology*, 21, 3200–3209, <https://doi.org/https://doi.org/10.1111/gcb.12982>, 2015.
- 505 Ceglar, A., Zampieri, M., Toreti, A., and Dentener, F.: Observed Northward Migration of Agro-Climate Zones in Europe Will Further Accelerate Under Climate Change, *Earth's Future*, 7, 1088–1101, <https://doi.org/10.1029/2019EF001178>, 2019.
- Chen, J., Zhu, R., Zhang, Q., Kong, X., and Sun, D.: Reduced-tillage management enhances soil properties and crop yields in a alfalfa-corn rotation: Case study of the Songnen Plain, China, *Scientific Reports*, 9, 1–10, <https://doi.org/10.1038/s41598-019-53602-7>, 2019.

- 510 Chenu, C., Angers, D. A., Barré, P., Derrien, D., Arrouays, D., and Balesdent, J.: Increasing organic stocks in agricultural soils: Knowledge gaps and potential innovations, *Soil and Tillage Research*, 188, 41–52, <https://doi.org/10.1016/j.still.2018.04.011>, 2019.
- Cooper, H. V., Sjögersten, S., Lark, R. M., and Mooney, S. J.: To till or not to till in a temperate ecosystem? Implications for climate change mitigation, *Environmental Research Letters*, 16, <https://doi.org/10.1088/1748-9326/abe74e>, 2021.
- Dal Ferro, N., Piccoli, I., Berti, A., Polese, R., and Morari, F.: Organic carbon storage potential in deep agricultural soil layers: Evidence from long-term experiments in northeast Italy, *Agriculture, ecosystems and environment*, 300, <https://doi.org/10.1016/j.agee.2020.106967>, 2020.
- 515 De Clercq, T., Heiling, M., Dercon, G., Resch, C., Aigner, M., Mayer, L., Mao, Y., Elsen, A., Steier, P., Leifeld, J., and Merckx, R.: Predicting soil organic matter stability in agricultural fields through carbon and nitrogen stable isotopes, *Soil Biology and Biochemistry*, 88, 29–38, <https://doi.org/10.1016/j.soilbio.2015.05.011>, 2015.
- 520 De Clercq, T., Merckx, R., Elsen, A., and Vandendriessche, H.: Impact of long-term compost amendments on soil fertility, soil organic matter fractions and nitrogen mineralization, *Acta Horticulturae*, 1146, 79–86, <https://doi.org/10.17660/ActaHortic.2016.1146.10>, 2016.
- Denef, K., Six, J., Paustian, K., and Merckx, R.: Importance of macroaggregate dynamics in controlling soil carbon stabilization: Short-term effects of physical disturbance induced by dry-wet cycles, *Soil Biology and Biochemistry*, 33, 2145–2153, [https://doi.org/10.1016/S0038-0717\(01\)00153-5](https://doi.org/10.1016/S0038-0717(01)00153-5), 2001.
- 525 Denef, K., Six, J., Merckx, R., and Paustian, K.: Carbon Sequestration in Microaggregates of No-Tillage Soils with Different Clay Mineralogy, *Soil Science Society of America Journal*, 68, 1935–1944, <https://doi.org/10.2136/sssaj2004.1935>, 2004.
- Denef, K., Zotarelli, L., Boddey, R. M., and Six, J.: Microaggregate-associated carbon as a diagnostic fraction for management-induced changes in soil organic carbon in two Oxisols, *Soil Biology and Biochemistry*, 39, 1165–1172, <https://doi.org/10.1016/j.soilbio.2006.12.024>, 2007.
- 530 Devine, S., Markewitz, D., Hendrix, P., and Coleman, D.: Soil aggregates and associated organic matter under conventional tillage, no-tillage, and forest succession after three decades, *PLoS ONE*, 9, 1–12, <https://doi.org/10.1371/journal.pone.0084988>, 2014.
- D’Hose, T., Cougnon, M., De Vlieghe, A., Vandecasteele, B., Viaene, N., Cornelis, W., Van Bockstaele, E., and Reheul, D.: The positive relationship between soil quality and crop production: A case study on the effect of farm compost application, *Applied Soil Ecology*, 75, 189–198, <https://doi.org/10.1016/j.apsoil.2013.11.013>, 2014.
- 535 Dignac, M. F., Derrien, D., Barré, P., Barot, S., Cécillon, L., Chenu, C., Chevallier, T., Freschet, G. T., Garnier, P., Guenet, B., Hedde, M., Klumpp, K., Lashermes, G., Maron, P. A., Nunan, N., Roumet, C., and Basile-Doelsch, I.: Increasing soil carbon storage: mechanisms, effects of agricultural practices and proxies. A review, *Agronomy for Sustainable Development*, 37, <https://doi.org/10.1007/s13593-017-0421-2>, 2017.
- Dimoyiannis, D.: Seasonal soil aggregate stability variation in relation to rainfall and temperature under Mediterranean conditions, *Earth Surface Processes and Landforms*, 34, 860–866, <https://doi.org/10.1002/esp.1785>, 2009.
- 540 Du, Z., Angers, D. A., Ren, T., Zhang, Q., and Li, G.: The effect of no-till on organic C storage in Chinese soils should not be overemphasized: A meta-analysis, *Agriculture, Ecosystems and Environment*, 236, 1–11, <https://doi.org/10.1016/j.agee.2016.11.007>, 2017.
- Elliott, E.: Aggregate Structure and Carbon, Nitrogen, and Phosphorus in Native and Cultivated Soils, *Soil Science Society of America Journal*, 50, 627–633, <https://doi.org/10.2136/sssaj1986.03615995005000030017x>, 1986.
- 545 Elliott, E. T., Palm, C. A., Reuss, D. E., and Monz, C. A.: Organic matter contained in soil aggregates from a tropical chronosequence: correction for sand and light fraction, *Agriculture, Ecosystems and Environment*, 34, 443–451, [https://doi.org/10.1016/0167-8809\(91\)90127-J](https://doi.org/10.1016/0167-8809(91)90127-J), 1991.

- Fontaine, S., Barot, S., Barré, P., Bdioui, N., Mary, B., and Rumpel, C.: Stability of organic carbon in deep soil layers controlled by fresh carbon supply, *Nature*, 450, 277–280, <https://doi.org/10.1038/nature06275>, 2007.
- 550 Fuentes, M., Govaerts, B., De León, F., Hidalgo, C., Dendooven, L., Sayre, K. D., and Etchevers, J.: Fourteen years of applying zero and conventional tillage, crop rotation and residue management systems and its effect on physical and chemical soil quality, *European Journal of Agronomy*, 30, 228–237, <https://doi.org/10.1016/j.eja.2008.10.005>, 2009.
- Fuentes, M., Hidalgo, C., Etchevers, J., de León, F., Guerrero, A., Dendooven, L., Verhulst, N., and Govaerts, B.: Conservation agriculture, increased organic carbon in the top-soil macro-aggregates and reduced soil CO₂ emissions, *Plant and Soil*, 355, 183–197, 555 <https://doi.org/10.1007/s11104-011-1092-4>, 2012.
- Ghimire, R., Norton, U., Bista, P., Obour, A. K., and Norton, J. B.: Soil organic matter, greenhouse gases and net global warming potential of irrigated conventional, reduced-tillage and organic cropping systems, *Nutrient Cycling in Agroecosystems*, 107, 49–62, <https://doi.org/10.1007/s10705-016-9811-0>, 2017.
- Giannakis, G., Panakoulia, S., Nikolaidis, N., and Paranychianakis, N.: Simulating Soil Fertility Restoration Using the CAST Model, *Procedia Earth and Planetary Science*, 10, 325–329, <https://doi.org/10.1016/j.proeps.2014.08.027>, 2014.
- 560 Giardini, L.: Ongoing Trials, in: *Productivity and Sustainability of Different Cropping Systems. 40 Years of Experiments in Veneto Region (Italy)*, pp. 89–97, Bologna, 2004.
- Gross, A. and Glaser, B.: Meta-analysis on how manure application changes soil organic carbon storage, *Scientific Reports*, 11, 1–13, <https://doi.org/10.1038/s41598-021-82739-7>, 2021.
- 565 Haddaway, N. R., Hedlund, K., Jackson, L. E., Kätterer, T., Lugato, E., Thomsen, I. K., Jørgensen, H. B., and Söderström, B.: How does tillage intensity affect soil organic carbon? A systematic review, *Environmental Evidence*, 6:30, 1–8, <https://doi.org/10.1186/s13750-017-0108-9>, 2017.
- Hobson, D., Harty, M., Tracy, S., and McDonnell, K.: The effect of tillage depth and traffic management on soil properties and root development during two growth stages of winter wheat (*Triticum aestivum* L.), *SOIL Discussions*, 2021, 1–32, <https://doi.org/10.5194/soil-2021-129>, 2021.
- 570 Hoffmann, S. and Kismányoky, T.: Soil fertility in a long-term fertilizer trial with different tillage systems, *Archives of Agronomy and Soil Science*, 46, 239–250, <https://doi.org/10.1080/03650340109366175>, 2001.
- Jastrow, J. D., Amonette, J. E., and Bailey, V. L.: Mechanisms controlling soil carbon turnover and their potential application for enhancing carbon sequestration, *Climatic Change*, 80, 5–23, <https://doi.org/10.1007/s10584-006-9178-3>, 2007.
- 575 Jat, H. S., Datta, A., Choudhary, M., Yadav, A. K., Choudhary, V., Sharma, P. C., Gathala, M. K., Jat, M. L., and McDonald, A.: Effects of tillage, crop establishment and diversification on soil organic carbon, aggregation, aggregate associated carbon and productivity in cereal systems of semi-arid Northwest India, *Soil and Tillage Research*, 190, 128–138, <https://doi.org/10.1016/j.still.2019.03.005>, 2019.
- Jirků, V., Kodešová, R., Mühlhanslová, M., and Žigová, A.: Seasonal variability of soil structure and soil hydraulic properties, in: *19th World Congress of Soil Science, Soil Solutions for a Changing World*, Brisbane, Australia, 2010.
- 580 Karlen, D. L., Wollenhaupt, N. C., Erbach, D. C., Berry, E. C., Swan, J. B., Eash, N. S., and Jordahl, J. L.: Crop residue effects on soil quality following 10-years of no-till corn, *Soil and Tillage Research*, 31, 149–167, [https://doi.org/10.1016/0167-1987\(94\)90077-9](https://doi.org/10.1016/0167-1987(94)90077-9), 1994.
- Karlen, D. L., Cambardella, C. A., Kovar, J. L., and Colvin, T. S.: Soil quality response to long-term tillage and crop rotation practices, *Soil and Tillage Research*, 133, 54–64, <https://doi.org/10.1016/j.still.2013.05.013>, 2013.

- Kätterer, T., Bolinder, M. A., Andrén, O., Kirchmann, H., and Menichetti, L.: Roots contribute more to refractory soil organic matter than above-ground crop residues, as revealed by a long-term field experiment, *Agriculture, Ecosystems and Environment*, 141, 184–192, <https://doi.org/10.1016/j.agee.2011.02.029>, 2011.
- Kay, B. D. and VandenBygaart, A. J.: Conservation tillage and depth stratification of porosity and soil organic matter, *Soil and Tillage Research*, 66, 107–118, [https://doi.org/10.1016/S0167-1987\(02\)00019-3](https://doi.org/10.1016/S0167-1987(02)00019-3), 2002.
- King, A. E., Congreves, K. A., Deen, B., Dunfield, K. E., Voroney, R. P., and Wagner-Riddle, C.: Quantifying the relationships between soil fraction mass, fraction carbon, and total soil carbon to assess mechanisms of physical protection, *Soil Biology and Biochemistry*, 135, 95–107, <https://doi.org/10.1016/j.soilbio.2019.04.019>, 2019.
- Kismányoky, T. and Tóth, Z.: Effect of mineral and organic fertilization on soil organic carbon content as well as on grain production of cereals in the IOSDV (ILTE) long-term field experiment, Keszthely, Hungary, *Archives of Agronomy and Soil Science*, 59, 1121–1131, <https://doi.org/10.1080/03650340.2012.712208>, 2013.
- Kong, A. Y. Y., Six, J., Bryant, D. C., Denison, R. F., and van Kessel, C.: The Relationship between Carbon Input, Aggregation, and Soil Organic Carbon Stabilization in Sustainable Cropping Systems, *Soil Science Society of America Journal*, 69, 1078–1085, <https://doi.org/10.2136/sssaj2004.0215>, 2005.
- Kotronakis, M., Giannakis, G. V., Nikolaidis, N. P., Rowe, E. C., Valstar, J., Paranychianakis, N. V., and Banwart, S. A.: Modeling the Impact of Carbon Amendments on Soil Ecosystem Functions Using the 1D-ICZ Model, vol. 142, Elsevier Inc., 1 edn., <https://doi.org/10.1016/bs.agron.2016.10.010>, 2017.
- Lavallee, J. M., Soong, J. L., and Cotrufo, M. F.: Conceptualizing soil organic matter into particulate and mineral-associated forms to address global change in the 21st century, *Global Change Biology*, 26, 261–273, <https://doi.org/10.1111/gcb.14859>, 2020.
- Lazcano, C., Zhu-Barker, X., and Decock, C.: Effects of Organic Fertilizers on the Soil Microorganisms Responsible for N₂O Emissions: A Review, *Microorganisms*, 9, <https://doi.org/10.3390/microorganisms9050983>, 2021.
- Lenth, R.: Estimated Marginal Means, aka Least-Squares Means., <https://cran.r-project.org/package=%0Aemmeans>, 2020.
- Li, Y., Li, Z., Chang, S. X., Cui, S., Jagadamma, S., Zhang, Q., and Cai, Y.: Residue retention promotes soil carbon accumulation in minimum tillage systems: Implications for conservation agriculture, *Science of the Total Environment*, 740, 140–147, <https://doi.org/10.1016/j.scitotenv.2020.140147>, 2020.
- Lin, Y., Ye, G., Kuzyakov, Y., Liu, D., Fan, J., and Ding, W.: Long-term manure application increases soil organic matter and aggregation, and alters microbial community structure and keystone taxa, *Soil Biology and Biochemistry*, 134, 187–196, <https://doi.org/10.1016/j.soilbio.2019.03.030>, 2019.
- Liu, X. J. A., Finley, B. K., Mau, R. L., Schwartz, E., Dijkstra, P., Bowker, M. A., and Hungate, B. A.: The soil priming effect: Consistent across ecosystems, elusive mechanisms, *Soil Biology and Biochemistry*, 140, 107–117, <https://doi.org/10.1016/j.soilbio.2019.107617>, 2020.
- Lugato, E., Simonetti, G., Morari, F., Nardi, S., Berti, A., and Giardini, L.: Distribution of organic and humic carbon in wet-sieved aggregates of different soils under long-term fertilization experiment, *Geoderma*, 157, 80–85, <https://doi.org/10.1016/j.geoderma.2010.03.017>, 2010.
- Luo, Z., Wang, E., and Sun, O. J.: Soil carbon change and its responses to agricultural practices in Australian agro-ecosystems: A review and synthesis, *Geoderma*, 155, 211–223, <https://doi.org/10.1016/j.geoderma.2009.12.012>, 2010.
- Meurer, K. H., Haddaway, N. R., Bolinder, M. A., and Kätterer, T.: Tillage intensity affects total SOC stocks in boreo-temperate regions only in the topsoil—A systematic review using an ESM approach, *Earth-Science Reviews*, 177, 613–622, <https://doi.org/https://doi.org/10.1016/j.earscirev.2017.12.015>, 2018.

- Mikha, M. M. and Rice, C. W.: Tillage and Manure Effects on Soil and Aggregate-Associated Carbon and Nitrogen, *Soil Science Society of America Journal*, 68, 809–816, <https://doi.org/10.2136/sssaj2004.8090>, 2004.
- 625 Minasny, B., Malone, B. P., McBratney, A. B., Angers, D. A., Arrouays, D., Chambers, A., Chaplot, V., Chen, Z. S., Cheng, K., Das, B. S., Field, D. J., Gimona, A., Hedley, C. B., Hong, S. Y., Mandal, B., Marchant, B. P., Martin, M., McConkey, B. G., Mulder, V. L., O'Rourke, S., Richer-de Forges, A. C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I., Stolbovoy, V., Stockmann, U., Sulaeman, Y., Tsui, C. C., Vågen, T. G., van Wesemael, B., and Winowiecki, L.: Soil carbon 4 per mille, *Geoderma*, 292, 59–86, <https://doi.org/10.1016/j.geoderma.2017.01.002>, 2017.
- 630 Modak, K., Biswas, D. R., Ghosh, A., Pramanik, P., Das, T. K., Das, S., Kumar, S., Krishnan, P., and Bhattacharyya, R.: Zero tillage and residue retention impact on soil aggregation and carbon stabilization within aggregates in subtropical India, *Soil and Tillage Research*, 202, 104 649, <https://doi.org/10.1016/j.still.2020.104649>, 2020.
- Mondal, S. and Chakraborty, D.: Global meta-analysis suggests that no-tillage favourably changes soil structure and porosity, *Geoderma*, 405, 115 443, <https://doi.org/10.1016/j.geoderma.2021.115443>, 2022.
- 635 Mühlbachová, G., Kusá, H., and Růžek, P.: Soil characteristics and crop yields under different tillage techniques, *Plant, Soil and Environment*, 61, 566–572, <https://doi.org/10.17221/567/2015-PSE>, 2015.
- Nimmo, J.: *Aggregation: Physical Aspects*, Published by Elsevier Inc., <https://doi.org/10.1016/b978-0-12-409548-9.05087-9>, 2013.
- Oades, J.: Soil organic matter and structural stability : mechanisms and implications for management A, *Plant and Soil*, 76, 319–337, <https://doi.org/10.1007/BF02205590>, 1984.
- 640 Ogle, S. M., Swan, A., and Paustian, K.: No-till management impacts on crop productivity, carbon input and soil carbon sequestration, *Agriculture, Ecosystems and Environment*, 149, 37–49, <https://doi.org/10.1016/j.agee.2011.12.010>, 2012.
- Pan, J., Wang, J., Zhang, R., Tian, D., Cheng, X., Wang, S., Chen, C., Yang, L., and Niu, S.: Microaggregates regulated by edaphic properties determine the soil carbon stock in Tibetan alpine grasslands, *Catena*, 206, 105 570, <https://doi.org/10.1016/j.catena.2021.105570>, 2021.
- 645 Panagea, I. S., Berti, A., Čermak, P., Diels, J., Elsen, A., Kusá, H., Piccoli, I., Poesen, J., Stoate, C., Tits, M., Toth, Z., and Wyseure, G.: Soil Water Retention as Affected by Management Induced Changes of Soil Organic Carbon: Analysis of Long-Term Experiments in Europe, <https://doi.org/10.3390/land10121362>, 2021.
- Paul, B. K., Vanlauwe, B., Ayuke, F., Gassner, A., Hoogmoed, M., Hurisso, T. T., Koala, S., Lelei, D., Ndabamenye, T., Six, J., and Pulleman, M. M.: Medium-term impact of tillage and residue management on soil aggregate stability, soil carbon and crop productivity, *Agriculture, Ecosystems and Environment*, 164, 14–22, <https://doi.org/10.1016/j.agee.2012.10.003>, 2013.
- 650 Paustian, K., Andrén, O., Janzen, H. H., Lal, R., Smith, P., Tian, G., Tiessen, H., Van Noordwijk, M., and Woormer, P. L.: Agricultural soils as a sink, *Soil Use and Management*, 13, 230–244, 1997.
- Peigné, J., Vian, J. F., Payet, V., and Saby, N. P.: Soil fertility after 10 years of conservation tillage in organic farming, *Soil and Tillage Research*, 175, 194–204, <https://doi.org/10.1016/j.still.2017.09.008>, 2018.
- 655 Piazza, G., Pellegrino, E., Moscatelli, M. C., and Ercoli, L.: Long-term conservation tillage and nitrogen fertilization effects on soil aggregate distribution, nutrient stocks and enzymatic activities in bulk soil and occluded microaggregates, *Soil and Tillage Research*, 196, 104 482, <https://doi.org/10.1016/j.still.2019.104482>, 2020.
- Piccoli, I., Chiarini, F., Carletti, P., Furlan, L., Lazzaro, B., Nardi, S., Berti, A., Sartori, L., Dalconi, M. C., and Morari, F.: Disentangling the effects of conservation agriculture practices on the vertical distribution of soil organic carbon. Evidence of poor carbon sequestration in North- Eastern Italy, *Agriculture, Ecosystems and Environment*, 230, 68–78, <https://doi.org/10.1016/j.agee.2016.05.035>, 2016.

- Piccoli, I., Lazzaro, B., Furlan, L., Berti, A., and Morari, F.: Examining crop root apparatus traits in a maize-soybean-winter wheat rotation under conservation agriculture management, *European Journal of Agronomy*, 122, 126–171, <https://doi.org/10.1016/j.eja.2020.126171>, 2021.
- Plaza-Bonilla, D., Cantero-Martínez, C., and Álvaro-Fuentes, J.: Tillage effects on soil aggregation and soil organic carbon profile distribution under Mediterranean semi-arid conditions, *Soil Use and Management*, 26, 465–474, <https://doi.org/10.1111/j.1475-2743.2010.00298.x>, 2010.
- Poeplau, C., Don, A., Six, J., Kaiser, M., Benbi, D., Chenu, C., Cotrufo, M. F., Derrien, D., Gioacchini, P., Grand, S., Gregorich, E., Griepentrog, M., Gunina, A., Haddix, M., Kuzyakov, Y., Kühnel, A., Macdonald, L. M., Soong, J., Trigalet, S., Vermeire, M. L., Rovira, P., van Wesemael, B., Wiesmeier, M., Yeasmin, S., Yevdokimov, I., and Nieder, R.: Isolating organic carbon fractions with varying turnover rates in temperate agricultural soils – A comprehensive method comparison, *Soil Biology and Biochemistry*, 125, 10–26, <https://doi.org/10.1016/j.soilbio.2018.06.025>, 2018.
- Powelson, D. S., Glendining, M. J., Coleman, K., and Whitmore, A. P.: Implications for soil properties of removing cereal straw: Results from long-term studies, *Agronomy Journal*, 103, 279–287, <https://doi.org/10.2134/agronj2010.0146s>, 2011.
- Pu, C., Kan, Z. R., Liu, P., Ma, S. T., Qi, J. Y., Zhao, X., and Zhang, H. L.: Residue management induced changes in soil organic carbon and total nitrogen under different tillage practices in the North China Plain, *Journal of Integrative Agriculture*, 18, 1337–1347, [https://doi.org/10.1016/S2095-3119\(18\)62079-9](https://doi.org/10.1016/S2095-3119(18)62079-9), 2019.
- R Core Team: R: A Language and Environment for Statistical Computing, <https://www.r-project.org/>, 2019.
- RStudio Team: RStudio: Integrated Development Environment for R, <http://www.rstudio.com/>, 2016.
- Rumpel, C. and Kögel-Knabner, I.: Deep soil organic matter—a key but poorly understood component of terrestrial C cycle, *Plant and Soil*, 338, 143–158, <https://doi.org/10.1007/s11104-010-0391-5>, 2011.
- Rusinamhodzi, L., Corbeels, M., van Wijk, M. T., Rufino, M. C., Nyamangara, J., and Giller, K. E.: A meta-analysis of long-term effects of conservation agriculture on maize grain yield under rain-fed conditions, *Agronomy for Sustainable Development*, 31, 657, <https://doi.org/10.1007/s13593-011-0040-2>, 2011.
- Sartori, F., Piccoli, I., Polese, R., and Berti, A.: Transition to conservation agriculture: how tillage intensity and covering affect soil physical parameters, *SOIL Discussions*, 2021, 1–20, <https://doi.org/10.5194/soil-2021-113>, 2021.
- Schils, R., Olesen, J. E., Kersebaum, K. C., Rijk, B., Oberforster, M., Kalyada, V., Khitrykau, M., Gobin, A., Kirchev, H., Manolova, V., Manolov, I., Trnka, M., Hlavinka, P., Paluoso, T., Peltonen-Sainio, P., Jauhiainen, L., Lorgeou, J., Marrou, H., Danalatos, N., Archontoulis, S., Fodor, N., Spink, J., Roggero, P. P., Bassu, S., Pulina, A., Seehusen, T., Uhlen, A. K., Żyłowska, K., Nieróbca, A., Kozyra, J., Silva, J. V., Maçãs, B. M., Coutinho, J., Ion, V., Takáč, J., Mínguez, M. I., Eckersten, H., Levy, L., Herrera, J. M., Hiltbrunner, J., Kryvobok, O., Kryvoshein, O., Sylvester-Bradley, R., Kindred, D., Topp, C. F., Boogaard, H., de Groot, H., Lesschen, J. P., van Bussel, L., Wolf, J., Zijlstra, M., van Loon, M. P., and van Ittersum, M. K.: Cereal yield gaps across Europe, *European Journal of Agronomy*, 101, 109–120, <https://doi.org/10.1016/j.eja.2018.09.003>, 2018.
- Schneider, F., Don, A., Hennings, I., Schmittmann, O., and Seidel, S. J.: The effect of deep tillage on crop yield – What do we really know?, *Soil and Tillage Research*, 174, 193–204, <https://doi.org/10.1016/j.still.2017.07.005>, 2017.
- Schrumpf, M., Kaiser, K., Guggenberger, G., Persson, T., Kögel-Knabner, I., and Schulze, E. D.: Storage and stability of organic carbon in soils as related to depth, occlusion within aggregates, and attachment to minerals, *Biogeosciences*, 10, 1675–1691, <https://doi.org/10.5194/bg-10-1675-2013>, 2013.

- Sheehy, J., Regina, K., Alakukku, L., and Six, J.: Impact of no-till and reduced tillage on aggregation and aggregate-associated carbon in Northern European agroecosystems, *Soil and Tillage Research*, 150, 107–113, <https://doi.org/10.1016/j.still.2015.01.015>, 2015.
- Six, J. and Paustian, K.: Aggregate-associated soil organic matter as an ecosystem property and a measurement tool, *Soil Biology and Biochemistry*, 68, A4–A9, <https://doi.org/10.1016/j.soilbio.2013.06.014>, 2014.
- 700 Six, J., Elliott, E., Paustian, K., and Doran, J. W.: Aggregation and Soil Organic Matter Accumulation in Cultivated and Native Grassland Soils, *Soil Science Society of America Journal*, 62, 1367–1377, <https://doi.org/10.2136/sssaj1998.03615995006200050032x>, 1998.
- Six, J., Elliott, E. T., and Paustian, K.: Soil macroaggregate turnover and microaggregate formation: A mechanism for C sequestration under no-tillage agriculture, *Soil Biology and Biochemistry*, 32, 2099–2103, [https://doi.org/10.1016/S0038-0717\(00\)00179-6](https://doi.org/10.1016/S0038-0717(00)00179-6), 2000a.
- Six, J., Paustian, K., Elliott, E. T., and Combrink, C.: Soil Structure and Organic Matter I. Distribution of Aggregate-Size Classes and
705 Aggregate-Associated Carbon, *Soil Science Society of America Journal*, 64, 681–689, <https://doi.org/10.2136/sssaj2000.642681x>, 2000b.
- Six, J., Callewaert, P., Lenders, S., De Gryze, S., Morris, S. J., Gregorich, E. G., Paul, E. A., and Paustian, K.: Measuring and Understanding Carbon Storage in Afforested Soils by Physical Fractionation, *Soil Science Society of America Journal*, 66, 1981–1987, <https://doi.org/10.2136/sssaj2002.1981>, 2002.
- Song, K., Xue, Y., Zheng, X., Lv, W., Qiao, H., Qin, Q., and Yang, J.: Effects of the continuous use of organic manure and chemical fertilizer
710 on soil inorganic phosphorus fractions in calcareous soil, *Scientific Reports*, 7, 1164, <https://doi.org/10.1038/s41598-017-01232-2>, 2017.
- Song, K., Zheng, X., Lv, W., Qin, Q., Sun, L., Zhang, H., and Xue, Y.: Effects of tillage and straw return on water-stable aggregates, carbon stabilization and crop yield in an estuarine alluvial soil, *Scientific Reports*, 9, 1–11, <https://doi.org/10.1038/s41598-019-40908-9>, 2019.
- Struijk, M., Whitmore, A. P., Mortimer, S. R., and Sizmur, T.: Obtaining more benefits from crop residues as soil amendments by application as chemically heterogeneous mixtures, *SOIL*, 6, 467–481, <https://doi.org/10.5194/soil-6-467-2020>, 2020.
- 715 Thomsen, I. K., Olesen, J. E., Møller, H. B., Sørensen, P., and Christensen, B. T.: Carbon dynamics and retention in soil after anaerobic digestion of dairy cattle feed and faeces, *Soil Biology and Biochemistry*, 58, 82–87, <https://doi.org/10.1016/j.soilbio.2012.11.006>, 2013.
- Tits, M., Elsen, A., Bries, J., and Vandendriessche, H.: Short-term and long-term effects of vegetable, fruit and garden waste compost applications in an arable crop rotation in Flanders, *Plant and Soil*, 376, 43–59, <https://doi.org/10.1007/s11104-012-1318-0>, 2014.
- Trumbore, S.: Radiocarbon and soil carbon dynamics, *Annual Review of Earth and Planetary Sciences*, 37, 47–66, <https://doi.org/10.1146/annurev.earth.36.031207.124300>, 2009.
- 720 Van den Putte, A., Govers, G., Diels, J., Gillijns, K., and Demuzere, M.: Assessing the effect of soil tillage on crop growth: A meta-regression analysis on European crop yields under conservation agriculture, *European Journal of Agronomy*, 33, 231–241, <https://doi.org/10.1016/j.eja.2010.05.008>, 2010.
- Varvel, G. E. and Wilhelm, W. W.: No-tillage increases soil profile carbon and nitrogen under long-term rainfed cropping systems, *Soil and
725 Tillage Research*, 114, 28–36, <https://doi.org/10.1016/j.still.2011.03.005>, 2011.
- Virto, I., Barré, P., Burlot, A., and Chenu, C.: Carbon input differences as the main factor explaining the variability in soil organic C storage in no-tilled compared to inversion tilled agrosystems, *Biogeochemistry*, 108, 17–26, <https://doi.org/10.1007/s10533-011-9600-4>, 2012.
- Vonk, W. J., van Ittersum, M. K., Reidsma, P., Zavattaro, L., Bechini, L., Guzmán, G., Pronk, A., Spiegel, H., Steinmann, H. H., Ruyschaert, G., and Hijbeek, R.: European survey shows poor association between soil organic matter and crop yields, *Nutrient Cycling in
730 Agroecosystems*, 118, 325–334, <https://doi.org/10.1007/s10705-020-10098-2>, 2020.
- Wang, H., Boutton, T. W., Xu, W., Hu, G., Jiang, P., and Bai, E.: Quality of fresh organic matter affects priming of soil organic matter and substrate utilization patterns of microbes, *Scientific Reports*, 5, <https://doi.org/10.1038/srep10102>, 2015.

- Webster, R.: Analysis of variance, inference, multiple comparisons and sampling effects in soil research, *European Journal of Soil Science*, 58, 74–82, <https://doi.org/10.1111/j.1365-2389.2006.00801.x>, 2007.
- 735 Wen, Y., Tang, Y., Wen, J., Wang, Q., Bai, L., Wang, Y., Su, S., Wu, C., Lv, J., and Zeng, X.: Variation of intra-aggregate organic carbon affects aggregate formation and stability during organic manure fertilization in a fluvo-aquic soil, *Soil Use and Management*, 37, 151–163, <https://doi.org/10.1111/sum.12676>, 2021.
- Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*, Springer: New York, NY, USA, 2016.
- Wiesmeier, M., Hübner, R., and Kögel-Knabner, I.: Stagnating crop yields: An overlooked risk for the carbon balance of agricultural soils?, *Science of the Total Environment*, 536, 1045–1051, <https://doi.org/10.1016/j.scitotenv.2015.07.064>, 2015.
- 740 Williams, D. M., Blanco-Canqui, H., Francis, C. A., and Galusha, T. D.: Organic farming and soil physical properties: An assessment after 40 years, *Agronomy Journal*, 109, 600–609, <https://doi.org/10.2134/agronj2016.06.0372>, 2017.
- Wortmann, C. S. and Shapiro, C. A.: The effects of manure application on soil aggregation, *Nutrient Cycling in Agroecosystems*, 80, 173–180, <https://doi.org/10.1007/s10705-007-9130-6>, 2008.
- 745 Xu, J., Han, H., Ning, T., Li, Z., and Lal, R.: Long-term effects of tillage and straw management on soil organic carbon, crop yield, and yield stability in a wheat-maize system, *Field Crops Research*, 233, 33–40, <https://doi.org/10.1016/j.fcr.2018.12.016>, 2019.
- Yang, X.-M. and Wander, M. M.: Temporal changes in dry aggregate size and stability: tillage and crop effects on a silty loam Mollisol in Illinois, *Soil and Tillage Research*, 49, 173–183, [https://doi.org/https://doi.org/10.1016/S0167-1987\(98\)00170-6](https://doi.org/https://doi.org/10.1016/S0167-1987(98)00170-6), 1998.
- Yin, Y., Wang, L., Liang, C., Xi, F., Pei, Z., and Du, L.: Soil Aggregate Stability and Iron and Aluminium Oxide Contents Under Different Fertiliser Treatments in a Long-Term Solar Greenhouse Experiment, *Pedosphere*, 26, 760–767, [https://doi.org/10.1016/S1002-0160\(15\)60086-8](https://doi.org/10.1016/S1002-0160(15)60086-8), 2016.
- Zhang, J., Wei, D., Zhou, B., Zhang, L., Hao, X., Zhao, S., Xu, X., He, P., Zhao, Y., Qiu, S., and Zhou, W.: Responses of soil aggregation and aggregate-associated carbon and nitrogen in black soil to different long-term fertilization regimes, *Soil and Tillage Research*, 213, <https://doi.org/10.1016/j.still.2021.105157>, 2021.
- 755 Zhang, P., Wei, T., Jia, Z., Han, Q., and Ren, X.: Soil aggregate and crop yield changes with different rates of straw incorporation in semiarid areas of northwest China, *Geoderma*, 230-231, 41–49, <https://doi.org/10.1016/j.geoderma.2014.04.007>, 2014.
- Zhao, H., Shar, A. G., Li, S., Chen, Y., Shi, J., Zhang, X., and Tian, X.: Effect of straw return mode on soil aggregation and aggregate carbon content in an annual maize-wheat double cropping system, *Soil and Tillage Research*, 175, 178–186, <https://doi.org/10.1016/j.still.2017.09.012>, 2018.
- 760 Zheng, H., Liu, W., Zheng, J., Luo, Y., Li, R., Wang, H., and Qi, H.: Effect of long-term tillage on soil aggregates and aggregate-associated carbon in black soil of northeast China, *PLoS ONE*, 13, 1–18, <https://doi.org/10.1371/journal.pone.0199523>, 2018.